

Time Limit: 2 sec / Memory Limit: 256 MB

Score : 300 points

Problem Statement

You are given an undirected connected graph with N vertices and M edges that does not contain self-loops and double edges.

The i -th edge ($1 \leq i \leq M$) connects Vertex a_i and Vertex b_i .

An edge whose removal disconnects the graph is called a *bridge*.

Find the number of the edges that are bridges among the M edges.

Notes

- A *self-loop* is an edge i such that $a_i = b_i$ ($1 \leq i \leq M$).
- *Double edges* are a pair of edges i, j such that $a_i = a_j$ and $b_i = b_j$ ($1 \leq i < j \leq M$).
- An undirected graph is said to be *connected* when there exists a path between every pair of vertices.

Constraints

- $2 \leq N \leq 50$
- $N - 1 \leq M \leq \min(N(N-1)/2, 50)$
- $1 \leq a_i < b_i \leq N$
- The given graph does not contain self-loops and double edges.
- The given graph is connected.

Input

Input is given from Standard Input in the following format:

```
N M
a1 b1
a2 b2
:
aM bM
```

Output

Print the number of the edges that are bridges among the M edges.

Sample Input 1

[Copy](#)

```
7 7
1 3
2 7
3 4
4 5
4 6
5 6
6 7
```

[Copy](#)


Sample Output 1

[Copy](#)

```
4
```

[Copy](#)

The figure below shows the given graph:



The edges shown in red are bridges. There are four of them.

Sample Input 2

[Copy](#)

```
3 3
1 2
1 3
2 3
```

[Copy](#)

Sample Output 2

[Copy](#)

```
0
```

[Copy](#)

It is possible that there is no bridge.

Sample Input 3

[Copy](#)

```
6 5
1 2
2 3
3 4
4 5
5 6
```

[Copy](#)

Sample Output 3

[Copy](#)

```
5
```

[Copy](#)

It is possible that every edge is a bridge.