

# Calculating the Radial Component of the Tidal Field, I think

Daniel Martens

May 20, 2016

We want to calculate:

$$S_{rr} = \hat{r}^i \hat{r}^j \nabla_i \nabla_j \nabla^{-2} \psi_\alpha \quad (1)$$

In the basis where:

$$\psi_\alpha = j_l(k_\alpha r) Y_{l_\alpha m_\alpha}(\theta, \phi) \quad (2)$$

Because we are in k-space, the inverse Laplacian operator is simple:

$$\begin{aligned} \nabla^2 \psi_\alpha &= -k_\alpha^2 \psi_\alpha \\ \nabla^{-2} \psi_\alpha &= \frac{-1}{k_\alpha^2} \psi_\alpha \end{aligned} \quad (3)$$

Before acting on the remaining terms with the directional gradients, it helps to first dot with the directional unit vectors:

$$\begin{aligned} \hat{r}^i \hat{r}^j \nabla_i \nabla_j &= \hat{r}^i (\nabla_i \hat{r}^j + [\hat{r}^j, \nabla_i]) \nabla_j \\ &= \hat{r}^i \nabla_i \hat{r}^j \nabla_j - \hat{r}^i \nabla_i \hat{r}^j \nabla_j = \frac{d^2}{dr^2} - \hat{r}^i \left( \frac{1}{r} (\delta_i^j - \hat{r}^j \hat{r}_i) \right) \nabla_j \\ &= \frac{d^2}{dr^2} \end{aligned} \quad (4)$$

So we see that:

$$S_{rr} = \frac{-1}{k_\alpha^2} Y_{l_\alpha m_\alpha}(\theta, \phi) \frac{d^2}{dr^2} [j_l(k_\alpha r)] \quad (5)$$

Using the recursion relation of the spherical bessel functions of the first kind:

$$2 \frac{dj_l(x)}{dx} = j_{l-1}(x) - j_{l+1}(x) \quad (6)$$

And the chain rule:

$$\frac{d}{dr} j_l(k_\alpha r) = k_\alpha \frac{d}{d(k_\alpha r)} j_l(k_\alpha r) \quad (7)$$

We end with our result:

$$S_{rr} = \frac{Y_{l_\alpha m_\alpha}(\theta, \phi)}{4} [2j_l(k_\alpha r) - j_{l+2}(k_\alpha r) - j_{l-2}(k_\alpha r)] \quad (8)$$