

Problem 8): For a public key encryption scheme $\Pi = (\text{Gen}, \text{Enc}, \text{Dec})$, we define **CPA** security according to the probability obtaining a secure result, as defined in the privacy experiment $\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}$. This experiment goes as follows

The LR-oracle experiment $\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n)$

1. $\text{Gen}(1^n)$ is run to obtain keys (pk, sk) .
2. A uniform bit $b \in \{0, 1\}$ is chosen.
3. The adversary \mathcal{A} is given input pk and oracle access to $\text{LR}_{pk, b}(\cdot, \cdot)$.
4. The adversary \mathcal{A} outputs a bit b' .
5. The adversary \mathcal{A} is defined to be 1 if $b' = b$, and 0 otherwise. If $\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n) = 1$, we say that \mathcal{A} **succeeds**.

Using this definition for the experiment $\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}$, we say that the encryption scheme Π is secure if the probability of \mathcal{A} succeeding, $\Pr[\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n) = 1]$ satisfies the condition

$$\Pr[\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n) = 1] \leq \frac{1}{2} + \text{negl}(n) \quad (8.1)$$

where $\text{negl}(n)$ is a function/value which is negligible on the order of n .

In detail, what we are seeking is indistinguishability of multiple encryptions. That is to say, if we have the plain-text of two different messages (denote them m_1 and m_2), which we encrypt using a public key (denote it pk), then an adversary \mathcal{A} having access to the cipher-text of both messages **and** the public key should not be able to distinguish the cipher-text of the messages under any circumstances. Using pk , the encryption algorithm (denoted Enc_{pk}) generates cipher-text from messages m_1 and m_2 . We use

$$\text{Enc}_{pk}(m_1) \quad \textbf{and} \quad \text{Enc}_{pk}(m_2)$$

to denote the cipher-text generated for these messages, respectively.

We denote both the information $(pk, \text{Enc}_{pk}(m_1), \& \text{Enc}_{pk}(m_2))$ available/provided to the adversary \mathcal{A} by

$$\mathcal{A}(pk, \text{Enc}_{pk}(m_1), \text{Enc}_{pk}(m_2)) \quad (8.2)$$

furthermore, we also use this notation to represent the outcome of running PubK on \mathcal{A} . When \mathcal{A} succeeds, then the expression in 8.2 yields the result

$$\mathcal{A}(pk, \text{Enc}_{pk}(m_1), \text{Enc}_{pk}(m_2)) = 1 \quad (8.3)$$

The expression in 8.2 yields

$$\mathcal{A}(pk, \text{Enc}_{pk}(m_1), \text{Enc}_{pk}(m_2)) = 0 \quad (8.4)$$

otherwise.

Since **CPA** security requires security over multiple encryptions using the same public key, we will formally define this security using **two** pairs of messages that are all being encrypted using the same public key. We denote the first pair of messages by $m_{1,0}$ and $m_{2,0}$. Similarly, the second pair of messages are denoted by $m_{1,1}$ and $m_{2,1}$. We now use the same notation as in 8.2 with these message pairs (*and their associated public key pk*) to represent the attack by \mathcal{A} . This gives

$$\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})), \quad (8.2 \text{ a})$$

for the first message pair; and

$$\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})), \quad (8.2 \text{ b})$$

for the second message pair.

Before proceeding, we point out that we can equivalently use the expression from 8.3 in place of the $\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n) = 1$ term from 8.1. More clearly, we may formally write this equivalence as

$$\text{PubK}_{\mathcal{A}, \Pi}^{\text{LR-cpa}}(n) = 1 \quad \longleftrightarrow \quad \mathcal{A}(pk, \text{Enc}_{pk}(m_1), \text{Enc}_{pk}(m_2)) = 1$$

This allows us to write a version of 8.1 for both and . For the first message pair (*represented in*), this gives the result

$$\Pr[\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] \leq \frac{1}{2} + \text{negl}_0(n), \quad (8.3)$$

where negl_0 represents the negligible function required to satisfy this expression as applied to this message pair (*we are making allowances in case the results in and use different negl functions*). Writing our expression for the second message pair In a similar fashion yields

$$\Pr[\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \leq \frac{1}{2} + \text{negl}_1(n) \quad (8.4)$$

where negl_1 represents the negligible function required to satisfy this expression as applied to this message pair just as before (*we will see later that any difference between these negl functions is inconsequential; however differentiating between the negl functions used in either case is required for mathematical rigor*).

We now subtract the equation in 8.4 from the equation in 8.3 to obtain

$$\begin{aligned}
& \left\{ \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] - \right. \\
& \quad \left. - \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \right\} \leq \\
& \leq \left(\frac{1}{2} + \text{negl}_0(n) \right) - \left(\frac{1}{2} + \text{negl}_1(n) \right) \\
& \leq \text{negl}_0(n) - \text{negl}_1(n)
\end{aligned}$$

$$\begin{aligned}
& \left\{ \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] - \right. \\
& \quad \left. - \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \right\} \leq \left(\frac{1}{2} + \text{negl}_0(n) \right) - \left(\frac{1}{2} + \text{negl}_1(n) \right)
\end{aligned}$$

$$\begin{aligned}
& \left\{ \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] - \right. \\
& \quad \left. - \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \right\} \leq \text{negl}_0(n) - \text{negl}_1(n)
\end{aligned}$$

$$\begin{aligned}
& \left\{ \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] - \right. \\
& \quad \left. - \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \right\} \leq \left(\frac{1}{2} + \text{negl}_0(n) \right) - \left(\frac{1}{2} + \text{negl}_1(n) \right) \\
& \left\{ \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,0}), \text{Enc}_{pk}(m_{2,0})) = 1] - \right. \\
& \quad \left. - \Pr [\mathcal{A}(pk, \text{Enc}_{pk}(m_{1,1}), \text{Enc}_{pk}(m_{2,1})) = 1] \right\} \leq \text{negl}_0(n) - \text{negl}_1(n)
\end{aligned}$$

Adapting the security

Since we will eventually make use of the security definition in 8.1, we differentiate the expressions in
and

that \mathcal{A} be unable to break