EVALUATION REPORT / BUILD OF 24 OCT 2026

# BANTAY-BAHA

Flood-linked dengue surge risk and storm-driven loss of hospital access, ranked by province for any typhoon track. Results of the full pipeline run on 501 verified source files.

PJDSC 2026

02

# WHAT WE BUILT, WHAT THE DATA SAYS

Bantay-Baha joins three graphs into one ranking: a flood-exposure layer (rain, river routing, flood hazard), a lead-lag outbreak graph learned with PCMCI+, and a road-plus-ferry access graph that a typhoon cuts. A quantile forecast of dengue gives the probability **R** that a province's next four weeks exceed its usual bad month; the access model gives the share **A** of people cut off from a hospital; priority is **P = R × A**. Every number below was computed by `build.bat` from the raw files; nothing is synthetic.

**0.54**
relative WIS vs seasonal-naive, 1-4 weeks (target below 1)

**85%**
90% interval coverage (target 80-95%)

**8**
PCMCI+ lead-lag links among 84 provinces (cross-correlation: 4,265)

**60**
pre-computed scenarios (4 typhoons x 15 tracks)

## FINDINGS

- **The forecast beats the bar.** Pooled multi-quantile XGBoost reaches relative WIS 0.42 at one week and 0.66 at four weeks over 34,050 rolling-origin forecasts in 2018-2019; 90% intervals cover 85%.
- **Two honest null results.** Lead-lag graph features do not improve the forecast (relative WIS graph / no-graph = 0.999), and adding rain, river, storm and flood features moves relative WIS from 0.542 to 0.544 (ratio 1.003). At 1-4 weeks, recent cases and the season carry the skill.
- **Surge probability is informative but under-confident.** Brier 0.089 vs 0.216 for the base rate; forecasts of 0.1-0.7 come true 10-20 points more often. It needs recalibration before the tier thresholds are trusted.
- **The ranking points at the right places.** Spearman(R, observed next-4-week incidence) = 0.69 for Ompong and 0.74 for Tisoy; 7 of Ompong's top-10 provinces by priority surged against a 40% base rate.
- **Access loss is large and robust to settings.** The typhoons' gale swaths cancel 42-100 of 108 RoRo routes and cut off 0.3 M-1.0 M people (roads cut by Sentinel-1 observed flood); Kendall's τ of the ranking under nine setting changes is at least 0.99.

> **Not yet in these results:** the MAP friction raster access method (data on disk, not wired into the scenarios); the DOH NHFR hospital list and PSA PSGC (manual downloads); the Chronos-2 / TabPFN-TS benchmarks. See the scorecard (page 13).





*Figure 1. The forecast target: DOH-Epi weekly dengue summed over the 85 harmonised units, with the four backtest typhoons. 2020 is used for access scenarios only.*

SUMMARY

03

# DATA AND ACQUISITION

501 files from 22 sources (7.0 GB) are listed in `data/raw/MANIFEST.csv` with size and sha256; all were verified on disk before this build. Open sources come from `scripts/download_data.py`, Earth Engine sources from `scripts/download_gee.py`, and the team shares them through a private bucket (`setup.bat`, every file checked against the manifest).

| FOLDER | DATASET | ROLE | FROM | FILES | MB |
|---|---|---|---|---|---|
| dengue_hdx | DOH-Epi dengue, province/city x week 2016–Jan 2021 | forecast target | HDX | 1 | 2 |
| census2020 | 2020 CPH population by municipality | population | HDX | 1 | 1 |
| cod_ab | COD-AB boundaries ADM0-4 (v03, 2025) | join keys, geometry | HDX | 2 | 370 |
| ibtracs | IBTrACS v04r01 Western Pacific | storm exposure, swaths | NOAA NCEI | 1 | 115 |
| chirps | CHIRPS v2.0 daily 0.05°, 2016–2020 | rain features, routing | IRI Data Library | 5 | 563 |
| imerg | GPM IMERG V07 daily 0.1° | rain features | Earth Engine | 60 | 68 |
| hydrorivers | HydroRIVERS v1.0 Australasia | river routing | HydroSHEDS | 2 | 139 |
| noah | UP NOAH flood hazard 5/25-yr, 139 provinces | static susceptibility, road proxy | UP NOAH (Drive) | 142 | 4,733 |
| copernicus_ems | EMSR312 / EMSR478 vector packages | observed flood (Ompong AOIs) | Copernicus EMS | 16 | 28 |
| worldpop | WorldPop 2020 100 m UN-adjusted | population grid | WorldPop | 1 | 159 |
| meta_hrsl | Meta HRSL under-5 / 60+ | vulnerability (not yet used) | HDX | 2 | 112 |
| osm | OpenStreetMap Philippines PBF | road network | Geofabrik | 1 | 611 |
| roroph | roroph RoRo routes (108) | ferry edges (2024–26 proxy) | pinasr/roroph | 2 | <1 |
| hotosm_health | HOT OSM health facilities | hospital nodes (fallback) | HDX | 1 | 2 |
| noah_facilities | UP NOAH critical facilities | hospital nodes (fallback) | UP NOAH | 1 | 4 |
| map_friction | MAP friction surface 2019 | raster access (wired next) | Earth Engine | 2 | 2 |
| gfd | Global Flood Database events | context (ends 2018) | Earth Engine | 1 | <1 |
| sentinel1 | Sentinel-1 scene inventory | flood-mapping plan | Earth Engine | 1 | <1 |
| cchain | Project CCHAIN PIDSR city-week | leptospirosis (stretch) | HDX | 4 | 7 |
| mendeley_parreno | Parreño national weekly notifiable diseases | cross-check | Mendeley Data | 1 | <1 |
| opendengue | OpenDengue V1.3 | national cross-check | OpenDengue | 3 | 110 |
| *s1_flood* | *Sentinel-1 flood masks per typhoon (UN-SPIDER)* | *observed flood* | *Earth Engine* | – | *pending* |
| *psgc* | *PSA PSGC 2Q 2026* | *pcode check* | *manual* | – | *pending* |
| *nhfr* | *DOH NHFR* | *hospital list* | *manual* | – | *pending* |

# STEP 0: ONE UNIT-WEEK TABLE

## HARMONISED UNITS

The DOH file has **126 reporting units, not 81 provinces**: provinces mixed with 45 highly urbanised and component cities, plus misspellings (NUEVA VISCAYA, SISQUIJOR, BASILIG CITY = Bislig). COD-AB uses current boundaries (Maguindanao split in 2022, Special Geographic Area in 2024). The reviewed crosswalk in `data/crosswalks/` folds cities into their province and undoes the post-2020 changes, giving **85 units**; all 126 dengue units resolve and every unit has data. A unit-week counts only if every member reported (NaN otherwise, never 0).

## POPULATION

The 2020 census table's municipal codes are old 9-digit PSGC and do not join to COD-AB for 54 municipalities (24.2 M people). Its own *Province* column already places cities in their province, so units are summed on it: 108.67 M people.





*Figure 2. Coverage of the 85 x 262 panel. Sentinel-1 is filled only for the four landfall windows (4 weeks each), by design.*

## LAYER 1 INPUTS

Rain: CHIRPS and IMERG agree (weekly r = 0.87). River routing accumulates CHIRPS rain down the HydroRIVERS network. NOAH 25-yr hazard covers 74 of 85 units; the median unit has 20% of its people in a hazard zone. Units with no map are **missing, not zero**: Basilan, Batanes, Biliran, Camiguin, Davao Occidental, Guimaras, Northern Samar, Romblon, Siquijor, Sulu, Tawi-Tawi.

Storm exposure: every Western Pacific track 2016-2020 from IBTrACS, gale radius from USA R34, else JMA R30, else a fixed 150 km (the fallback never triggered for the four backtest storms).

05

# METHODS AND EVALUATION PROTOCOL

## LAYER 2: LEAD-LAG OUTBREAK GRAPH

log(1 + incidence), 52-week seasonal difference, standardised inside the training window. PCMCI+ (Runge, 2020) with partial-correlation tests, lags 1-3 weeks, pc_alpha 0.01; lagged inter-province edges are kept when the Benjamini-Hochberg (1995) q-value is below 0.05. The FDR step matters: on pure noise with 85 series, PCMCI+ alone returned 60-90 false edges. PageRank (Page et al., 1999) on the reversed graph scores sentinels; Louvain (Blondel et al., 2008) finds communities; lagged cross-correlation with the same FDR is the baseline graph.

## FORECAST

One pooled XGBoost (Chen & Guestrin, 2016) per horizon (1-4 weeks) with the multi-quantile objective at the 23 CDC FluSight levels, on log(1 + incidence), back-transformed to cases. Features: lagged incidence and moving averages, same week last year, week of year, CHIRPS / IMERG / river rain lags, storm exposure, NOAH hazard, and leader signals from the graph. SHAP values (Lundberg & Lee, 2017) from the median model give the dashboard's "drivers". A second model predicts the 4-week cumulative incidence directly; **R** is its probability of exceeding the unit's 90th percentile of 4-week incidence in the training window.

## EVALUATION

**Rolling origin, never a random split.** From January 2018 to December 2019 the model, the lead-lag graph and the surge thresholds are refit every 13 weeks on data up to the refit date only, then forecast every origin until the next refit. Training targets never pass the refit date and 2020 is never a target. Scores: weighted interval score on case counts (Bracher et al., 2021), relative WIS against a seasonal-naive forecast with empirical error quantiles, 50% and 90% coverage, and an online conformal 90% band built from past errors only (Gibbs & Candès, 2021) as a check on the quantiles.

## LAYER 3: ACCESS

OSM highways (motorway to track) become a graph of junctions; RoRo routes are ferry edges (crossing time plus half the headway); hospitals are HOT OSM plus NOAH name-matched facilities (2,199) until NHFR arrives. WorldPop (Tatem, 2017) is summed to 1 km cells, each joined to the nearest road node. A multi-source Dijkstra gives hours to the nearest hospital. A storm cancels ferry routes crossing its gale swath and cuts roads in observed flood (Sentinel-1, observed tracks) or, for shifted tracks, cuts (high hazard) or slows (medium) roads in NOAH 25-yr hazard inside the swath. A cell is cut off if it loses its route, or its trip at least doubles and grows by more than an hour (after Petricola et al., 2022).

06

# FORECAST SKILL

| HORIZON | N | WIS XGB | WIS NO-GRAPH | WIS NAIVE | REL WIS | GRAPH / NO-GRAPH | COV 50 | COV 90 | COV 90 CONFORMAL |
|---|---|---|---|---|---|---|---|---|---|
| 1 wk | 8,640 | 14.7 | 14.7 | 35.1 | **0.42** | 0.998 | 0.47 | 0.89 | 0.88 |
| 2 wk | 8,555 | 18.1 | 18.0 | 35.4 | **0.51** | 1.002 | 0.44 | 0.86 | 0.87 |
| 3 wk | 8,470 | 21.0 | 21.1 | 35.6 | **0.59** | 0.998 | 0.42 | 0.84 | 0.86 |
| 4 wk | 8,385 | 23.6 | 23.6 | 35.9 | **0.66** | 0.999 | 0.40 | 0.82 | 0.85 |
| **all** | **34,050** | **19.3** | **19.3** | **35.5** | **0.54** | **0.999** | **0.43** | **0.85** | **0.87** |





*Figure 3. Left: mean WIS by horizon (labels: relative WIS vs seasonal-naive). Right: coverage against the nominal 50% and 90% lines and the proposal's 80-95% band.*

**Targets met.** Relative WIS is below 1 at every horizon (0.42 to 0.66) and 90% coverage is inside 80-95% at every horizon (four weeks is close to the floor at 0.82). The 50% intervals are a little narrow (0.40-0.47). The online conformal band reaches similar coverage, a sign the quantiles are not badly miscalibrated.





*Figure 4. Rolling-origin forecasts for Cagayan (Ompong's landfall province, dashed line), one and four weeks ahead.*

07

# ABLATIONS AND CALIBRATION

## WHAT CARRIES THE SKILL

| COMPARISON | RELATIVE WIS | READING |
|---|---|---|
| XGBoost vs seasonal-naive | 0.544 | 46% lower WIS than "same week last year" |
| with vs without graph leader features | 0.999 | no gain: the graph has 0 links in every 2018 window and 2-6 in 2019 |
| with vs without rain / river / storm / flood features | 1.003 | 0.544 vs 0.542 against seasonal-naive, same rows |

The graph ablation follows from the graph itself: with so few significant links, leader signals are empty for most provinces. The rain result has two likely reasons, not yet tested: rain moves with the season in a four-year panel, so the season features already carry it, and any effect of rain at lags longer than four weeks is outside this forecast horizon. Both are reported as findings rather than tuned away.





*Figure 5. Reliability of R over the 2018-2019 unit-weeks. Points above the diagonal: R is under-confident.*

## SURGE PROBABILITY R

Brier score 0.089 against 0.216 for always predicting the base rate. The base rate is 32%, not 10%, because 2019 was an epidemic year and the thresholds come from earlier, quieter years.

R is under-confident in the middle range: forecasts of 0.1-0.7 come true 10-20 points more often. Isotonic recalibration fitted on past rolling-origin forecasts only is the next step, before the tier thresholds (R ≥ 0.3, R ≥ 0.5) are relied on.

# THE LEAD–LAG GRAPH




Final window: 84 series, T = 156 weeks after seasonal differencing (Davao Occidental sits out with more than 10% missing weeks). PCMCI+ keeps **8** lagged inter-province links at q < 0.05. Plain lagged cross-correlation with the same FDR keeps **4,265**, because the 2019 national epidemic makes almost every pair co-move.

The proposal's sentinel idea is therefore **weakly supported at province-week level**. The links are statistical dependencies, not evidence of transmission, and several join distant provinces. Likely reasons: weekly province totals average out local waves, four years is short for 84 series, and reporting artefacts dominate small provinces. Synchrony studies such as van Panhuis et al. (2015) worked at longer time scales.

*Figure 6. The 8 PCMCI+ links learned on 2016–2019 (teal = leader).*

| LEADER | FOLLOWER | LAG (WK) | PARTIAL CORR | Q | BOOTSTRAP FREQ |
|---|---|---|---|---|---|
| Cavite | Sorsogon | 1 | 0.38 | 0.001 | 0.70 |
| Metropolitan Manila Second District | Bulacan | 3 | 0.36 | 0.006 | 0.70 |
| Quirino | Surigao del Sur | 1 | 0.36 | 0.007 | 0.65 |
| Ilocos Norte | Romblon | 1 | 0.33 | 0.020 | 0.75 |
| Metropolitan Manila Third District | Zamboanga del Norte | 1 | 0.33 | 0.029 | 0.85 |
| Sulu | Ilocos Sur | 1 | 0.32 | 0.037 | 0.55 |
| Camarines Sur | Pangasinan | 1 | 0.31 | 0.042 | 0.65 |
| Samar (Western Samar) | Antique | 1 | 0.31 | 0.045 | 0.55 |

09

# OBSERVED FLOOD: SENTINEL-1

Radar sees through typhoon clouds. For each typhoon, new open water is mapped on Google Earth Engine (Gorelick et al., 2017) by comparing Sentinel-1 VH backscatter before and after landfall, in 100 m tiles near the track: before = median of days -30 to -3, after = darkest image of days 0 to +10, **both from the same relative orbit**, so the viewing geometry matches. A pixel is new flood when VH drops by at least 3 dB and ends water-dark (at most -18 dB); permanent water (JRC), slopes over 5° and specks under 8 pixels are removed (after UN-SPIDER, n.d.).





*Figure 7. New-flood share of the land Sentinel-1 saw, per unit, in the landfall week, with each track.*

| TYPHOON | PEOPLE IN NEW FLOOD | UNITS >1% FLOODED | MAX UNIT SHARE | MOST PEOPLE FLOODED |
|---|---|---|---|---|
| Ompong | 0.5 M | 5 | 8.0% | Pampanga |
| Tisoy | 1.1 M | 14 | 9.6% | Isabela |
| Rolly | 0.6 M | 10 | 4.7% | Camarines Sur |
| Ulysses | 1.4 M | 14 | 10.2% | Cagayan |





*Figure 8. Ompong around Tuguegarao: Sentinel-1 new flood (teal) and Copernicus EMS observed flood (red outlines).*

## CHECKED AGAINST COPERNICUS EMS

Inside the EMS areas of interest, EMS mapped 108 km$^2$ of observed flood and Sentinel-1 flagged 46 km$^2$; 26 km$^2$ overlap. So **55%** of Sentinel-1 flood there is confirmed by EMS, and Sentinel-1 recovers **24%** of the EMS area.

The first version followed UN-SPIDER's after/before ratio on dB values and recovered almost none of the EMS flood: dividing negative dB values under-detects surfaces that were already dark, and mixing orbits compared different viewing geometries. Sampling backscatter at EMS flood points showed drops of 4-10 dB that the ratio missed. Low recall is expected: EMS used other imagery and dates, while one Sentinel-1 orbit passed here only on 16 and 22 September.

# ACCESS LOSS UNDER FOUR TYPHOONS

**Baseline.** 109.4 M people in 1 km cells; 97.3% have a road or ferry route to one of 2,199 hospitals, with a population-weighted mean trip of 0.24 h. Results are also written for 1,641 municipalities.

| TYPHOON | FERRY ROUTES CUT | PEOPLE CUT OFF | MEAN A | MEAN R | DEPLOY / PRE-POSITION | TOP PRIORITY |
|---|---|---|---|---|---|---|
| Ompong Mangkhut 2018 | 42 | 0.3 M | 0.2% | 0.22 | 0 / 21 | Pampanga |
| Tisoy Kammuri 2019 | 100 | 0.7 M | 0.6% | 0.08 | 0 / 5 | Camarines Norte |
| Rolly Goni 2020 | 42 | 0.4 M | 0.3% | 0.02 | 0 / 0 | Oriental Mindoro |
| Ulysses Vamco 2020 | 54 | 1.0 M | 0.8% | 0.04 | 0 / 2 | Cagayan |





*Figure 9. Access loss A per unit for the observed tracks. Roads are cut by Sentinel-1 observed flood; ferries crossing the gale swath are cancelled.*

> Roads are cut only where Sentinel-1 saw new water, so cut-offs are storm-specific; the radar misses floods under dense vegetation and in built-up areas and between passes, so these are lower bounds. With the NOAH 25-yr hazard proxy instead (build of 24 Oct, before Sentinel-1), the same tracks cut off 4.8-10.3 M people: an upper bound. The truth lies between. Ferry routes are 2024-2026 operations used as a proxy for 2018-2020. Shifted tracks always use the NOAH hazard proxy (no image of a storm that did not happen).

# PRIORITY: OMPONG 2018





*Figure 10. P = R x A for Ompong with its track and gale swath; top 5 labelled.*





*Figure 11. Risk vs access. Dotted lines are the provisional tier thresholds (R = 0.3, A = 10%); marker shape encodes the action tier.*

Ompong made landfall in Cagayan on 15 September 2018. The model, trained on data to the landfall week only, puts **Pampanga** first, with 0 provinces in the "deploy mobile clinics" tier and 21 in "pre-position supplies". Provinces with high R but little observed loss of access fall to "pre-position"; "deploy" needs at least 10% of people cut off, which observed flood alone did not produce for this storm (the hazard-map upper bound did, for 7 provinces).

The dashboard repeats this for every storm and 14 shifted variants of each track (±0.5° and ±1° north, ±1° east), so a health officer can ask "what if it comes in further south?" and get an answer in seconds.

# BACKTEST, SENSITIVITY, BENCHMARK

## DID HIGH-PRIORITY PROVINCES SEE MORE CASES?

| TYPHOON | ORIGIN | UNITS | SURGED | P(R, OBS) | P(P, OBS) | TOP-10 SURGED | BRIER R | NOTE |
|---|---|---|---|---|---|---|---|---|
| Ompong | 2018-09-16 | 85 | 40% | 0.69 | 0.21 | 70% | 0.177 | |
| Tisoy | 2019-12-08 | 85 | 5% | 0.74 | -0.05 | 0% | 0.041 | |
| Rolly | 2020-11-01 | 61 | 0% | 0.48 | 0.05 | 0% | 0.001 | 2020: COVID-depressed reporting, indicative only |
| Ulysses | 2020-11-15 | 85 | 1% | 0.35 | 0.12 | 0% | 0.016 | 2020: COVID-depressed reporting, indicative only |





*Figure 12. Spearman correlation of each ranking with observed incidence over the next four weeks (relative to each unit's threshold).*

R ranks provinces well for both 2018-2019 storms. P correlates less with cases, as it should: A measures access, not disease. After Ompong, 7 of the 10 highest-priority provinces surged against a 40% base rate. After Tisoy almost nothing surged (the 2019 season was over), so the top-10 rate is 0 and the Brier score is low for the right reason.

## SENSITIVITY: KENDALL'S $\tau$ OF P AGAINST THE BASE SETTINGS

| VARIANT | OMPONG τ | TISOY τ | ROLLY τ | ULYSSES τ | MEAN A (OMPONG) |
|---|---|---|---|---|---|
| cut-off ratio 1.5 | 1.000 | 0.997 | 1.000 | 1.000 | 0.23% |
| cut-off ratio 3 | 1.000 | 0.997 | 0.999 | 1.000 | 0.23% |
| ferry rule: port in swath | 1.000 | 1.000 | 1.000 | 1.000 | 0.23% |
| gale fallback 100 km | 1.000 | 1.000 | 1.000 | 1.000 | 0.23% |
| gale fallback 200 km | 1.000 | 1.000 | 1.000 | 1.000 | 0.23% |
| medium hazard slow x0.25 | 1.000 | 1.000 | 1.000 | 1.000 | 0.23% |
| medium hazard slow x0.75 | 1.000 | 1.000 | 1.000 | 1.000 | 0.23% |
| speed x0.8 | 1.000 | 0.998 | 0.987 | 0.997 | 0.23% |
| speed x1.2 | 1.000 | 0.996 | 0.986 | 0.998 | 0.23% |

Every variant changes A (speed ±20% moves mean A by about 4%) but none reorders the ranking: P's spread is dominated by R, which ranges 0.02-0.93 across provinces. The gale fallback never triggers for these storms.

## METRO MANILA BENCHMARK

Cutting every NCR road in NOAH 25-yr medium or high hazard leaves **29.7%** of NCR without a route within the cut-off rule. Tariverdi et al. (2023) report about 22% losing access to higher-level care at a 15 cm flood. The definitions differ, so this is an order-of-magnitude check, not a replication.

# PROPOSAL OBJECTIVES: SCORECARD

| OBJECTIVE (PROPOSAL) | RESULT | STATUS | NOTE |
|---|---|---|---|
| Province-week panel, 2016-2020, joined on PSGC | 85 harmonised units x 262 weeks; dengue, rain, river, storms, NOAH, Sentinel-1 | met | |
| Lead-lag graph, sentinels, communities; test sentinel features out of sample | PCMCI+ with FDR: 8 links; ablation run | met | result is null (graph / no-graph 0.999) |
| Relative WIS below 1.0 vs seasonal-naive, 2018-2019 | 0.544 overall, 0.66 at 4 weeks | met | |
| 90% interval coverage of 80-95% | 0.852 overall; 0.82-0.89 by horizon | met | |
| Access loss per municipality, two methods, four typhoons | network method, 1,641 municipalities, 4 storms | partly | raster (friction) method not wired in yet |
| Sensitivity bounds; Metro Manila benchmark | 9 variants, $\tau \geq 0.99$; NCR 30% vs ~22% | met | |
| Dashboard for any historical or shifted track in under a minute | 60 pre-computed scenarios; loads in about 6 s | met | |
| System Usability Scale from three target users | not done | not yet | needs the user sessions |
| Foundation-model benchmarks (Chronos-2, TabPFN-TS) | Chronos-2 script written, not run | not yet | needs the torch install |

## PROPOSAL CORRECTIONS FOUND IN THE DATA

- 126 dengue reporting units, not 81 provinces; deaths unreliable (2016 case fatality 3.9%); 2020 collapsed under COVID-19.
- An open subnational leptospirosis series exists (Project CCHAIN, 12 cities, 2008-2022).
- EMSR478 (Rolly) has damage grading only, no flood delineation; the Global Flood Database (Tellman et al., 2021) ends in 2018.
- Flood hazard maps come from the UP NOAH archive, not LiPAD; 11 units have no 25-yr map.
- HydroRIVERS (Lehner & Grill, 2013) places the Philippines in the Australasia file, not Asia.
- IBTrACS (Knapp et al., 2010) USA R34 is missing on up to 58% of fixes of the backtest storms; JMA R30 fills the gaps.
- roroph routes describe 2024-2026 operations with partly coarse port coordinates.

14

# LIMITATIONS AND NEXT STEPS

## THREATS TO VALIDITY

- **Two evaluation years.** 2018-2019 contains one epidemic; skill and calibration may differ in quieter or COVID-era years.
- **Surveillance artefacts.** Reporting delays and under-reporting are in the target; forecasts predict reported, not true, cases.
- **Observed flood.** Sentinel-1 sees each place every 6-12 days and misses flooding under dense vegetation and in built-up areas; shifted tracks fall back to static hazard maps.
- **Network gaps.** OSM completeness is uneven outside Luzon; ferries are 2024-2026 routes; hospitals are an OSM / NOAH fallback without facility level or capacity.
- **Provisional rules.** Action-tier thresholds, road speeds and the cut-off rule are assumptions, partly addressed by the sensitivity analysis.
- **Causality.** PCMCI+ links are conditional dependencies, not transmission routes.

## NEXT STEPS

1. Add a second radar or optical source between Sentinel-1 passes and compare observed-flood A with the hazard proxy.
2. Wire in the MAP friction raster method (Weiss et al., 2020) and report the network-vs-raster gap per province.
3. Recalibrate R and review the tiers with DOH and LGU users; run the System Usability Scale test.
4. Replace the hospital fallback with NHFR; validate pcodes against PSGC 2Q 2026.
5. Run the Chronos-2 benchmark (Ansari et al., 2025); extend to leptospirosis with the CCHAIN city series.

## REPRODUCIBILITY

`setup.bat` (or `download_data.bat`) → `build.bat` → `run_app.bat`. This report and the slides are regenerated with `python scripts/make_results.py`. Missing inputs are skipped, listed in `data/processed/status.json` and on the dashboard; re-running after they arrive needs no code change.

# REFERENCES


Ansari, A. F., Shchur, O., Küken, J., Auer, A., Han, B., Mercado, P., Rangapuram, S. S., Shen, H., Stella, L., Shen, X., Turkmen, C., Bohlke-Schneider, M., Erickson, N., Kapoor, S., Li, Z., Wang, B., Williams, A. R., Ahmed, A., Weber, S., … Wang, Y. (2025). *Chronos-2: From univariate to universal forecasting*. arXiv. https://arxiv.org/abs/2510.15821

Balbuena, A., Herbosa, T., & Gundran, C. P. (2010). Cases of leptospirosis after Typhoon Ondoy (international name: Ketsana) seen at the emergency department: The Philippine General Hospital experience. *WIT Transactions on Information and Communication Technologies, 43*, 605–612. https://doi.org/10.2495/RISK100501

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological), 57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Blondel, V. D., Guillaume, J.-L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment, 2008*(10), P10008. https://doi.org/10.1088/1742-5468/2008/10/P10008

Bracher, J., Ray, E. L., Gneiting, T., & Reich, N. G. (2021). Evaluating epidemic forecasts in an interval format. *PLoS Computational Biology, 17*(2), e1008618. https://doi.org/10.1371/journal.pcbi.1008618

Buczak, A. L., Baugher, B., Babin, S. M., Ramac-Thomas, L. C., Guven, E., Elbert, Y., Koshute, P. T., Velasco, J. M. S., Roque, V. G., Jr., Tayag, E. A., Yoon, I.-K., & Lewis, S. H. (2014). Prediction of high incidence of dengue in the Philippines. *PLoS Neglected Tropical Diseases, 8*(4), e2771. https://doi.org/10.1371/journal.pntd.0002771

Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. In *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining* (pp. 785–794). https://doi.org/10.1145/2939672.2939785

Copernicus Emergency Management Service. (2018). *EMSR312: Typhoon Mangkhut in the Philippines* [Rapid mapping activation]. European Commission. https://mapping.emergency.copernicus.eu/activations/EMSR312/

Department of Health Epidemiology Bureau. (2021). *DOH-Epi dengue data 2016–2021* [Data set]. Humanitarian Data Exchange. https://data.humdata.org/dataset/ac63a95e-7296-42fb-802b-7f7541c73e45

Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., & Michaelsen, J. (2015). The climate hazards infrared precipitation with stations—A new environmental record for monitoring extremes. *Scientific Data, 2*, 150066. https://doi.org/10.1038/sdata.2015.66

Gibbs, I., & Candès, E. (2021). Adaptive conformal inference under distribution shift. *Advances in Neural Information Processing Systems, 34*, 1660–1672.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment, 202*, 18–27. https://doi.org/10.1016/j.rse.2017.06.031

Huffman, G. J., Stocker, E. F., Bolvin, D. T., Nelkin, E. J., & Tan, J. (2023). *GPM IMERG final precipitation L3 half hourly 0.1 degree x 0.1 degree V07* [Data set]. NASA Goddard Earth Sciences Data and Information Services Center. https://doi.org/10.5067/GPM/IMERG/3B-HH/07

Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., & Neumann, C. J. (2010). The International Best Track Archive for Climate Stewardship (IBTrACS): Unifying tropical cyclone data. *Bulletin of the American Meteorological Society, 91*(3), 363–376. https://doi.org/10.1175/2009BAMS2755.1

Lehner, B., & Grill, G. (2013). Global river hydrography and network routing: Baseline data and new approaches to study the world's large river systems. *Hydrological Processes, 27*(15), 2171–2186. https://doi.org/10.1002/hyp.9740

# REFERENCES (CONTINUED)


Lundberg, S. M., & Lee, S.-I. (2017). A unified approach to interpreting model predictions. *Advances in Neural Information Processing Systems, 30*, 4765–4774.

Matsushita, N., Ng, C. F. S., Kim, Y., Suzuki, M., Saito, N., Ariyoshi, K., Salva, E. P., Dimaano, E. M., Villarama, J. B., Go, W. S., & Hashizume, M. (2018). The non-linear and lagged short-term relationship between rainfall and leptospirosis and the intermediate role of floods in the Philippines. *PLoS Neglected Tropical Diseases, 12*(4), e0006331. https://doi.org/10.1371/journal.pntd.0006331

OpenStreetMap contributors. (2026). *OpenStreetMap data extract: Philippines* [Data set]. Geofabrik. https://download.geofabrik.de/asia/philippines.html

Page, L., Brin, S., Motwani, R., & Winograd, T. (1999). *The PageRank citation ranking: Bringing order to the web* (Technical Report 1999-66). Stanford InfoLab.

Petricola, S., Reinmuth, M., Lautenbach, S., Hatfield, C., & Zipf, A. (2022). Assessing road criticality and loss of healthcare accessibility during floods: The case of Cyclone Idai, Mozambique 2019. *International Journal of Health Geographics, 21*, 14. https://doi.org/10.1186/s12942-022-00315-2

Philippine Statistics Authority. (2021). *Philippines: 2020 census of population and housing, population by municipality* [Data set]. Humanitarian Data Exchange. https://data.humdata.org/dataset/c1edf0c5-58c8-4d39-b566-937b1ad7907f

Philippine Statistics Authority & National Mapping and Resource Information Authority. (2025). *Philippines: Subnational administrative boundaries (COD-AB)* [Data set]. Humanitarian Data Exchange. https://data.humdata.org/dataset/caf116df-f984-4deb-85ca-41b349d3f313

Runge, J. (2020). Discovering contemporaneous and lagged causal relations in autocorrelated nonlinear time series datasets. In *Proceedings of the 36th Conference on Uncertainty in Artificial Intelligence* (PMLR 124, pp. 1388–1397). https://arxiv.org/abs/2003.03685

Runge, J., Nowack, P., Kretschmer, M., Flaxman, S., & Sejdinovic, D. (2019). Detecting and quantifying causal associations in large nonlinear time series datasets. *Science Advances, 5*(11), eaau4996. https://doi.org/10.1126/sciadv.aau4996

Tariverdi, M., Nunez-del-Prado, M., Leonova, N., & Rentschler, J. (2023). Measuring accessibility to public services and infrastructure criticality for disasters risk management. *Scientific Reports, 13*, 1569. https://doi.org/10.1038/s41598-023-28460-z

Tatem, A. J. (2017). WorldPop, open data for spatial demography. *Scientific Data, 4*, 170004. https://doi.org/10.1038/sdata.2017.4

Tellman, B., Sullivan, J. A., Kuhn, C., Kettner, A. J., Doyle, C. S., Brakenridge, G. R., Erickson, T. A., & Slayback, D. A. (2021). Satellite imaging reveals increased proportion of population exposed to floods. *Nature, 596*, 80–86. https://doi.org/10.1038/s41586-021-03695-w

UN-SPIDER. (n.d.). *Recommended practice: Flood mapping and damage assessment using Sentinel-1 SAR data in Google Earth Engine*. United Nations Office for Outer Space Affairs. https://www.un-spider.org/advisory-support/recommended-practices/recommended-practice-google-earth-engine-flood-mapping

van Panhuis, W. G., Choisy, M., Xiong, X., Chok, N. S., Akarasewi, P., Iamsirithaworn, S., Lam, S. K., Chong, C. K., Lam, F. C., Phommasak, B., Vongphrachanh, P., Bouaphanh, K., Rekol, H., Hien, N. T., Thai, P. Q., Duong, T. N., Chuang, J.-H., Liu, Y.-L., Ng, L.-C., … Burke, D. S. (2015). Region-wide synchrony and traveling waves of dengue across eight countries in Southeast Asia. *Proceedings of the National Academy of Sciences, 112*(42), 13069–13074. https://doi.org/10.1073/pnas.1501375112

Weiss, D. J., Nelson, A., Vargas-Ruiz, C. A., Gligorić, K., Bavadekar, S., Gabrilovich, E., Bertozzi-Villa, A., Rozier, J., Gibson, H. S., Shekel, T., Kamath, C., Lieber, A., Schulman, K., Shao, Y., Qarkaxhija, V., Nandi, A. K., Keddie, S. H., Rumisha, S., Amratia, P., … Gething, P. W. (2020). Global maps of travel time to healthcare facilities. *Nature Medicine, 26*, 1835–1838. https://doi.org/10.1038/s41591-020-0931-3