


Engineering tunable anharmonic potentials with light-atom interaction



C. McGarry^{1,2}, T. Chalermputitarak^{1,2}, K. Schwennicke³,
F. Scuccimarra^{1,2}, C. Valahu^{1,2}, P. Nagpal^{1,2}, V. Matsos^{1,2},
M. Millican^{1,2}, I. Kassal^{3,2,4}, T. R. Tan^{1,2,4}


cameron.mcgarry@sydney.edu.au



1. School of Physics, University of Sydney, NSW 2006, Australia
2. ARC Centre of Excellence for Engineered Quantum Systems, University of Sydney, Australia
3. School of Chemistry, University of Sydney, NSW 2006, Australia
4. Sydney Nano Institute, University of Sydney, NSW 2006, Australia



We have developed and implemented a scheme for realising anharmonic motion in a trapped ion system. We use this to investigate quantum tunnelling in a double well.

Theory: Quantum Signal Processing for generation of anharmonic potentials




Repeated application of $G_c(\theta, \kappa)$ allows for the realisation of any potential that can be realised by Fourier decomposition. Realising the sine equivalent allows the generation of any arbitrary potential.


Experiment: simulation of cosine potential



Wigner function during evolution (simulation)



Simulated potential and wavefunction



Application: Chemical dynamics

Target ro-vibrational Hamiltonian and non-linear chemical dynamics simulations. See poster 30 (Frank Scuccimarra)

$$H_{VC} = \sum_j \omega_j a_j^\dagger a_j + \sum_{n,m} C_{n,m} |n\rangle \langle m| \quad C_{n,m} = e_0^{(n,m)} + \sum_j e_j^{(n,m)} x_j + \underbrace{\sum_{j,k} e_{j,k}^{(n,m)} x_j x_k}_{\text{Anharmonicity}}$$

Vibrational coupling Hamiltonian