

Stock Assessment Update: Status of Widow Rockfish Stock (*Sebastodes entomelas*) off the U.S. West Coast in 2025

Michael Kinneen¹, Maurice Goodman¹, Anna Sulc¹, Laurinne Balstad², Raquel Ruiz Diaz¹, Kristina Randrup¹, William Patrone¹, Laura Spencer¹, Alaia Morell³, Alberto Rovellini¹, Allison Dedrick⁴, Nick Grunloh⁵, Madison Bargas⁶, Stephanie Hopkins⁷, Vladlena Gertseva⁸, Kiva L. Oken⁸, Ian G. Taylor⁸, Melissa A. Haltuch⁹ and Owen Hamel⁸

1. School of Aquatic & Fishery Sciences (SAFS), University of Washington, 1122 NE Boat St, Seattle, WA, 98195
2. Department of Environmental Science and Policy, University of California Davis, Wickson Hall, 350 N Quad, Davis, CA, 95616
3. UW Puget Sound Institute, Center for Urban Waters, 326 East D Street, Tacoma, WA, 98421
4. California Department of Fish and Wildlife, 1740 North Market Boulevard, Sacramento, CA, 95834
5. Southwest Fisheries Science Center, 8901 La Jolla Shores Drive, La Jolla, CA, 92037
6. Coastal Oregon Marine Experiment Station, Oregon State University, 430 Strand Agriculture Hall, Corvallis, OR, 97331
7. University of California, Santa Cruz, Institute of Marine Sciences' Fisheries Collaborative Program, 1156 High Street, Santa Cruz, CA, 95064
8. NOAA Fisheries Northwest Fisheries Science Center, 2725 Montlake Boulevard East, Seattle, WA, 98112
9. NOAA Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Building 4, Seattle, WA, 98115

Table of contents

Disclaimer	i
Executive Summary	ii
Stock	ii
Catches	iii
Data and Assessment	v
Stock biomass and dynamics	vi
Recruitment	x
Exploitation status	xii
Ecosystem considerations	xv
Reference points	xvi
Management performance	xviii
Harvest projections and decision table	xviii
Scientific uncertainty	lxii
Research and data needs	lxii
1 Introduction	1
1.1 Distribution and Stock Structure	1
1.2 Life History	1
1.3 Ecosystem Considerations	1
1.4 Fishery description	2
1.5 Management History	2
1.6 Management performance	2
1.7 Fisheries off Canada and Alaska	2
2 Data	3
2.1 Fishery-independent data	3
2.1.1 Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey	3
2.1.2 Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey	3
2.1.3 SWFSC and NWFSC/PWCC Midwater Trawl Survey	5
2.2 Fishery-dependent data	6
2.2.1 Landings	6
2.2.2 Fishery catch-per-unit-effort	6
2.2.3 Fishery length and age data	7
2.2.4 Discards	7
2.2.5 Biological data	8
2.3 Environmental and Ecosystem Data	10
3 Assessment model	11
3.1 History of modeling approaches	11
3.2 Responses to most recent past STAR Panel recommendations	11

3.3	Model Structure and Assumptions	11
3.3.1	Model Changes from the Last Assessment	11
3.3.2	Modeling Platform and Structure	13
3.3.3	Model Overview	13
3.3.4	Model Parameters	14
3.3.5	Key Assumptions and Structural Choices	16
3.4	Base Model Results	16
3.4.1	Parameter Estimates	16
3.4.2	Fits to the Data	18
3.4.3	Population Trajectory	20
3.5	Model Diagnostics	21
3.5.1	Convergence	21
3.5.2	Parameter Uncertainty	22
3.5.3	Sensitivity Analyses	22
3.5.4	Retrospective Analysis	23
3.5.5	Likelihood Profiles and key parameters	24
4	Management	25
4.1	Reference Points	25
4.2	Unresolved problems and major uncertainties	25
4.3	Harvest Projections and Decision Tables	27
4.4	Evaluation of Scientific Uncertainty	28
4.5	Regional management considerations	28
4.6	Research and Data Needs	29
4.7	Acknowledgements	30
5	Tables	31
5.1	Data	31
5.1.1	Fishery-dependent data	31
5.1.2	Fishery-independent data	59
5.2	Model results	62
5.3	Management	96
6	Figures	100
6.1	Data	100
6.1.1	Indices	100
6.1.2	Composition data	109
6.1.3	Biological data	116
6.2	Model	120
6.2.1	Bridging	120
6.2.2	Selectivity	122
6.2.3	Biology	125
6.2.4	Fits to data	127
6.2.5	Timeseries	140

6.3	Model Diganostics	145
6.3.1	Sensitivity analyses	145
6.3.2	Retrospective analysis	148
6.3.3	Likelihood profiles	149
6.4	Management	153
7	References	155
8	Appendix A: Annual fits to length and age composition	158

Please cite this publication as:

M. Kinneen, M. Goodman, A. Sulc, L. Balstad, R. Diaz, K. Randrup, W. Patrone, L. Spencer, A. Morell, A. Rovellini, A. Dedrick, N. Grunloh, M. Bargas, S. Hopkins, V. Gersteva, K. Oken, I. Taylor, M. Haltuch, O. Hamel (2025) Stock Assessment Update: Status of Widow Rockfish (*Sebastodes entomelas*) Along the U.S. West Coast in 2025. Pacific Fishery Management Council, Portland, Oregon.

Disclaimer

These materials do not constitute a formal publication and are for information only. They are in a pre-review, pre-decisional state and should not be formally cited or reproduced. They are to be considered provisional and do not represent any determination or policy of NOAA or the Department of Commerce.

total_catch 1 10,669 total_catch 1 9,824

Executive Summary

Stock

This is an update assessment of widow rockfish (*Sebastodes entomelas*) that reside in the waters off California, Oregon, and Washington from the U.S.-Canada border in the north to the U.S.-Mexico border in the south. The most recent benchmark was conducted in 2015 (Hicks and Wetzel 2015) and first updated in 2019 (Adams et al. 2019). This assessment represents the second update assessment of this stock. Widow rockfish inhabit water depths of 25–370 m from northern Baja California, Mexico to Southeastern Alaska. Although catches north of the U.S.-Canada border and south of the U.S.-Mexico border were not included in this assessment, it is not certain if those populations contribute to the biomass of widow rockfish off of the U.S. West Coast, possibly through adult migration and/or larval dispersion. Following the 2015 benchmark assessment (Hicks and Wetzel 2015), this update assessment is based on a single costwide area model.

Catches

Historically, fisheries have caught widow rockfish since the turn of the 20th century. Landings in the trawl fishery are estimated to have increased into the 1940s and remained relatively constant and small (below 1,000 mt per year) throughout the 1950s and into the 1960s before the foreign trawl fleet increased catches into the 1970s, with a peak at almost 5,000 mt in 1967. In the late 1970s a midwater trawl fishery developed for widow rockfish and catches increased rapidly with the discovery of large aggregations that form at night.

Total landings of widow rockfish peaked in the early 1980s, increasing from approximately 1,000 metric tons (mt) in 1978 to over 25,000 mt in 1981. The large landings in the early 1980s were curtailed with trip limits beginning in 1982, which resulted in a decline in landings throughout the 1980s and 1990s following sequential reductions in the trip limits. From 2000 to 2003, landings of widow rockfish dropped from over 4,000 mt to about 40 mt and remained low through 2016. Catches increased rapidly following the quota share reallocation in 2017, and have been near or above 10,000 mt in all years between 2018-2024. Midwater trawl gears in groundfish and Pacific Whiting (hake) fisheries account for the majority of the recent catch.

Widow rockfish are a desirable market species and it is believed that discarding was low historically. However, management restrictions (e.g., trip limits) resulted in a substantial amount of discarding beginning in the early 1980s. Trawl rationalization was introduced in 2011. Between 2011 and 2024 very little discarding of widow rockfish is estimated to have occurred. Recent discards in the model informed by data from the West Coast Observer Program (WCGOP), and total catches (discards plus landings) are reported in addition to landings.

Table i: Recent landings for the bottom trawl, midwater trawl, at-sea hake, net, and hook-and-line fisheries and the total landings across fisheries and the total mortality (discards + landings) (mt). 100% mortality is assumed for discards and catch sources are described below.

Year	Bottom Trawl	Midwater Trawl	At-Sea Hake	Net	Hook-and-line	Total Landings	Total Mortality
2016	9.6	588.0	440.8	0.0	0.8	1,039.2	1,039.4
2017	35.9	4,852.1	1,455.2	0.0	2.7	6,345.9	6,361.5
2018	35.9	9,374.3	1,081.3	0.0	1.5	10,493.0	10,522.8
2019	27.9	8,157.9	1,101.6	0.0	2.1	9,289.4	9,315.3
2020	73.6	7,532.2	746.7	0.0	2.7	8,355.2	8,379.6
2021	103.7	10,141.3	617.3	0.0	4.5	10,866.9	10,899.7
2022	126.8	10,839.8	1,119.0	0.1	8.7	12,094.4	12,129.7
2023	82.3	10,228.0	673.2	0.0	7.1	10,990.6	11,023.5
2024	27.6	9,160.8	533.8	0.0	12.9	9,735.1	9,764.1
2025	35.6	9,345.0	1,455.2	0.0	2.7	10,838.5	10,868.1




Figure i: Landings of widow rockfish from 1916 to 2024 for bottom trawl, midwater trawl, net, and hook- and-line fisheries, and catches of widow rockfish for the foreign (1966–1976) and Pacific Whiting (hake) fisheries (green).

Data and Assessment

This assessment uses the length- and age-structured modeling software Stock Synthesis (version 3.30.23.1), while the update assessment in 2019 used 3.30.13. The coastwide population was modeled assuming separate growth and mortality parameters for each sex (a two-sex model) from 1916 to 2024, and forecasted beyond 2024.

Five fishing fleets were specified within the model: 1) a shorebased bottom trawl fleet with coastwide catches from 1916–2024, 2) a shorebased midwater trawl fleet with coastwide catches from 1979–2024, 3) a mostly midwater trawl fleet that targets Pacific Hake/Whiting (*Merluccius productus*) and includes a foreign and at-sea fleet with catches from 1975–2024, a domestic shorebased fleet that targeted Pacific Hake with catches from 1991–2024, and foreign vessels that targeted Pacific Hake and rockfish between 1966–1976, 4) a net fishery consisting of catches mostly from California from 1981–2024, and 5) a hook-and-line fishery (predominantly longline) with coastwide catches from 1916–2024. Landings are summarized in Figure i.

Data from three fishery-independent surveys were also included in the model: 1) the National Marine Fisheries Service (NMFS) Southwest Fisheries Science Center (SWFSC) and Northwest Fisheries Science Center (NWFSC)/Pacific Whiting Conservation Cooperative (PWCC) Midwater Trawl Survey that provides pre-recruit indices of abundance from 2004–2024, 2) the NMFS Triennial Shelf Survey which was conducted from 1977–2004

in depths less than 500 meters, and 3) the NWFSC West Coast Groundfish Bottom Trawl Survey (WCGBTS) which has been surveying the entire U.S. West Coast in depths between 55 and 1,280 meters since 2003. Three historical fishery-dependent CPUE indices were included as in the 2015 and 2019 assessments, derived from 1) Oregon bottom trawl (1984-1999), 2) Pacific Whiting at-sea foreign (1977-1988), and 3) Pacific whiting at-sea domestic fleets (1983-1998).

The data used in the assessment model consisted of survey abundance indices, length compositions, discard data, and age compositions. Model-based biomass indices and length compositions were determined for the NMFS Triennial Shelf and NWFSC West Coast Groundfish Bottom Trawl Surveys. Length and age compositions were also available from the five fisheries. Age data for all years of the WCGBTS were input as age-at-length compositions. Discard data for the bottom trawl, midwater trawl, and hook-and-line fisheries were available in various years in the form of discarded biomass and length compositions.

The base model estimated parameters for length-based selectivity for all fleets and surveys, retention curves based on length for the bottom trawl and midwater trawl fleets, a length-at-age relationship, natural mortality for males and females assuming lognormal priors, and recruitment deviations starting in 1900. A Beverton-Holt stock-recruitment function was used to model productivity and the steepness parameter was fixed at 0.72 based on a steepness meta-analysis for west coast rockfishes.

Estimation uncertainty in the base model was determined using approximate asymptotic 95% confidence intervals based on maximum likelihood theory. Natural mortality and steepness are major sources of uncertainty. We used high and low combinations of these parameters to define a range of states of nature, with results presented in a decision table.

Stock biomass and dynamics

The predicted spawning biomass from the base model generally showed a slight decline over the time series until 1966 when the foreign fleet began. A short, but sharp decline occurred, followed by a steep increase (peaking in 1979) due to strong recruitment. The spawning biomass declined rapidly with the developing domestic midwater fishery in the late 1970s and early 1980s. The stock continued to decline until 2000. A combination of strong recruitment and low catches resulted in a steady increase in spawning biomass through 2016. A target fishery for widow rockfish was reestablished in 2017. The stock exhibits a decline in most recent years with the increased catches since 2017, and lack of evidence for recent large recruitment events.

The 2024 spawning stock biomass (SSB) relative to unfished equilibrium spawning biomass is at 55, above the target of 40%.

Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in the estimated spawning biomass is high, especially in the early years. Spawning biomass is estimated to be at 45,443 mt in 2024, with an asymptotic 95% confidence interval of 45,046 - 45,046. The corresponding confidence interval for the relative SSB is 55.3% - 55.3%, for which the lower bound is below the management target of 40%.




Figure ii: Estimated female spawning biomass time-series from the base model (solid line) with an approximate asymptotic 95% confidence interval (dashed lines).




Figure iii: Estimated relative spawning biomass (depletion) with approximate 95% asymptotic confidence intervals (dashed lines) for the base case assessment model.

Table ii: Recent trend in estimated female spawning biomass (mt) and relative spawning biomass (depletion).

Year	Spawning Biomass	95% Confidence Interval	Estimated Depletion (%)	95% Confidence Interval
2,016	64,261	64,261 - 64,261	0.788677	0.789 - 0.789
2,017	65,863	65,863 - 65,863	0.808343	0.808 - 0.808
2,018	63,934	63,934 - 63,934	0.784662	0.785 - 0.785
2,019	60,055	60,055 - 60,055	0.737058	0.737 - 0.737
2,020	57,126	57,126 - 57,126	0.701113	0.701 - 0.701
2,021	55,346	55,346 - 55,346	0.679266	0.679 - 0.679
2,022	52,884	52,884 - 52,884	0.649048	0.649 - 0.649
2,023	49,359	49,359 - 49,359	0.605791	0.606 - 0.606
2,024	45,046	45,046 - 45,046	0.552851	0.553 - 0.553
2,025	40,184	40,184 - 40,184	0.493175	0.493 - 0.493
2,026	34,268	34,268 - 34,268	0.420573	0.421 - 0.421

Recruitment

Recruitment deviations were estimated for the entire time series modeled. There is little information regarding recruitment prior to 1965, and the uncertainty in these estimates is expressed in the model.

There are several very large, but uncertain, estimates of recruitment (in descending order of magnitude) in 1970, 2008, 2016, and 1971. Other large recruitment events (in descending order of magnitude) occurred in 1978, 1981, 2010, and 1991.

The five lowest recruitments (in ascending order) occurred in 2012, 2019, 2020, 2011, and 1976. Estimates of recruitment appear to be episodic and characterized by periods of low recruitment.




Figure iv: Time-series of estimated recruitments (medians as open circles) for the base case model with approximate asymptotic 95% confidence interval (vertical bars). Estimated mean unfished equilibrium recruitment (R_0) is shown as the closed circle with a 95% confidence interval at the beginning of the time series.

Table iii: Recent estimated trend in widow rockfish recruitment with approximate 95% confidence intervals determined from the base model.

Year	Estimated recruitment (number in thousands)	95% Confidence Interval	Recruitment Deviations	95% Confidence Interval
2,016	98,971	98,971 - 98,971	1.2488800	1.249, 1.249
2,017	27,557	27,557 - 27,557	-0.0326351	-0.033, -0.033
2,018	10,482	10,482 - 10,482	-1.0141200	-1.014, -1.014
2,019	8,202	8,202 - 8,202	-1.2731400	-1.273, -1.273
2,020	8,835	8,835 - 8,835	-1.2137700	-1.214, -1.214
2,021	18,753	18,753 - 18,753	-0.4782710	-0.478, -0.478
2,022	20,353	20,353 - 20,353	-0.4115310	-0.412, -0.412
2,023	29,951	29,951 - 29,951	-0.0365446	-0.037, -0.037
2,024	25,888	25,888 - 25,888	-0.1894370	-0.189, -0.189
2,025	30,867	30,867 - 30,867	0.0000000	0, 0
2,026	29,941	29,941 - 29,941	0.0000000	0, 0

Exploitation status

The spawning biomass of widow rockfish reached a low in 2001 before increasing due to low catch levels.

The lower bound of the 95% confidence interval of the estimated depletion dipped below the overfished threshold (25% minimum stock size threshold) in the very late 1990s and early 2000s. After recovering after 2016, the lower bound of the 95% confidence interval dipped below the 40% management target in 2024 but remains above the overfished threshold. The current depletion point estimate is greater than the spawning biomass target.

Table iv: Recent trend in spawning potential ratio and summary exploitation rate. Harvest rate is defined as catch divided by age 4+ biomass.

Year	Estimated (1-SPR)/(1- SPR50%)	~95% Confidence Interval	Harvest rate (proportion)	~95% Confidence Interval
2,016	0.163	0.163 - 0.163	0.008	0.008, 0.008
2,017	0.742	0.742 - 0.742	0.050	0.05, 0.05
2,018	1.078	1.078 - 1.078	0.086	0.086, 0.086
2,019	1.060	1.06 - 1.06	0.083	0.083, 0.083
2,020	1.046	1.046 - 1.046	0.073	0.073, 0.073
2,021	1.214	1.214 - 1.214	0.097	0.097, 0.097
2,022	1.275	1.275 - 1.275	0.117	0.117, 0.117
2,023	1.256	1.256 - 1.256	0.120	0.12, 0.12
2,024	1.267	1.267 - 1.267	0.122	0.122, 0.122
2,025	1.438	1.438 - 1.438	0.154	0.154, 0.154




Figure v: Time-series of estimated summary harvest rate (catch divided by age 4+ biomass) for the base case model (round points) with approximate 95% asymptotic confidence intervals (gray lines).




Figure vi: Trend in estimated fishing intensity (relative to the SPR management target) through 2024 with 95% asymptotic confidence intervals. One minus SPR is used so that higher exploitation rates occur on the upper portion of the y-axis. The relative management target is plotted as a horizontal line and values above this reflect harvests in excess of the overfishing proxy based on SPR50%.




Figure vii: Phase plot of estimated relative fishing intensity relative biomass for the base case model. The relative (1- SPR) is $(1-SPR)$ divided by 0.5 (one minus the SPR target). Lines through the final point (2024) show 95% intervals based on the asymptotic uncertainty for each dimension. The shaded ellipse is a 95% region which accounts for the estimated correlation between the two quantities.

Ecosystem considerations

Rockfish (*Sebastodes* spp.) are an important component of the California Current ecosystem along the U.S. West Coast, comprising more than sixty-five species filling various niches in both soft and hard bottom habitats from the nearshore to the continental slope, as well as near bottom and pelagic zones. Widow rockfish frequently aggregate in the pelagic zone.

Recruitment is one mechanism by which the ecosystem may directly impact the population dynamics of widow rockfish. The specific pathways through which environmental conditions exert influence on widow rockfish dynamics are unclear; however, changes in water temperature and currents, distribution of prey and predators, and the amount and timing of upwelling are all possible linkages. Changes in the environment may also result in changes in age-at-maturity, fecundity, growth, and survival which can affect how the status of the stock and its susceptibility to fishing are determined. Unfortunately, there are few data available for widow rockfish that provide insights into these effects.

Fishing has effects on both the age structure of a population as well as the target species habitat. Fishing often targets larger, older fish, and years of fishing mortality results in a truncated age-structure when compared to unfished conditions. Rockfish are often associated with habitats containing living structures such as sponges and corals, and fishing may alter that habitat to a less desirable state. This assessment provides insight on the effects of fishing on age structure, and recent studies on essential fish habitat are beginning to characterize important locations for rockfish throughout their life history; however there is little current information available to evaluate the specific effects of fishing on the ecosystem issues specific to widow rockfish.

Reference points

Reference points were calculated using the estimated selectivities and average catch distribution among fleets in the most recent five years of the model (2019-2024). Total yields (landings plus discards) were 5,594 mt when using an SPR50% reference harvest rate and with a 95% confidence interval of 5,594 to 5,594 mt. The spawning biomass equivalent to 40% of the unfished spawning output (SB40%) was 32,592 mt. Catches between the late 1990's and 2016 were well below the point estimate of potential long-term yields calculated using an SPR50% reference point, and the population is estimated to have increased continuously throughout that time period. However, catches from 2017 through 2024 were above the point estimate of potential long-term yields using an SPR50% reference point, exceeding the upper bound of the confidence interval in all years since 2018, and exceeding the point estimate by on average NaN%.

The predicted spawning biomass from the base model generally showed a slight decline until the late 1970s, followed by steep increase above unfished equilibrium biomass and reaching a peak in 1979. This was followed by a steep decrease up to the mid-1980s, and then a more gradual decease through 2000 (Figure [iii](#)). Between 2001 and 2016, the spawning biomass increased continuously due to small catches, and several years of high recruitment (though with lower than average recruitment in other recent years). The spawning biomass relative to unfished equilibrium spawning biomass climbed above the target of 40% of unfished spawning biomass in the early 2000's. It is estimated to still be above the target, though the lower bound of the 95% confidence interval lies between the target and minimum stock size threshold (Figure [iii](#)). The fishing intensity (relative 1-SPR) exceeded the current estimates of the harvest rate limit (SPR50%) throughout

the 1980s and early 1990s, and has again since 2018 (Figure v). Exploitation rates on widow rockfish between 2001 and 2016 were predicted to be far below target levels. In recent years, the stock has experienced exploitation rates that have been above the target level while the biomass level has remained above the target level (Figure vii).

The equilibrium yield plot is shown in Figure viii, based on a steepness value fixed at 0.72. The predicted maximum sustainable yield under the assumptions of this assessment occurs near 25% of equilibrium unfished spawning biomass.

Table v: Summary of reference points and management quantities for the base case model.

Reference Point	Estimate	Lower Interval	Upper Interval
Unfished Spawning Biomass (mt)	81,479.20	81,479.20	81,479.20
Unfished Age 4+ Biomass (mt)	151,093.00	151,093.00	151,093.00
Unfished Recruitment (R0)	33,951.00	33,951.00	33,951.00
2026 Spawning Biomass (mt)	34,267.90	34,267.90	34,267.90
2026 Fraction Unfished	0.42	0.42	0.42
Reference Points Based SB40%			
Proxy Spawning Biomass (mt) SB40%	32,591.70	32,591.70	32,591.70
SPR Resulting in SB40%	0.46	0.46	0.46
Exploitation Rate Resulting in SB40%	0.09	0.09	0.09
Yield with SPR Based On SB40% (mt)	5,865.90	5,865.90	5,865.90
Reference Points Based on SPR Proxy for MSY			
Proxy Spawning Biomass (mt) (SPR50)	36,352.30	36,352.30	36,352.30
SPR50	0.50		
Exploitation Rate Corresponding to SPR50	0.08	0.08	0.08
Yield with SPR50 at SB SPR (mt)	5,594.01	5,594.01	5,594.01
Reference Points Based on Estimated MSY Values			
Spawning Biomass (mt) at MSY (SB MSY)	21,615.50	21,615.50	21,615.50
SPR MSY	0.34	0.34	0.34
Exploitation Rate Corresponding to SPR MSY	0.13	0.13	0.13
MSY (mt)	6,270.27	6,270.27	6,270.27

Management performance

Exploitation rates on widow rockfish exceeded MSY proxy target harvest rates during the 1980s and 1990s and spawning biomass is predicted to have fallen below the proxy management target of 40%. Exploitation rates decreased in the late 1990s due to management restrictions, and have increased in recent years. Predicted catches in the last decade have not exceeded the annual catch limit (ACL) set by management.

Table vi: Recent trend in total catch and commercial landings (mt) relative to the management guidelines. Estimated total catch reflects the commercial landings plus the model estimated dead discarded biomass.

Year	OFL (mt)	ABC (mt)	ACL (mt)	Landings (mt)	Total Mortality (mt)
2016	3,990	3,790.00	2,000	1,039.24	1,039.41
2017	14,130	13,508.28	13,508	6,345.89	6,361.45
2018	13,237	12,654.57	12,655	10,493.03	10,522.76
2019	12,375	11,830.50	11,831	9,289.43	9,315.27
2020	11,714	11,198.58	11,199	8,355.24	8,379.58
2021	15,749	14,725.32	14,725	10,866.89	10,899.73
2022	14,826	13,788.18	13,788	12,094.40	12,129.69
2023	13,633	12,624.16	12,624	10,990.57	11,023.50
2024	12,453	11,481.67	11,482	9,735.12	9,764.11
2025	12,254	11,236.92	11,237	10,838.45	10,868.09

Harvest projections and decision table

Two categories of parameters that greatly contributed to uncertainty in the results were natural mortality (an important estimated parameter) and steepness (not estimated in the model). A combination of these two factors was used as the axis of uncertainty to define low and high states of nature. This differed from the 2019 assessment which included a third factor, 2013 recruitment strength. The 12.5% and 87.5% quantiles for female and male natural mortality (independently) were chosen as low and high values (0.115 yr^{-1} and 0.134 yr^{-1} for females; 0.127 yr^{-1} and 0.146 yr^{-1} for males). Steepness is probably the most important factor since it was fixed in the base model and is not incorporated in the estimation uncertainty. The 12.5% and 87.5% quantiles from the steepness prior (without widow rockfish data) were used to define the low and high values

of steepness (0.536 and 0.904). The low combination of these two factors defined the low state of nature and the high combination of these two factors defined the high state of nature. The predictions of spawning biomass in 2025 from the low and high states of nature are close to the 12.5% and 87.5% lognormal quantiles from the base model.

This assessment synthesizes many sources of data and estimates recruitment variability, thus it is classified as a Category 1 stock assessment. Therefore, the sigma to determine the catch reduction to account for scientific uncertainty is 0.50.

A twelve year projection of the base model with catches equal to the current ACL in 2025 and 2026 (10,669 mt and 9,824 mt, respectively) and catches of 9,000 mt for all later years and a catch allocation equal to the percentages for each fleet in 2024 predicts spawning biomass will decrease over the projection period for all states of nature (Table [viii](#)).

Projections with catches based on the predicted annual catch limit (ACL) using the SPR rate of 50%, the 40:10 control rule, and a 0.45 P* adjustment using a sigma of 0.50 from 2027 onward suggest that the spawning biomass will decrease over the projection period for all states of nature (Table [viii](#)). Predicted ACL catches range from 3,231 mt in 2027 to 4,392 mt in 2036.

Projections with catches based on the predicted annual catch limit (ACL) using the SPR rate of 50%, the 40:10 control rule, and a 0.25 P* adjustment using a sigma of 0.50 from 2027 onward suggest that the spawning biomass will decrease over the projection period for all states of nature. Predicted ACL catches range from 1,790 mt in 2027 to 1,886 mt in 2036.

Table vii: Projection of potential OFL, landings, and catch, summary biomass (age-4 and older), spawning biomass, and depletion for the base case model projected with total catch equal to the predicted ABC. The predicted OFL is the calculated total catch determined by FSPR=50%.

Year	Adopted OFL (mt)	Adopted ACL (mt)	Assumed Catch (mt)	OFL (mt)	Buffer ABC (mt)	ACL (mt)	Spawning Biomass (mt)
2026		10,868.10	11,382.00	10,391.77	10,392.00		34,267.9
2027		3,257.04					28,679.9
2028			3,739.66		3,281.94		27,879.0
2029			3,990.00		3,483.38		27,817.3
2030			4,300.89		3,762.86		28,260.4
2031			4,613.63		4,054.96		28,980.0
2032			4,879.79		4,315.79		29,793.7
2033			5,078.83		4,515.05		30,572.9
2034			5,220.57		4,652.60		31,264.2

Year	Adopted OFL (mt)	Adopted ACL (mt)	Assumed Catch (mt)	OFL (mt)	Buffer ABC (mt)	ACL (mt)	Spawning Biomass (mt)
2035				5,324.31		4,755.63	31,868.4
2036				5,405.10		4,833.43	32,400.2

[1] 40603.1

[1] 0.2753312

[1] 0.3109781

	Label	Value	StdDev	(Val - 1.0)/Stddev
SSB_Virgin	SSB_Virgin	8.17335e+04	7.18804e+03	NA
SSB_Initial	SSB_Initial	8.17335e+04	7.18804e+03	NA
SSB_1916	SSB_1916	8.15912e+04	1.14375e+04	NA
SSB_1917	SSB_1917	8.15264e+04	1.16058e+04	NA
SSB_1918	SSB_1918	8.14375e+04	1.17401e+04	NA
SSB_1919	SSB_1919	8.13402e+04	1.18476e+04	NA
SSB_1920	SSB_1920	8.12694e+04	1.19349e+04	NA
SSB_1921	SSB_1921	8.11988e+04	1.20087e+04	NA
SSB_1922	SSB_1922	8.11387e+04	1.20736e+04	NA
SSB_1923	SSB_1923	8.10847e+04	1.21314e+04	NA
SSB_1924	SSB_1924	8.10258e+04	1.21818e+04	NA
SSB_1925	SSB_1925	8.09809e+04	1.22241e+04	NA
SSB_1926	SSB_1926	8.09265e+04	1.22581e+04	NA
SSB_1927	SSB_1927	8.08507e+04	1.22842e+04	NA
SSB_1928	SSB_1928	8.07805e+04	1.23029e+04	NA
SSB_1929	SSB_1929	8.07003e+04	1.23146e+04	NA
SSB_1930	SSB_1930	8.06172e+04	1.23201e+04	NA
SSB_1931	SSB_1931	8.05149e+04	1.23198e+04	NA
SSB_1932	SSB_1932	8.04146e+04	1.23141e+04	NA
SSB_1933	SSB_1933	8.03082e+04	1.23034e+04	NA
SSB_1934	SSB_1934	8.02026e+04	1.22879e+04	NA
SSB_1935	SSB_1935	8.00863e+04	1.22678e+04	NA
SSB_1936	SSB_1936	7.99570e+04	1.22432e+04	NA
SSB_1937	SSB_1937	7.98107e+04	1.22139e+04	NA
SSB_1938	SSB_1938	7.96574e+04	1.21799e+04	NA
SSB_1939	SSB_1939	7.95029e+04	1.21411e+04	NA
SSB_1940	SSB_1940	7.93385e+04	1.20971e+04	NA
SSB_1941	SSB_1941	7.91335e+04	1.20477e+04	NA
SSB_1942	SSB_1942	7.89073e+04	1.19925e+04	NA
SSB_1943	SSB_1943	7.86536e+04	1.19310e+04	NA
SSB_1944	SSB_1944	7.81848e+04	1.18631e+04	NA
SSB_1945	SSB_1945	7.74377e+04	1.17882e+04	NA
SSB_1946	SSB_1946	7.63178e+04	1.17057e+04	NA
SSB_1947	SSB_1947	7.54883e+04	1.16137e+04	NA

SSB_1948	SSB_1948	7.49685e+04	1.15119e+04	NA
SSB_1949	SSB_1949	7.45372e+04	1.14004e+04	NA
SSB_1950	SSB_1950	7.41498e+04	1.12794e+04	NA
SSB_1951	SSB_1951	7.37116e+04	1.11490e+04	NA
SSB_1952	SSB_1952	7.31747e+04	1.10096e+04	NA
SSB_1953	SSB_1953	7.26259e+04	1.08611e+04	NA
SSB_1954	SSB_1954	7.21028e+04	1.07042e+04	NA
SSB_1955	SSB_1955	7.15811e+04	1.05398e+04	NA
SSB_1956	SSB_1956	7.10466e+04	1.03691e+04	NA
SSB_1957	SSB_1957	7.04443e+04	1.01939e+04	NA
SSB_1958	SSB_1958	6.97812e+04	1.00162e+04	NA
SSB_1959	SSB_1959	6.92104e+04	9.83793e+03	NA
SSB_1960	SSB_1960	6.87408e+04	9.66109e+03	NA
SSB_1961	SSB_1961	6.83033e+04	9.48754e+03	NA
SSB_1962	SSB_1962	6.80829e+04	9.31773e+03	NA
SSB_1963	SSB_1963	6.79659e+04	9.15089e+03	NA
SSB_1964	SSB_1964	6.81774e+04	8.98392e+03	NA
SSB_1965	SSB_1965	6.84004e+04	8.81274e+03	NA
SSB_1966	SSB_1966	6.88946e+04	8.63134e+03	NA
SSB_1967	SSB_1967	6.70918e+04	8.44004e+03	NA
SSB_1968	SSB_1968	6.51849e+04	8.23451e+03	NA
SSB_1969	SSB_1969	6.49434e+04	8.01073e+03	NA
SSB_1970	SSB_1970	6.58081e+04	7.76822e+03	NA
SSB_1971	SSB_1971	6.68215e+04	7.50781e+03	NA
SSB_1972	SSB_1972	6.77827e+04	7.23014e+03	NA
SSB_1973	SSB_1973	6.93983e+04	6.95337e+03	NA
SSB_1974	SSB_1974	7.17657e+04	6.68608e+03	NA
SSB_1975	SSB_1975	7.54885e+04	6.42365e+03	NA
SSB_1976	SSB_1976	8.04256e+04	6.15544e+03	NA
SSB_1977	SSB_1977	8.50744e+04	5.82591e+03	NA
SSB_1978	SSB_1978	8.83275e+04	5.39149e+03	NA
SSB_1979	SSB_1979	8.90318e+04	4.86288e+03	NA
SSB_1980	SSB_1980	8.39459e+04	4.31170e+03	NA
SSB_1981	SSB_1981	7.17673e+04	3.83215e+03	NA
SSB_1982	SSB_1982	5.72993e+04	3.49545e+03	NA
SSB_1983	SSB_1983	4.53939e+04	3.31427e+03	NA
SSB_1984	SSB_1984	4.31866e+04	3.25897e+03	NA
SSB_1985	SSB_1985	4.25015e+04	3.29463e+03	NA
SSB_1986	SSB_1986	4.28427e+04	3.38447e+03	NA
SSB_1987	SSB_1987	4.31680e+04	3.51437e+03	NA
SSB_1988	SSB_1988	4.13322e+04	3.66243e+03	NA
SSB_1989	SSB_1989	4.01405e+04	3.82917e+03	NA
SSB_1990	SSB_1990	3.72364e+04	4.02339e+03	NA

SSB_1991	SSB_1991	3.43787e+04	3.96435e+03	NA
SSB_1992	SSB_1992	3.35886e+04	3.98356e+03	NA
SSB_1993	SSB_1993	3.27815e+04	3.99451e+03	NA
SSB_1994	SSB_1994	3.05962e+04	4.00511e+03	NA
SSB_1995	SSB_1995	2.96892e+04	4.16443e+03	NA
SSB_1996	SSB_1996	2.88725e+04	4.42712e+03	NA
SSB_1997	SSB_1997	2.87706e+04	4.81046e+03	NA
SSB_1998	SSB_1998	2.82713e+04	5.22389e+03	NA
SSB_1999	SSB_1999	2.82616e+04	5.65945e+03	NA
SSB_2000	SSB_2000	2.80012e+04	6.02689e+03	NA
SSB_2001	SSB_2001	2.78636e+04	6.32957e+03	NA
SSB_2002	SSB_2002	2.86987e+04	6.60256e+03	NA
SSB_2003	SSB_2003	3.06321e+04	6.90930e+03	NA
SSB_2004	SSB_2004	3.32294e+04	7.28781e+03	NA
SSB_2005	SSB_2005	3.61426e+04	7.71527e+03	NA
SSB_2006	SSB_2006	3.89338e+04	8.10148e+03	NA
SSB_2007	SSB_2007	4.14576e+04	8.40836e+03	NA
SSB_2008	SSB_2008	4.34987e+04	8.59697e+03	NA
SSB_2009	SSB_2009	4.54639e+04	8.75371e+03	NA
SSB_2010	SSB_2010	4.74191e+04	8.88022e+03	NA
SSB_2011	SSB_2011	4.97611e+04	9.05605e+03	NA
SSB_2012	SSB_2012	5.22302e+04	9.23262e+03	NA
SSB_2013	SSB_2013	5.53635e+04	9.49937e+03	NA
SSB_2014	SSB_2014	5.88355e+04	9.80524e+03	NA
SSB_2015	SSB_2015	6.21020e+04	1.00688e+04	NA
SSB_2016	SSB_2016	6.47722e+04	1.02288e+04	NA
SSB_2017	SSB_2017	6.63683e+04	1.02297e+04	NA
SSB_2018	SSB_2018	6.44022e+04	1.01146e+04	NA
SSB_2019	SSB_2019	6.04648e+04	1.00615e+04	NA
SSB_2020	SSB_2020	5.74851e+04	1.01182e+04	NA
SSB_2021	SSB_2021	5.56818e+04	1.03444e+04	NA
SSB_2022	SSB_2022	5.32215e+04	1.07051e+04	NA
SSB_2023	SSB_2023	4.97239e+04	1.10454e+04	NA
SSB_2024	SSB_2024	4.54432e+04	1.12164e+04	NA
SSB_2025	SSB_2025	4.06031e+04	1.11793e+04	NA
SSB_2026	SSB_2026	3.48367e+04	1.10160e+04	NA
SSB_2027	SSB_2027	2.98219e+04	1.08204e+04	NA
SSB_2028	SSB_2028	2.88808e+04	9.71976e+03	NA
SSB_2029	SSB_2029	2.86934e+04	8.87489e+03	NA
SSB_2030	SSB_2030	2.90271e+04	8.26927e+03	NA
SSB_2031	SSB_2031	2.96551e+04	7.88133e+03	NA
SSB_2032	SSB_2032	3.03957e+04	7.70607e+03	NA
SSB_2033	SSB_2033	3.11186e+04	7.71057e+03	NA

SSB_2034	SSB_2034	3.17661e+04	7.83212e+03	NA
SSB_2035	SSB_2035	3.23341e+04	8.01283e+03	NA
SSB_2036	SSB_2036	3.28340e+04	8.21543e+03	NA
Recr_Virgin	Recr_Virgin	3.41029e+04	5.74731e+03	NA
Recr_Initial	Recr_Initial	3.41029e+04	5.74731e+03	NA
Recr_1916	Recr_1916	3.38250e+04	2.10287e+04	NA
Recr_1917	Recr_1917	3.37947e+04	2.10042e+04	NA
Recr_1918	Recr_1918	3.37607e+04	2.09768e+04	NA
Recr_1919	Recr_1919	3.37233e+04	2.09465e+04	NA
Recr_1920	Recr_1920	3.36836e+04	2.09141e+04	NA
Recr_1921	Recr_1921	3.36403e+04	2.08785e+04	NA
Recr_1922	Recr_1922	3.35933e+04	2.08398e+04	NA
Recr_1923	Recr_1923	3.35421e+04	2.07974e+04	NA
Recr_1924	Recr_1924	3.34858e+04	2.07508e+04	NA
Recr_1925	Recr_1925	3.34248e+04	2.07001e+04	NA
Recr_1926	Recr_1926	3.33576e+04	2.06441e+04	NA
Recr_1927	Recr_1927	3.32831e+04	2.05823e+04	NA
Recr_1928	Recr_1928	3.32018e+04	2.05146e+04	NA
Recr_1929	Recr_1929	3.31126e+04	2.04404e+04	NA
Recr_1930	Recr_1930	3.30149e+04	2.03590e+04	NA
Recr_1931	Recr_1931	3.29076e+04	2.02697e+04	NA
Recr_1932	Recr_1932	3.27907e+04	2.01723e+04	NA
Recr_1933	Recr_1933	3.26631e+04	2.00659e+04	NA
Recr_1934	Recr_1934	3.25242e+04	1.99499e+04	NA
Recr_1935	Recr_1935	3.23723e+04	1.98232e+04	NA
Recr_1936	Recr_1936	3.22060e+04	1.96845e+04	NA
Recr_1937	Recr_1937	3.20232e+04	1.95323e+04	NA
Recr_1938	Recr_1938	3.18220e+04	1.93649e+04	NA
Recr_1939	Recr_1939	3.15995e+04	1.91802e+04	NA
Recr_1940	Recr_1940	3.13551e+04	1.89777e+04	NA
Recr_1941	Recr_1941	3.10873e+04	1.87566e+04	NA
Recr_1942	Recr_1942	3.07971e+04	1.85175e+04	NA
Recr_1943	Recr_1943	3.04917e+04	1.82669e+04	NA
Recr_1944	Recr_1944	3.01739e+04	1.80086e+04	NA
Recr_1945	Recr_1945	2.98423e+04	1.77424e+04	NA
Recr_1946	Recr_1946	2.95007e+04	1.74722e+04	NA
Recr_1947	Recr_1947	2.91888e+04	1.72238e+04	NA
Recr_1948	Recr_1948	2.89036e+04	1.69939e+04	NA
Recr_1949	Recr_1949	2.86480e+04	1.67864e+04	NA
Recr_1950	Recr_1950	2.84408e+04	1.66159e+04	NA
Recr_1951	Recr_1951	2.83064e+04	1.65014e+04	NA
Recr_1952	Recr_1952	2.82752e+04	1.64661e+04	NA
Recr_1953	Recr_1953	2.83917e+04	1.65424e+04	NA

Recr_1954	Recr_1954	2.86998e+04	1.67630e+04	NA
Recr_1955	Recr_1955	2.92387e+04	1.71574e+04	NA
Recr_1956	Recr_1956	3.00201e+04	1.77342e+04	NA
Recr_1957	Recr_1957	3.09737e+04	1.84379e+04	NA
Recr_1958	Recr_1958	3.18807e+04	1.90918e+04	NA
Recr_1959	Recr_1959	3.24044e+04	1.94214e+04	NA
Recr_1960	Recr_1960	3.23334e+04	1.92549e+04	NA
Recr_1961	Recr_1961	3.19458e+04	1.88119e+04	NA
Recr_1962	Recr_1962	3.19515e+04	1.86300e+04	NA
Recr_1963	Recr_1963	3.28968e+04	1.91044e+04	NA
Recr_1964	Recr_1964	3.42894e+04	1.99650e+04	NA
Recr_1965	Recr_1965	3.59014e+04	2.09062e+04	NA
Recr_1966	Recr_1966	3.77074e+04	2.16766e+04	NA
Recr_1967	Recr_1967	3.76019e+04	2.09672e+04	NA
Recr_1968	Recr_1968	3.52647e+04	1.86020e+04	NA
Recr_1969	Recr_1969	3.02323e+04	1.68110e+04	NA
Recr_1970	Recr_1970	1.22196e+05	3.60375e+04	NA
Recr_1971	Recr_1971	9.99569e+04	2.70295e+04	NA
Recr_1972	Recr_1972	1.33456e+04	5.87151e+03	NA
Recr_1973	Recr_1973	1.12335e+04	4.28359e+03	NA
Recr_1974	Recr_1974	1.51836e+04	5.10934e+03	NA
Recr_1975	Recr_1975	3.18991e+04	6.28601e+03	NA
Recr_1976	Recr_1976	1.02562e+04	3.62408e+03	NA
Recr_1977	Recr_1977	5.88467e+04	9.05473e+03	NA
Recr_1978	Recr_1978	8.99057e+04	1.23217e+04	NA
Recr_1979	Recr_1979	2.85920e+04	6.42269e+03	NA
Recr_1980	Recr_1980	4.95135e+04	8.90225e+03	NA
Recr_1981	Recr_1981	8.22999e+04	1.26407e+04	NA
Recr_1982	Recr_1982	4.31228e+04	8.65756e+03	NA
Recr_1983	Recr_1983	2.73924e+04	6.58182e+03	NA
Recr_1984	Recr_1984	5.39490e+04	9.77916e+03	NA
Recr_1985	Recr_1985	3.92101e+04	8.06922e+03	NA
Recr_1986	Recr_1986	1.83995e+04	5.75270e+03	NA
Recr_1987	Recr_1987	4.69997e+04	1.01871e+04	NA
Recr_1988	Recr_1988	2.96616e+04	8.17368e+03	NA
Recr_1989	Recr_1989	2.16998e+04	6.88000e+03	NA
Recr_1990	Recr_1990	3.35302e+04	9.16401e+03	NA
Recr_1991	Recr_1991	6.01034e+04	1.42847e+04	NA
Recr_1992	Recr_1992	2.24677e+04	7.04156e+03	NA
Recr_1993	Recr_1993	2.82104e+04	8.46385e+03	NA
Recr_1994	Recr_1994	2.54832e+04	8.17438e+03	NA
Recr_1995	Recr_1995	1.58665e+04	5.99677e+03	NA
Recr_1996	Recr_1996	1.16617e+04	5.01205e+03	NA

Recr_1997	Recr_1997	1.78565e+04	7.12398e+03	NA
Recr_1998	Recr_1998	3.20377e+04	1.08714e+04	NA
Recr_1999	Recr_1999	4.23974e+04	1.30900e+04	NA
Recr_2000	Recr_2000	3.49304e+04	1.09118e+04	NA
Recr_2001	Recr_2001	1.92600e+04	6.71262e+03	NA
Recr_2002	Recr_2002	1.71137e+04	5.80657e+03	NA
Recr_2003	Recr_2003	1.66403e+04	5.99689e+03	NA
Recr_2004	Recr_2004	5.48643e+04	1.40382e+04	NA
Recr_2005	Recr_2005	1.05972e+04	4.46585e+03	NA
Recr_2006	Recr_2006	4.27387e+04	1.08477e+04	NA
Recr_2007	Recr_2007	1.06383e+04	4.38395e+03	NA
Recr_2008	Recr_2008	1.00500e+05	2.20592e+04	NA
Recr_2009	Recr_2009	1.32014e+04	4.74708e+03	NA
Recr_2010	Recr_2010	6.57660e+04	1.43077e+04	NA
Recr_2011	Recr_2011	9.73072e+03	3.57881e+03	NA
Recr_2012	Recr_2012	6.41589e+03	2.37187e+03	NA
Recr_2013	Recr_2013	5.64989e+04	1.24612e+04	NA
Recr_2014	Recr_2014	2.34753e+04	6.87366e+03	NA
Recr_2015	Recr_2015	1.83953e+04	6.31604e+03	NA
Recr_2016	Recr_2016	1.00038e+05	2.48641e+04	NA
Recr_2017	Recr_2017	2.78917e+04	9.64807e+03	NA
Recr_2018	Recr_2018	1.06147e+04	4.32580e+03	NA
Recr_2019	Recr_2019	8.28856e+03	3.59146e+03	NA
Recr_2020	Recr_2020	8.90733e+03	4.10245e+03	NA
Recr_2021	Recr_2021	1.88891e+04	8.97132e+03	NA
Recr_2022	Recr_2022	2.04638e+04	1.18661e+04	NA
Recr_2023	Recr_2023	3.01145e+04	1.82039e+04	NA
Recr_2024	Recr_2024	2.60457e+04	1.59171e+04	NA
Recr_2025	Recr_2025	3.10454e+04	1.96163e+04	NA
Recr_2026	Recr_2026	3.01561e+04	1.91579e+04	NA
Recr_2027	Recr_2027	2.91668e+04	1.86714e+04	NA
Recr_2028	Recr_2028	2.89518e+04	1.84882e+04	NA
Recr_2029	Recr_2029	2.89077e+04	1.84016e+04	NA
Recr_2030	Recr_2030	2.89860e+04	1.83928e+04	NA
Recr_2031	Recr_2031	2.91295e+04	1.84333e+04	NA
Recr_2032	Recr_2032	2.92928e+04	1.84973e+04	NA
Recr_2033	Recr_2033	2.94465e+04	1.85654e+04	NA
Recr_2034	Recr_2034	2.95794e+04	1.86279e+04	NA
Recr_2035	Recr_2035	2.96926e+04	1.86828e+04	NA
Recr_2036	Recr_2036	2.97896e+04	1.87305e+04	NA
SPRratio_1916	SPRratio_1916	1.17788e-02	1.94525e-03	-
5.08018e+02				
SPRratio_1917	SPRratio_1917	1.82936e-02	3.04149e-03	-

3.22771e+02			
SPRratio_1918	SPRratio_1918	2.10278e-02	3.51914e-03
2.78185e+02			-
SPRratio_1919	SPRratio_1919	1.45724e-02	2.46068e-03
4.00470e+02			-
SPRratio_1920	SPRratio_1920	1.49250e-02	2.53282e-03
3.88925e+02			-
SPRratio_1921	SPRratio_1921	1.23983e-02	2.11629e-03
4.66666e+02			-
SPRratio_1922	SPRratio_1922	1.07786e-02	1.84930e-03
5.34916e+02			-
SPRratio_1923	SPRratio_1923	1.18406e-02	2.03890e-03
4.84653e+02			-
SPRratio_1924	SPRratio_1924	7.62557e-03	1.32015e-03
7.51712e+02			-
SPRratio_1925	SPRratio_1925	9.44319e-03	1.63860e-03
6.04513e+02			-
SPRratio_1926	SPRratio_1926	1.43340e-02	2.48843e-03
3.96100e+02			-
SPRratio_1927	SPRratio_1927	1.19105e-02	2.07438e-03
4.76330e+02			-
SPRratio_1928	SPRratio_1928	1.35517e-02	2.36304e-03
4.17449e+02			-
SPRratio_1929	SPRratio_1929	1.31514e-02	2.29725e-03
4.29578e+02			-
SPRratio_1930	SPRratio_1930	1.70437e-02	2.97687e-03
3.30198e+02			-
SPRratio_1931	SPRratio_1931	1.51827e-02	2.65727e-03
3.70612e+02			-
SPRratio_1932	SPRratio_1932	1.52258e-02	2.66752e-03
3.69173e+02			-
SPRratio_1933	SPRratio_1933	1.30579e-02	2.29168e-03
4.30663e+02			-
SPRratio_1934	SPRratio_1934	1.38535e-02	2.43203e-03
4.05483e+02			-
SPRratio_1935	SPRratio_1935	1.49742e-02	2.62896e-03
3.74683e+02			-
SPRratio_1936	SPRratio_1936	1.69209e-02	2.96979e-03
3.31027e+02			-
SPRratio_1937	SPRratio_1937	1.58401e-02	2.78241e-03
3.53707e+02			-
SPRratio_1938	SPRratio_1938	1.28597e-02	2.26233e-03
4.36338e+02			-

SPRratio_1939 4.77701e+02	SPRratio_1939 1.17511e-02 2.06876e-03	-
SPRratio_1940 3.02569e+02	SPRratio_1940 1.84776e-02 3.24396e-03	-
SPRratio_1941 2.81337e+02	SPRratio_1941 1.98494e-02 3.48390e-03	-
SPRratio_1942 2.48198e+02	SPRratio_1942 2.24487e-02 3.93859e-03	-
SPRratio_1943 7.33271e+01	SPRratio_1943 7.36277e-02 1.26334e-02	-
SPRratio_1944 3.66255e+01	SPRratio_1944 1.40692e-01 2.34620e-02	-
SPRratio_1945 2.08650e+01	SPRratio_1945 2.29866e-01 3.69103e-02	-
SPRratio_1946 2.90089e+01	SPRratio_1946 1.71619e-01 2.85562e-02	-
SPRratio_1947 5.23691e+01	SPRratio_1947 9.95716e-02 1.71939e-02	-
SPRratio_1948 7.04067e+01	SPRratio_1948 7.51892e-02 1.31353e-02	-
SPRratio_1949 8.96384e+01	SPRratio_1949 5.97076e-02 1.04898e-02	-
SPRratio_1950 7.88097e+01	SPRratio_1950 6.77196e-02 1.18295e-02	-
SPRratio_1951 5.98324e+01	SPRratio_1951 8.81882e-02 1.52394e-02	-
SPRratio_1952 6.08775e+01	SPRratio_1952 8.68781e-02 1.49993e-02	-
SPRratio_1953 7.00285e+01	SPRratio_1953 7.61821e-02 1.31920e-02	-
SPRratio_1954 7.30583e+01	SPRratio_1954 7.33737e-02 1.26834e-02	-
SPRratio_1955 7.00531e+01	SPRratio_1955 7.66192e-02 1.31812e-02	-
SPRratio_1956 5.37281e+01	SPRratio_1956 9.88456e-02 1.67725e-02	-
SPRratio_1957 4.19732e+01	SPRratio_1957 1.24717e-01 2.08534e-02	-
SPRratio_1958 4.52669e+01	SPRratio_1958 1.16576e-01 1.95159e-02	-
SPRratio_1959 4.71014e+01	SPRratio_1959 1.12690e-01 1.88383e-02	-
SPRratio_1960	SPRratio_1960 1.35524e-01 2.23368e-02	-

3.87018e+01			
SPRratio_1961	SPRratio_1961	1.14926e-01	1.90570e-02
4.64435e+01			-
SPRratio_1962	SPRratio_1962	1.28138e-01	2.10192e-02
4.14792e+01			-
SPRratio_1963	SPRratio_1963	7.51567e-02	1.25596e-02
7.36364e+01			-
SPRratio_1964	SPRratio_1964	1.02492e-01	1.67859e-02
5.34680e+01			-
SPRratio_1965	SPRratio_1965	4.51720e-02	7.51914e-03
1.26986e+02			-
SPRratio_1966	SPRratio_1966	5.52072e-01	6.91075e-02
6.48162e+00			-
SPRratio_1967	SPRratio_1967	6.13026e-01	7.46739e-02
5.18218e+00			-
SPRratio_1968	SPRratio_1968	3.64849e-01	5.23581e-02
1.21309e+01			-
SPRratio_1969	SPRratio_1969	1.46723e-01	2.32852e-02
3.66446e+01			-
SPRratio_1970	SPRratio_1970	1.45389e-01	2.26319e-02
3.77613e+01			-
SPRratio_1971	SPRratio_1971	1.80275e-01	2.68549e-02
3.05242e+01			-
SPRratio_1972	SPRratio_1972	1.51607e-01	2.21469e-02
3.83075e+01			-
SPRratio_1973	SPRratio_1973	1.84147e-01	2.56643e-02
3.17894e+01			-
SPRratio_1974	SPRratio_1974	1.43781e-01	1.96101e-02
4.36621e+01			-
SPRratio_1975	SPRratio_1975	1.33356e-01	1.74042e-02
4.97950e+01			-
SPRratio_1976	SPRratio_1976	1.83093e-01	2.21101e-02
3.69472e+01			-
SPRratio_1977	SPRratio_1977	1.38925e-01	1.58611e-02
5.42886e+01			-
SPRratio_1978	SPRratio_1978	1.88641e-01	1.97806e-02
4.10178e+01			-
SPRratio_1979	SPRratio_1979	7.89572e-01	5.39993e-02
3.89687e+00			-
SPRratio_1980	SPRratio_1980	1.32497e+00	5.50862e-02
SPRratio_1981	SPRratio_1981	1.55413e+00	4.72262e-02
SPRratio_1982	SPRratio_1982	1.64328e+00	4.32561e-02
SPRratio_1983	SPRratio_1983	1.36409e+00	6.20350e-02
			5.86907e+00

SPRratio_1984	SPRratio_1984	1.36239e+00	6.34946e-02	5.70737e+00
SPRratio_1985	SPRratio_1985	1.28982e+00	6.70777e-02	4.32061e+00
SPRratio_1986	SPRratio_1986	1.29459e+00	6.76460e-02	4.35487e+00
SPRratio_1987	SPRratio_1987	1.42644e+00	6.34460e-02	6.72124e+00
SPRratio_1988	SPRratio_1988	1.33898e+00	7.10435e-02	4.77143e+00
SPRratio_1989	SPRratio_1989	1.44646e+00	6.89498e-02	6.47513e+00
SPRratio_1990	SPRratio_1990	1.44165e+00	6.37377e-02	6.92918e+00
SPRratio_1991	SPRratio_1991	1.26710e+00	7.49052e-02	3.56587e+00
SPRratio_1992	SPRratio_1992	1.26194e+00	7.68908e-02	3.40665e+00
SPRratio_1993	SPRratio_1993	1.40956e+00	7.29368e-02	5.61531e+00
SPRratio_1994	SPRratio_1994	1.32300e+00	8.37820e-02	3.85519e+00
SPRratio_1995	SPRratio_1995	1.36307e+00	8.73006e-02	4.15885e+00
SPRratio_1996	SPRratio_1996	1.32091e+00	9.72645e-02	3.29939e+00
SPRratio_1997	SPRratio_1997	1.34255e+00	1.04852e-01	3.26697e+00
SPRratio_1998	SPRratio_1998	1.19479e+00	1.24997e-01	1.55838e+00
SPRratio_1999	SPRratio_1999	1.12740e+00	1.37211e-01	9.28479e-01
SPRratio_2000	SPRratio_2000	9.87969e-01	1.49141e-01	-
8.06709e-02				
SPRratio_2001	SPRratio_2001	6.24194e-01	1.25371e-01	-
2.99755e+00				
SPRratio_2002	SPRratio_2002	1.81579e-01	4.70621e-02	-
1.73902e+01				
SPRratio_2003	SPRratio_2003	1.80714e-02	4.95902e-03	-
1.98009e+02				
SPRratio_2004	SPRratio_2004	3.52530e-02	9.35287e-03	-
1.03150e+02				
SPRratio_2005	SPRratio_2005	6.62870e-02	1.70374e-02	-
5.48037e+01				
SPRratio_2006	SPRratio_2006	6.66740e-02	1.67759e-02	-
5.56348e+01				
SPRratio_2007	SPRratio_2007	6.86305e-02	1.68964e-02	-
5.51225e+01				
SPRratio_2008	SPRratio_2008	7.18004e-02	1.72018e-02	-
5.39596e+01				
SPRratio_2009	SPRratio_2009	4.76235e-02	1.12504e-02	-
8.46529e+01				
SPRratio_2010	SPRratio_2010	4.36794e-02	1.00657e-02	-
9.50077e+01				
SPRratio_2011	SPRratio_2011	4.87426e-02	1.08127e-02	-
8.79756e+01				
SPRratio_2012	SPRratio_2012	5.96679e-02	1.28535e-02	-
7.31576e+01				

SPRratio_2013 4.93418e+01	SPRratio_2013 9.16516e-02 1.84093e-02	-
SPRratio_2014 3.42895e+01	SPRratio_2014 1.30693e-01 2.53520e-02	-
SPRratio_2015 3.18315e+01	SPRratio_2015 1.44426e-01 2.68782e-02	-
SPRratio_2016 2.89600e+01	SPRratio_2016 1.61198e-01 2.89641e-02	-
SPRratio_2017 2.72133e+00	SPRratio_2017 7.36208e-01 9.69351e-02	-
SPRratio_2018 01	SPRratio_2018 1.07014e+00 1.15216e-01	6.08731e-
SPRratio_2019 01	SPRratio_2019 1.05353e+00 1.21258e-01	4.41493e-
SPRratio_2020 01	SPRratio_2020 1.04078e+00 1.27910e-01	3.18803e-
SPRratio_2021	SPRratio_2021 1.20989e+00 1.35541e-01	1.54850e+00
SPRratio_2022	SPRratio_2022 1.27030e+00 1.43380e-01	1.88516e+00
SPRratio_2023	SPRratio_2023 1.24828e+00 1.59211e-01	1.55945e+00
SPRratio_2024	SPRratio_2024 1.25706e+00 1.77648e-01	1.44703e+00
SPRratio_2025	SPRratio_2025 1.41766e+00 1.83504e-01	2.27603e+00
SPRratio_2026	SPRratio_2026 1.49286e+00 1.96536e-01	2.50772e+00
SPRratio_2027 4.22837e-01	SPRratio_2027 9.72732e-01 6.44878e-02	-
SPRratio_2028 6.21686e-01	SPRratio_2028 9.61994e-01 6.11333e-02	-
SPRratio_2029 7.57138e-01	SPRratio_2029 9.57833e-01 5.56932e-02	-
SPRratio_2030 7.41735e-01	SPRratio_2030 9.57977e-01 5.66544e-02	-
SPRratio_2031 5.43238e-01	SPRratio_2031 9.59684e-01 7.42143e-02	-
SPRratio_2032 3.93321e-01	SPRratio_2032 9.62617e-01 9.50438e-02	-
SPRratio_2033 3.15859e-01	SPRratio_2033 9.65106e-01 1.10472e-01	-
SPRratio_2034 2.80481e-01	SPRratio_2034 9.66145e-01 1.20702e-01	-
SPRratio_2035 2.57561e-01	SPRratio_2035 9.67133e-01 1.27609e-01	-
SPRratio_2036 2.53934e-01	SPRratio_2036 9.66683e-01 1.31203e-01	-
F_1916	F_1916 5.18135e-04 7.57618e-05	-

1.31924e+04			
F_1917	F_1917	8.06852e-04	1.19247e-04
8.37922e+03			-
F_1918	F_1918	9.28490e-04	1.38426e-04
7.21736e+03			-
F_1919	F_1919	6.41739e-04	9.63492e-05
1.03723e+04			-
F_1920	F_1920	6.57355e-04	9.93635e-05
1.00574e+04			-
F_1921	F_1921	5.45501e-04	8.29515e-05
1.20487e+04			-
F_1922	F_1922	4.73922e-04	7.24386e-05
1.37982e+04			-
F_1923	F_1923	5.20841e-04	7.99597e-05
1.24998e+04			-
F_1924	F_1924	3.34810e-04	5.15942e-05
1.93756e+04			-
F_1925	F_1925	4.14706e-04	6.40961e-05
1.55951e+04			-
F_1926	F_1926	6.31423e-04	9.78355e-05
1.02148e+04			-
F_1927	F_1927	5.24901e-04	8.15100e-05
1.22620e+04			-
F_1928	F_1928	5.98186e-04	9.30429e-05
1.07413e+04			-
F_1929	F_1929	5.81634e-04	9.05872e-05
1.10327e+04			-
F_1930	F_1930	7.53461e-04	1.17459e-04
8.50722e+03			-
F_1931	F_1931	6.71016e-04	1.04688e-04
9.54577e+03			-
F_1932	F_1932	6.73255e-04	1.05080e-04
9.51013e+03			-
F_1933	F_1933	5.79500e-04	9.04600e-05
1.10482e+04			-
F_1934	F_1934	6.14083e-04	9.58395e-05
1.04277e+04			-
F_1935	F_1935	6.63562e-04	1.03521e-04
9.65346e+03			-
F_1936	F_1936	7.49083e-04	1.16797e-04
8.55547e+03			-
F_1937	F_1937	7.03092e-04	1.09548e-04
9.12202e+03			-

F_1938	F_1938	5.70879e-04	8.88600e-05	-
1.12472e+04				
F_1939	F_1939	5.23438e-04	8.13652e-05	-
1.22838e+04				
F_1940	F_1940	8.27220e-04	1.28375e-04	-
7.78325e+03				
F_1941	F_1941	8.91573e-04	1.38139e-04	-
7.23264e+03				
F_1942	F_1942	1.01456e-03	1.56904e-04	-
6.36686e+03				
F_1943	F_1943	3.40830e-03	5.26013e-04	-
1.89461e+03				
F_1944	F_1944	6.71520e-03	1.03623e-03	-
9.58556e+02				
F_1945	F_1945	1.14413e-02	1.76987e-03	-
5.58547e+02				
F_1946	F_1946	8.29976e-03	1.29188e-03	-
7.67641e+02				
F_1947	F_1947	4.64730e-03	7.24790e-04	-
1.37330e+03				
F_1948	F_1948	3.47757e-03	5.40996e-04	-
1.84201e+03				
F_1949	F_1949	2.74126e-03	4.24603e-04	-
2.34868e+03				
F_1950	F_1950	3.11885e-03	4.80379e-04	-
2.07520e+03				
F_1951	F_1951	4.10752e-03	6.28995e-04	-
1.58331e+03				
F_1952	F_1952	4.04705e-03	6.16372e-04	-
1.61583e+03				
F_1953	F_1953	3.53705e-03	5.35391e-04	-
1.86119e+03				
F_1954	F_1954	3.40096e-03	5.11039e-04	-
1.95014e+03				
F_1955	F_1955	3.55718e-03	5.30205e-04	-
1.87935e+03				
F_1956	F_1956	4.62967e-03	6.84160e-04	-
1.45488e+03				
F_1957	F_1957	5.90964e-03	8.66141e-04	-
1.14772e+03				
F_1958	F_1958	5.49607e-03	7.99244e-04	-
1.24431e+03				
F_1959	F_1959	5.29548e-03	7.62971e-04	-

1.30373e+03			
F_1960	F_1960	6.42280e-03	9.15514e-04
1.08527e+03			-
F_1961	F_1961	5.37985e-03	7.58201e-04
1.31182e+03			-
F_1962	F_1962	6.01616e-03	8.35496e-04
1.18969e+03			-
F_1963	F_1963	3.43420e-03	4.68937e-04
2.12516e+03			-
F_1964	F_1964	4.73874e-03	6.32869e-04
1.57262e+03			-
F_1965	F_1965	2.03428e-03	2.65434e-04
3.75975e+03			-
F_1966	F_1966	3.31062e-02	4.19889e-03
2.30274e+02			-
F_1967	F_1967	3.80389e-02	4.81592e-03
1.99746e+02			-
F_1968	F_1968	1.92471e-02	2.42862e-03
4.03832e+02			-
F_1969	F_1969	6.84134e-03	8.38864e-04
1.18393e+03			-
F_1970	F_1970	6.74103e-03	7.87811e-04
1.26078e+03			-
F_1971	F_1971	8.48039e-03	9.38641e-04
1.05634e+03			-
F_1972	F_1972	7.05019e-03	7.38153e-04
1.34518e+03			-
F_1973	F_1973	8.78367e-03	8.76145e-04
1.13134e+03			-
F_1974	F_1974	6.17020e-03	5.61577e-04
1.76971e+03			-
F_1975	F_1975	5.37263e-03	4.25455e-04
2.33780e+03			-
F_1976	F_1976	7.89314e-03	5.67640e-04
1.74778e+03			-
F_1977	F_1977	6.49398e-03	4.22912e-04
2.34920e+03			-
F_1978	F_1978	9.48165e-03	5.54908e-04
1.78501e+03			-
F_1979	F_1979	5.74916e-02	2.99749e-03
3.14433e+02			-
F_1980	F_1980	1.45874e-01	7.14472e-03
1.19546e+02			-

F_1981	F_1981	2.13365e-01	1.10506e-02	-
7.11849e+01				
F_1982	F_1982	2.36635e-01	1.40334e-02	-
5.43964e+01				
F_1983	F_1983	1.30708e-01	9.29970e-03	-
9.34752e+01				
F_1984	F_1984	1.32041e-01	9.52306e-03	-
9.11429e+01				
F_1985	F_1985	1.13541e-01	8.47898e-03	-
1.04548e+02				
F_1986	F_1986	1.17956e-01	9.01939e-03	-
9.77942e+01				
F_1987	F_1987	1.62743e-01	1.28872e-02	-
6.49683e+01				
F_1988	F_1988	1.37643e-01	1.21068e-02	-
7.12292e+01				
F_1989	F_1989	1.69781e-01	1.62332e-02	-
5.11434e+01				
F_1990	F_1990	1.73372e-01	1.41027e-02	-
5.86151e+01				
F_1991	F_1991	1.19636e-01	1.07052e-02	-
8.22368e+01				
F_1992	F_1992	1.18286e-01	1.11782e-02	-
7.88781e+01				
F_1993	F_1993	1.62019e-01	1.63183e-02	-
5.13523e+01				
F_1994	F_1994	1.34775e-01	1.53206e-02	-
5.64746e+01				
F_1995	F_1995	1.35168e-01	1.74267e-02	-
4.96270e+01				
F_1996	F_1996	1.25865e-01	1.81102e-02	-
4.82676e+01				
F_1997	F_1997	1.35952e-01	2.17150e-02	-
3.97904e+01				
F_1998	F_1998	1.04814e-01	1.94623e-02	-
4.59958e+01				
F_1999	F_1999	9.77416e-02	1.94693e-02	-
4.63426e+01				
F_2000	F_2000	8.24625e-02	1.76610e-02	-
5.19528e+01				
F_2001	F_2001	4.12784e-02	9.35206e-03	-
1.02514e+02				
F_2002	F_2002	8.24711e-03	1.90947e-03	-

5.19387e+02			
F_2003	F_2003	7.10318e-04	1.59507e-04
6.26488e+03			-
F_2004	F_2004	1.37689e-03	3.01209e-04
3.31538e+03			-
F_2005	F_2005	2.66461e-03	5.67355e-04
1.75787e+03			-
F_2006	F_2006	2.78498e-03	5.76666e-04
1.72928e+03			-
F_2007	F_2007	2.99975e-03	6.04799e-04
1.64848e+03			-
F_2008	F_2008	3.07204e-03	6.04919e-04
1.64804e+03			-
F_2009	F_2009	2.06554e-03	3.95672e-04
2.52212e+03			-
F_2010	F_2010	1.87267e-03	3.49950e-04
2.85220e+03			-
F_2011	F_2011	2.20216e-03	3.98018e-04
2.50692e+03			-
F_2012	F_2012	2.46182e-03	4.31771e-04
2.31034e+03			-
F_2013	F_2013	4.14207e-03	7.04848e-04
1.41287e+03			-
F_2014	F_2014	5.86336e-03	9.67444e-04
1.02759e+03			-
F_2015	F_2015	7.04018e-03	1.12927e-03
8.79293e+02			-
F_2016	F_2016	8.40338e-03	1.31287e-03
7.55288e+02			-
F_2017	F_2017	4.96841e-02	7.59219e-03
1.25170e+02			-
F_2018	F_2018	8.55249e-02	1.33570e-02
6.84641e+01			-
F_2019	F_2019	8.27456e-02	1.37616e-02
6.66530e+01			-
F_2020	F_2020	7.23536e-02	1.29970e-02
7.13737e+01			-
F_2021	F_2021	9.65786e-02	1.84102e-02
4.90719e+01			-
F_2022	F_2022	1.16363e-01	2.40923e-02
3.66771e+01			-
F_2023	F_2023	1.19037e-01	2.73596e-02
3.21994e+01			-

F_2024	F_2024	1.20756e-01	3.09734e-02	-
2.83871e+01				
F_2025	F_2025	1.49776e-01	4.27753e-02	-
1.98765e+01				
F_2026	F_2026	1.60113e-01	5.19318e-02	-
1.61729e+01				
F_2027	F_2027	6.31838e-02	1.07845e-02	-
8.68666e+01				
F_2028	F_2028	6.25498e-02	9.41691e-03	-
9.95497e+01				
F_2029	F_2029	6.36649e-02	8.95371e-03	-
1.04575e+02				
F_2030	F_2030	6.58041e-02	9.62413e-03	-
9.70681e+01				
F_2031	F_2031	6.81801e-02	1.04740e-02	-
8.89649e+01				
F_2032	F_2032	7.02232e-02	1.12619e-02	-
8.25596e+01				
F_2033	F_2033	7.15552e-02	1.20934e-02	-
7.67730e+01				
F_2034	F_2034	7.21756e-02	1.29191e-02	-
7.18182e+01				
F_2035	F_2035	7.24565e-02	1.36864e-02	-
6.77710e+01				
F_2036	F_2036	7.23958e-02	1.40217e-02	-
6.61549e+01				
Bratio_1917	Bratio_1917	9.97465e-01	1.11841e-01	-
2.26632e-02				
Bratio_1918	Bratio_1918	9.96378e-01	1.13969e-01	-
3.17775e-02				
Bratio_1919	Bratio_1919	9.95187e-01	1.15676e-01	-
4.16055e-02				
Bratio_1920	Bratio_1920	9.94321e-01	1.17068e-01	-
4.85067e-02				
Bratio_1921	Bratio_1921	9.93458e-01	1.18252e-01	-
5.53234e-02				
Bratio_1922	Bratio_1922	9.92722e-01	1.19303e-01	-
6.10016e-02				
Bratio_1923	Bratio_1923	9.92061e-01	1.20248e-01	-
6.60221e-02				
Bratio_1924	Bratio_1924	9.91340e-01	1.21085e-01	-
7.15178e-02				
Bratio_1925	Bratio_1925	9.90791e-01	1.21806e-01	-

7.56003e-02		
Bratio_1926	Bratio_1926 9.90126e-01	1.22409e-01 -
8.06633e-02		
Bratio_1927	Bratio_1927 9.89198e-01	1.22901e-01 -
8.78920e-02		
Bratio_1928	Bratio_1928 9.88340e-01	1.23290e-01 -
9.45756e-02		
Bratio_1929	Bratio_1929 9.87358e-01	1.23586e-01 -
1.02291e-01		
Bratio_1930	Bratio_1930 9.86341e-01	1.23799e-01 -
1.10332e-01		
Bratio_1931	Bratio_1931 9.85090e-01	1.23939e-01 -
1.20299e-01		
Bratio_1932	Bratio_1932 9.83863e-01	1.24013e-01 -
1.30126e-01		
Bratio_1933	Bratio_1933 9.82560e-01	1.24027e-01 -
1.40610e-01		
Bratio_1934	Bratio_1934 9.81269e-01	1.23988e-01 -
1.51070e-01		
Bratio_1935	Bratio_1935 9.79846e-01	1.23898e-01 -
1.62665e-01		
Bratio_1936	Bratio_1936 9.78264e-01	1.23761e-01 -
1.75632e-01		
Bratio_1937	Bratio_1937 9.76474e-01	1.23578e-01 -
1.90372e-01		
Bratio_1938	Bratio_1938 9.74598e-01	1.23349e-01 -
2.05934e-01		
Bratio_1939	Bratio_1939 9.72709e-01	1.23075e-01 -
2.21745e-01		
Bratio_1940	Bratio_1940 9.70697e-01	1.22753e-01 -
2.38714e-01		
Bratio_1941	Bratio_1941 9.68189e-01	1.22382e-01 -
2.59932e-01		
Bratio_1942	Bratio_1942 9.65421e-01	1.21960e-01 -
2.83527e-01		
Bratio_1943	Bratio_1943 9.62317e-01	1.21482e-01 -
3.10191e-01		
Bratio_1944	Bratio_1944 9.56582e-01	1.20944e-01 -
3.58994e-01		
Bratio_1945	Bratio_1945 9.47441e-01	1.20340e-01 -
4.36751e-01		
Bratio_1946	Bratio_1946 9.33740e-01	1.19661e-01 -
5.53736e-01		

Bratio_1947 6.42696e-01	Bratio_1947 9.23591e-01 1.18889e-01	-
Bratio_1948 7.01312e-01	Bratio_1948 9.17230e-01 1.18022e-01	-
Bratio_1949 7.52100e-01	Bratio_1949 9.11953e-01 1.17068e-01	-
Bratio_1950 7.99667e-01	Bratio_1950 9.07214e-01 1.16031e-01	-
Bratio_1951 8.54092e-01	Bratio_1951 9.01853e-01 1.14914e-01	-
Bratio_1952 9.20841e-01	Bratio_1952 8.95284e-01 1.13718e-01	-
Bratio_1953 9.90973e-01	Bratio_1953 8.88569e-01 1.12446e-01	-
Bratio_1954 1.06052e+00	Bratio_1954 8.82169e-01 1.11106e-01	-
Bratio_1955 1.13220e+00	Bratio_1955 8.75786e-01 1.09711e-01	-
Bratio_1956 1.20761e+00	Bratio_1956 8.69246e-01 1.08275e-01	-
Bratio_1957 1.29308e+00	Bratio_1957 8.61877e-01 1.06817e-01	-
Bratio_1958 1.38799e+00	Bratio_1958 8.53765e-01 1.05358e-01	-
Bratio_1959 1.47416e+00	Bratio_1959 8.46781e-01 1.03936e-01	-
Bratio_1960 1.54960e+00	Bratio_1960 8.41035e-01 1.02585e-01	-
Bratio_1961 1.62165e+00	Bratio_1961 8.35682e-01 1.01328e-01	-
Bratio_1962 1.66668e+00	Bratio_1962 8.32986e-01 1.00208e-01	-
Bratio_1963 1.69769e+00	Bratio_1963 8.31554e-01 9.92203e-02	-
Bratio_1964 1.68582e+00	Bratio_1964 8.34142e-01 9.83844e-02	-
Bratio_1965 1.67121e+00	Bratio_1965 8.36870e-01 9.76116e-02	-
Bratio_1966 1.62135e+00	Bratio_1966 8.42917e-01 9.68843e-02	-
Bratio_1967 1.87503e+00	Bratio_1967 8.20860e-01 9.55397e-02	-
Bratio_1968	Bratio_1968 7.97529e-01 9.39785e-02	-

2.15444e+00			
Bratio_1969	Bratio_1969	7.94575e-01	9.26303e-02
2.21769e+00			-
Bratio_1970	Bratio_1970	8.05154e-01	9.14722e-02
2.13011e+00			-
Bratio_1971	Bratio_1971	8.17553e-01	9.03161e-02
2.02010e+00			-
Bratio_1972	Bratio_1972	8.29313e-01	8.90873e-02
1.91595e+00			-
Bratio_1973	Bratio_1973	8.49080e-01	8.82837e-02
1.70949e+00			-
Bratio_1974	Bratio_1974	8.78044e-01	8.81276e-02
1.38385e+00			-
Bratio_1975	Bratio_1975	9.23593e-01	8.90587e-02
8.57938e-01			-
Bratio_1976	Bratio_1976	9.83997e-01	9.12177e-02
1.75438e-01			-
Bratio_1977	Bratio_1977	1.04087e+00	9.32591e-02
01			4.38291e-
Bratio_1978	Bratio_1978	1.08068e+00	9.39187e-02
01			8.58994e-
Bratio_1979	Bratio_1979	1.08929e+00	9.22241e-02
01			9.68222e-
Bratio_1980	Bratio_1980	1.02707e+00	8.56314e-02
01			3.16098e-
Bratio_1981	Bratio_1981	8.78064e-01	7.32057e-02
1.66566e+00			-
Bratio_1982	Bratio_1982	7.01050e-01	5.96916e-02
5.00824e+00			-
Bratio_1983	Bratio_1983	5.55389e-01	4.96036e-02
8.96328e+00			-
Bratio_1984	Bratio_1984	5.28383e-01	4.73370e-02
9.96298e+00			-
Bratio_1985	Bratio_1985	5.20001e-01	4.66172e-02
1.02966e+01			-
Bratio_1986	Bratio_1986	5.24175e-01	4.68068e-02
1.01657e+01			-
Bratio_1987	Bratio_1987	5.28155e-01	4.71677e-02
1.00036e+01			-
Bratio_1988	Bratio_1988	5.05695e-01	4.64383e-02
1.06444e+01			-
Bratio_1989	Bratio_1989	4.91114e-01	4.63600e-02
1.09768e+01			-

Bratio_1990	Bratio_1990	4.55582e-01	4.60491e-02	-
1.18225e+01				
Bratio_1991	Bratio_1991	4.20619e-01	4.39622e-02	-
1.31791e+01				
Bratio_1992	Bratio_1992	4.10952e-01	4.32186e-02	-
1.36295e+01				
Bratio_1993	Bratio_1993	4.01077e-01	4.23752e-02	-
1.41338e+01				
Bratio_1994	Bratio_1994	3.74341e-01	4.13701e-02	-
1.51235e+01				
Bratio_1995	Bratio_1995	3.63243e-01	4.21344e-02	-
1.51125e+01				
Bratio_1996	Bratio_1996	3.53251e-01	4.41909e-02	-
1.46353e+01				
Bratio_1997	Bratio_1997	3.52005e-01	4.77362e-02	-
1.35745e+01				
Bratio_1998	Bratio_1998	3.45896e-01	5.19280e-02	-
1.25963e+01				
Bratio_1999	Bratio_1999	3.45778e-01	5.62146e-02	-
1.16379e+01				
Bratio_2000	Bratio_2000	3.42591e-01	5.99322e-02	-
1.09692e+01				
Bratio_2001	Bratio_2001	3.40908e-01	6.29509e-02	-
1.04699e+01				
Bratio_2002	Bratio_2002	3.51126e-01	6.53152e-02	-
9.93451e+00				
Bratio_2003	Bratio_2003	3.74780e-01	6.76258e-02	-
9.24529e+00				
Bratio_2004	Bratio_2004	4.06558e-01	7.04435e-02	-
8.42437e+00				
Bratio_2005	Bratio_2005	4.42200e-01	7.36209e-02	-
7.57665e+00				
Bratio_2006	Bratio_2006	4.76351e-01	7.62757e-02	-
6.86522e+00				
Bratio_2007	Bratio_2007	5.07229e-01	7.80313e-02	-
6.31504e+00				
Bratio_2008	Bratio_2008	5.32201e-01	7.85340e-02	-
5.95664e+00				
Bratio_2009	Bratio_2009	5.56245e-01	7.86266e-02	-
5.64383e+00				
Bratio_2010	Bratio_2010	5.80167e-01	7.82985e-02	-
5.36196e+00				
Bratio_2011	Bratio_2011	6.08821e-01	7.82842e-02	-

4.99691e+00		
Bratio_2012	Bratio_2012 6.39031e-01 7.81272e-02	-
4.62028e+00		
Bratio_2013	Bratio_2013 6.77365e-01 7.85966e-02	-
4.10495e+00		
Bratio_2014	Bratio_2014 7.19845e-01 7.92412e-02	-
3.53547e+00		
Bratio_2015	Bratio_2015 7.59810e-01 7.93856e-02	-
3.02561e+00		
Bratio_2016	Bratio_2016 7.92480e-01 7.85220e-02	-
2.64282e+00		
Bratio_2017	Bratio_2017 8.12009e-01 7.62804e-02	-
2.46448e+00		
Bratio_2018	Bratio_2018 7.87954e-01 7.45033e-02	-
2.84614e+00		
Bratio_2019	Bratio_2019 7.39779e-01 7.47845e-02	-
3.47961e+00		
Bratio_2020	Bratio_2020 7.03324e-01 7.61804e-02	-
3.89439e+00		
Bratio_2021	Bratio_2021 6.81260e-01 7.91179e-02	-
4.02867e+00		
Bratio_2022	Bratio_2022 6.51158e-01 8.44896e-02	-
4.12881e+00		
Bratio_2023	Bratio_2023 6.08366e-01 9.08166e-02	-
4.31237e+00		
Bratio_2024	Bratio_2024 5.55992e-01 9.60409e-02	-
4.62311e+00		
Bratio_2025	Bratio_2025 4.96774e-01 9.94438e-02	-
5.06041e+00		
Bratio_2026	Bratio_2026 4.26222e-01 1.02357e-01	-
5.60565e+00		
Bratio_2027	Bratio_2027 3.64868e-01 1.04478e-01	-
6.07910e+00		
Bratio_2028	Bratio_2028 3.53352e-01 9.20464e-02	-
7.02524e+00		
Bratio_2029	Bratio_2029 3.51060e-01 8.24112e-02	-
7.87442e+00		
Bratio_2030	Bratio_2030 3.55143e-01 7.56557e-02	-
8.52357e+00		
Bratio_2031	Bratio_2031 3.62827e-01 7.17325e-02	-
8.88264e+00		
Bratio_2032	Bratio_2032 3.71888e-01 7.07532e-02	-
8.87751e+00		

Bratio_2033 8.56705e+00	Bratio_2033 3.80733e-01 7.22848e-02	-
Bratio_2034 8.11257e+00	Bratio_2034 3.88654e-01 7.53579e-02	-
Bratio_2035 7.64102e+00	Bratio_2035 3.95604e-01 7.90989e-02	-
Bratio_2036 7.20933e+00	Bratio_2036 4.01720e-01 8.29869e-02	-
SSB_unfished	SSB_unfished 8.17335e+04 7.22592e+03	NA
Totbio_unfished	Totbio_unfished 1.57283e+05 1.41425e+04	NA
SmryBio_unfished	SmryBio_unfished 1.51584e+05 1.34472e+04	NA
Recr_unfished	Recr_unfished 3.41029e+04 5.74731e+03	NA
SSB_Btgt	SSB_Btgt 3.26934e+04 2.89037e+03	NA
SPR_Btgt	SPR_Btgt 4.58333e-01 7.07303e-18	NA
annF_Btgt	annF_Btgt 8.75296e-02 4.32807e-03	NA
Dead_Catch_Btgt	Dead_Catch_Btgt 5.90211e+03 7.00270e+02	NA
SSB_SPR	SSB_SPR 3.64657e+04 3.22387e+03	NA
annF_SPR	annF_SPR 7.61335e-02 3.80371e-03	NA
Dead_Catch_SPR	Dead_Catch_SPR 5.62826e+03 6.65721e+02	NA
SSB_MSY	SSB_MSY 2.16668e+04 1.88801e+03	NA
SPR_MSY	SPR_MSY 3.36540e-01 1.60767e-03	NA
annF_MSY	annF_MSY 1.31509e-01 6.78851e-03	NA
Dead_Catch_MSY	Dead_Catch_MSY 6.31025e+03 7.56278e+02	NA
Ret_Catch_MSY	Ret_Catch_MSY 6.29147e+03 7.54037e+02	NA
B_MSY/SSB_unfished	B_MSY/SSB_unfished 2.65090e-01 1.78080e-03	NA
ForeCatch_2025	ForeCatch_2025 1.06686e+04 2.99174e-04	NA
ForeCatch_2026	ForeCatch_2026 9.82360e+03 8.85529e-04	NA
ForeCatch_2027	ForeCatch_2027 3.45573e+03 1.79537e+03	NA
ForeCatch_2028	ForeCatch_2028 3.44496e+03 1.60076e+03	NA
ForeCatch_2029	ForeCatch_2029 3.61838e+03 1.50526e+03	NA
ForeCatch_2030	ForeCatch_2030 3.87612e+03 1.45348e+03	NA
ForeCatch_2031	ForeCatch_2031 4.15228e+03 1.38604e+03	NA
ForeCatch_2032	ForeCatch_2032 4.40352e+03 1.29854e+03	NA
ForeCatch_2033	ForeCatch_2033 4.59801e+03 1.21212e+03	NA
ForeCatch_2034	ForeCatch_2034 4.73289e+03 1.13835e+03	NA
ForeCatch_2035	ForeCatch_2035 4.83358e+03 1.07948e+03	NA
ForeCatch_2036	ForeCatch_2036 4.90151e+03 8.75662e+02	NA
OFLCatch_2025	OFLCatch_2025 5.66709e+03 1.80582e+03	NA
OFLCatch_2026	OFLCatch_2026 4.56858e+03 1.67315e+03	NA
OFLCatch_2027	OFLCatch_2027 3.81851e+03 1.57579e+03	NA
OFLCatch_2028	OFLCatch_2028 3.87475e+03 1.40826e+03	NA
OFLCatch_2029	OFLCatch_2029 4.09795e+03 1.33035e+03	NA
OFLCatch_2030	OFLCatch_2030 4.38880e+03 1.29217e+03	NA

OFLCatch_2031	OFLCatch_2031	4.68820e+03	1.23669e+03	NA
OFLCatch_2032	OFLCatch_2032	4.94779e+03	1.16321e+03	NA
OFLCatch_2033	OFLCatch_2033	5.14510e+03	1.09669e+03	NA
OFLCatch_2034	OFLCatch_2034	5.28694e+03	1.04637e+03	NA
OFLCatch_2035	OFLCatch_2035	5.39061e+03	1.00853e+03	NA
OFLCatch_2036	OFLCatch_2036	5.47043e+03	9.77302e+02	NA
ForeCatchret_2025	ForeCatchret_2025	1.06373e+04	5.33005e+00	NA
ForeCatchret_2026	ForeCatchret_2026	9.79484e+03	4.89634e+00	NA
ForeCatchret_2027	ForeCatchret_2027	3.44709e+03	1.79083e+03	NA
ForeCatchret_2028	ForeCatchret_2028	3.43634e+03	1.59672e+03	NA
ForeCatchret_2029	ForeCatchret_2029	3.60926e+03	1.50145e+03	NA
ForeCatchret_2030	ForeCatchret_2030	3.86629e+03	1.44978e+03	NA
ForeCatchret_2031	ForeCatchret_2031	4.14168e+03	1.38249e+03	NA
ForeCatchret_2032	ForeCatchret_2032	4.39224e+03	1.29521e+03	NA
ForeCatchret_2033	ForeCatchret_2033	4.58621e+03	1.20902e+03	NA
ForeCatchret_2034	ForeCatchret_2034	4.72073e+03	1.13543e+03	NA
ForeCatchret_2035	ForeCatchret_2035	4.82115e+03	1.07672e+03	NA
ForeCatchret_2036	ForeCatchret_2036	4.88891e+03	8.73417e+02	NA
ln(SPB)_1916	ln(SPB)_1916	1.13095e+01	1.40181e-01	NA
ln(SPB)_1970	ln(SPB)_1970	1.10945e+01	1.18043e-01	NA
ln(SPB)_2024	ln(SPB)_2024	1.07242e+01	2.46823e-01	NA
CumNorm				
SSB_Virgin		NA		
SSB_Initial		NA		
SSB_1916		NA		
SSB_1917		NA		
SSB_1918		NA		
SSB_1919		NA		
SSB_1920		NA		
SSB_1921		NA		
SSB_1922		NA		
SSB_1923		NA		
SSB_1924		NA		
SSB_1925		NA		
SSB_1926		NA		
SSB_1927		NA		
SSB_1928		NA		
SSB_1929		NA		
SSB_1930		NA		
SSB_1931		NA		
SSB_1932		NA		
SSB_1933		NA		
SSB_1934		NA		

SSB_1935	NA
SSB_1936	NA
SSB_1937	NA
SSB_1938	NA
SSB_1939	NA
SSB_1940	NA
SSB_1941	NA
SSB_1942	NA
SSB_1943	NA
SSB_1944	NA
SSB_1945	NA
SSB_1946	NA
SSB_1947	NA
SSB_1948	NA
SSB_1949	NA
SSB_1950	NA
SSB_1951	NA
SSB_1952	NA
SSB_1953	NA
SSB_1954	NA
SSB_1955	NA
SSB_1956	NA
SSB_1957	NA
SSB_1958	NA
SSB_1959	NA
SSB_1960	NA
SSB_1961	NA
SSB_1962	NA
SSB_1963	NA
SSB_1964	NA
SSB_1965	NA
SSB_1966	NA
SSB_1967	NA
SSB_1968	NA
SSB_1969	NA
SSB_1970	NA
SSB_1971	NA
SSB_1972	NA
SSB_1973	NA
SSB_1974	NA
SSB_1975	NA
SSB_1976	NA
SSB_1977	NA

SSB_1978	NA
SSB_1979	NA
SSB_1980	NA
SSB_1981	NA
SSB_1982	NA
SSB_1983	NA
SSB_1984	NA
SSB_1985	NA
SSB_1986	NA
SSB_1987	NA
SSB_1988	NA
SSB_1989	NA
SSB_1990	NA
SSB_1991	NA
SSB_1992	NA
SSB_1993	NA
SSB_1994	NA
SSB_1995	NA
SSB_1996	NA
SSB_1997	NA
SSB_1998	NA
SSB_1999	NA
SSB_2000	NA
SSB_2001	NA
SSB_2002	NA
SSB_2003	NA
SSB_2004	NA
SSB_2005	NA
SSB_2006	NA
SSB_2007	NA
SSB_2008	NA
SSB_2009	NA
SSB_2010	NA
SSB_2011	NA
SSB_2012	NA
SSB_2013	NA
SSB_2014	NA
SSB_2015	NA
SSB_2016	NA
SSB_2017	NA
SSB_2018	NA
SSB_2019	NA
SSB_2020	NA

SSB_2021	NA
SSB_2022	NA
SSB_2023	NA
SSB_2024	NA
SSB_2025	NA
SSB_2026	NA
SSB_2027	NA
SSB_2028	NA
SSB_2029	NA
SSB_2030	NA
SSB_2031	NA
SSB_2032	NA
SSB_2033	NA
SSB_2034	NA
SSB_2035	NA
SSB_2036	NA
Recr_Virgin	NA
Recr_Initial	NA
Recr_1916	NA
Recr_1917	NA
Recr_1918	NA
Recr_1919	NA
Recr_1920	NA
Recr_1921	NA
Recr_1922	NA
Recr_1923	NA
Recr_1924	NA
Recr_1925	NA
Recr_1926	NA
Recr_1927	NA
Recr_1928	NA
Recr_1929	NA
Recr_1930	NA
Recr_1931	NA
Recr_1932	NA
Recr_1933	NA
Recr_1934	NA
Recr_1935	NA
Recr_1936	NA
Recr_1937	NA
Recr_1938	NA
Recr_1939	NA
Recr_1940	NA

Recr_1941	NA
Recr_1942	NA
Recr_1943	NA
Recr_1944	NA
Recr_1945	NA
Recr_1946	NA
Recr_1947	NA
Recr_1948	NA
Recr_1949	NA
Recr_1950	NA
Recr_1951	NA
Recr_1952	NA
Recr_1953	NA
Recr_1954	NA
Recr_1955	NA
Recr_1956	NA
Recr_1957	NA
Recr_1958	NA
Recr_1959	NA
Recr_1960	NA
Recr_1961	NA
Recr_1962	NA
Recr_1963	NA
Recr_1964	NA
Recr_1965	NA
Recr_1966	NA
Recr_1967	NA
Recr_1968	NA
Recr_1969	NA
Recr_1970	NA
Recr_1971	NA
Recr_1972	NA
Recr_1973	NA
Recr_1974	NA
Recr_1975	NA
Recr_1976	NA
Recr_1977	NA
Recr_1978	NA
Recr_1979	NA
Recr_1980	NA
Recr_1981	NA
Recr_1982	NA
Recr_1983	NA

Recr_1984	NA
Recr_1985	NA
Recr_1986	NA
Recr_1987	NA
Recr_1988	NA
Recr_1989	NA
Recr_1990	NA
Recr_1991	NA
Recr_1992	NA
Recr_1993	NA
Recr_1994	NA
Recr_1995	NA
Recr_1996	NA
Recr_1997	NA
Recr_1998	NA
Recr_1999	NA
Recr_2000	NA
Recr_2001	NA
Recr_2002	NA
Recr_2003	NA
Recr_2004	NA
Recr_2005	NA
Recr_2006	NA
Recr_2007	NA
Recr_2008	NA
Recr_2009	NA
Recr_2010	NA
Recr_2011	NA
Recr_2012	NA
Recr_2013	NA
Recr_2014	NA
Recr_2015	NA
Recr_2016	NA
Recr_2017	NA
Recr_2018	NA
Recr_2019	NA
Recr_2020	NA
Recr_2021	NA
Recr_2022	NA
Recr_2023	NA
Recr_2024	NA
Recr_2025	NA
Recr_2026	NA

Recr_2027	NA
Recr_2028	NA
Recr_2029	NA
Recr_2030	NA
Recr_2031	NA
Recr_2032	NA
Recr_2033	NA
Recr_2034	NA
Recr_2035	NA
Recr_2036	NA
SPRratio_1916	0.000000e+00
SPRratio_1917	0.000000e+00
SPRratio_1918	0.000000e+00
SPRratio_1919	0.000000e+00
SPRratio_1920	0.000000e+00
SPRratio_1921	0.000000e+00
SPRratio_1922	0.000000e+00
SPRratio_1923	0.000000e+00
SPRratio_1924	0.000000e+00
SPRratio_1925	0.000000e+00
SPRratio_1926	0.000000e+00
SPRratio_1927	0.000000e+00
SPRratio_1928	0.000000e+00
SPRratio_1929	0.000000e+00
SPRratio_1930	0.000000e+00
SPRratio_1931	0.000000e+00
SPRratio_1932	0.000000e+00
SPRratio_1933	0.000000e+00
SPRratio_1934	0.000000e+00
SPRratio_1935	0.000000e+00
SPRratio_1936	0.000000e+00
SPRratio_1937	0.000000e+00
SPRratio_1938	0.000000e+00
SPRratio_1939	0.000000e+00
SPRratio_1940	0.000000e+00
SPRratio_1941	0.000000e+00
SPRratio_1942	0.000000e+00
SPRratio_1943	0.000000e+00
SPRratio_1944	6.515260e-294
SPRratio_1945	6.039260e-97
SPRratio_1946	2.866540e-185
SPRratio_1947	0.000000e+00
SPRratio_1948	0.000000e+00

SPRratio_1949	0.000000e+00
SPRratio_1950	0.000000e+00
SPRratio_1951	0.000000e+00
SPRratio_1952	0.000000e+00
SPRratio_1953	0.000000e+00
SPRratio_1954	0.000000e+00
SPRratio_1955	0.000000e+00
SPRratio_1956	0.000000e+00
SPRratio_1957	0.000000e+00
SPRratio_1958	0.000000e+00
SPRratio_1959	0.000000e+00
SPRratio_1960	0.000000e+00
SPRratio_1961	0.000000e+00
SPRratio_1962	0.000000e+00
SPRratio_1963	0.000000e+00
SPRratio_1964	0.000000e+00
SPRratio_1965	0.000000e+00
SPRratio_1966	4.559400e-11
SPRratio_1967	1.098600e-07
SPRratio_1968	3.737410e-34
SPRratio_1969	3.223990e-294
SPRratio_1970	2.851800e-312
SPRratio_1971	7.043010e-205
SPRratio_1972	2.672895e-321
SPRratio_1973	5.165940e-222
SPRratio_1974	0.000000e+00
SPRratio_1975	0.000000e+00
SPRratio_1976	4.684250e-299
SPRratio_1977	0.000000e+00
SPRratio_1978	0.000000e+00
SPRratio_1979	4.874200e-05
SPRratio_1980	1.000000e+00
SPRratio_1981	1.000000e+00
SPRratio_1982	1.000000e+00
SPRratio_1983	1.000000e+00
SPRratio_1984	1.000000e+00
SPRratio_1985	9.999920e-01
SPRratio_1986	9.999930e-01
SPRratio_1987	1.000000e+00
SPRratio_1988	9.999990e-01
SPRratio_1989	1.000000e+00
SPRratio_1990	1.000000e+00
SPRratio_1991	9.998190e-01

SPRratio_1992	9.996710e-01
SPRratio_1993	1.000000e+00
SPRratio_1994	9.999420e-01
SPRratio_1995	9.999840e-01
SPRratio_1996	9.995150e-01
SPRratio_1997	9.994560e-01
SPRratio_1998	9.404290e-01
SPRratio_1999	8.234200e-01
SPRratio_2000	4.678520e-01
SPRratio_2001	1.360880e-03
SPRratio_2002	5.206180e-68
SPRratio_2003	0.000000e+00
SPRratio_2004	0.000000e+00
SPRratio_2005	0.000000e+00
SPRratio_2006	0.000000e+00
SPRratio_2007	0.000000e+00
SPRratio_2008	0.000000e+00
SPRratio_2009	0.000000e+00
SPRratio_2010	0.000000e+00
SPRratio_2011	0.000000e+00
SPRratio_2012	0.000000e+00
SPRratio_2013	0.000000e+00
SPRratio_2014	6.471870e-258
SPRratio_2015	1.351640e-222
SPRratio_2016	1.182160e-184
SPRratio_2017	3.251040e-03
SPRratio_2018	7.286490e-01
SPRratio_2019	6.705720e-01
SPRratio_2020	6.250620e-01
SPRratio_2021	9.392490e-01
SPRratio_2022	9.702960e-01
SPRratio_2023	9.405550e-01
SPRratio_2024	9.260550e-01
SPRratio_2025	9.885780e-01
SPRratio_2026	9.939240e-01
SPRratio_2027	3.362070e-01
SPRratio_2028	2.670740e-01
SPRratio_2029	2.244830e-01
SPRratio_2030	2.291240e-01
SPRratio_2031	2.934830e-01
SPRratio_2032	3.470410e-01
SPRratio_2033	3.760550e-01
SPRratio_2034	3.895540e-01

SPRratio_2035	3.983730e-01
SPRratio_2036	3.997730e-01
F_1916	0.000000e+00
F_1917	0.000000e+00
F_1918	0.000000e+00
F_1919	0.000000e+00
F_1920	0.000000e+00
F_1921	0.000000e+00
F_1922	0.000000e+00
F_1923	0.000000e+00
F_1924	0.000000e+00
F_1925	0.000000e+00
F_1926	0.000000e+00
F_1927	0.000000e+00
F_1928	0.000000e+00
F_1929	0.000000e+00
F_1930	0.000000e+00
F_1931	0.000000e+00
F_1932	0.000000e+00
F_1933	0.000000e+00
F_1934	0.000000e+00
F_1935	0.000000e+00
F_1936	0.000000e+00
F_1937	0.000000e+00
F_1938	0.000000e+00
F_1939	0.000000e+00
F_1940	0.000000e+00
F_1941	0.000000e+00
F_1942	0.000000e+00
F_1943	0.000000e+00
F_1944	0.000000e+00
F_1945	0.000000e+00
F_1946	0.000000e+00
F_1947	0.000000e+00
F_1948	0.000000e+00
F_1949	0.000000e+00
F_1950	0.000000e+00
F_1951	0.000000e+00
F_1952	0.000000e+00
F_1953	0.000000e+00
F_1954	0.000000e+00
F_1955	0.000000e+00
F_1956	0.000000e+00

F_1957	0.000000e+00
F_1958	0.000000e+00
F_1959	0.000000e+00
F_1960	0.000000e+00
F_1961	0.000000e+00
F_1962	0.000000e+00
F_1963	0.000000e+00
F_1964	0.000000e+00
F_1965	0.000000e+00
F_1966	0.000000e+00
F_1967	0.000000e+00
F_1968	0.000000e+00
F_1969	0.000000e+00
F_1970	0.000000e+00
F_1971	0.000000e+00
F_1972	0.000000e+00
F_1973	0.000000e+00
F_1974	0.000000e+00
F_1975	0.000000e+00
F_1976	0.000000e+00
F_1977	0.000000e+00
F_1978	0.000000e+00
F_1979	0.000000e+00
F_1980	0.000000e+00
F_1981	0.000000e+00
F_1982	0.000000e+00
F_1983	0.000000e+00
F_1984	0.000000e+00
F_1985	0.000000e+00
F_1986	0.000000e+00
F_1987	0.000000e+00
F_1988	0.000000e+00
F_1989	0.000000e+00
F_1990	0.000000e+00
F_1991	0.000000e+00
F_1992	0.000000e+00
F_1993	0.000000e+00
F_1994	0.000000e+00
F_1995	0.000000e+00
F_1996	0.000000e+00
F_1997	0.000000e+00
F_1998	0.000000e+00
F_1999	0.000000e+00

F_2000	0.000000e+00
F_2001	0.000000e+00
F_2002	0.000000e+00
F_2003	0.000000e+00
F_2004	0.000000e+00
F_2005	0.000000e+00
F_2006	0.000000e+00
F_2007	0.000000e+00
F_2008	0.000000e+00
F_2009	0.000000e+00
F_2010	0.000000e+00
F_2011	0.000000e+00
F_2012	0.000000e+00
F_2013	0.000000e+00
F_2014	0.000000e+00
F_2015	0.000000e+00
F_2016	0.000000e+00
F_2017	0.000000e+00
F_2018	0.000000e+00
F_2019	0.000000e+00
F_2020	0.000000e+00
F_2021	0.000000e+00
F_2022	9.796410e-295
F_2023	1.024670e-227
F_2024	1.639650e-177
F_2025	3.506650e-88
F_2026	4.138630e-59
F_2027	0.000000e+00
F_2028	0.000000e+00
F_2029	0.000000e+00
F_2030	0.000000e+00
F_2031	0.000000e+00
F_2032	0.000000e+00
F_2033	0.000000e+00
F_2034	0.000000e+00
F_2035	0.000000e+00
F_2036	0.000000e+00
Bratio_1917	4.909590e-01
Bratio_1918	4.873250e-01
Bratio_1919	4.834070e-01
Bratio_1920	4.806560e-01
Bratio_1921	4.779400e-01
Bratio_1922	4.756790e-01

Bratio_1923	4.736800e-01
Bratio_1924	4.714930e-01
Bratio_1925	4.698680e-01
Bratio_1926	4.678550e-01
Bratio_1927	4.649810e-01
Bratio_1928	4.623260e-01
Bratio_1929	4.592630e-01
Bratio_1930	4.560730e-01
Bratio_1931	4.521230e-01
Bratio_1932	4.482330e-01
Bratio_1933	4.440890e-01
Bratio_1934	4.399600e-01
Bratio_1935	4.353910e-01
Bratio_1936	4.302920e-01
Bratio_1937	4.245090e-01
Bratio_1938	4.184210e-01
Bratio_1939	4.122560e-01
Bratio_1940	4.056640e-01
Bratio_1941	3.974580e-01
Bratio_1942	3.883870e-01
Bratio_1943	3.782080e-01
Bratio_1944	3.598000e-01
Bratio_1945	3.311460e-01
Bratio_1946	2.898800e-01
Bratio_1947	2.602110e-01
Bratio_1948	2.415540e-01
Bratio_1949	2.259950e-01
Bratio_1950	2.119520e-01
Bratio_1951	1.965270e-01
Bratio_1952	1.785670e-01
Bratio_1953	1.608490e-01
Bratio_1954	1.444530e-01
Bratio_1955	1.287760e-01
Bratio_1956	1.135990e-01
Bratio_1957	9.799150e-02
Bratio_1958	8.257080e-02
Bratio_1959	7.021930e-02
Bratio_1960	6.061930e-02
Bratio_1961	5.243950e-02
Bratio_1962	4.778920e-02
Bratio_1963	4.478290e-02
Bratio_1964	4.591520e-02
Bratio_1965	4.733960e-02

Bratio_1966	5.247170e-02
Bratio_1967	3.039450e-02
Bratio_1968	1.560300e-02
Bratio_1969	1.328800e-02
Bratio_1970	1.658130e-02
Bratio_1971	2.168670e-02
Bratio_1972	2.768580e-02
Bratio_1973	4.368000e-02
Bratio_1974	8.320160e-02
Bratio_1975	1.954630e-01
Bratio_1976	4.303680e-01
Bratio_1977	6.694120e-01
Bratio_1978	8.048280e-01
Bratio_1979	8.335330e-01
Bratio_1980	6.240360e-01
Bratio_1981	4.789010e-02
Bratio_1982	2.750860e-07
Bratio_1983	1.598770e-19
Bratio_1984	1.128240e-23
Bratio_1985	3.727330e-25
Bratio_1986	1.439020e-24
Bratio_1987	7.496420e-24
Bratio_1988	9.478670e-27
Bratio_1989	2.531890e-28
Bratio_1990	1.537300e-32
Bratio_1991	6.012580e-40
Bratio_1992	1.391970e-42
Bratio_1993	1.227140e-45
Bratio_1994	5.955310e-52
Bratio_1995	7.032080e-52
Bratio_1996	8.750840e-49
Bratio_1997	2.951090e-42
Bratio_1998	1.144000e-36
Bratio_1999	1.359800e-31
Bratio_2000	2.753920e-28
Bratio_2001	6.067990e-26
Bratio_2002	1.501700e-23
Bratio_2003	1.191670e-20
Bratio_2004	1.835340e-17
Bratio_2005	1.788180e-14
Bratio_2006	3.339610e-12
Bratio_2007	1.356390e-10
Bratio_2008	1.291860e-09

Bratio_2009	8.338250e-09
Bratio_2010	4.125270e-08
Bratio_2011	2.917400e-07
Bratio_2012	1.918180e-06
Bratio_2013	2.023130e-05
Bratio_2014	2.035660e-04
Bratio_2015	1.240710e-03
Bratio_2016	4.110950e-03
Bratio_2017	6.860640e-03
Bratio_2018	2.212730e-03
Bratio_2019	2.511160e-04
Bratio_2020	4.924210e-05
Bratio_2021	2.806030e-05
Bratio_2022	1.824230e-05
Bratio_2023	8.081680e-06
Bratio_2024	1.892150e-06
Bratio_2025	2.095310e-07
Bratio_2026	1.040140e-08
Bratio_2027	6.065900e-10
Bratio_2028	1.075550e-12
Bratio_2029	1.728310e-15
Bratio_2030	7.832400e-18
Bratio_2031	3.310900e-19
Bratio_2032	3.466870e-19
Bratio_2033	5.376090e-18
Bratio_2034	2.504670e-16
Bratio_2035	1.087070e-14
Bratio_2036	2.831720e-13
SSB_unfished	NA
Totbio_unfished	NA
SmryBio_unfished	NA
Recr_unfished	NA
SSB_Btgt	NA
SPR_Btgt	NA
annF_Btgt	NA
Dead_Catch_Btgt	NA
SSB_SPR	NA
annF_SPR	NA
Dead_Catch_SPR	NA
SSB_MSY	NA
SPR_MSY	NA
annF_MSY	NA
Dead_Catch_MSY	NA

Ret_Catch_MSY	NA
B_MSY/SSB_unfished	NA
ForeCatch_2025	NA
ForeCatch_2026	NA
ForeCatch_2027	NA
ForeCatch_2028	NA
ForeCatch_2029	NA
ForeCatch_2030	NA
ForeCatch_2031	NA
ForeCatch_2032	NA
ForeCatch_2033	NA
ForeCatch_2034	NA
ForeCatch_2035	NA
ForeCatch_2036	NA
OFLCatch_2025	NA
OFLCatch_2026	NA
OFLCatch_2027	NA
OFLCatch_2028	NA
OFLCatch_2029	NA
OFLCatch_2030	NA
OFLCatch_2031	NA
OFLCatch_2032	NA
OFLCatch_2033	NA
OFLCatch_2034	NA
OFLCatch_2035	NA
OFLCatch_2036	NA
ForeCatchret_2025	NA
ForeCatchret_2026	NA
ForeCatchret_2027	NA
ForeCatchret_2028	NA
ForeCatchret_2029	NA
ForeCatchret_2030	NA
ForeCatchret_2031	NA
ForeCatchret_2032	NA
ForeCatchret_2033	NA
ForeCatchret_2034	NA
ForeCatchret_2035	NA
ForeCatchret_2036	NA
ln(SPB)_1916	NA
ln(SPB)_1970	NA
ln(SPB)_2024	NA

Table viii: Summary table of 12-year projections beginning in 2027 for alternate states of nature based on the axis of uncertainty (a combination of M, h). Columns range over low, mid, and high state of nature, and rows range over different assumptions of total catch levels (discards + retained). Catches in 2025 and 2026 are allocated using the percentage of landings for each fleet in 2019-2024.

			State of nature					
			Low		Base		High	
Management decision	Year	catch (mt)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)
Constant catch (9,000 K)	2027	9,000	17,024	19.5	34,918	42.7	38,407	46.6
	2028	9,007	13,360	15.3	30,977	37.9	34,351	41.7
	2029	9,068	10,428	12.0	27,812	34.0	31,280	38.0
	2030	9,067	7,962	9.1	25,315	31.0	29,056	35.3
	2031	7,419	5,757	6.6	23,321	28.5	27,478	33.4
	2032	6,404	4,389	5.0	21,690	26.5	26,373	32.0
	2033	5,494	3,396	3.9	20,279	24.8	25,570	31.0
	2034	4,662	2,631	3.0	18,981	23.2	24,949	30.3
	2035	3,901	2,019	2.3	17,736	21.7	24,442	29.7
	2036	3,187	1,509	1.7	16,511	20.2	24,011	29.1
ACLP*=0.25, sigma=0.50	2027	1,790	24,754	28.4	29,822	36.5	41,750	50.7
	2028	1,750	24,698	28.3	29,349	35.9	41,257	50.1
	2029	1,791	25,148	28.9	29,596	36.2	41,522	50.4
	2030	1,863	25,952	29.8	30,369	37.2	42,422	51.5
	2031	1,939	26,957	30.9	31,461	38.5	43,786	53.2
	2032	1,991	28,047	32.2	32,692	40.0	45,448	55.2
	2033	1,993	29,155	33.5	33,935	41.5	47,258	57.4
	2034	1,969	30,261	34.7	35,143	43.0	49,121	59.6
	2035	1,929	31,364	36.0	36,298	44.4	50,970	61.9
	2036	1,886	32,473	37.3	37,406	45.8	52,765	64.1

			State of nature					
			Low		Base		High	
Management decision	Year	catch (mt)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)
ACLp*=0.45, sigma=0.50	2027	3,231	24,754	28.4	34,918	42.7	41,750	50.7
	2028	3,204	23,939	27.5	33,924	41.5	40,507	49.2
	2029	3,351	23,652	27.1	33,609	41.1	40,062	48.6
	2030	3,574	23,687	27.2	33,792	41.3	40,237	48.8
	2031	3,808	23,860	27.4	34,288	42.0	40,838	49.6
	2032	4,020	24,056	27.6	34,944	42.8	41,693	50.6
	2033	4,180	24,202	27.8	35,642	43.6	42,657	51.8
	2034	4,279	24,280	27.9	36,319	44.4	43,638	53.0
	2035	4,350	24,303	27.9	36,957	45.2	44,593	54.1
	2036	4,392	24,286	27.9	37,550	45.9	45,497	55.2

Scientific uncertainty

Spawning biomass is estimated to be at 40,603 mt in 2025, with a sigma of 0.2753. OFL is estimated to be 5,667 mt in 2019 with a coefficient of variation of 0.311

Research and data needs

There are many areas of research that could be improved to benefit the understanding and assessment of widow rockfish:

- **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for widow rockfish. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life-history of widow rockfish may reduce that uncertainty.
- **Coastwide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for widow rockfish off of the west coast of the U.S. and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and

Alaska to determine if there are similarities with the U.S. West Coast observations would help to define the connectivity between widow rockfish north and south of the U.S.-Canada border.

- **Historical landings and discards:** Although progress has been done in reconstructing historical landings of rockfish on the U.S. West Coast, the historical landings and discards continue to be uncertain for widow rockfish and improvements would increase the certainty that fishing removals are applied appropriately. Because landings are assumed to be known exactly in the assessment model, uncertainty in the predictions does not include uncertainty in the landings. A thorough look at historical landings, species compositions, and discarding practices would reduce the potential uncertainty that is not entirely accounted for.




Figure viii: Equilibrium yield curve for the base case model and associated target and limit reference points. Values are based on most recent fishery selectivity and distribution with steepness fixed at 0.720. The fraction unfished is relative to unfished spawning biomass.

1 Introduction

This is an assessment of widow rockfish that inhabit the waters off California, Oregon, and Washington from the U.S.-Canadian border in the north to the U.S.-Mexico border in the south, and does not include Puget Sound.

Sebastodes entomelas (widow rockfish) is named after its black-lined gut cavity (*ento* meaning within and *melas* meaning black). It has been referred to as buda, beccafico (Italian bird), and viuva (widow) prior to the 1930s. More recently, the widow rockfish is also called brownie, belinda bass, brown bomber, and soft brown.

1.1 Distribution and Stock Structure

Widow rockfish inhabit water depths of 25–370 m from northern Baja California, Mexico to Southeastern Alaska, and are most abundant from British Columbia to Northern California. Although catches north of the U.S.-Canada border or south of the U.S.-Mexico border were not included in this assessment, it is possible that these populations contribute to the biomass of widow rockfish off of the U.S. West Coast through adult migration and/or larval dispersion.

This assessment is based on a single coastwide area model. There is little evidence of genetically separate stocks along the U.S. coast and past assessments have used a single area versus a two-area assessment model and results were found to be similar (He et al. 2011). There is some evidence of biological differences between areas. For example, widow rockfish collected off California tend to mature at a smaller length than widow rockfish collected off of Oregon (Barss and Echeverria 1987). This may be due to environmental or anthropogenic effects rather than genetic differences.

1.2 Life History

This section is not required for an update assessment; please refer to the most recent full assessment (Hicks and Wetzel 2015) for additional information.

1.3 Ecosystem Considerations

This section is not required for an update assessment; please refer to the most recent full assessment (Hicks and Wetzel 2015) for additional information.

1.4 Fishery description

This section is not required for an update assessment; please refer to the most recent full assessment (Hicks and Wetzel 2015) for additional information.

1.5 Management History

This section is not required for an update assessment; please refer to the most recent full assessment (Hicks and Wetzel 2015) for additional information.

1.6 Management performance

Total mortality estimates from the WCGOP and from the stock assessment may differ due to the use of different methods. Investigation into how these methods differ is beyond the scope of a benchmark assessment. Table 3 shows that recent landings have been below recommended catch levels. Landings are a considerable amount below the ACL, and it is unlikely that total mortality has exceeded the ACL in the last 10 years.

1.7 Fisheries off Canada and Alaska

This section is not required for an update assessment; please refer to the most recent full assessment (Hicks and Wetzel 2015) for additional information.

2 Data

Many sources of data were available for this assessment (Figure 30), including indices of abundance, landings, discards, and length and age observations from fishery-dependent and fishery-independent sources. Data used in this assessment are described below. No new data sources were considered in this update assessment.

2.1 Fishery-independent data

Data from three fishery-independent surveys were used in this assessment: 1) the SWFSC and NWFSC/PWCC Midwater Trawl Survey (hereafter, “juvenile survey”); 2) the Alaska Fisheries Science Center (AFSC)/NWFSC Triennial Shelf Trawl Survey (hereafter, “triennial survey”); and 3) the NWFSC West Coast Groundfish Bottom Trawl Survey (hereafter, “WCGBTS”). Depth and latitude strata used to analyze the catch-rates, length compositions, and age compositions are the same as in the 2019 assessment (Table 10 & Table 12).

2.1.1 Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey

The Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey (Triennial Survey) was first conducted by the Alaska Fisheries Science Center (AFSC) in 1977 and was conducted every three years, ending 2004. The survey’s design and sampling methods are most recently described in (Weinberg et al. 2002).

The time series suggests a possible slightly increasing trend in biomass from 1980–1983, although is relatively flat until the end of the period in 2001 and 2004 when the index declines significantly. The index and length compositions for this historical Triennial survey were unchanged from the previous assessment. Please refer to to Hicks and Wetzel (2015) for details on those data inputs.

2.1.2 Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey

The Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (WCG-BTS)) has been conducted annually, since 2003. It is based on a random-grid design; covering the coastal waters between depths of 55–1,280 m (Bradburn et al. 2011). No survey occurred in 2020 due to Coronavirus disease (COVID-19). Widow rockfish are not

commonly caught in the WCGBTS. Higher catch rates occur north of 40° N latitude and catches are rare south of 36° N latitude (Figure 1). Few large fish are found shallower than 100 m and few small fish are found in the deeper water of the slope. There is no clear trend in length with latitude other than smaller fish tend to occur south of approximately 36° N latitude, and there appears to be some very small fish found near 39° N latitude.

For this assessment, geostatistical models of biomass density were fit to survey data using the R package [Species Distribution Models with Template Model Builder \(sdmTMB\)](#) (Anderson et al. 2022). This is an updated approach compared to the 2015 assessment (non-spatial delta-GLMM) and the 2019 update assessment (VAST delta-lognormal model). The sdmTMB model used a 200 knot mesh of the survey area (Thorson et al. 2015). The prediction grid was truncated to only include available survey locations in depths between 55–500 m to limit extrapolating beyond the data and edge effects. Tweedie, delta-binomial, delta-gamma, and mixture distributions, which allow for extreme catch events, were investigated. The positive catch weight model includes survey pass ('first' for early season or 'second' for late season) and year. Vessel-year effects, which have traditionally been included in index standardization for this survey, were not included as the estimated variance for the random effect was close to zero. Vessel-year effects were more prominent when models did not include spatial effects and were included for each unique combination of vessel and year in the data to account for the random selection of commercial vessels used during sampling (Helser et al. 2004; Thorson and Ward 2014).

Results are only shown for the delta-gamma and delta-lognormal distributions, which reported the best diagnostics among the explored models (Figure 5). Both models converged (positive definite Hessian matrix) but predicted data from both models showed slightly right-heavy tails compared to null expectations, with the gamma model having stronger divergence. Spatiotemporal estimates of positive catch for the delta-lognormal distribution were then converted into annual indices using `sdmtmb::get_index()` function informed by the spatial bounds of the survey (Anderson et al. 2022).

Overall, the delta-lognormal index estimates is more comparable to the 2019 spatiotemporal VAST-based index than the delta-gamma index, and less influenced by extreme catch events, particularly in 2013 and 2016; for these reasons, in addition to better model performance observed above, the delta-lognormal sdmTMB-based index was used for the base model in this assessment. The delta-lognormal mean value (2262.824) was slightly lower than the means of the index values used in the 2015 assessment (2701.12) and the 2019 update assessment (3301.765). Comparisons of the different error structures, design-based estimate and the VAST index used in 2019 are in Figure 8.

Length, age, and conditional age-at-length compositions were processed using the `nwfscSurvey` package in R publicly available on GitHub (Wetzel et al. 2025). Length compositions were created by expanding to the tow and summing to give a strata specific composition (Table 7). Strata definition was retained from the benchmark assessment.

The strata compositions were combined to a coastwide composition using a design-based index of abundance from each strata. The design based index is constructed by taking the average catch per unit effort (CPUE) defined as catch per area swept across tows in each stratum and year. The sum of strata specific composition data was then calculated, weighting by the average CPUE per stratum multiplied by the area of each stratum. Age distributions were included in the model as conditional-age-at-length (CAAL) observations. The marginal age-compositions were also included, but only for easier viewing of strong cohorts. The CAAL data were not expanded and were binned according to length, age, sex, and year.

The input sample sizes for length and marginal age-composition data were calculated based on (Stewart and Hamel 2014). The input sample size of CAAL data was set at the number of fish at each length by sex and by year. Expanded length frequencies from this survey show intermittent years of small fish; the 2018–2024 period generally suggests most fish are around 40–45cm in length (Figure 9). Strong cohorts are not immediately apparent and it seems that ageing error may result in some variability between years. Conditional age-at-length proportions (Figure 11) show relatively consistent length-at-age with few outliers.

2.1.3 SWFSC and NWFSC/PWCC Midwater Trawl Survey

An updated coastwide pre-recruit index of abundance for 2001–2024 for widow rockfish was created using data from three midwater trawl surveys targeting young-of-the-year (YOY) rockfish (SWFSC and NWFSC/PWCC Midwater Trawl Survey (Juvenile Survey)), provided by Tanya Rogers (SWFSC, pers. comm.). All surveys used identical gear, enabling the construction of a consistent coastwide index spanning from 36°N to the U.S./Canada border since 2004. Sampling in 2020 was limited due to the COVID-19 pandemic and excluded from all models. In 2010 and 2012, coverage was incomplete, so these years excluded from the final model to align with the 2019 assessment. Following the 2015 and 2019 assessments, data from 2001–2003 were also excluded due to limited spatial coverage (36°30' to 38°20' N latitude).

The index was built using a spatial GLM with the sdmTMB package (Anderson et al. 2022), modeling 100-day standardized catch-per-tow as a function of year (fixed effect), Julian date (GAM smoother, $k = 4$), spatial random field, and spatiotemporal random effects. Models with Tweedie, delta-lognormal, and delta-gamma error structures were compared; DHARMA residuals and simulation-based diagnostics indicated the Tweedie model performed best. The index shows a strong increasing trend in juvenile abundance from 2017 to 2023, with a slight decline in 2024. Recent values remain high relative to the previous decade, and uncertainty estimates support the robustness of this trend.

2.2 Fishery-dependent data

2.2.1 Landings

Widow rockfish is primarily caught by bottom trawl and midwater trawl gears (Table 1). This species also commonly bycaught by fishery targeting Pacific hake. Minimal amounts are also taken by non-trawl gears such as hook-and-line and net (Table 9).

The definitions of fishing fleets have not been changed from those in the 2015 and 2019 assessments. Five fishing fleets were specified within the model: 1) a shorebased bottom trawl fleet with coastwide catches from 1916–2024, 2) a shorebased midwater trawl fleet with coastwide catches from 1979–2024, 3) a mostly midwater trawl fleet that targets Pacific Hake/Whiting (*Merluccius productus*) and includes a foreign and at-sea fleet with catches from 1975–2024, a domestic shorebased fleet that targeted Pacific Hake with catches from 1991–2024, and foreign vessels that targeted Pacific Hake and rockfish between 1966–1976, 4) a net fishery consisting of catches mostly from California from 1981–2024, and 5) a hook-and-line fishery (predominantly longline) with coastwide catches from 1916–2024. As in previous assessments, catches from Puget Sound and those from commercial shrimp trawls, commercial pots, and recreational fisheries were excluded (as these are generally minimal).

Landings from the 1916-2018 period were carried forward into this assessment with slight modifications to the midwater and bottom trawl catches from California. Because PacFIN appear to underestimate midwater trawl catches in California in 1979-1980 when midwater trawl fishery for widow rockfish developed (Edward Dick, Pers. Comms.) we adjusted midwater and bottom trawl catches from California in these years to reflect the ratio of California midwater to bottom trawl catches in 1981-1982.

Recent catches (2019-onward) were extracted from PacFIN for commercial shorebased data and NORPAC for at-sea hake fishery bycatch, and were otherwise appended onto the 1916-2018 landings and apportioned among fleets using the same criteria as those documented in the 2015 assessment Hicks and Wetzel (2015).

2.2.2 Fishery catch-per-unit-effort

Three fishery-dependent CPUE indices were included as in the last assessment (Adams et al. 2019). Indices were derived from 1) Oregon bottom trawl (1984-1999), 2) Pacific Whiting at-sea foreign (1977-1988), and 3) Pacific whiting at-sea domestic fleets (1983-1998). These were not updated; see Adams et al. (2019) for details.

2.2.3 Fishery length and age data

Biological data from commercial fisheries that caught widow rockfish were extracted from PacFIN on March 25, 2025 and from the NORPAC database. Lengths and age samples taken during port sampling in California, Oregon, and Washington were used to generate length and age compositions. The data were classified into bottom trawl, midwater trawl, hake trawl, net, and hook-and-line fleets using pacfintools package in R. For each fleet, the raw observations were expanded to the trip level, to account for differences in samples sizes relative catch weights among trips (first stage expansion). The expanded length observations were then further expanded to state level, to account for differences in sampling intensity of widow rockfish landings among states combined into a single fleet (second stage expansion).

Table 4 shows the number of landings sampled and Table 5 shows the number of lengths taken for each year, gear, and fleet from the three states. Table 6 shows these numbers for the at-sea fleet. Expanded length compositions for bottom trawl, midwater trawl, hake fisheries, net, and hook-and-line are shown in Figure 12 to Figure 16. Age compositions for the five fleets are shown in Figure 17 and Figure 21. Occasional cohorts appear to move through the population, indicating that widow rockfish population dynamics may be characterized by episodic recruitment events.

2.2.4 Discards

Data on discards on Widow Rockfish are available from three different sources. Historical sources included (Pikitch and Rogers 1989) and Enhanced Data Collected Project (EDCP, Sampson 2002). These historical sources were not reanalyzed for this update assessment, and discard amounts were not changed from the last assessment. Sex-specific length frequencies were also available from (Pikitch and Rogers 1989). Discard estimates are shown in Table 9 and range from 463 to 1,847 mt. Length compositions for discards show a wide range of sizes being discarded, with a peak around 40 cm (Figure 22).

The West Coast Groundfish Observer Program (WCGOP) provided information on recent discard of widow rockfish between 2002–2023. Since 2011, under trawl rationalization, 100% observer coverage is required for the limited entry trawl sectors, which resulted in a large increase in data and ability to determine discard behavior.

Discard length data for 2004–2023 was provided by WCGOP. In 2015 and 2019 assessments, these data were used to estimate retention curves for bottom trawl and hook-and-like fleets. WCGOP discard lengths from 2004–2017 are unchanged for the bottom trawl fleet, and new data are added for all consecutive years through 2023.

Major changes occurred only in the treatment of the hook-and-line discard data; discards from the hook-and-line fleet were added to the removals for this fleet. The hook-and-line removals of widow rockfish are extremely minimal (Table 1) and comprised only approximately 0.2% of the total removals over the last twenty years, with discard being a small fraction of that. Discard lengths for the hook & line fleet provided by WCGOP this year were corrected to omit nearshore fixed gear fleets, which were included with the hook-and-line fleet in the previous benchmark and update assessments erroneously. This change resulted in changes to the discard length distribution and years for which data was available. The hook-and-line removals comprised only approximately 0.2% of the total removals over the last twenty years (Table 1), with discard being a small fraction of that. The biological samples of the discard amount are also scarce, with input sample sizes not exceeding 6 and averaging around 3 per year. With this limited data, the model was unable to reliably estimate retention parameters and exhibited substantial sensitivity to even slight changes in discard amounts within the hook-and-line fleet. Therefore, in this assessment, we added the hook-and-line discard to hook-and-line landings.

Discard length composition data, became available this year (but was not available for previous assessments). These newly available data for midwater fleets (hake, rockfish) beginning in 2016, were not included in this updated assessment as this would require major changes to the model structure which are beyond the scope of this update assessment but will need to be considered in the next full assessment.

Figure 22 shows discard totals in metric tons for each year, the source of the discard data, and whether it was included in the assessment. Coefficients of variation (CV) are set to 5% for years/fleets with 100% observer coverage. For the non-catch share sectors, total discards by fleet were calculated by summarizing the observed discards (d) and observed retention (r) by fleet on a coastwide basis. Using the observed landings (R), the total discards were calculated as $D_{y,f} = \frac{d_{y,f}}{r_{y,f}} R_{y,f}$ where y and f indicate year and fleet, respectively. CV's for the WCGOP data were calculated by bootstrapping vessels within ports because the observer program randomly chooses vessels within ports to be observed. For the hook-and-line fleet, no estimates of observation error were available for the years 2011–2014, 2019 and 2021. A median CV for the hook-and-line fleet, calculated across years where CV's were available, was used for these years. Length compositions of the discards for the bottom trawl fleet can be seen in Figure 23.

2.2.5 Biological data

The approach to the estimation of all biological parameters was the same as in Adams et al. (2019).

2.2.5.1 Weight-length relationship

Weight-at-length data were updated for this assessment. Following the approach used in the 2015 benchmark, data used to estimate the length-weight relationship for widow rockfish were gathered from commercial catch sampling schemes (PACFIN, ASHOP) and fishery-independent surveys (Triennial and WCGBT Survey).

The following relationships between weight and length for females and males were used in the current assessment:

$$\text{Females: } \text{weight} = 1.7355 \times 10^{-5} \cdot \text{Length}^{2.9617}$$

$$\text{Males: } \text{weight} = 1.4824 \times 10^{-5} \cdot \text{Length}^{3.0047}$$

where weight is measured in kilograms and length in cm. These parameters were used in the assessment as fixed.

2.2.5.2 Maturity schedule

Maturity parameters in this update assessment were carried over from 2015 benchmark and 2019 update assessments. Please refer to Hicks and Wetzel (2015) for further details.

2.2.5.3 Fecundity

Fecundity parameters remain unchanged from previous assessment. Please refer to Hicks and Wetzel (2015) for further details.

2.2.5.4 Natural Mortality

In this assessment, natural mortality (M) is estimated for females and males, while using the Hamel and Cope (2022) prior. The prior on M has been updated to reflect the most recent guidance from Hamel and Cope (2022); the log-mean therefore remains unchanged while the log-SD has been set to 0.31. Using a maximum age of 54 the point estimate and median of the prior on M is 0.10.

2.2.5.5 Length-at-age

Growth parameters were fully estimated within the assessment model, for females and males separately following the same formulation in 2015 and 2019 assessment models.

2.2.5.6 Ageing bias and imprecision

Ageing error matrices were unchanged from the previous assessment. Please refer to Hicks and Wetzel (2015), Table 14, and Figure 29 for futher details.

2.3 Environmental and Ecosystem Data

This assessment did not use any environmental or ecosystem data related to the stock.

3 Assessment model

3.1 History of modeling approaches

This section is not required for an update assessment; please refer to the most recent full assessment (Adams et al. 2019) for additional information.

3.2 Responses to most recent past STAR Panel recommendations

This section is not required for update assessments.

3.3 Model Structure and Assumptions

3.3.1 Model Changes from the Last Assessment

The assessment followed the same model structure as the 2015 base assessment (Hicks and Wetzel 2015) and 2019 update assessment (Adams et al. 2019). The changes made to the previous assessment model include:

- **Adding most recent data from commercial fleets** (as described in Data section of this report).
- **Updating WCGBTS index using current methodology** (as described in Data section of this report).
- **Updating the prior for natural mortality** based on Hamel and Cope (2022).
- **Updating length-weight parameters estimated** while including most recent data.
- **Extending the main period for estimating recruitment deviations and updating recruitment bias adjustment parameters** based on Methot and Taylor (2011).
- **Adding a block to the retention curve for the midwater trawl fleet and a block to hake fleet selectivity to account for recent changes in fleet behavior.** Adding a block to the retention curve for midwater trawl allowed for recent increases in midwater trawl discards (from 2017 forward), and improved fit to discard amounts modeled in the assessment. Adding a block to hake fleet selectivity accounted for a change in fish mean length (2020-2024) due to shifts in the spatial distribution of the hake fleet and to improve fit to length compositions. Neither change had a discernible influence on the estimated SSB. See “Model Bridging” section for detail.

- **Adding hook-and-line discards to landings in the hook-and-line fleet.** Removals using hook-and-line gear (a fleet in the model) comprised only approximately 0.2% of the total removals over the last twenty years (Table 1), with discard being a small fraction of that. The change to the treatment of hook-and-line discard was necessary since discard lengths for the hook-and-line fleet provided by WCGOP this year were corrected to omit nearshore fixed gear fleets samples, which were erroneously included with the hook-and-line fleet in the previous benchmark and update assessments. This change resulted in changes to the discard length distribution and years for which data was available (Figure below). With decreased sample sizes of hook-and-line discard length data available, the model was unable to reliably estimate retention parameters and exhibited substantial sensitivity to even slight changes in discard amounts within the hook-and-line fleet. Therefore, in this assessment, we added the hook-and-line discard to hook-and-line landings. Correction of the hook-and-line discard lengths translated into lower estimated recruitment in the 2010s than that of in 2019 model (higher recruitment estimates in the 2019 assessment were previously informed by erroneous estimates of smaller fish, which were corrected in this assessment). Lower recruitment in turn caused a decrease in stock size in recent years (see “Model Bridging”).

3.3.1.1 Model Bridging

The exploration of models began by bridging from the 2019 update assessment to SS3 version 3.30.2, which produced no discernible difference. We then focused on data bridging. Updating the catch series did not have a substantial effect on the historical biomass, and estimated the stock biomass increasing from 2000 to 2020 before decreasing up to the current period. Updating the model discards had the most significant impact on the absolute stock biomass (SSB), with a small decrease in absolute SSB prior to 1980 and a large decrease in absolute SSB from the mid 2010s through the current year (Figure 31). Updating model indices had a similar effect, while updating the age and length composition data increased the absolute stock biomass in the current period. Updates to discards likewise had the largest effect on relative SSB (fraction of unfished biomass); changes in relative SSB owing to all other datasets were in general small.

None of the other model bridging steps had a substantial effect on the estimates of stock biomass (Figure 31). The model bridging changes included (1) updating the prior for natural mortality (M) to follow that recommended by Hamel and Cope (2022), (2) updating fixed parameters of the male and female length-weight curve by fitting to data from the NWFSC WCGBT, ASHOP, and the triennial survey outside the model, (3) re-fitting the bias adjustment ramp for recruitment deviates, (4) adding a block on retention in the midwater trawl fishery from 2011-2016 and allowing estimation of retention in the final years of the model, which was previously fixed at 0.99, (5) adding a block to hake fleet selectivity from 2020-2024 (5) re-fitting the model using the MLE

as initial values following a jittering analysis which revealed the previous MLE was a local minima. These changes collectively resulted in a very small decrease in relative SSB post-1990, and pairwise correlations among all shared parameters in these models were high (>0.999). Bridging from the previous prior on M to the Hamel and Cope (2022) prior is detailed in the “priors” section.

To estimate discards in the model, time blocks for changes in selectivity and retention in discard data were used. Except for the aforementioned addition of blocks to the midwater trawl and hake fleets, the same structure for time blocks was used as in the previous assessment (Adams et al. 2019). The addition of midwater trawl and hake fleets block was necessitated by poor fits to midwater trawl discards and the length composition data from the hake fleet in from 2020 - 2024, respectively. The poor fit to hake fleet length composition data may be due to the high variability in the distribution of the hake / pacific whiting fleet between years, as the fleet moves to avoid bycatch (Holland and Martin 2019). Poor fit to midwater trawl discards is thought to be due to (1) the previous decision to fix late-year retention to 0.99 which represented an overestimate of the (very low) discards in 2011-2016, and (2) the reallocation of quota shares and subsequent increase in midwater trawl discards in 2017. These choices are described in more detail in the “fits to data” section.

3.3.2 Modeling Platform and Structure

For this update assessment, new versions of the previously used software were used. Stock Synthesis v3.30.13 was used to estimate the parameters in the 2019 model. R4SS, version 1.35.3, along with R version 3.5.3 were used to investigate and plot the 2019 model fits. For the update, Stock Synthesis v3.30.2 and R4SS, version 1.51.0, along with R version 4.5.0 were used. A summary of the data sources used in the model (details discussed above) is shown in Figure 40.

3.3.3 Model Overview

The model is a two-sex, age-structured model starting in 1916 with an accumulated age group at 40 years. Sex-specific growth and natural mortality were estimated. The lengths in the population were tracked by 1 cm intervals and the length data were binned into 2 cm intervals. A curvilinear ageing imprecision relationship was estimated and used to model ageing error. Fecundity was assumed to be proportional to body weight, thus spawning biomass was used as the measure of spawning output.

3.3.3.1 Model Fleets and Areas

The assessment uses a single-area model, consistent with previous assessments (Adams et al. 2019). Multiple fisheries encounter Widow Rockfish. The definitions of fishing fleets have not been changed from previous assessments (Adams et al. 2019).

3.3.4 Model Parameters

3.3.4.1 Estimated and Fixed Parameters

There were 214 estimated parameters in the base model. These included one parameter for recruitment (R_0), 10 sex-specific parameters for growth, two sex-specific natural mortality parameters, four parameters for extra variability on the survey indices (survey indices were fixed at zero), four parameters for the catchability of the hake series and the Triennial Shelf survey (the catchabilities for other surveys were calculated analytically), 49 parameters for selectivity, retention, and time blocking of the fleets, eight parameters for survey selectivity, 125 recruitment deviations, and 12 forecast recruitment deviations.

Fixed parameters in the model were as follows. Consistent with (Adams et al. 2019), steepness was fixed at 0.72 as was the mean of the current west coast rockfish steepness prior as described above. A sensitivity analysis and a likelihood profile were done for steepness. The standard deviation of recruitment deviates was fixed at 0.60. Maturity at age was fixed as described in (Adams et al. 2019). Length-weight parameters were fixed at estimates using length-weight observations from the NWFSC WCGBT survey (Figure 24 and Table 16).

The final base model assumed asymptotic selectivity (using the double-normal formulation in SS3) for each fishery, except for the midwater trawl fishery. The NWFSC and Triennial surveys both used spline curves. All selectivity curves were length based and the same shape as the 2019 update, with the exception of the NWFSC survey fleet. The NWFSC survey fleet selectivity was estimated to be slightly lower at lengths greater than 45 cm, compared to the selectivity estimated in 2019.

Time blocks were used for selectivity and retention parameters in the bottom trawl, midwater trawl, hake and hook-and-line fisheries as indicated in Table 15.

3.3.4.2 Priors

The prior on natural mortality (M) in the previous assessment was defined as a lognormal with mean on the log-scale of $\ln(5.4/A_{\max})$ and $SD(\ln(M)) = 0.4384343$ following analysis of the data in (Then et al. 2015) by Owen Hamel and the authors. In the current assessment the prior on M has been updated to reflect guidance from (Hamel and Cope

2022); the log-mean therefore remains unchanged while the log-SD has been set to 0.31. Using a maximum age of 54 the point estimate and median of the prior on M is 0.10.

The prior for steepness (h) assumes a beta distribution with parameters based on an update of the Dorn rockfish prior (commonly used in past West Coast rockfish assessments) conducted by Jim Thorson (pers. comm, NWFSC, NOAA) which was reviewed and endorsed by the SSC in 2015. During the stock assessment review, it was decided that the steepness prior should be developed without the past Widow Rockfish data to avoid using the same data to inform both the model prior and likelihood. Without Widow Rockfish, the prior used for the 2015 assessment was a beta distribution with mean = 0.798 and SD = 0.132 (corresponding beta parameters $\alpha = 6.59$ and $\beta = 1.667$). The 2019 update assessments used the current West Coast rockfish steepness prior with mean = 0.72 and SD = 0.16 ($\alpha = 4.95$, $\beta = 1.93$) which was approved for use in all rockfish stock assessments for 2019. This update assessment uses the same prior.

3.3.4.3 Recruitment deviations

The specification of when to estimate recruitment deviations is an assumption that likely affects model uncertainty. It was decided to estimate recruitment deviations from 1900–2024 to appropriately quantify uncertainty. The earliest length-composition data occur in 1976 and the earliest age data were in 1978. The most informed years for estimating recruitment deviations based on available composition data were from about the mid-1970s to about 2014. The period from 1900–1970 was fit using an early series with little or no bias adjustment, the main period of recruitment deviates was 1971–2020 with an upward and downward ramping of bias adjustment, and 2021 onward was fit using forecast recruitment deviates with little bias adjustment. (Methot and Taylor 2011) summarize the reasoning behind varying levels of bias adjustment based on the information available to estimate the deviates. The standard deviation of recruitment variability (sigma- R) was assumed to be 0.6 in the 2015 assessment, based on iteratively tuning to a value slightly less than the observed variability of recruitment deviations in the period 1975–2010 in 2015 (Figure 37).

Survey indices and total discards were fitted assuming a lognormal likelihood.

3.3.4.4 Sample weights

As previous full assessment used the McAllister & Ianelli method (Hicks and Wetzel 2015) and changing the weighting method is outside the TORs for an update, Francis weighting method is presented as a sensitivity.

The 2019 update assessment weighted composition data via the lambdas (Adams et al. 2019). In the current assessment, the method to weight the composition datasets in SS3

was to use variance adjustment factors as the weighting factors. The fleet and data-type (length or age) factor was entered as variance adjustments factors until the harmonic mean of the effective sample sizes matched the mean of the adjusted input sample sizes (McAllister and Ianelli 1997). Once the weighting was determined, lambda factors for all fleets with both marginal length and marginal age compositions were down-weighted by 0.5 to account for the potential double use of data since length and age are observed from the same fish.

3.3.5 Key Assumptions and Structural Choices

This section is not required for an update assessment; please refer to the most recent full assessment (Adams et al. 2019) for additional information.

3.4 Base Model Results

The base model parameter estimates along with approximate asymptotic standard errors are shown in Table 16 and the likelihood components are shown in Table 17. Estimates of key derived parameters and approximate 95% asymptotic confidence intervals are shown in Table 18.

3.4.1 Parameter Estimates

The estimates of natural mortality (0.124599 yr^{-1} and 0.136775 yr^{-1} for females and males, respectively) were higher than suggested by the medians of the prior distributions used in this assessment and the 2015 assessment. Fixing M at lower values than those estimates resulted in a pattern of reduced recruitment immediately before the fishery started. This suggests that the model is attempting to reduce the number of observations of older fish in the data. The estimates of M fall within the 95% confidence interval of the prior distribution (0.055– 0.184), and are shown in Figure 27.

Estimating M is difficult in stock assessments, and the estimated values may represent model misspecification instead of the actual life-history trait. However, in alternative models to the base, the estimates of M were rarely less than 0.14 yr^{-1} (Table 20). Uncertainty in the estimated M was also much less than the range of the prior (Figure 27). The assumption that appeared to have the largest effect on M was introducing dome-shaped selectivity in the midwater trawl fleet, which made M smaller (Table 20).

Selectivity curves were estimated for commercial and survey fleets and parameter estimates are provided in `?@tbl-param-ests-select`. The estimated selectivity, retention, and

keep (the product of selectivity and retention) curves for the trawl and hook-and-line fleets are shown in Figure 32. The selectivity curves showed a shift to larger fish in 2002 for the bottom trawl fishery and a shift to smaller fish in 2003 for the hook-and-line fishery. The bottom trawl shift is consistent with the introduction of the Rockfish Conservation Area (RCA) and gear restrictions (shoreward of the 18 RCA) that virtually eliminated fishing in shelf habitats where smaller Widow Rockfish would more likely be encountered. Around this same time, the fixed-gear RCA specifications began preventing fishing between 30 and 100 m.

The retention curves (Figure 47) showed a shift to retaining a lower percentage of fish since trip limits were introduced. The asymptote of the retention curve for the bottom trawl fishery sequentially decreased as more management restrictions were introduced to about 50% retention of larger fish in the 1998-2010 period. In recent years, bottom trawl retention is estimated at approximately 99%.

Midwater trawl and hook-and-line fisheries estimated an asymptote to retention just above 80% for the period 1983-2010. Both the selectivity for the hake fleet and the selectivity of the net fleet did not support dome-shaped selectivity (Figure 46). The estimated selectivity curves for the Triennial and NWFSC WCGBT surveys were similar to each other except that the triennial survey selected larger fish (Figure 33). The NWFSC WCGBT survey exhibited a more pronounced dome-shaped selectivity compared to the 2015 assessment.

In the 2015 assessment, additional survey variability (process error added directly to each year's input variability) for the triennial and NWFSC WCGBT surveys was not estimated in the model because the estimate was zero (Hicks and Wetzel 2015). To avoid bound issues in estimation of the Hessian, the authors fixed these at zero because the model results included reasonable estimates of variance. We retained the same modelling approach for the update assessment. The additional standard deviation added to the fishery-dependent indices was quite large, ranging from 0.16 for the bottom trawl index and 0.58 for the foreign at-sea hake fleet. The additional variability on the juvenile survey was the highest, at 1.69, giving the index very little weight in the model.

The estimates of maximum size for both females and males (Table 16) were not unexpected given the data in Figure 28. Estimates of k were slightly different in the model, but that is expected when accounting for selectivity. Estimated growth curves are shown in (Figure 35).

Estimates of recruitment suggest that the Widow Rockfish population is characterized by variable recruitment with occasional strong recruitment events and periods of low recruitment (Figure 36, Table 18). There is little information regarding recruitment prior to 1965.

3.4.2 Fits to the Data

Fits to data are discussed for survey abundance indices, discard data (biomass and length compositions), length composition data for the fisheries and surveys, marginal age compositions for the fisheries, and conditional age-at-length observations for the NWFSC WCGBT survey.

Survey indices and total discards were fitted assuming a lognormal likelihood. The five indices of abundance (three survey series, and two fishery indices of abundance) are shown in Figure 38. The Triennial Shelf Survey treatment was consistent with (Adams et al. 2019). Extra standard error was estimated for all of the series except for the two survey series (Table 16). None of the series showed patterns in residuals, and with the large amount of error, none of the series showed serious lack of fit. The recent NWFSC WCGBT survey showed a general increase from 2003 - 2015 followed by a general decrease from 2016 - 2024, which was also estimated in the base model (Figure 38, middle right). Low (2015, 2021, 2022) and high (2016) estimates of abundance did not fit well in the model.

The 2019 model applied four time blocks to midwater trawl retention, with constant fixed discard rates of 1% in both early years (1916-1981) and late years (2011 onward). With the addition of new discard data since the 2017 reallocation, the use of a fixed discard rate from 2011 onward resulted in poor fits to the discard data. Therefore the current assessment utilizes an additional block on retention from 2011-2016, with the final block beginning in 2017; retention is estimated in both the 2011-2016 and 2017-2024 time blocks, though it remains fixed for the earliest time block (1916-1981) at the value used in the previous assessment. The first time block with discard data was 1983 to 2001. Predicted discards for all three years of the Pikitch data (1985–1987) were underfit, but within the confidence limits (Figure 39). EDCP data in 1997 and 1998 were underfit. The second time block was from 2002-2010, which contained only one observation in 2002 (and was fit exactly, as expected). Discard rates were overestimated in 2012-2013 and underestimated in 2015-2016, though they were generally well fit in in the 2017-2024 time blocks.

Hook-and-line discard biological data was recently revised by WCGOP. Previous assessments combined length compositioon data form the nearshore pot and net fisheries to the Hook-and-line fleet in the model. Nearshore pot and net gears samples are now removed, according to current best practices. This update to the data resulted in two challenges: i) reduced sample sizes for Hook-and-line length composition data and ii) length compositions which differed significantly from the 2019 and 2015 assessments. As a result, retention parameters for the hook-and-line fleet were poorly informed, and the decision was made to combine hook-and-line discard and retained catch. Biological samples for hook-and-line discards were then removed from the model, and discards were

not estimated for the hook-and-line fleet. Hook-and-line removals were approximately 0.2% of the total Widow Rockfish removals in the last two decades.

Fits to the length-composition data are displayed in two different ways: the Pearson residuals-at-length are shown for each year for all types of length compositions, and also compared across fleets. More detailed plots of fitted lines drawn over the plotted proportions at length are shown in Appendix A. Pearson residuals for the fisheries (Figure 40, Figure 41) do not show consistent patterns, but they do show that some fleets are not fitting some cohorts. Each fleet also shows that there are periods where older fish are underfit, and periods when older fish are overfit. With a peaked length frequency distribution, it is common for these patterns to appear given shifts in the expected distribution due to sampling error, and time-varying parameters that are assumed time-invariant. The net fishery observed some very large fish in the first two years of data, but did not observe those fish in later years. This pattern was not seen in any other fishery. There were also years where females showed positive residuals (filled circle, observed > expected) and males showed negative residuals (e.g., Figure 40, early years of bottom trawl and midwater trawl). It is uncertain if this pattern is related to growth, sexing error, or to sex-specific selectivity (e.g., when Widow Rockfish aggregate, sexes possibly may be aggregating separately). Overall, the fits to commercial fishery length compositions showed some patterns that the 2015 assessment deemed to require complicated modelling assumptions to alleviate. However, the residuals were mostly less than 2 in absolute value, especially for fleets that were well informed by data. The model fits the marginal length composition integrated among years for each fleet well (Figure 42).

The fits to bottom trawl discard length frequencies were generally good except in the years since trawl rationalisation began (2011). These recent years observed small fish, which the estimated selectivity of the trawl fleet did not allow for. There were no other years that showed small fish being caught by the trawl fleet. Attempting to explain these small fish with additional time blocks on selectivity and retention did not help because explaining the small fish in the discards worsened the fits to the landed and larger fish. Discards are extremely small in this time period, so it is unlikely that a misfit here will have a lot of effect on the model. Combining the discard length frequencies over years may not be appropriate for the bottom trawl fishery due to the likely changes in discarding practices, but Figure 40 shows the prediction of discarding smaller females than observed and a more peaked observed distribution of discarded males than predicted.

The Triennial Shelf and NWFSC WCGBT surveys length frequencies showed underfitting of older fish in some years and underfitting of younger fish in others (Figure 44). The combined length frequencies across years were bimodal with a valley around 37 cm, and the model showed an indication of a bimodal distribution but was unable to adequately capture both peaks (Figure 42). The nonparametric selectivity pattern helped to reduce this pattern, but selectivity may be even more complicated for the surveys.

Age data were fitted as marginal age compositions for the fishing fleets and as conditional

age-at-length for the NWFSC WCGBT survey, which were expanded by tow and then by strata. Raw observations of age-at-length, which assumes that within each length bin the observed ages are a random sample of fish, were not used as they are inconsistent with the length compositions which are expanded. Using expanded age-at-length ensures that as the length bin size is increased, it approaches the expanded marginal age composition. Pearson residuals for the commercial fleets are shown in Figure 40 and Figure 41. For the trawl fisheries in Figure 40, there are diagonal patterns that mostly correspond to cohorts ageing through the years. However, there are instances where the diagonal seems to shift, such as the filled circles of the midwater trawl fishery on the lower left of the plot (years 1981–1991). The patterns match the length compositions residuals in some cases. The bottom trawl fishery shows the largest residuals in the most recent years, which could indicate a change in selectivity. The net and hook-and-line fits to age compositions (Figure 41) showed larger residuals than the trawl fisheries. As with the fits to the length compositions, the net fishery showed the inability to match the large number of older fish observed in the early years. There appears to be a strong shift in residuals in 1988 when a lack of fit to potentially a cohort appears. The residuals were typically less than 2 for fits to the age data. However, the female age compositions occasionally produced some large residuals that were not consistently seen in the male age compositions. Aggregating across years shows that the fit to age comps was good for the trawl fleets and less so for the net and hook-and-line fleets, which had smaller sample sizes (Figure 45). The aggregated data also showed that the predictions were often unable to fit the peak in the data.

The observed and expected age-at-length are shown in Figure 46 for the twelve years of the NWFSC WCGBT survey observations. The fits generally match the observations with some misfit at larger lengths. The standard deviation of age-at-length was variable and often the expectation was higher than the observations at larger lengths. Plots with the residuals for individual observations showed reasonably good fits to the conditional age-at-length data from the NWFSC shelf/combo survey (Figure 49). Some outliers are apparent, with large residuals mostly at smaller lengths for a given age.

3.4.3 Population Trajectory

The predicted spawning biomass (in metric tons) is given in Table 19 and plotted in Figure 50. The predicted spawning biomass from the base model generally showed a slight decline over the time series until 1966 when the foreign fleet began. A short, but sharp decline occurred, followed by a steep increase due to strong recruitment. The spawning biomass declined rapidly with the developing domestic midwater fishery in the late 1970s and early 1980s. The stock continued to decline until 2000 when a combination of strong recruitment and low catches resulted in a quick increase from 2000 - 2018. The stock declined rapidly from 2019 until the end of the time series, due to a combination of increased catches and low recruitment.

The 2025 spawning biomass relative to unfished equilibrium spawning biomass is above the target of 40% of unfinished spawning biomass (49.7%), with a low of `depletion[which.min(depletion[["Deplete"]]), "Deplete"]%` in `depletion[which.min(depletion[["Deplete"]]), "Yr"]` (Figure 52). This suggests that Widow Rockfish was not overfished, as was inferred from previous assessments (Williams et al. 2000). Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in the estimated spawning biomass is high, especially in the early years. It should be noted that while the stock is currently above the 40% target, the target does fall within the approximate 95% confidence interval (Figure 52).

Recruitment deviations were estimated for the entire time series that was modeled (Figure 36 and discussed in Section 3.3.1) and provide a more realistic portrayal of uncertainty. There are very large, but uncertain, estimates of recruitment in 1970, 2008, 2016, and 1971 (in descending order of magnitude). Other large recruitment events (in descending order of magnitude) occurred in 1978, 1981, 2010, 1991, and 1997. The five lowest recruitments (in ascending order) occurred in 2012, 2019, 2020, 2011, and 1976. The 2008 and 2016 year classes were estimated as 2 of the 4 strongest year classes. The 2019 update assessment estimated the 2013 recruitment as the strongest year class over the duration of the fishery, however the current assessment does not support this. It may be worthwhile to investigate the periods of strong and weak year classes further to see if it is an artifact of the data, a consistent autocorrelation, or a result of the environment. The input bias adjustment ramp matched the estimated (Figure 37).

The stock-recruit curve resulting from a fixed value of steepness is shown in Figure 37 with estimated recruitments also shown. The stock is predicted to have never fallen to low enough levels that the steepness is obvious. However, the lowest levels of predicted spawning biomass showed some of the smallest recruitments and very few above average recruitments. Steepness was not estimated in this model, but sensitivities to alternative values of steepness are discussed below.

3.5 Model Diagnostics

3.5.1 Convergence

Due to it consistently hitting the lower bound when estimated, the decision was made to fix the log-ascending width of the hook-and-line fleet selectivity curve at -5. Model convergence was determined by examining the final gradient, checking that the Hessian was positive (semi-)definite, and initializing the minimizer from perturbed values around the maximum likelihood estimates (MLE) to determine if the model found a better minimum (“jittering”). Initial jittering analyses indicated that the model had converged

to a false (local) minimum, though the difference in log-likelihoods at the previous estimates and the new MLE was small (<0.5). After accepting the new MLE as the base model, jittering was repeated 100 times with a jitter coefficient of 0.10 and (using a convergence threshold of 0.01) a better minimum was not found. 2% of the jittered models achieved the minimum negative log-likelihood and `rpaste0(jitters$njitters_within_2, "%")` were within two likelihood units. Through the jittering done as explained above and likelihood profiles, we are confident that the base case as presented represents the best fit to the data given the assumptions made. There were no difficulties in inverting the Hessian to obtain the parameter variance-covariance matrix. Likelihood profile runs which fixed M, R₀ and h at more extreme values did not initially converge. This was addressed by using the parameter estimates from models with less extreme values as starting values in subsequent models. Convergence was defined as the lowest negative log-likelihood achieved with jittering where the Hessian matrix was invertible.

3.5.2 Parameter Uncertainty

Parameter estimates are shown in `?@tbl-bio-params`, Table 16 and Table 19 along with approximate asymptotic standard errors. The only parameters with an absolute value of correlation greater than 0.95 were the female and male natural mortality parameters, which is expected. Estimates of key derived quantities are given in Table 18 along with approximate 95% asymptotic confidence intervals. There is a reasonable amount of uncertainty in the estimates of biomass. The confidence interval of the 2025 estimate of depletion is 30.12%–69.17% and above the management target of 40% of the unfished spawning biomass.

3.5.3 Sensitivity Analyses

Sensitivity analysis was performed to determine the model behavior under different assumptions than those of the base case model. 8 sensitivity analyses were conducted to explore the potential differences in model structure and assumptions, including:

1. Fixed natural mortality at 0.1 for both sexes (2015 assessment prior)
2. Fixed natural mortality at 0.124 yr⁻¹ for females and 0.129 yr⁻¹ for males (2011 assessment prior)
3. Forcing asymptotic selectivity on the midwater trawl fleet
4. Fitting logistic curves for NWFSC WCGBT survey selectivities
5. Weighting the composition data using the Francis method
6. Updated Washington catch reconstruction
7. Inclusion of previously excluded shrimp trawl data
8. Exclusion of triennial survey data

Likelihood values and estimates of key parameters are shown in Table 20. Predicted spawning biomass trajectories, estimated recruitment deviations and comparisons of model estimates for 2025 are shown in Figure 57, Figure 58 and Figure 59. The estimates of current stock depletion ranged from 49.14%-153.59% across the sensitivity runs, with fixing natural mortality to 0.1 (2015 assessment prior) resulting in the lowest estimate and forcing asymptotic selectivity on the midwater trawl fleet resulting in the highest estimate. Generally, the trajectory of the spawning biomass was qualitatively similar across all tested models, e.g., peak around late 1970s and late 2010s, projected decrease in biomass in 2025 followed by some recovery into the 2030s; the quantitative magnitude of these trends did vary across cases.

Fixing M at values lower than the base case estimate resulted in decreases in estimated spawning biomass, while fixing steepness across the values tested resulted in similar or increased estimated spawning biomass. The relative spawning biomass in 2025 changed to 86.73% with an M of 0.124 yr^{-1} and 0.129 yr^{-1} for females and males, respectively (2011 assessment prior), and to 49.14% with an M of 0.1 yr^{-1} (2015 assessment prior).

Forcing asymptotic selectivity on the midwater fleet increased estimated biomass in 2025 by 153.59% compared to base model, while forcing logistic selectivity on the NWFSC WCGBT resulted in similar estimated spawning biomass to the base model. Including shrimp trawl data and updating WA catch reconstruction had almost no impact on the estimated spawning biomass. Excluding the triennial survey data lead to slight increases in estimated spawning biomass.

The alternative weighting using the Francis method generally increased the estimate of spawning biomass across the timeseries, but the estimated biomass for 2025 was similar between the Francis weighted model and the base model (117.66% compared to base model).

3.5.4 Retrospective Analysis

A 5-year retrospective analysis was conducted by running the model using data only through 2020, 2021, 2022, 2023, and 2024 progressively (Figure 60). The initial scale of the spawning population was basically unchanged for all of these retrospectives. Removing 4–5 years of data led to slightly lower estimates of fishing mortality (F) and slightly higher spawning biomass over the last 15 years. In contrast, removing only 1–2 years resulted in higher F and lower biomass estimates. Despite these minor differences, population trends from all retrospective runs were very close, and there were no consistent patterns as years were removed. No concerning patterns were observed in the retrospective analysis.

3.5.5 Likelihood Profiles and key parameters

Likelihood profiles were conducted for R₀, steepness (even though it was not estimated in the base case) and over male and female natural mortality values simultaneously. These likelihood profiles were conducted by fixing the parameter at specific values and removing the prior on the parameter being profiled. Without the original prior distribution the MLE estimates from the base case will likely be different than the MLE in the likelihood profile, but this displays what information the data have. There was some difficulty in achieving model convergence for many parameterizations in the likelihood profile. In some cases jittering was required.

For profiles of natural mortality, the negative log-likelihood was minimized at a value of 0.132 for males, and a value of 0.121 for females. Profiles for natural mortality for each sex are illustrated in (Figure 63). For steepness, the negative log-likelihood was minimized at a steepness of 0.825, but the 95% confidence interval extends over the entire range of possible steepness values. Profiles for steepness are illustrated in (Figure 62). For R₀, the total likelihood supported the estimated value, though there was variable support for each likelihood component across the range of R₀ evaluated. As R₀ increased, natural mortality also increased and the relative spawning biomass in 2024 was less depleted. Profiles for R₀ are illustrated in Figure 61.

4 Management

4.1 Reference Points

Reference points were calculated using the estimated selectivities and catch distribution among fleets in the most recent year of the model (2024). Sustainable total yields (landings plus discards) were 5,594 mt when using an SPR50% reference harvest rate and with a 95% confidence interval of 5,594 to 5,594 mt. The spawning biomass equivalent to 40% of the unfished spawning output (SB40%) was 32,592 mt. Catches between the late 1990's and 2016 were well below the point estimate of potential long-term yields calculated using an SPR50% reference point, and the population is estimated to have increased continuously throughout that time period. However, catches from 2017 through 2024 were above the point estimate of potential long-term yields using an SPR50% reference point, exceeding the upper bound of the confidence interval in all years since 2018, and exceeding the point estimate by on average 83%.

The predicted spawning biomass from the base model generally showed a slight decline until the late 1970s, followed by steep increase above unfished equilibrium biomass and reaching a peak in 1979. This was followed by a steep decrease up to the mid-1980s, and then a more gradual decrease through 2000 (Figure 52). Between 2001 and 2016, the spawning biomass increased continuously due to small catches, and several years of high recruitment (though with lower than average recruitment in other recent years). The spawning biomass relative to unfished equilibrium spawning biomass climbed above the target of 40% of unfished spawning biomass in the early 2000's. It is estimated to still be above the target, though the lower bound of the 95% confidence interval lies between the target and minimum stock size threshold (Figure 52). The fishing intensity (relative 1-SPR) exceeded the current estimates of the harvest rate limit (SPR50%) throughout the 1980s and early 1990s, and has again since 2018 (Figure 55). Exploitation rates on widow rockfish between 2001 and 2016 were predicted to be far below target levels. In recent years, the stock has experienced exploitation rates that have been above the target level while the biomass level has remained above the target level (Figure 56).

The equilibrium yield plot is shown in Figure 65, based on a steepness value fixed at 0.720. The predicted maximum sustainable yield under the assumptions of this assessment occurs near 25% of equilibrium unfished spawning biomass.

4.2 Unresolved problems and major uncertainties

This is a reconfiguration of a long line of stock assessments for widow rockfish on the U.S. West Coast and although scientifically credible advice is provided by synthesizing

many sources of data, there remain data and structural assumptions that contribute to uncertainty in the estimates. Major sources of uncertainty include landings, discards, natural mortality, and recruitment, which are discussed below.

Discards of widow rockfish are even more uncertain than landings, but because widow rockfish is a marketable species, historical discard rates were likely lower than less desirable or smaller species. In this assessment, we assumed that discarding was nearly negligible before management restrictions began in 1982. Once trip limits were introduced, discarding tended to be an all or none event, and detecting large, but rare, discard events with far less than 100% observer coverage has a low probability. From 2002 onward, the WCGOP has provided data on discards from vessels that were randomly selected for observer coverage, thus some uncertainty is present in the total amount discarded. The implementation of trawl rationalization in 2011 resulted in almost 100% observer coverage for the trawl fleet and very little incentive to discard widow rockfish, though the 2017 reallocation of quota shares is likely to have influenced discarding practices. The open access fixed-gear fleet is not monitored by the full observer coverage required under trawl rationalization and data from the 2015 assessment show that discarding of widow rockfish has occurred on fixed gear vessels in recent years (limited entry vessel fishing with fixed gear are subject to 100% observer coverage). Uncertainty in recent discards is greatly reduced because of observer coverage, but it is unknown what historical discarding may have been. The model assumes a discard rate of 1% pre-1982, which is arbitrary, but reasonable. Discard mortality is assumed to be 100%, which may overestimate actual mortality (Jarvis & Lowe, 2007), but given the low number of discards, is likely to have a minimal effect on assessment results.

There may also be uncertainty in the ability of bottom trawl surveys to be a reliable measure of widow abundance, which spend a significant portion of their time in mid-water (Wilkins 1986). Uncertainty in the widow rockfish NWFSC WCGBTS estimates was high, and inter-annual variability greater than might be expected for a long-lived rockfish species, so fits to this index were poor. Multiple surveys are used in the assessment, but further consideration of additional surveys is reasonable.

Widow rockfish is a relatively long-lived fish, and natural mortality is likely to be lower than many species of fish, such as gadoids. Ages above 50 years have been observed and it is expected that natural mortality could be less than 0.10 yr^{-1} (the median of the prior for natural mortality used in this assessment). However, even with length and age data available back to the late 1970s, natural mortality was estimated at 0.125 for females and 0.137 yr^{-1} for males, with a small amount of uncertainty (e.g., a 6.6% coefficient of variation for females). This assessment attempts to capture that uncertainty by estimating natural mortality (M) and integrating that uncertainty into the derived biomass estimates, as well as additional uncertainty by including levels outside of the predicted interval in a decision table.

Model sensitivities and profiles over M showed that current stock status was highly

sensitive to the assumption about natural mortality. Notably, the estimated natural mortality values for both sexes from this assessment are lower than those from the 2015 and 2019 assessments, for which natural mortality was estimated above 0.15 yr^{-1} and 0.14 yr^{-1} , respectively. The estimates of M varied slightly depending on the weight given to age and length data, or removing recent years of data, but M was always estimated above 0.12 yr^{-1} . The likelihood profile over natural mortality provides support for values up to or above 0.14 yr^{-1} , but with greater curvature than in the 2019 assessment, suggesting additional data has reduced the support for higher natural mortality values.

Steepness was fixed at 0.720 in the base model, but a likelihood profile showed that it would be estimated at a value less than that. Estimates of M increased with lower steepness, while unfished equilibrium spawning biomass increased and current spawning biomass decreased. Sustainable yields at the SPR50% reference harvest rate ranged from approximately 2641 to 5992 mt depending on the value of steepness.

4.3 Harvest Projections and Decision Tables

Two categories of parameters that greatly contributed to uncertainty in the results were natural mortality (an important estimated parameter) and steepness (not estimated in the model). A combination of these two factors was used as the axis of uncertainty to define low and high states of nature. This differed from the 109 assessment which included a third factor, 2013 recruitment strength. The 12.5% and 87.5% quantiles for female and male natural mortality (independently) were chosen as low and high values (0.115 yr^{-1} and 0.134 yr^{-1} for females; 0.127 yr^{-1} and 0.146 yr^{-1} for males). Steepness is probably the most important factor since it was fixed in the base model and is not incorporated in the estimation uncertainty. The 12.5% and 87.5% quantiles from the steepness prior (without widow rockfish data) were used to define the low and high values of steepness (0.536 and 0.904). The low combination of these three factors defined the low state of nature and the high combination of these three factors defined the high state of nature. The predictions of spawning biomass in 2025 from the low and high states of nature are close to the 12.5% and 87.5% lognormal quantiles from the base model.

A twelve year projection of the base model with catches equal to the current ACL in 2025 and 2026 (10,669 mt and 9824 mt, respectively) and catches of 9,000 mt for all later years and a catch allocation equal to the percentages for each fleet in 2024 predicts spawning biomass will decrease over the projection period for all states of nature (Table 23, Figure 66).

Projections with catches based on the predicted annual catch limit (ACL) using the SPR rate of 50%, the 40:10 control rule, and a 0.45 P* adjustment using a sigma of 0.50 from 2027 onward suggest that the spawning biomass will decrease over the projection period

for all states of nature. Predicted ACL catches range from 3,321 mt in 2027 to 4,392 mt in 2036 Table 23, Figure 66).

Projections with catches based on the predicted annual catch limit (ACL) using the SPR rate of 50%, the 40:10 control rule, and a 0.25 P* adjustment using a sigma of 0.50 from 2021 onward suggest that the spawning biomass will decrease over the projection period for all states of nature. Predicted ACL catches range from 1,790 mt in 2027 to 1,886 mt in 2036 Table 23, Figure 66).

This assessment synthesizes many sources of data and estimates recruitment variability, thus it is classified as a Category 1 stock assessment. Therefore, the sigma for P* to determine the catch reduction to account for scientific uncertainty is 0.50.

4.4 Evaluation of Scientific Uncertainty

Spawning biomass is estimated to be at 40,603 mt in 2025, with a sigma of 0.2753. OFL is estimated to be 5,667.1 mt in 2019 with a coefficient of variation of 0.3187.

4.5 Regional management considerations

Widow rockfish have shown latitudinal differences in life-history parameters, which has led past assessment authors to pursue a two-area model. Modelling a stock with two areas is difficult because it requires many assumptions about recruitment distribution, movement, and connectivity, while also splitting data into two areas that reduces sample sizes when compared to a coastwide model. The upside is that it can result in a better model that more accurately predicts regional status. This assessment is a coastwide model because not enough is known about the assumptions that would have to be made for a two-area model.

It is still important to consider regional differences when making management decisions. Following recent cohorts through time with survey data showed that older fish showed up in the north after younger fish were observed in the south (Figure 1). This may indicate connectivity between the north and the south and that this is truly one stock. However, more investigation is needed.

Widow rockfish are managed on a coastwide basis and observed more often in the NWFSC WCGBT bottom trawl survey north of latitude 40° 10' N. Bottom trawl catches in California have historically been as large as in Oregon and larger than in Washington, but recently catches in California have been small. Rockfish Conservation Areas (RCAs) cover a significant proportion of widow rockfish habitat, but a midwater trawl fishery is

beginning to re-develop that can fish in these areas. Future assessments and management of widow rockfish may want to monitor where catches are being taken to make sure that specific areas are not being overexploited. In addition, research on the connectivity along the coast as well as regional differences would help to inform the potential for overfishing specific areas.

4.6 Research and Data Needs

There are many areas of research that could be improved to benefit the understanding and assessment of widow rockfish. Below, we specifically identify five topics that we believe are most important (order does not indicate importance).

- **Historical landings and discards:** The historical landings and discards are uncertain for widow rockfish and improvements would increase the certainty that fishing removals are applied appropriately. Because landings are assumed to be known exactly in the assessment model, uncertainty in the predictions does not include uncertainty in the landings. A thorough look at historical landings, species compositions, and discarding practices would reduce the potential uncertainty that is not entirely accounted for.
- **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for widow rockfish. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life-history of widow rockfish may reduce that uncertainty.
- **Basin-wide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for widow rockfish off of the west coast of the U.S. and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and Alaska to determine if there are similarities with the U.S. West Coast observations would help to define the connectivity between widow rockfish north and south of the U.S.-Canada border.

4.7 Acknowledgements

Thank you to the University of Washington Stock Assessment course instructors Vlada Gertseva, Melissa Haltuch, Kiva Oken and Ian Taylor for their guidance in preparing our model. We want to acknowledge Tanya Rodgers (juvenile survey data), Ian Taylor and Chantel Wetzel (NWFSC survey data), XXX (ASHOP data), XXX (PACFIN data) and XXX (catch reconstruction and composition bridging).

5 Tables

5.1 Data

5.1.1 Fishery-dependent data

Table 1: Landings for bottom trawl, midwater trawl, net, and hook-and-line (mt) fisheries from Washington, Oregon, and California.

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1916	6.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	71.8	0.3	0.0
1917	9.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	111.9	0.3	0.0
1918	11.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	128.5	0.3	0.0
1919	7.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	88.6	0.3	0.0
1920	8.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	90.7	0.4	0.0
1921	6.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	75.1	0.4	0.0
1922	5.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	65.2	0.4	0.0
1923	6.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	71.7	0.4	0.0
1924	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	46.2	0.4	0.0
1925	3.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	58.7	0.4	0.0
1926	8.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	85.5	0.4	0.0
1927	11.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	66.4	0.5	0.0
1928	16.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	72.0	0.8	0.0
1929	23.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	62.1	1.3	0.0
1930	20.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	90.4	1.2	0.0
1931	20.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	78.6	0.9	0.0
1932	21.7	0.4	0.0	0.0	0.0	0.0	0.0	0.0	77.7	0.3	0.0
1933	34.3	0.2	0.0	0.0	0.0	0.0	0.0	0.0	50.9	0.5	0.0
1934	30.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	59.7	0.5	0.0
1935	28.9	0.2	0.7	0.0	0.0	0.0	0.0	0.0	67.9	0.5	0.0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1936	23.4	0.7	1.1	0.0	0.0	0.0	0.0	0.0	84.3	1.2	0.0
1937	33.6	1.3	0.9	0.0	0.0	0.0	0.0	0.0	66.3	1.3	0.0
1938	32.2	0.0	1.1	0.0	0.0	0.0	0.0	0.0	49.6	1.0	0.0
1939	38.8	1.9	1.0	0.0	0.0	0.0	0.0	0.0	34.2	0.7	0.0
1940	30.6	43.7	1.0	0.0	0.0	0.0	0.0	0.0	43.9	1.5	0.0
1941	24.8	67.3	1.4	0.0	0.0	0.0	0.0	0.0	34.1	1.9	0.0
1942	5.4	126.1	1.8	0.0	0.0	0.0	0.0	0.0	10.2	3.1	0.0
1943	28.3	439.2	1.2	0.0	0.0	0.0	0.0	0.0	18.0	3.9	0.0
1944	148.6	770.7	2.0	0.0	0.0	0.0	0.0	0.0	38.0	1.4	0.0
1945	353.4	1,196.6	3.4	0.0	0.0	0.0	0.0	0.0	66.8	1.1	0.0
1946	353.2	735.0	0.8	0.0	0.0	0.0	0.0	0.0	69.7	1.3	0.0
1947	98.1	452.8	0.2	0.0	0.0	0.0	0.0	0.0	91.3	0.7	0.0
1948	139.4	297.3	0.1	0.0	0.0	0.0	0.0	0.0	39.6	1.2	0.0
1949	75.1	254.7	0.0	0.0	0.0	0.0	0.0	0.0	43.9	0.6	0.0
1950	70.9	286.8	1.8	0.0	0.0	0.0	0.0	0.0	63.4	0.8	0.0
1951	249.4	252.9	2.0	0.0	0.0	0.0	0.0	0.0	49.1	0.6	0.0
1952	236.6	264.2	0.2	0.0	0.0	0.0	0.0	0.0	39.9	0.6	0.0
1953	242.6	211.5	1.2	0.0	0.0	0.0	0.0	0.0	13.7	0.3	0.0
1954	155.8	267.3	3.1	0.0	0.0	0.0	0.0	0.0	21.3	0.4	0.0
1955	166.3	277.5	2.5	0.0	0.0	0.0	0.0	0.0	18.2	0.4	0.0
1956	196.8	361.3	0.7	0.0	0.0	0.0	0.0	0.0	41.8	0.3	0.0
1957	233.1	489.5	0.1	0.0	0.0	0.0	0.0	0.0	37.4	0.6	0.0
1958	284.3	380.4	0.2	0.0	0.0	0.0	0.0	0.0	36.6	0.1	0.0
1959	229.9	412.8	0.1	0.0	0.0	0.0	0.0	0.0	28.6	0.2	0.0
1960	180.0	608.6	0.2	0.0	0.0	0.0	0.0	0.0	21.9	0.2	0.0
1961	118.4	543.1	0.2	0.0	0.0	0.0	0.0	0.0	15.0	0.5	0.0
1962	115.9	623.8	2.0	0.0	0.0	0.0	0.0	0.0	15.4	0.4	0.0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1963	221.2	190.2	2.1	0.0	0.0	0.0	0.0	0.0	19.6	0.4	0.0
1964	104.1	480.9	3.2	0.0	0.0	0.0	0.0	0.0	13.0	0.1	0.0
1965	155.9	80.6	2.2	0.0	0.0	0.0	0.0	0.0	20.2	0.6	0.0
1966	123.0	455.8	0.6	0.0	0.0	0.0	0.0	0.0	37.4	0.4	0.0
1967	141.9	743.9	0.6	0.0	0.0	0.0	0.0	0.0	31.9	1.1	0.0
1968	155.0	240.6	16.7	0.0	0.0	0.0	0.0	0.0	19.0	1.0	0.0
1969	223.5	229.3	16.7	0.0	0.0	0.0	0.0	0.0	17.6	2.3	0.0
1970	257.3	27.7	3.0	0.0	0.0	0.0	0.0	0.0	9.0	0.9	0.0
1971	316.2	50.6	11.7	0.0	0.0	0.0	0.0	0.0	10.2	1.8	0.0
1972	411.9	51.8	14.1	0.0	0.0	0.0	0.0	0.0	17.8	2.3	0.0
1973	428.1	20.9	32.4	0.0	0.0	0.0	0.0	0.0	15.8	2.5	0.0
1974	426.4	7.3	6.5	0.0	0.0	0.0	0.0	0.0	41.3	3.1	0.0
1975	429.9	9.0	12.0	0.0	0.0	0.0	0.0	0.0	28.4	1.6	0.0
1976	467.3	56.0	36.2	0.0	0.0	0.0	0.0	0.0	39.5	2.2	0.0
1977	459.0	340.0	125.8	0.0	0.0	0.0	0.0	0.0	38.1	2.6	0.0
1978	538.9	340.1	336.7	0.0	0.0	0.0	0.0	0.0	157.4	3.8	0.0
1979	916.9	519.4	305.0	1,398.5	3,746.0	2,199.8	0.0	0.0	97.1	6.4	0.0
1980	2,109.2	410.8	338.4	3,217.2	8,460.7	6,969.4	0.0	3.4	55.9	3.7	0.0
1981	2,660.2	1,527.1	681.2	2,627.4	13,861.9	6,183.5	15.5	3.2	67.5	4.0	0.0
1982	3,656.7	782.8	522.0	7,008.1	8,184.4	5,458.0	38.1	37.1	180.6	5.9	0.0
1983	3,667.1	1,403.6	1,554.6	205.1	1,495.6	1,656.5	280.0	14.5	23.5	10.2	0.0
1984	1,434.6	1,428.5	381.8	1,378.6	3,982.8	1,064.6	324.8	26.6	22.8	3.8	0.0
1985	1,363.0	895.1	317.6	1,281.6	3,423.4	1,214.6	585.8	40.2	26.1	1.1	0.0
1986	1,640.4	1,230.1	716.1	362.2	3,150.5	1,834.1	500.8	0.0	81.5	1.9	0.0
1987	2,261.1	1,185.5	698.4	0.0	5,114.5	3,013.1	584.6	0.0	52.4	2.7	0.0
1988	1,585.3	1,152.8	1,290.3	0.0	4,305.6	1,785.0	220.7	0.0	72.3	1.0	0.2
1989	1,838.3	2,027.5	647.7	0.0	4,957.7	2,726.9	253.6	0.1	44.7	0.4	0.0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1990	1,812.7	2,289.3	1,210.4	0.0	3,352.8	1,021.1	411.2	0.0	126.9	7.3	0.2
1991	996.4	1,989.2	878.9	0.0	1,779.9	260.2	234.8	0.0	89.7	5.2	0.3
1992	917.4	2,709.5	646.5	0.0	1,183.8	282.5	45.4	0.0	165.8	9.2	0.5
1993	1,088.3	3,457.0	1,109.8	1.2	1,706.8	547.9	51.6	0.0	63.7	44.7	0.5
1994	557.9	2,600.7	644.1	210.0	1,564.4	387.5	58.4	0.0	71.7	9.6	0.4
1995	1,361.1	2,386.7	339.0	292.7	1,283.4	700.7	57.6	0.0	19.0	7.2	0.1
1996	1,056.8	2,292.1	237.9	238.8	998.2	609.4	16.1	0.0	21.6	11.0	0.1
1997	1,032.5	2,502.8	241.7	253.6	1,453.1	735.8	16.4	0.0	22.4	15.6	0.0
1998	686.2	1,641.1	188.4	81.6	493.4	307.8	48.7	0.0	62.4	24.1	0.0
1999	485.0	945.0	182.7	100.1	1,634.2	315.9	10.0	0.0	29.0	14.7	0.1
2000	34.2	19.6	2.9	680.8	2,604.8	379.4	6.8	0.0	11.9	2.5	0.0
2001	9.3	28.8	1.0	310.3	1,092.4	287.1	7.0	0.0	6.4	0.7	0.0
2002	8.7	6.0	2.4	40.0	151.7	59.8	0.0	0.0	0.4	0.1	0.0
2003	3.1	0.3	0.2	0.4	0.0	9.3	0.4	0.0	0.3	0.6	0.0
2004	5.9	2.4	0.1	7.5	0.0	21.3	0.0	0.0	0.2	0.1	0.0
2005	2.7	0.2	0.2	5.2	0.0	27.6	0.1	0.0	0.4	0.8	0.1
2006	3.8	2.0	0.3	3.6	0.0	9.3	0.0	0.0	0.8	0.0	0.0
2007	2.7	1.8	0.3	1.0	0.0	0.5	2.9	0.0	1.6	0.3	0.0
2008	0.2	1.7	0.2	29.2	0.0	12.9	0.0	0.0	1.2	0.0	0.0
2009	1.9	2.1	0.2	2.3	0.0	34.1	0.2	0.0	0.4	0.0	0.0
2010	1.2	2.9	0.7	9.0	0.0	45.7	0.0	0.0	0.0	0.1	0.0
2011	1.1	10.0	7.2	0.0	12.4	31.5	0.0	0.0	0.0	0.0	0.0
2012	2.3	27.0	12.0	0.0	5.9	41.5	0.0	0.0	0.2	0.1	0.0
2013	4.8	44.0	2.4	0.0	204.5	36.6	0.0	0.0	0.9	0.1	0.0
2014	2.7	46.1	22.5	0.0	259.7	46.9	0.0	0.0	1.7	0.1	0.0
2015	1.8	10.4	0.0	0.0	386.7	92.5	0.0	0.0	1.6	0.4	0.0
2016	0.4	8.4	0.8	0.0	574.5	13.5	0.0	0.0	0.7	0.0	0.0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
2017	1.0	34.3	0.6	51.9	4,770.3	29.9	0.0	0.0	2.3	0.3	0.1
2018	21.4	14.5	0.1	215.9	7,605.9	1,552.4	0.0	0.0	1.5	0.0	0.0
2019	10.3	17.6	0.0	142.6	6,752.5	1,262.8	0.0	0.0	2.1	0.0	0.0
2020	8.7	64.9	0.0	84.8	5,834.1	1,613.3	0.0	0.0	2.3	0.2	0.3
2021	20.1	83.4	0.2	169.3	8,395.5	1,576.5	0.0	0.0	4.2	0.0	0.4
2022	60.0	66.8	0.0	499.1	8,794.5	1,546.2	0.0	0.0	8.4	0.0	0.3
2023	35.5	46.8	0.0	493.4	8,108.4	1,626.3	0.0	0.0	6.1	1.0	0.0
2024	16.0	11.6	0.0	0.0	7,449.8	1,711.0	0.0	0.0	11.9	0.9	0.1

Table 2: Catch (mt) from the foreign & domestic at-sea fleet and the domestic shoreside hake fleet. Catches (mt) from the Pacific whiting at-sea fishery as determined by onboard observers.

Year	Foreign & Domestic	Shoreside hake		
	At-sea	CA	OR	WA
1966	3,670.0	0.0	0.0	0.0
1967	3,902.0	0.0	0.0	0.0
1968	1,956.0	0.0	0.0	0.0
1969	358.0	0.0	0.0	0.0
1970	554.0	0.0	0.0	0.0
1971	701.0	0.0	0.0	0.0
1972	421.0	0.0	0.0	0.0
1973	656.0	0.0	0.0	0.0
1974	418.0	0.0	0.0	0.0
1975	391.2	0.0	0.0	0.0
1976	718.5	0.0	0.0	0.0
1977	119.3	0.0	0.0	0.0
1978	191.9	0.0	0.0	0.0
1979	197.9	0.0	0.0	0.0
1980	272.0	0.0	0.0	0.0
1981	227.9	0.0	0.0	0.0
1982	157.5	0.0	0.0	0.0
1983	131.5	0.0	0.0	0.0
1984	294.7	0.0	0.0	0.0
1985	182.6	0.0	0.0	0.0
1986	256.8	0.0	0.0	0.0
1987	181.3	0.0	0.0	0.0
1988	231.6	0.0	0.0	0.0

Year	Foreign & Domestic	Shoreside hake		
	At-sea	CA	OR	WA
1989	212.0	0.0	0.0	0.0
1990	230.2	0.0	0.0	0.0
1991	471.3	42.7	39.0	9.3
1992	389.6	13.5	42.1	6.2
1993	173.2	0.4	91.2	11.0
1994	370.7	2.1	210.8	28.6
1995	228.6	7.2	192.1	36.8
1996	252.2	5.7	475.1	104.7
1997	215.5	7.2	133.9	22.1
1998	268.5	40.4	278.0	28.1
1999	191.8	12.7	166.4	15.2
2000	205.4	7.7	70.9	4.7
2001	174.0	9.2	26.4	9.0
2002	154.9	1.2	2.6	1.4
2003	14.5	0.4	7.6	4.6
2004	21.2	7.4	12.4	8.5
2005	80.1	5.2	59.1	13.6
2006	143.0	3.6	11.3	35.3
2007	146.0	1.0	46.1	35.3
2008	115.2	29.2	36.1	37.5
2009	26.6	2.3	46.6	59.8
2010	44.6	9.0	35.3	17.5
2011	38.4	0.0	79.9	19.5
2012	79.2	0.0	85.1	17.1
2013	31.2	0.0	115.1	29.2
2014	56.2	0.0	250.1	35.9

Year	Foreign & Domestic	Shoreside hake		
	At-sea	CA	OR	WA
2015	37.4	0.0	267.9	80.8
2016	193.8	0.0	187.6	59.3
2017	481.3	0.0	791.1	182.8
2018	207.6	0.0	720.3	153.4
2019	198.9	0.0	605.6	297.0
2020	86.1	0.0	474.4	186.2
2021	115.5	0.0	419.7	82.1
2022	187.1	0.0	728.8	203.1
2023	206.5	0.0	368.8	98.0
2024	66.6	0.0	408.3	59.0

Table 3: Recent trend in the overfishing limits (OFLs), the acceptable biological catches (ABCs), the annual catch limits (ACLs), the total landings, and total mortality all in metric tons (mt).

Year	OFL (mt)	ABC (mt)	ACL (mt)	Landings (mt)	Total Mortality (mt)
2016	3,990	3,790.00	2,000	1,039.24	1,039.41
2017	14,130	13,508.28	13,508	6,345.89	6,361.45
2018	13,237	12,654.57	12,655	10,493.03	10,522.76
2019	12,375	11,830.50	11,831	9,289.43	9,315.27
2020	11,714	11,198.58	11,199	8,355.24	8,379.58
2021	15,749	14,725.32	14,725	10,866.89	10,899.73
2022	14,826	13,788.18	13,788	12,094.40	12,129.69
2023	13,633	12,624.16	12,624	10,990.57	11,023.50
2024	12,453	11,481.67	11,482	9,735.12	9,764.11
2025	12,254	11,236.92	11,237	10,838.45	10,868.09

Table 4: Number of trips sampled for length data by gear and state for non-whiting fisheries.

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1971	0	0	0	0	0	2	0	0	0	0	0
1977	0	0	0	25	0	0	0	0	0	0	0
1978	50	0	0	0	0	0	10	0	0	0	0
1979	32	0	0	0	0	0	4	0	3	0	0
1980	101	0	0	6	0	19	0	0	1	0	1
1981	72	3	0	59	20	31	0	0	6	0	0
1982	88	7	0	89	34	41	1	0	11	0	0
1983	158	16	0	46	10	25	18	0	9	0	0
1984	146	20	0	29	12	22	25	0	4	0	0
1985	149	20	0	25	35	16	81	0	5	0	0
1986	108	17	0	25	28	27	59	0	16	0	0
1987	88	29	0	49	74	36	37	0	3	0	0
1988	79	30	7	37	42	14	43	0	2	0	0
1989	81	49	14	30	67	16	81	0	7	0	0
1990	80	58	11	39	62	30	74	0	8	0	0
1991	74	76	20	17	63	15	23	0	12	0	0
1992	55	98	22	5	41	9	31	0	53	1	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1993	60	69	28	5	49	8	19	0	39	0	0
1994	54	67	13	2	21	16	34	0	38	0	0
1995	53	47	17	11	14	16	14	0	7	0	0
1996	49	33	17	11	12	13	4	0	10	0	0
1997	54	49	16	10	21	19	2	0	20	0	0
1998	41	43	26	3	11	8	5	0	15	0	0
1999	38	28	21	5	19	11	1	0	3	1	0
2000	14	0	3	16	44	19	0	0	8	1	0
2001	12	6	2	10	38	11	0	0	2	3	0
2002	22	8	7	1	15	10	1	0	2	0	0
2003	7	0	1	0	0	5	0	0	0	0	0
2004	5	1	1	0	0	12	0	0	0	0	0
2005	4	2	0	0	0	10	0	0	1	0	0
2006	7	3	2	0	0	8	0	0	4	1	0
2007	7	16	4	0	0	3	0	0	4	1	0
2008	5	18	5	0	0	12	0	0	2	0	0
2009	19	30	0	0	0	14	0	0	0	0	0
2010	18	22	1	0	0	11	0	0	0	2	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
2011	6	14	9	0	1	6	0	0	1	0	0
2012	14	19	5	0	4	7	0	0	3	1	0
2013	20	21	1	0	6	6	0	0	9	1	0
2014	18	20	3	0	5	7	0	0	12	2	0
2015	37	23	0	0	18	4	0	0	9	7	2
2016	27	14	0	0	7	1	0	0	2	4	2
2017	22	41	0	3	33	3	0	0	5	2	3
2018	31	25	7	10	60	4	0	0	3	4	7
2019	34	33	1	2	48	12	0	0	7	3	2
2020	29	18	0	2	31	5	0	0	13	8	1
2021	42	18	2	4	39	7	0	0	10	2	0
2022	13	10	0	12	46	4	0	0	2	5	2
2023	20	7	0	7	51	7	0	0	3	7	5
2024	27	13	0	0	52	9	0	0	16	9	4

Table 5: Number of lengths of widow rockfish by gear and state for non-whiting fisheries.

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1971	0	0	0	0	0	408	0	0	0	0	0
1977	0	0	0	96	0	0	0	0	0	0	0
1978	303	0	0	0	0	0	66	0	0	0	0
1979	436	0	0	0	0	0	34	0	7	0	0
1980	727	0	0	13	0	1,900	0	0	1	0	2
1981	444	250	0	1,340	1,746	3,100	0	0	19	0	0
1982	932	792	0	3,144	3,960	4,100	1	0	84	0	0
1983	1,352	478	0	1,411	321	2,500	103	0	31	0	0
1984	1,722	2,394	0	1,278	1,525	2,199	126	0	11	0	0
1985	1,853	2,233	0	1,176	3,971	1,600	557	0	8	0	0
1986	1,740	1,425	0	1,032	2,788	2,650	321	0	120	0	0
1987	998	865	0	1,744	2,198	1,942	262	0	11	0	0
1988	763	916	350	1,230	1,239	700	334	0	3	0	0
1989	1,007	1,099	700	1,325	1,843	800	450	0	23	0	0
1990	1,202	1,320	550	1,510	1,454	1,500	612	0	37	0	0
1991	1,596	1,569	997	761	1,442	750	268	0	75	0	0
1992	1,470	1,982	1,100	222	1,760	450	231	0	689	2	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1993	1,682	1,410	1,400	231	1,156	400	275	0	238	0	0
1994	1,359	1,464	650	112	557	842	410	0	554	0	0
1995	1,539	1,066	850	519	296	800	175	0	22	0	0
1996	1,364	845	704	437	316	650	132	0	80	0	0
1997	2,063	1,231	557	382	620	950	80	0	212	0	0
1998	1,368	1,013	865	125	291	400	179	0	318	0	0
1999	1,420	727	952	240	514	550	1	0	104	20	0
2000	263	0	101	641	1,147	950	0	0	65	1	0
2001	139	98	2	349	960	550	0	0	4	20	0
2002	318	185	136	39	319	500	2	0	74	0	0
2003	234	0	46	0	0	208	0	0	0	0	0
2004	26	18	3	0	0	508	0	0	0	0	0
2005	27	48	0	0	0	399	0	0	4	0	0
2006	79	58	7	0	0	461	0	0	36	1	0
2007	12	302	104	0	0	250	0	0	64	1	0
2008	8	274	76	0	0	1,086	0	0	27	0	0
2009	170	316	0	0	0	1,079	0	0	0	0	0
2010	205	233	100	0	0	903	0	0	0	5	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
2011	32	246	93	0	30	550	0	0	17	0	0
2012	136	353	241	0	95	688	0	0	9	7	0
2013	153	365	39	0	215	486	0	0	102	1	0
2014	134	324	106	0	150	700	0	0	242	4	0
2015	263	295	0	0	530	400	0	0	45	11	2
2016	143	254	0	0	210	100	0	0	38	4	24
2017	316	864	0	158	949	125	0	0	73	3	23
2018	645	161	12	507	1,492	350	0	0	32	7	10
2019	566	346	50	90	1,149	600	0	0	47	6	7
2020	593	228	0	83	759	233	0	0	134	15	1
2021	850	226	8	183	890	307	0	0	67	3	0
2022	272	185	0	502	1,025	180	0	0	11	7	3
2023	376	135	0	316	1,125	571	0	0	18	97	13
2024	540	164	0	0	1,255	830	0	0	268	86	10

Table 6: Number of hauls or trips and number of lengths sampled from the at-sea hake and shoreside hake fisheries.

Year	Number of hauls (at-sea) or trips (shoreside)		Number of lengths	
	Domestic at-sea	Shoreside	Domestic at-sea	Shoreside
1992	214	0	1,474	0
1993	239	0	1,468	0
1994	361	3	3,458	78
1995	304	19	1,789	570
1996	332	18	2,620	540
1997	397	30	2,841	869
1998	481	32	2,431	975
1999	598	52	3,070	1,551
2000	571	33	2,845	1,004
2001	522	19	1,758	576
2002	369	3	1,204	70
2003	291	2	665	26
2004	512	19	1,670	380
2005	1,228	1	5,538	50
2006	1,295	14	6,104	594
2007	1,491	21	10,658	860
2008	1,138	36	7,324	966
2009	400	24	1,976	845
2010	980	43	4,734	1,214
2011	982	43	3,605	1,286
2012	914	46	4,779	1,291
2013	901	40	3,808	1,160
2014	773	50	3,970	1,452

Year	Number of hauls (at-sea) or trips (shoreside)		Number of lengths	
	Domestic at-sea	Shoreside	Domestic at-sea	Shoreside
2015	522	36	2,312	1,313
2016	801	49	3,934	1,465
2017	997	57	5,406	1,353
2018	461	65	2,245	1,283
2019	469	73	2,642	1,536
2020	214	37	902	839
2021	310	61	1,776	1,279
2022	333	88	1,489	1,745
2023	469	68	1,738	1,525
2024	83	60	251	1,231

Table 7: Number of landings sampled for ages by gear and state for non-whiting fisheries.

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1978	7	0	0	0	0	0	0	0	0	0	0
1979	11	0	0	0	0	0	0	0	0	0	0
1980	27	0	0	0	0	18	0	0	0	0	0
1981	14	3	0	30	20	31	0	0	0	0	0
1982	87	6	0	71	34	40	1	0	4	0	0
1983	151	16	0	45	10	25	5	0	2	0	0
1984	144	20	0	29	12	22	10	0	2	0	0
1985	137	20	0	25	33	16	65	0	3	0	0
1986	106	17	0	22	28	27	53	0	3	0	0
1987	84	27	0	49	62	36	27	0	0	0	0
1988	67	29	6	34	41	14	39	0	2	0	0
1989	75	49	14	30	66	16	75	0	3	0	0
1990	70	58	11	32	62	30	65	0	2	0	0
1991	65	76	20	17	63	15	19	0	9	0	0
1992	45	93	22	4	26	9	21	0	15	0	0
1993	28	67	28	0	49	8	6	0	3	0	0
1994	28	67	13	2	21	15	7	0	1	0	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1995	8	45	17	3	13	16	0	0	0	0	0
1996	36	32	14	6	11	13	2	0	1	0	0
1997	42	46	11	10	20	19	0	0	9	0	0
1998	27	42	14	2	11	8	2	0	3	0	0
1999	29	27	19	3	18	10	0	0	0	0	0
2000	8	0	2	9	42	19	0	0	3	0	0
2001	2	6	0	4	35	10	0	0	0	0	0
2002	17	8	2	1	15	10	1	0	0	0	0
2003	3	0	0	0	0	5	0	0	0	0	0
2004	3	0	1	0	0	12	0	0	0	0	0
2005	0	2	0	0	0	10	0	0	0	0	0
2006	6	3	1	0	0	8	0	0	2	1	0
2007	6	16	4	0	0	3	0	0	3	1	0
2008	5	18	5	0	0	12	0	0	0	0	0
2009	8	29	0	0	0	14	0	0	0	0	0
2010	7	21	1	0	0	11	0	0	0	2	0
2011	0	5	7	0	1	5	0	0	0	0	0
2012	0	8	5	0	0	7	0	0	0	1	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
2013	0	7	1	0	3	5	0	0	0	0	0
2014	0	4	2	0	1	7	0	0	0	0	0
2015	0	22	0	0	14	4	0	0	0	0	1
2016	0	13	0	0	6	1	0	0	0	0	2
2017	0	36	0	0	31	3	0	0	0	0	3
2018	0	25	7	0	46	4	0	0	0	0	7
2019	0	16	1	0	34	12	0	0	0	0	1
2020	0	15	0	0	25	5	0	0	0	2	0
2021	0	12	2	0	31	6	0	0	0	0	0
2022	0	8	0	0	45	4	0	0	0	0	2
2023	0	5	0	0	48	7	0	0	0	1	5
2024	0	7	0	0	42	7	0	0	0	0	0

Table 8: Number of landings sampled for ages by gear and state for non-whiting fisheries.

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1980	404	0	0	0	0	1,775	0	0	0	0	0
1981	205	109	0	598	600	3,050	0	0	0	0	0
1982	834	174	0	2,382	1,019	3,944	1	0	18	0	0
1983	1,283	475	0	1,365	321	2,480	55	0	3	0	0
1984	1,678	600	0	1,278	360	2,194	89	0	5	0	0
1985	1,762	589	0	1,176	963	1,591	477	0	4	0	0
1986	1,704	680	0	913	939	2,594	188	0	5	0	0
1987	968	805	0	1,742	1,837	1,940	186	0	0	0	0
1988	692	886	298	1,132	1,209	695	290	0	3	0	0
1989	919	1,099	695	1,323	1,794	799	403	0	6	0	0
1990	1,051	1,310	550	1,309	1,447	1,497	533	0	8	0	0
1991	1,308	1,566	991	761	1,413	748	164	0	23	0	0
1992	676	1,889	1,097	82	574	450	87	0	91	0	0
1993	472	1,361	1,398	0	1,155	400	57	0	3	0	0
1994	516	1,463	650	54	556	749	58	0	1	0	0
1995	167	1,027	850	68	276	800	0	0	0	0	0
1996	873	827	699	158	292	649	88	0	7	0	0
1997	892	1,164	549	187	593	949	0	0	55	0	0

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
1998	1,019	987	699	82	291	400	84	0	46	0	0
1999	1,026	706	950	133	479	500	0	0	0	0	0
2000	157	0	100	353	1,067	948	0	0	12	0	0
2001	43	98	0	132	858	485	0	0	0	0	0
2002	294	179	99	21	319	488	2	0	0	0	0
2003	87	0	0	0	0	208	0	0	0	0	0
2004	7	0	3	0	0	506	0	0	0	0	0
2005	0	48	0	0	0	399	0	0	0	0	0
2006	74	58	6	0	0	361	0	0	5	1	0
2007	11	302	54	0	0	150	0	0	23	1	0
2008	8	274	75	0	0	600	0	0	0	0	0
2009	81	315	0	0	0	759	0	0	0	0	0
2010	54	231	50	0	0	539	0	0	0	5	0
2011	0	63	84	0	30	250	0	0	0	0	0
2012	0	80	73	0	0	163	0	0	0	7	0
2013	0	190	26	0	90	153	0	0	0	0	0
2014	0	91	52	0	30	229	0	0	0	0	0
2015	0	152	0	0	69	195	0	0	0	0	1

53

Year	Bottom Trawl			Midwater Trawl			Net		Hook-and-Line		
	CA	OR	WA	CA	OR	WA	CA	WA	CA	OR	WA
2016	0	156	0	0	36	28	0	0	0	0	24
2017	0	209	0	0	223	100	0	0	0	0	23
2018	0	161	12	0	495	200	0	0	0	0	10
2019	0	55	49	0	176	597	0	0	0	0	1
2020	0	61	0	0	134	233	0	0	0	3	0
2021	0	53	8	0	135	300	0	0	0	0	0
2022	0	44	0	0	281	129	0	0	0	0	3
2023	0	28	0	0	312	320	0	0	0	2	13
2024	0	33	0	0	248	340	0	0	0	0	0

Table 9: Discard totals (mt) for four fleets derived from Pikitch data, EDCP data, and WCGOP data. Italics indicate years that were not fitted to because they were simply added to the landings (Shoreside hake) or omitted because they were outside of the main study period.

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Bottom Trawl	1981	Pikitch	900.1900	54.26%	CV	2019 report	No
Bottom Trawl	1982	Pikitch	1,450.7400	44.12%	CV	2019 report	No
Bottom Trawl	1983	Pikitch	1,847.1500	43.91%	CV	2019 report	No
Bottom Trawl	1984	Pikitch	586.3600	55.78%	CV	2019 report	No

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Bottom Trawl	1985	Pikitch	462.9000	49.53%	CV	2019 data	Yes
Bottom Trawl	1986	Pikitch	534.8000	53.11%	CV	2019 data	Yes
Bottom Trawl	1987	Pikitch	1,035.5000	42.57%	CV	2019 data	Yes
Bottom Trawl	1988	Pikitch	1,177.0900	43.38%	CV	2019 report	No
Bottom Trawl	1989	Pikitch	1,217.7400	44.70%	CV	2019 report	No
Bottom Trawl	1990	Pikitch	1,010.9500	51.53%	CV	2019 report	No
Bottom Trawl	1991	Pikitch	1,219.2500	42.20%	CV	2019 report	No
Bottom Trawl	1992	Pikitch	1,217.5100	44.62%	CV	2019 report	No
Bottom Trawl	1993	Pikitch	1,430.1800	46.57%	CV	2019 report	No
Bottom Trawl	1994	Pikitch	1,177.7100	43.11%	CV	2019 report	No
Bottom Trawl	1995	EDCP	924.8000	83.18%	CV	2019 data	Yes
Bottom Trawl	1996	EDCP	3,084.5000	67.07%	CV	2019 data	Yes
Bottom Trawl	1997	EDCP	3,353.3000	75.06%	CV	2019 data	Yes
Bottom Trawl	1998	EDCP	42.6000	48.80%	CV	2019 data	Yes
Bottom Trawl	1999	EDCP	4.8000	68.78%	CV	2019 data	Yes
Bottom Trawl	2002	WCGOP	13.2200	43.07%	CV	2019 data	Yes
Bottom Trawl	2003	WCGOP	1.2100	81.96%	CV	2019 data	Yes
Bottom Trawl	2004	WCGOP	5.1300	75.89%	CV	2019 data	Yes
Bottom Trawl	2005	WCGOP	10.1700	44.61%	CV	2019 data	Yes

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Bottom Trawl	2006	WCGOP	0.0300	135.56%	CV	2019 data	Yes
Bottom Trawl	2007	WCGOP	13.8600	61.57%	CV	2019 data	Yes
Bottom Trawl	2008	WCGOP	3.9000	44.54%	CV	2019 data	Yes
Bottom Trawl	2009	WCGOP	26.5700	33.77%	CV	2019 data	Yes
Bottom Trawl	2010	WCGOP	22.7400	54.32%	CV	2019 data	Yes
Bottom Trawl	2011	WCGOP	0.0800	5.00%	CV	2019 data	Yes
Bottom Trawl	2012	WCGOP	0.0100	5.00%	CV	2019 data	Yes
Bottom Trawl	2013	WCGOP	2.4300	5.00%	CV	2019 data	Yes
Bottom Trawl	2014	WCGOP	0.0900	5.00%	CV	2019 data	Yes
Bottom Trawl	2015	WCGOP	0.0300	5.00%	CV	2019 data	Yes
Bottom Trawl	2016	WCGOP	0.0200	5.00%	CV	2019 data	Yes
Bottom Trawl	2017	WCGOP	0.2600	5.00%	CV	2019 data	Yes
Bottom Trawl	2018	WCGOP	0.0143	5.00%	CV	2025 data	Yes
Bottom Trawl	2019	WCGOP	0.7832	5.00%	CV	2025 data	Yes
Bottom Trawl	2020	WCGOP	0.2763	5.00%	CV	2025 data	Yes
Bottom Trawl	2021	WCGOP	0.1440	5.00%	CV	2025 data	Yes
Bottom Trawl	2022	WCGOP	0.0750	5.00%	CV	2025 data	Yes
Bottom Trawl	2023	WCGOP	0.1184	5.00%	CV	2025 data	Yes
Midwater	1981	Pikitch	6,479.8800	23.24%	CV	2019 report	No

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Midwater	1982	Pikitch	5,722.2500	22.84%	CV	2019 report	No
Midwater	1984	Pikitch	1,737.5700	23.33%	CV	2019 report	No
Midwater	1985	Pikitch	1,502.0000	24.09%	CV	2019 data	Yes
Midwater	1986	Pikitch	1,321.2000	23.64%	CV	2019 data	Yes
Midwater	1987	Pikitch	1,798.4000	26.20%	CV	2019 data	Yes
Midwater	1988	Pikitch	1,615.8300	24.82%	CV	2019 report	No
Midwater	1989	Pikitch	1,981.8600	25.26%	CV	2019 report	No
Midwater	1990	Pikitch	1,205.4400	24.51%	CV	2019 report	No
Midwater	1991	Pikitch	565.9400	24.33%	CV	2019 report	No
Midwater	1992	Pikitch	356.0000	25.00%	CV	2019 report	No
Midwater	1993	Pikitch	569.8600	25.34%	CV	2019 report	No
Midwater	1994	Pikitch	536.8000	25.43%	CV	2019 report	No
Midwater	1995	Pikitch	663.2400	23.81%	CV	2019 report	No
Midwater	1996	Pikitch	465.6600	24.84%	CV	2019 report	No
Midwater	1997	Pikitch	663.1400	24.10%	CV	2019 report	No
Midwater	1998	Pikitch	217.1500	25.53%	CV	2019 report	No
Midwater	1997	EDCP	1.0000	83.26%	CV	2019 data	Yes
Midwater	1998	EDCP	18.7000	80.00%	CV	2019 data	Yes
Midwater	2002	WCGOP	39.4000	40.71%	CV	2019 data	Yes

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Midwater	2012	WCGOP	0.0000	5.00%	CV	2025 data	Yes
Midwater	2013	WCGOP	0.0020	5.00%	CV	2025 data	Yes
Midwater	2014	WCGOP	0.0136	5.00%	CV	2025 data	Yes
Midwater	2015	WCGOP	0.8800	5.00%	CV	2025 data	Yes
Midwater	2016	WCGOP	1.5600	5.00%	CV	2025 data	Yes
Midwater	2017	WCGOP	9.7500	5.00%	CV	2025 data	Yes
Midwater	2018	WCGOP	37.2300	5.00%	CV	2025 data	Yes
Midwater	2019	WCGOP	18.7800	5.00%	CV	2025 data	Yes
Midwater	2020	WCGOP	45.4400	5.00%	CV	2025 data	Yes
Midwater	2021	WCGOP	36.3800	5.00%	CV	2025 data	Yes
Midwater	2022	WCGOP	47.6000	5.00%	CV	2025 data	Yes
Midwater	2023	WCGOP	17.3700	5.00%	CV	2025 data	Yes
Hook & Line	2007	WCGOP	0.0160	31.8%	CV	2025 data	Yes, in landings
Hook & Line	2008	WCGOP	0.0120	28.3%	CV	2025 data	Yes, in landings
Hook & Line	2010	WCGOP	0.0080	44.9%	CV	2025 data	Yes, in landings
Hook & Line	2011	WCGOP	0.0110	27.4%	CV	2025 data	Yes, in landings
Hook & Line	2012	WCGOP	0.0010	44.9%	CV	2025 data	Yes, in landings
Hook & Line	2013	WCGOP	0.0010	10.0%	CV	2025 data	Yes, in landings
Hook & Line	2014	WCGOP	0.0090	27.4%	CV	2025 data	Yes, in landings

Fleet	Year	Source	Discards	Error	Error Stat	Source 1	Used in Assessment
Hook & Line	2015	WCGOP	0.0550	55.8%	CV	2025 data	Yes, in landings
Hook & Line	2016	WCGOP	0.1880	15.8%	CV	2025 data	Yes, in landings
Hook & Line	2017	WCGOP	0.0370	47.0%	CV	2025 data	Yes, in landings
Hook & Line	2018	WCGOP	0.1380	35.5%	CV	2025 data	Yes, in landings
Hook & Line	2019	WCGOP	0.0120	44.9%	CV	2025 data	Yes, in landings
Hook & Line	2021	WCGOP	0.0000	44.9%	CV	2025 data	Yes, in landings
Hook & Line	2022	WCGOP	0.0270	54.7%	CV	2025 data	Yes, in landings
Hook & Line	2023	WCGOP	0.0020	64.8%	CV	2025 data	Yes, in landings

5.1.2 Fishery-independent data

Table 10: Depth ranges and limits of the southern latitude in the Triennial survey for the different years.

Years	Depth range (m)	Southern latitude
1977	91-457	34.05
1980-1986	55-366	36.80
1989-1992	55-366	34.50
1995-2004	55-500	34.50

Table 11: Stratifications used for the two surveys.

Triennial					
Strata	Area (km2)	Depth1	Depth2	Latitude1	Latitude2
A	33,730.25	55	183	34.5	49
B	11,062.63	183	400	34.5	49

Table 12: Stratifications used for the two surveys.

NWFSC WCGBT					
Strata	Area (km2)	Depth1	Depth2	Latitude1	Latitude2
A	10,687.86	55	183	34.5	40.5
B	3,394.82	183	400	34.5	40.5
C	23,042.39	55	183	40.5	49.0
D	7,667.81	183	400	40.5	49.0

Table 13: Number of positive tows, lengths, and ages in each year from the Triennial survey (Tri) and the NWFSC WCGBT survey (NW).

Year	Number of positive tows		Number of tows with lengths		Number of lengths		Number of tows with ages		Number of ages	
	Tri	NW	Tri	NW	Tri	NW	Tri	NW	Tri	NW
1977	80		1		9					
1980	38		3		166		1		22	
1983	70		5		385					
1986	46		8		317					
1989	38		20		713					
1992	50		10		708					
1995	43		43		500					
1998	59		58		738					
2001	28		28		130					
2003		20		18		216				
2004	36	12	33	12	219	84		12		43
2005		20		20		78		18		65
2006		26		26		172		26		89
2007		27		27		92		27		83
2008		17		17		26		15		20
2009		31		31		141		31		123
2010		28		28		240		28		116
2011		31		31		313		31		152
2012		32		32		181		32		91
2013		18		18		364		18		246
2014		29		28		349		28		264
2015		21		21		149		21		93
2016		40		40		888		40		556
2017		30		30		310		30		213

Year	Number of positive tows		Number of tows with lengths		Number of lengths		Number of tows with ages		Number of ages	
	Tri	NW	Tri	NW	Tri	NW	Tri	NW	Tri	NW
2018		34		34		410		34		353
2019		23		23		219		23		161
2021		18		17		66		17		66
2022		18		18		125		18		109
2023		30		29		159		29		110
2024		35		35		485		35		347

Table 14: Ageing error for two labs that was used in the assessment model.

True Age	Standard Deviation CAP	Standard Deviation SWFSC
0.5	0.145	0.111
1.5	0.145	0.111
2.5	0.187	0.147
3.5	0.233	0.187
4.5	0.283	0.233
5.5	0.338	0.284
6.5	0.398	0.341
7.5	0.463	0.406
8.5	0.533	0.478
9.5	0.612	0.560
10.5	0.697	0.651
11.5	0.790	0.755
12.5	0.892	0.871
13.5	1.003	1.001
14.5	1.124	1.148
15.5	1.256	1.313
16.5	1.401	1.499

True Age	Standard Deviation CAP	Standard Deviation SWFSC
17.5	1.558	1.708
18.5	1.731	1.943
19.5	1.919	2.207
20.5	2.124	2.504
21.5	2.349	2.839
22.5	2.594	3.215
23.5	2.861	3.638
24.5	3.154	4.113
25.5	3.473	4.649
26.5	3.821	5.250
27.5	4.202	5.927
28.5	4.618	6.689
29.5	5.072	7.545
30.5	5.568	8.508
31.5	6.109	9.592
32.5	6.700	10.810
33.5	7.346	12.181
34.5	8.052	13.723
35.5	8.822	15.456
36.5	9.663	17.407
37.5	10.582	19.600
38.5	11.585	22.067
39.5	12.680	24.842
40.5	13.877	27.964

5.2 Model results

Table 15: Specifications of the base assessment model for widow rockfish.

<i>Starting year</i>	1916
<i>Population characteristics</i>	
Maximum age	40
Genders	2
Population length bins	6-60 cm by 1 cm bins
Summary biomass (mt)	Age 4+
<i>Data characteristics</i>	
Data lengths	8-56 cm by 2 cm bins
Data ages	14611
First reference age for growth calcs	3
Second reference age for growth calcs	40
First mature age	3
Starting year of estimated recruitment	1900
<i>Fishery characteristics</i>	
Catchability	Analytical estimate
Fishery selectivity (not midwater trawl)	Asymptotic Double Normal
Midwater trawl fishery selectivity	Dome-shaped Double Normal
Juvenile survey selectivity	Double Normal
Triennial survey selectivity	Cubic spline with 3 nodes
NWFSC WCGBT survey selectivity	Cubic spline with 3 nodes
<i>Fishery time blocks</i>	
Bottom trawl selectivity	1916 - 2001, 2002 -
Bottom trawl retention	1916 - 1981, 2011 -
Midwater trawl selectivity	1916-1982, 1983-2001, 2002-2010, 2011-
Midwater trawl retention	1916-1982, 1983-2001, 2002-2010, 2011-2016, 2017 -
Hake trawl selectivity	1916 - 2019, 2020 -
Hook-and-line selectivity	1916-2002, 2003-

Table 16: Parameter estimates, parameter bounds (low, high), estimation status, estimated standard deviation (SD), prior information [distribution(mean, SD)] used in the base model.

Label	Value	Bounds	Status	SD	Prior
NatM_- uniform_Fem_- GP_1	0.125	(0.01, 0.3)	ok	0	lognor- mal(0.100, 0.310)
L_at_Amin_- Fem_GP_1	20.6	(10, 40)	ok	0	none
L_at_Amax_- Fem_GP_1	49.5	(35, 60)	ok	0	none
VonBert_K_- Fem_GP_1	0.181	(0.01, 0.4)	ok	0	none
CV_young_- Fem_GP_1	0.116	(0.01, 0.4)	ok	0	none
CV_old_Fem_- GP_1	0.0482	(0.01, 0.4)	ok	0	none
Wtlen_1_Fem_- GP_1	1.59e-05	(-3, 3)	fixed	0	none
Wtlen_2_Fem_- GP_1	2.99	(-3, 10)	fixed	0	none
Mat50%_Fem_- GP_1	5.47	(-3, 50)	fixed	0	none
Mat_slope_- Fem_GP_1	-0.775	(-3, 3)	fixed	0	none
Eggs/kg_inter_- Fem_GP_1	1	(-1, 1)	fixed	0	none
Eggs/kg_slope_- wt_Fem_GP_1	0	(0, 1)	fixed	0	none
NatM_- uniform_Mal_- GP_1	0.137	(0.01, 0.3)	ok	0	lognor- mal(0.100, 0.310)
L_at_Amin_- Mal_GP_1	21	(10, 40)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
L_at_Amax_- Mal_GP_1	43.6	(35, 60)	ok	0	none
VonBert_K_- Mal_GP_1	0.245	(0.01, 0.4)	ok	0	none
CV_young_- Mal_GP_1	0.0942	(0.01, 0.4)	ok	0	none
CV_old_Mal_- GP_1	0.0569	(0.01, 0.4)	ok	0	none
Wtlen_1_Mal_- GP_1	1.45e-05	(-3, 3)	fixed	0	none
Wtlen_2_Mal_- GP_1	3.01	(-3, 10)	fixed	0	none
CohortGrowDev	1	(0, 2)	fixed	0	none
FracFemale_- GP_1	0.5	(1e-06, 1)	fixed	0	none
SR_LN(R0)	10.4	(1, 20)	ok	0	none
SR_BH_stEEP	0.72	(0.2, 1)	fixed	0	beta(0.720, 0.160)
SR_sigmaR	0.6	(0, 2)	fixed	0	none
SR_regime	0	(-5, 5)	fixed	0	none
SR_autocorr	0	(0, 0.5)	fixed	0	none
Early_InitAge_- 16	-0.00142	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_- 15	-0.00161	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_- 14	-0.00181	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_- 13	-0.00204	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_- 12	-0.0023	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_InitAge_-11	-0.00258	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_-10	-0.00289	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_9	-0.00323	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_8	-0.00361	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_7	-0.00401	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_6	-0.00444	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_5	-0.00491	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_4	-0.00542	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_3	-0.00597	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_2	-0.00658	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_InitAge_1	-0.00725	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_RecrDev_1916	-0.00799	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_RecrDev_1917	-0.00881	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_RecrDev_1918	-0.00971	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_RecrDev_1919	-0.0107	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_RecrDev_1920	-0.0118	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_- RecrDev_1921	-0.013	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1922	-0.0143	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1923	-0.0158	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1924	-0.0174	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1925	-0.0191	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1926	-0.0211	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1927	-0.0232	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1928	-0.0256	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1929	-0.0281	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1930	-0.031	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1931	-0.0341	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1932	-0.0375	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1933	-0.0413	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1934	-0.0454	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1935	-0.0499	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1936	-0.0549	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_- RecrDev_1937	-0.0604	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1938	-0.0665	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1939	-0.0733	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1940	-0.0808	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1941	-0.0891	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1942	-0.0982	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1943	-0.108	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1944	-0.118	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1945	-0.128	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1946	-0.138	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1947	-0.147	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1948	-0.156	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1949	-0.164	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1950	-0.171	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1951	-0.175	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1952	-0.175	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_- RecrDev_1953	-0.17	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1954	-0.159	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1955	-0.139	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1956	-0.112	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1957	-0.0794	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1958	-0.0493	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1959	-0.0319	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1960	-0.0332	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1961	-0.0445	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1962	-0.0438	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1963	-0.013	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1964	0.0405	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1965	0.0985	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1966	0.159	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1967	0.172	(-5, 5)	dev	0	normal(0.00, 0.60)
Early_- RecrDev_1968	0.123	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_- RecrDev_1969	-0.0191	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1970	1.39	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1971	1.2	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1972	-0.804	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1973	-0.967	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1974	-0.657	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1975	0.0929	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1976	-1.04	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1977	0.698	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1978	1.12	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1979	-0.0286	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1980	0.526	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1981	1.05	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1982	0.431	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1983	0.0109	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1984	0.697	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Main_- RecrDev_1985	0.38	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1986	-0.378	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1987	0.558	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1988	0.105	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1989	-0.203	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1990	0.246	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1991	0.844	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1992	-0.135	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1993	0.0968	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1994	0.00952	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1995	-0.458	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1996	-0.759	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1997	-0.333	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1998	0.256	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_1999	0.537	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2000	0.345	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Main_- RecrDev_2001	-0.248	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2002	-0.374	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2003	-0.416	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2004	0.758	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2005	-0.902	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2006	0.476	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2007	-0.924	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2008	1.31	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2009	-0.725	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2010	0.874	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2011	-1.04	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2012	-1.45	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2013	0.723	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2014	-0.171	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2015	-0.432	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2016	1.25	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Main_- RecrDev_2017	-0.0326	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2018	-1.01	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2019	-1.27	(-5, 5)	dev	0	normal(0.00, 0.60)
Main_- RecrDev_2020	-1.21	(-5, 5)	dev	0	normal(0.00, 0.60)
Late_RecrDev_- 2021	-0.478	(-5, 5)	dev	0	normal(0.00, 0.60)
Late_RecrDev_- 2022	-0.412	(-5, 5)	dev	0	normal(0.00, 0.60)
Late_RecrDev_- 2023	-0.0365	(-5, 5)	dev	0	normal(0.00, 0.60)
Late_RecrDev_- 2024	-0.189	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2025	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2026	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2027	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2028	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2029	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2030	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2031	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2032	0	(-5, 5)	dev	0	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
ForeRecr_2033	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2034	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2035	0	(-5, 5)	dev	0	normal(0.00, 0.60)
ForeRecr_2036	0	(-5, 5)	dev	0	normal(0.00, 0.60)
LnQ_base_- BottomTrawl(1)	-5.99	(-25, 25)	fixed	0	none
Q_extraSD_- BottomTrawl(1)	0.165	(0, 2)	ok	0	none
LnQ_base_- Hake(3)	-11.1	(-20, 2)	ok	0	none
Q_extraSD_- Hake(3)	0.371	(0, 2)	ok	0	none
LnQ_base_- JuvSurvey(6)	-1.63	(-25, 25)	fixed	0	none
Q_extraSD_- JuvSurvey(6)	1.68	(0, 2)	ok	0	none
LnQ_base_- Triennial(7)	-2.06	(-4, 4)	ok	0	none
Q_extraSD_- Triennial(7)	0	(0, 2)	fixed	0	none
LnQ_base_- NWFSC(8)	-3.13	(-25, 25)	fixed	0	none
Q_extraSD_- NWFSC(8)	0	(0, 2)	fixed	0	none
LnQ_base_- ForeignAtSea(9)	-11.4	(-25, 25)	fixed	0	none
Q_extraSD_- ForeignAtSea(9)	0.578	(0, 2)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
LnQ_base_-					
Hake(3)_-	0.467	(1e-04, 2)	ok	0	normal(0.500, 0.500)
BLK10add_-					
1991					
LnQ_base_-					
Triennial(7)_-	0.161	(1e-04, 2)	ok	0	normal(0.500, 0.500)
BLK9add_1995					
Size_DblN_-					
peak_-	43.4	(10, 59)	ok	0	none
BottomTrawl(1)					
Size_DblN_-					
top_logit_-	2.5	(-5, 10)	ok	0	none
BottomTrawl(1)					
Size_DblN_-					
ascend_se_-	4.59	(-4, 12)	ok	0	none
BottomTrawl(1)					
Size_DblN_-					
descend_se_-	9	(-2, 10)	fixed	0	none
BottomTrawl(1)					
Size_DblN_-					
start_logit_-	-9	(-9, 10)	fixed	0	none
BottomTrawl(1)					
Size_DblN_-					
end_logit_-	8	(-9, 9)	fixed	0	none
BottomTrawl(1)					
Retain_L_infl_-					
BottomTrawl(1)	3.6	(-5, 60)	ok	0	none
Retain_L_-					
width_-	0.955	(0.01, 8)	ok	0	none
BottomTrawl(1)					
Retain_L_-					
asymptote_-					
logit_-	4.6	(-10, 10)	fixed	0	none
BottomTrawl(1)					

Label	Value	Bounds	Status	SD	Prior
Retain_L_-maleoffset_-BottomTrawl(1)	0	(-10, 10)	fixed	0	none
Size_DblN_-peak_Midwater-Trawl(2)	36.9	(10, 59)	ok	0	none
Size_DblN_-top_logit_MidwaterTrawl(2)	-9.43	(-10, 10)	ok	0	none
Size_DblN_as-cend_se_MidwaterTrawl(2)	2.87	(-4, 12)	ok	0	none
Size_DblN_descend_se_MidwaterTrawl(2)	3.91	(-2, 10)	ok	0	none
Size_DblN_-start_logit_MidwaterTrawl(2)	-9	(-9, 10)	fixed	0	none
Size_DblN_-end_logit_MidwaterTrawl(2)	-1.35	(-9, 9)	ok	0	none
Retain_L_infl_-Midwater-Trawl(2)	-5	(-5, 60)	fixed	0	none
Retain_L_-width_MidwaterTrawl(2)	1.2	(0.01, 8)	fixed	0	none
Retain_L_-asymptote_-logit_Midwater-Trawl(2)	5.77	(-10, 10)	ok	0	none
Retain_L_male-offset_MidwaterTrawl(2)	0	(-10, 10)	fixed	0	none
Size_DblN_-peak_Hake(3)	33.5	(10, 59)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
Size_DblN_- top_logit_- Hake(3)	2.5	(-5, 10)	ok	0	none
Size_DblN_- ascend_se_- Hake(3)	2.1	(-4, 12)	ok	0	none
Size_DblN_- descend_se_- Hake(3)	9	(-2, 10)	fixed	0	none
Size_DblN_- start_logit_- Hake(3)	-9	(-9, 10)	fixed	0	none
Size_DblN_- end_logit_- Hake(3)	8	(-9, 9)	fixed	0	none
Size_DblN_- peak_Net(4)	42.6	(10, 59)	ok	0	none
Size_DblN_- top_logit_Net(4)	2.51	(-5, 10)	ok	0	none
Size_DblN_- ascend_se_- Net(4)	3.56	(-4, 12)	ok	0	none
Size_DblN_- descend_se_- Net(4)	9	(-2, 10)	fixed	0	none
Size_DblN_- start_logit_- Net(4)	-9	(-9, 10)	fixed	0	none
Size_DblN_- end_logit_- Net(4)	8	(-9, 9)	fixed	0	none
Size_DblN_- peak_HnL(5)	23.6	(10, 59)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
Size_DblN_- top_logit_- HnL(5)	2.5	(-5, 10)	ok	0	none
Size_DblN_- ascend_se_- HnL(5)	-5	(-5, 12)	fixed	0	none
Size_DblN_- descend_se_- HnL(5)	9	(-2, 10)	fixed	0	none
Size_DblN_- start_logit_- HnL(5)	-9	(-9, 10)	fixed	0	none
Size_DblN_- end_logit_- HnL(5)	8	(-9, 9)	fixed	0	none
SizeSpline_- Code_- Triennial(7)	0	(0, 2)	fixed	0	none
SizeSpline_- GradLo_- Triennial(7)	0.119	(-0.001, 1)	ok	0	none
SizeSpline_- GradHi_- Triennial(7)	0.0381	(-1, 1)	ok	0	none
SizeSpline_- Knot_1_- Triennial(7)	24	(8, 56)	fixed	0	none
SizeSpline_- Knot_2_- Triennial(7)	34	(8, 56)	fixed	0	none
SizeSpline_- Knot_3_- Triennial(7)	48	(8, 56)	fixed	0	none
SizeSpline_- Val_1_- Triennial(7)	-1.82	(-10, 10)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
SizeSpline_- Val_2_- Triennial(7)	-1	(-10, 10)	fixed	0	none
SizeSpline_- Val_3_- Triennial(7)	0.435	(-10, 10)	ok	0	none
SizeSpline_- Code_- NWFSC(8)	0	(0, 2)	fixed	0	none
SizeSpline_- GradLo_- NWFSC(8)	0.467	(-0.001, 1)	ok	0	none
SizeSpline_- GradHi_- NWFSC(8)	-0.109	(-1, 1)	ok	0	none
SizeSpline_- Knot_1_- NWFSC(8)	24	(8, 56)	fixed	0	none
SizeSpline_- Knot_2_- NWFSC(8)	34	(8, 56)	fixed	0	none
SizeSpline_- Knot_3_- NWFSC(8)	48	(8, 56)	fixed	0	none
SizeSpline_- Val_1_- NWFSC(8)	-2.23	(-10, 10)	ok	0	none
SizeSpline_- Val_2_- NWFSC(8)	-1	(-10, 10)	fixed	0	none
SizeSpline_- Val_3_- NWFSC(8)	-0.116	(-10, 10)	ok	0	none
SizeSel_P1_- ForeignAtSea(9)	1	(-2, 60)	fixed	0	none

Label	Value	Bounds	Status	SD	Prior
SizeSel_P2_- ForeignAtSea(9)	-1	(-2, 60)	fixed	0	none
mi- nage@sel=1_Ju- vSurvey(6)	0	(0, 1)	fixed	0	none
max- age@sel=1_Ju- vSurvey(6)	0	(0, 1)	fixed	0	none
mi- nage@sel=1_- NWFSC(8)	0	(0, 1)	fixed	0	none
max- age@sel=1_- NWFSC(8)	40	(0, 50)	fixed	0	none
Size_DblN_- peak_Bottom- Trawl(1)_- BLK4rep1_1916	39	(10, 59)	ok	0	none
Size_DblN_as- cend_se_Bot- tomTrawl(1)_- BLK4rep1_1916	3.43	(-4, 12)	ok	0	none
Retain_L_infl_- Bottom- Trawl(1)_- BLK2rep1_1982	27.2	(-5, 50)	ok	0	none
Retain_L_infl_- Bottom- Trawl(1)_- BLK2rep1_1990	27.5	(-5, 50)	ok	0	none
Retain_L_- width_Bottom- Trawl(1)_- BLK2rep1_1982	0.97	(0.01, 5)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
Retain_L_- width_Bottom- Trawl(1)_- BLK2repl_1990	1.83	(0.01, 5)	ok	0	none
Retain_L_- asymptote_- logit_Bottom- Trawl(1)_- BLK1repl_1982	1.72	(-10, 10)	ok	0	none
Retain_L_- asymptote_- logit_Bottom- Trawl(1)_- BLK1repl_1990	0.785	(-10, 10)	ok	0	none
Retain_L_- asymptote_- logit_Bottom- Trawl(1)_- BLK1repl_1998	0.108	(-10, 10)	ok	0	none
Size_DblN_- peak_Midwater- Trawl(2)_- BLK7repl_1916	38.7	(10, 59)	ok	0	none
Size_DblN_- peak_Midwater- Trawl(2)_- BLK7repl_1983	38	(10, 59)	ok	0	none
Size_DblN_- peak_Midwater- Trawl(2)_- BLK7repl_2002	37.4	(10, 59)	ok	0	none
Size_DblN_as- cend_se_Mid- waterTrawl(2)_- BLK7repl_1916	3.37	(-4, 12)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
Size_DblN_as-cend_se_Mid-waterTrawl(2)_-BLK7repl_1983	3.08	(-4, 12)	ok	0	none
Size_DblN_as-cend_se_Mid-waterTrawl(2)_-BLK7repl_2002	2.8	(-4, 12)	ok	0	none
Size_DblN_descend_se_Mid-waterTrawl(2)_-BLK7repl_1916	4.24	(-2, 10)	ok	0	none
Size_DblN_descend_se_Mid-waterTrawl(2)_-BLK7repl_1983	3.06	(-2, 10)	ok	0	none
Size_DblN_descend_se_Mid-waterTrawl(2)_-BLK7repl_2002	-1.42	(-2, 10)	ok	0	none
Size_DblN_end_logit_Mid-waterTrawl(2)_-BLK7repl_1916	-1.93	(-9, 9)	ok	0	none
Size_DblN_end_logit_Mid-waterTrawl(2)_-BLK7repl_1983	-0.424	(-9, 9)	ok	0	none
Size_DblN_end_logit_Mid-waterTrawl(2)_-BLK7repl_2002	1.56	(-9, 9)	ok	0	none

Label	Value	Bounds	Status	SD	Prior
Retain_L_- asymptote_- logit_Midwater- Trawl(2)_-	4.59	(-10, 10)	fixed	0	none
BLK12repl_- 1916					
Retain_L_- asymptote_- logit_Midwater- Trawl(2)_-	1.66	(-10, 10)	ok	0	none
BLK12repl_- 1983					
Retain_L_- asymptote_- logit_Midwater- Trawl(2)_-	1.85	(-10, 10)	ok	0	none
BLK12repl_- 2002					
Retain_L_- asymptote_- logit_Midwater- Trawl(2)_-	8.95	(-10, 10)	ok	0	none
BLK12repl_- 2011					
Size_DblN_- peak_Hake(3)_- BLK11repl_- 1916	42.8	(10, 59)	ok	0	none
Size_DblN_- top_logit_- Hake(3)_-	2.5	(-5, 10)	ok	0	none
BLK11repl_- 1916					
Size_DblN_- ascend_se_- Hake(3)_-	3.72	(-4, 12)	ok	0	none
BLK11repl_- 1916					

Label	Value	Bounds	Status	SD	Prior
Size_DblN_-peak_HnL(5)_-BLK5repl_1916	37.2	(15, 59)	ok	0	none
Size_DblN_-ascend_se_-HnL(5)_-BLK5repl_1916	3.75	(-4, 12)	ok	0	none

Table 17: Likelihood components and other quantities related to the minimization of the base case model.

component	log.likelihood	lambdas
Total	19,053	
Survey	13.015	
Discard	16,793.300	
Length composition	860.879	
Age composition	1,366.410	
Recruitment	17.868	1
Forecast Recruitment	0.605	1
Priors	0.976	1
Softbounds	0.010	

Table 18: Estimates of key derived parameters and reference points with approximate 95% asymptotic confidence intervals.

Reference Point	Estimate	95% Confidence Interval
Unfished Spawning Biomass (mt)	81479.2	81479.2 - 81479.2
Unfished Age 4+ Biomass (mt)	151093	151093 - 151093
Unfished Recruitment (R0)	33951	33951 - 33951
2026 Spawning Biomass (mt)	34267.9	34267.9 - 34267.9
2026 Fraction Unfished	0.421	0.421 - 0.421

Reference Point	Estimate	95% Confidence Interval
Reference Points Based SB40%		
Proxy Spawning Biomass (mt) SB40%	32591.7	32591.7 - 32591.7
SPR Resulting in SB40%	0.458	0.458 - 0.458
Exploitation Rate Resulting in SB40%	0.087	0.087 - 0.087
Yield with SPR Based On SB40% (mt)	5865.9	5865.9 - 5865.9
Reference Points Based on SPR Proxy for MSY		
Proxy Spawning Biomass (mt) (SPR50)	36352.3	36352.3 - 36352.3
SPR50	0.5	
Exploitation Rate Corresponding to SPR50	0.076	0.076 - 0.076
Yield with SPR50 at SB SPR (mt)	5594.01	5594.01 - 5594.01
Reference Points Based on Estimated MSY Values		
Spawning Biomass (mt) at MSY (SB MSY)	21615.5	21615.5 - 21615.5
SPR MSY	0.337	0.337 - 0.337
Exploitation Rate Corresponding to SPR MSY	0.131	0.131 - 0.131
MSY (mt)	6270.27	6270.27 - 6270.27

Table 19: Time series of population estimates from the base case model.

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
1916	156,405.0	81,337.8	150,770.0	0.998	33,675.0	78.4	0.012	0.0
1917	156,282.0	81,273.0	150,651.0	0.997	33,645.0	121.9	0.018	0.0
1918	156,118.0	81,184.3	150,492.0	0.996	33,611.2	140.2	0.021	0.0
1919	155,940.0	81,087.1	150,319.0	0.995	33,574.1	96.8	0.015	0.0
1920	155,808.0	81,016.5	150,192.0	0.994	33,534.7	99.1	0.015	0.0
1921	155,675.0	80,946.1	150,065.0	0.993	33,491.7	82.1	0.012	0.0
1922	155,557.0	80,886.1	149,954.0	0.993	33,445.1	71.3	0.011	0.0
1923	155,448.0	80,832.3	149,851.0	0.992	33,394.2	78.3	0.012	0.0
1924	155,327.0	80,773.6	149,738.0	0.991	33,338.4	50.3	0.008	0.0
1925	155,228.0	80,729.0	149,647.0	0.991	33,277.8	62.3	0.009	0.0
1926	155,109.0	80,674.9	149,538.0	0.990	33,211.0	94.7	0.014	0.0
1927	154,951.0	80,599.3	149,389.0	0.989	33,137.0	78.7	0.012	0.0
1928	154,800.0	80,529.5	149,249.0	0.988	33,056.4	89.6	0.014	0.0
1929	154,629.0	80,449.7	149,089.0	0.987	32,967.8	87.0	0.013	0.0
1930	154,449.0	80,367.0	148,922.0	0.986	32,871.0	112.6	0.017	0.0
1931	154,232.0	80,265.2	148,720.0	0.985	32,764.5	100.1	0.015	0.0
1932	154,015.0	80,165.4	148,518.0	0.984	32,648.5	100.3	0.015	0.0

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
1933	153,782.0	80,059.5	148,303.0	0.983	32,521.9	86.2	0.013	0.0
1934	153,547.0	79,954.5	148,086.0	0.981	32,384.0	91.2	0.014	0.0
1935	153,286.0	79,838.9	147,846.0	0.980	32,233.3	98.4	0.015	0.0
1936	152,997.0	79,710.3	147,579.0	0.978	32,068.4	110.9	0.017	0.0
1937	152,673.0	79,564.9	147,279.0	0.977	31,887.2	103.9	0.016	0.0
1938	152,329.0	79,412.5	146,962.0	0.975	31,687.5	84.2	0.013	0.0
1939	151,976.0	79,259.0	146,639.0	0.973	31,466.7	77.0	0.012	0.0
1940	151,597.0	79,095.7	146,292.0	0.971	31,224.2	121.4	0.019	0.0
1941	151,139.0	78,891.9	145,869.0	0.968	30,958.5	130.5	0.020	0.0
1942	150,634.0	78,667.0	145,403.0	0.965	30,670.3	148.0	0.023	0.0
1943	150,072.0	78,414.9	144,884.0	0.962	30,367.4	495.4	0.074	0.0
1944	149,137.0	77,947.7	143,995.0	0.957	30,052.2	970.0	0.141	0.0
1945	147,727.0	77,202.4	142,635.0	0.948	29,722.9	1,637.1	0.231	0.0
1946	145,692.0	76,084.4	140,651.0	0.934	29,384.1	1,171.1	0.172	0.0
1947	144,150.0	75,256.9	139,163.0	0.924	29,074.7	648.8	0.100	0.0
1948	143,120.0	74,739.3	138,188.0	0.917	28,792.2	482.1	0.075	0.0
1949	142,225.0	74,310.5	137,347.0	0.912	28,539.3	377.7	0.060	0.0
1950	141,392.0	73,925.8	136,562.0	0.907	28,334.8	427.2	0.068	0.0
1951	140,467.0	73,490.5	135,682.0	0.902	28,202.9	559.0	0.088	0.0

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
1952	139,384.0	72,956.8	134,635.0	0.895	28,173.9	546.5	0.087	0.0
1953	138,300.0	72,411.3	133,577.0	0.889	28,292.3	473.9	0.076	0.0
1954	137,288.0	71,891.7	132,575.0	0.882	28,602.1	452.2	0.074	0.0
1955	136,319.0	71,373.7	131,596.0	0.876	29,142.4	469.5	0.077	0.0
1956	135,389.0	70,843.1	130,626.0	0.869	29,925.0	606.5	0.099	0.0
1957	134,425.0	70,244.8	129,588.0	0.862	30,880.1	768.0	0.125	0.0
1958	133,465.0	69,585.9	128,516.0	0.854	31,789.4	708.3	0.117	0.0
1959	132,778.0	69,019.3	127,689.0	0.847	32,316.7	678.1	0.113	0.0
1960	132,373.0	68,553.9	127,139.0	0.841	32,250.0	818.8	0.136	0.0
1961	132,100.0	68,120.6	126,757.0	0.836	31,866.5	683.8	0.115	0.0
1962	132,215.0	67,904.5	126,830.0	0.833	31,875.4	765.0	0.129	0.0
1963	132,458.0	67,791.8	127,094.0	0.832	32,822.9	437.6	0.075	0.0
1964	133,175.0	68,007.8	127,825.0	0.835	34,218.0	607.2	0.103	0.0
1965	133,852.0	68,235.6	128,422.0	0.837	35,831.1	261.9	0.045	0.0
1966	135,005.0	68,735.0	129,381.0	0.844	37,641.8	4,293.1	0.553	0.0
1967	132,567.0	66,937.7	126,697.0	0.822	37,538.7	4,830.3	0.614	0.0
1968	130,137.0	65,036.4	124,037.0	0.798	35,199.0	2,392.6	0.366	0.0
1969	130,510.0	64,800.5	124,305.0	0.795	30,148.0	852.2	0.147	0.0
1970	132,932.0	65,670.9	126,546.0	0.806	122,104.0	854.8	0.146	0.0

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
1971	136,418.0	66,690.3	128,934.0	0.818	99,816.4	1,095.5	0.181	0.0
1972	141,484.0	67,657.7	130,794.0	0.830	13,320.1	923.7	0.152	0.0
1973	148,767.0	69,278.7	131,902.0	0.850	11,215.3	1,160.5	0.185	0.0
1974	157,293.0	71,650.7	146,791.0	0.879	15,161.1	907.1	0.144	0.0
1975	165,185.0	75,378.3	162,950.0	0.925	31,862.0	876.6	0.134	0.0
1976	170,154.0	80,320.8	167,719.0	0.986	10,238.1	1,325.4	0.183	0.0
1977	171,765.0	84,976.6	168,324.0	1.043	58,765.3	1,094.3	0.139	0.0
1978	171,608.0	88,238.2	166,600.0	1.083	89,766.7	1,581.1	0.189	0.0
1979	170,321.0	88,952.4	164,736.0	1.092	28,540.3	9,479.0	0.790	0.1
1980	161,806.0	83,875.6	151,073.0	1.029	49,429.6	22,055.2	1.326	0.1
1981	143,006.0	71,703.8	131,750.0	0.880	82,151.5	28,135.3	1.555	0.2
1982	121,213.0	57,240.0	114,399.0	0.703	43,035.3	27,100.1	1.644	0.2
1983	103,336.0	45,335.4	93,702.3	0.556	27,329.0	12,263.6	1.365	0.1
1984	102,409.0	43,126.7	91,703.5	0.529	53,811.9	12,125.2	1.363	0.1
1985	102,307.0	42,439.2	95,704.1	0.521	39,098.8	10,882.4	1.291	0.1
1986	102,854.0	42,776.9	96,829.1	0.525	18,344.0	11,439.1	1.296	0.1
1987	101,995.0	43,097.9	94,394.1	0.529	46,840.9	15,386.7	1.428	0.2
1988	96,484.5	41,257.3	90,863.6	0.506	29,542.4	12,529.8	1.340	0.1
1989	92,739.4	40,060.4	88,075.0	0.492	21,609.6	14,984.0	1.448	0.2

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
1990	85,719.3	37,150.1	79,251.2	0.456	33,375.3	13,764.8	1.443	0.2
1991	79,673.9	34,289.5	74,925.9	0.421	59,765.9	8,982.3	1.268	0.1
1992	78,087.7	33,494.8	73,213.5	0.411	22,334.1	8,679.2	1.263	0.1
1993	76,856.8	32,682.5	70,224.1	0.401	28,020.3	11,405.1	1.411	0.2
1994	73,254.6	30,490.8	65,813.9	0.374	25,293.4	8,896.1	1.325	0.1
1995	72,242.1	29,573.9	68,210.7	0.363	15,743.1	9,252.1	1.365	0.1
1996	70,312.5	28,744.2	66,058.3	0.353	11,571.3	8,347.8	1.323	0.1
1997	68,494.1	28,626.1	65,014.6	0.351	17,708.9	8,879.4	1.345	0.1
1998	65,195.9	28,110.1	62,664.7	0.345	31,771.8	6,605.5	1.198	0.1
1999	63,295.9	28,082.8	60,568.6	0.345	42,041.6	5,957.5	1.131	0.1
2000	61,712.1	27,806.7	57,558.2	0.341	34,645.0	4,779.6	0.992	0.1
2001	61,546.4	27,655.3	55,743.9	0.339	19,104.9	2,318.4	0.628	0.0
2002	64,415.3	28,477.1	58,276.1	0.350	16,975.7	484.4	0.183	0.0
2003	69,416.3	30,395.4	64,782.6	0.373	16,516.2	46.4	0.018	0.0
2004	74,640.7	32,974.4	71,448.3	0.405	54,391.7	99.1	0.036	0.0
2005	79,250.6	35,867.1	75,788.2	0.440	10,517.6	203.3	0.067	0.0
2006	83,414.2	38,639.0	78,813.8	0.474	42,326.2	220.4	0.067	0.0
2007	87,353.8	41,145.7	80,807.3	0.505	10,550.3	244.2	0.069	0.0
2008	91,584.6	43,173.3	87,900.0	0.530	99,446.3	272.1	0.072	0.0

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
2009	95,878.5	45,124.7	89,444.4	0.554	13,077.9	186.1	0.048	0.0
2010	101,109.0	47,066.2	94,748.8	0.578	65,190.9	178.8	0.044	0.0
2011	107,035.0	49,388.7	95,626.4	0.606	9,723.1	212.2	0.049	0.0
2012	113,738.0	51,835.3	109,112.0	0.636	6,447.6	270.8	0.060	0.0
2013	119,724.0	54,937.3	112,627.0	0.674	57,313.9	470.3	0.092	0.0
2014	124,637.0	58,374.1	122,250.0	0.716	23,638.6	722.7	0.132	0.0
2015	127,873.0	61,607.6	123,964.0	0.756	18,329.3	879.8	0.146	0.0
2016	130,134.0	64,260.8	122,724.0	0.789	98,970.8	1,039.4	0.163	0.0
2017	132,291.0	65,863.2	127,220.0	0.808	27,556.6	6,361.5	0.742	0.1
2018	129,454.0	63,933.6	122,344.0	0.785	10,482.1	10,522.8	1.078	0.1
2019	123,043.0	60,054.8	111,974.0	0.737	8,201.8	9,315.3	1.060	0.1
2020	118,467.0	57,126.2	115,125.0	0.701	8,834.6	8,379.6	1.046	0.1
2021	113,776.0	55,346.0	112,130.0	0.679	18,753.2	10,899.7	1.214	0.1
2022	105,127.0	52,883.9	103,484.0	0.649	20,352.7	12,129.7	1.275	0.1
2023	94,075.9	49,359.3	91,819.4	0.606	29,950.8	11,023.5	1.256	0.1
2024	83,492.9	45,045.9	80,049.9	0.553	25,887.9	9,764.1	1.267	0.1
2025	74,415.4	40,183.5	70,404.7	0.493	30,867.0	10,868.1	1.438	0.2
2026	65,168.5	34,267.9	60,347.2	0.421	29,940.6	10,868.1	1.554	0.2
2027	57,339.2	28,679.9	52,688.0	0.352	28,796.8	3,257.0	0.964	0.1

Year	Total Biomass (mt)	Spawning Biomass (mt)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploitation Rate
2028	58,413.1	27,879.0	53,350.2	0.342	28,604.3	3,281.9	0.954	0.1
2029	60,259.6	27,817.3	55,344.7	0.341	28,589.1	3,483.4	0.951	0.1
2030	62,400.8	28,260.4	57,604.9	0.347	28,697.0	3,762.9	0.952	0.1
2031	64,512.9	28,980.0	59,733.5	0.356	28,866.9	4,055.0	0.955	0.1
2032	66,423.2	29,793.7	61,636.2	0.366	29,051.3	4,315.8	0.959	0.1
2033	68,070.9	30,572.9	63,261.2	0.375	29,220.7	4,515.1	0.962	0.1
2034	69,478.3	31,264.2	64,639.7	0.384	29,365.5	4,652.6	0.964	0.1
2035	70,698.9	31,868.4	65,831.2	0.391	29,488.0	4,755.6	0.965	0.1
2036	71,771.8	32,400.2	66,878.1	0.398	29,592.9	4,833.4	0.966	0.1

Table 20: Quantities of interest from the sensitivity analyses.

	Base model	M = 0.1	M = 0.124 female, M = 0.129 male male	MWT asymptotic selex	WCGBTS logistic selex	Including shrimp trawl	New WA catch recon.	Excluding triennial	Francis weighting
Difference from Base Model Likelihood									
Total	7664.49	-7,642.780	-7,659.090	-7,616.870	-7,664.360	-7,664.490	-7,664.490	-7,759.400	-8,891.930
Survey	13.022	-10.413	-12.700	-11.850	-12.784	-13.022	-13.022	-14.298	-12.598
Length	854.968	-853.375	-856.292	-843.347	-843.171	-854.968	-854.968	-947.215	-1,476.972
Age	1366.28	-1,356.610	-1,360.490	-1,330.780	-1,378.150	-1,366.280	-1,366.280	-1,366.710	-1,961.063
Discards	5410.79	-5,410.080	-5,410.630	-5,410.970	-5,410.800	-5,410.790	-5,410.790	-5,410.810	-5,410.740
Recruitment	17.84	-9.918	-17.179	-19.850	-17.791	-17.840	-17.840	-18.630	-29.161
Forecast recruitment	0.601	-0.463	-0.584	-0.664	-0.639	-0.601	-0.601	-0.638	-0.892
Parameter priors	0.983	-1.907	-1.217	0.594	-1.034	-0.983	-0.983	-1.111	-0.495
Parameter values									
Natural mortality (female)	0.125	0.100	0.124	0.152	0.123	0.125	0.125	0.127	0.135
Length at Amin (female)	20.652	20.666	20.707	21.313	20.417	20.652	20.652	20.511	20.685

95

	Base model	M = 0.1	M = 0.124 female, M = 0.129 male	MWT asymptotic selex	WCGBTS logistic selex	Including shrimp trawl	New WA catch recon.	Excluding triennial	Francis weighting
Virgin recruitment (thousands)	34.103	19.899	31.238	56.483	33.014	34.103	34.103	36.393	44.556
ln(R0)	10.437	9.898	10.349	10.942	10.405	10.437	10.437	10.502	10.704
SSB unfished (mt)	151584	139,734.000	145,723.000	174,575.000	150,977.000	151,584.000	151,584.000	156,464.000	170,203.000
SB0 (thousand mt)	81.734	72.835	75.759	88.400	81.910	81.734	81.734	84.482	90.227
SSB 2025 (thousand mt)	40.603	19.951	35.214	62.364	38.632	40.603	40.603	44.929	47.773
B ratio 2025	0.497	0.274	0.465	0.705	0.472	0.497	0.497	0.532	0.529
SPR ratio 2025	1.257	1.657	1.312	0.852	1.308	1.257	1.257	1.193	1.110

Table 21: Estimated data weights derived using the McAllister-Ianelli and Francis methods.

Fleet	Composition data type	McAllister Ianelli weighting (base model)	Francis weighting (alternative)
BottomTrawl	Length	0.050976	0.045830
Hake	Length	0.027980	0.019866
HnL	Length	0.169713	0.147854
MidwaterTrawl	Length	0.043618	0.037046
Net	Length	0.115959	0.121900
NWFSC	Length	0.101232	0.086482
Triennial	Length	0.092531	0.091771
BottomTrawl	Age	0.206741	0.230360
Hake	Age	0.180223	0.236138
HnL	Age	0.517972	0.524997
MidwaterTrawl	Age	0.131374	0.132665
Net	Age	0.179957	0.204433
NWFSC	Age	0.124533	0.129401

Table 22: Projection of potential OFL, landings, and catch, summary biomass (age-4 and older), spawning biomass, and depletion for the base case model projected with total catch equal to the predicted ABC. The predicted OFL is the calculated total catch determined by F SPR=50%

Year	Adopted OFL (mt)	Adopted ACL (mt)	Assumed Catch (mt)	OFL (mt)	Buffer	ABC (mt)	ACL (mt)	Spawning Biomass (mt)	Fraction Unfished
2026			10,868.10	11,382.00	10,391.77	10,392.00		34,267.9	0.421
2027			3,257.04					28,679.9	0.352
2028				3,739.66		3,281.94		27,879.0	0.342
2029				3,990.00		3,483.38		27,817.3	0.341
2030				4,300.89		3,762.86		28,260.4	0.347
2031				4,613.63		4,054.96		28,980.0	0.356
2032				4,879.79		4,315.79		29,793.7	0.366
2033				5,078.83		4,515.05		30,572.9	0.375
2034				5,220.57		4,652.60		31,264.2	0.384
2035				5,324.31		4,755.63		31,868.4	0.391
2036				5,405.10		4,833.43		32,400.2	0.398

Table 23: Summary table of 12-year projections beginning in 2027 for alternate states of nature based on the axis of uncertainty (a combination of M, h, and 2013 recruitment strength). Columns range over low, mid, and high state of nature, and rows range over different assumptions of total catch levels (discards + retained). Catches in 20XX and 20XX are allocated using the percentage of landings for each fleet in 20XX.

			State of nature					
			Low		Base		High	
Management decision	Year	catch (mt)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)
Constant catch (9,000 K)	2027	9,000	17,024	19.5	34,918	42.7	38,407	46.6
	2028	9,007	13,360	15.3	30,977	37.9	34,351	41.7
	2029	9,068	10,428	12.0	27,812	34.0	31,280	38.0
	2030	9,067	7,962	9.1	25,315	31.0	29,056	35.3
	2031	7,419	5,757	6.6	23,321	28.5	27,478	33.4
	2032	6,404	4,389	5.0	21,690	26.5	26,373	32.0
	2033	5,494	3,396	3.9	20,279	24.8	25,570	31.0
	2034	4,662	2,631	3.0	18,981	23.2	24,949	30.3
	2035	3,901	2,019	2.3	17,736	21.7	24,442	29.7
	2036	3,187	1,509	1.7	16,511	20.2	24,011	29.1
ACLP*=0.25, sigma=0.50	2027	1,790	24,754	28.4	29,822	36.5	41,750	50.7
	2028	1,750	24,698	28.3	29,349	35.9	41,257	50.1
	2029	1,791	25,148	28.9	29,596	36.2	41,522	50.4
	2030	1,863	25,952	29.8	30,369	37.2	42,422	51.5
	2031	1,939	26,957	30.9	31,461	38.5	43,786	53.2
	2032	1,991	28,047	32.2	32,692	40.0	45,448	55.2
	2033	1,993	29,155	33.5	33,935	41.5	47,258	57.4
	2034	1,969	30,261	34.7	35,143	43.0	49,121	59.6
	2035	1,929	31,364	36.0	36,298	44.4	50,970	61.9
	2036	1,886	32,473	37.3	37,406	45.8	52,765	64.1

			State of nature					
			Low		Base		High	
Management decision	Year	catch (mt)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)	Spawning biomass (mt)	Depletion (%)
ACLp*=0.45, sigma=0.50	2027	3,231	24,754	28.4	34,918	42.7	41,750	50.7
	2028	3,204	23,939	27.5	33,924	41.5	40,507	49.2
	2029	3,351	23,652	27.1	33,609	41.1	40,062	48.6
	2030	3,574	23,687	27.2	33,792	41.3	40,237	48.8
	2031	3,808	23,860	27.4	34,288	42.0	40,838	49.6
	2032	4,020	24,056	27.6	34,944	42.8	41,693	50.6
	2033	4,180	24,202	27.8	35,642	43.6	42,657	51.8
	2034	4,279	24,280	27.9	36,319	44.4	43,638	53.0
	2035	4,350	24,303	27.9	36,957	45.2	44,593	54.1
	2036	4,392	24,286	27.9	37,550	45.9	45,497	55.2

6 Figures

6.1 Data

6.1.1 Indices




Figure 1: Observations of two cohorts (2008, top and 2010, bottom) from the NWFSC WCGBT survey data. Darker circles indicate more observations (possibly within the same tow).




Figure 2: Fishery-dependent indices of abundance from the 2011 assessment scaled to the mean of their own series.




Figure 3: Survey tow locations in 2004, showing the difference in station design for the NWFSC WCBTS survey relative to the Triennial trawl survey (Figure from Stewart (2007)).




Figure 4: Distribution of dates of operation for the triennial survey (1980-2004). Solid bars show the mean date for each survey year, points represent individual hauls dates, but are jittered to allow better delineation of the distribution of individual points (Figure from (Stewart 2007)).




Figure 5: QQ plots for delta-gamma (panel 1) and delta-lognormal (panel 2, used in base model) sdmTMB index.




Figure 6: Model-based survey estimates for the Triennial with estimated 95% confidence intervals. Based estimates and 95% confidence intervals are shown in blue for comparison.




Figure 7: Expanded length compositions weighted by estimated numbers from the GLMM in each strata for the Triennial survey.




Figure 8: Comparison of the estimated index of relative abundance for the West Coast Groundfish Bottom Trawl Survey using a GLMM (gray line, 2015 stock assessment), VAST (black line, 2019 stock assessment), a delta-gamma sdmTMB index (blue line) and a delta-lognormal sdmTMB index (red line, used in base model). The error bars give 5 and 95% intervals for each survey.




Figure 9: Expanded length compositions for the WCGBTS




Figure 10: Expanded marginal age compositions from the WCGBTS.




Figure 11: Conditional age-at-length from WCGBTS observations for females (red) and males (blue).

6.1.2 Composition data




Figure 12: Expanded length compositions for the bottom trawl fishery. The area of the circle is proportional to the proportion-at-length.




Figure 13: Expanded length compositions for the midwater trawl fishery. The area of the circle is proportional to the proportion-at-length.




Figure 14: Expanded length compositions for the hake fishery. The area of the circle is proportional to the proportion-at-length.




Figure 15: Expanded length compositions for the net fishery. The area of the circle is proportional to the proportion-at-length.




Figure 16: Expanded length compositions for the hook-and-line fishery. The area of the circle is proportional to the proportion-at-length.




Figure 17: Expanded age compositions for the bottom trawl fishery. The area of the circle is proportional to the proportion-at-age.




Figure 18: Expanded age compositions for the midwater trawl fishery. The area of the circles is proportional to the proportion-at-age.




Figure 19: Expanded age compositions for the hake fishery. The area of the circles is proportional to the proportion-at-age.




Figure 20: Expanded age compositions for the net fishery. The area of the circles is proportional to the proportion-at-age.




Figure 21: Expanded age compositions for the hook-and-line fishery. The area of the circles is proportional to the proportion-at-age.




Figure 22: Length compositions for discards from the Pikitch study. The discard length comps were fitted into the model.




Figure 23: Length compositions of the discards for the bottom trawl.

6.1.3 Biological data




Figure 24: Fits to weight-at-length observations (dashed line) for females (left) and males (right) using observations from all data sources. The weight-at-length curve from the 2019 assessment is indicated with a solid line.




Figure 25: Maturity-at-length (left) and maturity-at-age (right) from data reported by Barss & Echeverria (1987). Circles are proportional to the number of observations at that length or age. Lines are estimated logistic curves fitted to the data. The bars at the bottom are the number of samples by each state. The purple line is the estimated maturity-at-age using all data with each state equally weighted, and is used in the assessment model with maturity-at-age for ages 2 and lower set equal to zero.




Figure 26: Number at age observed from all data for female and male widow rockfish.




Figure 27: The prior for natural mortality (black, M , yr $^{-1}$) and the estimated M for females (red) and males (blue) with asymptotic approximation to the sampling distribution. The means of each distribution are shown with triangles.




Figure 28: Length-at-age observations (points, slightly jittered) and predicted length-at-age von Bertalanffy curves for female (top) and male (bottom) widow rockfish collected from all fishery (BDS and At-Sea) and survey (Triennial and NWFSC) data.




Figure 29: Estimated ageing error for the Cooperative Ageing Project lab and the SWFSC.

6.2 Model

6.2.1 Bridging




Figure 30: Data sources by type and year that were used in the base model.




Figure 31: Bridging from the 2019 assessment model with updated data and bridging steps.

6.2.2 Selectivity




Figure 32: Estimated selectivity for different fleets and surveys.




Figure 33: Estimated selectivity curves for 2018 of the hake fleet (topleft), net fishing fleets (topright), the triennial survey (bottomleft), and the NWFSC survey (bottomright).




Figure 34: Estimated selectivity for Bottom Trawl (top row) and Midwater Trawl (bottom row) for males (left column) and females (right column)

6.2.3 Biology




Figure 35: Length at age (top-left panel) with estimated coefficient of variation (CV, thick line) and calculated standard deviation (SD, thin line) versus length at age in the top-right panel and versus age in the lower-left panel.




Figure 36: Estimates of recruitment (upper) and recruitment deviates (lower) with approximate asymptotic 95% confidence intervals (vertical lines) from the MLE estimates.

6.2.4 Fits to data




Figure 37: Estimated and input recruitment bias adjustment ramp. Red line shows current settings for bias adjustment ramp. Red line shows current settings for bias adjustment specified in the model. Blue line shows least squares estimate of alternative bias adjustment relationship for recruitment deviations.




Figure 38: Fits (blue lines) to the abundance estimates (black points) for the base model. A separate q is estimated for the Hake series starting in 1991. Juvenile survey index (in numbers) is on the bottom. 95% confidence intervals are shown in the input standard errors. Thicker lines (if present) indicate input uncertainty before addition of the estimated additional uncertainty parameter.




Figure 39: Predicted (blue line) and observed (open circles) discards for the bottom trawl (top) and midwater trawl (bottom) fleets from the base model. 95% confidence intervals are shown for the observations.




Figure 40: Pearson residuals for fits to length frequency data (left) and age frequency data (right) for landings from the trawl commercial fleets (rows). Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 41: Pearson residuals for fits to length frequency data (left) and age frequency data (right) for landings from the net and hook-and-line commercial fleets (rows). Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 42: Combined length frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).




Figure 43: Pearson residuals for fits to the discard length frequencies from the bottom trawl (left) and hook-and-line (right) fleets. Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 44: Pearson residuals for fits to the triennial survey length frequency data (left) and NWFSC WCGBT (shelf/slope) survey length frequency data (right). Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 45: Combined age frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).




Figure 46: Observed and expected age-at-length with 95% confidence intervals (left) and observed and expected standard deviation of age-at-length with 95% confidence intervals (right) for the NWFSC WCGBT survey data.




Figure 47: Estimated retention curves by time block (if applicable) for each of the five fleets in the model.




Figure 48: Combined age frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).




Figure 49: Combined age frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).

6.2.5 Timeseries




Figure 50: Predicted spawning biomass (thousand mt) for widow rockfish using the base assessment. The solid line is the MLE estimate, and the dashed lines depict the approximate asymptotic 95% confidence intervals.




Figure 51: Predicted summary biomass (age 4+) from the base model.




Figure 52: Predicted relative spawning biomass from the widow rockfish base case assessment. The solid blue line is the MLE estimate, and the dashed lines depict the approximate asymptotic 95% confidence intervals. The red lines show the equilibrium level (100%), the management target of 40% of unfished biomass, and the minimum stock size threshold of 25% of unfished biomass.




Figure 53: Predicted spawning biomass (2011 onward) or spawning output (2000-2009) from past assessments in comparison with the current assessment.




Figure 54: Estimated recruitment and the assumed stock-recruit relationship (black line).
The dashed line shows the effect of the bias correction for the lognormal distribution.




Figure 55: Plot of the summary fishing mortality for each year of the model with 95% confidence intervals.




Figure 56: Phase plot of relative $(1-SPR)/(1-SPR_{50\%})$ (y-axis) and depletion (x-axis) for widow rockfish.

6.3 Model Diganostics

6.3.1 Sensitivity analyses




Figure 57: Estimates of recruitment deviations for sensitivity models.




Figure 58: Spawning biomass (with 95% confidence interval around the base model) for the base model and sensitivity runs. (no rec dev)



Figure 59: Comparision of model estimated quantities for 2025, given as percent of the base model estimated value, by sensitivity run.

6.3.2 Retrospective analysis




Figure 60: Five-year retrospective estimates of spawning biomass (top) and recruitment deviations (bottom).

6.3.3 Likelihood profiles




Figure 61: Likelihood components in the likelihood profile for unfished equilibrium recruitment (R_0)




Figure 62: Likelihood components in the likelihood profile for steepness (h).




Figure 63: Likelihood components in the likelihood profile for natural mortality (M).
Note: male and female natural mortality are set to the same value.




Figure 64: Time series of recruitment estimates for models with different fixed values of natural mortality (M)

6.4 Management




Figure 65: Estimated plot of equilibrium yield vs the fraction of unfished biomass.




Figure 66: Base model, low state of nature, and high state of nature spawning biomass trajectories under three catch scenarios: constant catch of 9000 mt, $ACL = p*0.25$, $ACL = p*0.45$ for 2027 to 2036. The shaded areas indicate the 12.5% and 87.5% lognormal quantiles of spawning biomass.

7 References

- Adams, G., Kapur, M., McQuaw, K., Hamel, O., Stephens, A., and Wetzel, C. 2019. Status of Widow Rockfish (*Sebastodes entomelas*) Along the U.S. West Coast in 2019. Pacific Fishery Management Council, Portland, Oregon.
- Anderson, S.C., Ward, E.J., English, P.A., and Barnett, L.A. 2022. sdmTMB: An R package for fast, flexible, and user-friendly generalized linear mixed effects models with spatial and spatiotemporal random fields. *BioRxiv*: 2022–03.
- Barss, W., and Echeverria, T.W. 1987. Maturity of widow rockfish *Sebastodes entomelas* from the northeastern Pacific, 1977–82. pp. 13–18.
- Bradburn, M.J., Keller, A.A., and Horness, B.H. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition.
- Hamel, O.S., and Cope, J.M. 2022. Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research* **256**: 106477.
- He, X., Pearson, D., Dick, E.J., Field, J., Ralston, S., and MacCall, A. 2011. Status of the widow rockfish resource in 2011. Pacific Fishery Management Council, Portland, OR.
- Helser, T.E., Punt, A.E., and Methot, R.D. 2004. A generalized linear mixed model analysis of a multi-vessel fishery resource survey. *Fisheries Research* **70**(2-3): 251–264.
- Hicks, A., and Wetzel, C. 2015. The Status of Widow Rockfish (*Sebastodes entomelas*) Along the U.S. West Coast in 2015. Pacific Fishery Management Council, Portland, OR.
- Holland, D.S., and Martin, C. 2019. Bycatch quotas, risk pools, and cooperation in the Pacific whiting fishery. *Frontiers in Marine Science* **6**: 600.
- McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling-importance resampling algorithm. *Canadian Journal of Fisheries and Aquatic Sciences* **54**(2): 284–300.
- Methot, R.D., and Taylor, I.G. 2011. Adjusting for bias due to variability of estimated recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(10): 1744–1760.
- Pikitch, E.K., and Rogers, J.B. 1989. Life history characteristics of commercially important groundfish species off California, Oregon and Washington.
- Stewart, I.J., and Hamel, O.S. 2014. Bootstrapping of sample sizes for length-or age-composition data used in stock assessments. *Canadian journal of fisheries and aquatic sciences* **71**(4): 581–588.
- Then, A.Y., Hoenig, J.M., Hall, N.G., Hewitt, D.A., and Handling editor: Ernesto Jardim. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science* **72**(1): 82–92.
- Thorson, J.T., Shelton, A.O., Ward, E.J., and Skaug, H.J. 2015. Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. *ICES Journal of Marine Science* **72**(5): 1297–1310.

- Thorson, J.T., and Ward, E.J. 2014. Accounting for vessel effects when standardizing catch rates from cooperative surveys. *Fisheries Research* **155**: 168–176.
- Weinberg, J.R., Rago, P.J., Wakefield, W.W., and Keith, C. 2002. Estimation of tow distance and spatial heterogeneity using data from inclinometer sensors: An example using a clam survey dredge. *Fisheries Research* **55**(1-3): 49–61.
- Wetzel, C.R., Johnson, K.F., and Hicks, A.C. 2025. nwfscSurvey: Northwest Fisheries Science Center Survey. Available from <https://github.com/pfmc-assessments/nwfscSurvey>.

8 Appendix A: Annual fits to length and age composition




Figure A1: Fits to the retained length compositions for the bottom trawl fleet.




Figure A2: Fits to the discarded length compositions for the bottom trawl fleet.




Figure A3: Fits to the retained length compositions for the midwater trawl fleet.




Figure A4: Fits to the retained length compositions for the hake fleet.




Figure A5: Fits to the retained length compositions for the net fleet.




Figure A6: Fits to the retained length compositions for the hook-and-line fleet.




Figure A7: Fits to the length compositions for the triennial survey.




Figure A8: Fits to the length compositions for the NWFSC WCGBT survey.




Figure A9: Fits to the retained age compositions for the bottom trawl fleet.




Figure A10: Fits to the retained age compositions for the midwater trawl fleet.




Figure A11: Fits to the retained age compositions for the hake fleet.




Figure A12: Fits to the retained age compositions for the net fleet.




Figure A13: Fits to the retained age compositions for the hook-and-line fleet.