

Status of widow rockfish stock off the U.S. West Coast in 2025



Michael Kinneen¹, Maurice C. Goodman¹, Anna Sulc¹, Laurinne Balstad², Raquel Ruiz Diaz¹, Kristina Randrup¹, William Patrone¹, Laura Spencer¹, Alaia Morell³, Alberto Rovellini¹, Allison Dedrick⁴, Nick Grunloh⁵, Madison Bargas⁶, Stephanie Hopkins⁷, Vladlena Gertseva⁸, Kiva L. Oken⁸, Ian G. Taylor⁸, Melissa A. Haltuch⁹ and Owen Hamel⁸

1. School of Aquatic & Fishery Sciences (SAFS), University of Washington, 1122 NE Boat St, Seattle, WA, 98195
2. Department of Environmental Science and Policy, University of California Davis, Wickson Hall, 350 N Quad, Davis, CA, 95616
3. UW Puget Sound Institute, Center for Urban Waters, 326 East D Street, Tacoma, WA, 98421
4. California Department of Fish and Wildlife, 1740 North Market Boulevard, Sacramento, CA, 95834
5. Southwest Fisheries Science Center, 8901 La Jolla Shores Drive, La Jolla, CA, 92037
6. Coastal Oregon Marine Experiment Station, Oregon State University, 430 Strand Agriculture Hall, Corvallis, OR, 97331
7. University of California, Santa Cruz, Institute of Marine Sciences' Fisheries Collaborative Program, 1156 High Street, Santa Cruz, CA, 95064
8. NOAA Fisheries Northwest Fisheries Science Center, 2725 Montlake Boulevard East, Seattle, WA, 98112
9. NOAA Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Building 4, Seattle, WA, 98115



U.S. Department of Commerce
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northwest Fisheries Science Center

Table of contents

Disclaimer	i
Executive summary	ii
Stock	ii
Catches	ii
Data and Assessment	iv
Stock biomass and dynamics	v
Recruitment	viii
Exploitation status	x
Ecosystem considerations	xii
Reference points	xii
Management performance	xiv
Harvest projections and decision table	xiv
Scientific uncertainty	xvi
Unresolved problems and major uncertainties	xvi
Research and Data Needs	xvii
1 Introduction	1
1.1 Distribution and Stock Structure	1
1.2 Life History	1
1.3 Ecosystem Considerations	1
1.4 Fishery description	2
1.5 Management History	3
1.6 Management performance	3
1.7 Fisheries off Canada and Alaska	3
2 Data	4
2.1 Fishery-independent data	4
2.1.1 Triennial Survey	4
2.1.2 West Coast Groundfish Bottom Trawl Survey	4
2.1.3 Juvenile Survey	6
2.2 Fishery-dependent data	7
2.2.1 Landings	7
2.2.2 Fishery catch-per-unit-effort	8
2.2.3 Fishery length and age data	8
2.2.4 Discards	9
2.2.5 Biological data	10
2.3 Environmental and Ecosystem Data	11
3 Assessment model	12
3.1 History of modeling approaches	12
3.2 Responses to most recent past STAR Panel recommendations	12
3.3 Responses to SSC Groundfish Subcommittee requests	12
3.4 Model Structure and Assumptions	12
3.4.1 Changes from the Last Assessment and Model Bridging	12
3.4.2 Modeling Platform and Structure	16
3.4.3 Model Overview	16
3.4.4 Model Parameters	17
3.4.5 Key Assumptions and Structural Choices	19

3.5	Base Model Results	20
3.5.1	Parameter Estimates	21
3.5.2	Fits to the Data	22
3.5.3	Population Trajectory	23
3.6	Model Diagnostics	24
3.6.1	Convergence	24
3.6.2	Parameter Uncertainty	24
3.6.3	Sensitivity Analyses	24
3.6.4	Retrospective Analysis	25
3.6.5	Likelihood Profiles and key parameters	26
4	Management	27
4.1	Reference Points	27
4.2	Unresolved problems and major uncertainties	28
4.3	Harvest Projections and Decision Tables	29
4.4	Evaluation of Scientific Uncertainty	30
4.5	Regional management considerations	30
4.6	Research and Data Needs	30
5	Tables	32
5.1	Management performance	32
5.2	Data	33
5.2.1	Fishery-dependent data	33
5.2.2	Fishery-independent data	48
5.3	Model results	51
5.4	Model diagnostics	70
5.5	Management	71
6	Figures	73
6.1	Data	73
6.1.1	Indices	74
6.1.2	Composition data	79
6.1.3	Biological data	83
6.2	Model	84
6.2.1	Bridging	84
6.2.2	Selectivity	98
6.2.3	Biology	100
6.2.4	Fits to data	102
6.2.5	Time series	110
6.3	Model Diagnostics	115
6.3.1	Sensitivity analyses	115
6.3.2	Retrospective analysis	119
6.3.3	Likelihood profiles	121
6.4	Management	126

7	References	127
---	------------	-----

Please cite this publication as:

Kinneen, M., Goodman, M. C., Sulc, A., Balstad, L., Diaz, R., Randrup, K., Patrone, W., Spencer, L., Morell, A., Rovellini, A., Dedrick, A., Grunloh, N., Bargas, M., Hopkins, S., Gersteva, V., Oken, K., Taylor, I., Haltuch, M., & Hamel, O. (2025) Status of widow rockfish stock off the U.S. West Coast in 2025. Pacific Fishery Management Council, Portland, Oregon.

Disclaimer

These materials do not constitute a formal publication and are for information only. They are in a pre-review, pre-decisional state and should not be formally cited or reproduced. They are to be considered provisional and do not represent any determination or policy of NOAA or the Department of Commerce.

Executive summary

Stock

This is an assessment of widow rockfish (*Sebastes entomelas*) that reside in the waters off California, Oregon, and Washington from the U.S.-Canada border in the north to the U.S.-Mexico border in the south. Widow rockfish inhabit water depths of 25–370 m from northern Baja California, Mexico to Southeastern Alaska. Although catches north of the U.S.-Canada border and south of the U.S.-Mexico border were not included in this assessment, it is not certain if those populations contribute to the biomass of widow rockfish off of the U.S. West Coast, possibly through adult migration and/or larval dispersion. Following the 2015 benchmark assessment (Hicks and Wetzel 2015), this update assessment is based on a single coastwide area model.

Catches

Historically, fisheries have caught widow rockfish since the turn of the 20th century. Landings in the trawl fishery are estimated to have increased into the 1940s and remained relatively constant and small (below 1,000 mt per year) throughout the 1950s and into the 1960s before catches increased from a foreign trawl fleet in the 1970s, with a peak at almost 5,000 mt in 1967. Catches by a midwater trawl fleet increased rapidly in the late 1970s following the discovery that widow rockfish form large aggregations at night.

Total landings of widow rockfish peaked in the early 1980s, increasing from approximately 1,000 metric tons (mt) in 1978 to over 25,000 mt in 1981. The large landings in the early 1980s were curtailed with trip limits beginning in 1982, which resulted in a decline in landings throughout the 1980s and 1990s following sequential reductions in the trip limits. From 2000 to 2003, landings of widow rockfish dropped from over 4,000 mt to about 40 mt and remained low through 2016. Landings increased rapidly following the quota share reallocation in 2017, and have been near or above 10,000 mt in all years between 2018 and 2024. Midwater trawl gears targeting rockfish and bycatch in the Pacific hake/whiting (*Merluccius productus*, hereafter “hake”) fisheries account for the majority of the recent catch.

Widow rockfish are a desirable market species and it is believed that discarding was low historically. However, management restrictions (e.g., trip limits) resulted in a substantial amount of discarding beginning in the early 1980s. Trawl rationalization was introduced in 2011. Between 2011 and 2024 very little discarding of widow rockfish has occurred. Catches for the past ten years are in Table i and for the entire time series in Figure i.

Table i: Recent total catch for the bottom trawl, midwater trawl, hake fleets. Catch combines landings and discards. The bottom trawl catch includes catches from fleets that were previously separated as net, and hook-and-line fisheries as described in the model bridging section.

Year	BottomTrawl (mt)	MidwaterTrawl (mt)	Hake (mt)	Total Catch (mt)
2015	14.6	480.1	386.2	880.8
2016	11.3	589.6	440.8	1,041.7
2017	39.3	4,861.9	1,455.2	6,356.3
2018	39.2	9,411.5	1,081.3	10,532.0
2019	30.6	8,177.4	1,101.6	9,309.6
2020	76.6	7,577.7	746.7	8,401.0
2021	108.4	10,147.9	617.3	10,873.6
2022	136.4	10,860.9	1,119.0	12,116.3
2023	89.7	10,234.1	673.2	10,996.9
2024	40.9	9,172.1	533.8	9,746.8




Figure i: Total catches (landings and discards combined) of widow rockfish from 1916 to 2024 for bottom trawl, midwater trawl, and hake fleets.

Data and Assessment

The most recent benchmark assessment was conducted in 2015 (Hicks and Wetzel 2015) which was then updated in 2019 (Adams et al. 2019). This assessment began as a second update of the 2015 benchmark stock assessment and was initially reviewed by the Pacific Fishery Management Council (PFMC) Science and Statistical Committee (SSC) Groundfish Subcommittee (GFSC) in August 2025. The GFSC requested additional explorations and changes to be reviewed in October 2025 and then again in January 2026 due to impacts of the 2025 United States federal government shutdown on the October review. Many of the changes made after the August 20205 model are outside the scope of the update assessment terms of reference, making this assessment closer to a benchmark assessment.

This assessment uses the length- and age-structured modeling software Stock Synthesis (version 3.30.24.1). The coastwide population was modeled assuming separate growth and mortality parameters for females and males from 1916 to 2024.

The fleet structure in this assessment was streamlined compared to the 2015 benchmark and the 2019 update assessments to focus on major fisheries that remove more than 97% of the catch. Those include: 1) a coastwide shore-based bottom trawl fleet (1916–2024), 2) a coastwide shore-based midwater trawl fleet that targets rockfish (1979–2024), and 3) a mostly-midwater-trawl fleet that targets hake and includes an at-sea fleet (1975–2024) and a domestic shore-based fleet (1991–2024). The bottom trawl fleet includes minimal catches made by net and hook-and-line gears, as well as widow rockfish catches made within historical foreign fishery targeting Pacific Ocean Perch between 1966–1976. The model includes catch, age, and length data for all three fishing fleets. Catch in each fleet represents total removals. The discard amount for bottom trawl and midwater trawl fleets were reconstructed outside the model and added to the landings within the corresponding fleets.

There are three historical fishery-dependent CPUE indices retained from the 2015 benchmark: 1) Oregon bottom trawl (1984–1999), 2) at-sea foreign hake (1977–1988), and 3) at-sea domestic hake fleets (1983–1998). Data from three fishery-independent surveys were also included in the model: 1) length composition and an index for the Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey (Triennial Survey) were retained from the 2015 benchmark (1977–2004), 2) length and conditional age-at-length compositions and an index for the Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (WCGBTS) (2003–2024), and 3) a recruitment index from the National Marine Fisheries Service (NMFS) SWFSC and NWFSC/PWCC Midwater Trawl Survey (Juvenile Survey) (2004–2024).

The base model estimated parameters for length-based selectivity for all fleets and surveys, a sex-specific length-at-age relationship, sex-specific natural mortality, and recruitment deviations. A Beverton-Holt stock-recruitment function was used to model productivity, and the steepness parameter was fixed at 0.72 based on a steepness meta-analysis (Thorson et al. 2019) for west coast rockfishes. The model also uses widow rockfish fecundity-at-length

relationships from Dick et al. (2017) instead of assuming spawning output being proportional to body weight as in 2015 and 2019 models.

Natural mortality is the major source of uncertainty that is proposed as the axis of uncertainty to define a states of nature in a decision table.

Stock biomass and dynamics

The time series of estimated spawning output and the fraction of unfished spawning output are in Figure ii and Figure iii, respectively, and for the most recent year in Table ii. The spawning output declined rapidly with the developing domestic midwater fishery in the late 1970s and early 1980s, and remained low until 2000. A combination of strong recruitment and low catches resulted in a steady increase in spawning output from 2000 through 2016. A target fishery for widow rockfish was reestablished in 2017. The stock exhibits a decline in most recent years with the increased catches since 2017.

The 2025 spawning output relative to unfished equilibrium spawning output is 50%, above the target of 40% (95% confidence interval of 30 - 70%).

Spawning output is estimated to be at 10,572 billion eggs in 2025 (95% confidence interval of 5,127 - 16,017 billions of eggs). The uncertainty in the estimated spawning output is high, especially in the early years.




Figure ii: Estimated female spawning output time-series from the base model (solid line) with asymptotic 95% confidence interval (dashed lines).




Figure iii: Estimated relative spawning output (depletion) with asymptotic 95% confidence interval (dashed lines) for the base model.

Table ii: Recent trend in estimated female spawning output (billions of eggs) and fraction of unfished spawning output (relative spawning output).

Year	Spawning output	Lower Interval (mt)	Upper Interval (mt)	Fraction Unfished	Lower Interval	Upper Interval
2015	14,522	10,011	19,034	0.688	0.532	0.844
2016	15,089	10,522	19,655	0.715	0.560	0.869
2017	15,436	10,876	19,996	0.731	0.580	0.882
2018	14,925	10,427	19,424	0.707	0.560	0.854
2019	13,816	9,386	18,247	0.654	0.510	0.799
2020	12,948	8,537	17,359	0.613	0.469	0.758
2021	12,457	7,967	16,947	0.590	0.442	0.738
2022	12,033	7,340	16,726	0.570	0.412	0.728
2023	11,644	6,668	16,621	0.552	0.380	0.723
2024	11,195	5,945	16,444	0.530	0.345	0.716
2025	10,572	5,127	16,017	0.501	0.304	0.698

Recruitment

Recruitment deviations were estimated for the entire time series modeled. There is little information regarding recruitment prior to 1965, and the uncertainty in these estimates is expressed in the model. Estimates of recruitment appear to be episodic and characterized by periods of low recruitment, with several very large, but uncertain, year classes (Figure iv, Figure v).

The 2019 update assessment (Adams et al. 2019) estimated the largest recruitment on record in 2013, and above-average recruitment in 2014. With more parsimonious fleet structure in this assessment, and the additional years of data, 2013 is now estimated as only the tenth-largest recruitment on record, and 2014 is estimated as below average.




Figure iv: Time-series of estimated recruitments (medians as open circles) for the base model with asymptotic 95% confidence interval (vertical bars). Estimated unfished equilibrium recruitment (R_0) is indicated with a filled circle.

Table iii: Recent estimated trend in widow rockfish recruitment with asymptotic 95% confidence intervals determined from the base model.

Year	Recruitment (1,000s)	Lower Interval (1,000s)	Upper Interval (1,000s)	Recruitment Deviations	Lower Interval	Upper Interval
2015	21,973	11,088	43,542	-0.412	-1.010	0.186
2016	126,249	77,149	206,597	1.331	1.014	1.648
2017	47,765	24,880	91,701	0.339	-0.191	0.868
2018	17,416	7,712	39,328	-0.690	-1.426	0.046
2019	16,236	6,719	39,237	-0.774	-1.596	0.048
2020	22,868	8,487	61,615	-0.446	-1.414	0.522
2021	37,217	13,346	103,783	0.023	-1.019	1.064
2022	37,214	12,833	107,917	0.004	-1.091	1.099
2023	48,077	16,299	141,815	0.248	-0.868	1.365
2024	42,674	14,564	125,041	0.136	-0.970	1.241

Exploitation status

The population declined throughout the 1980s and 1990s when fishing intensity was above the management target. The population then increased between 2000 and 2016 when fishing intensity was well below the management target. Fishing intensity has increased substantially since 2017, and is estimated to have been above the management target since 2018, though the relative spawning output remains above the management target of 40%. Recent fishing intensities are in Table iv, and a full time series of relative spawning output and fishing intensity is in Figure v.

Table iv: Estimated recent trend in relative fishing intensity ($((1-SPR)/(1-SPR_{50\%}))$) and exploitation rate (as the proportion of age 4+ biomass) with asymptotic 95% confidence intervals for both quantities.

Year	$(1-SPR)/(1-SPR_{50\%})$	Lower Interval (SPR)	Upper Interval (SPR)	Exploitation Rate	Lower Interval (Rate)	Upper Interval (Rate)
2015	0.160	0.104	0.217	0.007	0.005	0.010
2016	0.182	0.120	0.244	0.009	0.006	0.011
2017	0.789	0.600	0.978	0.053	0.038	0.068
2018	1.151	0.930	1.371	0.092	0.064	0.119
2019	1.141	0.907	1.375	0.089	0.061	0.118
2020	1.127	0.878	1.375	0.077	0.050	0.103
2021	1.277	1.016	1.537	0.098	0.062	0.134
2022	1.301	1.023	1.579	0.115	0.069	0.161
2023	1.238	0.928	1.548	0.114	0.063	0.164
2024	1.214	0.870	1.558	0.110	0.057	0.164




Figure v: Phase plot of relative fishing intensity ($(1-SPR)/(1-SPR_{50\%})$) versus fraction unfished (spawning output, B/B_0). Lines through the final point show marginal asymptotic 95% intervals for each dimension. The shaded ellipse is an approximate 95% confidence region which accounts for the estimated covariance between the two quantities.

Ecosystem considerations

Recruitment is a key mechanism by which the ecosystem may directly impact the population dynamics of widow rockfish; however, the specific pathways through which environmental conditions exert influence on widow rockfish dynamics are unclear. Changes in the environment may result in changes in other population processes, as well. Unfortunately, there are few data available for widow rockfish that provide insights into these effects.

Fishing has effects on habitats, in addition to the population itself. Rockfish are often associated with habitats containing living structures such as sponges and corals which fishing may threaten. Recent studies on essential fish habitat are beginning to characterize important locations for rockfish throughout their life history; however there is little current information available to evaluate the specific effects of fishing on the ecosystem characteristics pertinent to the management of widow rockfish.

Reference points

Reference points were calculated using the estimated selectivity parameters and average catch distribution among fleets in the most recent five years of the model (2019-2024). A list of estimates of the current state of the population, as well as reference points based on 1) a target unfished spawning output of 40%, 2) a spawning potential ratio (SPR) of 0.5, and 3) the model estimate of maximum sustainable yield (MSY), are all listed in Table [v](#). The SPR is expected lifetime reproductive output under a given fishing intensity as a proportion of the unfished expected lifetime reproductive output.

While the estimate of population scale (total spawning output) has remained relatively steady across the 2015, 2019, and 2025 assessments, unfished recruitment (R_0) and natural mortality (M), which are positively correlated, have declined from 2015 (female M of 0.157 yr^{-1}) to 2019 (0.144 yr^{-1}) to 2025 (0.135 yr^{-1}). Thus, although the population scale is estimated consistently, additional years of data (in particular, age composition data) following the re-opening of the fishery have led the model to estimate that the population is less productive, which results in lower yields at all three reference points (proxy based on spawning output, proxy based on fishing intensity, and model estimate of MSY).

Table v: Summary of reference points and management quantities for the base model.

Reference Point	Estimate	Lower Interval	Upper Interval
Unfished Spawning output (billions of eggs)	21,110.8	17,760.2	24,461.4
Unfished Age 4+ Biomass (mt)	156,376.0	131,787.1	180,964.9
Unfished Recruitment (R0)	40,470.3	28,185.9	52,754.7
2025 Spawning output (billions of eggs)	10,571.7	5,126.8	16,016.6
2025 Fraction Unfished	0.501	0.304	0.698
Reference Points Based SO40%			
Proxy Spawning output (billions of eggs) SO40%	8,444.3	7,104.1	9,784.6
SPR Resulting in SO40%	0.458	0.458	0.458
Exploitation Rate Resulting in SO40%	0.083	0.076	0.091
Yield with SPR Based On SO40% (mt)	6,238.8	4,973.6	7,504.1
Reference Points Based on SPR Proxy for MSY			
Proxy Spawning output (billions of eggs) (SPR50)	9,418.7	7,923.8	10,913.6
SPR50	0.5		
Exploitation Rate Corresponding to SPR50	0.073	0.067	0.079
Yield with SPR50 at SO SPR (mt)	5,930.5	4,732.1	7,128.9
Reference Points Based on Estimated MSY Values			
Spawning output (billions of eggs) at MSY (SO MSY)	5,393.1	4,540.5	6,245.8
SPR MSY	0.328	0.324	0.332
Exploitation Rate Corresponding to SPR MSY	0.127	0.116	0.138
MSY (mt)	6,745.5	5,358.5	8,132.4

Management performance

Since annual catch limits for widow rockfish increased in 2017 and new fishing opportunities for use of midwater trawl gear became available, attainment of the annual catch limit has been high (Table [vi](#)). Specifically, attainment of the annual catch limit (ACL) has exceeded 70% every year since 2018, averaged 77% from 2017-2024, and was as high as 88% in 2022.

Table vi: Recent trend in total catch and commercial landings (mt) relative to the management guidelines. Estimated total catch reflects the commercial landings plus the model estimated dead discarded biomass.

Year	OFL (mt)	ABC (mt)	ACL (mt)	Catch (mt)
2015	4,137	3,929	2,000	881
2016	3,990	3,790	2,000	1,042
2017	14,130	13,508	13,508	6,356
2018	13,237	12,655	12,655	10,532
2019	12,375	11,830	11,831	9,310
2020	11,714	11,199	11,199	8,401
2021	15,749	14,725	14,725	10,874
2022	14,826	13,788	13,788	12,116
2023	13,633	12,624	12,624	10,997
2024	12,453	11,482	11,482	9,747

Harvest projections and decision table

Projections for overfishing limit (OFL) (mt), ABC (mt), ACL (mt), the buffer between the OFL and ABC, estimated spawning output (billions of eggs), and fraction of unfished spawning output with adopted OFL and ACL and assumed catch for the first two years of the projection period. The predicted OFL is the calculated total catch determined by FSPR=50% are provided in Table [vii](#).

Table vii: Potential overfishing limit (OFL) (mt), Acceptable Biological Catch (ABC) (mt), ACL (mt), the buffer between the OFL and ABC, estimated spawning output (billions of eggs), and fraction of unfished spawning output with adopted OFL and ACL and assumed catch for the first two years of the projection period. The predicted OFL is the calculated total catch determined by FSPR=50%.

Year	Adopted OFL (mt)	Adopted ACL (mt)	Assumed Catch (mt)	OFL (mt)	Buffer	ABC (mt)	ACL (mt)	Spawning output (billions of eggs)	Fraction Unfished
2025	12,254	11,237	10,669	-	-	-	-	10,571.70	0.501
2026	11,382	10,392	9,824	-	-	-	-	9,630.18	0.456
2027	-	-	-	4,916	0.935	4,596	4,596	8,799.27	0.417
2028	-	-	-	5,172	0.93	4,810	4,810	8,763.01	0.415
2029	-	-	-	5,548	0.926	5,137	5,137	8,869.58	0.420
2030	-	-	-	5,866	0.922	5,409	5,409	9,054.47	0.429
2031	-	-	-	6,047	0.917	5,545	5,545	9,245.68	0.438
2032	-	-	-	6,117	0.913	5,585	5,585	9,403.42	0.445
2033	-	-	-	6,130	0.909	5,572	5,572	9,520.54	0.451
2034	-	-	-	6,123	0.904	5,535	5,535	9,605.99	0.455
2035	-	-	-	6,116	0.9	5,504	5,504	9,671.68	0.458
2036	-	-	-	6,116	0.896	5,480	5,480	9,725.54	0.461

Uncertainty in natural mortality (estimated for both sexes) contributed greatly to uncertainty in the results. Natural mortality is proposed to be used as the axis of uncertainty to define low and high states of nature for the decision table. This differed from the 2019 assessment which included two more factors, steepness (not estimated in the model) and 2013 recruitment strength. Steepness was fixed in the base model at the mean of a prior has not changed since 2019, and the 2013 year class is no longer a major source of uncertainty, and there is no recent similarly large estimate of recruitment. The 12.5% and 87.5% quantiles for female and male natural mortality (independently) are proposed to be used as low and high values (0.126 yr^{-1} and 0.144 yr^{-1} for females; 0.137 yr^{-1} and 0.157 yr^{-1} for males). The decision table with projections under alternative catch streams is to be developed during the review meeting.

Scientific uncertainty

The model estimate of the log-scale standard deviation of the OFL in 2025 is 0.293. This is less than the default SSC value of 0.5 for a category 1 assessment, so harvest projections assume an initial sigma of 0.5.

Unresolved problems and major uncertainties

Major sources of uncertainty include natural mortality, stock productivity, and spatial dynamics which are discussed below.

This assessment attempts to capture uncertainty in M by estimating it inside the model (thereby propagating uncertainty in M into uncertainty in spawning output and other derived quantities), and by incorporating variation in M in a decision table. Model sensitivities and profiles over M showed that current stock status was highly sensitive to the assumption about natural mortality.

Widow rockfish is a relatively long-lived fish, and their natural mortality (M) is likely to be lower than many managed fish stocks (e.g., gadoids). Ages above 50 years have been observed and it is expected that natural mortality could be less than 0.10 yr^{-1} (the median of the prior for natural mortality used in this assessment). However, even with length and age data available back to the late 1970s, M was estimated at 0.135 for females and 0.147 yr^{-1} for males, with a small amount of uncertainty (e.g., a 6.1% coefficient of variation for females).

Notably, the estimated M for both sexes from this assessment are lower than those from the 2015 and 2019 assessments, for which natural mortality was estimated above 0.15 yr^{-1} and 0.14 yr^{-1} , respectively. The estimates of M varied slightly depending on the weight given to age and length data, or removing recent years of data, but M was always estimated above 0.12 yr^{-1} . The likelihood profile over natural mortality provides support for values up to or above 0.14 yr^{-1} , but with greater curvature than in the 2019 assessment, suggesting additional data has reduced the support for higher natural mortality values. A contributing factor to this change is the increased mean age and frequency of older fish in catch observed by the WCGBTS survey in recent years. The successful rebounding of the stock as a result of reduced fishing pressure from 2003-2016 allowed more individuals to reach older ages than observed in the earlier years of the WCGBTS. However, consideration should be given in future assessments to structural changes which may improve the fits to WCGBTS age data.

Steepness was fixed at 0.720 in the base model, but a likelihood profile showed that it would be estimated at a value less than that. Estimates of M increased with lower steepness, while unfished spawning output increased and current spawning output decreased. Sustainable yields at the $\text{SPR}_{50\%}$ reference harvest rate ranged from approximately 2641 to 5992 mt depending on the value of steepness.

A greater understanding of the spatial dynamics of widow rockfish would help with interpretation of patterns in the existing data and help improve future assessment models. In particular, there are opposing trends in the age distribution sampled in recent years from the target fishery which is concentrated in a subset of the range of the stock and the coastwide age distribution WCGBTS. These differences could be due to some combination of localized depletion, spatial variability in age relative to benthic vs pelagic habitat, spatial distribution relative to untrawlable habitat, and the rate of mixing among areas.

Research and Data Needs

There are many areas of research that could be improved to benefit the understanding and assessment of widow rockfish, including the following:

- **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for widow rockfish. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life-history of widow rockfish may reduce that uncertainty. Investigating the ageing error and bias would help to understand the influences that the age data have on this assessment, particularly the influence of age data on natural mortality.
- **Sex-specific selectivity:** The midwater and bottom trawl length-composition data fits showed divergent residual patterns between male and female fish. The underlying mechanism driving this pattern is unclear, and could be related to growth, sexing error, or to sex-specific selectivity (e.g., when widow rockfish aggregate, sexes possibly may be aggregating separately). Sex-specific selectivity for these two fleets could be explored or included to address this.
- **Coastwide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for widow rockfish off of the west coast of the U.S. and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and Alaska to determine if there are similarities with the U.S. West Coast observations would help to define the connectivity between widow rockfish north and south of the U.S.-Canada border. Research into spatial dynamics within the U.S. west coast (as noted above under Unresolved problems and major uncertainties), is also important.

1 Introduction

This is an assessment of widow rockfish that inhabit the waters off California, Oregon, and Washington from the U.S.-Canadian border in the north to the U.S.-Mexico border in the south, and does not include Puget Sound.

Sebastes entomelas (widow rockfish) is named after its black-lined gut cavity (*ento* meaning within and *melas* meaning black). It has been referred to as buda, beccafico (Italian bird), and viuva (widow) prior to the 1930s. More recently, the widow rockfish is also called brownie, belinda bass, brown bomber, and soft brown.

1.1 Distribution and Stock Structure

Widow rockfish inhabit water depths of 25–370 m from northern Baja California, Mexico to Southeastern Alaska, and are most abundant from British Columbia to Northern California. Although catches north of the U.S.-Canada border or south of the U.S.-Mexico border were not included in this assessment, it is possible that these populations contribute to the biomass of widow rockfish off of the U.S. West Coast through adult migration and/or larval dispersion.

This assessment is based on a single coastwide area model. There is little evidence of genetically separate stocks along the U.S. coast and past assessments have used a single area versus a two-area assessment model and results were found to be similar (He et al. 2011). There is some evidence of biological differences between areas. For example, widow rockfish collected off California tend to mature at a smaller length than widow rockfish collected off of Oregon (Barss and Echeverria 1987). This could be due to environmental or anthropogenic effects rather than genetic differences.

1.2 Life History

This section is not required for an update assessment; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) for information.

1.3 Ecosystem Considerations

This section is not required for an update assessment; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) for information.

1.4 Fishery description

Widow rockfish were lightly exploited by bottom trawl and hook-and-line gears prior to the 1980s. After many attempts to start trawl fisheries off the west coast of the United States in the late 1800s, the availability of otter trawl nets and the diesel engine in the mid-1920s helped trawl fisheries expand (Douglas et al. 1998). The trawl fisheries really became established during World War II when demand increased for shark livers and bottomfish. A mink food fishery also developed during World War II (Jones and Harry Jr 1960). Foreign fleets began fishing for rockfish in the mid-1960s until the EEZ was implemented in 1977 (Rogers 2003). Longline catches of widow rockfish are present from the turn of the century and continue in recent years, mainly from fisheries targeting sablefish and halibut.

In the late 1960s and early 1970s, it is reported that foreign fishing vessels caught large numbers of widow rockfish (Rogers 2003). In the late 1970s a domestic midwater trawl fishery began developing off of Oregon when it was realized that widow rockfish form dense aggregations at night (Gunderson 1984). The fishery expanded very quickly, with landings from trawl, net, and hook-and-line gears increasing more than 20 times by the early 1980s. As early as 1982, trip limits were imposed to keep catches below recommended annual levels. Trip limits became more restrictive over the years until widow rockfish was declared overfished in 2001. In 2002, harvest guidelines were greatly reduced resulting in decreased catch. Since 2017, catches have steadily increased to levels approximately equal to those in the late 1980's and early 1990's. Widow rockfish catch history by fleet is shown in Figure 6.

The majority of current catches are taken by the midwater trawl fleet, though a significant portion is caught as bycatch by the fishery targeting hake. While historically a major fleet, the catches from the bottom trawl gears have declined due to a combination of regulatory changes, fleet consolidation and reduced ex-vessel value (Jeff Lackey, pers. comm.). The spatial efforts of the midwater trawl fleet has also shifted significantly in recent years. Currently, fishing is concentrated around central Oregon and primarily off the shelf, with the fleet focusing on several well established primary and secondary spots. This change in distribution of efforts is largely due to the fleet attempting to avoid bycatch, reduce catches of undersized fish and minimize distance from processing facilities in central Oregon. If mid-water vessels have remaining yellowtail rockfish quota later in the season, they may be more inclined to target some of the Washington grounds where a mix of widow and yellowtail rockfishes is found. The industry has also noted that the fleet is dynamic throughout the season, with fishers responding to changes in oceanographic conditions, feed distribution and target species distribution (Jeff Lackey, pers. comm.).

Historical discarding practices are not well known, but it is believed that little discarding occurred prior to management restrictions. With the introduction of trip limits, limited data from the mid-1980s show occasional very high discard rates of widow rockfish from tows that occurred near the end of a trip. In recent years, discards from all fleets have been low,

likely due to a combination of changes in fleet behavior, increased observer coverage and the introduction of the IFQ system in 2011.

1.5 Management History

Management history prior to 2015 is detailed in the most recent benchmark assessment (Hicks and Wetzel 2015). In 2017, the NMFS implemented a quota share (QS) reallocation rule, which re-established a target fishery for widow rockfish by allocating quotas among permit holders based on historical allocations, removing daily vessel limits, and allowing the trading of QS (NMFS 2017).

1.6 Management performance

Table 8 shows that recent catches have been below recommended catch levels. Total mortality estimates from the West Coast Groundfish Observer Program (WCGOP) and from the stock assessment might differ due to the use of different estimation methods. Investigation into how these methods differ is beyond the scope of an update assessment.

1.7 Fisheries off Canada and Alaska

This section is not required for an update assessment; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) for information.

2 Data

Many sources of data were available for this assessment (Figure 7), including indices of abundance, landings, discards, and length and age observations from fishery-dependent and fishery-independent sources. Data used in this assessment are described below. No new data sources were considered in this update assessment.

2.1 Fishery-independent data

Data from three fishery-independent surveys were used in this assessment: 1) the SWFSC and NWFSC/PWCC Midwater Trawl Survey (hereafter, “juvenile survey”); 2) the Alaska Fisheries Science Center (AFSC)/NWFSC Triennial Shelf Trawl Survey (hereafter, “triennial survey”); and 3) the NWFSC West Coast Groundfish Bottom Trawl Survey (hereafter, “WCGBTS”). Depth and latitude strata used to analyze the catch-rates, length compositions, and age compositions are the same as in the 2019 update assessment (Adams et al. 2019) and shown in Table 15.

2.1.1 Triennial Survey

The Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey (Triennial Survey) was first conducted by the Alaska Fisheries Science Center (AFSC) in 1977 and was conducted every three years, ending 2004. The survey's design and sampling methods are most recently described in (Weinberg et al. 2002).

The time series suggests a possible slightly increasing trend in biomass from 1980–1983, although is relatively flat until the end of the period in 2001 and 2004 when the index declines significantly. The index and length compositions for this historical Triennial survey were unchanged from the previous assessment. Please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) and most recent update assessment (Adams et al. 2019) for details on those data inputs.

2.1.2 West Coast Groundfish Bottom Trawl Survey

The Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (WCGBTS) has been conducted annually, since 2003. It is based on a random-grid design; covering the coastal waters between depths of 55–1,280 m (Keller et al. 2017). No survey occurred in 2020 due to the COVID-19 pandemic. Widow rockfish are not commonly caught in the

WCG BTS. Higher catch rates occur north of 40° N latitude and catches are rare south of 36° N latitude. Few large fish are found shallower than 100 m and few small fish are found in the deeper water of the slope. There is no clear trend in length with latitude other than smaller fish tend to occur south of approximately 36° N latitude, and there appears to be some very small fish found near 39° N latitude.

For this assessment, geostatistical models of biomass density were fit to survey data using the R package [Species Distribution Models with Template Model Builder](#) (`sdmTMB`) (Anderson et al. 2022). This is an updated approach compared to the 2015 benchmark assessment (non-spatial delta-GLMM) and the 2019 update assessment (VAST delta-lognormal model). The `sdmTMB` model used a 200 knot mesh of the survey area (Thorson et al. 2015). The prediction grid was truncated to only include available survey locations in depths between 55–500 m to limit extrapolating beyond the data and edge effects. Tweedie, delta-binomial, delta-gamma, and mixture distributions, which allow for extreme catch events, were investigated. The positive catch weight model includes survey pass ('first' for early season or 'second' for late season) and year. Vessel-year effects, which have traditionally been included in index standardization for this survey, were not included as the estimated variance for the random effect was close to zero. Vessel-year effects were more prominent when models did not include spatial effects and were included for each unique combination of vessel and year in the data to account for the random selection of commercial vessels used during sampling (Helser et al. 2004; Thorson and Ward 2014).

Results are shown for the delta-gamma and delta-lognormal distributions, which reported the best diagnostics among the explored models (Figure 9). Both models converged (positive definite Hessian matrix) but predicted data from both models showed slightly right-heavy tails compared to the assumed likelihoods, with the gamma model having stronger divergence. Spatiotemporal estimates of biomass from the delta-lognormal model were then converted into annual indices using the `sdmtmb::get_index()` function by integrating across the spatial domain of the survey (Anderson et al. 2022).

Overall, the delta-lognormal index estimates is more comparable to the 2019 spatiotemporal VAST-based index than the delta-gamma index, and less influenced by extreme catch events, particularly in 2013 and 2016; for these reasons, in addition to better model performance observed above, the delta-lognormal `sdmTMB`-based index was used for the base model in this assessment. The delta-lognormal mean value (2262.824) was slightly lower than the means of the index values used in the 2015 benchmark assessment (2701.12) and the 2019 update assessment (3301.765). However, since these are used as relative indices, these differences in mean values have no impact in themselves on the outcome of the assessment. Comparisons of the different error structures, design-based estimate and the VAST index used in 2019 are in Figure 10.

Length, age, and conditional age-at-length compositions were processed using the `nwfscSurvey` package in R publicly available on GitHub (Wetzel et al. 2025). Length compositions were created by expanding to the tow and summing to give a strata specific composition

Table 13. Strata definition was retained from the benchmark assessment. The strata compositions were combined to a coastwide composition using a design-based index of abundance from each stratum. The design based index is constructed by taking the average catch per unit effort (CPUE) defined as catch per area swept across tows in each stratum and year. The sum of strata specific composition data was then calculated, weighting by the average CPUE per stratum multiplied by the area of each stratum. Age distributions were included in the model as conditional-age-at-length (CAAL) observations. The marginal age-compositions were also included, but only for easier viewing of strong cohorts. The CAAL data were not expanded and were binned according to length, age, sex, and year.

The input sample sizes for length and marginal age-composition data were calculated based on (Stewart and Hamel 2014). The input sample size of CAAL data was set at the number of fish at each length by sex and by year. Expanded length frequencies from this survey show intermittent years of small fish; the 2018–2024 period generally suggests most fish are around 40–45cm in length (Figure 11). Strong cohorts are not immediately apparent and it seems that ageing error may result in some variability between years. Conditional age-at-length proportions (Figure 12) show relatively consistent length-at-age with few outliers.

2.1.3 Juvenile Survey

An updated coastwide pre-recruit index of abundance for 2001–2024 for widow rockfish was created using data from three midwater trawl surveys targeting young-of-the-year (YOY) rockfish (Juvenile Survey), provided by Tanya Rogers (SWFSC, pers. comm.). All surveys used identical gear, enabling the construction of a consistent coastwide index spanning from 36°N to the U.S./Canada border since 2004. Sampling in 2020 was limited due to the COVID-19 pandemic and excluded from all models. In 2010 and 2012, coverage was incomplete, so these years excluded from the final model to align with the 2019 assessment. Following the 2015 and 2019 assessments, data from 2001–2003 were also excluded due to limited spatial coverage (36°30' to 38°20' N latitude).

The index was built using a spatial GLM with the sdmTMB package (Anderson et al. 2022), modeling 100-day standardized catch-per-tow as a function of year (fixed effect), Julian date (GAM smoother, $k = 4$), spatial random field, and spatiotemporal random effects. Models with Tweedie, delta-lognormal, and delta-gamma error structures were compared; DHARMA residuals and simulation-based diagnostics indicated the Tweedie model performed best. The index shows a moderately strong increasing trend in juvenile abundance from 2017 to 2023, with a slight decline in 2024 (Figure 41). Recent values remain high relative to the previous decade, and uncertainty estimates support the robustness of this trend.

2.2 Fishery-dependent data

Widow rockfish is primarily caught by bottom trawl and midwater trawl gears (Table 9). This species also commonly bycaught by fishery targeting Pacific hake/whiting (*Merluccius productus*, hereafter “hake”). Minimal amounts have been historically taken by non-trawl gears such as net and hook-and-line, which represented 1.4% and 1.1% of catches, respectively.

The fleet structure in this assessment was streamlined compared to the 2015 benchmark and the 2019 update assessments. The catches in the model are divided among three fishing fleets:

1. A coastwide shore-based bottom trawl fleet (1916–2024). This fleet also includes minimal catches made by net and hook-and-line gears, as well as widow rockfish catches made within historical foreign fishery targeting Pacific Ocean Perch between 1966–1976.
2. A coastwide shore-based midwater trawl fleet that targets rockfish (1979–2024).
3. A mostly-midwater trawl fleet that targets hake and includes an at-sea fleet (1975–2024) and a domestic shore-based fleet (1991–2024).

Catches by fleet and year are provided in Table 9. As in previous assessments, catches from Puget Sound and those from commercial shrimp trawls, commercial pots, and recreational fisheries were excluded (as these are generally minimal).

2.2.1 Landings

Landings from the 1916–2018 period were carried forward into this assessment with two modifications. One was made to the midwater and bottom trawl catches from California. Because PacFIN appears to underestimate midwater trawl catches in California in 1979–1980 when midwater trawl fishery for widow rockfish developed (Edward Dick, SWFSC, pers. comm.) we adjusted midwater and bottom trawl catches from California in these years to reflect the ratio of California midwater to bottom trawl catches in 1981–1982.

The second update was the use of new catch reconstruction for widow rockfish, provided by the Washington Department of Fish and Wildlife (WDFW). New Washington historical landings represent the best scientific information available. The three main sources used in this reconstruction included the US Fish Commission Report (UFSC), Washington Bound Volumes, and Washington Statistical Bulletin. The historical species composition was based on the various historical reports and interviews of fishermen and dockside samplers. The landings between 1981 and 2000 were also provided by WDFW, since WDFW developed and used an improved method for apportioning unidentified rockfish (URCK) category in fish

tickets to the individual species landings. This improved approach relaxed the borrowing rules for missing data used in the WDFW species allocation algorithm that feeds into PacFIN.

Recent catches (2019-onward) were extracted from PacFIN for commercial shorebased data and NORPAC for at-sea hake fishery bycatch, and were otherwise appended onto the 1916-2018 landings and apportioned among fleets using the same criteria as those documented in the 2015 benchmark assessment Hicks and Wetzel (2015).

2.2.2 Fishery catch-per-unit-effort

Three fishery-dependent CPUE indices were included as in the most recent update assessment (Adams et al. 2019). Indices were derived from 1) Oregon bottom trawl (1984-1999), 2) hake at-sea foreign (1977-1988), and 3) hake at-sea domestic fleets (1983-1998). These were not updated; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) and most recent update assessment (Adams et al. 2019) for information.

2.2.3 Fishery length and age data

Biological data from commercial fisheries that caught widow rockfish were extracted from PacFIN on March 25, 2025 and from the NORPAC database on February 3, 2025. Lengths and age samples taken during port sampling in California, Oregon, and Washington were used to generate length and age compositions. The data were classified into bottom trawl, midwater trawl, and hake trawl. For each fleet, the raw observations were expanded to the trip level, to account for differences in samples sizes relative to catch weights among trips (first stage expansion). The expanded length observations were then further expanded to state level, to account for differences in sampling intensity of widow rockfish landings among states combined into a single fleet (second stage expansion).

Table 10 shows the number of trips sampled and Table 11 shows the number of lengths taken for each year and fleet for non-hake fleets from the three states. Table 12 shows these numbers for the shoreside and at-sea hake fisheries. Expanded length compositions for bottom trawl, midwater trawl and hake fleets are shown in Figure 13 to Figure 15. Table 13 and Table 14 show the number of trips and number ages used in the. Age compositions for the three fleets are shown in Figure 16 and Figure 18. Occasional cohorts appear to move through the population, indicating that widow rockfish population dynamics may be characterized by episodic recruitment events.

2.2.4 Discards

Data on discards of widow rockfish are available from three different sources. Historical sources included Pikitch et al. (1988) and Enhanced Data Collection Project (EDCP, Sampson and Lee (2002)). The West Coast Groundfish Observer Program (WCGOP) provided information on recent discard of widow rockfish between 2002–2023. Since 2011, under trawl rationalization, 100% observer coverage is required for the limited entry trawl sectors, which resulted in a large increase in data and ability to determine discard behavior.

For this assessment, we estimated bottom trawl discards and midwater trawl discards externally, using historical sources as well as WCGOP data, and added calculated discard amounts to landings of the corresponding fleets by year. The methods used to reconstruct discard amounts in different time periods are detailed below.

1. For the years 1982–1989, landings in the bottom trawl and midwater trawl fleets were scaled using fleet-specific discard rates estimated from the Pikitch study.
2. For the years 1990–2001, landings in the bottom trawl and midwater trawl fleets were scaled using fleet-specific discard rates estimated from the EDCP data.
3. For the years 2002–2023, GEMM estimates of fleet-specific discards amounts were added to the landed catch (bottom trawl + other, and midwater trawl).
4. The 2024 values were set to the average from the GEMM estimates for the period 2021–2023.

The treatment of the discard changed from the 2015 benchmark and 2019 update assessments. Then, discards were estimated with retention functions within the model, which was standard practice in 2015, following a long period of large discards due to trip limits. However, discard rates have been low for widow rockfish for more than 10 years so the benefit of using retention functions to extrapolate beyond the range of observed data is reduced and the biological data from the discards are no longer as likely to be informative about dynamics of the total population. As a part of model bridging, we illustrated that treating discard as catch (rather than estimating them in the model) did not have much impact on model results. Also, this approach avoids hard-to-diagnose issues that come from estimating retention curves (especially with limited amounts of data). For some species, the discard data may provide important information about recent recruitment because the smaller fish are being discarded. However, for this stock, removing the discard amounts and length-compositions from the likelihood had little impact on recruitment estimates.

Widow rockfish bycatch within the hake fishery represents total catch, with discards already accounted for.

2.2.5 Biological data

The approach to the estimation of all biological parameters was the same as in the 2019 update assessment, Adams et al. (2019).

2.2.5.1 Weight-length relationship

Weight-at-length data were updated for this assessment. Following the approach used in the 2015 benchmark, data used to estimate the length-weight relationship for widow rockfish were gathered from commercial catch sampling schemes (PACFIN, ASHOP) and fishery-independent surveys (Triennial Survey and WCGBTS).

The following relationships between weight and length for females and males were used in the current assessment:

$$\text{Females:} \quad \text{weight} = 1.5885 \times 10^{-5} \cdot \text{length}^{2.9873}$$

$$\text{Males:} \quad \text{weight} = 1.4507 \times 10^{-5} \cdot \text{length}^{3.0123}$$

where weight is measured in kilograms and length in cm. These parameters were used in the assessment as fixed.

2.2.5.2 Maturity schedule

Maturity parameters in this update assessment were carried over from 2015 benchmark and 2019 update assessments; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) and most recent update assessment (Adams et al. 2019) for information.

2.2.5.3 Fecundity

The model uses widow rockfish specific fecundity-at-length relationships from the meta-analysis from Dick et al. (2017) instead of assuming spawning output being proportional to body weight as in 2015 and 2019 models. Accounting for size-dependent fecundity in rockfish stocks using the meta-analysis from Dick et al. (2017) has become standard practice in recent years and reflects the best available science. The fecundity parameters for widow rockfish are reported at <https://github.com/EJDick-NOAA/Rockfish-Fecundity> as follows:

$a = 1.10961\text{E-}08$, $b = 4.545$

The value for a above produces spawning output values in billions of eggs from SS3, given length in cm.

2.2.5.4 Natural Mortality

In this assessment, natural mortality (M) is estimated for females and males, while using the Hamel and Cope (2022) prior. The prior on M has been updated to reflect the most recent guidance from Hamel and Cope (2022); the log-mean therefore remains unchanged while the log-SD has been set to 0.31. Using a maximum age of 54 the point estimate and median of the prior on M is 0.10.

2.2.5.5 Length-at-age

Growth parameters were fully estimated within the assessment model, for females and males separately following the same formulation in 2015 and 2019 assessment models.

2.2.5.6 Ageing bias and imprecision

Ageing error matrices were unchanged from the previous assessment; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) and most recent update assessment (Adams et al. 2019) for information.

2.3 Environmental and Ecosystem Data

This assessment did not use any environmental or ecosystem data related to the stock.

3 Assessment model

3.1 History of modeling approaches

This section is not required for an update assessment; please refer to the most recent benchmark assessment (Hicks and Wetzel 2015) and most recent update assessment (Adams et al. 2019) for information.

3.2 Responses to most recent past STAR Panel recommendations

There are no recommendations from the most recent STAR Panel to address.

3.3 Responses to SSC Groundfish Subcommittee requests

Responses to the most recent requests from the Science and Statistical Committee (SSC) and Pacific Fishery Management Council (PFMC) are provided in a separate document.

3.4 Model Structure and Assumptions

3.4.1 Changes from the Last Assessment and Model Bridging

The last benchmark stock assessment for widow rockfish was conducted in 2015 (Hicks and Wetzel 2015), and it was updated in 2019 (Adams et al. 2019).

In this assessment, the changes made to the 2019 model (Adams et al. 2019) occurred in two stages. The first round of changes was reviewed in August 2025. The model reviewed at the August 2025 Pacific Fishery Management Council (PFMC) Science and Statistical Committee (SSC) Groundfish Subcommittee (GFSC) meeting followed the 2015 benchmark assessment in its structure and assumptions, as required by the Terms of Reference for update assessments (with the exception of removing discard length composition data for the hook-and-line fleet, as discussed and approved at the review meeting).

The second round of changes followed after the August 2025 review meeting, when we explored a larger range of options which go beyond update assessment Terms of Reference and no longer match the 2015 benchmark. The second round of changes resulted in more

parsimonious model that was raised to the current best practices in West Coast groundfish stock assessments.

The changes made to the model are listed in two separate groups, according to stages described above.

Changes made from the 2019 update assessment for August 2025 update:

The exploration of models began by bridging from the 2019 update assessment to current SS3 version, which produced no discernible difference.

We then focused on data bridging (Figure 25).

- **Adding most recent data from commercial fleets** (as described in “Data” section of this report). Updating the catch series did not have a substantial effect on the historical biomass; the stock biomass increasing from 2000–2020 before decreasing up to the current period.
- **Updating WCG BTS index using current methodology** (as described in “Data” section of this report).
- **Updating the log-SD of the prior for natural mortality** based on Hamel and Cope (2022). The log-mean remained unchanged while the log-SD has been decreased from 0.44 to 0.31.
- **Updating length-weight parameters estimated** by including most recent data.
- **Extending the main period for estimating recruitment deviations and updating recruitment bias adjustment parameters** based on Methot and Taylor (2011).
- **Adding a block to hake fleet selectivity to account for recent changes in fleet behavior.** Adding a block to hake fleet selectivity accounted for a change in fish mean length (2020–2024) due to shifts in the spatial distribution of the hake fleet and improved the fit to the length compositions (Holland and Martin 2019). This change had a discernible influence on the estimated spawning biomass.
- **Adding hook-and-line discards to landings in the hook-and-line fleet.** WCGOP corrected discard lengths for the hook-and-line fleet to omit nearshore fixed gear samples, which were erroneously included with the hook-and-line fleet in the previous benchmark (Hicks and Wetzel 2015) and update assessments (Adams et al. 2019). This resulted in changes to the discard length distribution and years for which data was available (Figure 19). Because available hook-and-line discard length samples were too scarce, this assessment model was unable to reliably estimate retention parameters. Therefore, in this assessment, we added the hook-and-line discard to hook-and-line landings.

Updates to discards, particularly the removal of hook-and-line length composition data and adding hook-and-line discards to landings, had the most significant impact on the absolute stock biomass: absolute spawning biomass before 1980 was slightly lower than the 2019

update assessment estimates, and noticeably lower from the mid-2010s through the current year.

In the 2015 and 2019 assessments, the hook-and-line discard length data consisted primarily of small fish from nearshore fixed gear Figure 19, and the model interpreted the presence of those small fish as a recruitment signal, with the 2019 update assessment estimated the 2013 recruitment as the strongest year class over the duration of the fishery. Those discard lengths interacted in the model with discard amounts so that the model exhibited substantial sensitivity to even slight changes in discard data within the hook-and-line fleet. Figure 22, Figure 23 and Figure 24 show intermediate bridging steps in updating hook-and-line discard data. The slight decrease in 2011-2017 discard amounts (compared to those used in the 2019 assessment) caused decrease in estimated recruitment (Figure 24) leading to lower spawning biomass (Figure 22). Adding discards for years 2018-2023 caused recruitment to be estimated unreasonably high bringing spawning biomass well above unfished level Figure 23. Such a disproportionate effect of a small amount of data from the fleet that contributes only 1.1% of catch is most likely related to how discard was modeled in the 2015 assessment. Recent assessments' exploration revealed that there are some hard-to-diagnose issues that come from estimating retention parameters (especially with limited amounts of data). With the discard length data as well as retention parameters removed, and the hook-and-line discards added to landings, the model no longer estimated extremely high recruitment in the 2010s, which resulted in lower estimates of spawning biomass and stock productivity in recent years.

Updates to survey indices and composition data, following updates to discards, lead to minor changes in spawning biomass estimates (Figure 25). The rest of the changes also had only minor impact on recent relative spawning biomass estimates (Figure 25) and collectively resulted in a very small effect in relative spawning biomass post-1990 (Figure 26).

Changes made from August 2025 model for the January 2026 model:

After August 2025 review, we explored changes to the model that were outside the Terms of Reference for an update assessment. Those changes included:

- **Streamlining the fleet structure** (as described in "Data" section of this report). The landings from fleets with minimal and mostly historical catches (hook-and-line and net) were added to the bottom trawl landings with hook-and-line and net composition data removed from the model.
- **Use external estimates of discards rather than estimate within the model.** Methods, used to reconstruct discards in the bottom trawl and midwater trawl fleets, based on the Pikitch, EDCP and WCGOP data are described in "Data" section of this report. All retention parameters were removed from the model.
- **Updating the data weighting approach.** The data weighting approach was changed to match recent practices, including the use of the Francis rather than McAllister-Ianelli approach, and no longer including a 50% additional reduction in weights. The

Francis method performed better in a simulation analysis (Punt 2017) and has been the preferred approach in most recent stock assessments. The 50% multiplier on the data weights used in the previous assessment was an approach more common in 2015 to account for the potential double use of data since length and age are observed from the same fish. However, the sum of the adjusted input sample sizes for length and age data combined are already far lower than the number of sampled fish because they are primarily based on trips or hauls and then further reduced by the data-weighting algorithm. This 50% adjustment is no longer common practice and has not been used any other recent assessments.

- **Use a fecundity relationship rather than assume the number of eggs is proportional to body weight.** Use fecundity-at-length based on the meta-analysis from Dick et al. (2017) has become standard practice in recent years and reflects the best available science.
- **Various additional refinements were made.** These refinements had a minimal impact on the model.
 - The landings from Washington were updated to use the new Washington catch reconstruction.
 - The historical catch within foreign fishery that targeted Pacific Ocean Perch during 1966–1976 period was moved from the hake fleet to the bottom trawl fleet.
 - The bias adjustment ramp was updated after the retuning and also to modify the estimated values to have full bias adjustment starting in 1970.
 - The start of the final block on midwater trawl selectivity was moved from 2011 to 2017.

Comparisons of summary biomass, fraction unfished and estimated recruitment for major bridging steps within the second round of changes are shown in Figure 27, Figure 28 and Figure 29. None of the updates resulted in large changes in estimated population trajectories and model outputs, but several important quantities were sensitive to a few of the changes.

Combining discards with landings in bottom trawl and midwater trawl fleets did not reveal unexpected or unreasonable patterns with a disproportionate effect of changes on model results (as it was in case of hook-and-line discard data). Simplifying the fleet structure did not bring about large changes in model results either, but interacted with the Francis weighting approach. Extending the block on midwater trawl selectivity did not cause any noticeable change.

The use of the Francis weighting approach (instead of McAllister-Ianelli) and using $\lambda = 1$ (and not further down-weighting fishery composition data) resulted in the biggest change from the August 2025 base model, causing the estimates of natural mortality to go up, which translated into increases in the estimated productivity of the stock, including equilibrium catch and estimated overfishing limits (OFLs). Relative to the August 2025 model, the updated data weighting method upweighted length data from the bottom trawl fleet and

age data from all three commercial fleets (bottom trawl, midwater trawl, hake); all other composition data sources were downweighted (Table 17). For bottom trawl, in particular, this upweighting is likely in part due to the removal of discard composition data, as Stock Synthesis does not apply separate data weights to discarded and retained composition data from the same fleet. The discard compositions are very noisy and difficult to fit, so their removal allowed for upweighting of the remaining data. The decrease in the weight of WCG BTS conditional age-at-length data by approximately one-third, paired with upweighting of other data, is likely the major driver of the increased estimate of natural mortality.

With the change in fecundity relationship spawning output is now expressed in billions of eggs and no longer directly comparable with spawning biomass expressed in metric tons in previous models. This change did not have much impact on age 4+ biomass (Figure 27), but the spawning output time series was impacted by the changes in relative contributions of larger/older fish which are disproportionately removed from the population, illustrated in Figure 30, which leads to lower estimates of fraction of unfished and estimated OFLs. In particular, the stock status (fraction of unfished spawning output) reached a low point of 0.232 in 1998 (below the 0.25 overfished threshold) as compared to a minimum of 0.326 in 2001 in the August 2025 base model. The addition of the fecundity relationship also resulted in slightly lower estimates of OFLs and fraction unfished at the end of the time series compared to the previous step in this bridging, though still higher than the August 2025 estimates. Note that the model reviewed in October 2025 included a fecundity relationship associated with “unobserved rockfish” rather than the parameters specific to widow rockfish. This is the only difference in model configuration between the October 2025 and current base models, although the difference between the two fecundity relationships is smaller than the difference between either one and the weight-length relationship used to estimate spawning output in previous assessments.

Comparison of the time series estimates from the 2019 update assessment model and this model are shown in Figure 31 to Figure 34.

3.4.2 Modeling Platform and Structure

Stock Synthesis statistical catch-at-age modelling framework, version 3.30.24.1, was used for this assessment; and r4ss version 1.52.1 and R version 4.5.0 to investigate and plot the model fits.

3.4.3 Model Overview

The model is a two-sex, age-structured model starting in 1916 with an accumulated age group at 40 years. Sex-specific growth and natural mortality were estimated. The length

frequency distributions range between 8 and 56 cm, binned into 2 cm intervals. Population 2 cm length bins are defined at a wider range, between 6 and 60 cm. Ageing error was retained from the 2015 benchmark assessment (Hicks and Wetzel 2015). The model uses widow rockfish fecundity-at-length relationships instead of assuming spawning output being proportional to body weight as in the 2015 and 2019 models. With the change in fecundity relationship, spawning output is now expressed in billions of eggs and not in metric tons, as in 2015 and 2019 models.

3.4.3.1 Model Fleets and Areas

The assessment uses a single-area model, consistent with the previous benchmark and update assessments. The fleet structure in this assessment was streamlined compared to the 2015 benchmark and the 2019 update assessments. In this assessment, the catches are divided among three fishing fleets:

1. A coastwide shore-based bottom trawl fleet (1916–2024). This fleet also includes minimal catches made by net and hook-and-line gears, which represented 1.4% and 1.1% of the input catches, respectively. Removals from net and hook-and-line fisheries were previously treated as separate fleets. The bottom trawl fleet also includes widow rockfish catches made within historical foreign fishery targeting Pacific Ocean Perch between 1966–1976.
2. A coastwide shore-based midwater trawl fleet that targets rockfish (1979–2024).
3. A mostly-midwater trawl fleet that targets hake and includes an at-sea fleet (1975–2024) and a domestic shore-based fleet (1991–2024).

The following fishery-independent surveys are also included in the model:

1. Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast Triennial Shelf Survey (Triennial Survey).
2. Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey (WCGBTS) (2003–2024).
3. recruitment index from the National Marine Fisheries Service (NMFS) SWFSC and NWFSC/PWCC Midwater Trawl Survey (Juvenile Survey) (2004–2024).

3.4.4 Model Parameters

3.4.4.1 Estimated and Fixed Parameters

There are 194 estimated parameters in the base model (relative to 214 in the August 2025 base model which included additional parameters for selectivity and retention). These included one parameter for recruitment (R_0), 10 sex-specific parameters for sex-specific growth,

two sex-specific natural mortality parameters, four parameters for the catchability (q) of the hake and the Triennial Survey of abundance (catchability for other indices were calculated analytically using the “float” option in SS3), four parameters each for extra variability for the hake, bottom trawl, juvenile and foreign-at-sea indices of abundance, 36 parameters for selectivity, 125 recruitment deviations (16 of which set up the initial age structure), and 12 forecast recruitment deviations.

Fixed parameters in the model included steepness, standard deviation of recruitment deviates, maturity at age, fecundity-at-length, and length-weight parameters. Consistent with the previous assessment (Adams et al. 2019), steepness was fixed at 0.72, matching the mean of the current west coast rockfish steepness prior (Thorson et al. 2019), described below in the “Priors” section. A likelihood profile were done for steepness. The standard deviation of recruitment deviates was fixed at 0.60. Maturity at age was fixed as described in the previous update assessment, Adams et al. (2019). Fecundity-at-length parameters were fixed at values from Dick et al. (2017). Length-weight parameters were fixed at estimates using length-weight observations from the WCGBTS (Figure 20 and Table 18).

All selectivity curves in the base model were length based and had the same shape as the 2015 benchmark and the 2019 update assessments. The base model assumed asymptotic selectivity (using the double-normal formulation in SS3) for each fishery, except for the midwater trawl fishery (Hicks and Wetzel 2015). The WCGBTS and Triennial Survey both used spline curves. The WCGBTS survey fleet selectivity was estimated to be slightly lower at lengths greater than 45 cm, compared to the selectivity estimated in 2019.

Time blocks were used for selectivity parameters in the bottom trawl, midwater trawl, and hake fisheries as indicated in Figure 37.

3.4.4.2 Priors

The prior on natural mortality (M) in the previous assessment was defined as a lognormal with mean on the log-scale of $\ln(5.4/A_{\max})$ and $SD(\ln(M)) = 0.44$ following Hamel (2015) and Then et al. (2015). In the current assessment, the prior on M has been updated to reflect guidance from Hamel and Cope (2022). With this update, the log-mean remains unchanged while the log-SD has been set to 0.31. Using a maximum age of 54 the point estimate and median of the prior on M is 0.10.

The prior for steepness (h) has not changed since the 2019 update assessment (Adams et al. 2019). It assumes a beta distribution with parameters based on Thorson et al. (2019), with mean = 0.72 and SD = 0.16 ($\alpha = 4.95$, $\beta = 1.93$) which was approved for use in all rockfish stock assessments.

3.4.4.3 Recruitment deviations

Recruitment deviations from 1900–2024 were estimated to appropriately quantify uncertainty; inappropriate specifications can have a large effect on model uncertainty. The standard deviation of recruitment variability (σ_R) was assumed to be 0.6 in the 2015 assessment, based on iteratively tuning to a value slightly less than the observed variability of recruitment deviations in the period 1975–2010 in 2015 (Figure 40). The earliest length-composition data occur in 1976 and the earliest age data were in 1978. The most informed years for estimating recruitment deviations based on available composition data were from about the mid-1970s to about 2019. The period from 1900–1970 was fit using an early series with little or no bias adjustment, the main period of recruitment deviates was 1971–2020 using a bias adjustment ramp, and 2021 onward was fit using forecast recruitment deviates with little bias adjustment. Methot and Taylor (2011) summarize the reasoning behind varying levels of bias adjustment based on the information available to estimate the deviates. The bias adjustment settings were tuned to the estimates produced by `r4ss::SS_fitbiasramp()` with the exception that the first year with full bias adjustment was set to 1970 (instead of the estimate of 1975, which occurs after two large, reasonably well-informed recruitments in 1970 and 1971).

3.4.4.4 Sample weights

The base model uses the Francis (2011) weighting method. This is a departure from the August 2025 base model, which followed the most recent benchmark assessment (Hicks and Wetzel 2015) in using the McAllister and Ianelli (1997) method. The data weighting approach was changed to match recent practices, including the use of the Francis rather than McAllister-Ianelli approach. The Francis method performed better in a simulation analysis (Punt 2017) and has been the preferred approach in most recent stock assessments.

An additional change from the previous assessments was the removal of a 50% additional reduction in weights (via the lambda factors) for all fleets with both marginal length and marginal age compositions, which down-weighted the likelihood contribution by 50% to account for the potential double use of data since length and age are observed from the same fish. This 50% adjustment is no longer common practice and has not been used any other recent assessments. The adjusted input sample sizes are based on trips or hauls rather than the number of sampled fish, and the sum of the adjusted input sample sizes for length and age data combined are already far lower than the number of sampled fish, so no further reduction is necessary to avoid double use.

3.4.5 Key Assumptions and Structural Choices

Following recommendation of Pacific Fishery Management Council (PFMC) Science and Statistical Committee (SSC) Groundfish Subcommittee (GFSC) meeting in August 2025,

for this update assessment we explored a larger range of options which go beyond update assessment Terms of Reference, which resulted in more parsimonious model that was raised to the current best practices in West Coast groundfish stock assessments.

The major structural choice was related to fleet structure, which was simplified to focus on major fisheries that remove more than 97% of catch. Those included bottom trawl, midwater trawl and hake fleets. The fisheries with minimal and mostly historical catches, such as hook-and line and net, were added to the bottom trawl fleet. In the 2015 benchmark and 2019 update assessments catches were divided among five fleets (bottom trawl, midwater trawl, catches within fleet targeting hake, net and hook-and line).

Also, the 2015 benchmark and 2019 update assessments estimated discards using retention curves for the bottom trawl, midwater trawl, and hook-and-line fleets based on discard biomass and length composition data from the WCGOP. Changes to the underlying discard length composition data for the hook-and-line fleet from WCGOP illustrated that very large recruitment events previously estimated for 2013 and 2014 were mainly driven by very small amounts of composition data for a minor fleet responsible for less than one-tenth of a percent of the catch in the last decade. This led to implausible population dynamics, so the decision was made to consolidate fleets and add minimal (and primarily historical) catches from net and hook-and-line fisheries to the shore-based bottom trawl fleet. In this assessment, the discards in bottom trawl and midwater trawl fleets were estimated externally, based on historical and the WCGOP data, and added to landings within the corresponding fleets. With that, all retention parameters became obsolete and thus were removed from the model. Discard rates have been low for widow rockfish for more than 10 years so the benefit of using retention functions to extrapolate beyond the range of observed data is reduced and the biological data from the discards are no longer as likely to be informative about dynamics of the total population. This change also helps avoid hard-to-diagnose issues that come from estimating retention curves (especially with limited amounts of data), as we encountered with the hook-and-line fleet, earlier in the assessment process. Removing the discard amounts and length-compositions from the likelihood had little impact on recruitment estimates.

Other structural choices were related to updating model to the current assessment practices, and included using up-to-date weighting approach and treatment of fecundity. Details on that are provided “Changes from the Last Assessment and Model Bridging” section of this report.

3.5 Base Model Results

As a supplement to the model results figures included in this report and described below, a full set of [diagnostic plots](#) created by the `r4ss` package (Taylor et al. 2021) is available on the assessment GitHub repository (github.com/mcgoodman/widow_rockfish_2025) along with the Stock Synthesis [input files](#).

3.5.1 Parameter Estimates

The base model parameter estimates along with asymptotic standard errors are shown in Table 18. Estimates of key derived parameters and asymptotic 95% confidence intervals are shown in Table 21.

The estimates of natural mortality (0.135 yr^{-1} and 0.147 yr^{-1} for females and males, respectively). These estimates continue gradual decrease in M from those of the 2019 update assessment (Female 0.1444, Male: 0.1549; Adams et al. (2019), Figure 35) and the 2015 benchmark assessment (Female: 0.1572, Male: 0.1705; Hicks and Wetzel (2015)). However, they were higher than suggested by the medians of the prior distributions used in this assessment and the previous assessments. Fixing M at lower values than those estimates resulted in a pattern of below average recruitment in the early part of the time series which is not well informed by age or length composition data (Figure 64). This suggests that the model is attempting to reduce the number of observations of older fish in the data. The estimates of M fall within the 95% quantile range of the prior distribution (0.055–0.184), and are shown in Figure 35.

Updating age composition data resulted in the largest decrease in M among the various intermediate models bridging from the 2019 assessment. This is broadly consistent with observations of older individuals in updated age composition data. For instance, the maximum age for widow rockfish in the WCGBTS data prior to 2019 was 37 yrs with a 99th percentile of 29, while data collected since include individuals up to 51 with a 99th percentile of 34. Updating that natural mortality prior resulted in a comparatively small decrease in M (Figure 35).

Estimating M is difficult in stock assessments, and the estimated values may represent model misspecification instead of the actual life-history trait. However, in alternative models to the base, the estimates of M were not higher than 0.13 yr^{-1} (Table 20), except when forcing asymptotic selectivity on the midwater trawl fleet (Table 20). Uncertainty in the estimated M was also much less than the range of the prior (Figure 35).

Selectivity curves were estimated for commercial and survey fleets and parameter estimates are provided in Table 18. The estimated selectivity are shown in Figure 36. The selectivity curves showed a strong dome-shaped selectivity for the midwater fleet, while the selectivity for the hake fleet did not support dome-shaped selectivity (Figure 36). The estimated selectivity curves for the Triennial and WCGBTS were similar to each other except that the triennial survey selected larger fish (Figure 37). The WCGBTS exhibited a more pronounced dome-shaped selectivity compared to the 2019 assessment.

In the 2015 benchmark assessment, additional survey variability (observation error added directly to each year's input variability) for the Triennial Survey and WCGBTS was not estimated in the model because initial runs estimated the it at zero (Hicks and Wetzel

2015). To avoid bound issues in estimation of the Hessian, their values were fixed at zero. We retained the same modelling approach for the update assessment. The additional standard deviation added to the fishery-dependent indices was quite large: 0.2 for the bottom trawl index and 0.6 for the foreign at-sea hake fleet. The additional variability on the juvenile survey was the highest, at 1.27, giving the index very little weight in the model.

The estimates of maximum size for both females and males (Table 18) did not differ substantially from Adams et al. (2019). Estimates of k were slightly different in the model, but that is expected when accounting for selectivity. Estimated growth curves are shown in (Figure 38).

Estimates of recruitment suggest that the widow rockfish population is characterized by variable recruitment with occasional strong recruitment events and periods of low recruitment (Figure 39, Table 21). There is little information regarding recruitment prior to 1965.

3.5.2 Fits to the Data

Fits to data are provided for survey abundance indices, length composition data for the fisheries and surveys, marginal age compositions for the fisheries, and conditional age-at-length observations for the WCG BTS.

Survey indices and total discards were fitted assuming a lognormal likelihood. The six indices of abundance (three survey series, and three fishery indices of abundance) are shown in Figure 41. The Triennial Survey treatment was consistent with Adams et al. (2019). Extra standard error was estimated for all of the series except for the two survey series (Table 18). None of the series showed patterns in residuals, and with the large amount of error, none of the series showed serious lack of fit. The recent WCG BTS showed a general increase from 2003 - 2015 followed by a general decrease from 2016 - 2024, which was also estimated in the base model (Figure 41). The model did not fit the indices variability well, particularly years with relatively low (2015, 2021, 2022) or high (2016) estimates of abundance, as might be expected for a long-lived rockfish species.

Annual fits to the length-composition data are depicted using Pearson residuals-at-length for all fleets in Figures 42. More detailed plots of fitted and observed proportions at length are shown in Appendix A. The combined WCG BTS length frequencies across years displayed a weakly bimodal distribution with a valley around 37 cm, which the model approximated well (Figure 43). Fits to the Triennial Survey length frequencies (which have not been updated in this assessment) show underestimation of 35-45 cm males and overestimation of similar length females, as in Hicks and Wetzel (2015).

Age data were fitted as marginal age compositions for the fishing fleets and as conditional age-at-length for the WCG BTS, which were expanded by tow and then by strata, following the

2015 benchmark assessment (Hicks and Wetzel (2015)). Pearson residuals for the commercial fleets are shown in Figures 42. Aggregating across years shows that the fit to age comps was good (Figure 45). The observed and expected age-at-length are shown in Figure 46 for the WCGBTS observations. Plots with the residuals for individual observations showed reasonably good fits to the conditional age-at-length data from the WCGBTS (Figure 47).

3.5.3 Population Trajectory

The predicted spawning output (billions of eggs) is given in Table 19 and plotted in Figure 48. The predicted spawning output from the base model generally showed a slight decline over the time series until 1966 when the foreign fleet began. A short, but sharp decline occurred, followed by a steep increase due to strong recruitment. The spawning output declined rapidly with the developing domestic midwater fishery in the late 1970s and early 1980s. The stock continued to decline until 2000 when a combination of strong recruitment and low catches resulted in a quick increase from 2000 - 2018. The stock declined rapidly from 2019 until the end of the time series, due to a combination of increased catches and low recruitment.

The 2025 spawning output relative to unfished spawning output is above the target of 40% of unfished spawning output (50.1%), with a low of 22% in 1998 (Figure 50). Approximate confidence intervals based on the asymptotic variance estimates show that the uncertainty in the estimated spawning output is high, especially in the early years. It should be noted that while the stock is currently above the 40% target, the target does fall within the approximate 95% confidence interval (Figure 50).

Recruitment deviations (Figure 39 and discussed in Section 3.3.1) provide a more realistic portrayal of uncertainty. Estimates of recruitment appear to be episodic and characterized by periods of low recruitment, with several very large, but uncertain, estimates. The 2019 update assessment estimated the 2013 recruitment as the strongest year class over the duration of the fishery, however the current assessment does not support this. It may be worthwhile to investigate the periods of strong and weak year classes further to see if it is an artifact of the data, a consistent autocorrelation, or a result of the environment. The input bias adjustment ramp matched the estimated ramp (Figure 40).

The stock-recruit curve resulting from a fixed value of steepness is shown in Figure 40 with estimated annual recruitment also shown. The stock is predicted to have never fallen to low enough levels that the steepness is obvious. However, the lowest levels of predicted spawning output showed some of the smallest recruitment events and very few above average recruitment events. Steepness was not estimated in this model, but sensitivities to alternative values of steepness were explored via likelihood profile analysis and discussed below.

3.6 Model Diagnostics

3.6.1 Convergence

A number of tests were performed to verify convergence of the model. Following conventional AD Model Builder methods (Fournier et al. 2012), we checked that the Hessian matrix for the base model was positive-definite. There were no difficulties in inverting the Hessian to obtain asymptotic variance estimates. The maximum component of the final gradient is very low.

To confirm that the reported estimates were from the global best fit, we evaluated the model's ability to recover similar likelihood estimates when initialized from dispersed starting points (jitter option in SS3). Starting parameters were jittered using a setting of 0.1 for 100 iterations. This perturbs the initial values used for minimization with the intention of causing the search to traverse a broader region of the likelihood surface. The majority (57 out of 100) returned to the same objective function value as the proposed model. The remaining runs exhibited worse fit than the proposed model. The spread of this search indicates that the jitter was sufficient to search a large portion of the likelihood surface, and that the model is in a global optimum.

We also completed a set of standard model diagnostics that included retrospective analysis and likelihood profiles described below.

3.6.2 Parameter Uncertainty

Parameter estimates are shown in Table 18 and Table 19 along with approximate asymptotic standard errors. The only parameters with an absolute value of correlation greater than 0.95 were the female and male natural mortality parameters, which is expected. Estimates of key derived quantities are given in Table 21 along with approximate 95% asymptotic confidence intervals. There is a reasonable amount of uncertainty in the estimates of biomass. The confidence interval of the 2025 estimate of the fraction of unfished spawning output is 30.35%–69.8% and mostly above the management target of 40% of the unfished spawning output.

3.6.3 Sensitivity Analyses

Sensitivity analysis was performed to determine the model behavior under different assumptions than those of the base case model. Broad range of sensitivity runs were conducted at multiple stages of the assessment. Here, we present only several of them:

1. Fixed natural mortality at 0.1 for both sexes (the median of the prior used in the assessment).
2. Fixed natural mortality at 0.144 yr^{-1} for females and 0.155 yr^{-1} for males (2019 assessment estimated values).
3. Fixed natural mortality at 0.157 yr^{-1} for females and 0.17 yr^{-1} for males (2015 assessment estimated values).
4. Estimating natural mortality using the prior based on the maximum age of 50 years old.
5. Estimating natural mortality using the prior based on the maximum age of 45 years old.
6. Using double-normal selectivity curve for WCGBTS, fixing it asymptotic.
7. Using double-normal selectivity curve for WCGBTS, allowed it to be dome-shaped.
8. Adding small amount of recreational catches as reported by WCGOP.

Differences in likelihood and estimates of key parameters and derived quantities are shown in Table 20. Predicted trajectories of spawning output, fraction unfished, recruitments and recruitment deviations comparisons of model estimates for 2025 are shown in Figure 54, Figure 55, Figure 56, and Figure 57.

Fixing M at values lower than the base case estimate resulted in decreases in estimated spawning output in recent years. Fixing M at 0.1, the median of a prior informed by maximum age of 54 years, resulted in largest decrease in spawning output and associated management quantities. The unfished spawning output in sensitivity runs with M fixed at the estimates from the 2019 and 2015 assessments is very similar among these runs. However, the lower the natural mortality is, the lower the final assessment year spawning output is, since the less productive stock is impacted more by fishery removals. The models with higher natural mortality need to scale up estimated recruitments, in order to provide for a sufficient amount of fish to sustain the catches over time, while having the same recruitment deviations.

The model results were not sensitive to the alternative values of the median of the M prior, and estimated M values were similar in all runs. This is an indication that the data provide a stronger signal about M than the prior.

The last three runs (changing the selectivity pattern for WCGBTS and adding recreational catches) did not have a noticeable impact.

3.6.4 Retrospective Analysis

A 10-year retrospective analysis (Figure 58) was conducted by running the model removing the most recent 1 to 10 years of data (i.e., “Data -1 Years” corresponds to data through 2023 instead of 2024). Population trends from all retrospective runs were very close, and

no concerning patterns were observed in the retrospective analysis. There is a general trend of increasing scale and fraction unfished as years are removed. This result is expected because the fraction of older fish in the WCGBTS increased substantially in recent years, and sequentially peeling off individual years of WCGBTS age data leads to increasing estimates of natural mortality and unfished recruitment. Additionally, as years are pulled away the estimate of the size of the 2016 year class decreases towards average and the sizes of year classes since 2016 increase towards average. This is also expected as information about those recent year classes is removed.

Also, estimated relative spawning output from the base model was compared with previous assessments Figure 59. The current assessment follows a similar trend to previous assessments over the overlapping years, and estimated spawning output falls within the mid-range of previous assessments.

3.6.5 Likelihood Profiles and key parameters

Likelihood profiles were conducted for unfished virgin recruitment (R_0), steepness (h) and sex-specific natural mortality (M) values simultaneously. These likelihood profiles were conducted by fixing the parameter at specific values and removing the prior on the parameter being profiled. Without the original prior distribution the MLE estimates from the base case will likely be different than the MLE in the likelihood profile, but this displays what information the data have.

For profiles of natural mortality, the negative log-likelihood was minimized at a value of 0.147 for males, and a value of 0.135 for females. Profiles for natural mortality for each sex are illustrated in (Figure 62, Figure 63). Profiles over female natural mortality indicate that the recruitment deviation prior and age data most strongly influence the estimate of natural mortality, with survey indices also being somewhat influential. Length data provides very little information. Profiles over male natural mortality behaved similarly. For steepness, the negative log-likelihood was minimized at a steepness of 0.77, however the 95% confidence interval extends over the entire range of possible steepness values. Profiles for steepness are illustrated in (Figure 61). For R_0 , the negative log-likelihood was minimized at a value of 10.6, which supports the base model estimates of R_0 at 10.608. Profiles for R_0 are illustrated in Figure 60.

4 Management

4.1 Reference Points

Reference points were calculated using the estimated selectivity parameters and average catch distribution among fleets in the most recent five years of the model (2019-2024). A list of estimates of the current state of the population, as well as reference points based on 1) a target unfished spawning output of 40%, 2) a SPR of 0.5, and 3) the model estimate of maximum sustainable yield (MSY), are all listed in Table 21. The SPR is expected lifetime reproductive output under a given fishing intensity as a proportion of the unfished expected lifetime reproductive output.

The estimated trajectory in spawning output from the base model generally showed a slight decline until the late 1970s, followed by steep increase above unfished equilibrium biomass and reaching a peak in 1979. This was followed by a steep decrease up to the mid-1980s, and then a more gradual decrease through 2000 (Figure 50). Between 2001 and 2016, the spawning output increased continuously due to small catches and several years of high recruitment (though with lower than average recruitment in other recent years). The spawning output relative to unfished equilibrium spawning output climbed above the target of 40% of unfished spawning output in the early 2000s. It is estimated to still be above the target, though the lower bound of the 95% confidence interval lies between the target and minimum stock size threshold (Figure 50). The fishing intensity (relative 1-SPR) exceeded the current estimates of the harvest rate limit ($SPR_{50\%}$) throughout the 1980s and early 1990s, and has again since 2018 (Figure 52). Exploitation rates on widow rockfish between 2001 and 2016 were predicted to be far below target levels. In recent years, the stock has experienced exploitation rates that have been above the target level while the biomass level has remained above the target level (Figure 53).

While the estimate of population scale has remained relatively steady across the 2015, 2019, and 2025 assessments, natural mortality (M) have declined from 2015 (female M of 0.157 yr^{-1}) to 2019 (0.144 yr^{-1}) to 2025 (0.135 yr^{-1}), reflecting the increase of the number of older fish in the population throughout rebuilding years. Thus, although the population scale is estimated consistently, additional years of data (in particular, age composition data) following the re-opening of the fishery have led the model to estimate that the population is less productive, which results in lower yields at all three reference points (proxy based on spawning output, proxy based on fishing intensity, and model estimate of MSY). Also, the model did not support large 2013 year class previously estimated in the 2019 assessment, which contributed to decrease in of stock productivity and estimated OFLs.

The equilibrium yield plot is shown in Figure 65, based on a steepness value fixed at 0.720. The predicted maximum sustainable yield under the assumptions of this assessment occurs near 27% of equilibrium unfished spawning output, however this represents only 12% higher

yield with considerably more risk than under the 50% SPR policy, which occurs near 45% of unfished spawning output.

4.2 Unresolved problems and major uncertainties

Major sources of uncertainty include landings, discards, natural mortality, and recruitment, which are discussed below.

Because widow rockfish is a marketable species, historical discard rates were likely lower than less desirable or smaller species. This assessment assumed that discarding was nearly negligible before management restrictions began in 1982. Once trip limits were introduced, discarding tended to be an all or none event; rare, but large, discard events are unlikely to have been detected given low observer coverage. From 2002 onward, the WCGOP has provided data on discards from vessels that were randomly selected for observer coverage, thus some uncertainty is present in the total amount discarded. The implementation of trawl rationalization in 2011 resulted in almost 100% observer coverage for the trawl fleet and very little incentive to discard widow rockfish. Discard mortality is assumed to be 100%, which may overestimate actual mortality (Jarvis & Lowe, 2007).

There may also be uncertainty in the ability of bottom trawl surveys to reliably estimate abundances of widow rockfish, which spend a significant portion of their time in mid-water (Wilkins 1986). Multiple surveys are used in the assessment, but further consideration of additional surveys is reasonable.

This assessment attempts to capture uncertainty in M by estimating it inside the model (thereby propagating uncertainty in M into uncertainty in spawning output and other derived quantities), and by incorporating variation in M in a decision table. Model sensitivities and profiles over M showed that current stock status was highly sensitive to the assumption about natural mortality.

Widow rockfish is a relatively long-lived fish, and their natural mortality (M) is likely to be lower than many managed fish stocks (e.g., gadoids). Ages above 50 years have been observed and it is expected that natural mortality could be less than 0.10 yr^{-1} (the median of the prior for natural mortality used in this assessment). However, even with length and age data available back to the late 1970s, M was estimated at 0.135 for females and 0.147 yr^{-1} for males, with a small amount of uncertainty (e.g., a 6.1% coefficient of variation for females).

Notably, the estimated M for both sexes from this assessment are lower than those from the 2015 and 2019 assessments, for which natural mortality was estimated above 0.15 yr^{-1} and 0.14 yr^{-1} , respectively. The estimates of M varied slightly depending on the weight given to age and length data, or removing recent years of data, but M was always estimated

above 0.12 yr^{-1} . The likelihood profile over natural mortality provides support for values up to or above 0.14 yr^{-1} , but with greater curvature than in the 2019 assessment, suggesting additional data has reduced the support for higher natural mortality values. A contributing factor to this change may be the increased mean age and frequency of older fish in catch observed by the WCG BTS survey in recent years. The successful rebounding of the stock as a result of reduced fishing pressure from 2003-2016 may have allowed year classes to age and become fully observed by WCG BTS.

Steepness was fixed at 0.720 in the base model, but a likelihood profile showed that it would be estimated at a value less than that. Estimates of M increased with lower steepness, while unfished spawning output increased and current spawning output decreased. Sustainable yields at the $\text{SPR}_{50\%}$ reference harvest rate ranged from approximately 2641 to 5992 mt depending on the value of steepness.

4.3 Harvest Projections and Decision Tables

Projections for overfishing limit (OFL) (mt), ABC (mt), ACL (mt), the buffer between the OFL and ABC, estimated spawning output (billions of eggs), and fraction of unfished spawning output with adopted OFL and ACL and assumed catch for the first two years of the projection period. The predicted OFL is the calculated total catch determined by $\text{FSPR}=50\%$ are provided in Table 22.

Uncertainty in natural mortality (estimated for both sexes) contributed greatly to uncertainty in the results. Natural mortality is proposed to be used as the axis of uncertainty to define low and high states of nature. This differed from the 2019 assessment which included two more factors, steepness (not estimated in the model) and 2013 recruitment strength. Steepness was fixed in the base model at the mean of a prior has not changed since 2019, and the 2013 year class is no longer a major source of uncertainty, and there is no recent similarly large estimate of recruitment. To define state of nature, the 12.5% and 87.5% quantiles for female and male natural mortality (independently) are proposed to be used as low and high values (0.126 yr^{-1} and 0.144 yr^{-1} for females; 0.137 yr^{-1} and 0.157 yr^{-1} for males).

The decision table with projections under alternative catch streams is to be provided after the review meeting.

4.4 Evaluation of Scientific Uncertainty

This assessment synthesizes many sources of data and estimates recruitment variability, thus it is classified as a Category 1 stock assessment. Therefore, the sigma to determine the catch reduction to account for scientific uncertainty is 0.50.

Spawning is estimated to be at 10,572 billion eggs in 2025, with a sigma of 0.2628. OFL is estimated to be 6,134 mt in 2025 with a coefficient of variation of 0.2934.

4.5 Regional management considerations

Widow rockfish are managed on a coastwide basis and observed more often in the WCGBTS north of latitude 40° 10' N. Bottom trawl catches in California have historically been as large as in Oregon and larger than in Washington, but recently catches in California have been small. Future assessments and management of widow rockfish may want to monitor where catches are being taken to make sure that specific areas are not being overexploited. In addition, research on the connectivity along the coast as well as regional differences would help to inform the potential for overfishing specific areas.

4.6 Research and Data Needs

There are many areas of research that could be improved to benefit the understanding and assessment of widow rockfish, including the following:

- **Natural mortality:** Uncertainty in natural mortality translates into uncertain estimates of status and sustainable fishing levels for widow rockfish. The collection of additional age data, re-reading of older age samples, reading old age samples that are unread, and improved understanding of the life-history of widow rockfish may reduce that uncertainty. Investigating the ageing error and bias would help to understand the influences that the age data have on this assessment, particularly the influence of age data on natural mortality.
- **Sex-specific selectivity:** The midwater and bottom trawl length-composition data fits showed divergent residual patterns between male and female fish. The underlying mechanism driving this pattern is unclear, and could be related to growth, sexing error, or to sex-specific selectivity (e.g., when widow rockfish aggregate, sexes possibly may be aggregating separately). Sex-specific selectivity for these two fleets could be explored or included to address this.

- **Coastwide understanding of stock structure, biology, connectivity, and distribution:** This is a stock assessment for widow rockfish off of the west coast of the U.S. and does not consider data from British Columbia or Alaska. Further investigating and comparing the data and predictions from British Columbia and Alaska to determine if there are similarities with the U.S. West Coast observations would help to define the connectivity between widow rockfish north and south of the U.S.-Canada border. Research into spatial dynamics within the U.S. west coast (as noted above under Unresolved problems and major uncertainties), is also important.

5 Tables

Note: table numbering is continuous with executive summary

5.1 Management performance

Table 8: Recent trend in the OFL, ABC, ACL, total landings, total mortality all in metric tons (mt).

Year	OFL (mt)	ABC (mt)	ACL (mt)	Catch (mt)
2015	4,137	3,929.00	2,000	880.84
2016	3,990	3,790.00	2,000	1,041.65
2017	14,130	13,508.28	13,508	6,356.30
2018	13,237	12,654.57	12,655	10,531.98
2019	12,375	11,830.50	11,831	9,309.60
2020	11,714	11,198.58	11,199	8,400.97
2021	15,749	14,725.32	14,725	10,873.65
2022	14,826	13,788.18	13,788	12,116.29
2023	13,633	12,624.16	12,624	10,996.91
2024	12,453	11,481.67	11,482	9,746.77

5.2 Data

5.2.1 Fishery-dependent data

Table 9: Total removals (mt) used in widow rockfish assessment, by year and by fleet. Total removals represent landings and discards combined. Bottom trawl fleet also includes minimal catches made by net and hook-and-line gears, as well as widow rockfish catches made within historical foreign fishery targeting Pacific Ocean Perch between 1966–1976.

Yr	Bottom trawl	Midwater trawl	Hake
1916	78	0	0
1917	122	0	0
1918	140	0	0
1919	97	0	0
1920	99	0	0
1921	82	0	0
1922	71	0	0
1923	78	0	0
1924	50	0	0
1925	62	0	0
1926	95	0	0
1927	79	0	0
1928	90	0	0
1929	87	0	0
1930	113	0	0
1931	100	0	0
1932	100	0	0
1933	86	0	0
1934	91	0	0
1935	98	0	0

Yr	Bottom trawl	Midwater trawl	Hake
1936	110	0	0
1937	103	0	0
1938	83	0	0
1939	76	0	0
1940	121	0	0
1941	129	0	0
1942	145	0	0
1943	494	0	0
1944	965	0	0
1945	1,635	0	0
1946	1,170	0	0
1947	649	0	0
1948	485	0	0
1949	379	2	0
1950	432	0	0
1951	562	0	0
1952	554	0	0
1953	476	0	0
1954	454	0	0
1955	470	0	0
1956	610	0	0
1957	771	0	0
1958	713	0	0
1959	683	0	0
1960	823	0	0
1961	688	0	0
1962	771	0	0

Yr	Bottom trawl	Midwater trawl	Hake
1963	446	0	0
1964	611	0	0
1965	269	0	0
1966	4,294	0	0
1967	4,828	0	0
1968	2,398	0	0
1969	849	0	0
1970	853	0	0
1971	1,094	0	0
1972	922	0	0
1973	1,163	0	0
1974	907	0	0
1975	846	0	31
1976	1,198	0	130
1977	1,044	0	123
1978	1,426	0	195
1979	1,880	7,573	200
1980	2,924	18,710	272
1981	4,994	23,022	228
1982	6,278	26,022	158
1983	8,561	4,359	131
1984	4,397	8,198	295
1985	3,852	7,500	183
1986	5,174	6,961	258
1987	5,875	10,426	183
1988	5,382	7,791	232
1989	5,914	9,891	212

Yr	Bottom trawl	Midwater trawl	Hake
1990	8,079	5,191	230
1991	5,796	2,406	556
1992	6,159	1,731	448
1993	7,996	2,674	266
1994	5,387	2,540	584
1995	5,685	2,690	429
1996	4,980	2,244	733
1997	5,213	2,871	357
1998	3,616	1,051	587
1999	2,272	2,405	371
2000	106	4,284	284
2001	73	1,983	210
2002	20	253	159
2003	6	14	23
2004	14	37	50
2005	9	46	144
2006	10	47	158
2007	33	24	193
2008	8	79	180
2009	26	96	135
2010	25	72	89
2011	22	60	130
2012	46	62	164
2013	57	268	147
2014	85	331	308
2015	15	480	386
2016	11	590	441

Yr	Bottom trawl	Midwater trawl	Hake
2017	39	4,862	1,455
2018	39	9,412	1,081
2019	31	8,177	1,102
2020	77	7,578	747
2021	108	10,148	617
2022	136	10,861	1,119
2023	90	10,234	673
2024	41	9,172	534

Table 10: Number of trips sampled for length data by gear and state for non-hake fisheries.

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
1971	0	0	0	0	0	2
1977	0	0	0	25	0	0
1978	50	0	0	0	0	0
1979	32	0	0	0	0	0
1980	101	0	0	6	0	19
1981	72	3	0	59	20	31
1982	88	7	0	89	34	41
1983	158	16	0	46	10	25
1984	146	20	0	29	12	22
1985	149	20	0	25	35	16
1986	108	17	0	25	28	27
1987	88	29	0	49	74	36
1988	79	30	7	37	42	14
1989	81	49	14	30	67	16
1990	80	58	11	39	62	30
1991	74	76	20	17	63	15
1992	55	98	22	5	41	9
1993	60	69	28	5	49	8
1994	54	67	13	2	21	16
1995	53	47	17	11	14	16
1996	49	33	17	11	12	13
1997	54	49	16	10	21	19
1998	41	43	26	3	11	8
1999	38	28	21	5	19	11
2000	14	0	3	16	44	19

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
2001	12	6	2	10	38	11
2002	22	8	7	1	15	10
2003	7	0	1	0	0	5
2004	5	1	1	0	0	12
2005	4	2	0	0	0	10
2006	7	3	2	0	0	8
2007	7	16	4	0	0	3
2008	5	18	5	0	0	12
2009	19	30	0	0	0	14
2010	18	22	1	0	0	11
2011	6	14	9	0	1	6
2012	14	19	5	0	4	7
2013	20	21	1	0	6	6
2014	18	20	3	0	5	7
2015	37	23	0	0	18	4
2016	27	14	0	0	7	1
2017	22	41	0	3	33	3
2018	31	25	7	10	60	4
2019	34	33	1	2	48	12
2020	29	18	0	2	31	5
2021	42	18	2	4	39	7
2022	13	10	0	12	46	4
2023	20	7	0	7	51	7
2024	27	13	0	0	52	9

Table 11: Number of lengths of widow rockfish by gear and state for non-hake fisheries.

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
1971	0	0	0	0	0	408
1977	0	0	0	96	0	0
1978	303	0	0	0	0	0
1979	436	0	0	0	0	0
1980	727	0	0	13	0	1,900
1981	444	250	0	1,340	1,746	3,100
1982	932	792	0	3,144	3,960	4,100
1983	1,352	478	0	1,411	321	2,500
1984	1,722	2,394	0	1,278	1,525	2,199
1985	1,853	2,233	0	1,176	3,971	1,600
1986	1,740	1,425	0	1,032	2,788	2,650
1987	998	865	0	1,744	2,198	1,942
1988	763	916	350	1,230	1,239	700
1989	1,007	1,099	700	1,325	1,843	800
1990	1,202	1,320	550	1,510	1,454	1,500
1991	1,596	1,569	997	761	1,442	750
1992	1,470	1,982	1,100	222	1,760	450
1993	1,682	1,410	1,400	231	1,156	400
1994	1,359	1,464	650	112	557	842
1995	1,539	1,066	850	519	296	800
1996	1,364	845	704	437	316	650
1997	2,063	1,231	557	382	620	950
1998	1,368	1,013	865	125	291	400
1999	1,420	727	952	240	514	550
2000	263	0	101	641	1,147	950

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
2001	139	98	2	349	960	550
2002	318	185	136	39	319	500
2003	234	0	46	0	0	208
2004	26	18	3	0	0	508
2005	27	48	0	0	0	399
2006	79	58	7	0	0	461
2007	12	302	104	0	0	250
2008	8	274	76	0	0	1,086
2009	170	316	0	0	0	1,079
2010	205	233	100	0	0	903
2011	32	246	93	0	30	550
2012	136	353	241	0	95	688
2013	153	365	39	0	215	486
2014	134	324	106	0	150	700
2015	263	295	0	0	530	400
2016	143	254	0	0	210	100
2017	316	864	0	158	949	125
2018	645	161	12	507	1,492	350
2019	566	346	50	90	1,149	600
2020	593	228	0	83	759	233
2021	850	226	8	183	890	307
2022	272	185	0	502	1,025	180
2023	376	135	0	316	1,125	571
2024	540	164	0	0	1,255	830

Table 12: Number of hauls or trips and number of lengths sampled from the at-sea hake and shoreside hake fisheries.

Year	Number of hauls (at-sea) or trips (shoreside)		Number of lengths	
	Domestic at-sea	Shoreside	Domestic at-sea	Shoreside
1992	214	0	1,474	0
1993	239	0	1,468	0
1994	361	3	3,458	78
1995	304	19	1,789	570
1996	332	18	2,620	540
1997	397	30	2,841	869
1998	481	32	2,431	975
1999	598	52	3,070	1,551
2000	571	33	2,845	1,004
2001	522	19	1,758	576
2002	369	3	1,204	70
2003	291	2	665	26
2004	512	19	1,670	380
2005	1,228	1	5,538	50
2006	1,295	14	6,104	594
2007	1,491	21	10,658	860
2008	1,138	36	7,324	966
2009	400	24	1,976	845
2010	980	43	4,734	1,214
2011	982	43	3,605	1,286
2012	914	46	4,779	1,291
2013	901	40	3,808	1,160
2014	773	50	3,970	1,452

Year	Number of hauls (at-sea) or trips (shoreside)		Number of lengths	
	Domestic at-sea	Shoreside	Domestic at-sea	Shoreside
2015	522	36	2,312	1,313
2016	801	49	3,934	1,465
2017	997	57	5,406	1,353
2018	461	65	2,245	1,283
2019	469	73	2,642	1,536
2020	214	37	902	839
2021	310	61	1,776	1,279
2022	333	88	1,489	1,745
2023	469	68	1,738	1,525
2024	83	60	251	1,231

Table 13: Number of trips sampled for ages by gear and state for non-hake fisheries.

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
1978	7	0	0	0	0	0
1979	11	0	0	0	0	0
1980	27	0	0	0	0	18
1981	14	3	0	30	20	31
1982	87	6	0	71	34	40
1983	151	16	0	45	10	25
1984	144	20	0	29	12	22
1985	137	20	0	25	33	16
1986	106	17	0	22	28	27
1987	84	27	0	49	62	36
1988	67	29	6	34	41	14
1989	75	49	14	30	66	16
1990	70	58	11	32	62	30
1991	65	76	20	17	63	15
1992	45	93	22	4	26	9
1993	28	67	28	0	49	8
1994	28	67	13	2	21	15
1995	8	45	17	3	13	16
1996	36	32	14	6	11	13
1997	42	46	11	10	20	19
1998	27	42	14	2	11	8
1999	29	27	19	3	18	10
2000	8	0	2	9	42	19
2001	2	6	0	4	35	10
2002	17	8	2	1	15	10

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
2003	3	0	0	0	0	5
2004	3	0	1	0	0	12
2005	0	2	0	0	0	10
2006	6	3	1	0	0	8
2007	6	16	4	0	0	3
2008	5	18	5	0	0	12
2009	8	29	0	0	0	14
2010	7	21	1	0	0	11
2011	0	5	7	0	1	5
2012	0	8	5	0	0	7
2013	0	7	1	0	3	5
2014	0	4	2	0	1	7
2015	0	22	0	0	14	4
2016	0	13	0	0	6	1
2017	0	36	0	0	31	3
2018	0	25	7	0	46	4
2019	0	16	1	0	34	12
2020	0	15	0	0	25	5
2021	0	12	2	0	31	6
2022	0	8	0	0	45	4
2023	0	5	0	0	48	7
2024	0	7	0	0	42	7

Table 14: Number of age samples by gear and state for non-hake fisheries.

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
1978	107	0	0	0	0	0
1979	269	0	0	0	0	0
1980	404	0	0	0	0	1,775
1981	205	109	0	598	600	3,050
1982	834	174	0	2,382	1,019	3,944
1983	1,283	475	0	1,365	321	2,480
1984	1,678	600	0	1,278	360	2,194
1985	1,762	589	0	1,176	963	1,591
1986	1,704	680	0	913	939	2,594
1987	968	805	0	1,742	1,837	1,940
1988	692	886	298	1,132	1,209	695
1989	919	1,099	695	1,323	1,794	799
1990	1,051	1,310	550	1,309	1,447	1,497
1991	1,308	1,566	991	761	1,413	748
1992	676	1,889	1,097	82	574	450
1993	472	1,361	1,398	0	1,155	400
1994	516	1,463	650	54	556	749
1995	167	1,027	850	68	276	800
1996	873	827	699	158	292	649
1997	892	1,164	549	187	593	949
1998	1,019	987	699	82	291	400
1999	1,026	706	950	133	479	500
2000	157	0	100	353	1,067	948
2001	43	98	0	132	858	485
2002	294	179	99	21	319	488

Year	Bottom Trawl			Midwater Trawl		
	CA	OR	WA	CA	OR	WA
2003	87	0	0	0	0	208
2004	7	0	3	0	0	506
2005	0	48	0	0	0	399
2006	74	58	6	0	0	361
2007	11	302	54	0	0	150
2008	8	274	75	0	0	600
2009	81	315	0	0	0	759
2010	54	231	50	0	0	539
2011	0	63	84	0	30	250
2012	0	80	73	0	0	163
2013	0	190	26	0	90	153
2014	0	91	52	0	30	229
2015	0	152	0	0	69	195
2016	0	156	0	0	36	28
2017	0	209	0	0	223	100
2018	0	161	12	0	495	200
2019	0	55	49	0	176	597
2020	0	61	0	0	134	233
2021	0	53	8	0	135	300
2022	0	44	0	0	281	129
2023	0	28	0	0	312	320
2024	0	33	0	0	248	340

5.2.2 Fishery-independent data

Table 15: Stratifications used for the WCGBTS.

Strata	Area (km2)	Depth1	Depth2	Latitude1	Latitude2
A	10,687.86	55	183	34.5	40.5
B	3,394.82	183	400	34.5	40.5
C	23,042.39	55	183	40.5	49.0
D	7,667.81	183	400	40.5	49.0

Table 16: Number of positive tows, lengths, and ages in each year from the Triennial survey (Tri) and the WCGBTS (NW).

Year	Number of positive tows		Number of tows with lengths		Number of lengths		Number of tows with ages		Number of ages	
	Tri	NW	Tri	NW	Tri	NW	Tri	NW	Tri	NW
1977	80		1		9					
1980	38		3		166		1		22	
1983	70		5		385					
1986	46		8		317					
1989	38		20		713					
1992	50		10		708					
1995	43		43		500					
1998	59		58		738					
2001	28		28		130					
2003		20		18		216				
2004	36	12	33	12	219	84		12		43
2005		20		20		78		18		65
2006		26		26		172		26		89
2007		27		27		92		27		83
2008		17		17		26		15		20
2009		31		31		141		31		123
2010		28		28		240		28		116
2011		31		31		313		31		152
2012		32		32		181		32		91
2013		18		18		364		18		246
2014		29		28		349		28		264
2015		21		21		149		21		93
2016		40		40		888		40		556
2017		30		30		310		30		213

Year	Number of positive tows		Number of tows with lengths		Number of lengths		Number of tows with ages		Number of ages	
	Tri	NW	Tri	NW	Tri	NW	Tri	NW	Tri	NW
2018		34		34		410		34		353
2019		23		23		219		23		161
2021		18		17		66		17		66
2022		18		18		125		18		109
2023		30		29		159		29		110
2024		35		35		485		35		347

Table 17: Comparison of key data weighting quantities from the August 2025 (Aug.) and the proposed base model (New). 'Lambda' is the likelihood multiplier applied in the August 2025 model (Lambda = 1 for all fleets in the new proposed base model). 'Weight x lambda' is the product of the estimated data weighting and the lambda. 'Sum N input' is the sum of the input sample sizes across all observations (which differs for BottomTrawl because discard data has been removed). 'Sum N adj.' is that sum after adjustment by the data weighting and lambda. 'CAAL' is conditional age-at-length data. Entries with — indicate that fleet does not exist in the new proposed base model, and the composition data has been removed.

Type	Fleet	Lambda (Aug.)	Weight x lambda (Aug.)	Weight x lambda (New)	Sum N input (Aug.)	Sum N input (New)	Sum N adj. x lambda (Aug.)	Sum N adj. x lambda (New)
Length	BottomTrawl	0.5	0.028	0.055	14976.0	14336.4	422.2	785.0
Length	MidwaterTrawl	0.5	0.104	0.039	14908.6	14908.6	1547.2	587.2
Length	Hake	0.5	0.055	0.022	23864.6	23864.6	1317.5	532.2
Length	Net	0.5	0.248	—	1196.8	—	296.4	—
Length	HnL	0.5	0.186	—	1033.2	—	193.2	—
Length	Triennial	1.0	0.372	0.092	459.0	459.0	171.0	42.1
Length	WCG BTS	1.0	0.635	0.083	989.1	989.1	628.2	82.1
Age	BottomTrawl	0.5	0.083	0.194	8289.2	8289.2	688.3	1610.2
Age	MidwaterTrawl	0.5	0.140	0.151	8358.2	8358.2	1168.2	1258.1
Age	Hake	0.5	0.120	0.262	6051.2	6051.2	730.1	1582.5
Age	Net	0.5	0.250	—	737.8	—	184.7	—
Age	HnL	0.5	0.276	—	125.4	—	34.7	—
CAAL	WCG BTS	1.0	0.296	0.106	3243.6	3243.6	963.5	346.2

5.3 Model results

Table 18: Parameter estimates, parameter bounds (low, high), estimation status, estimated standard deviation (SD), prior information [distribution(mean, SD)] used in the base model.

Label	Value	Bounds	Status	SD	Prior
NatM_uniform_Fem_GP_1	0.135	(0.01, 0.3)	ok	0.00824	lognormal(0.100, 0.310)
L_at_Amin_Fem_GP_1	18.9	(10, 40)	ok	0.852	none
L_at_Amax_Fem_GP_1	49.4	(35, 60)	ok	0.365	none
VonBert_K_Fem_GP_1	0.194	(0.01, 0.4)	ok	0.00941	none
CV_young_Fem_GP_1	0.142	(0.01, 0.4)	ok	0.0185	none
CV_old_Fem_GP_1	0.0471	(0.01, 0.4)	ok	0.00397	none
Wtlen_1_Fem_GP_1	1.59e-05	(-3, 3)	fixed	0	none
Wtlen_2_Fem_GP_1	2.99	(-3, 10)	fixed	0	none
Mat50%_Fem_GP_1	5.47	(-3, 50)	fixed	0	none
Mat_slope_Fem_GP_1	-0.775	(-3, 3)	fixed	0	none
Eggs_scalar_Fem_GP_1	1.11e-08	(-1, 1)	fixed	0	none
Eggs_exp_len_Fem_GP_1	4.54	(0, 5)	fixed	0	none
NatM_uniform_Mal_GP_1	0.147	(0.01, 0.3)	ok	0.00867	lognormal(0.100, 0.310)
L_at_Amin_Mal_GP_1	20.3	(10, 40)	ok	0.639	none
L_at_Amax_Mal_GP_1	43.6	(35, 60)	ok	0.313	none
VonBert_K_Mal_GP_1	0.267	(0.01, 0.4)	ok	0.0135	none
CV_young_Mal_GP_1	0.0915	(0.01, 0.4)	ok	0.0113	none

Label	Value	Bounds	Status	SD	Prior
CV_old_Mal_GP_1	0.0548	(0.01, 0.4)	ok	0.00392	none
Wtlen_1_Mal_GP_1	1.45e-05	(-3, 3)	fixed	0	none
Wtlen_2_Mal_GP_1	3.01	(-3, 10)	fixed	0	none
CohortGrowDev	1	(0, 2)	fixed	0	none
FracFemale_GP_1	0.5	(1e-06, 1)	fixed	0	none
SR_LN(R0)	10.6	(6, 15)	ok	0.155	none
SR_BH_steep	0.72	(0.2, 1)	fixed	0	beta(0.720, 0.160)
SR_sigmaR	0.6	(0, 2)	fixed	0	none
SR_regime	0	(-5, 5)	fixed	0	none
SR_autocorr	0	(0, 0.5)	fixed	0	none
Early_InitAge_16	-0.00131	(-5, 5)	dev	0.6	normal(0.00, 0.60)
Early_InitAge_15	-0.00149	(-5, 5)	dev	0.6	normal(0.00, 0.60)
Early_InitAge_14	-0.0017	(-5, 5)	dev	0.6	normal(0.00, 0.60)
Early_InitAge_13	-0.00192	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_12	-0.00217	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_11	-0.00245	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_10	-0.00276	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_9	-0.00309	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_8	-0.00345	(-5, 5)	dev	0.599	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_InitAge_7	-0.00384	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_6	-0.00427	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_5	-0.00472	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_4	-0.00522	(-5, 5)	dev	0.599	normal(0.00, 0.60)
Early_InitAge_3	-0.00577	(-5, 5)	dev	0.598	normal(0.00, 0.60)
Early_InitAge_2	-0.00638	(-5, 5)	dev	0.598	normal(0.00, 0.60)
Early_InitAge_1	-0.00705	(-5, 5)	dev	0.598	normal(0.00, 0.60)
Early_RecrDev_1916	-0.00779	(-5, 5)	dev	0.598	normal(0.00, 0.60)
Early_RecrDev_1917	-0.00861	(-5, 5)	dev	0.598	normal(0.00, 0.60)
Early_RecrDev_1918	-0.00952	(-5, 5)	dev	0.597	normal(0.00, 0.60)
Early_RecrDev_1919	-0.0105	(-5, 5)	dev	0.597	normal(0.00, 0.60)
Early_RecrDev_1920	-0.0116	(-5, 5)	dev	0.597	normal(0.00, 0.60)
Early_RecrDev_1921	-0.0128	(-5, 5)	dev	0.596	normal(0.00, 0.60)
Early_RecrDev_1922	-0.0142	(-5, 5)	dev	0.596	normal(0.00, 0.60)
Early_RecrDev_1923	-0.0157	(-5, 5)	dev	0.596	normal(0.00, 0.60)
Early_RecrDev_1924	-0.0173	(-5, 5)	dev	0.595	normal(0.00, 0.60)
Early_RecrDev_1925	-0.0191	(-5, 5)	dev	0.595	normal(0.00, 0.60)
Early_RecrDev_1926	-0.0211	(-5, 5)	dev	0.594	normal(0.00, 0.60)
Early_RecrDev_1927	-0.0234	(-5, 5)	dev	0.593	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_RecrDev_1928	-0.0258	(-5, 5)	dev	0.593	normal(0.00, 0.60)
Early_RecrDev_1929	-0.0285	(-5, 5)	dev	0.592	normal(0.00, 0.60)
Early_RecrDev_1930	-0.0314	(-5, 5)	dev	0.591	normal(0.00, 0.60)
Early_RecrDev_1931	-0.0347	(-5, 5)	dev	0.59	normal(0.00, 0.60)
Early_RecrDev_1932	-0.0383	(-5, 5)	dev	0.589	normal(0.00, 0.60)
Early_RecrDev_1933	-0.0423	(-5, 5)	dev	0.588	normal(0.00, 0.60)
Early_RecrDev_1934	-0.0466	(-5, 5)	dev	0.587	normal(0.00, 0.60)
Early_RecrDev_1935	-0.0515	(-5, 5)	dev	0.586	normal(0.00, 0.60)
Early_RecrDev_1936	-0.0568	(-5, 5)	dev	0.584	normal(0.00, 0.60)
Early_RecrDev_1937	-0.0627	(-5, 5)	dev	0.583	normal(0.00, 0.60)
Early_RecrDev_1938	-0.0693	(-5, 5)	dev	0.581	normal(0.00, 0.60)
Early_RecrDev_1939	-0.0768	(-5, 5)	dev	0.579	normal(0.00, 0.60)
Early_RecrDev_1940	-0.0851	(-5, 5)	dev	0.577	normal(0.00, 0.60)
Early_RecrDev_1941	-0.0942	(-5, 5)	dev	0.574	normal(0.00, 0.60)
Early_RecrDev_1942	-0.104	(-5, 5)	dev	0.572	normal(0.00, 0.60)
Early_RecrDev_1943	-0.115	(-5, 5)	dev	0.569	normal(0.00, 0.60)
Early_RecrDev_1944	-0.126	(-5, 5)	dev	0.566	normal(0.00, 0.60)
Early_RecrDev_1945	-0.137	(-5, 5)	dev	0.563	normal(0.00, 0.60)
Early_RecrDev_1946	-0.148	(-5, 5)	dev	0.561	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_RecrDev_1947	-0.159	(-5, 5)	dev	0.558	normal(0.00, 0.60)
Early_RecrDev_1948	-0.169	(-5, 5)	dev	0.555	normal(0.00, 0.60)
Early_RecrDev_1949	-0.178	(-5, 5)	dev	0.553	normal(0.00, 0.60)
Early_RecrDev_1950	-0.185	(-5, 5)	dev	0.551	normal(0.00, 0.60)
Early_RecrDev_1951	-0.189	(-5, 5)	dev	0.55	normal(0.00, 0.60)
Early_RecrDev_1952	-0.188	(-5, 5)	dev	0.55	normal(0.00, 0.60)
Early_RecrDev_1953	-0.179	(-5, 5)	dev	0.551	normal(0.00, 0.60)
Early_RecrDev_1954	-0.161	(-5, 5)	dev	0.554	normal(0.00, 0.60)
Early_RecrDev_1955	-0.131	(-5, 5)	dev	0.558	normal(0.00, 0.60)
Early_RecrDev_1956	-0.0904	(-5, 5)	dev	0.565	normal(0.00, 0.60)
Early_RecrDev_1957	-0.043	(-5, 5)	dev	0.573	normal(0.00, 0.60)
Early_RecrDev_1958	-0.00283	(-5, 5)	dev	0.578	normal(0.00, 0.60)
Early_RecrDev_1959	0.0118	(-5, 5)	dev	0.578	normal(0.00, 0.60)
Early_RecrDev_1960	-0.00448	(-5, 5)	dev	0.571	normal(0.00, 0.60)
Early_RecrDev_1961	-0.0304	(-5, 5)	dev	0.562	normal(0.00, 0.60)
Early_RecrDev_1962	-0.0321	(-5, 5)	dev	0.556	normal(0.00, 0.60)
Early_RecrDev_1963	0.0125	(-5, 5)	dev	0.555	normal(0.00, 0.60)
Early_RecrDev_1964	0.0819	(-5, 5)	dev	0.558	normal(0.00, 0.60)
Early_RecrDev_1965	0.137	(-5, 5)	dev	0.559	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Early_RecrDev_1966	0.191	(-5, 5)	dev	0.549	normal(0.00, 0.60)
Early_RecrDev_1967	0.199	(-5, 5)	dev	0.533	normal(0.00, 0.60)
Early_RecrDev_1968	0.192	(-5, 5)	dev	0.489	normal(0.00, 0.60)
Early_RecrDev_1969	-0.0265	(-5, 5)	dev	0.528	normal(0.00, 0.60)
Main_RecrDev_1970	1.37	(-5, 5)	dev	0.25	normal(0.00, 0.60)
Main_RecrDev_1971	1.22	(-5, 5)	dev	0.228	normal(0.00, 0.60)
Main_RecrDev_1972	-0.833	(-5, 5)	dev	0.402	normal(0.00, 0.60)
Main_RecrDev_1973	-0.929	(-5, 5)	dev	0.351	normal(0.00, 0.60)
Main_RecrDev_1974	-0.584	(-5, 5)	dev	0.301	normal(0.00, 0.60)
Main_RecrDev_1975	0.158	(-5, 5)	dev	0.177	normal(0.00, 0.60)
Main_RecrDev_1976	-1.18	(-5, 5)	dev	0.319	normal(0.00, 0.60)
Main_RecrDev_1977	0.673	(-5, 5)	dev	0.128	normal(0.00, 0.60)
Main_RecrDev_1978	1.02	(-5, 5)	dev	0.112	normal(0.00, 0.60)
Main_RecrDev_1979	-0.0252	(-5, 5)	dev	0.176	normal(0.00, 0.60)
Main_RecrDev_1980	0.451	(-5, 5)	dev	0.147	normal(0.00, 0.60)
Main_RecrDev_1981	1.03	(-5, 5)	dev	0.118	normal(0.00, 0.60)
Main_RecrDev_1982	0.409	(-5, 5)	dev	0.153	normal(0.00, 0.60)
Main_RecrDev_1983	-0.19	(-5, 5)	dev	0.195	normal(0.00, 0.60)
Main_RecrDev_1984	0.45	(-5, 5)	dev	0.125	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Main_RecrDev_1985	0.283	(-5, 5)	dev	0.129	normal(0.00, 0.60)
Main_RecrDev_1986	-0.626	(-5, 5)	dev	0.229	normal(0.00, 0.60)
Main_RecrDev_1987	0.205	(-5, 5)	dev	0.139	normal(0.00, 0.60)
Main_RecrDev_1988	-0.158	(-5, 5)	dev	0.177	normal(0.00, 0.60)
Main_RecrDev_1989	-0.391	(-5, 5)	dev	0.201	normal(0.00, 0.60)
Main_RecrDev_1990	-0.00759	(-5, 5)	dev	0.162	normal(0.00, 0.60)
Main_RecrDev_1991	0.652	(-5, 5)	dev	0.108	normal(0.00, 0.60)
Main_RecrDev_1992	-0.23	(-5, 5)	dev	0.186	normal(0.00, 0.60)
Main_RecrDev_1993	0.101	(-5, 5)	dev	0.165	normal(0.00, 0.60)
Main_RecrDev_1994	-0.0208	(-5, 5)	dev	0.196	normal(0.00, 0.60)
Main_RecrDev_1995	-0.407	(-5, 5)	dev	0.271	normal(0.00, 0.60)
Main_RecrDev_1996	-0.608	(-5, 5)	dev	0.339	normal(0.00, 0.60)
Main_RecrDev_1997	-0.149	(-5, 5)	dev	0.299	normal(0.00, 0.60)
Main_RecrDev_1998	0.462	(-5, 5)	dev	0.216	normal(0.00, 0.60)
Main_RecrDev_1999	0.524	(-5, 5)	dev	0.201	normal(0.00, 0.60)
Main_RecrDev_2000	0.585	(-5, 5)	dev	0.165	normal(0.00, 0.60)
Main_RecrDev_2001	-0.354	(-5, 5)	dev	0.25	normal(0.00, 0.60)
Main_RecrDev_2002	-0.33	(-5, 5)	dev	0.217	normal(0.00, 0.60)
Main_RecrDev_2003	-0.273	(-5, 5)	dev	0.23	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Main_RecrDev_2004	0.754	(-5, 5)	dev	0.117	normal(0.00, 0.60)
Main_RecrDev_2005	-1.03	(-5, 5)	dev	0.341	normal(0.00, 0.60)
Main_RecrDev_2006	0.689	(-5, 5)	dev	0.123	normal(0.00, 0.60)
Main_RecrDev_2007	-1.15	(-5, 5)	dev	0.379	normal(0.00, 0.60)
Main_RecrDev_2008	1.13	(-5, 5)	dev	0.112	normal(0.00, 0.60)
Main_RecrDev_2009	-0.759	(-5, 5)	dev	0.341	normal(0.00, 0.60)
Main_RecrDev_2010	0.561	(-5, 5)	dev	0.159	normal(0.00, 0.60)
Main_RecrDev_2011	-0.941	(-5, 5)	dev	0.341	normal(0.00, 0.60)
Main_RecrDev_2012	-1.29	(-5, 5)	dev	0.352	normal(0.00, 0.60)
Main_RecrDev_2013	0.414	(-5, 5)	dev	0.174	normal(0.00, 0.60)
Main_RecrDev_2014	-0.0184	(-5, 5)	dev	0.237	normal(0.00, 0.60)
Main_RecrDev_2015	-0.412	(-5, 5)	dev	0.305	normal(0.00, 0.60)
Main_RecrDev_2016	1.33	(-5, 5)	dev	0.162	normal(0.00, 0.60)
Main_RecrDev_2017	0.339	(-5, 5)	dev	0.27	normal(0.00, 0.60)
Main_RecrDev_2018	-0.69	(-5, 5)	dev	0.376	normal(0.00, 0.60)
Main_RecrDev_2019	-0.774	(-5, 5)	dev	0.419	normal(0.00, 0.60)
Main_RecrDev_2020	-0.446	(-5, 5)	dev	0.494	normal(0.00, 0.60)
Late_RecrDev_2021	0.0227	(-5, 5)	dev	0.531	normal(0.00, 0.60)
Late_RecrDev_2022	0.00382	(-5, 5)	dev	0.559	normal(0.00, 0.60)

Label	Value	Bounds	Status	SD	Prior
Late_RecrDev_2023	0.248	(-5, 5)	dev	0.57	normal(0.00, 0.60)
Late_RecrDev_2024	0.136	(-5, 5)	dev	0.564	normal(0.00, 0.60)
ForeRecr_2025	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2026	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2027	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2028	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2029	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2030	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2031	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2032	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2033	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2034	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2035	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
ForeRecr_2036	0	(-5, 5)	dev	0.6	normal(0.00, 0.60)
LnQ_base_BottomTrawl(1)	-6.17	(-25, 25)	fixed	0	none
Q_extraSD_BottomTrawl(1)	0.204	(0, 2)	ok	0.0662	none
LnQ_base_Hake(3)	-11.1	(-20, 2)	ok	0.183	none
Q_extraSD_Hake(3)	0.386	(0, 2)	ok	0.0891	none
LnQ_base_JuvSurvey(6)	-5.04	(-25, 25)	fixed	0	none

Label	Value	Bounds	Status	SD	Prior
Q_extraSD_JuvSurvey(6)	1.27	(0, 2)	ok	0.314	none
LnQ_base_Triennial(7)	-1.88	(-4, 4)	ok	0.505	none
Q_extraSD_Triennial(7)	0	(0, 2)	fixed	0	none
LnQ_base_WCGBTS(8)	-3.34	(-25, 25)	fixed	0	none
Q_extraSD_WCGBTS(8)	0	(0, 2)	fixed	0	none
LnQ_base_ForeignAtSea(9)	-11.5	(-25, 25)	fixed	0	none
Q_extraSD_ForeignAtSea(9)	0.595	(0, 2)	ok	0.155	none
LnQ_base_Hake(3)_BLK10add_1991	0.6	(1e-04, 2)	ok	0.228	normal(0.500, 0.500)
LnQ_base_Triennial(7)_BLK9add_1995	0.249	(1e-04, 2)	ok	0.358	normal(0.500, 0.500)
Size_DbIN_peak_BottomTrawl(1)	42.1	(10, 59)	ok	1.71	none
Size_DbIN_top_logit_BottomTrawl(1)	2.5	(-5, 10)	ok	167	none
Size_DbIN_ascend_se_BottomTrawl(1)	4.06	(-4, 12)	ok	0.326	none
Size_DbIN_descend_se_BottomTrawl(1)	9	(-2, 10)	fixed	0	none
Size_DbIN_start_logit_BottomTrawl(1)	-9	(-9, 10)	fixed	0	none
Size_DbIN_end_logit_BottomTrawl(1)	8	(-9, 9)	fixed	0	none
Size_DbIN_peak_MidwaterTrawl(2)	37.1	(10, 59)	ok	1.2	none
Size_DbIN_top_logit_MidwaterTrawl(2)	-9.25	(-10, 10)	ok	18.1	none
Size_DbIN_ascend_se_MidwaterTrawl(2)	2.8	(-4, 12)	ok	0.495	none
Size_DbIN_descend_se_MidwaterTrawl(2)	3.94	(-2, 10)	ok	1.55	none

Label	Value	Bounds	Status	SD	Prior
Size_DbIN_start_logit_MidwaterTrawl(2)	-9	(-9, 10)	fixed	0	none
Size_DbIN_end_logit_MidwaterTrawl(2)	-1.17	(-9, 9)	ok	2.02	none
Size_DbIN_peak_Hake(3)	34.5	(10, 59)	ok	2.09	none
Size_DbIN_top_logit_Hake(3)	2.5	(-5, 10)	ok	168	none
Size_DbIN_ascend_se_Hake(3)	2.51	(-4, 12)	ok	1.03	none
Size_DbIN_descend_se_Hake(3)	9	(-2, 10)	fixed	0	none
Size_DbIN_start_logit_Hake(3)	-9	(-9, 10)	fixed	0	none
Size_DbIN_end_logit_Hake(3)	8	(-9, 9)	fixed	0	none
SizeSpline_Code_Triennial(7)	0	(0, 2)	fixed	0	none
SizeSpline_GradLo_Triennial(7)	0.127	(-0.001, 1)	ok	0.0677	none
SizeSpline_GradHi_Triennial(7)	0.094	(-1, 1)	ok	0.202	none
SizeSpline_Knot_1_Triennial(7)	24	(8, 56)	fixed	0	none
SizeSpline_Knot_2_Triennial(7)	34	(8, 56)	fixed	0	none
SizeSpline_Knot_3_Triennial(7)	48	(8, 56)	fixed	0	none
SizeSpline_Val_1_Triennial(7)	-2.17	(-10, 10)	ok	0.756	none
SizeSpline_Val_2_Triennial(7)	-1	(-10, 10)	fixed	0	none
SizeSpline_Val_3_Triennial(7)	0.435	(-10, 10)	ok	0.509	none
SizeSpline_Code_WCGBTS(8)	0	(0, 2)	fixed	0	none
SizeSpline_GradLo_WCGBTS(8)	0.447	(-0.001, 1)	ok	0.299	none

Label	Value	Bounds	Status	SD	Prior
SizeSpline_GradHi_WCGBTS(8)	-0.098	(-1, 1)	ok	0.151	none
SizeSpline_Knot_1_WCGBTS(8)	24	(8, 56)	fixed	0	none
SizeSpline_Knot_2_WCGBTS(8)	34	(8, 56)	fixed	0	none
SizeSpline_Knot_3_WCGBTS(8)	48	(8, 56)	fixed	0	none
SizeSpline_Val_1_WCGBTS(8)	-2.19	(-10, 10)	ok	0.665	none
SizeSpline_Val_2_WCGBTS(8)	-1	(-10, 10)	fixed	0	none
SizeSpline_Val_3_WCGBTS(8)	0.05	(-10, 10)	ok	0.354	none
minage@sel=1_JuvSurvey(6)	0	(0, 1)	fixed	0	none
maxage@sel=1_JuvSurvey(6)	0	(0, 1)	fixed	0	none
minage@sel=1_WCGBTS(8)	0	(0, 1)	fixed	0	none
maxage@sel=1_WCGBTS(8)	40	(0, 50)	fixed	0	none
Size_DbIN_peak_BottomTrawl(1)_BLK4repl_1916	39.4	(10, 59)	ok	0.604	none
Size_DbIN_ascend_se_BottomTrawl(1)_BLK4repl_-1916	3.47	(-4, 12)	ok	0.161	none
Size_DbIN_peak_MidwaterTrawl(2)_BLK7repl_1916	38.6	(10, 59)	ok	1.32	none
Size_DbIN_peak_MidwaterTrawl(2)_BLK7repl_1983	38.2	(10, 59)	ok	0.593	none
Size_DbIN_peak_MidwaterTrawl(2)_BLK7repl_2002	37.8	(10, 59)	ok	2.19	none
Size_DbIN_ascend_se_MidwaterTrawl(2)_BLK7repl_-1916	3.39	(-4, 12)	ok	0.381	none

Label	Value	Bounds	Status	SD	Prior
Size_DbIN_ascend_se_MidwaterTrawl(2)_BLK7repl_1983	2.99	(-4, 12)	ok	0.191	none
Size_DbIN_ascend_se_MidwaterTrawl(2)_BLK7repl_2002	2.96	(-4, 12)	ok	0.715	none
Size_DbIN_descend_se_MidwaterTrawl(2)_BLK7repl_1916	4.11	(-2, 10)	ok	1.35	none
Size_DbIN_descend_se_MidwaterTrawl(2)_BLK7repl_1983	2.57	(-2, 10)	ok	0.82	none
Size_DbIN_descend_se_MidwaterTrawl(2)_BLK7repl_2002	-1.82	(-2, 10)	ok	5.29	none
Size_DbIN_end_logit_MidwaterTrawl(2)_BLK7repl_1916	-1.67	(-9, 9)	ok	3.15	none
Size_DbIN_end_logit_MidwaterTrawl(2)_BLK7repl_1983	-0.609	(-9, 9)	ok	0.307	none
Size_DbIN_end_logit_MidwaterTrawl(2)_BLK7repl_2002	2	(-9, 9)	ok	2.97	none
Size_DbIN_peak_Hake(3)_BLK11repl_1916	41.4	(10, 59)	ok	0.67	none
Size_DbIN_top_logit_Hake(3)_BLK11repl_1916	2.5	(-5, 10)	ok	168	none
Size_DbIN_ascend_se_Hake(3)_BLK11repl_1916	3.35	(-4, 12)	ok	0.18	none

Table 19: Time series of population estimates from the base case model.

Year	Total Biomass (mt)	Spawning output (billions of eggs)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploita- tion Rate
1916	161,653	21,079	156,039	0.999	40,150	78	0.011	0.001
1917	161,529	21,062	155,919	0.998	40,114	122	0.018	0.001
1918	161,366	21,040	155,761	0.997	40,074	140	0.021	0.001
1919	161,191	21,015	155,590	0.995	40,029	97	0.014	0.001
1920	161,061	20,997	155,466	0.995	39,982	99	0.015	0.001
1921	160,930	20,979	155,341	0.994	39,930	82	0.012	0.001
1922	160,814	20,964	155,232	0.993	39,873	71	0.011	0.000
1923	160,705	20,951	155,130	0.992	39,811	78	0.012	0.001
1924	160,584	20,936	155,016	0.992	39,743	50	0.007	0.000
1925	160,483	20,926	154,923	0.991	39,669	62	0.009	0.000
1926	160,360	20,913	154,810	0.991	39,587	95	0.014	0.001
1927	160,196	20,894	154,656	0.990	39,496	79	0.012	0.001
1928	160,039	20,877	154,510	0.989	39,397	90	0.013	0.001
1929	159,859	20,857	154,342	0.988	39,288	87	0.013	0.001
1930	159,669	20,837	154,166	0.987	39,168	113	0.017	0.001
1931	159,441	20,811	153,952	0.986	39,035	100	0.015	0.001
1932	159,210	20,787	153,738	0.985	38,890	100	0.015	0.001
1933	158,963	20,760	153,508	0.983	38,732	86	0.013	0.001
1934	158,708	20,734	153,273	0.982	38,558	91	0.014	0.001
1935	158,426	20,705	153,013	0.981	38,368	98	0.015	0.001
1936	158,113	20,673	152,723	0.979	38,158	110	0.016	0.001
1937	157,761	20,637	152,396	0.978	37,926	103	0.015	0.001
1938	157,385	20,598	152,049	0.976	37,668	83	0.012	0.001
1939	156,995	20,560	151,690	0.974	37,381	76	0.011	0.001

Year	Total Biomass (mt)	Spawning output (billions of eggs)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploita- tion Rate
1940	156,573	20,519	151,303	0.972	37,066	121	0.018	0.001
1941	156,065	20,467	150,833	0.970	36,720	129	0.020	0.001
1942	155,503	20,410	150,314	0.967	36,342	145	0.022	0.001
1943	154,877	20,345	149,735	0.964	35,942	494	0.073	0.003
1944	153,870	20,224	148,778	0.958	35,536	965	0.140	0.006
1945	152,386	20,030	147,349	0.949	35,111	1,635	0.231	0.011
1946	150,273	19,738	145,293	0.935	34,674	1,170	0.172	0.008
1947	148,649	19,520	143,727	0.925	34,265	649	0.099	0.005
1948	147,527	19,381	142,663	0.918	33,889	485	0.076	0.003
1949	146,524	19,266	141,719	0.913	33,556	381	0.060	0.003
1950	145,570	19,161	140,819	0.908	33,295	432	0.068	0.003
1951	144,510	19,044	139,807	0.902	33,146	562	0.089	0.004
1952	143,285	18,900	138,621	0.895	33,165	554	0.088	0.004
1953	142,049	18,751	137,410	0.888	33,427	476	0.077	0.003
1954	140,893	18,608	136,258	0.881	34,011	454	0.074	0.003
1955	139,793	18,464	135,132	0.875	35,000	470	0.077	0.003
1956	138,759	18,314	134,031	0.868	36,431	610	0.100	0.005
1957	137,738	18,144	132,894	0.859	38,160	771	0.126	0.006
1958	136,799	17,956	131,784	0.851	39,678	713	0.118	0.005
1959	136,235	17,790	131,011	0.843	40,221	683	0.115	0.005
1960	136,076	17,651	130,649	0.836	39,536	823	0.137	0.006
1961	136,179	17,523	130,627	0.830	38,492	688	0.117	0.005
1962	136,775	17,459	131,225	0.827	38,409	771	0.130	0.006
1963	137,557	17,433	132,099	0.826	39,883	446	0.077	0.003
1964	138,806	17,506	133,400	0.829	41,870	611	0.103	0.005
1965	139,961	17,598	134,463	0.834	43,338	269	0.046	0.002

Year	Total Biomass (mt)	Spawning output (billions of eggs)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploita- tion Rate
1966	141,515	17,767	135,790	0.842	44,813	4,294	0.565	0.032
1967	139,388	17,366	133,427	0.823	44,144	4,828	0.626	0.036
1968	137,219	16,920	131,100	0.801	42,755	2,398	0.365	0.018
1969	137,810	16,867	131,665	0.799	33,638	849	0.143	0.006
1970	140,465	17,081	134,144	0.809	132,665	853	0.142	0.006
1971	144,015	17,333	136,657	0.821	115,417	1,094	0.176	0.008
1972	148,722	17,582	138,742	0.833	14,778	922	0.147	0.007
1973	154,967	17,897	139,575	0.848	13,445	1,163	0.180	0.008
1974	163,294	18,278	153,365	0.866	19,026	907	0.140	0.006
1975	172,030	18,930	169,843	0.897	40,135	877	0.130	0.005
1976	177,385	19,921	174,863	0.944	10,545	1,328	0.178	0.008
1977	178,784	21,043	175,189	0.997	67,871	1,167	0.146	0.007
1978	177,862	22,027	172,780	1.043	96,292	1,621	0.191	0.009
1979	175,689	22,514	170,390	1.066	33,987	9,653	0.806	0.057
1980	165,542	21,529	155,511	1.020	54,487	21,906	1.369	0.141
1981	145,738	18,616	135,524	0.882	95,410	28,244	1.609	0.208
1982	123,718	14,792	117,077	0.701	50,092	32,458	1.758	0.277
1983	100,648	10,753	91,564	0.509	26,190	13,051	1.471	0.143
1984	99,416	9,717	89,273	0.460	48,775	12,890	1.461	0.144
1985	99,603	9,292	93,651	0.440	40,931	11,535	1.383	0.123
1986	100,260	9,234	95,486	0.437	16,462	12,393	1.400	0.130
1987	98,450	9,212	92,532	0.436	37,769	16,484	1.520	0.178
1988	91,405	8,784	86,810	0.416	26,017	13,404	1.435	0.154
1989	85,979	8,498	82,638	0.403	20,481	16,017	1.552	0.194
1990	76,555	7,698	72,080	0.365	29,400	13,501	1.528	0.187
1991	69,298	6,971	65,711	0.330	55,560	8,759	1.363	0.133

Year	Total Biomass (mt)	Spawning output (billions of eggs)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploita- tion Rate
1992	66,407	6,690	62,609	0.317	22,763	8,338	1.364	0.133
1993	63,909	6,371	58,834	0.302	31,286	10,935	1.524	0.186
1994	59,272	5,685	53,376	0.269	26,847	8,511	1.460	0.159
1995	57,816	5,292	54,246	0.251	17,867	8,803	1.513	0.162
1996	55,913	4,920	52,019	0.233	14,290	7,957	1.486	0.153
1997	54,579	4,762	51,421	0.226	22,377	8,441	1.507	0.164
1998	52,260	4,638	49,740	0.220	40,872	5,255	1.270	0.106
1999	52,422	4,826	49,539	0.229	44,046	5,048	1.203	0.102
2000	52,554	4,992	48,217	0.236	47,344	4,674	1.111	0.097
2001	53,361	5,124	47,520	0.243	18,657	2,265	0.710	0.048
2002	57,404	5,445	51,638	0.258	19,454	432	0.199	0.008
2003	63,793	5,997	58,922	0.284	21,170	43	0.020	0.001
2004	70,648	6,708	67,787	0.318	60,944	101	0.041	0.001
2005	76,532	7,532	73,117	0.357	10,504	199	0.072	0.003
2006	81,633	8,395	77,054	0.398	60,168	215	0.068	0.003
2007	86,164	9,204	79,957	0.436	9,736	250	0.073	0.003
2008	91,308	9,899	87,447	0.469	96,802	268	0.073	0.003
2009	96,025	10,518	89,298	0.498	14,774	257	0.067	0.003
2010	101,612	11,117	96,410	0.527	55,845	186	0.047	0.002
2011	106,926	11,740	97,798	0.556	12,540	212	0.051	0.002
2012	113,179	12,384	109,498	0.587	8,954	273	0.062	0.002
2013	118,105	13,082	112,865	0.620	49,443	472	0.099	0.004
2014	121,949	13,820	119,665	0.655	32,350	723	0.141	0.006
2015	123,844	14,522	120,504	0.688	21,973	881	0.160	0.007
2016	124,597	15,089	118,504	0.715	126,249	1,042	0.182	0.009
2017	125,747	15,436	120,077	0.731	47,765	6,356	0.789	0.053

Year	Total Biomass (mt)	Spawning output (billions of eggs)	Total Biomass 4+ (mt)	Fraction Unfished	Age-0 Recruits (1,000s)	Total Mortality (mt)	(1-SPR)/(1- SPR_50%)	Exploita- tion Rate
2018	122,629	14,925	114,902	0.707	17,416	10,532	1.151	0.092
2019	116,555	13,816	104,377	0.654	16,236	9,310	1.141	0.089
2020	114,538	12,948	109,729	0.613	22,868	8,401	1.127	0.077
2021	113,380	12,457	110,824	0.590	37,217	10,874	1.277	0.098
2022	108,443	12,033	105,476	0.570	37,214	12,116	1.301	0.115
2023	100,976	11,644	96,857	0.552	48,077	10,997	1.238	0.114
2024	93,859	11,195	88,467	0.530	42,674	9,747	1.214	0.110
2025	88,274	10,572	82,550	0.501	36,894	10,669	1.351	0.129
2026	82,615	9,630	76,358	0.456	36,267	9,824	1.385	0.129
2027	79,012	8,799	73,427	0.417	35,624	4,596	0.958	0.063
2028	81,326	8,763	76,229	0.415	35,594	4,810	0.955	0.063
2029	83,523	8,870	78,501	0.420	35,682	5,137	0.952	0.065
2030	85,224	9,054	80,251	0.429	35,832	5,409	0.949	0.067
2031	86,366	9,246	81,389	0.438	35,981	5,545	0.946	0.068
2032	87,094	9,403	82,102	0.445	36,101	5,585	0.943	0.068
2033	87,573	9,521	82,560	0.451	36,187	5,572	0.941	0.067
2034	87,928	9,606	82,897	0.455	36,250	5,535	0.937	0.067
2035	88,237	9,672	83,192	0.458	36,297	5,504	0.935	0.066
2036	88,530	9,726	83,475	0.461	36,335	5,480	0.932	0.066

5.4 Model diagnostics

Table 20: Sensitivity analysis results: differences in negative log-likelihood, estimates of key parameters, and estimates of derived quantities between the base model and several alternative models (columns). See main text for details on each sensitivity analysis. Red values indicate negative log-likelihoods that were lower (fit better to that component) than the base model.

Label	January 2026 base model	M=0.1 (the median of a prior)	M fixed at 2019 model estimates	M fixed at 2015 model estimates	M prior based on max age 50	M prior based on max age 45	WCG-BTS selectivity double-normal, asymptotic	WCG-BTS selectivity double-normal, allowed to be dome	Recreational catches added
Diff. in likelihood from base model									
Total	0	23.01	0.85	3.49	-0.48	-0.97	1.65	1.65	0.01
Index	0	2.563	0.167	0.952	0.005	0.016	0.06	0.06	0.002
Length comp	0	-0.588	-0.471	-0.222	-0.004	-0.011	1.781	1.781	-0.004
Age comp	0	11.582	1.405	2.533	0.043	0.109	-0.101	-0.101	0.015
Parm priors	0	-1.255	0.491	1.425	-0.435	-0.884	-0.006	-0.006	-0.001
Estimates of key parameters									
M Female	0.135	0.1	0.144	0.157	0.136	0.137	0.135	0.135	0.135
M Male	0.147	0.1	0.155	0.17	0.148	0.149	0.147	0.147	0.147
Estimates of derived quantities									
Unfished age 4+ bio 1000 mt	156.4	153.1	159.2	168.3	156.7	157.1	155.9	155.9	156.5
B0 trillions of eggs	21.11	21.1	20.83	21.56	21.11	21.11	21.04	21.04	21.12
B2025 trillions of eggs	10.57	4.61	11.47	13.39	10.67	10.81	10.39	10.39	10.57
Fraction unfished 2025	0.501	0.218	0.55	0.621	0.506	0.512	0.494	0.494	0.5
Relative fishing intensity 2024	1.214	1.719	1.122	0.989	1.205	1.193	1.226	1.226	1.214
2027 OFL mt	4916	1307	5809	7400	4995	5108	4831	4831	4914
2027 ACL mt	4596	351	5431	6919	4671	4776	4517	4517	4595
Equil. catch at MSY mt	6746	5073	7249	8244	6789	6851	6720	6720	6748
Equil. catch at SPR targ. mt	5931	4548	6360	7193	5967	6020	5909	5909	5933

5.5 Management

Table 21: Estimates of key derived parameters and reference points with approximate 95% asymptotic confidence intervals.

Reference Point	Estimate	Lower Interval	Upper Interval
Unfished Spawning output (billions of eggs)	21,110.8	17,760.2	24,461.4
Unfished Age 4+ Biomass (mt)	156,376.0	131,787.1	180,964.9
Unfished Recruitment (R0)	40,470.3	28,185.9	52,754.7
2025 Spawning output (billions of eggs)	10,571.7	5,126.8	16,016.6
2025 Fraction Unfished	0.501	0.304	0.698
Reference Points Based SO40%			
Proxy Spawning output (billions of eggs) SO40%	8,444.3	7,104.1	9,784.6
SPR Resulting in SO40%	0.458	0.458	0.458
Exploitation Rate Resulting in SO40%	0.083	0.076	0.091
Yield with SPR Based On SO40% (mt)	6,238.8	4,973.6	7,504.1
Reference Points Based on SPR Proxy for MSY			
Proxy Spawning output (billions of eggs) (SPR50)	9,418.7	7,923.8	10,913.6
SPR50	0.5		
Exploitation Rate Corresponding to SPR50	0.073	0.067	0.079
Yield with SPR50 at SO SPR (mt)	5,930.5	4,732.1	7,128.9
Reference Points Based on Estimated MSY Values			
Spawning output (billions of eggs) at MSY (SO MSY)	5,393.1	4,540.5	6,245.8
SPR MSY	0.328	0.324	0.332
Exploitation Rate Corresponding to SPR MSY	0.127	0.116	0.138
MSY (mt)	6,745.5	5,358.5	8,132.4

Table 22: Potential OFL, ABC, ACL, the buffer between the OFL and ABC, estimated spawning biomass (SB) (mt), and fraction of unfished SB with adopted OFL and ABC and assumed catch for the first two years of the projection period. The predicted OFL is the calculated total catch determined by FSPR=50%.

Year	Adopted OFL (mt)	Adopted ACL (mt)	Assumed Catch (mt)	OFL (mt)	Buffer	ABC (mt)	ACL (mt)	Spawning output (billions of eggs)	Fraction Unfished
2025	12,254	11,237	10,669	-	-	-	-	10,571.70	0.501
2026	11,382	10,392	9,824	-	-	-	-	9,630.18	0.456
2027	-	-	-	4,916	0.935	4,596	4,596	8,799.27	0.417
2028	-	-	-	5,172	0.93	4,810	4,810	8,763.01	0.415
2029	-	-	-	5,548	0.926	5,137	5,137	8,869.58	0.420
2030	-	-	-	5,866	0.922	5,409	5,409	9,054.47	0.429
2031	-	-	-	6,047	0.917	5,545	5,545	9,245.68	0.438
2032	-	-	-	6,117	0.913	5,585	5,585	9,403.42	0.445
2033	-	-	-	6,130	0.909	5,572	5,572	9,520.54	0.451
2034	-	-	-	6,123	0.904	5,535	5,535	9,605.99	0.455
2035	-	-	-	6,116	0.9	5,504	5,504	9,671.68	0.458
2036	-	-	-	6,116	0.896	5,480	5,480	9,725.54	0.461

6 Figures

Note: figure numbering is continuous with executive summary

6.1 Data




Figure 6: Total catches (landings and discards combined) of widow rockfish from 1916 to 2024 for bottom trawl, midwater trawl, and hake fleets.




Figure 7: Data sources by type and year that were used in the base model.

6.1.1 Indices




Figure 8: Fishery-dependent indices of abundance from the 2011 assessment scaled to the mean of their own series.




Figure 9: QQ plots for delta-gamma (panel 1) and delta-lognormal (panel 2, used in base model) sdmTMB index.




Figure 10: Comparison of the estimated index of relative abundance for the West Coast Groundfish Bottom Trawl Survey using a GLMM (gray line, 2015 stock assessment), VAST (black line, 2019 stock assessment), a delta-gamma sdmTMB index (blue line) and a delta-lognormal sdmTMB index (red line, used in base model). The error bars give 5 and 95% intervals for each survey.




Figure 11: Expanded length compositions for the WCGBTs




Figure 12: Conditional age-at-length from WCG BTS observations for females (red) and males (blue).

6.1.2 Composition data




Figure 13: Expanded length compositions for the bottom trawl fishery. The area of the circle is proportional to the proportion-at-length.




Figure 14: Expanded length compositions for the midwater trawl fishery. The area of the circle is proportional to the proportion-at-length.




Figure 15: Expanded length compositions for the hake fishery. The area of the circle is proportional to the proportion-at-length.




Figure 16: Expanded age compositions for the bottom trawl fishery. The area of the circle is proportional to the proportion-at-age.




Figure 17: Expanded age compositions for the midwater trawl fishery. The area of the circles is proportional to the proportion-at-age.




Figure 18: Expanded age compositions for the hake fishery. The area of the circles is proportional to the proportion-at-age.




Figure 19: Comparison between discard data for the hook-and-line fleet from the 2019 assessment, which included nearshore fixed gear fleets, and discards queried from WCGOP for this assessment.

6.1.3 Biological data




Figure 20: Fits to weight-at-length observations (dashed line) for females (left) and males (right) from the previous and current assessments. Fits to the current assessment use observations from all data sources (points)




Figure 21: Number at age observed from all data for female and male widow rockfish.

6.2 Model

6.2.1 Bridging




Figure 22: Spawning biomass with intermediate steps in hook-and-like discard data updates.




Figure 23: Fraction unfished with intermediate steps in hook-and-like discard data updates.




Figure 24: Fraction unfished with intermediate steps in hook-and-like discard data updates.




Figure 25: Time series of spawning biomass (SB) estimates from model runs bridging from the 2019 assessment (dark blue) to the August 2025 base model (red). For illustration purposes, the addition of midwater and bottom trawl discards (without updating hook-and-line discards) precedes the addition of hook-and-line discards to hook-and-line landings. Some bridging steps with minimal impact on SB are grouped (including updating mortality priors, length/weight regression estimates, stock-recruitment bias ramp estimation, and the addition of blocks on midwater and hake retention).




Figure 26: Time series of relative spawning biomass (fraction unfished) estimates from model runs bridging from the 2019 assessment (dark blue) to the August 2025 base model (red).




Figure 27: Time series of relative spawning biomass (fraction unfished) estimates from model runs bridging from the August 2025 model (blue) to the base model (red).




Figure 28: Time series of relative spawning biomass (fraction unfished) estimates from model runs bridging from the August 2025 model (blue) to the base model (red).




Figure 29: Time series of estimated recruitment from model runs bridging from the August 2025 model (blue) to the base model (red).




Figure 30: Two fecundity-at-length relationships (blue) are contrasted with weight-at-length for females (red), along with the mature numbers at length at the start and end of the base model (green). The exponent of the weight-at-length curve is 2.987 compared to 4.043 for the unobserved rockfish fecundity (used in the October 2025 model) and 4.545 for the Widow rockfish-specific fecundity (used in the January 2026 base model). Comparing these relationships to the number of mature fish in each length bin in 1916 and 2025 shows that when the fecundity relationships are used, the spawning output is more dependent on the larger fish (over 45 cm) which are disproportionately removed from the population due to selective fishing on larger sizes. This combination of factors leads to lower estimates of fraction of unfished when one of the fecundity relationships is used.




Figure 31: Time series of fraction of unfished spawning output estimates and asymptotic 95% confidence intervals estimates from 2019 and 2025 base models.




Figure 32: Time series of age 4+ biomass from the 2019 and 2025 base models.




Figure 33: Estimates of age-0 recruits (in millions of individuals) from 2019 and 2025 base models.




Figure 34: Estimates of age-0 (log-)recruitment deviations from 2019 and 2025 base models, with asymptotic 95% confidence intervals.




Figure 35: The prior for natural mortality (grey, M , yr⁻¹) and the estimated M for females (red) and males (blue) from the 2015 base model, 2019 base model, and selected intermediate models bridging to the August 2025 base model. Prior 95% quantiles are based on the assumed lognormal distribution. Confidence intervals for estimated male and female M are shown only for the 2015, 2019, and 2025 base models and are based on a normal asymptotic approximation to the sampling distribution. Depicted bridging models are selected to highlight changes with a notable impact on M .

6.2.2 Selectivity




Figure 36: Estimated selectivity for different fleets and surveys.




Figure 37: Estimated selectivity curves by time block for bottom trawl, midwater trawl, and hake fleets.

6.2.3 Biology




Figure 38: Length at age (top-left panel) with estimated coefficient of variation (CV, thick line) and calculated standard deviation (SD, thin line) versus length at age in the top-right panel and versus age in the lower-left panel.




Figure 39: Estimates of recruitment (upper) and recruitment deviates (lower) with approximate asymptotic 95% confidence intervals (vertical lines) from the MLE estimates.

6.2.4 Fits to data




Figure 40: Estimated and input recruitment bias adjustment ramp. Red line shows current settings for bias adjustment ramp. Red line shows current settings for bias adjustment specified in the model. Blue line shows least squares estimate of alternative bias adjustment relationship for recruitment deviations.




Figure 41: Fits (blue lines) to the abundance estimates (black points) for the base model. A separate q is estimated for the Hake series starting in 1991. Except the juvenile survey (counts) all indices are biomass-based. 95% confidence intervals are shown assuming a normal distribution of the log-estimates. Thicker lines (if present) indicate input uncertainty before addition of the estimated additional uncertainty parameter. The y-axis of the juvenile survey is truncated to highlight interannual variability in counts.




Figure 42: Pearson residuals for fits to length frequency data (left) and age frequency data (right) for landings from the trawl commercial fleets (rows). Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 43: Combined length frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).




Figure 44: Pearson residuals for fits to the triennial survey length frequency data (left) and WCGBTS length frequency data (right). Filled circles indicate that the fitted proportion was less than the observed proportion. Red indicates females, blue males, and gray unsexed.




Figure 45: Combined age frequencies for all years from fishery (retained catch) and survey length frequency data (points). Fits are shown by the red line (females) and blue line (males).




Figure 46: Observed and expected age-at-length with 95% confidence intervals (left) and observed and expected standard deviation of age-at-length with 95% confidence intervals (right) for the WCGBTS.




Figure 47: Combined age and length frequency data for all years from fishery (retained catch) and survey length frequency data (points) for females (red) and males (blue).

6.2.5 Time series




Figure 48: Predicted spawning output (billions of eggs) for widow rockfish using the base model. The solid line is the MLE estimate, and the dashed lines depict the approximate asymptotic 95% confidence intervals.




Figure 49: Predicted summary biomass (age 4+) from the base model.




Figure 50: Predicted relative spawning output from the widow rockfish base model. The solid blue line is the MLE estimate, and the dashed lines depict the approximate asymptotic 95% confidence intervals. The red lines show the equilibrium level (100%), the management target of 40% of unfished biomass, and the minimum stock size threshold of 25% of unfished biomass.




Figure 51: Estimated recruitment and the assumed stock-recruit relationship (black line). The dashed line shows the effect of the bias correction for the lognormal distribution.




Figure 52: Plot of the summary fishing mortality for each year of the model with 95% confidence intervals.




Figure 53: Phase plot of relative fishing intensity $(1-SPR)/(1-SPR_{50\%})$ (y-axis) and fraction unfished spawning output (x-axis) for widow rockfish.

6.3 Model Diagnostics

6.3.1 Sensitivity analyses




Figure 54: Spawning output (with 95% confidence interval around the base model) for the base model and sensitivity runs.




Figure 55: Comparison of fraction unfished for the base model and sensitivity runs.




Figure 56: Estimates of recruitment for base model and sensitivity models.




Figure 57: Estimates of recruitment deviations for base model and sensitivity models.

6.3.2 Retrospective analysis




Figure 58: Ten-year retrospective estimates of spawning output (top) and recruitment deviations (bottom).




Figure 59: Estimated fraction of unfished spawning output from past assessments in comparison with the current assessment. The units of spawning output differ among assessments but the ratio remains comparable.

6.3.3 Likelihood profiles





Figure 60: Likelihood components in the likelihood profile for unfished equilibrium recruitment (R_0)

Figure 61: Likelihood components in the likelihood profile for steepness (h).




Figure 62: Likelihood components in the likelihood profile for female natural mortality (M). Male natural mortality are set to the same value.




Figure 63: Likelihood components in the likelihood profile for male natural mortality (M). Female natural mortality is set to the same value.




Figure 64: Time series of recruitment deviations for models with different fixed values of natural mortality (M)

6.4 Management




Figure 65: Estimated plot of equilibrium yield vs the fraction of unfished biomass.

7 References

- Adams, G., Kapur, M., McQuaw, K., Hamel, O., Stephens, A., and Wetzel, C. 2019. Status of Widow Rockfish (*Sebastes entomelas*) Along the U.S. West Coast in 2019. Pacific Fishery Management Council, Portland, Oregon.
- Anderson, S.C., Ward, E.J., English, P.A., and Barnett, L.A. 2022. sdmTMB: An R package for fast, flexible, and user-friendly generalized linear mixed effects models with spatial and spatiotemporal random fields. *BioRxiv*: 2022–03.
- Barss, W., and Echeverria, T.W. 1987. Maturity of widow rockfish *Sebastes entomelas* from the northeastern Pacific, 1977-82. pp. 13–18.
- Dick, E., Beyer, S., Mangel, M., and Ralston, S. 2017. A meta-analysis of fecundity in rockfishes (genus *sebastes*). *Fisheries Research* **187**: 73–85. Elsevier.
- Douglas, D.A., and others. 1998. Species composition of rockfish in catches by oregon trawlers, 1963-93.
- Fournier, D.A., Skaug, H.J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M.N., Nielsen, A., and Sibert, J. 2012. AD Model Builder: Using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optimization Methods and Software* **27**(2): 233–249.
- Francis, R.C. 2011. Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(6): 1124–1138.
- Gunderson, D.R. 1984. The great widow rockfish hunt of 1980-1982. *North American Journal of Fisheries Management* **4**(4B): 465–468. Taylor & Francis.
- Hamel, O.S. 2015. A method for calculating a meta-analytical prior for the natural mortality rate using multiple life history correlates. *ICES Journal of Marine Science* **72**(1): 62–69.
- Hamel, O.S., and Cope, J.M. 2022. Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research* **256**: 106477.
- He, X., Pearson, D., Dick, E.J., Field, J., Ralston, S., and MacCall, A. 2011. Status of the widow rockfish resource in 2011. Pacific Fishery Management Council, Portland, OR.
- Helser, T.E., Punt, A.E., and Methot, R.D. 2004. A generalized linear mixed model analysis of a multi-vessel fishery resource survey. *Fisheries Research* **70**(2-3): 251–264.
- Hicks, A., and Wetzel, C. 2015. The Status of Widow Rockfish (*Sebastes entomelas*) Along the U.S. West Coast in 2015. Pacific Fishery Management Council, Portland, OR.
- Holland, D.S., and Martin, C. 2019. Bycatch quotas, risk pools, and cooperation in the Pacific whiting fishery. *Frontiers in Marine Science* **6**: 600.
- Jones, W.G., and Harry Jr, G.Y. 1960. The oregon trawl fishery for mink food 1948-1957. *Fish Commission of Oregon Research Briefs* **8**: 14–30.
- Keller, A.A., Wallace, J.R., and Methot, R.D. 2017. The northwest fisheries science center's west coast groundfish bottom trawl survey: History, design, and description. NOAA Technical Memorandum NMFS NOAA-TM-NMFS-NWFSC-136: 38 pp.
- McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling-importance resampling algorithm. *Canadian Journal of Fisheries and Aquatic Sciences* **54**(2): 284–300.
- Methot, R.D., and Taylor, I.G. 2011. Adjusting for bias due to variability of estimated

- recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(10): 1744–1760.
- NMFS. 2017. Magnuson-Stevens Act Provisions; Fisheries Off West Coast States; Pacific Coast Groundfish Fishery; Widow Rockfish Reallocation in the Individual Fishing Quota Fishery.
- Pikitch, E.K., Erickson, D.L., and Wallace, J.R. 1988. An evaluation of the effectiveness of trip limits as a management tool.
- Punt, A.E. 2017. Some insights into data weighting in integrated stock assessments. *Fisheries Research* **192**: 52–65.
- Rogers, J.B. 2003. Species allocation of sebastes and sebastolobus sp. Caught by foreign countries from 1965 through 1976 off washington, oregon, and california, USA.
- Sampson, D.B., and Lee, Y.-W. 2002. Evaluating the accuracy of trawl logbook estimates of groundfish discards. *In* International institute of fisheries economics and trade.
- Stewart, I.J., and Hamel, O.S. 2014. Bootstrapping of sample sizes for length-or age-composition data used in stock assessments. *Canadian journal of fisheries and aquatic sciences* **71**(4): 581–588.
- Taylor, I.G., Doering, K.L., Johnson, K.F., Wetzel, C.R., and Stewart, I.J. 2021. Beyond visualizing catch-at-age models: Lessons learned from the r4ss package about software to support stock assessments. *Fisheries Research* **239**: 105924. Available from <https://doi.org/10.1016/j.fishres.2021.105924>.
- Then, A.Y., Hoenig, J.M., Hall, N.G., Hewitt, D.A., and Handling editor: Ernesto Jardim. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science* **72**(1): 82–92.
- Thorson, J.T., Dorn, M.W., and Hamel, O.S. 2019. Steepness for west coast rockfishes: Results from a twelve-year experiment in iterative regional meta-analysis. *Fisheries Research* **217**: 11–20. Elsevier.
- Thorson, J.T., Shelton, A.O., Ward, E.J., and Skaug, H.J. 2015. Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. *ICES Journal of Marine Science* **72**(5): 1297–1310.
- Thorson, J.T., and Ward, E.J. 2014. Accounting for vessel effects when standardizing catch rates from cooperative surveys. *Fisheries Research* **155**: 168–176.
- Weinberg, J.R., Rago, P.J., Wakefield, W.W., and Keith, C. 2002. Estimation of tow distance and spatial heterogeneity using data from inclinometer sensors: An example using a clam survey dredge. *Fisheries Research* **55**(1-3): 49–61.
- Wetzel, C.R., Johnson, K.F., and Hicks, A.C. 2025. nwfsSurvey: Northwest Fisheries Science Center Survey. Available from <https://github.com/pfmc-assessments/nwfsSurvey>.
- Wilkins, M.E. 1986. Development and evaluation of methodologies for assessing and monitoring the abundance of widow rockfish, sebastes entomelas. *Fishery Bulletin* **84**(2).

Acronyms

ABC Acceptable Biological Catch. [xv](#), [32](#), [72](#)

ACL annual catch limit. [xiv](#), [xv](#), [32](#), [72](#)

Juvenile Survey SWFSC and NWFSC/PWCC Midwater Trawl Survey. [iv](#), [6](#)

NMFS National Marine Fisheries Service. [iv](#), [3](#)

OFL overfishing limit. [xv](#), [xvi](#), [30](#), [32](#), [72](#)

SB spawning biomass. [72](#)

sdmTMB [Species Distribution Models with Template Model Builder](#). [5](#)

SPR spawning potential ratio. [xii](#), [27](#)

Triennial Survey Alaska Fisheries Science Center/Northwest Fisheries Science Center West Coast
Triennial Shelf Survey. [iv](#), [10](#), [18](#), [21](#), [22](#)

WCGBTS Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey.
[iv](#), [xvi](#), [xvii](#), [4](#), [10](#), [13](#), [16](#), [18](#), [21](#), [22](#), [23](#), [25](#), [26](#), [29](#), [30](#)

WCGOP West Coast Groundfish Observer Program. [3](#), [20](#), [28](#)