

Determining the causal effects of lipid levels on risk of dementia: a triangulation of new and existing evidence



Luke A McGuinness

A dissertation submitted to the University of Bristol in accordance with the requirements for award of the degree of

Doctor of Philosophy

in the Faculty of Health Sciences

December 2021

Word count: 48,351

Abstract

Background: Identification of causal relationships between modifiable targets and risk of dementia is central to the development of evidence-based prevention strategies. Blood lipid levels have been implicated in the aetiology of dementia by genetic linkage and functional cell biology studies, but current epidemiological evidence has yet to reach a consensus on their role in dementia risk. This thesis sought to triangulate the causal effects of lipid levels (total cholesterol, LDL-c, HDL-c, and triglycerides) on risk of incident dementia (all-cause dementia, Alzheimer's disease, and vascular dementia) using existing and new evidence.

Methods: Four distinct analyses were conducted. Firstly, a systematic review of 81 published and preprinted studies was used to summarise the existing evidence base. Preprints were searched using a new research tool created as part of this thesis. The existing evidence was then supplemented via two primary studies: i) a cohort study of the association of lipid regulating agents and dementia incidence in the Clinical Practice Research Datalink (CPRD); and ii) an individual participant data meta-analysis of the association of blood lipids with dementia incidence in previously unanalysed cohorts accessed through the Dementia Platform UK. Finally, a novel quantitative triangulation framework was proposed, building on recent developments in risk-of-bias assessment and bias-/indirectness-adjusted meta-analyses. This framework was then used to quantitatively triangulate the evidence identified and produced by this thesis.

Results: The systematic review did not identify a consistent effect of blood lipids on any dementia outcome across study designs, though there was some suggestion of a protective effect of LDL-c lowering on all-cause dementia and Alzheimer's disease in observational studies of statin use. The analysis of lipid regulating agents in the CPRD provided weak evidence for a protective effect of statins on all-cause dementia and Alzheimer's disease but suggested a harmful association with vascular dementia. However, the use of control outcomes illustrated this finding was likely due to confounding by indication related to vascular factors. The

individual participant data meta-analysis suffered from a low response rate to data access requests, but in the three cohorts analysed, there was some evidence only for the association of triglycerides and vascular dementia. Finally, the triangulation analysis integrating the results of the previous three studies did not provide strong evidence for the causal effect of blood lipids on dementia outcomes.

Conclusions: This thesis provides new evidence concerning the role of blood lipids as a modifiable risk factor for dementia and highlights the uncertainty that still remains in relation to this causal question. In addition, it has developed new evidence synthesis methods and tools, specifically around the inclusion of preprints in systematic reviews and the quantitative triangulation of evidence sources.

For Brendan McHugh

Acknowledgements

There are many people without whose input and support this work would have been severely diminished:

First and foremost, I would like to thank my supervisors - Julian Higgins, Yoav Ben-Shlomo, Liz Coulthard and George Davey-Smith - for all their support over the past three years. Thanks also to Rupert Payne and Emma Anderson for their engaging discussion and useful feedback during my annual progress reviews. Sharen O'Keffe helped me at numerous points throughout my PhD, and I was very lucky to receive her support. Similarly, Mark Newbury of the Dementia Platform UK worked tirelessly to support my applications for data access - I am extremely grateful for his help and commitment in chasing cohort owners.

Thanks must also go to those who contributed their time and energy to my research projects: Matthew Lee, Lena Schmidt, Alexandra McAleenan, and Vincent Cheng. Francesca Spiga kindly shared an advance copy of her review of tools to assess Mendelian randomisation studies, and Rebecca Turner shared her code for bias-/indirectness-adjusted meta-analyses that underpins the quantitative triangulation framework presented in Chapter 7. Thanks to Vivienne for proof-reading.

A particular thank you to Emily Hennessy, with whom I co-wrote the guidance paper for preparing primary research for incorporation into systematic reviews. Similarly, thanks to Athena Sheppard for her support in assessing the impact of journal policies on open-data sharing.

Thanks also to all those who raised issues or otherwise provided feedback on the open source tools built as part of this thesis: @abannachbrown, @ailengt, @AJFOWLER, @danielskatz, @earcanal, @Jamesohare1, @jannisborn, @paulsharpeY, @rachaelm-burke, @rdboyes, @seabbs, @vanAmsterdam, @yochannah, and @zaddyzad.. A special mention must go to Randall Boyes and Alex J Fowler for their input on the `robvis` tool.

I have been supported throughout my studies by an NIHR Doctoral Research Fellowship (DRF-2018-11-ST2-048). The views expressed in this thesis are my own and do not necessarily represent those of the NIHR or the Department of Health and Social Care.

On a personal note, there are several people without whom I would not have survived the process of writing a dissertation during COVID times. Thanks to Mum and Dad for your never-ending love and support (I promise I'll get a real job now!). Thank also to Doug for introducing me to the joys of running, Ben for always being around to chat, and Christin for encouraging regular coffee-breaks.

Finally, deepest thanks to Ciara Gardiner: holder-of-forts, maker-of-spreadsheets, and all-round fiancee-extraordinaire. I can't wait for the next adventure with you.

Canynge Hall, Bristol

1 December 2021

Author's declaration

I declare that the work in this dissertation was carried out in accordance with the requirements of the University's Regulations and Code of Practice for Research Degree Programmes and that it has not been submitted for any other academic award. Except where indicated by specific reference in the text, the work is the candidate's own work. Work done in collaboration with, or with the assistance of, others, is indicated as such. Any views expressed in the dissertation are those of the author.

SIGNED: DATE:

Contents

List of Figures	xiii
List of Tables	xv
List of Abbreviations	xvii
1 Background, Theoretical framework, Aims & Objectives	1
Lay summary	2
1.1 Introduction	2
1.2 Dementia	3
1.2.1 Definition and underlying pathologies	3
1.2.2 Diagnostic criteria	4
1.2.3 Public health importance	6
1.2.4 Treatments	6
1.2.5 Risk factors	7
1.3 Serum lipids	8
1.3.1 Lipid fractions	8
1.3.2 Statins	9
1.3.3 Other lipid regulating agents (LRA)	11
1.4 Evidence for the association between blood lipids and dementia . .	13
1.4.1 Basic science	13
1.4.2 Observational studies	13
1.4.3 Randomised controlled trials	14
1.4.4 Mendelian randomisation	15

Contents

1.5	Theoretical framework: Evidence synthesis	16
1.5.1	Inclusion of preprints	17
1.5.2	Individual participant data meta-analysis	18
1.5.3	Triangulating across study designs	19
1.6	Thesis overview	20
1.6.1	Aims and objectives	20
1.6.2	Structure	21
1.7	Outputs from this thesis	22
1.7.1	Contributions to the scientific literature	22
1.7.2	Presentations/Talks	24
1.7.3	Software	25
1.8	Summary	26
2	medrxivr: an R package for systematically searching biomedical preprints	27
	Lay summary	28
2.1	Introduction	28
2.2	Development	30
2.2.1	Success criteria	30
2.2.2	Alternative medRxiv/bioRxiv interfaces	30
2.2.3	Early versions	31
2.2.4	Package infrastructure	32
2.3	Usage	32
2.3.1	Installation	33
2.3.2	Importing preprint metadata	33
2.3.3	Performing a search	35
2.3.4	Refining a search	36
2.3.5	Exporting to a bibliography file	38
2.3.6	Downloading the PDFs of relevant records	38
2.4	Discussion	39

2.4.1	Reception and future plans	39
2.4.2	Use cases	39
2.4.3	Limitations of <code>medrxivr</code>	40
2.4.4	Role of open source tools in evidence synthesis	41
2.5	Summary	42
3	Systematic review of existing evidence on the association between blood lipids and dementia outcomes: Methods	43
	Lay summary	43
3.1	Introduction	44
3.2	Methods	45
3.2.1	Protocol	45
3.2.2	Contributions	45
3.2.3	Eligibility criteria	45
3.2.4	Information sources and search strategy	47
3.2.5	Study selection	49
3.2.6	Validation of screening process	50
3.2.7	Data extraction	51
3.2.8	Risk-of-bias assessment	53
3.2.9	Analysis methods	57
3.3	Summary	62
4	Systematic review of existing evidence on the association between blood lipids and dementia outcomes: Results	63
	Lay summary	63
4.1	Introduction	64
4.2	Search results	64
4.2.1	Initial search and validation of search filters	64
4.2.2	Screening results	64
4.2.3	Validation of screening	67
4.2.4	Characteristics of included studies	68

4.3 Primary analyses	77
4.3.1 All-cause dementia	77
4.3.2 Alzheimer's disease	82
4.3.3 Vascular dementia	87
4.3.4 Risk of bias	89
4.4 Additional analyses	90
4.4.1 Dose response meta-analysis of lipid levels	90
4.4.2 Heterogeneity and publication bias	91
4.4.3 Added evidential value of including preprints	92
4.5 Discussion	93
4.5.1 Summary of findings	93
4.5.2 Comparison with previous reviews	95
4.5.3 Inclusion of Mendelian randomisation studies	96
4.5.4 Inclusion of preprints	99
4.5.5 Open data sharing	102
4.5.6 Strengths and limitations	102
4.6 Summary	104
5 Primary analysis of lipid-regulating agents and dementia outcomes	107
Lay summary	108
5.1 Introduction	108
5.2 Methods	109
5.2.1 Study protocol	109
5.2.2 Data source	109
5.2.3 Cohort definition	111
5.2.4 Exposures	112
5.2.5 Outcomes	113
5.2.6 Covariates	113
5.2.7 Missing data	116
5.2.8 Estimation methods	116

5.2.9	Immortal time bias and time-varying treatment indicators	117
5.2.10	Time axis	119
5.2.11	Sensitivity analyses	119
5.3	Results	123
5.3.1	Patient characteristics	123
5.3.2	Missing data	128
5.3.3	Primary analysis	129
5.3.4	Sensitivity analyses	131
5.4	Discussion	139
5.4.1	Summary of findings	139
5.4.2	Interpretation of results	139
5.4.3	Strengths and limitations	144
5.4.4	Enabling easy synthesis of this analysis	146
5.4.5	Summary	146
6	Individual participant data meta-analysis of blood lipid levels and dementia outcomes	149
	Lay Summary	150
6.1	Introduction	150
6.2	Methods	151
6.2.1	Eligibility criteria	151
6.2.2	Applying for data access	152
6.2.3	Primary analysis	153
6.3	Results	156
6.3.1	Data access	156
6.3.2	Included data sources	159
6.3.3	Covariate definition & missing data	161
6.3.4	Analytical results	161
6.4	Discussion	168
6.4.1	Summary of findings	168

Contents

6.4.2	Limitations	169
6.4.3	Strengths	173
6.4.4	Reflections on the process	173
6.4.5	Future work	174
6.5	Summary	175
7	Aetiological triangulation across new and existing evidence sources	177
	Lay summary	178
7.1	Introduction	178
7.2	Data sources	179
7.3	Qualitative (narrative) triangulation	182
7.4	Quantitative triangulation	185
7.4.1	Methods	185
7.4.2	Results	195
7.5	Discussion	201
7.5.1	Summary of findings	201
7.5.2	Limitations	202
7.5.3	Strengths	205
7.5.4	Future research	206
7.5.5	Outputs	207
7.6	Summary	208
8	Discussion	209
8.1	Lay summary	210
8.2	Introduction	210
8.2.1	Chapter summary	210
8.3	Summary of clinical findings	211
8.3.1	Effect of blood lipids on dementia outcomes	211
8.3.2	Comparison with new evidence	212
8.3.3	Implications for clinical practice	212

8.3.4	Implications for public health	213
8.4	Summary of methodological contributions	214
8.4.1	Inclusion of preprinted evidence	214
8.4.2	Bias	214
8.4.3	Triangulation framework	215
8.5	Overall strengths and limitations	216
8.5.1	Strengths	216
8.5.2	Weaknesses	217
8.6	Lessons learned	218
8.7	Future work	218
8.7.1	Generation of new RCT evidence	219
8.7.2	Evidence on vascular dementia	219
8.7.3	Preprinted evidence	220
8.7.4	Reviewing Mendelian randomisation studies	221
8.7.5	Systematic reviews and quantitative triangulation	221
8.8	Overall conclusions	222
9	Bibliography	223
Appendices		
A	Appendices by Chapter	253
A.1	Chapter 1	254
A.1.1	Publications beyond the scope of this thesis	254
A.2	Chapter 2	256
A.2.1	Code for publication rate analysis	256
A.3	Chapter 3	258
A.3.1	Deviations from protocol	258
A.3.2	Web of Science databases searched	259
A.3.3	Search strategy	259
A.3.4	Code to search preprints	271

A.3.5 Calculating Gwet's AC1	273
A.3.6 Risk-of-bias assessment tool for Mendelian randomisation studies	274
A.4 Chapter 4	277
A.4.1 Dose response plots by fraction	277
A.5 Chapter 5	278
A.5.1 Amendments to protocol	278
A.5.2 Minimum follow-up period	279
A.6 Chapter 6	281
A.6.1 Covariate matrix	281
A.6.2 Dementia definitions across cohorts	281
A.6.3 Comparison with previous analysis of the Whitehall II cohort	283
A.7 Chapter 7	283
A.7.1 Calculation of average adjustment values from previous expert elicitation exercise	283
A.7.2 Comparison of two scenarios of bias	286
A.8 Chapter 8	288
A.8.1 Reproducible research	288
A.8.2 Catalogue of failures	288
B Other Appendices	291
B.1 Software used to create this thesis	292
B.2 Producing risk-of-bias visualisations with <code>robvis</code>	295
B.2.1 Simple visualisations	297
B.3 Copies of papers arising from this thesis	298

List of Figures

1.1	Statin mechanism of action	10
2.1	Role of <code>medrxivr</code> in a systematic review workflow	29
2.2	Overview of <code>medrxivr</code> data sources	35
4.1	PRISMA flow diagram	66
4.2	Geographical distribution of study cohorts	76
4.3	Meta-analysis of hypercholesterolemia on all-cause dementia	78
4.4	Random-effects meta-analysis of four lipid fractions on all-cause dementia	79
4.5	Random-effects meta-analysis of statins on all-cause dementia	80
4.6	Random-effects meta-analysis of statins on all-cause dementia	81
4.7	Random-effects meta-analysis of statins on all-cause dementia	82
4.8	Meta-analysis of hypercholesterolemia on Alzheimer's disease	83
4.9	Random-effects meta-analysis of four lipid fractions on Alzheimer's disease	84
4.10	Summary of duplication across two sample Mendelian randomisation studies	85
4.11	Random-effects meta-analysis of statins on Alzheimer's disease	86
4.12	Random-effects meta-analysis of genetically lowered LDL-c via HMGCR inhibition on Alzheimer's disease	87
4.13	Meta-analysis of hypercholesterolemia on vascular dementia	87
4.14	Random-effects meta-analysis of four lipid fractions on Alzheimer's disease	88
4.15	(ref:obsStatinDVaD-scrap)	89
4.16	Dose-response meta-analysis of total cholesterol	91

List of Figures

4.17	Study designs by year of publication	94
4.18	Overview of assumptions in Mendelian randomisation analyses	97
4.19	Growth of preprint repositories over time	100
5.1	Decision tree for assigning dementia subtypes	113
5.2	shortcap	118
5.3	Causal diagram for ideal negative outcome	121
5.4	Attrition of CPRD participants	124
5.5	Results from primary analyses of CPRD data	130
5.6	Complete case vs. imputed data analysis	132
5.7	Results of control outcome analysis	133
5.8	Comparison of different combinations of covariates	134
5.9	Sensitivity analysis: grouped year of entry	135
5.10	Comparison of pregnancy analysis	137
5.11	Analysis stratified by statin properties	138
5.12	Confounding by indication causal diagram	141
5.13	Comparison of code used in this analysis with those used in the Smeeth et al. 2010 paper	144
6.1	Flowchart	157
6.2	IPD meta-analysis of all-cause dementia, stratified by lipid fraction	163
6.3	IPD meta-analysis of all-cause dementia, stratified by lipid fraction	164
6.4	Comparison of partially and maximally adjusted results	165
6.5	shortcap	166
6.6	shortcap	167
6.7	shortcap	168
6.8	Comparison of results across the two analysis of the Whitehall II cohort	172
7.1	Illustration of calculating the absolute direction of biases, based on bias type.	190
7.2	Bias direction plot for a single study	196

List of Figures

7.3 Bias direction plot, summarising internal biases	198
7.4 Random effects meta-analysis using unadjusted and bias-/indirectness-adjusted results.	199
7.5 shortcap	200
7.6 shortcap	201
A.1 shortcap	277
A.2 shortcap	278
A.3 shortcap	279
A.4 shortcap	280
A.5 Comparison of results across the two analysis of the Whitehall II cohort	283
A.6 Comparison of bias-/indirectness-adjusted meta-analyses under two different scenarios of prior distributions of bias. Scenario 1: position value between the distributions assigned to “High” and “Moderate” bias levels is the same as between the “Moderate” and “Low” levels. Scenario 2: position value between the distributions assigned to “High” and “Moderate” bias levels is twice that between “Moderate” and “Low” levels	287
A.7 Comparison of bias-/indirectness-adjusted meta-analyses under two different scenarios of prior distributions of bias.	287
B.1 shortcap	297
B.2 shortcap	297

List of Tables

1.1	Overview of the DSM-5 criteria for dementia and vascular dementia.	5
1.2	Classification of blood lipid levels	9
1.3	Overview of common statins	11
1.4	Summary of available treatments for hyperlipidaemia.	12
3.1	searchOverview	48
3.2	Ranges for Gwet's AC1	51
4.1	Inter-rater agreement	67
4.2	Inter-rater agreement	67
4.3	Characteristics of included studies	69
5.1	Example of CPRD Read code hierarchy	110
5.2	Covariates adjusted for.	114
5.3	Crude rates, stratified by outcome and drug class of interest.	126
5.4	Patient characteristics by drug class	127
5.5	Participants who stopped, switched or added treatments by initial treatment type	128
5.6	Frequency of diagnoses by grouped year of cohort entry	136
5.7	Summary of statin properties (lipophilicity vs hydrophilicity).	137
6.1	Exclusion reasons for cohorts providing data	159
6.2	Summary of characteristics of IPD cohorts	162
7.1	Summary of research designs included in this thesis	181
7.2	Mapping of risk-of-bias judgements across assessment tools	188

List of Tables

7.3	Prior distributions mapped to different extents of bias.	192
7.4	Prior distributions mapped to different extents of bias.	193
7.5	singleIndirect	197
A.1	Publication status of medRxiv preprints	258
A.2	Summary of Web of Science databases searched.	259
A.3	searchOverviewAppendix	260
A.4	Medline search strategy	260
A.5	Embase search strategy	263
A.6	PyscINFO search strategy	265
A.7	CENTRAL search strategy	267
A.8	Web of Science search strategy	269
A.9	Mendelian randomisation risk-of-bias assessment tool	274
A.10	missingMatrix	281
A.11	Definition of dementia outcomes in IPD meta-analysis	282
A.12	addTable1	284
A.13	addTable1	285
A.14	addTable1	285
A.15	turnerEstimates	286
B.1	biasData	296
B.2	biasDirectionData	296

List of Abbreviations

AC	Agreement coefficient
ACE	Acetylcholinesterase
AD	Alzheimer's disease
ADGC	Alzheimer's Disease Genetics Consortium
API	Application programming interface
ApoE	Apolipoprotein E
BAS	Bile acid sequestrants
BMI	Body mass index
CAD	Coronary arterial disease
CaPS	Caerphilly Prospective Study
CBS	Coronary bypass surgery
CCI	Charlson co-morbidity index
CIND	Cognitive impairment not dementia
CKD	Chronic kidney disease
CPRD	Clinical Practice Research Datalink
CRAN	Comprehensive R Archive Network
CSV	Comma separated value
CVD	Cardiovascular disease
DOI	Digital object identifier
DPUK	Dementia Platform UK
DSM	Diagnostic and Statistical Manual of Mental Disorders
EHR	Electronic health record
EPIC	European Prospective Investigation of Cancer
FAGs	Fatty acid groups
GLGC	Global Lipids Genetic Consortium

List of Abbreviations

GWAS	Genome wide association studies
HDL	High density lipoprotein
HES	Hospital episode statistics
HMG-CoA	3-hydroxy-3-methylglutaryl-coenzyme-A
HMGCR	HMG-CoA reductase
HR	Hazard ratio
ICD	International Classification of Diseases
IGAP	International Genomics of Alzheimer's Project
IHD	Ischemic heart disease
IMD	Index of multiple deprivation
IPD	Individual participant data
IV	Instrumental variable
JUPITER	Justification for the Use of Statin in Prevention: An Intervention Trial Evaluating Rosuvastatin
LDL-c	Low density lipoprotein cholesterol
LRA	Lipid regulating agent
MCI	Mild cognitive impairment
MICE	Multiple imputation by chained equations
MMSE	Mini Mental State Exam
MoCA	Montreal Cognitive Assessment
MR	Mendelian randomization
NAG	Nicotinic acid groups
NINDS-ADRA	National Institute of Neurological Disorders and Stroke - Alzheimer's Disease and Related Disorders Association
NINDS-AIREN	National Institute of Neurological Disorders and Stroke - Association Internationale pour la Recherche en l'Enseignement en Neurosciences
NOS	Newcastle-Ottowa Scale
NRSE	Non-randomised studies of exposures
NRSI	Non-randomised studies of interventions
OR	Odds ratio

List of Abbreviations

PAD	Peripheral arterial disease
PCSK9i	PCSK9 inhibitors
PDF	Portable document format
PR	Proportional hazards
PROSPER	Prospective Study of Pravastatin in the Elderly
PYAR	Participant-years-at-risk
RCT	Randomised controlled trial
REML	Restricted maximum likelihood
RoB-ME	Risk of bias due to missing evidence
RR	Risk/rate ratio
SD	Standard deviation
SNP	Single nucleotide polymorphism
TC	Total cholesterol
TG	Triglycerides
VaD	Vascular dementia

*Alles in allem genommen haben wir hier offenbar
einen eigenartigen Krankheitsprozeß vor uns.*

~

*Considering everything, it seems we are dealing here
with a special illness.*

— Alois Alzheimer, 1907^{1,2}

1

Background, Theoretical framework, Aims & Objectives

Contents

Lay summary	2
1.1 Introduction	2
1.2 Dementia	3
1.3 Serum lipids	8
1.4 Evidence for the association between blood lipids and dementia	13
1.5 Theoretical framework: Evidence synthesis	16
1.6 Thesis overview	20
1.7 Outputs from this thesis	22
1.8 Summary	26

Lay summary

Around 850,000 people in the UK live with dementia, and by 2040, nearly twice as many will have the condition. Despite many promising candidates, no cure for dementia currently exists, meaning the focus is on finding ways to prevent the condition. The best way to do this is to find risk factors (characteristics that influence a person's chance of developing a disease) for dementia that we can easily change. Avoiding a risk factor does not guarantee that a person will not develop dementia but makes it less likely. A key risk factor for dementia may be the levels of lipids (fatty substances such as cholesterol) in a person's blood, though not all existing research agrees. The aim of this thesis is to use all available evidence to assess whether blood lipid levels are in fact a risk factor for dementia.

This introductory chapter provides background information on both dementia and blood lipids, and on the potential link between them. It introduces the theory used to frame the research presented here, and then maps the formal aims and objectives of the research project to the relevant chapters of this thesis. Finally, it summarises the outputs (journal articles, presentations and software) that were created as part of this thesis.

1.1 Introduction

This chapter provides an overview of the broad context of this thesis, introducing the core concepts used throughout and providing some background on each. It briefly discusses the underlying pathologies and diagnosis of dementia, its public health importance, and the current state of treatment and prevention research. It then provides background on blood lipids and lipid-modifying treatments, and summarises the types of evidence used to examine the effect of these exposures on dementia outcomes.

The chapter introduces evidence synthesis as the key framework used to guide the research presented in the remaining chapters. Finally, it outlines the aims, objectives and structure of this thesis, and briefly summarises the contributions to the scientific literature that arose from this research.

1.2 Dementia

1.2.1 Definition and underlying pathologies

Defined by the Diagnostic and Statistical Manual of Mental Disorders as a “major neurocognitive disorder”, dementia is a progressive disease which impairs cognitive functions including speech, memory and executive reasoning.³ At advanced stage, the condition causes severe behavioral and personality changes,⁴ culminating in reduced motor control that affects patients’ ability to swallow or breathe.⁵ The condition has several distinct underlying causes, including Alzheimer’s disease and vascular dementia.⁶

Alzheimer’s disease is the most common cause of dementia, accounting for approximately 60-80% of cases. Characterised by substantial cognitive impairment and difficulty with high level executive function, it is an insidious disease, with initial onset thought to occur up to 15 years prior to symptomatic presentation.⁷ Much remains unknown about Alzheimer’s pathogenesis, despite research implicating the “amyloid hypothesis”,⁷ as a potential mechanism of disease. Under this hypothesis, the build-up of amyloid plaques (composed mainly of amyloid- β peptide) and neurofibrillary tangles (composed mainly of tau protein) triggers a range of physiological changes, including inflammation and cell death, that result in cognitive impairment.⁷

Vascular dementia (VaD) is the second largest underlying pathology of dementia, accounting for ~10-25% of cases. Vascular dementia is caused by a range of cerebrovascular disorders, and as a result, presentation of symptoms can vary widely.⁸ Similarly, due to the varied underlying pathophysiology, vascular dementia can onset either quite rapidly following a cerebrovascular event such as a stroke or over a long time-frame due to a series of small infarcts.⁹ Vascular dementia regularly co-occurs in patients with Alzheimer’s disease.⁸ This presentation is described as “mixed” dementia,¹⁰ and occurs in approximately 25% of cases.⁶

The remaining 10-30% of cases are caused by other dementia subtypes (e.g. Lewy body or frontotemporal dementias) or other neurological diseases (e.g. Parkinson's disease).⁶

1.2.2 Diagnostic criteria

Dementia is difficult to diagnose, primarily due to its slow onset, in addition to the confusion of initial symptoms with normal ageing.⁷ Dementia is diagnosed on the basis of behavioral and cognitive changes as assessed by an experienced clinician, using one of several diagnostic criteria.

One of the most commonly used sets of criteria for diagnosing dementia cases are the Diagnostic and Statistical Manual of Mental Disorders (DSM) criteria (Table 1.1).³ Outlined in Table 1.1, these form the broad definition of a dementia diagnoses, and are supported by a detailed patient history, evidence from carers and family members, and objective assessments of cognitive ability using neurocognitive tests.

Many cognitive assessment tools exist for the purpose of informing a diagnoses of dementia,¹¹ with two of the best known of these being the Mini Mental State Exam (MMSE) and Montreal Cognitive Assessment (MoCA) scale. The distinction between these memory scales and diagnostic criteria presented above should be noted. For example, the MMSE is used to provide evidence for part A of the criteria presented in 1.1. Taken alone, it does not indicate the absence or presence of dementia, instead merely indicating cognitive impairment which could be due to another cause (for example, temporary delirium as a result of an infection or surgery).

Table 1.1: Overview of the DSM-5 criteria for dementia and vascular dementia.³

Criterion	Major neurocognitive event (previously dementia)
A	Evidence of significant cognitive decline from a previous level of performance in one or more cognitive domains: [*] - Learning and memory - Language - Executive function - Complex attention - Perceptual-motor - Social cognition
B	The cognitive deficits interfere with independence in everyday activities. At a minimum, assistance should be required with complex instrumental activities of daily living, such as paying bills or managing medications.
C	The cognitive deficits do not occur exclusively in the context of a delirium.
D	The cognitive deficits are not better explained by another mental disorder (eg, major depressive disorder, schizophrenia).

* From DSM: Evidence of decline is based on concern of the individual, a knowledgeable informant, or the clinician that there has been a significant decline in cognitive function and a substantial impairment in cognitive performance, preferably documented by standardized neuropsychological testing or, in its absence, another quantified clinical assessment.

Differentiating between the underlying causes of a dementia diagnosis is challenging but necessary, as whether the patient has Alzheimer's disease or vascular dementia will affect expected progression and potential treatment options available (see Section 1.2.4). Cause-specific criteria exist for the diagnosis of dementia subtypes. For example, the NINCDS-ADRDA criteria are commonly used to assess patients for Alzheimer's disease,¹² while vascular dementia is diagnosed using the NINCDS-AIREN criteria.¹³

1.2.3 Public health importance

Dementia is quickly becoming a critically important public health issue. Despite the age-specific incidence and prevalence of dementia remaining relatively constant over time,¹⁴ an ageing population is set to create a dementia epidemic, particularly in Westernised countries.¹⁵ While approximately 525,000 patients have received a dementia diagnosis, the true number of people currently living with dementia in the UK is thought to be closer to 850,000, with this figure expected to double by 2040.¹⁶ Globally, the prevalence of dementia is expected to reach 75 million by 2030.¹⁴ Dementia is a leading cause of death in the UK,¹⁷ and one of the few without a proven treatment.

Dementia also has a substantial economic impact. In 2015, the estimated total cost of dementia in England was £24.2 billion. Health care costs alone were £3.8 billion, with the remainder being divided between unpaid care and social care costs.¹⁸ Globally, the cost of dementia care is expected to rise to \$1tr by 2030.¹⁹ As such, the urgent need to reduce the burden of dementia, both at the personal and system level, is clear.

1.2.4 Treatments

Developing treatments for dementia is regularly deemed to be one of the most challenging markets in the pharmaceutical world, with trials of seemingly promising therapeutics being regularly abandoned due to futility.²⁰ At present, there are no known curative treatments for dementia, regardless of the underlying cause, though several available therapeutics can help alleviate the symptoms of Alzheimer's disease. The most common of these are acetylcholinesterase (ACE) inhibitors, which inhibit the degradation of the neurotransmitter acetylcholine by competitively binding the ACE enzyme. Acetylcholine plays a key role in controlling the cholinergic synapses, which are highly concentrated in regions of the brain (such as the neocortex) that

control higher level brain functions such as memory and attention.²¹ Commonly prescribed ACE inhibitors include donepezil and galantamine.²² ACE inhibitors increase the availability of the neurotransmitter, and have shown clinical effect in easing the behavioural and memory-related symptoms of Alzheimer's disease.²³ ACE inhibitors represent only a stop-gap treatment, treating the symptoms rather than the underlying pathology which may continue to progress.²⁴

1.2.5 Risk factors

Given the substantial burden that dementia represents and the absence of any curative therapies, as detailed in the above sections, the assessment of easily modifiable targets for their utility in the prevention of dementia should be prioritized.²⁵ To date, a substantial amount of research has been produced examining putative risk factors for dementia.²⁶⁻²⁸

The benefits of a prevention-based approach based on addressing these risk factors are well-studied. Reducing the prevalence of the seven most important risk factors for dementia (obesity, hypertension,²⁹ diabetes, smoking, physical inactivity, and low educational attainment) by 10-20% per decade is estimated to result in a reduction in dementia prevalence of 8-15% by 2050.³⁰

In this context, lipid levels represent a promising target for preventative treatment, due to the ready availability of lipid-modifying treatments which could be repurposed.³¹ Determining whether variations in lipid levels are causative for dementia may prove critical in reducing the future burden of the condition.

This thesis will focus on blood lipids as the primary risk factor of interest. The next section provides an overview of blood lipid fractions and therapeutic interventions that modify them, while Section 1.4 provides an overview of the existing evidence for an association between lipids and dementia outcomes.

1.3 Serum lipids

1.3.1 Lipid fractions

The blood lipid profile contains a range of component parts, or fractions. However, this thesis will only consider the two most important fractions, triglycerides (TG) and cholesterol, which are either absorbed from food (exogenous lipids) or produced internally (endogenous lipids).²⁶

Triglycerides are the simplest and most common type of lipids found across the body. They are used to store unused calories from food, and to move energy around the body.³² In contrast, cholesterol is primarily used to create cell walls and certain sex hormones.³³ As lipids are not water soluble, within the blood stream, cholesterol is transported in lipoprotein structures of varying densities. Low-density-lipoprotein cholesterol (LDL-c), commonly known as the “bad” cholesterol, transports fat to cells, acting as an energy conveyor. In contrast, High density-lipoprotein cholesterol (HDL-c), transports cholesterol to the liver to be broken down and excreted.²⁶

In addition to the individual fractions, total serum cholesterol (TC) is a commonly-used summary measure to estimate the total amount of lipid present in the blood. The level of TC is derived from measurements of the individual HDL-c, LDL-c and TG levels using the Friedwald formula:³⁴

$$TC \approx LDLc + HDLc + kTG \quad (1.1)$$

where k is 0.20 if measurements are in milligrams per decilitre (mg/dl) and 0.45 if measured in millimole per litre ($mmol/l$). Widely-used ranges for the acceptable levels of different types of lipids are based on the National Cholesterol Education Program (NCEP)³⁵, and are outlined in Table 1.2.

Elevated LDL-c in the bloodstream, a condition also known as hypercholesterolaemia or hyperlipidaemia,³⁶ can lead to atherosclerosis,³⁷ the build-up of fatty deposits in the blood vessels. These deposits constrict blood flow and can lead to vascular

complications. Alternatively, part of the deposit can detach from the artery walls, forming a clot that can lead to a heart attack or stroke.³⁷

Table 1.2: Classification of blood lipid levels according to the National Cholesterol Education Program guidelines.³⁵

Fraction	Measure (mg/dL)	Classification
LDL cholesterol	<100	Optimal
	100-129	Near/above optimal
	130-159	Borderline high
	160-189	High
	>190	Very high
HDL cholesterol	<40	Low
	>60	High
Triglycerides	<150	Normal
	150-199	Borderline high
	200-499	High
Total cholesterol	>500	Very high
	<200	Desirable
	200-239	Borderline high
	>240	High

1.3.2 Statins

Statins are a commonly prescribed method of lipid regulation.³⁸ Statins inhibit the conversion of 3-hydroxy-3-methylglutaryl-coenzyme-A (HMG-CoA) into mevalonate, by competitively binding with HMG-CoA reductase (HMGCR). This conversion is a key rate-limiting step in the cholesterol biosynthesis pathway (see Figure 1.1), enabling statins to reduce effectively the production of LDL cholesterol.




Figure 1.1: Overview of statins mechanism of action, inhibiting HMG-CoA reductase which controls the conversion of HMG-CoA to mevalonate, the rate-limiting step in cholesterol biosynthesis.

Several statin treatments have been widely available for some time (see Table 1.3). Depending on the statin and dosage prescribed, the average reduction in LDL-c concentrations ranges from 15% with low-intensity regimen (e.g. ravastatin 5 mg/day) up to 60% with a high-intensity regimen (e.g. rosuvastatin 80 mg/day).^{38,39} Statins also vary with regard to their lipophilicity (the extent to which they are lipid soluble), affecting their localisation within the body, with hydrophilic statins being concentrated in the liver and lipophilic statins circulating more widely.⁴⁰ This may create a divide in the pleiotropic affects of statins with differing lipophilicity, particularly given the ability of lipophilic statins to permeate the blood brain barrier.⁴¹

Table 1.3: Overview of commonly-prescribed statins, summarising their approval date (US), properties and lipid-lowering effect.

Name	Brand name	Year approved	Properties	Lipid-lowering effect
Atorvastatin	Lipitor	1996	Lipophilic	+++
Pravastatin	Lipostat	1989	Hydrophilic	+
Rosuvastatin	Crestor	2003	Hydrophilic	++++
Simvastatin	Zocor	1992	Lipophilic	++

1.3.3 Other lipid regulating agents (LRA)

There are several other interventions that can be used to modify a persons lipid profile, each of which act via different mechanisms. However, in general, these treatments are either used as adjunct (additional) treatments to statin therapy or are used in situations where statins are contra-indicated or not tolerated.

The most commonly used non-statin therapeutic is ezetimibe,⁴² which prevents intestinal absorption of cholesterol. However, when used alone, it has a limited LDL-c lowering effect, leading to the creation of combined statin/ezetimibe therapies (both compounds contained in a single pill, as opposed to complimentary treatments).⁴³

Fibrates provide a second example of non-statin therapy. They are used to treat hypertriglyceridaemia by reducing production of triglyceride carrying compounds in the liver. They are commonly used in patients with mixed hyperlipidaemia if treatment with statins has failed to sufficiently control cholesterol levels.

Finally, PCSK9 inhibitors (or PCSK9i) are a relatively new treatment with strong lipid lowering effects, lauded as a potential alternative to statins.⁴⁴ Their mechanism of action is to bind to and inhibit PCSK9, which breaks down LDL-c receptors on the surface of the liver, thus allowing more LDL-c to be internalised and broken down.

1.3 - Serum lipids

Other therapies targeting triglycerides exist, including nicotinic acids⁴⁵ and omega-3-fatty acids,⁴⁶ but they far less effective in LDL-c lowering than the therapies described above.

Table 1.4: Summary of available treatments for hyperlipidaemia.

Treatment	Effect	Mechanism of action	Examples
HMG CoA reductase inhibitors (statins)	Lowers LDL-c & TG Raises HDL-c	Inhibits cholesterol biosynthesis pathway in the liver	Atorvastatin, Simvastatin, Pravastatin
Ezetimibe	Lowers LDL-c	Prevents absorption of cholesterol from diet	
Bile acid sequestrants	Lowers LDL-c	Prevent bile acid reabsorption in the gastro-intestinal tract, increasing conversion of cholesterol to bile acids	Colestipol
Proprotein convertase subtilisin kexin 9 (PCSK9) inhibitors	Lowers LDL-c	Bind to PCSK9 protein, preventing it from breaking down LDL receptors on hepatic cells, increasing cholesterol uptake	Evolocumab, Alirocumab

1.4 Evidence for the association between blood lipids and dementia

This section provides an overview of the varying sources of evidence on the relationship between blood lipid levels and dementia risk.

1.4.1 Basic science

A role for lipids in the aetiology of dementia is supported by both genetic linkage studies and functional cell biology studies. The generation of the amyloid plaques found in the brains of Alzheimer's patients is cholesterol dependent,^{47,48} while the most established genetic risk factor for late-onset dementia, apolipoprotein E (*ApoE*), is involved in cerebral cholesterol transport. Several other genes involved in cholesterol transport have also been found to be associated with increased AD susceptibility.⁴⁹⁻⁵¹

Despite these results, evidence from the diverse range of epidemiological studies on this topic has been inconclusive.

1.4.2 Observational studies

By far the largest source of evidence on the relationship between blood lipids and dementia outcomes comes from observational designs. Several studies have examined the relationships between concentrations of serum lipids (total cholesterol (TC), low density lipoprotein cholesterol (LDL-c), high density lipoprotein cholesterol (HDL-c) and triglycerides) and both Alzheimer's disease and vascular dementia and reported extremely varied results. In some studies, a high serum cholesterol concentration has been found to be associated with an increase in susceptibility to AD,⁵²⁻⁵⁶ however others have shown no association,⁵⁷⁻⁶⁰ or a reduced susceptibility.^{61,62} With regard to vascular dementia, decreased levels of HDL-c appear to be associated

1.4 - Evidence for the association between blood lipids and dementia

with increased risk,⁶² while for LDL-c, studies have reported both positive and negative associations.^{62,63}

1.4.3 Randomised controlled trials

In terms of the central research of this thesis, RCTs of statin therapy can be used to provide indirect evidence for the effect of reducing blood LDL-c levels on dementia risk. However, RCTs may be infeasible if the outcome of interest is one with a long prodromal period, such as dementia (see Section 1.2.1), as they would require extremely long and costly follow-up.⁶⁴ It is no surprise then that the two previous trials providing evidence on the effect of statins on dementia risk, identified by a recent Cochrane review,⁶⁵ are in fact trials of statins for the prevention of coronary related outcomes.

While being widely cited, these studies have major limitations that reduce their utility as a source of evidence on the effect of statin treatment in assessing the impact of lipid-lowering treatment on dementia risk. Firstly, the criteria used to determine a dementia outcome are unclear. One of the trials, the Prospective Study of Pravastatin in the Elderly (PROSPER) trial,⁶⁶ reported not on dementia outcomes but on the change in cognitive scores over a mean of 3.2 years. As highlighted in Section 1.2.2, a “change in score” alone is insufficient to diagnose a dementia outcome. The second trial, the Medical Research Council/British Health Foundation Protection Study,⁶⁷ found no effect of simvastatin on dementia (OR: 1.00, 95%CI: 0.61-1.65), but did not report how the outcome was assessed/recorded within the trial.

Additionally, the two trials did not make any effort to assign an underlying pathology to each case, instead reporting an all-cause dementia outcome. As discussed in Section 1.2.1, the different underlying pathologies of dementia have different mechanisms of action, and so it is not guaranteed that the effect of statins would be consistent across them.

1.4 - Evidence for the association between blood lipids and dementia

Both trials were also limited by the relatively short follow-up period examined. This limited follow-up was expected, as the primary aim of both trials was to assess the effect of statins on short-term coronary-related outcomes.^{66,67} The PROSPER trial had a mean follow-up of 3.2 years, while the MRC/BHF Protection Study estimated risk at 5 years of follow-up. Given the long lag time between non-symptomatic onset of dementia and clinical presentation, it is likely that these durations are insufficient to fully capture the onset of dementia. Finally, as they included only patients at high vascular risk, their generalisability to other settings is limited.⁶⁵

1.4.4 Mendelian randomisation

Newer methodological approaches, such as Mendelian randomisation (MR),⁶⁸ have also been used to examine the effect of varying lipid levels on dementia risk. Mendelian randomisation attempts to combat the risk of reverse causation and residual confounding inherent to observational studies by using natural random variation in participants' genomes.⁶⁹ In brief, MR uses genetic variants that are both strongly associated with the exposure of interest and are independent from potential confounders to strengthen causal inference.⁶⁸ The analytic method relies on several assumptions about the instrumental variable (IV),⁷⁰ namely that:

1. the IV is associated with the exposure of interest (the relevance assumption);
2. the IV and outcome do not share a common cause (the independence assumption); and
3. the IV does not affect the outcome other than via the exposure (the exclusion restriction assumption).

Recent MR studies indicated that genetically determined low levels of LDL-c may cause a reduction in AD risk.^{71,72} However, the effect was attenuated in sensitivity analysis that exclude the region surrounding the *ApoE* gene, the strongest known risk factor for Alzheimer's disease.⁷³ Inclusion of *ApoE4* variants invalidates the

1.5 - Theoretical framework: Evidence synthesis

exclusion restriction criteria (Assumption 3, above), as the risk reduction observed may be driven by variants in this region via a pathway independent of lipid levels. This was supported by further MR studies where *ApoE4* variants were intentionally excluded.⁷⁴ Despite the increasing number of MR studies examining this topic, no systematic review of this study design as a source of evidence has been performed.

In summary, multiple sources of evidence exist on the relationship between statins and dementia. In the next section, I introduce the synthesis of diverse sources of evidence as the theoretical framework used in this thesis.

1.5 Theoretical framework: Evidence synthesis

Evidence synthesis is the process of finding and integrating information from several sources to examine a research question.⁷⁵ A common type of evidence synthesis is a systematic review, either with or without a meta-analysis.⁷⁶

The results of an evidence synthesis exercise can be used to provide a more definitive answer to that question or, failing that, to highlight gaps in the existing evidence base. The ability to identify these gaps is particularly useful in guiding future research to address questions that have yet to be answered.

This thesis seeks to use an evidence synthesis framework to assess the effect of lipids, and treatments that influence lipid levels, on dementia outcomes. Specifically, this thesis considers three concepts within the umbrella term of evidence synthesis:

- Inclusion of preprints
- Individual participant data meta-analysis
- Triangulation across evidence sources

These three elements are expanded on below and are used to frame the research presented in the subsequent Chapters.

1.5.1 Inclusion of preprints

The importance of including grey literature, defined as literature not published in peer-reviewed journals,⁷⁷ in systematic reviews is widely acknowledged. Meta-research studies have demonstrated that systematic reviews excluding grey literature sources overestimate the effect of interventions.^{78–80} Common, well-accepted forms of grey literature include conference abstracts, government reports and theses.^{77,81}

However, the role of preprints in evidence syntheses is less well defined. Defined by the Committee on Publication Ethics as ‘scholarly manuscript[s] posted by the author(s) in an openly accessible platform, usually before or in parallel with the peer review process’,⁸² preprints serve several purposes. They are used to establish primacy when submitting to a journal where the peer-review process may take several months.⁸³ In addition, they allow for the rapid disseminate research findings, as occurred during the COVID-19 pandemic.⁸⁴ Finally, they are a source of information on manuscripts that may not have been accepted elsewhere, helping to combat publication bias or the “file-drawer” effect.⁸⁵

One of the major criticisms of using preprints as an evidence source is that they have not yet undergone formal peer review.^{86,87} However, this approach assigns substantial weight to peer-review as a indicator of “quality”, and is at odds with the acceptance of non-reviewed conference proceedings as an evidence source.^{81,88} The argument for including preprints as an evidence source is further strengthened by results that demonstrate preprinted studies seldom change following peer review. Meta-studies of the concordance between preprinted and published studies showed that results were broadly comparable between the two, indicating that while the numerical results may change, the overall interpretation of the results were consistent in the majority of cases.^{89–91} This indicates that preprints should be considered a reliable reflection of a given study.

In this thesis, preprints are considered an important source of evidence, in contrast to previous reviews on this topic. However, as with grey literature,⁸⁸ there are several logistical problems with carrying out systematic searches in preprint repositories.

1.5 - Theoretical framework: Evidence synthesis

As such, to enable the inclusion of preprints in the systematic review described in Chapters 3, a new tool addressing these issues is presented in Chapter 2.

1.5.2 Individual participant data meta-analysis

Individual participant data (IPD) meta-analyses are commonly considered to be the gold standard in evidence synthesis methodology.^{92,93} IPD methods seek to obtain the raw data from each study identified in a systematic review, rather than basing the meta-analysis on summary results extracted from the literature.⁹²

In the context of this thesis, if lipids are found to have a causal role in development of dementia, evidence-based preventative strategies would be best informed by identifying the types of individuals who are most likely to receive benefit from treatment with lipid-modifying agents.⁹⁴⁻⁹⁶ However, if primary studies do not present results stratified by covariates of interest, meta-analyses of summary-level data on this topic often have limited ability to examine research questions related to exposure-covariate interactions.⁹² In terms of this thesis, participant age and sex are considered to be of particular interest.^{94,97}

An IPD meta-analysis of lipid levels on dementia outcomes would overcome this limitation of summary-level data, as access to the raw data allows for an analysis that investigates these interactions.⁹⁸ This approach has the added benefit of allowing a common set of inclusion criteria and a common statistical model to be applied across all datasets, potentially eliminating some important sources of heterogeneity.⁹⁹

Despite their advantages, IPD meta-analysis are rarely performed.¹⁰⁰ Factors limiting their uptake include the increased time and effort they require when compared to a summary-level analysis, and the low success rate associated with obtaining the raw data.^{101,102} The data underlying primary studies are frequently not publicly available,^{103,104} and the availability of data “available on request from authors” declines rapidly over time.¹⁰⁵ Several systematic barriers to open data sharing have been identified¹⁰⁶. Of particular concern for biomedical IPD analyses are legal

1.5 - Theoretical framework: Evidence synthesis

issues surrounding the sharing of medical data, motivated by concerns around patient privacy.¹⁰⁷

In response to these limitations, new collaborative initiatives have developed to enable rapid access to relevant data in a secure supported workshop. The most import in relation to this thesis is the Dementia Platform UK (DPUK),¹⁰⁸ which aims to provide access to several dementia-related datasets via a single simplified application process.

I will attempt to obtain the raw data from relevant primary studies identified by the systematic review in Chapters 3 & 4. Any data obtained will be combined with that available from the DPUK portal as part of an IPD meta-analysis in Chapter 6. This will enable the assessment of the effect of lipids on dementia stratified by key variables such as sex.

1.5.3 Triangulating across study designs

As illustrated in Section 1.4, several diverse epidemiological methods have been used to examine the effect of varying blood lipid levels on dementia risk. However, each method is limited by its own biases. Aetiological triangulation is a developing evidence synthesis method that seeks to exploit these inherent differences in study design, and as a result, in biases.¹⁰⁹ If several sources of evidence are available and point towards identical conclusions about an exposure-outcome relationship, and these sources are at risk of unrelated biases, this strengthens our confidence in the result. The ideal scenario is where predicted sources of bias are likely to be in competing directions, one to strengthen the effect of the exposure and the other to attenuate it.¹⁰⁹ As such, triangulating these results can provide a middle-ground between the competing directions of bias. A triangulation approach can also prove useful in helping to prospectively design new studies that are at risk of different sources of bias to those already observed in the published literature.¹¹⁰ However, existing methods for triangulation are limited to a qualitative discussion of the

different identified results with respect to their biases,¹⁰⁹ an approach which faces issues in interpretability when assessing many results.

To address this limitation, this thesis seeks to develop and apply a novel quantitative triangulation approach to answer causal questions on the effect of blood lipids on dementia outcomes. All existing evidence, regardless of study design, is first identified and assessed using domain-based tools by the systematic review presented in Chapters 3 & 4. The existing evidence base is then supplemented by two primary analyses, presented in Chapters 5 and 6. Finally all existing and new evidence is incorporated into the quantiatative triangulation framework in Chapter 7.

1.6 Thesis overview

1.6.1 Aims and objectives

The over-arching aim of this thesis is to explore the relationship between blood lipid levels, and by extension treatments that modify blood lipid levels such as statins, and the subsequent risk of dementia and related outcomes

The specific research objectives that this thesis seeks to address are:

- To create a tool that allows for the inclusion of health related preprints in evidence syntheses in a systematic and reproducible manner
- To review all available evidence across multiple diverse study designs to assess the effect of lipids and lipid regulating agents on dementia risk
- To examine whether there is evidence for an effect of lipid-regulating agents on dementia and related outcomes in a large scale population-based cohort, the Clinical Practice Research Datalink (CPRD)
- To produce evidence on lipid-covariate interactions as part of an IPD meta-analysis

- To propose a generalised framework for the quantitative triangulation across diverse evidence sources

1.6.2 Structure

Chapters are self-contained, presenting the methods and results of that specific research project. They are bookended by introductory and discussion sections which place the methods and results in context. Each chapter is prefaced by a “lay” or plain English summary.

- **Chapter 1:** Background information on dementia and blood lipid levels. This chapter provides an introduction to the topics covered in this thesis to non-subject area experts, and discusses the motivation for the remainder of the thesis.
- **Chapter 2:** This chapter introduces a new tool, `medrxivr`, which was used to developed to allow for systematic searches of the health-related preprint repositories.
- **Chapters 3 & 4:** These chapters describe, respectively, the methods and results of a comprehensive systematic review and meta-analysis. This review examined all available evidence on the effect of blood lipids and interventions that modified blood lipids on dementia outcomes.
- **Chapter 5:** This chapter examines the relationship between lipid-regulating agent use and dementia outcomes in the Clinical Practice Research Datalink, a large primary care electronic health record database.
- **Chapter 6:** This chapter describes an individual participant data analysis of several longitudinal cohort studies to describe the relationship between blood serum lipids and dementia outcomes, stratified by important covariates such as sex.

- **Chapter 7:** This chapter integrates the evidence identified and produced by the preceding chapters as part of a quantitative triangulation framework.
- **Chapter 8:** This chapter summarise the key clinical and methodological findings of the thesis and discusses their implications for practice. The overall strengths and weaknesses of this project are discussed in detail, and further avenues of research are suggested.

1.7 Outputs from this thesis

The outputs of this thesis are detailed below, and include peer-reviewed papers, presentations, and open-source evidence synthesis tools.

1.7.1 Contributions to the scientific literature

During the course of this thesis, I have made several contributions to the scientific literature. Those arising from or directly related to the contents of this submission are presented below.

McGuinness L. A., Higgins J. P. T., Walker, V. M., Davies, N. M., Martin, R. M., Coulthard, E., Davey-Smith, G., Kehoe, P. G., and Ben-Shlomo, Y. (2021) Association of lipid-regulating drugs with dementia and related conditions: an observational study of data from the Clinical Practice Research Datalink. medRxiv 10.1101/2021.10.21.21265131

Preprinted manuscript of the analysis of lipid-regulating agents and dementia outcomes in the CPRD, which is presented in Chapter 5.

McGuinness, L. A., and L Schmidt. (2020) *medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R*. *Journal of Open Source Software* 5.54 2651. DOI: 10.21105/joss.02651

A paper introducing the open-source preprint search tool described in Chapter 2. As is common for journal articles describing software, the paper is intentionally short providing only a broad overview of the tool while extensive documentation is available from the project website (see Section 2.1 for more details).

Hennessy, E. A., Acabchuk, R., Arnold, P. A., Dunn, A. G., Foo, Y. Z., Johnson, B. T., Geange, S. R., Haddaway, N. R., Nakagawa, S., Mapanga, W., Mengersen, K., Page, M., Sánchez-Tójar, A. Welch, V., and McGuinness L. A. (2021). *Ensuring Prevention Science Research is Synthesis-Ready for Immediate and Lasting Scientific Impact*. *Prevention Science* . DOI: 10.1007/s11121-021-01279-8

The experience of extracting data for the systematic review in Chapters 3 & 4 inspired a practical guide for researchers. This piece was co-written with Dr. Emily Hennessy (see Author Declarations in the front materials).

McGuinness, L. A., and Higgins J. P. T. (2020) “Risk-of-bias VISualization (*robvis*): An R package and Shiny web app for visualizing risk-of-bias assessments.” *Research Synthesis Method*). DOI: 10.1002/jrsm.1411

The tool used to visualise the risk-of-bias assessments in Chapters 3 & 4 has been published in Research Synthesis Methods. See Appendix B.2 for more details on this tool. Note that this publication does not describe the recently-developed functionality for producing bias direction plots, as described in Chapter 7.

McGuinness, L. A., and Sheppard A. L. 2020. “A Descriptive Analysis of the Data Availability Statements Accompanying Medrxiv Preprints and a Compar-

1.7 - Outputs from this thesis

ison with Their Published Counterparts.” PLOS ONE 16(5): e0250887. DOI: 10.1371/journal.pone.0250887

Using the tool described in Chapter 2, I led a “research-on-research” study to assess the concordance between the openness of data availability statements accompanying a sample of medRxiv preprints and their published counterparts.

For information on additional contributions to the scientific literature not directly related to this thesis, see Appendix A.1.1.

1.7.2 Presentations/Talks

“Identifying and triangulating all available evidence on the effect of blood lipids and statins on dementia outcomes”: Poster presentation, Alzheimer’s Association International Conference 2021.

“medrxivr: A new tool for searching for and retrieving records and PDFs from the medRxiv preprint repository”: Accepted oral presentation abstract, Cochrane Colloquium 2020 (note: event was cancelled due to the COVID-19 pandemic).

“On the shoulders of giants”: advantages and challenges to building on established evidence synthesis packages, using the {robris} package as a case study”: Oral presentation, Evidence Synthesis and Meta-Analysis in R Conference (ESMARConf) 2021.

“RoB 2.0: A revised tool to assess risk of bias in randomised trials”: Webinar, co-presented with Dr. Theresa Moore as part of the Evidence Synthesis Ireland Methods Series.

1.7.3 Software

medrxvir

An R package that allows users to search and retrieve bibliographic data from the medRxiv¹¹¹ and bioRxiv¹¹² preprint repositories. See Chapter 2 for more details. Install a stable version of the package from the Comprehensive R Archive Network (CRAN), or alternatively install the development version from GitHub, using:

```
# CRAN version
install.packages("medrxivr")

# Development version
devtools::install_github("ropensci/medrxivr")
```

triangulate

An R package containing function to implement the quantitative triangulation approach detailed in Chapter 7. At present, only the development version of the package is available, which can be installed from GitHub using:

```
# Development version
devtools::install_github("mcguinlu/triangulate")
```

robvis

An R package and associated shiny web application that allows users to visualize the results of the risk-of-bias assessments performed as part of a systematic review. See Appendix B.2 for more details. Install a stable version of the package from CRAN, or alternatively install the development version from GitHub, using:

```
# CRAN version
install.packages("robvis")
```

```
# Development version  
devtools::install_github("mcguinlu/robvis")
```

1.8 Summary

- In this introductory chapter, I provided background information on the core elements of the central research question and framed the research presented in the context of a theoretical framework of evidence synthesis. I outlined the central aim of the thesis and provided an overview of the individual analyses performed.
- I also described the contributions of this thesis to the scientific literature. Finally, I discussed the research tools and software developed to support the analyses in this thesis, one of which (the preprint search tool) is introduced more fully in the following chapter.

Why are open source statistical programming languages the best?

Because they R.

— Chris Beely, 2013¹¹³

2

medrxivr: an R package for systematically searching biomedical preprints

Contents

Lay summary	28
2.1 Introduction	28
2.2 Development	30
2.3 Usage	32
2.4 Discussion	39
2.5 Summary	42

Lay summary

Preprints are copies of academic manuscripts that are posted online in advance of being formally published by an academic journal. They represent an important source of scientific literature. A new software program called `medrxivr` was created as part of this thesis to allow researchers to find preprints related to their research in a transparent and reproducible way. Development of this tool was an essential part of this thesis, as preprints represent a key source of information needed for the research reported in future chapters.

2.1 Introduction

Preprints represent an increasingly important source of scientific information (see Section 1.5.1). As a result, repositories of preprinted articles should be considered a distinct but complementary information source when reviewing the evidence base as part of a systematic review. The two key repositories in the health science are bioRxiv, established in 2013,¹¹² and medRxiv, which launched in 2019 and was designed to replace the “Epidemiology” and “Clinical Trial” categories of bioRxiv.¹¹¹

Searching these preprints as part of the systematic review described in Chapters 3 & 4 was a necessity, as many of the existing reviews on the topic of lipids and dementia have not considered this important source of evidence. At the time of writing, however, the bioRxiv and medRxiv websites allow only simple search queries as opposed to the complex searches containing Boolean logic operators (AND/OR/NOT) that information specialists use to query other major databases.^{114,115} Additionally, the best available extraction mechanism for obtaining references for all records returned by a search involved going through each record and downloading individual citations one-by-one. As the scale of these preprint databases increase, particularly in light of the massive expansion of the medRxiv repository as a result of COVID, this already time-consuming and error-prone method is no longer feasible.

This chapter outlines the development and key functionality of `medrxivr` (version 0.0.5), a tool I created to facilitate the systematic searching of medRxiv and




Figure 2.1: Role of `medrxivr` in a systematic review workflow - `medrxivr` allows for systematic searching of biomedical preprints as part of the initial literature searching. Following title and abstract screening, reviewers can then programmatically retrieve a copy of the PDF of included records to facilitate the full-text screening stage (similar to Endnote's "Find Full Text" feature).

bioRxiv preprints. The factors that necessitated the development of this tool in the context of this thesis are outlined, and the use of `medrxivr` in my own projects and by other researchers is discussed. As the majority of work on this aspect of my thesis is represented by lines of code or online documentation (available at <https://github.com/ropensci/medrxivr> and <https://docs.ropensci.org/medrxivr/> respectively), this chapter is an intentionally short, high-level summary of my work on this project. The GitHub repository for the `medrxivr` contains a complete record of the development of this tool, including discussion with other members of the systematic review community.¹¹⁶

2.2 Development

2.2.1 Success criteria

I developed the tool to meet three success criteria,¹¹⁷ influenced both by the functionality required to perform systematic searches as part of the review presented in the subsequent chapters, discussion with information specialist colleagues, and an informal survey of the evidence synthesis and health librarian communities on Twitter. The criteria were as follows:

1. reliable, reproducible and transparent search functionality, allowing for Boolean (AND/OR/NOT) operator logic;
2. support for bulk export of references returned by the search to a file type that can be readily imported into a reference manager (e.g., *.bib* or *.ris*); and
3. automated retrieval of the full-text PDFs of relevant records, similar to the Find Full Text feature offered by EndNote.

2.2.2 Alternative medRxiv/bioRxiv interfaces

Prior to development of this tool, I conducted an audit of existing tools for accessing medRxiv and bioRxiv metadata. While none addresses the success criteria described above, two of these tools are useful to consider to highlight the additional functionality that `medrxivr` contributes.

The first, a platform called Rxivist¹¹⁸, allows users to search preprints using keywords. However, the core functionality of the Rxivist platform is focused around exploring the number of times a preprint has been downloaded and/or shared on Twitter, to allow researchers to find the most popular papers related to their topic. The search interface¹¹⁹ does not allow for complex search strategies using Boolean operators and there is no option to batch-export the results of a search.

The second tool, `search.bioPreprint`, allows users to search for terms across a range of preprint servers, including medRxiv and bioRxiv, but also journals which use a post-publication peer-review process such as F1000Research.¹²⁰ However, similar to the Rxivist platform, this tool is designed for researchers aiming to keep up to date with recent developments in their fields rather than systematically assess the entirety of the available literature. As such, the platform only returns the most recent 1,000 records by publication date.

Finally, neither tool provides an easy way to programmatically download a copy of the PDF of relevant preprints as part of the preparation for the full-text screening stage of a systematic review.

2.2.3 Early versions

Work on the `medrxivr` tool began in Summer 2019, and initially consisted of a development of set of R scripts to allow for searching medRxiv and bioRxiv as part of the systematic search outlined in Chapter 3. Following interest from other researchers in using the *ad-hoc* web-scraping scripts, additional development work took place in 2019/2020, allowing for improved searching and exporting functionality and I released the initial version of the `medrxivr` R package in February 2020.

Early versions of the tool had a reliance on scraping data directly from the repository website. Web-scraping is a fragile mechanism for extracting data, as it is entirely dependent on consistent website design and underlying code structure remaining unchanged.^{121,122}. In the case of `medrxivr`, as the medRxiv/bioRxiv websites are regularly updated, ensuring the web-scraping performed as expected required that I regularly update or fix the script.

However, an application programming interface (API) for the medRxiv and bioRxiv repositories was made public in early 2020 by the institution responsible for managing these preprint repositories, the Cold Springs Harbor Laboratory. This allowed for newer versions of the `medrxivr` package to engage in active “fault

prevention” and provide a more robust interface to the data by removing the reliance of web-scraping.¹²²

2.2.4 Package infrastructure

I wrote the `medrxvir` package in R using RStudio,¹²³ and followed development best-practice, including detailed documentation, a robust unit testing framework (99% of all code lines within the package are formally tested across multiple platforms including Windows, MacOS, and Linux), and in-depth code review by two experienced, independent reviewers.

2.3 Usage

The `medrxivr` R package is split into two component parts:

- an interface to the Cold Springs Harbor Laboratory API, which imports medRxiv and bioRxiv metadata into R; and
- a collection of functions for working with the imported metadata, with an explicit focus on searching this data as part of a systematic review or evidence synthesis project.

The standard workflow is to download a copy of all metadata contained in the repository, and then to perform searches on this local copy. This is a workaround as the Cold Springs Harbor Laboratory API does not provide any functionality to search the database.

While the package allows users to interact with and search both medRxiv and bioRxiv metadata, as the process is identical for both, searching the medRxiv database is used as an illustrative example throughout this chapter.

2.3.1 Installation

`medrxivr` has been released to the Comprehensive R Archive Network (CRAN), and can be installed with the following code:

```
install.packages("medrxivr")
```

Alternatively, the development version of the package can be installed from GitHub:

```
# install.packages("devtools")
devtools::install_github("ropensci/medrxivr")
```

2.3.2 Importing preprint metadata

Prior to searching the metadata, it must first be imported in R. In `medrixvr`, I have provided two separate but related methods for users to import the data (Figure 2.2). The first of these methods, accessed via the `mx_api_content()` function, creates a local copy of all data available from the medRxiv API at the time the function is run.

```
# Get a copy of the database from the live medRxiv API endpoint
mx_data <- mx_api_content()
```

This provides an up-to-the-minute reflection of the medRxiv preprint repository. However, this approach has two limitations. Firstly, as the API returns results as a series of pages limited to 100 records per page, downloading the entire database requires a time-intensive process of cycling through each page. Secondly, the API can become unavailable, either during peak usage times or planned maintenance windows. To address these limitations, I provide a second method of accessing

medRxiv data, called via the `mx_snapshot()` function, which allows users to access a maintained static snapshot of the database.

```
# Import a copy of the medRxiv data from the snapshot  
mx_data <- mx_snapshot()
```

This snapshot is created each morning at 6am using a process known as “git-scraping”,¹²⁴ whereby the entire database is downloaded using the `mx_api_content()` function and saved as a comma separated value (CSV) file in an online repository (Figure 2.2). Calling `mx_snapshot()` imports this CSV into R, and has the advantage of both faster loading of the data into R (as it is imported as a single file and does not require cycling through the output of the API) and an absence of any reliance on the API. The one limitation of this approach is that the snapshot (by its nature) will not contain details of records added to the database since it was taken. However, given that the number of records added each day is relatively low, this should pose minor issues.




Figure 2.2: Overview of `medrxivr` data sources - Users can either access the API directly via `mx_api_content()`, or can import a maintained snapshot of the database, taken each morning at 6am, via the `mx_snapshot()` function. Note: due to the size of bioRxiv, only a maintained snapshot of the medRxiv repository is available via `mx_snapshot()`.

2.3.3 Performing a search

Once a local copy of the metadata is created, the first step in searching it is to create a search strategy. Search terms to be combined with the OR operator are contained in vectors (`c(...)`), while topics to be combined with the AND operator are contained in lists (`list(...)`).

```
# Create the search query

topic1 <- c("dementia", "alzheimer's") # Combined with OR
topic2 <- c("lipids", "statins")        # Combined with OR

myquery <- list(topic1, topic2)          # Combined with AND
```

For example, when written in standard syntax, the search contained in the `myquery` object above would be: “((dementia **OR** alzheimer’s) **AND** (lipids **OR** statins))”. There is no limit to the number of search terms that can be included in each topic, nor in the number of topics that can be searched for. Search terms can also contain common syntax used by systematic reviewers and health librarians, including the NEAR operator for the identification of co-localised terms and wild-cards which allow for alternate spellings (e.g. “randomisation” vs “randomization”).

Once a strategy has been defined, it is passed along with the local copy of the database to the `mx_search()` function.

```
# Run the search

results <- mx_search(mx_data,
                      myquery)
```

2.3.4 Refining a search

An important argument of the `mx_search()` is `report`, which outputs a structured table with each search strategy presented on an individual line and the number of records associated with this strategy.¹²⁵

```

results <- mx_search(mx_data,
                      myquery,
                      report = TRUE)

## Found 1 record(s) matching your search.

##
## Total topic 1 records: 224
## dementia: 224
## alzheimer's: 0
##
## Total topic 2 records: 119
## lipids: 90
## statins: 33

```

This allows users to discover which terms in their search are contributing most to the total number of results returned. This is important as part of developing a search strategy,¹¹⁴ as it allows for the key terms related to each topic to be discovered. It also aids in identifying misspelled or case-sensitive search terms, which will frequently return no results. As an example, in the search presented above, the term “alzheimer’s” returns no records. This is expected, as “Alzheimer’s” is a proper noun and so should be capitalised, but serves to illustrate the usefulness of the reporting function.

In hindsight, the decision to make the search case-sensitive was a poor one as users reported being unaware of this setting when searching. However, changing the default behaviour of `mx_search()` to be case-insensitive is problematic, as it could result in different numbers of hits based solely on which version of the package the user has installed. To address this, `mx_search()` now contains an `autocaps` argument, which when set to `TRUE` will automatically search for both capitalised and uncapitalised versions of the search term.

2.3.5 Exporting to a bibliography file

In line with my second success criteria (Section 2.2.1), one of the key features of the `medrxivr` is the ability for users to export easily the results of their systematic search to a reference manager. While it is a seemingly simple request, this is is one of the key ways in which `medrxivr` is set apart for other preprint search tools, including the native medRxiv/bioRxiv website search functionality.

For example, the results of our simple search above can be exported to the "`medrxiv_export.bib`" file using the following code:

```
mx_export(results,
          file = "medrxiv_export.bib",
          report = TRUE)
```

2.3.6 Downloading the PDFs of relevant records

`medrxivr` also allows users to download the full text papers for records that are deemed eligible for full-text screening (see Figure 2.1). `mx_download()` takes the list of included records and saves the PDF for each to a folder specified by the user. This functionality is similar to the “Find Full Text” feature offered by EndNote.

```
mx_download(results, # Search results, less excluded records
            "pdf/") # Directory to save PDFs to
```

2.4 Discussion

2.4.1 Reception and future plans

The tool has been well received by the community (as of December 2021, `medrxivr` has been downloaded more than 6100 times), and several use cases have been reported. It has been used to investigate the role of preprints in the response to the 2019 coronavirus outbreak,¹²⁶ perform searches of preprints as part of a systematic review,^{127,128} and examine how data-sharing behaviour is affected by journal policies (see 1.7).¹²⁹

The package has been accepted into the rOpenSci suite of packages, a collection of “carefully vetted, staff- and community-contributed R software tools that lower barriers to working with scientific data sources on the web”.¹³⁰ As part of this process, following rigorous peer-review, an associated article introducing the tool was published by the Journal of Open Source Software.¹³¹ The entire review discussion is publicly available and can be viewed online.¹³² The tool has also been well received by the open-source community, demonstrated by the engagement of other developers in contributing to important new functionality and suggesting bug-fixes.

Lobbying of the Cold Springs Harbor Laboratory to develop the API to allow for direct searching of the database has been ongoing. This would negate the current need to download a local copy of the relevant preprint database before searching it, which is currently the rate limiting step for performing searches. For example, as of January 2021, downloading a copy of the bioRxiv database takes approximately an hour.

2.4.2 Use cases

In addition to being used to search systematically health-related preprint servers, as illustrated in the systematic review presented in Chapters 3 & 4, `medrxivr` has other uses.

2.4 - Discussion

For example, I led a descriptive analysis of the change in data availability statements between preprinted and published versions of the same manuscript, stratified by journal data sharing policy access, underpinned by preprint meta-data provided by `medrixvr`. By comparing the preprinted and published versions of the data availability statement, I could examine the same manuscript - same content, authors and funders - under two different publication policies (preprint server vs peer-review journal). This comparison provides evidence on whether stricter policies which require data sharing as a condition of publication result in increased data availability. The analysis found some evidence that data availability statements more frequently described open data on publication when the journal mandated data sharing, compared to when the journal did not mandate data sharing. This study has since been published,¹²⁹ and a copy is included in Appendix B.3.

Secondly, using `medrxivr`, an analysis of the publication rate for medRxiv preprints was performed (see Appendix A.2.1). 87 of the 129 records (67.4%) posted on medRxiv in July 2019 were published by 30th July 2021 (i.e. allowing for a two-year lag between preprint posting and publication). This finding agrees with previous work demonstrating that two-thirds of bioRxiv preprints are published in a peer-reviewed journal within two years of posting.¹³³ This analysis further illustrates that a substantial number of preprints are never formally published but remain accessible as preprints.

In summary, the meta-research/methodological analyses described above illustrate that easy access to medRxiv/bioRxiv metadata has applications beyond the inclusion of preprints in systematic review.

2.4.3 Limitations of `medrxivr`

While searching of the medRxiv and bioRxiv databases was crucial for the systematic review element of my thesis, there are some important limitations to note here. A key example is that the tool only searches the available metadata of preprint records

(the title, abstract and keywords), rather than the full text of preprints, and so some relevant records might be missed. However, this approach echoes that used by other search platforms such as OvidSP, and while some relevant records may be missed (reduced sensitivity), limiting the search to the metadata fields prevents non-relevant records from being returned (high specificity). A key example of the reduced specificity when searching the full text, identified during development of `medrxivr`, is that a search for “dementia” would return a record where the only occurrence of this term is in the title of one of the references.¹³⁴

There is also the potential that the cross-section of literature posted on medrxiv/bioRxiv is substantially different to those suffering from publication bias (studies or analyses that are not published for a range of reasons including results that are not deemed “novel” or are not statistically significant).¹³⁵ This is because simply lowering the barriers to publication may well encourage authors to publish “null” results, but due to the effort involved in writing up a distributable manuscript, it is unlikely to completely address the “file drawer” effect.⁸⁵

It is probably too early (and likely too methodologically difficult) to tell whether the increased popularity and acceptance of preprint repositories will have any effect of the availability of research that was not considered “publishable” at other venues.

2.4.4 Role of open source tools in evidence synthesis

Part of the motivation for creating the `medrxivr` tool was a belief that the development and distribution of open source scripts and tools should be a fundamental part of evidence synthesis research.^{136,137} In the case of `medrxivr`, it is likely that several other evidence synthesists had written personal scripts that have a similar, or related, functionality - in fact, following development of the tool, I identified one other researcher that has done so (Nicholas Fraser, author of the `rbiorxiv` package, which allows for importing medRxiv metadata into R but does not provide search functionality).¹³⁸ If these scripts continue to be developed in

private and are never shared or publicised, this will inevitably hamper the efforts of the evidence synthesis community, not only in terms of duplication of time and effort but also due to lost opportunities for collaboration.¹³⁷ Creating and sharing well-documented packages, the recognised standard for sharing code in R, represents one way to reduce this inefficiency.¹³⁹

2.5 Summary

- In this Chapter, I have introduced a new tool, `medrxivr`, for performing complex systematic searches of the medRxiv and bioRxiv preprint repositories.
- I have outlined the motivation for developing this tool in relation to this thesis - more specifically, that it was used to perform systematic and reproducible searches of key literature sources used in the comprehensive systematic review described in Chapters 3 & 4.
- I have contrasted `medrxivr` with other available interfaces to medRxiv/bioRxiv data to highlight the added functionality it offers. I have also discussed the tools reception to date, its limitations, and the important role of open-source tools like `medrxivr` in evidence synthesis.

“It is surely a great criticism of our profession that we have not organised a critical summary by speciality or sub-speciality, up-dated periodically, of all relevant RCTS.”

— Archibald Cochrane, 2000¹⁴⁰

3

Systematic review of existing evidence on the association between blood lipids and dementia outcomes: Methods

Contents

Lay summary	43
3.1 Introduction	44
3.2 Methods	45
3.3 Summary	62

Lay summary

Systematic reviews are a type of research study that aim to collect and combine all existing evidence to provide the best possible answer to an important research question. Well-performed reviews involve multiple steps including: searching for existing studies; assessment of the studies against predefined inclusion criteria; collection of data from each study; and assessment of each study's method and results.

This chapter presents the methods used to perform a systematic review of primary studies that have examined the relationship between the levels of blood lipids (such as cholesterol and triglycerides) and dementia outcomes. In addition, the review examined the relationship between treatments that change blood lipid levels, such as statins, and dementia outcomes. The results of this systematic review are then presented in Chapter 4.

3.1 Introduction

In this chapter, I describe a comprehensive systematic review of the relationship between blood lipid levels (and treatments that modify them) and the subsequent risk of dementia and related outcomes.

This analysis sought to address two specific aims. Firstly, as discussed in the introduction to this thesis (Section 1.4), several diverse forms of evidence on the relationship between lipids and dementia exist. These include randomised controlled trials, observational studies of different design, and Mendelian randomisation studies. However, based on a scoping review of existing literature, no previous evidence synthesis exercise has attempted to examine the association of lipids/statins with dementia outcomes across these distinct evidence types. Collating these diverse evidence sources is important, as, if the observed association between lipids and dementia is constant across them, it increases our confidence in the association. As such, the primary aim of this analysis was to systematically review all available literature describing prospective analyses, regardless of study design.

Secondly, I explicitly sought to include health-related preprint servers as a potential evidence source in this review, as they are infrequently considered by evidence synthesists but can report relevant unpublished analyses. As a sensitivity analysis to the review presented in this chapter, I sought to quantify the additional evidential value of including preprints, making use of the preprint search tool presented in Chapter 2.

Given the size of the review, I have separated the methods and results into two chapters for ease of reading. This chapter details the methodology used, while Chapter 4 presents the results.

3.2 Methods

3.2.1 Protocol

A pre-specified protocol for this analysis was registered on the Open Science Framework platform and is available for inspection.¹⁴¹ Deviations from the protocol are summarised in Appendix A.3.1.

3.2.2 Contributions

In line with best-practice guidance, secondary reviewers were used to check the accuracy of screening, data extraction and risk-of-bias assessment processes. Due to the scale of the project, this review was performed in conjunction with a team of secondary reviewers (see Acknowledgements and Author Declaration in the front matter).

3.2.3 Eligibility criteria

Inclusion criteria

I sought to include studies that examined blood lipid levels as a risk factor for dementia outcomes, defined either as binary hypercholesterolemia variable or by category/1-standard-deviation increase in a specific lipid fraction (total cholesterol, high- and low-density lipoprotein cholesterol and triglycerides). I also aimed to include studies examining the effect of lipid-regulating agents (LRA) as a source of indirect evidence. LRA are treatments used to modify blood lipids levels, and common examples include statins, fibrates, and ezetimibe.

Eligible study designs were randomised controlled trials of LRA, cohort studies examining blood lipids or LRA, and genetic instrumental variable studies (more commonly known as Mendelian randomisation studies) examining the effect of genetically increased/decreased blood lipid levels.

Eligible studies screened participants for dementia at baseline and excluded any prevalent cases. Alternatively, where no baseline screening was employed, participants were assumed to be dementia free if less than 50 years of age at baseline. Studies of any duration were included to allow for exploration of the effect of length of follow-up on the effect estimate. No limits were placed on the sample size of included studies. Eligible studies defined dementia outcomes according to recognised criteria. Examples of acceptable criteria included the International Classification of Diseases (ICD),¹⁴² National Institute of Neurological Disorders and Stroke Association-Internationale pour la Recherche en l’Enseignement en Neurosciences (NINDS-AIREN),¹³ or Diagnostic and Statistical Manual of Mental Disorders (DSM) criteria.³ Studies utilising electronic health records were the sole exception to this, as it was assumed that health care professionals employed their own expertise when entering the outcome into the EHR.

Conference abstracts with no corresponding full-text publication were eligible, and where required, I contacted authors to obtain relevant additional information

not available from the abstract. No limitations were imposed on publication status, date, venue or language.

Exclusion criteria

Due to the significant impact of a memory-related outcome such as dementia on exposure recall, case-control studies were excluded, though case-control studies where historical records are used to determine the exposure status were eligible for inclusion. Cross-sectional studies, qualitative studies, case reports/series and narrative reviews were also excluded, as were studies that measure change in continuous cognitive measures (e.g. MoCA score) without an attempt to map these scores to ordinal groups (e.g. no dementia/dementia).¹⁴³ Previous systematic reviews were not eligible for inclusion, but their reference lists were screened to identify any potentially relevant articles.

Studies with outcomes not directly related to the clinical syndrome of dementia (e.g., neuroimaging) and studies where there was no screening for dementia at baseline were excluded. An exception to the latter criteria was made if the sample was initially assessed in mid-life (i.e. below the age of 50) were excluded. Additionally, studies implementing a “multi-domain intervention” where a lipid-regulating agent is included in each arm were excluded. For example, a study examining exercise + statins versus statins alone would be excluded, but a study examining exercise + statins vs exercise alone would be eligible as the effect of statins can be estimated. Finally, studies using a dietary intervention, for example an omega-3 fatty acid enriched diet, were excluded as it is difficult to disentangle the effect of other elements contained within the diet. Note that this is distinct from studies which delivered a simple tablet-based omega-3 intervention, which would have been eligible for inclusion.

3.2.4 Information sources and search strategy

I systematically searched several electronic bibliographic databases to identify potentially relevant entries (hereafter referred to as “records”). The following databases were searched in June 2019 from inception onwards: Medline, EMBASE, Psychinfo, Cochrane Central Register of Controlled Trials (CENTRAL), and Web of Science Core Collection. As the contents of the Web of Science Core Collection can vary by institution,¹¹⁵ the specific databases and date ranges for each database searched via this platform are listed in Appendix A.3.2. The search strategy used in each database was developed in an iterative manner using a combination of free text and controlled vocabulary (MeSH/EMTREE)⁸¹ terms to identify studies which have examined the relationship between blood lipids levels and dementia, incorporating input from an information specialist. The strategy included terms related to lipids, lipid modifying treatments, and dementia, and was designed for MEDLINE before being adapted for use in the other bibliography databases listed. A high-level outline of the strategy is presented in the Table 3.1 below and the full search strategies for each database are presented in Appendix A.3.3.

Table 3.1: Summary of systematic search by topic. The full search strategy including all terms and the number of hits per term is included in Appendix A.3.3.

No.	Concept
1	Dementia
2	Lipids
3	Lipid-modifying treatments
4	1 AND 2
5	1 AND 3
6	4 OR 5
7	Animals NOT (Animals AND Humans)
8	6 NOT 7
9	Observational filter
10	Randomised controlled trial (RCT) filter
11	Mendelian randomisation/Instrumental variable filter
12	OR/ 9-11
13	8 AND 12

For all topics, search queries were comprised of relevant free text & controlled vocabulary terms.

When searching the bibliographic databases, study design filters (Lines 9-11 in Table 3.1) were employed to try to reduce the screening load. To ensure that the study design filters were not excluding potentially relevant records, a random sample of 500 records identified by the main search but excluded by the filters (defined as “8 NOT 12” in Table 3.1) was screened.

I also searched clinical trial registries, for example ClinicalTrials.gov, to identify relevant randomised controlled trials. In addition, I searched the bioRxiv and medRxiv preprint repositories using the tool developed in Chapter 2 to identify potentially relevant preprinted studies (see Appendix A.3.4 for the code used to search these preprint repositories).

Grey literature was searched via ProQuest, OpenGrey and Web of Science Conference Proceedings Citation Index, while theses were accessed using the Open Access Theses

3.2 - Methods

and Dissertations portal. In addition, the abstracts list of relevant conferences (e.g. the proceedings of the Alzheimer’s Association International Conference, published in the journal *Alzheimer’s & Dementia*) were searched by hand. Finally, the reference lists of included studies were searched by hand, while forward and reverse citation searching (“snowballing”) was performed using Google Scholar.^{144,145}

3.2.5 Study selection

Records were imported into Endnote and de-duplicated using the method outlined in Bramer et al. (2016).¹⁴⁶ In summary, this method uses multiple stages to identify potential duplicates. The initial step involves automatic deletion of records matching on multiple fields (“Author” + “Year” + “Title” + “Journal”), and is followed by manual review of articles with less overlap (e.g. those identified as duplicates based on the “Title” field alone).

Following de-duplication of records, screening (both title/abstract and full-text) was performed using a combination of Endnote and Rayyan. Endnote is a citation management tool,¹⁴⁷ and Rayyan is a web-based screening application.¹⁴⁸ Title and abstract screening to remove obviously irrelevant records was performed, with a random ~10% sample of excluded records being screened in duplicate to ensure consistency with the inclusion criteria. Additionally, I re-screened the same ~10% sample with 3 month lag to assess intra-rater consistency.

Similarly, I completed all full-text screening, with a random ~20% being screened by a second reviewer. Additionally, any records I identified as being difficult to assess against the inclusion criteria were also screened by the second reviewer. Reasons for exclusion at this stage were recorded. Disagreements occurring during either stage of the screening process were resolved through discussion with a senior colleague. A PRIMSA flow diagram was produced to document how records moved through the review.¹⁴⁹

3.2.6 Validation of screening process

Inter- and intra-rater reliability during the screening stages were assessed for a 10% sub-sample of records. Intra-rater reliability involved a single reviewer applying the inclusion criteria to the same set of records while blinded to their previous decisions (i.e. assessment of consistency), while inter-rater reliability involved two reviewers independently screening the same set of records (i.e. assessment of accuracy).

Rater reliability was assessed using Gwet's agreement coefficient (AC1).¹⁵⁰ This measure was chosen over other methods such as percent agreement (number of agreements divided by total number of assessments), as it accounts for chance agreement between reviewers but does not suffer from bias due to severely imbalanced marginal totals in the same way that Cohen's *kappa* value does.¹⁵⁰ Given the small number of included studies in this review as a proportion of the total number screened, this is a useful characteristic. The formulae used to calculate the coefficient are given in Appendix A.3.5.

Interpretation of agreement coefficients is a widely debated topic, and while arbitrary cut-off values may mislead readers,¹⁵³ they provide a useful rubric by which to assess inter-rater agreement. Here, I used guidelines based on a stricter interpretation of the Cohen's *kappa* coefficient,¹⁵⁴ presented in Table 3.2.

Table 3.2: Suggested ranges to aid in interpretation of Gwet's AC1 inter-rater reliability metric¹⁵⁴

Kappa	Interpretation
0 – 0.20	None
0.21 – 0.39	Minimal
0.40 – 0.59	Weak
0.60 – 0.79	Moderate
0.80 – 0.90	Strong
> 0.90	Almost perfect

Intra- and inter-rater reliability was assessed against these cut-offs. If this assessment demonstrates an issue with the screening process (defined as an AC1 of less than 0.9), it indicates the need for a larger proportion of records to be dual-screened.

3.2.7 Data extraction

Data extraction was performed using a piloted data extraction form. I extracted all data in the first instance, which was subsequently checked for accuracy by a second member of the review team. Extracted items included:

- Article metadata: year of publication, author, journal)
- Study characteristics: study location, data source, exposure, outcomes, diagnostic criteria used)
- Patient characteristics: age, sex, baseline cognition scores, baseline education scores
- Results: exposure, outcome, effect measure, effect estimate, error estimate, p-value).

Grouping multiple reports into studies

As part of the data extraction process, multiple records resulting from a single analysis were included and grouped into single units, hereafter called studies. This was most common in cases where multiple records reported on the same analysis but at different time points. In this case, the result corresponding to the longest follow-up was used. Grouping records into studies builds out the most comprehensive account of a given analysis by incorporating information from all available records.

3.2 - Methods

This was particularly relevant to preprints and published papers reporting the same study, which were not considered to be duplicate records but instead different reports of the same study. This decision was taken due to the potential for the published version to offer some information that the preprint did not, and vice versa.

Combining across groups

In line with best practice, where summary data was presented across two groups (e.g. age at baseline stratified by hypercholesterolemia status), the following approach was used to combine the groups:¹⁵⁵

$$N = N_1 + N_2 \quad (3.1)$$

$$Mean = \frac{(N_1 M_1 + N_2 M_2)}{(N_1 + N_2)} \quad (3.2)$$

$$SD = \sqrt{\frac{(N_1 - 1)SD_1^2 + (N_2 - 1)SD_2^2 + \frac{N_1 N_2}{N_1 + N_2}(M_1^2 + M_2^2 - 2M_1 M_2)}{N_1 + N_2 - 1}} \quad (3.3)$$

where N_i , M_i and SD_i denote the number of participants, mean and standard deviation in the i th subgroup. This approach was implemented in a systematic manner, with the raw group data being extracted and a cleaning script employed to combine the groups for analysis.

Harmonisation of cholesterol measures

Where necessary, lipid levels reported in $mmol/L$ were converted in mg/dL using the following formula:

$$mg/dL = mmol/L \times Z \quad (3.4)$$

where $Z = 38.67$ for total cholesterol, LDL-c and HDL-c, and $Z = 88.57$ for triglycerides.¹⁵⁶ The choice of mg/dL was influenced by the widely-used categories of lipids levels on the mg/dL scale, as shown in Table 1.2 in Section 1.3.1.

Following up with authors

Where additional data, not included in the report of an analysis, were required either for data extraction or risk-of-bias assessment, the corresponding author of the study was contacted. This approach was taken due to the potentially large impact of additional information obtained through contact with study authors on the results of the review.¹⁵⁷

3.2.8 Risk-of-bias assessment

A key aim of the review presented here is to identify different sources of evidence at risk of a diverse range of biases, and to contrast and compare findings across them (see Section 7.1 for an overview of triangulation and Chapter 7 for the results of this analysis). To enable this triangulation exercise, detailed and structured risk-of-bias assessment formed an important part of this review.

There has been a recent movement within the evidence synthesis community away from examining *methodological quality* to assessing *risk of bias*,^{158,159} and thus directly evaluating the internal validity of a study. Internal validity is defined here as the absence of systematic error (or bias) in a study, which may influence its results.^{160,161}

This move was prompted by an unclear definition of methodological quality which could include facets such as unclear reporting, in addition to challenges in the

comparison of results from different tools. As part of this shift, the focus moved from checklist or score based tools towards domain-based methods, in which different potential sources of bias in a study are assessed in order. Finally, the new tools move from assessing bias at the study level to considering separately each individual numerical result reported. For example, a study may report on the efficacy of an intervention at six months and two years follow-up. In this case, the proportion of missing outcome data may not be an issue at six months but may introduce bias after two years of follow-up. In this case, assigning a single risk-of-bias judgement to the study as a whole masks the different biases applicable to each unique result.

In this review, domain-based tools were used to assess the risk of bias for each result in each included study. The study design-specific tools are introduced and discussed in more detail in the following sections.

Randomised controlled trials

Randomised controlled trials were assessed using the RoB 2 tool.¹⁶² The tool assesses the risk of bias across five domains:

- bias arising from the randomisation process,
- bias due to deviations from intended intervention,
- bias due to missing outcome data,
- bias in measurement of the outcome,
- bias in selection of the reported result.

Acceptable judgements for each domain include: “low risk”, “some concerns”, “high risk”. Each of the five domains contains a series of signaling questions or prompts which guide the user through the tool. Once a domain-level judgement for each domain has been assigned, an overall judgement, using the same three levels of risk of bias, is assigned to the result.

Non-randomised studies of interventions/exposures

For non-randomised studies of interventions (NRSI), I used the ROBINS-I (Risk Of Bias In Non-randomised Studies - of Interventions) tool.¹⁵⁹ This tool assess the risk of bias across seven domains:

- bias due to confounding,
- bias due to selection of participants,
- bias in classification of interventions,
- bias due to deviations from intended interventions,
- bias due to missing data,
- bias in measurement of outcomes, and
- bias in selection of the reported result.

Similar to RoB 2, it has a number of signaling questions which inform the domain-level judgement, with acceptable judgements including “low”, “moderate”, “serious” and “critical”. In the context of the tool, observational studies are assessed in reference to an idealised randomised controlled trial. Under this approach, the (rare) overall judgement of “low” indicates that the results should be considered equivalent to that produced by a randomised controlled trial.

While a risk-of-bias tool for non-randomised studies of exposures (NRSE) is currently under development,¹⁶³ it was insufficiently developed at the time the risk-of-bias assessments for this review were performed. Instead, I used a version of the ROBINS-I tool informed by the preliminary ROBINS-E tool (Risk of Bias In Non-randomised Studies – of Exposure), which I had previously applied in a published review.¹⁶⁴ The version had no signaling questions and so judgements, using the same four levels of risk of bias as ROBINS-I, were made at the domain level. The motivation for using this tool above other established tools such as the Newcastle-Ottawa scale (NOS)¹⁶⁵ was two-fold. In the first instance, as mentioned in the introduction to this section, using a domain-based tool has distinct advantages over better-developed checklist-type tools including the NOS. Additionally, using a domain-based tool for

non-randomised studies of exposures enabled better comparison with risk-of-bias assessments performed for the other study designs as part of this review.

Mendelian randomisation studies

At present, no formalised risk-of-bias assessment tool for Mendelian randomisation studies is available. Assessment of the risk of bias in Mendelian randomisation studies was informed by the approach used in a previous systematic review of Mendelian randomisation,¹⁶⁶ as identified by a review of risk-of-bias assessments in systematic reviews of Mendelian randomisation studies.¹⁶⁷ Advanced results from this review were obtained from contact with the review authors. A copy of this tool is available in Appendix A.3.6, but in summary, results were assessed for bias arising from weak instruments, genetic and other confounding, pleiotropy and population stratification. Acceptable judgements for each of the five domains in the tool were “low”, “moderate” and “high” risk of bias.

Risk of bias due to missing evidence

In addition to assessing the risk of bias within each result contributing to a synthesis, I also assessed risk of bias due to missing evidence at the analysis level. This assessment examines evidence missing due to selective non-reporting as distinct from the selective reporting of a single result from multiple planned analyses. The assessment was performed using the forthcoming RoB-ME (Risk of Bias due to Missing Evidence in a synthesis) tool.¹⁶⁸ The tool is in development stages, and as part of this review, I piloted the tool and provided feedback to the developers.

3.2.9 Analysis methods

An initial qualitative synthesis of evidence was performed, summarising the data extracted from studies stratified by study design. Where individual studies were deemed comparable, they were incorporated into a quantitative analysis or “meta-analysis”.

Analysis overview

Results were not combined across different study designs (i.e. RCTs were not combined in a meta-analysis with results from observational studies). The summary effect estimates produced across individual study designs are discussed, but are compared and contrasted more fully as part of the triangulation exercise presented in Chapter 8. Similarly, analyses are presented separately for each dementia outcome of interest (all-cause dementia, Alzheimer’s disease, vascular dementia).

The range of effect measures presented by studies (odds ratios, risk ratios, hazard ratios, etc.) are not directly interchangeable in the context of systematic review. As such, different reported effect estimates can be one potential problem that precludes a meta-analysis of all studies.¹⁶⁹ However if the outcome is rare, as is the case for dementia outcomes, the estimated odds, risk and hazard ratios will approximate each other.¹⁷⁰ As such, I did not stratify the analyses by reported effect measure.

Random-effects meta-analysis

I used a random-effects model in all meta-analyses.^{171,172} Random-effects meta-analysis does not assume one true underlying effect, but rather allows for a distribution of true effects with variance τ^2 . The result in the i th study is denoted as y_i . The weight assigned to this result is then given as the inverse of the variance of that result (v_i) plus the estimate of between-result variance (τ^2):

$$w_i = \frac{1}{v_i + \tau^2} \quad (3.5)$$

Once the weights are calculated for each result, the overall estimate (\hat{y}) and variance ($Var(\hat{y})$) can be estimated:

$$\hat{y} = \frac{\sum y_i w_i}{\sum w_i} \quad (3.6)$$

$$Var(\hat{y}) = \frac{1}{\sum w_i} \quad (3.7)$$

Results included in each meta-analysis were stratified into subgroups on the basis of the overall risk of bias judgement, and summary estimates for each subgroup, in addition to an overall effect estimate, are displayed in each forest plot. Additional descriptive statistics are presented, while prediction intervals are shown as a dotted line banding the overall effect estimate. Finally, where at least 10 results are available, a test of subgroup differences between studies at different levels of risk of bias was performed (see subsequent Section 3.2.9).¹⁷³ All models were implemented using the `metafor` R package.¹⁷⁴

Dose-response analyses

Several of the included studies presented data on multiple categories of lipid levels, but provided an overall effect estimate based on a comparison of only two of these categories (e.g. for example, highest vs lowest quartile). While this allows for easy interpretation of the resulting effect estimate, it ignores any potential non-linear relationships between the exposure and outcome, in addition to discarding useful information contained in the interim groups. In order to address this limitation,

3.2 - Methods

I performed a dose-response meta-analysis in those studies reporting more than two categories of lipid levels.

Studies were excluded from this analysis if the number of categories was less than three or if the necessary information for synthesis (cut-off points, number of participants and number of events per category) was not available. When the highest reported category was open ended (e.g LDL-c ≥ 200 mg/dL), I calculated the category midpoint by assuming the width of the highest category was the same as the one immediately below it. Similarly, when the lowest category was open-ended (e.g LDL-c ≤ 100 mg/dL), I set the lower boundary for this category to zero (though this is unlikely to occur naturally).

I took a two-stage dose-response meta-analysis approach, where study-specific trends are estimated before being pooled across studies.¹⁷⁵ To estimate within-study trends, a restricted cubic spline model was fitted within each study. This approach allows for a non-linear relationship between lipid levels and dementia, for example a U or J-shaped relationship, where low and high levels of a lipid fraction can have different effects versus the “normal” reference dose.^{176,177} Reference doses were defined *a priori* as the cut-off of the “Normal”/“Optimal” categories for each fraction, as detailed in Table 1.2. Under this approach, the reference dose was defined as 200 mg/dL for total cholesterol, 100 mg/dL for LDL-c, 40 mg/dL for HDL-c, and 150 mg/dL for triglycerides.

Restricted cubic spline models use “knots” to separate the data into subsets or “windows”, in which the trend is modelled. Further restrictions are then imposed, in that the curves in each window must join up “smoothly” at the knot location, and that the trend is assumed to be linear before the first knot and after the last knot.¹⁷⁸ The locations of the knots in the model were identified using fixed percentiles (5th, 50th, 95th) of the exposure data.¹⁷⁶ Once the within-study trends had been estimated, they were combined in a multivariate random-effects meta-analysis using the restricted maximum likelihood (REML) method.^{177,179,180} All dose-response analyses were implemented using the `dosresmeta` R package.¹⁸¹

Additional analyses

Where there was evidence of heterogeneity between results included in a meta-analysis, I investigated this further using meta-regression against reported characteristics. *A priori*, I was interested in the effect that the age at baseline, sex and risk-of-bias judgement had on the results. Syntheses with greater than 10 results were assessed for heterogeneity across these covariates.¹⁷³

Finally, I investigated the potential for small study effects, which may be caused by publication bias, both visually using funnel plots and formally using Egger's regression test.¹⁸²

Visualisation of results

Evidence maps are useful way to explore the distribution of research cohorts included in a systematic review.¹⁸³ As part of the initial descriptive synthesis, the location of each individual study contributing to the evidence base was visualised on a world map.

One of the limitations of current risk-of-bias assessment practice in systematic reviews is that they are often divorced from the results to which they refer, and are infrequently incorporated into the analysis.^{184,185} In response to this criticism, I developed a new visualisation tool which was designed to allow for the production of “paired” forest plots - a risk-of-bias assessment is presented alongside it’s corresponding numerical result - as recommended by the ROB2 publication.¹⁶² In addition, the risk of bias due to missing evidence in each synthesis is shown beside the overall summary diamond. This tool was developed as an adjunct to this thesis to aid in creating standardised risk-of-bias figures.¹⁸⁶ A summary of this tool is contained in Appendix B.2, and all forest plots presented in this Chapter

were created using this tool. Unless otherwise stated in the figure caption, only a single effect measure (hazard ratio/odds ratio/etc) is represented in the forest plot.

Assessment of added value of including preprints

Preprints are considered a valuable evidence source within this thesis (see Introduction, Section 1.5.1). As an adjunct analysis to this review, meta-analyses including a preprint result were re-analysed using a fixed effect model. The weight assigned to the preprinted result was extracted and used to assess the additional evidential value provided by the preprint. In addition, in meta-analysis including at least one preprinted result, I assessed the impact of including the preprinted results on the summary estimate using a leave-one-out approach.

Additionally, I performed a follow-up analysis, allowing for a two-year lag, to investigate whether identified preprints had been subsequently published (in which case preprints provided a snapshot into the future, and a systematic review update would capture the published version of the preprint) or not (in which case preprints provide a distinct evidence source to conventional bibliographic databases).

3.3 Summary

- In this chapter, I have presented the methods underpinning a comprehensive systematic review of the existing literature on the association of dementia incidence with blood lipids and lipid-regulating agents.
- I have detailed how this review will find, extract, critically assess and synthesize the results of existing studies. Additionally, I have highlighted how this review will examine the impact of including preprinted results, as enabled by the research tool presented in the previous chapter.

3.3 - Summary

- In the following chapter, I present the results of this review, and detail how the identified evidence is used throughout the remainder of the thesis.

Hold

— Hold

4

Systematic review of existing evidence on the association between blood lipids and dementia outcomes: Results

Contents

Lay summary	63
4.1 Introduction	64
4.2 Search results	64
4.3 Primary analyses	77
4.4 Additional analyses	90
4.5 Discussion	93
4.6 Summary	104

Lay summary

Using the methods outlined in the previous chapter, here I present the results of a systematic review of primary studies that have examined the relationship between the levels of blood lipids (such as cholesterol and triglycerides) and dementia outcomes. In addition, it examined the relationship between treatments that change blood lipid levels, such as statins, and dementia outcomes.

My review included 81 primary studies. I found that statins appear to reduce the risk of all-cause dementia and Alzheimer's disease, but were not associated with risk of vascular dementia. Lipid levels were not associated with any dementia outcome. However, problems with the methods used in some studies included in the review meant that I was less confident in their results. In addition, I found that several results were missing from my analysis as authors decided not to report them.

4.1 Introduction

Following the methods outlined in the previous chapter, I present the results of the comprehensive systematic review here. The use of the evidence identified by the review in future chapters is discussed. Finally, I make recommendations for future methodological work around the inclusion of preprints and Mendelian randomisation studies in evidence synthesis exercises.

4.2 Search results

4.2.1 Initial search and validation of search filters

The database search identified 23,447 records, of which 7,338 were duplicated records. To ensure the accuracy of the study design filters, a random sample of 500 records removed by the filters were screened, and no eligible records were identified. As expected, many of those excluded by the filters were commentaries/educational articles, or described basic science studies.

4.2.2 Screening results

Following de-duplication, the titles and abstracts of 16,109 records were assessed for eligibility. Of these, 387 were deemed potentially eligible. The full text records for these records were then accessed and screened.

The PRISMA flow diagram presented in Figure 4.1 illustrates the movement of articles through the review. To highlight the contribution of preprint archives to the review, the flow diagram delineates between those records captured through databases searches (presented on the left of the diagram) and those captured by the search tool described previously (presented in grey on the right of the diagram).

Common reasons for exclusion at the full-text stage included studies reporting on ineligible exposures (Number of studies (N) = 70 ; most commonly an ineligible lipid fraction) or outcomes (N = 67; e.g. change in cognitive scores), or the use of an ineligible study design (N = 56).

4.2 - Search results




Figure 4.1: PRISMA flow diagram illustrating how records moved through the systematic review process. The different contributions of standard bibliographic databases and preprint servers to the review are indicated.

4.2.3 Validation of screening

For the assessment of the intra-/inter-rater reliability, the estimated values of Gwet's agreement coefficient, $AC1$, were interpreted against the categories presented in Table 3.2. For the inter-rater reliability, agreement was “almost perfect” ($AC1 = 0.97$, $kappa = 0.54$, Table 4.1). Similarly for intra-rater reliability, agreement was “almost perfect” ($AC1 = 0.99$, $kappa = 0.65$, Table 4.2). The discrepancy between the $AC1$ and $kappa$ coefficients illustrates the sensitivity of $kappa$ to imbalanced marginals, caused in this sample by a large imbalance towards exclusion.¹⁸⁷

Table 4.1: Inter-rater agreement on a subset of records, indicating high accuracy.

		Initial screening decision		
		Exclude	Include	Total
Second reviewer decision	Exclude	1244	9	1253
	Include	26	22	48
	Total	1270	31	1301

Table 4.2: Intra-rater agreement on subset of records, indicating high consistency.

		Initial screening decision		
		Exclude	Include	Total
Same reviewer decision (with 3 month lag)	Exclude	1266	14	1280
	Include	4	17	21
	Total	1270	31	1301

Those records which were excluded in the initial screening, but were included by the second reviewer ($N = 26$, Table 4.1) were investigated. This discrepancy between the two reviewers was explained in all cases by differing interpretations of the inclusion criteria, most commonly around the definition of cognitive decline versus dementia and of eligible lipids fractions.

4.2.4 Characteristics of included studies

Following full-text screening, 81 unique studies (described across 95 reports) met the criteria for inclusion in this review.^{53–62,67,71,72,74,188–254} Table 4.3 presents a summary of the characteristics of each study.

The majority of included studies described non-randomised analyses, with the only two included randomised controlled trials (the Heart Protection Study/British Heart Foundation trial⁶⁷ and the JUPITER trial¹⁸⁸) both reporting on the effect of statin use on incidence of all-cause dementia in older adults. A similarly small number of Mendelian randomisation studies were identified, several of which employed a two-sample approach using summary statistics from the same published genome wide association studies (GWAS), leading to complications in the synthesis of these estimates due to the potential for double-counting (see Section 4.3.2).

Of the 31 non-randomised studies examining treatments that modify lipid levels, all examined statin use (N = 31; 100%), while a small number also reported on other non-statins agents such as fibrates (N = 2; 6.5%). In the 43 non-randomised studies of exposure, hypercholesterolemia (N = 19; 44.2%) and total cholesterol levels (N = 21; 48.8%) were the most frequently reported risk factors.

In terms of outcomes, the vast majority of studies examined either all-cause dementia (N = 59; 72.8%) or Alzheimer's disease (N = 51; 63%), with only a small proportion examining vascular dementia (N = 15; 18.5%). Some other outcome classifications such as vascular-component or mixed dementia were also investigated, but were much rarer.

Three included reports were preprinted (denoted in the Table 4.3 using an asterisk),^{253,255,256} one of which had subsequently been published and was captured by the primary literature search.²⁵⁴ All three included preprints were obtained from the bioRxiv preprint server and all described a Mendelian randomisation analysis.

4.2 - Search results

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterik.

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Randomised controlled trials							
HPS 2002 ⁶⁷	United Kingdom	20536	>70	24.8	Simvastatin	Dementia	NR
JUPITER 2009 ¹⁸⁸	Multiple	17902	66 (median) 60-71 (range)	38	Rosuvastatin	Dementia	NR
Non-randomised studies of interventions							
Ancelin 2012 ¹⁸⁹	France	7056	NR	67	Fibrate; Statin	Dementia; AD	DSM-IV; NINCDS-ADRDA
Arvanitakis 2008 ¹⁹⁰	United States	929	74.9 (NR)	68.7	Statin	AD	Consortium to Establish a Registry for Alzheimer's Disease (CERAD)
Bettermann 2012 ¹⁹¹	United States	3069	78.6 (3.3)	46.2	Non-statin LRA ; Statin	Dementia; AD; Vascular component	Consensus panel - criteria not reported
Beydoun 2011 ¹⁹²	United States	1604	57.6 (18.4)	38.5	Statin	Dementia	DSM-III-R
Chao 2015 ¹⁹³	Taiwan	256265	73.2 (7.4)	50.3	Statin	Non-vascular dementia	ICD-9
Chen 2014 ¹⁹⁴	Taiwan	18100	67 (8.6)	47.9	Statin	Dementia; AD; Non-AD	ICD-9
Chitnis 2015 ¹⁹⁵	United States	8062	74.47 (9.21)	53.04	Statin	Dementia	ICD-9
Chou 2014 ¹⁹⁶	Taiwan	33398	>60	53.9	Statin	Dementia; AD; VaD; Non-vascular dementia	ICD-9
Chuang 2015 ¹⁹⁷	Taiwan	123300	54 (13)	49.1	Statin	Dementia	ICD-9
Cramer 2008 ¹⁹⁸	United States	1674	70 (6.8)	58	Statin	Dementia/CIND	DSM-IV
Gnjidic 2016 ¹⁹⁹	Sweden	2056	>60	NR	Statin	Dementia	DSM-IV
Haag 2009 ²⁰⁰	Netherlands	6992	69.4 (9.1)	60	Non-statin LRA ; Statin	AD	NINCDS-ADRDA

4.2 - Search results

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterisk. (*continued*)

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Hendrie 2015 ²⁰¹	United States	974	76.6 (4.9)	69.7	Statin	Dementia; AD	DSM-IV; Consortium to Establish a Registry for Alzheimer's Disease (CERAD) EHR codelist
Hippisley-Cox 2010 ²⁰²	United Kingdom	2004692	46 (14)	51	Statin	Dementia	
Jick 2000 ²⁰³	United Kingdom	1364	50-89	61	Non-statin LRA ; Statin	Dementia	EHR codelist
Li 2004 ²⁰⁵	United States	2356	75.1 (6.1)	59.8	Non-statin LRA; Statin	Dementia; AD	DSM-IV; NINCDS-ADRDA
Li 2010 ²⁰⁴	United States	3392	75 (6.2)	59	Statin	AD	NINCDS-ADRDA
Liao 2013 ²⁰⁶	NR	5221	NR	NR	Statin	Dementia	NR
Liu 2019 ²⁰⁷	Taiwan	2012	74 (7.5)	NR	Statin	Dementia	ICD
Pan 2018 ²⁰⁸	Taiwan	14807	65 (13)	43	Statin	Dementia	ICD-9
Parikh 2011 ²⁰⁹	United States	377838	75.53 (6.07)	2	Statin	Dementia	ICD-9
Rea 2005 ²¹⁰	United States	2798	NR	NR	Non-statin LRA ; Statin	Dementia; AD; Mixed; VaD	NINCDS; NINCDS-ADRDA; Combination; State of California Alzheimer's Disease Diagnostic and Treatment Centers
Redelmeier 2019 ²¹¹	Canada	28815	76 (NR)	61.3	Statin	Dementia	ICD-9
Reitz 2004 ²¹²	United States	1168	78.4 (6.2)	68.3	Statin	VaD; AD	Cohort criteria; NINCDS-ADRDA
Smeeth 2009 ²¹³	United Kingdom	729529	50 (NR)	40-81	Statin	Dementia; AD; Non-AD	EHR codelist
Solomon 2010 ²¹⁴	Finland	17597	68 (5.8)	57	Statin	Dementia	EHR codelist

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterisk. (*continued*)

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Sparks 2008 ²¹⁵	United States	2068	75 (3.8)	54	Statin	AD	NINCDS-ADRDA
Szwast 2007 ²¹⁶	United States	1416	77.3 (5.3)	69.3	Statin	Dementia	DSM-IV
Yang 2015 ²¹⁷	Taiwan	45973	82 (5.3)	48	Fibrate; LRA (exlc. statin + fibrates); Statin	Dementia	ICD-9
Zamrini 2004 ²¹⁸	United States	3397	73 (NR)	0	Statin	AD	ICD-9
Zandi 2005 ²¹⁹	United States	3308	NR	NR	Non-statin LRA; Statin	Dementia; AD	DSM-III-R; NINCDS-ADRDA
Non-randomised studies of exposures							
Ancelin 2013 ²²⁰	France	7053	74 (5.3)	61.1	Hypercholesterolemia	AD; Dementia	NINCDS-ADRDA; DSM-IV
Batty 2014 ²²¹	United Kingdom	103764	47.3 (18.1)	55	Non-HDL-c; Hypercholesterolemia	Dementia	ICD
Benn 2017 ⁷⁴	NR	111194	56 (median) 46-66 (range)	55	LDL-c	AD; VaD; Dementia	ICD; ICD-10
Beydoun 2011 ¹⁹²	United States	1604	57.6 (18.4)	38.5	TC	Dementia	DSM-III-R
Bruce 2017 ²²²	Australia	217	63.6 (8.4)	45.6	HDL-c; TC; TG	Dementia	NR
Chiang 2007 ²²³	Taiwan	785	58 (7.4)	41.4	TC; TG	Dementia; AD; VaD	ICD-9; NR
Dodge 2011 ²²⁴	United States	822	71.6 (4.7)	64.4	Hypercholesterolemia	Dementia; AD	DSM-III-R; Consortium to Establish a Registry for Alzheimer's Disease (CERAD)
Forti 2010 ²²⁵	Italy	749	73 (6.1)	53	Hypercholesterolemia; TG	Dementia; AD; VaD	DSM-IV; NINCDS-ADRDA; NINCDS-AIREN
Gottesman 2017 ²²⁶	United States	15407	54.2 (5.8)	55	TC	Dementia	Combination
Gustafson 2012 ²²⁷	Sweden	NR	NR	100	TC	AD	NR

4.2 - Search results

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterisk. (*continued*)

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Hayden 2006 ²²⁸	United States	3308	74.0 (6.4)	58.2	Hypercholesterolemia	Dementia; AD; VaD	DSM-III-R; NINCDS-ADRDA;
Kimm 2011 ²²⁹	South Korea	848505	53 (9.3)	42.2	TC	AD; VaD; Dementia	ICD-10
Kivipelto 2001 ²³⁰	Finland	1499	50.4 (6.0)	62	Hypercholesterolemia	AD	NINCDS-ADRDA
Kivipelto 2005 ⁵³	Finland	1449	50.6 (6.0)	62	Hypercholesterolemia	Dementia	DSM-IV
Kuo 2015 ²³¹	Taiwan	67066	62.1(11.4)	48.4	Hypercholesterolemia	Dementia	ICD-9
Li 2005 ⁵⁷	United States	2141	74.9 (5.9)	60.5	HDL-c; TC	Dementia; AD	DSM-IV; NINCDS-ADRDA
Mainous 2005 ⁵⁸	United States	6558	NR	NR	Hypercholesterolemia	Dementia; AD	ICD-9
Mielke 2005 ⁵⁹	Sweden	382	NR	70	TC; TCG	Dementia	DSM-III-R
Mielke 2010 ⁶¹	France	1460	38-60 (range)	100	Hypercholesterolemia; TC	Dementia; AD	DSM-III-R; NINCDS-ADRDA
Mielke 2012 ²³²	United States	99	74 (2.5)	100	HDL-c; TC; TG	Dementia; AD	DSM-IV; NINCDS-ADRDA
Muller 2007 ²³³	United States	542	NR	NR	HDL-c; TG	Dementia; AD	DSM-IV; NINCDS-ADRDA
Noale 2013 ²³⁴	Italy	5632	71.3(5.3)	56.3	Hypercholesterolemia; TG	Dementia	DSM-III-R
Notkola 1998 ²³⁵	Finland	444	40-59 (range)	0	Hypercholesterolemia	AD	Combination
Peters 2009 ²³⁶	Multiple	3336	>80	60.4	HDL-c; TC	Dementia	DSM-IV
Raffaitin 2009 ²³⁷	France	7087	73.4 (4.9)	61	Hypercholesterolemia; TG	Dementia; AD; VaD	DSM-IV; NINCDS-ADRDA; Combination
Rantanen 2017 ²³⁸	Finland	3309	42 (median) 39-46 (range)	0	TC; Hypercholesterolemia	Dementia; AD; VaD	NR
Reitz 2004 ²¹²	United States	1168	78.4 (6.2)	68.3	HDL-c; LDL-c; Non-HDL-c; TC; TG	VaD; AD	Cohort criteria; NINCDS-ADRDA

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterisk. (*continued*)

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Reitz 2010 ⁶²	United States	1130	75.7(6.3)	65.7	HDL-c; LDL-c; TC	AD	NINCDS-ADRDA
Ronnemaa 2011 ²³⁹	United States	2268	49.6 (0.6)	0	Hypercholesterolemia	AD; VaD; Dementia	NINCDS-ADRDA; ADDTG; DSM-IV
Schilling 2017 ⁵⁴	France	9294	73.8 (5.3)	61	HDL-c; LDL-c; TC; TG	Dementia; AD; Mixed	DSM-IV; NINCDS-AIREN
Solomon 2007 ²⁴⁰	Finland	1449	50.4 (6.0)	62.1	Hypercholesterolemia	Dementia	NR
Solomon 2009 ⁵⁵	United States	9844	43 (1.7)	54	TC	AD; VaD	ICD-9
Strand 2013 ²⁴¹	Norway	48793	42.6 (4.3)	49	TC	Dementia; AD	ICD
Su 2017 ²⁴²	United Kingdom	212085	NR	NR	NR	NR	NR
Svensson 2019 ²⁴³	Japan	781	54.1 (5.6)	NR	HDL-c; Hypercholesterolemia	Dementia	DSM-IV
Tan 2003 ⁶⁰	United States	1026	76.1 (5.3)	63	HDL-C; TC	AD	NINCDS-ADRDA
Tynkkynen 2016 ²⁴⁴	Finland	13725	48.4 (13.3)	51.6	HDL-c	Dementia; AD	ICD-10
Tynkkynen 2018 ²⁴⁵	Multiple	22623	57 (9.2)	47	HDL-c; LDL-c; TC; TG	Dementia; AD	ICD-10
Wang 2012 ²⁴⁶	Taiwan	1230400	60 (13)	52	Hypercholesterolemia	AD	ICD-9
Whitmer 2005 ⁵⁶	United States	8845	68 (2.6)	53.7	Hypercholesterolemia	Dementia	ICD-9
Yamada 2009 ²⁴⁷	Japan	1637	>60	100	NR	NR	NR
Yoshitake 1995 ²⁴⁸	Japan	828	74 (5.9)	59.5	HDL-c; LDL-c; TC; TG	VaD; AD	NINCDS-AIREN; NINCDS-ADRDA
Zimmetbaum 1992 ²⁴⁹	United States	350	79 (median) 75-85 (range)	64.5	HDL-c; LDL-c; TC; TG	Dementia	DSM-III-R
Mendelian randomisation studies							
Andrews 2019 ^{250,255 *}	Multiple	54162	NR	NR	HDL-c; LDL-c; TC; TG	AD	NR

Table 4.3: Characteristics of included studies, stratified by study design. Note that three studies reported on multiple analytical designs within a single study report (Beydoun 2011,¹⁹² Reitz 2010,²¹² and Benn 2017⁷⁴), and these have been duplicated across the relevant sub-sections. Preprinted studies are denoted using an asterisk. (*continued*)

Study	Location	N	Age at baseline	Female (%)	Exposures	Outcomes	Diagnostic criteria
Benn 2017 ⁷⁴	Multiple	111194; 54162	NR	NR	HMGCR; LDL-c; PCSK-9	AD; VaD; Dementia	NR
Burgess 2017 ²⁵¹	Multiple	21165	NR	NR	HDL-c; LDL-c; TG	AD	NR
Larsson 2017 ⁷¹	Multiple	54162	NR	NR	LDL-c	AD	NR
Mukherjee 2013 ²⁵²	Multiple	54162	NR	NR	HDL-c; LDL-c; TG	AD	NR
Ostergaard 2015 ⁷²	Multiple	54162	NR	NR	HDL-c; LDL-c; TC; TG	AD	NR
So 2017 ²⁵³ *	Multiple	54162	NR	NR	HMGCR	AD	NR
Zhu 2018 ^{254,257} *	Multiple	54162	NR	NR	HDL-c; LDL-c; TG	AD	NR

* Denotes preprinted study.

Abbreviations: AD - Alzheimer's disease; DSM - Diagnostic and Statistical Manual (Roman numerals indicate edition); EHR - Electronic code list; ICD - International Classification of Disease (numbers indicate edition); HDL-c - high density lipoprotein cholesterol; LDL-c - low density lipoprotein cholesterol; NINCDS-ADRDA - National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association; NINCDS-AIREN - National Institute of Neurological Disorders and Stroke and Association Internationale pour la Recherche et l'Enseignement en Neurosciences; NR - Not reported; TC - total cholesterol; TG - triglycerides; VaD - vascular dementia.

As illustrated in Figure 4.2, the majority of reports described studies conducted in high-income countries, with the most highly represented region being North America. Of interest, several of the included studies were conducted in Taiwan ($N = 10$; 12.35%), all but one of which made use of the Taiwan National Health Insurance database.



Figure 4.2: Geographical distribution of study cohorts

Finally, there were several eligible studies reported as conference abstracts that did not present numerical results. These reports were included in the analysis to enable assessment of risk of bias due to missing evidence (see Section 3.2.8).

4.3 Primary analyses

In the following sections, analyses are grouped first by outcome (all-cause dementia, Alzheimer's disease and vascular dementia) and are then further stratified by exposure category (lipids, lipid regulating agents).

4.3.1 All-cause dementia

Lipids

Across all outcomes, lipid levels were categorised in a number of ways. The most common categorisation was hypercholesterolemia at baseline, defined most frequently as a total cholesterol measurement of greater than 6.5 mmol/L.

Eleven studies reported on the association of hypercholesterolemia with all-cause dementia and provided weak evidence for an effect (HR: 1.11, 95%CI: 0.98-1.25, Figure 4.3).

4.3 - Primary analyses




Figure 4.3: Random-effects meta-analysis of non-randomised studies examining the effect of hypercholesterolemia on all-cause dementia

Several studies analysed individual lipid fractions by estimating the risk of dementia per 1 standard deviation increase in that fraction (Figure 4.4). Weak evidence for an effect on all-cause dementia was found for total cholesterol ($N = 5$; HR: 0.97, 95%CI: 0.88-1.07), LDL-c ($N = 2$; HR: 0.97, 95%CI: 0.86-1.08), HDL-c ($N = 4$; HR: 1.05, 95%CI: 0.96-1.14) and triglycerides ($N = 3$; HR: 0.90, 95%CI: 0.74-1.09).

4.3 - Primary analyses



Figure 4.4: Random-effects meta-analysis of four lipid fractions (total cholesterol, HDL, LDL, and triglycerides) on all-cause dementia risk, standardised per 1-SD increase in the lipid fraction. Note that Tynkkynen *et al*²⁴⁵ reported results from multiple cohorts within a single study and the results for each cohort are presented separately here.

Finally, there were no identified Mendelian randomisation analysis examining the effect of lower lipid levels, as determined by any genetic instrument, on all-cause dementia risk.

Statins

The two randomised controlled trials provided weak evidence (OR: 1.07, 95%CI: 0.70-1.66) of an effect on statin use on all-cause dementia risk (Figure 4.5).




Figure 4.5: Random-effects meta-analysis of randomised controlled trials examining statins on all-cause dementia. D1-D5 and O refer to the five risk-of-bias domains and single overall judgement in the RoB2 tool, while the risk of bias due to missing evidence is shown beside the overall summary diamond in each forest plot presented above.

In contrast, a meta-analysis of 15 prospective observational studies provided some evidence of a protective effect of statins use on all-cause dementia risk (HR: 0.77, 95%CI: 0.69-0.87, Figure 4.6).

4.3 - Primary analyses




Figure 4.6: Random-effects meta-analysis of non-randomised studies examining the effect of statin use on all-cause dementia

Finally, a single Mendelian randomisation analysis was identified examining the effect of lowered LDL-c levels on the risk of all-cause dementia via genetic inhibition of the 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMGCR), emulating statin treatment (see Section 1.3.2 for more details of the statin mechanism of action). This analysis provided weak evidence of an effect (RR: 0.90, 95%CI: 0.29-2.81).

Fibrates

Two studies examined the effect of fibrate use on all-cause dementia and found weak evidence for an effect (HR: 0.89, 95%CI: 0.75-1.07, Figure 4.7).




Figure 4.7: Random-effects meta-analysis of non-randomised studies examining the effect of fibrate use on all-cause dementia

4.3.2 Alzheimer's disease

Lipids

Nine studies reported on the association of hypercholesterolemia with Alzheimer's disease and provided weak evidence for an effect (HR: 0.99, 95%CI: 0.78-1.25, Figure 4.8)

4.3 - Primary analyses




Figure 4.8: Random-effects meta-analysis of non-randomised studies examining the effect of hypercholesterolemia on Alzheimer's disease

Similarly to all-cause dementia, several studies analysed individual lipid fractions by estimating the risk of dementia per 1 standard deviation increase in that fraction (Figure 4.9). Weak evidence for an effect on all-cause dementia was found for total cholesterol ($N = 5$; HR: 1.00, 95%CI: 0.94-1.06), LDL-c ($N = 3$; HR: 1.06, 95%CI: 0.98-1.16), HDL-c ($N = 4$; HR: 0.99, 95%CI: 0.91-1.07) or triglycerides ($N = 3$; HR: 1.00, 95%CI: 0.84-1.18).

4.3 - Primary analyses




Figure 4.9: Random-effects meta-analysis of four lipid fractions (total cholesterol, HDL, LDL, and triglycerides) on Alzheimer's disease risk, standardised per 1-SD increase in the lipid fraction. Note that Tynkkynen *et al*²⁴⁵ reported results from multiple cohorts within a single study and the results for each cohort are presented separately here.

4.3 - Primary analyses

Finally, there were several identified Mendelian randomisation studies examining the effect of genetically lowered LDL-c on Alzheimer's disease risk (Figure 4.10). However, all of these studies took a two-sample approach, making use of summary statistics from the Global Lipid Genetics Consortium (GLGC) and International Genomics of Alzheimer's Project (IGAP) consortia. Synthesis of these results would result in a falsely precise estimate caused by multiple counting of the same participants, and as a result, no meta-analysis of these studies was performed.

However, across each analysis using varying number of SNPs, no evidence for an effect was observed (Figure 4.10).




Figure 4.10: Summary of duplication across Mendelian randomisation studies which used summary statistics from the Global Lipid Genetics Consortium (GLGC) and the International Genomics of Alzheimer's Project (IGAP). Note that the Alzheimer's Disease Genetics Consortium (ADGC) is a sub-cohort within IGAP.

Statins

There were no randomised trials of the use of statins or any other lipid regulating agents on Alzheimer's disease, though several observational studies reported on this outcome and provided evidence for a protective effect (N = 11; HR: 0.83, 95%CI: 0.69-0.99; Figure 4.11).

4.3 - Primary analyses




Figure 4.11: Random-effects meta-analysis of non-randomised studies examining the effect of statin use on Alzheimer's disease

Two Mendelian randomisation studies looked at lipid-lowering specifically as a result of HMGCR inhibition, mediated by SNPs in the gene (rs172338484 and rs12916).^{74,253} The first used a one sample approach (SNP-exposure and SNP-outcome associations are estimated using the same dataset) in a large Copenhagen-based cohort, while the second made use of summary level data obtained from the GLGC (SNP-exposure) and the IGAP (SNP-outcome). Meta-analysis of these estimates provided weak evidence of an effect (RR: 0.76, 95%CI: 0.51-1.14, Figure 4.12).

4.3 - Primary analyses




Figure 4.12: Random-effects meta-analysis of genetically lowered LDL-c via HMGCR inhibition on Alzheimer's disease

4.3.3 Vascular dementia

Lipids

Three studies reported on the association of hypercholesterolemia with vascular dementia and provided weak evidence for an effect (HR: 1.20, 95%CI: 1.00-1.44, Figure 4.13)




Figure 4.13: Random-effects meta-analysis of non-randomised studies examining the effect of hypercholesterolemia on vascular dementia

Few studies investigated the effect of individual lipid fractions. Only for total cholesterol was there greater than one result reported (Figure 4.14), and a meta-analysis of these found weak evidence for an effect (N = 2; HR: 1.05, 95%CI:

4.3 - Primary analyses

0.79-1.41). A single study provided evidence on the other three fractions,²⁴⁸ and similarly found minimal evidence of an effect LDL-c (HR: 1.12, 95%CI: 0.83-1.51), HDL-c (HR: 0.83, 95%CI: 0.60-1.14) or triglycerides (HR: 1.00, 95%CI: 0.75-1.34).




Figure 4.14: Random-effects meta-analysis of four lipid fractions (total cholesterol, HDL, LDL, and triglycerides) on Alzheimer's disease risk, standardised per 1-SD increase in the lipid fraction. Note that Tynkkyen *et al*²⁴⁵ reported results from multiple cohorts within a single study and the results for each cohort are presented separately here.

Finally, no Mendelian randomisation analyses examining the effect of genetically determined lipid levels on vascular dementia risk were identified.

Statins

As noted in Section 4.2.4 above, there was substantially less literature available on the association of my risk factors of interest and vascular dementia. There were no available randomised trials for this outcome. Three prospective cohort studies examined statin use and vascular dementia, though meta-analysis of these studies provided weak evidence for an effect (N = 3; HR: 0.99, 95%CI: 0.79-1.25; Figure 4.15).




Figure 4.15: Random-effects meta-analysis of non-randomised studies examining effect of statin use on vascular dementia

A single Mendelian randomisation analysis was identified that examined the effect of genetically lowered LDL-c levels via HMGCR inhibition on the risk of vascular dementia, which provided some evidence for an effect (RR: 0.44, 95%CI: 0.21-0.91).

4.3.4 Risk of bias

As shown in the forest plots in the above sections, the risk-of-bias assessments are presented alongside their corresponding numerical result. This section presents only a brief summary of the biases observed in each study design, as a detailed discussion of the sources and directions of bias in each result is presented in Chapter 7.

For the two randomised controlled trials, both were judged to be at low risk of bias. In contrast, many of the non-randomised studies of statin use were at serious risk of bias primarily due to poor controlling for confounding, immortal time bias, and missing outcome data. Similarly, non-randomised studies of exposures suffered from incomplete adjustment for potentially important confounders, and concerns over the selection of the reported result from among several analyses (e.g. examination of lipids as a binary or continuous variable). Finally, bias was introduced into Mendelian randomisation studies via the potential for horizontal pleiotropy and population stratification.⁷⁰

4.4 - Additional analyses

Following best practice, any result judged to be at critical risk of bias should be excluded from any quantitative analysis.¹⁵⁹ Three observational studies were excluded on this basis, predominantly due to a lack of adjustment for any potentially important confounders (i.e. the study reported unadjusted estimates).^{58,231,235}

The risk of bias due to missing evidence in each synthesis is shown beside the overall summary diamond in each forest plot presented above. For randomised controlled trials and non-randomised studies of interventions, the risk of bias due to missing evidence was assessed to be minimal. However, there was substantial evidence that results were selectively reported in studies examining the effect of lipid fractions on dementia outcomes (Figures 4.4, 4.9 and 4.14).

4.4 Additional analyses

4.4.1 Dose response meta-analysis of lipid levels

There were 13 studies initially considered eligible for the dose-response meta-analysis, as they appear to provide data on risk of dementia outcomes across several categories of lipid exposure. However, following data extraction, five were excluded as they did not report the relevant information needed (most commonly, the cut-off measures for each category).

Across the remaining eight studies, a sufficient number of results ($n \geq 3$) were identified only for the total cholesterol-Alzheimer's, LDL-Alzheimer's, and total cholesterol-dementia strata. This analysis provided weak evidence for a non-linear effect of lipid levels on dementia outcomes, and Figure 4.16 illustrates this for the total cholesterol-Alzheimer's strata. Similar figures for the other analysed lipid-outcome strata are presented in Appendix A.4.1.




Figure 4.16: Dose-response meta-analysis of total cholesterol on Alzheimer's disease

4.4.2 Heterogeneity and publication bias

Investigation of potential sources of heterogeneity was complicated by two factors. In the first instance, a minority of meta-analysis (4 out of 18, 22%) included more than 10 results, the recommended minimum required for meta-regression. Secondly, poor reporting of study characteristics of interest including education level and baseline cognitive ability precluded the use of several results in a meta-regression analysis.

Age and sex were assessed as potential causes of heterogeneity in the meta-analyses of statin use and hypercholesterolemia on all cause dementia and Alzheimer's disease, but I found weak evidence for variation in the observed effect estimates due to these factors.

Similarly, assessment of small-study effects, for which publication bias could be one

potential reason, was hindered by the relatively small number of results included in a given meta-analysis. However, all analyses assessed provided weak evidence of small-study effects.

4.4.3 Added evidential value of including preprints

As shown in the PRISMA flow diagram (Figure 4.1), the number of hits returned by preprint searching was not substantial (*bioRxiv* = 256, *medRxiv* = 0). From these hits, three preprinted reports of eligible studies were included in the review, of which two described unique studies not captured by the main search.^{253,255}

One preprint (So *et al*²⁵³) provided additional evidential value in a single meta-analysis (5.6% of the 18 meta-analyses performed in this chapter). This meta-analysis of Mendelian randomisation studies examined the effects of lipid-lowering via HMGCR mutations on incidence of Alzheimer's disease (Figure 4.12). To assess the evidence added to the meta-analysis through inclusion of the preprint, the data was re-analysed using a fixed-effect model. Examination of the weight assigned to each result in the analysis illustrates that a large proportion (78%) of the weight is given to the preprinted result. However, the inclusion of the preprinted study did not have a substantial impact on the result of the meta-analysis (RR: 0.76, 95%CI: 0.51-1.14) compared with that reported by the single published study (RR: 0.59, 95%CI: 0.25-1.39), other than a slight increase in precision.

The other two preprints identified^{255,256} reported on the effect of LDL-c on Alzheimer's disease using data from GLCC and IGAP consortia. As illustrated in Figure 4.10, these consortia were previously analysed in several published reports, and so these preprints did not add to new information to the evidence base.

For one of the three preprints identified, the published version of the preprint was captured by the main search. Investigation of the publication status of the two unpublished preprints found that, allowing for a two-year lag (i.e. up to July 2021), only one had been subsequently published.^{250,255}

4.5 Discussion

This review presented a summary of the available evidence on the association between lipids, and treatments that modify lipids such as statins, and the subsequent risk of dementia. This discussion seeks to summarise the key findings in terms of literature sources and results as reported. A detailed comparison across the evidence sources, exposure measures and sources of bias reported here is presented as part of the triangulation exercise in Chapter 7.

4.5.1 Summary of findings

There was some indication of a protective effect of statins on all-cause and Alzheimer's disease dementia when looking solely at observational studies. This finding was not supported by evidence from the two available RCTs, or by studies that emulated statin treatment using a genetic proxy, suggesting that these findings may be a result of differing exposure windows (e.g. mid-life lipid lowering in non-randomised studies versus life-course lipid exposure in Mendelian randomisation studies and late-life lipid reduction in RCTs) or alternatively due to biases within the non-randomised studies.

Across dementia outcomes, the majority of studies were non-randomised studies of lipids or lipid regulating agents. This distribution of evidence between analytical designs is to be expected. Randomised controlled trials of dementia are particularly challenging, as the follow-up made necessary by the long latent period of the condition makes trials logistically difficult and financially expensive. Similarly, Mendelian randomisation is a comparatively new study design, and studies employing it in relation to dementia outcomes only appear in the evidence base in recent years, as illustrated in Figure 4.17. This recent increase is likely driven by the availability

4.5 - Discussion

of summary genome wide association studies (GWAS) that form the basis of the two-sample Mendelian randomisation approach.




Figure 4.17: Included study designs for any dementia outcome by year of publication

A central finding of this review is the absence of studies examining vascular dementia as an outcome, most noticeable when comparing the evidence base for statins in dementia/Alzheimer's (Figures 4.6 & 4.11) with that available for vascular dementia (Figure 4.15). This is particularly interesting given that lipids and statins are strongly related to the prevention of vascular disease. A potential explanation for this observation may be publication bias or the "file-drawer effect",⁸⁵ though there was weak evidence of a small-study effects for this outcome (of which publication bias is one potential cause).¹⁸² Similarly, only one Mendelian randomisation study examined this outcome, likely due to the absence (until recently) of vascular dementia GWAS which precludes a two-sample approach.

Of note, this review did not include the commonly cited PROSPER RCT, which examined the effect of pravastatin on cardiovascular disease risk and reported on cognitive outcomes as one of several secondary outcomes.²⁵⁸ While widely cited in relation to the effect of statins on dementia risk and included in the Cochrane

4.5 - Discussion

review of RCTs on this topic,⁶⁵ the trial reported solely on the change in a range of cognitive measures (MMSE, Stroop test, Picture-Word Learning test and others) over follow-up. Though a useful indicator of general cognitive decline, it is not equivalent to a dementia diagnosis using recognised criteria, as cognitive tests should feed into a broader diagnostic pathway (see Section 1.2.2). As such, this trial did not meet the inclusion criteria for this review.

Risk of bias across the individual results was generally high. The causes and expected impact of these biases on the results are discussed in more detail in Chapter 7. Of particular interest to this chapter, however, was the high risk of bias due to missing evidence observed for observational studies of lipid levels. In many cases, estimates were known to be missing from meta-analyses due to preferential reporting of significant results, leading to high risk of bias due to missing evidence. These missing estimates were most commonly identified through differing descriptions of the same analysis in conference abstract versus the final publication.^{247,259} In addition, some authors stated outright that non-significant results were not reported (e.g. “The other lipid variables not significantly associated with dementia and Alzheimer’s disease … were not reported in the table.”²²⁰). However, as all identified missing results are likely to be non-significant, they would not be expected to have a substantial impact on their respective meta-analysis (which provided weak evidence of an effect), other than increasing the precision of the summary estimate.

Finally in terms of generalisability, despite a large proportion of the included studies being conducted in the Western world (Figure 4.2), the applicability of the results to other populations is aided by the inclusion of several studies which made use of data from the Taiwan health insurance database.

4.5.2 Comparison with previous reviews

While conducting this review, I identified several previous systematic reviews of this topic.²⁶⁰⁻²⁶⁵ However, to my knowledge, this review is the first to use established

domain based assessments tools (for example, the RoB 2 tool for randomised controlled trials)¹⁶² to assess the risk of bias in included studies. The majority of the highly cited reviews on this topic either do not formally consider risk of bias in the observational studies they include,^{260,266} or used a non-domain-based assessment tool (e.g. the Newcastle-Ottawa Scale).²⁶³

I identified one previous review of Mendelian randomisation studies examining risk factors for Alzheimer's disease. However, this review was conducted prior to the majority of Mendelian randomisation studies included in this review being published. In addition this previous review extracted the results of analyses which did not account for SNPs in the *ApoE4* genetic region (see the following section for a discussion of the bias this introduces).

Despite these differences in time scales and methodology, the duplication of work across systematic reviews (including this one) is substantial. In retrospect, an alternative approach to conducting a further systematic review from scratch, known as an umbrella review or review-of-reviews,^{267,268} could have been employed. This study design uses other systematic reviews rather than primary studies as the unit of analysis. Furthermore, it would have enabled more efficient identification of relevant primary studies to which the methods which sets this review apart could have been applied.

4.5.3 Inclusion of Mendelian randomisation studies

One of the particular strengths of this review is the inclusion and critical assessment of Mendelian randomisation studies as a source of evidence.

Mendelian randomisation is a powerful analytical technique, using natural variation in participants' genomes to identify causal links between a genetically determined risk factor and an outcome, given that the three core assumptions detailed in Figure 4.18 are valid. These assumptions are namely that:

- the genetic variant associates with the risk factor of interest (relevance assumption);
- the variant-exposure association has no unmeasured confounders (independence assumption);
- the variants affect the outcome only through their effect on the risk factor of interest (exclusion restriction assumption).




Figure 4.18: Summary of the assumptions in Mendelian randomisation analyses: (I) *relevance* - the genetic variant associates with the risk factor of interest; (II) *independence* - the variant-exposure association has no unmeasured confounders; and (III) *exclusion restriction* - variants affect the outcome only through their effect on the risk factor of interest (i.e. there is no horizontal pleiotropy).

However, inclusion of Mendelian randomisation as an acceptable study design in this review was complicated by a number of factors. Firstly, this study design is relatively new, particularly when compared to randomised trials or cohort studies. Figure 4.17 demonstrates that Mendelian randomisation studies only begin to appear in the evidence base much later than NRSE/NRSI, likely due to the limited availability of large scale GWAS datasets needed for two-sample Mendelian randomisation analyses. As such, the process and tools for systematically assessing this study design are not as well developed. A key example of this is the absence of validated search filters for Mendelian randomisation studies. This limitation is further complicated by the varying terminology used to describe the method, particularly in the early

4.5 - Discussion

years of its application, which led to me including general terms for instrumental variable analyses in my search.

Additionally, there is currently no widely used risk-of-bias assessment tool for Mendelian randomisation studies. While a recent commentary provided a checklist for interpreting Mendelian randomisation studies, this guide includes reporting items in their quality checklist.⁷⁰ Reporting quality is an important consideration for Mendelian randomisation studies, as reflected by the recent release of the STROBE MR reporting guidelines,²⁶⁹ but it is a separate consideration to bias, as discussed in Section 3.2.8. Similarly, a previous review of Mendelian randomisation studies used the Q-Genie tool which was designed to assess the quality of GWAS included in a meta-analysis.²⁷⁰ While this tool assesses the underlying GWAS used, it does not address the additional methodological considerations of the Mendelian randomisation analysis itself. For this review, I used the best available author-devised tool, sourced from a recent review of systematic reviews of Mendelian randomisation studies.¹⁶⁷

As a further stumbling block, Mendelian randomisation lends itself to the analysis of multiple exposure-outcome comparisons as part of a single study, particularly when using a two-sample summary data design. This is particularly relevant to the consideration of bias due to missing evidence. As an example, through snowballing and other measures, I identified at least one relevant Mendelian randomisation study that had not been captured by the search strategy.⁷¹ This study examined lipid fractions as one of many risk factors for Alzheimer’s disease, though the search would not have been expected to find it given the absence of any lipid-related keywords in the title or abstract. Studies examining multiple risk factors such as this can introduce bias into a systematic review, as it is commonly only those risk factors with a statistically significant result that are reported in the abstract and so are captured by a systematic search. These studies are described as “unknown unknown’s” in the context of the RoB-ME tool, and are considered to be particularly challenging (as opposed to an analysis that was insufficiently reported to be included in the statistical analysis, or the “known unknown’s”).

4.5 - Discussion

Useful future work to improve the methodology for inclusion of Mendelian randomisation studies in systematic reviews should involve the development of a validated search filter for this study design.^{271,272} Alternatively, in better-resourced reviews, a broader search (e.g. “risk factor” AND “dementia” AND “Mendelian randomisation”) followed by manual review of studies that examined multiple risk factors would be advisable. This was not feasible in the context of this review, given the large number of records to be screened even when using study design filters ($N = 16,109$). Additionally, the value of methods that support traditional bibliographic database searches, such as snowballing (forwards and backwards citation chasing) and communication with relevant topic experts, should not be underestimated. Finally, development of a risk-of-bias assessment tool for this study design by a panel of methodologists and analysts would be of substantial benefit.

One item of particular interest is the attenuation of any effects identified by Mendelian randomisation studies following the adjustment for/exclusion of genetic variation in the *ApoE4* gene region. As discussed in the introduction (see Section 1.4.1), an increasing number of *ApoE4* alleles is a major independent risk factor for Alzheimer’s disease and so violates the exclusion restriction criteria of Mendelian randomisation studies (Figure 4.18). In all cases, excluding variants in the *ApoE* region attenuates the observed effect to the null. A clear example of this is Benn *et al* (2017), where *ApoE* variants were not sufficiently identified and excluded, leading the published paper to report a protective effect of LDL-c on Alzheimer’s (RR: 0.83, 95%CI: 0.75-0.92).⁷⁴ Following several rapid responses, the data was re-analysed to exclude a larger area around *ApoE* which attenuated the finding to the null.²⁷³

4.5.4 Inclusion of preprints

As highlighted in Section 1.5.1, this review explicitly sought to synthesize all available evidence, irrespective of publication status (preprinted vs. published). Using the tool described in Chapter 2.1, two preprint servers related to health and

4.5 - Discussion

biomedical sciences were searched as part of this review. The small number of studies returned by the preprint searches (or the absence of any relevant hits in the medRxiv database - see Figure 4.1) is largely due to the timing of the preprint searches. The searches for this review were performed in mid-July 2019, but the medRxiv repository, an offshoot of the Epidemiology and Clinical Trials categories of the bioRxiv preprint server, only registered its first preprint 25th June 2019. As such, at the point it was searched, the medRxiv database contained only a very small number of records ($N = 148$; Figure 4.19).




Figure 4.19: Growth of preprint repositories over time. Given the relative sizes of the preprint repositories at the time the searches for this review were conducted (bioRxiv $N = 56,007$, medRxiv $N = 148$), the number of hits returned by each is expected.

4.5 - Discussion

Three relevant preprints from bioRxiv were identified. Of note, all three described a Mendelian randomisation study, potentially indicating that more biologically-focused study designs are over-represented in the bioRxiv repository. The added evidential value of including these preprints was described in Section 4.4.3 and indicated that results available only via preprinted reports can contribute additional evidence to a meta-analysis. However, the scale of this contribution is questionable. Preprinted evidence was incorporated into only a single meta-analysis in this review, and inclusion of the preprinted result did not substantially impact the summary effect estimate.

Of the three identified preprints, two were subsequently published as of September 2021. This fits well with the analysis presented in Chapter 2 that, allowing for a two-year lag, approximately two-thirds of preprints are published. It also illustrates the dual advantages of preprinted reports to evidence syntheses. Firstly, preprints provide an advance snapshot of the literature, capturing articles that will eventually be published but were not available at the time of the main search. For example one eligible preprint in this review was initially posted on bioRxiv in July 2019²⁵⁵ and was subsequently published in 2021 following peer-review.²⁵⁰ Secondly, inclusion of preprints allows for results that may never be formally published to be included in an evidence synthesis exercise, as is the case with a second preprint included in this review.²⁵³ Both of these aspects illustrate that inclusion of preprints is a necessity if the aim is to find all relevant literature on a topic at the time of searching.

Recently, inclusion of preprints in systematic reviews has become significantly more widespread. This is largely due to the role of preprint servers, in particular medRxiv, as a key evidence dissemination venue during the early stages of the COVID-19 pandemic.⁸⁴ How well this adoption of preprints will transfer to other less-urgent topics, where the speed of research does not put the same focus on preprinted articles, is currently unknown.

4.5.5 Open data sharing

As discussed in Section 4.4.1, many primary studies did not report important information required for the dose-response meta-analysis, and so could not be included in the synthesis. This limitation was compounded by the expected low response rate to requests for further information from primary authors. Contacting authors is worthwhile, as it can substantially change the conclusion of a systematic review²⁷⁴ and is not too costly to systematic reviewers,²⁷⁵ though a preferable option is that the authors of primary studies readily deposit all relevant study data at the point of publication.

Based on my experience of extracting data for this review, I co-authored a guidance article to aid primary prevention scientists in preparing and sharing their data so that it can easily be incorporated into a evidence synthesis exercise, using a trial of mindfulness interventions as a case study.²⁷⁶ A copy of this publication is available in Appendix B.3. In an attempt to apply my own guidance, I have invested a substantial amount of time and effort into making the data obtained by this review openly available to other researchers, via a GitHub repository.

4.5.6 Strengths and limitations

Strengths

There are several aspects where this review is distinct from those already available in the published literature. While several reviews of this research topic exist,^{260–263} the overlap between the list of studies included in each is not complete. As part of this review, I have not only performed an original search of primary literature databases, but have also screened the reference lists of comparable reviews to ensure no relevant study has been omitted.

Secondly, this review employed a structured approach to risk-of-bias assessment using a domain-based tool. This represents an important strength of this review, as the

detailed risk of bias assessments are used to inform the quantitative triangulation analysis presented in Chapter 7.

Thirdly, as discussed at length in the section above, in contrast to other available reviews and enabled by the tool described in Chapter 2, this review systematically searched preprinted health-related preprints. Finally, as a secondary element, I used this review to pilot new research synthesis methodologies, in particular a new visualisation approach for risk-of-bias assessments and a forthcoming tool for assessing the risk-of-bias due to missing evidence.

Limitations

The primary limitation of this review is that several included studies used data from EHR databases, which come with serious concerns regarding validity.^{277–279} Relatedly, several studies which made use of electronic health record databases did not report the specific code lists used, potentially introducing substantial heterogeneity into the analysis. An empirical example of the effect of differing EHR code lists is presented as part of the analysis in Chapter 5 (see Section 5.3.4).

In addition, the fact that only a sample of records were dual screened at the title/abstract and full-text stages is a potential limitation, as there is a chance that some eligible records could have been excluded. However, evidence from assessments of inter- and intra-rater reliability indicate that this is not a major concern.

One particular limitation with regards to the risk-of-bias assessment is the fact that the ROBINS-E assessments were performed using an adapted version of the ROBINS-I tool. This meant that there were no signaling questions to guide the domain-level risk-of-bias assessments, which may have influenced the accuracy with which the judgements were assigned. However, there is no published empirical evidence supporting the need for signaling questions and assessment of inter-rater reliability across the different tools did not indicate a specific problem with the

ROBINS-E assessments. In fact, low agreement was common across the tools, though this is expected based on the available literature.²⁸⁰

A final limitation is the potential for missing evidence on the basis of the results. Evidence for this limitation came from the ROB-ME assessments and was supported by empirical evidence that some studies containing relevant results were missed by the search (see Section @Ref(rev-discussion-MR) above for a fuller discussion of this issue with respect to Mendelian randomisation studies). Unfortunately, this is probably a common limitation across all systematic reviews, based on the way in which increased search sensitivity must be balanced with a manageable workload.

4.6 Summary

- In this chapter, I presented the results of a comprehensive systematic review of existing evidence on the association between lipid levels and dementia use. The review included both direct (studies that examined lipid levels directly) and indirect (studies examining lipid regulating agents such as statins) forms of evidence. In contrast to previous reviews, I also included preprinted evidence, facilitated by the research tool introduced in Chapter 2.
- I found no consistent relationship between lipids or lipid-regulating agents and dementia across different evidence sources. However, there was some indication of a protective effect of statins on all-cause and Alzheimer's disease dementia in non-randomised studies of interventions.
- The findings from this review are used throughout the subsequent chapters. In Chapter 5, this summary of the evidence guided the choice of analysis approach, ensuring that the new analysis was at risk of a different source of bias and provided evidence on an under-studied outcome (vascular dementia). In Chapter 6, prospective cohorts identified by the review were contacted in an attempt to obtain individual participant data. Finally, the results

4.6 - Summary

identified here are used as a key source of evidence for the triangulation exercise presented in Chapter 8.

When dealing with human beings controlled experiments frequently prove to be impracticable, so for a scientific basis for our assumptions we turn to past history to reconstruct the suspected causal chain of events - and then our statistical troubles may begin.

— Harold F. Dorn, 1953²⁸¹

5

Primary analysis of lipid-regulating agents and dementia outcomes

Contents

Lay summary	108
5.1 Introduction	108
5.2 Methods	109
5.3 Results	123
5.4 Discussion	139

Lay summary

Electronic health record (EHR) databases are large collections of medical data, used to manage patient administration and care. Under these systems, whenever a patient attends their GP, their clinical data is recorded in a central database using a standardised coding system. These databases have several advantages over traditional methods of data collection, including the number of people they contain and the relatively low cost of data collection via routine care. This is particularly important when studying diseases such as dementia, which may begin to develop in patients long before symptoms are seen.

This analysis makes use of the Clinical Practice Research Datalink (CPRD), which contains the electronic medical records of more than 3 million people from general practices across England. Using this data, the analysis presented in this chapter examined whether treatments which lower cholesterol levels (also known as lipid-regulating agents or LRA) of which statins are a prime example, affect the risk of all-cause dementia and related outcomes (Alzheimer's disease, vascular dementia and other dementias).

Little evidence for an effect of lipid-regulating agents effect on the risk of Alzheimer's disease was found, with the exception of a slightly increased risk in those prescribed a certain type of lipid-regulating agent called fibrates. In contrast, I found an increased risk of vascular and other (i.e. non-Alzheimer's) dementia was associated with lipid-regulating agent use.

This increased risk in outcomes with a vascular element (e.g. vascular dementia) is unexpected, and is very likely to be due to the presence of bias in the analysis. This bias, called "confounding by indication", is caused when those who are prescribed a statin are more at risk of vascular dementia for a range of reasons, which makes it appear as if statins are harmful. Despite this limitation, the analysis presented provides an important source of information which will be used in later chapters.

5.1 Introduction

In this chapter, I present an analysis of a large population-based electronic health record dataset to investigate the relationship between lipid-regulating agent (LRA) use and dementia outcomes. The analysis aims to address two important limitations

of the current evidence base as identified by the systematic review presented in Chapters 3 & 4.

Firstly, it explicitly examines vascular dementia as an outcome. The systematic review presented in the previous chapter identified an evidence gap around the effect of lipid-regulating agents on the risk of vascular dementia. As triangulation exercises require as many diverse sources of evidence as possible, this analysis provides a source of information on this outcome.

Secondly, and in a similar vein, the analysis intentionally takes a different analytical approach to that most commonly used to examine the effect of statins on dementia as identified by the systematic review. Specifically, this involved a concerted effort to address immortal time bias through use of a Cox proportional hazards analysis, incorporating a time-varying treatment indicator.²⁸² By employing this approach, the analysis provides an evidence source at risk of a distinct bias. The results from this analysis will be incorporated into the triangulation exercise presented in Chapter 8.

This chapter represents an extended version of a preprinted manuscript, a copy of which is available in Appendix B.3.

5.2 Methods

5.2.1 Study protocol

An *a priori* protocol for this study was published,²⁸³ and amendments to this are recorded in Appendix A.5.1.

5.2.2 Data source

Previously known as the General Practice Research Database, the Clinical Practice Research Datalink (CPRD) is a large population-based EHR database.²⁸⁴ The

5.2 - Methods

database has been collecting primary care data from participating practices across England since 1987.^{285,286} It contains the primary care records for more than 10 million primary care patients in England, and is broadly representative of the UK population in terms of age, sex and ethnicity.^{284,287}

To avoid the ambiguity of interpreting free-text clinical notes and to allow for easy analysis of the resulting data, the CPRD primarily collects data using a predefined coding system known as Read codes.²⁸⁸ All clinical events, including clinical test results and diagnoses, can be identified by a specific Read code. The codes use a nested approach (see Table 5.1), with the initial characters defining broad diagnostic topics (e.g. Eu... - Mental and behavioural disorders), while subsequent characters provide additional information on the specific condition diagnosed (e.g. Eu001 - Dementia in Alzheimer's disease with late onset).

Table 5.1: Example of CPRD Read code hierarchy, showing how “Dementia in Alzheimer’s disease with late onset” (*Eu001*) is nested under the top-level of “Mental disorders” (*Eu...*). Broad topics are specified using the initial two alpha-numeric characters of the Read code, while subsequent characters are used to define specific conditions and context.

Level	Read code	Term
1	E....	Mental disorders
2	Eu...	Mental and behavioral disorders
3	Eu0..	Organic mental disorder
4	Eu00.	Dementia in Alzheimer’s disease
5	Eu001	Dementia in Alzheimer’s disease with late onset

The index events, exposures and outcomes used in this analysis were identified using predetermined code lists, which are available for inspection from the archived repository accompanying this analysis (data/code availability is discussed in Section 5.4.4).

5.2.3 Cohort definition

This analysis included all participants registered at a participating practice between 1 January 1995 and 29 February 2016 who had a flag for “research quality” data (as defined by the CPRD). Records pre-dating the 1995 cut-off were included in the original CPRD extract obtained for this analysis. However, these older records were excluded from the analysis as data quality and reliability are thought to be higher after this date.²⁸⁹ Additionally, individuals with less than 12 months of continuous records prior to cohort entry were excluded, making the effective start date of the cohort 1 January 1996.

Participants were included in the study cohort if their record contained any of the following index events:

- a Read code for a diagnosis of hypercholesterolemia or related condition;
- a Read code for prescription of a lipid-regulating agent (such as statins);
- a total cholesterol test result of >4 mmol/L;
- an LDL-c test result of >2 mmol/L.

The blood lipid cut-offs were based on NIHR-recommended levels at the time the protocol was written. These index events allowed me to define a population of participants who were either at risk of hypercholesterolemia, as indicated by the elevated total or LDL cholesterol test results, or had already been diagnosed with it, as indicated by a diagnostic code/related prescription.

The index date for a participant was defined as the date where the first relevant code or test result was recorded on their clinical record, and participants were followed up until the earliest of:

- an outcome of interest;
- death;
- end of follow-up (29 February 2016);
- last registration date with their GP practice;

- the last CPRD collection date for their practice.

Participants were ineligible for the cohort if they were less than 40 years of age (as these patients are less likely to be prescribed a LRA), had less than 12 months of “research quality” data, were simultaneously prescribed more than one lipid-regulating agent (due to the difficult of assigning these to a single exposure group), or were diagnosed with an outcome of interest before or on the date of the index event (i.e. had less than one full day of follow-up).

5.2.4 Exposures

I considered seven lipid-regulating drug classes based on groupings in the British National Formulary,²⁹⁰ namely: statins, fibrates, bile acid sequestrants, ezetimibe, nicotinic acid groups, ezetimibe and statin (representing one treatment containing both drugs, rather than the two classes being prescribed concurrently), and omega-3 fatty acid groups.

A participants drug class was assigned based on their first recorded prescription, and any drug switching was ignored in an effort to mimic an intention-to-treat approach. We did however examine how often the initial drug class altered according to one of three criteria:

- **stopped:** defined as the last prescription of the primary class being followed by at least six months of observation;
- **added:** defined as a second drug class being prescribed before the last prescription of the initial class; and
- **switched:** defined as a second drug class being prescribed after the last prescription of the initial class.

5.2.5 Outcomes

I considered five outcomes as part of this analysis: probable Alzheimer's disease, possible Alzheimer's disease, vascular dementia, other dementias, and a composite all-cause dementia outcome. When two or more outcomes were coded in a participant's clinical record, a decision tree was used to differentiate between them (Figure 5.1). The diagnosis date of the outcome was determined by the first record of a relevant code.




Figure 5.1: Decision tree for assigning dementia subtypes, based on the presence of Read codes in the patient's record. Note that an outcome of “probable” or “Possible” Alzheimer's disease (AD) requires the absence of any vascular outcome codes.

5.2.6 Covariates

A range of additional variables were included in the analysis. These were intended to address the different distributions of potential confounding factors between those who were prescribed a lipid-regulating agent and those who were not. These are discussed in detail below and summarised in Table 5.2.

5.2 - Methods

Charlson index implemented using Read code lists.²⁹² Code lists based on those by Taylor et al.²⁹³

Table 5.2: Definition of covariates adjusted for in the full-adjusted model. The code lists used to define the majority of these covariates were originally created for use in a previously published analysis.²⁹¹ while others were built on or adapted from previous published work,^{292–294}.

Covariate	How was the covariate defined?
Previous history of coronary arterial disease	Presence of one or more relevant Read codes on record.
Previous history of coronary bypass surgery	Presence of one or more relevant Read codes on record.
Previous history of cerebrovascular disease (including stroke)	Presence of one or more relevant Read codes on record.
Chronic illness, including cancer and arthritis	Charlson index implemented using Read code lists. ²⁹² Code lists based on those by Taylor et al. ²⁹³
Socioeconomic position	2010 English Index of Multiple Deprivation (IMD) at the twentile level, where 1 represents the least deprived and 20 the most deprived.
Consultation rate	Calculated by dividing the total number of clinic visits by the length of the patient record prior to the index date to give an average annual rate.
Alcohol status	Recorded value (current, former or never).
Smoking status	Most recent of recorded value (current, former or never) or Read code indicating a recorded value. Code lists based on those by Wright et al. ²⁹⁴
Body Mass Index	Recorded value if available, or a calculated value using the last recorded height and weight measurements. Measurements taken before the age of 25 were excluded to ensure adult measurements were used.
Peripheral arterial disease	Presence of one or more relevant Read codes on record.
Hypertension	Presence of one or more relevant Read codes on record.
Baseline total cholesterol	Continuous value recorded as test result ("enttype==163 & test_data1==3")
Baseline LDL cholesterol	Continuous value recorded as test result ("enttype==177 & test_data1==3")
Chronic kidney disease	Presence of one or more relevant Read codes on record.
Type 1 Diabetes	Presence of one or more relevant Read codes on record.
Type 2 Diabetes	Presence of one or more relevant Read codes on record.

Demographic covariates adjusted for included age and gender. Age in years was

calculated at date of entry into the cohort, using the 1st of January of a patient's birth year (the exact date of birth was not provided by CPRD), and was adjusted for via its use as the time axis for the Cox model (see Section 5.2.10). Socioeconomic status was proxied using the Index of Multiple Deprivation (IMD) 2010. The IMD draws on seven domains (income; employment; education, skills and training; health and disability; crime; barriers to housing and services; living environment) to create an overall deprivation score for each of 32844 statistical geography areas in England. To help preserve patient privacy, IMD score is only available from the CPRD in twentiles, with 1 indicating the least deprived and 20 indicating the most deprived. Smoking and alcohol use was determined at index, and was categorised as current, former, or never use.

Body mass index (a summary measure calculated as $weight/height^2$), baseline total cholesterol and baseline LDL cholesterol measures were obtained, using the last recorded value prior to the index date. A variable indicating grouped year of entry into the cohort (≤ 2000 , 2001-2005, 2006-2010, > 2010) was included to allow for changes in prescribing trends across the lifetime of the cohort. To assess healthcare utilisation, I adjusted for the average annual number of consultations between the beginning of a patient's data and their entry into the cohort.

Finally, presence of a range of related conditions at baseline were accounted for, including cardiovascular disease, coronary bypass surgery, coronary artery disease, peripheral arterial disease, hypertension, chronic kidney disease, and Type 1 and Type 2 diabetes. In addition to adjusting for these covariates individually, a Charlson co-morbidity index (CCI) score was calculated for each participant. The CCI is a weighted index that uses presence and severity of a number of conditions to enable adjustment for the general health of a participant in terms of their mortality risk.²⁹⁵ The conditions considered under the index are: AIDS; cancer; cerebrovascular disease; chronic pulmonary disease; congestive heart disease; dementia; diabetes; diabetes with complications; hemiplegia; metastatic tumour; mild liver disease; moderate liver disease; myocardial infarction; peptic ulcer disease; peripheral vascular disease; renal disease; and rheumatological disease. Inclusion of

this index allowed me to attempt to adjust for the general health of patients included in the analysis.

code lists for all covariates can be found in the archived data repository accompanying this analysis (see Section 5.4.4).

5.2.7 Missing data

Missing data are a recognised problem in electronic health records databases.²⁹⁶ These databases are created from administrative data, collected primarily for the purposes of patient management and care rather than academic research.

In this analysis, missing data was handled using a multiple imputation approach.²⁹⁷ Variables with missing observations were identified for inclusion in the imputation model. Nominal variables with missing values were modelled using multinomial logistic regression, while continuous variables were modelled using linear regression. As per best practice, all variables used in the analytic model, including the outcome, were included in the imputation model.²⁹⁸ Using the MICE (Multiple Imputation by Chained Equations) command in STATA16, 20 imputed datasets were created.

Missing data was only considered problematic for variables where a numerical test result was expected (e.g. BMI), or where a code existed for the absence of the condition (e.g. categorical smoking status). This approach was necessary, as absence of a code for other treatments or conditions (e.g. statin use or dementia) was assumed to indicate absence of the treatment/condition rather than being considered missing.²⁹⁶

5.2.8 Estimation methods

A Cox proportional hazards (PR) model was used to estimate the effect of statins on dementia outcomes. Cox PR models are defined in general terms as:

$$h(t) = h_o(t) \times \exp(b_1x_1 + b_2x_2 + \dots + b_px_p) \quad (5.1)$$

where:

- t is the survival time;
- $h(t)$ is the hazard function; and
- x_1, x_2, \dots, x_p are the covariates which determine the hazard function, while b_1, b_2, \dots, b_p are the coefficients for each covariate.
- $h_o(t)$ is the baseline hazard - when all x_i are zero, the $\exp()$ function resolves to 1.

A Cox PR model was chosen for this analysis as it inherently accounts for the length of time participants spend in each exposure group. Using this approach, time-at-risk can be properly attributed to the appropriate exposure group, thus mitigating the impact of immortal time bias. This is discussed in detail in the following section.


5.2.9 Immortal time bias and time-varying treatment indicators

Immortal time bias describes two distinct but related types of bias, considered here in relation to statin use. The first presentation, the selection bias aspect (Panel A, Figure 5.2), occurs when time prior to statin initiation is excluded leading to the statin and control groups being followed up from different time points.²⁹⁹ Following the example displayed in the figure, the unexposed group are followed from the date of an index event (e.g. diagnosis of hypercholesterolemia), while the statin group is followed from date of a statin initiation. In this scenario, the time between the index event and statin initiation is missing, and any events that occur in the exposed group prior to the prescription will be inappropriately excluded from the analysis.

5.2 - Methods

The second presentation of immortal time bias is as a type of misclassification bias (Panel B, Figure 5.2). It occurs when the exposure time prior to statin initiation, and any events occurring within it, is inappropriately assigned to the exposed group.

A



B




Figure 5.2: Diagram illustrating the two presentations of immortal time bias, as a selection bias (Panel A) and a misclassification bias (Panel B).

This second presentation appears to be common in the existing literature on the relationship of statins and dementia, as several of the studies included in the systematic review performed as part of this thesis were identified as being at risk of immortal time bias following formal risk of bias assessment using the ROBINS-I tool (see Section 4.3.4).

Following a recommended approach to addressing the second form of immortal time bias, I employed a time-varying indicator of treatment status to correctly allocate time-at-risk to the exposed and unexposed groups.²⁹⁹ Under this approach,

all patients are followed from a common index date, defined as earliest of: (a) date of raised cholesterol test results; (b) hypercholesterolemia diagnosis; or (c) LRA prescription. Patients start in the unexposed group and contribute time-at-risk until they are prescribed a lipid-regulating agent, at which point they move into the exposed group. Note, patients for whom prescription of a lipid-regulating agent was the index event only contribute time to the exposed group (i.e. they enter the cohort and move into the exposed group on the same day).

5.2.10 Time axis

As part of a Cox proportional hazard model, there is the option to use either absolute time in cohort or participants age as the time scale of interest.^{300–302} A model using age as the time axis inherently accounts, or adjusts, for participants age as a potential confounder of the exposure-outcome relationship. As such, the main analyses presented all used age as the time axis.

5.2.11 Sensitivity analyses

The primary analysis examined the effect of a lipid-regulating agent on dementia risk, stratified by outcome and drug class. To assess the robustness of the results, a number of sensitivity analyses were performed. These are described in the following sections.

Complete case vs imputed data

Using multiple imputation to handle missing data is an alternative to a “complete case” approach,³⁰³ where participants missing any covariate are dropped from the dataset. As a recommended sensitivity analysis,³⁰⁴ I performed and compared

5.2 - Methods

the results of both methods, to investigate the impact of multiple imputation on the results.

Control outcomes

In addition to the primary outcomes of interest (described in Section 5.2.5), I extracted data on three additional control outcomes. The inclusion of control outcomes in observational analyses are a useful technique to assess the strength of uncontrolled confounding,³⁰⁵ and these outcomes are usually classed as either “negative” or “positive” outcomes.

Negative outcomes are defined as those without a likely causal path between the exposure and outcome (see Figure 5.3 for a directed acyclic graph, or DAG, describing an ideal negative outcome). Conversely, positive control outcomes are those with a known causal association with the exposure of interest, ideally sourced from large well conducted randomised controlled trials. Positive control outcomes are useful in observational epidemiology, as if the analysis can reproduce a known result for the control outcome, confidence in the result for the outcome of interest is increased. Due to the wealth of data available on statins as a lipid-regulating agent, I chose three control outcomes in reference to this drug class: back pain (negative control), ischaemic heart disease (positive protective control), and Type 2 diabetes (positive harmful control).




Figure 5.3: Causal diagram (directed acyclic graph) showing relationship between exposure X , outcome Y , confounders (measured C and unmeasured U) and an ideal negative outcome N . Note the absence of any arrow between X and N . In this scenario, any association observed between X and N is due to the presence of uncontrolled confounders U (assuming C has been adjusted for).

Despite observational analyses suggesting a link between statins and muscular pain (as opposed to more serious complications such as myopathy),³⁰⁶ systematic reviews of the adverse events of statin use³⁸ and N-of-1 trials explicitly exploring the association of statin use with muscle pain³⁰⁷ have found little evidence supporting an effect. This suggests that muscular backpain would be suitable for use as a negative control outcome in this analysis. Under this approach, if statin use is found to be associated with muscular backpain in this analysis, this suggests the presence of residual confounding and reduces my confidence in the results for the dementia outcomes.

Similarly, incident ischemic heart disease and Type 2 diabetes were included as a protective and harmful positive control outcome, respectively. The protective effect of lipid-lowering treatment, via statins, on the risk of ischemic heart disease is well-established,³⁸ while there is growing evidence for an increased risk of Type 2 diabetes with statin use.^{38,308,309} Similar to the negative outcome, if the analysis strategy can reproduce these known associations, this will provide evidence that potential confounders have been sufficiently adjusted for.

Impact of additional covariates

To observe the effect of adjusting for additional covariates, I ran two additional models unadjusted except for: (a) age; and (b) age and gender. The results of these models was then compared the results with the fully adjusted model.

Sensitivity cohorts

Two sensitivity cohorts were also created. The first stratified by year of entry into the cohort in an attempt to assess for time period effects. The second removed participants who may have been pregnant (coded as under 55) to assess the robustness of the estimates, as statins are contraindicated in pregnancy.³¹⁰

Statin properties

As detailed in the introduction, the properties of statins may be important due to the ability of lipophilic statins to cross the blood brain barrier (see Section 1.3.2).⁴¹ As such, I expected that any effects of statins on dementia outcomes would be stronger in the lipophilic as compared to the hydrophilic statin subgroup. To investigate this, I further stratified the statin exposure group into lipophilic (Atorvastatin, Lovastatin, Simvastatin, Cerivastatin) and hydrophilic (Pravastatin, Rosuvastatin, Fluvastatin) statins.

Impact of using different code lists for defining dementia outcomes

As part of an exploratory analysis of the effect of the choice of code lists on the analysis, I created an alternative Alzheimer's disease and non-Alzheimer's

5.3 - Results

dementia outcome using code lists from a published study by Smeeth *et al.*²¹³ The intended purpose of this analysis was to assess the robustness of my results to the choice of code list.

This published analysis used a propensity matching approach to estimate the association of statins with a range of outcomes in The Health Improvement Network database, an alternative source of English electronic health records which has substantial overlap with the CPRD.³¹¹ The code lists used in this analysis were obtained through correspondence with the authors of that study, and are available for inspection (see Section 5.4.4).

5.3 Results

5.3.1 Patient characteristics

Of the 3,179,733 participants included in the extract, 1,684,564 met the inclusion criteria (Figure 5.4), with a total follow-up of 10,835,685 patient years at risk.

5.3 - Results




Figure 5.4: Attrition of CPROD participants as the eligibility criteria were applied. The largest cause of attrition was the absence of an index event of interest.

5.3 - Results

The median participant age at index was 57 years (inter-quartile range (IQR):48-67years) and participants were followed up for a median of 5.9 years (IQR:2.7-9.7years). During follow-up, an all-cause dementia diagnosis was recorded for 41,830 patients (12,647 probable AD, 9,954 possible AD, 8,466 vascular dementia, 10,763 other dementias).

The number of events, time-at-risk and crude rates for each drug class, tabulated by dementia outcome, are shown in Table 5.3. A substantial majority (98.1%) of participants prescribed a lipid-regulating agent were prescribed a statin. I excluded the “Ezetimibe and statins” (N =127) and “Nicotinic acid groups” (N = 165) classes from subsequent class-based subgroup analyses based on the extremely small number of participants in these groups. Note that the “Ezetimibe and statins” treatment group represent those prescribed a single treatment containing both ezetimibe and statins, rather than those where the two treatments were prescribed concurrently.

Table 5.3: Summary of number of events, years at risk and crude rates per 100,000 participant-years-at-risk stratified by dementia outcome and drug class of interest.

Exposure Group	Any dementia			Possible AD			Probable AD			Vascular dementia			Other dementia		
	Events	PYAR	Rate *	Events	PYAR	Rate *	Events	PYAR	Rate *	Events	PYAR	Rate *	Events	PYAR	Rate *
No LRA (unexposed)	18,608	5,872,717	317	6,368	5,818,047	109	2,637	5,800,964	45	4,813	5,811,594	83	4,790	5,808,285	82
By drug class															
Statins	22,920	4,871,568	470	6,190	4,758,526	130	5,773	4,753,437	121	5,871	4,755,258	123	5,086	4,747,237	107
Omega-3 FAGs	19	8,034	236	4	7,927	50	7	7,950	88	4	7,938	50	4	7,925	50
Fibrates	141	38,003	371	49	37,102	132	21	36,835	57	36	37,001	97	35	36,983	95
Ezetimibe	32	6,604	485	8	6,429	124	7	6,425	109	12	6,444	186	5	6,393	78
BAS	106	36,370	291	28	35,808	78	19	35,726	53	26	35,768	73	33	35,808	92
Ezetimibe + Statins †	0	986	-	0	986	-	0	986	-	0	986	-	0	986	-
NAG	4	1,403	-	0	1,379	-	2	1,391	-	1	1,389	-	1	1,382	-
Total	41,830	10,835,686	386	12,647	10,666,205	119	8,466	10,643,714	80	10,763	10,656,378	101	9,954	10,644,999	94

* Crude rate per 100,000 participant-years-at-risk

† One treatment containing both drugs, rather than the two classes being prescribed concurrently

Abbreviations: AD - Alzheimer's disease; BAS - Bile acid sequestrants; NAG - Nicotinic acid groups; Omega-3 FGs - Omega-3 Fatty acid groups; PYAR - Participant-years-at-risk.

5.3 - Results

Table 5.4: Patient characteristics by drug class. Summary statistics are presented as “% (N)” unless otherwise specified in the variable name.

	Whole Sample	None	Statins	Bile acid sequestrants	Ezetimibe	Fibrates	Omega-3 Fatty Acid Groups
Sample size (N)	1,684,564	1,087,704	585,528	5,396	763	3,889	992
Year of cohort entry (median)	2006	2007	2004	2005	2004	2001	2005
Female	53.0% (893174)	56.2% (610950)	47.1% (276043)	66.4% (3585)	54.5% (416)	38.6% (1500)	52.6% (522)
Age at cohort entry (median)	57	54	62	57	60	58	56
CAD	0.4% (7133)	0.1% (589)	1.1% (6465)	0.1% (6)	0.9% (7)	1.4% (53)	1.3% (13)
CBS	0.3% (5699)	0.1% (682)	0.8% (4926)	0.1% (4)	0.4% (3)	2.0% (78)	0.6% (6)
CVD	2.1% (34899)	1.1% (11619)	3.9% (22977)	1.6% (86)	2.6% (20)	4.4% (170)	1.7% (17)
Charlson (ever > 0)	30.6% (516135)	25.1% (272642)	40.7% (238403)	42.5% (2292)	41.7% (318)	50.8% (1976)	40.4% (401)
IMD-2010 (median)	9	8	9	8	9	10	10
Consultation rate (mean/SD)	5.4 (5.4)	5.0 (5.0)	6.2 (6.1)	8.6 (7.4)	7.4 (6.6)	7.1 (6.2)	8.0 (8.0)
Alcohol (ever)	85.9% (1447151)	86.6% (941648)	84.7% (496110)	82.8% (4468)	84.0% (641)	82.9% (3223)	82.0% (813)
Smoking (ever)	51.1% (861355)	47.1% (511826)	58.6% (343074)	55.2% (2978)	57.5% (439)	60.2% (2341)	53.7% (533)
BMI (mean/SD)	27.0 (5.3)	26.7 (5.2)	27.7 (5.3)	26.8 (5.8)	28.1 (5.7)	29.0 (5.2)	26.9 (5.5)
PAD	0.7% (12613)	0.4% (4039)	1.4% (8424)	0.9% (47)	0.9% (7)	1.9% (75)	1.0% (10)
Hypertension	16.0% (269804)	11.5% (124604)	24.4% (143101)	12.8% (692)	23.9% (182)	25.8% (1002)	15.7% (156)
Total cholesterol (mean/SD)	5.7 (10.1)	5.5 (6.4)	6.2 (15.3)	5.3 (1.3)	7.1 (26.5)	6.4 (5.6)	5.6 (1.6)
LDL cholesterol (mean/SD)	3.6 (4.9)	3.4 (5.3)	4.0 (3.7)	3.1 (1.0)	3.9 (1.1)	3.3 (1.8)	3.2 (1.0)
CKD	0.1% (1295)	0.1% (740)	0.1% (545)	0.1% (6)	0.1% (1)	0.0% (0)	0.3% (3)
Type 1 Diabetes	0.2% (4037)	0.1% (785)	0.5% (3196)	0.3% (14)	1.0% (8)	0.8% (31)	0.1% (1)
Type 2 Diabetes	2.9% (48557)	1.1% (11797)	6.1% (35941)	2.3% (123)	5.4% (41)	15.8% (614)	2.8% (28)

Note: The ‘Nicotinic acid groups’ (n=165) and ‘Ezetimibe and Statins’ (n=127) subgroups are not shown, but are included in the whole sample column

Abbreviations: BMI - Body mass index; CAD - Coronary arterial disease; CBS - Coronary bypass surgery; CKD - Chronic kidney disease; CVD - Index of multiple deprivation; LRA - Lipid regulating agent; PAD - Peripheral arterial disease; SD - Standard deviation.

5.3 - Results

The distribution of baseline characteristics across the remaining seven drug classes can be seen in Table 5.4. Note due to the experimental design, the median year of entry is expected to be later for those not prescribed an LRA, as this exposure group is more likely to include those who entered into the cohort towards the end of study window (as these people had less follow-up time in which to be prescribed an LRA).

The stopping, addition and switching of drug classes was common across all drug classes (Table 5.5).

Table 5.5: Participants who stopped, switched or added treatments by initial treatment type.

	Whole Sample	Statins	Bile acid seques-trants	Ezetimibe	Ezetimibe & Statins	Fibrates	Nicotinic acid groups	Omega-3 Fatty Acid Groups
Stopped	6.9% (115899)	19.1% (111798)	56.1% (3028)	19.7% (150)	12.6% (16)	12.3% (478)	44.8% (74)	35.8% (355)
Added	1.6% (27441)	4.4% (25990)	3.6% (192)	19.0% (145)	3.9% (5)	21.6% (841)	3.6% (6)	26.4% (262)
Switched	0.9% (14935)	2.0% (11996)	11.3% (612)	34.6% (264)	64.6% (82)	44.0% (1713)	45.5% (75)	19.5% (193)

Definitions: Stopped - last prescription of the primary drug class followed by at least six months of observation with no further prescriptions; Added - second drug class prescribed before the last prescription of the initial class; Switched - second drug class being prescribed after the last prescription of the initial class.

5.3.2 Missing data

Full covariate information was available for 450,234 participants (26.7%). Six key variables had some missing data: IMD 2010 score was missing for 625,788 participants (37.1%), because it is only recorded for English practices; alcohol status was missing for 269,526 participants (16%); smoking status was missing for 84,424 participants (5%); BMI, or a calculated BMI from height and weight measurements, was missing for 266,672 participants (15.8%); baseline total cholesterol was missing for 119,675 participants (7.1%); and baseline LDL cholesterol was missing for 787,289 participants (46.7%).

5.3.3 Primary analysis

The results of the primary analysis using the fully adjusted Cox proportional hazards model with participant age as the time scale are presented for each drug/outcome combination in Figure 5.5.

For each outcome, the overall “Any drug” estimate was driven by the statin subgroup, based on its large size relative to the other drug classes.

5.3 - Results




Figure 5.5: Results from primary analyses of CPRD data using the fully adjusted model and participant age as the time scale.

Alzheimer's disease

My results show little evidence was found for an effect of lipid-regulating agents on probable (HR:0.98, 95%CI:0.94-1.01) and possible (HR:0.97, 95%CI:0.93-1.01) Alzheimer's disease when compared to no treatment, with the sole exception of fibrates on probable Alzheimer's disease (HR:1.52, 95%CI:1.13-2.03).

Non-Alzheimer's disease dementias

In contrast to the findings for Alzheimer's disease outcomes, an association between lipid-regulating agents and an increased risk of a subsequent diagnosis of vascular dementia (HR:1.81, 95%CI:1.73-1.89) or other dementias (HR:1.19, 95%CI:1.15-1.24) was observed. Again this effect was driven mainly by the statin subgroup, but there was some evidence that ezetimibe was associated with an increased risk of vascular (HR:2.33, 95%CI:1.11-4.89) and other (HR:1.88, 95%CI:1.01-3.5) dementia.

All-cause dementia

For the composite all-cause dementia outcome, I found treatment with a lipid-regulating agent was associated with a slightly increased risk (HR:1.17, 95%CI:1.14-1.19), which lies between the associations for the Alzheimer and non-Alzheimer dementia outcomes as would be expected. There was also some evidence that fibrates were associated with increased risk of all-cause dementia (HR:1.28, 95%CI:1.08-1.52).

5.3.4 Sensitivity analyses

The results of the series of sensitivity analyses performed are described in the following sections.

Complete case versus imputed data

In almost all cases, the use of imputed data resulted in a marginal attenuation of the effects observed when using a complete cases analysis. It should be noted that due to the large amount of missing data (e.g. 787,289 participants (46.7%) were missing a baseline LDL cholesterol measure), the number of participants included in the complete case analysis was substantially smaller than that included when using imputed data. In this case, though the overall position of the effect estimates does not change substantially when using the imputed dataset, there is a noticeable gain in power.²⁹⁷




Figure 5.6: Comparison of analyses using the complete case versus imputed cohorts.

Control outcomes

Following the primary analysis, the fully adjusted model was used to estimate the effect of treatment with a statin on the two control outcomes of back pain (negative) and ischemic heart disease (positive). The results of this analysis are presented in Figure 5.7.

For the negative control, there was some evidence that treatment with a statin was associated with an increased risk of back pain (HR: 1.04, 95%CI: 1.03-1.05), suggesting there may be some residual confounding. However, statin prescription was also associated with a substantially increased risk of ischemic heart disease (HR: 1.62, 95%CI: 1.59-1.64) and Type 2 diabetes (HR: 1.50, 95%CI: 1.48-1.51).




Figure 5.7: Results of control outcome analysis.

Impact of additional covariates

The results of three models adjusted for age only, age and sex, and the full covariates respectively, are presented in Figure 5.8. These models were used to estimate the impact of adjustment for additional covariates. Note that obtaining a completely unadjusted model is not possible, as age was used in the Cox model as the time scale.

5.3 - Results

Adjustment for additional covariates beyond age and sex had a limited impact on the observed effect estimates, with the exception of the probable AD outcome. In this case, adjustment for the full set of covariates attenuated to the null the protective effect observed when adjusting only for age and sex.




Figure 5.8: Results of three models adjusting for a different set of covariates. Note that the x-axis cutoffs (0.5,2) are different compared to other plots (0.3,3) to enable greater comparison between the different models.

Sensitivity cohorts: Entry year

When stratifying based on year of entry to the cohort, I observed no variation in risk by time period in any subgroup except for probable Alzheimer's disease (Figure 5.9).




Figure 5.9: Analysis of any lipid-regulating agent on probable AD outcome, stratified by grouped year of cohort entry.

On the assumption that this variation could be caused by changes in the frequency of probable AD diagnoses in the cohort over time, I performed a *post-hoc* investigation of the frequency of each dementia outcome by year of entry (Table 5.6). While the frequency of outcomes declines in more recent strata, likely due to the limited follow-up inherent to these groups, this decline in frequency is relatively constant

5.3 - Results

across the dementia subtypes.

Table 5.6: Frequency of diagnoses by grouped year of cohort entry

Year of cohort entry	No dementia	Probable AD	Possible AD	Vascular dementia	Other dementia	Total
<=2000	148550 (95.9%)	1775 (1.1%)	1677 (1.1%)	1345 (0.9%)	1585 (1.0%)	154932
2001-2005	613009 (96.3%)	6672 (1.0%)	5711 (0.9%)	4857 (0.8%)	6073 (1.0%)	636322
2006-2010	523913 (98.1%)	3355 (0.6%)	2169 (0.4%)	1890 (0.4%)	2506 (0.5%)	533833
2010<	357262 (99.4%)	845 (0.2%)	397 (0.1%)	374 (0.1%)	599 (0.2%)	359477
Total	1642734 (97.5%)	12647 (0.8%)	9954 (0.6%)	8466 (0.5%)	10763 (0.6%)	1684564

Sensitivity cohorts: Pregnancy

In the second sensitivity cohort, removing patients who may have been pregnant (coded as aged 55 and under at index) from the analysis had minimal effect on the effect estimates (Figure 5.10).




Figure 5.10: Comparison of analysis using main cohort and a cohort with potentially pregnancy women removed.

Statin properties

In the cohort, statins with lipophilic properties were much more frequently prescribed than hydrophilic statins (Table 5.7). Additionally, there is evidence for an increasing tendency to favour hydrophilic statins in recent years with the proportion of lipophilic statins prescribed falling from 18.2% in 1996-2000 to <1% in 2011-2016.

Table 5.7: Summary of statin properties (lipophilicity vs hydrophilicity) by grouped year of prescription.

Prescription Year Group	Hydrophilic	Lipophilic	Total
<=2000	7037 (18.2%)	31531 (81.8%)	38568
2001-2005	21427 (10.3%)	187018 (89.7%)	208445
2006-2010	3566 (1.6%)	217726 (98.4%)	221292
2010<	1115 (0.9%)	119035 (99.1%)	120150

When stratifying by statin properties, hydrophilic statins were less harmful in the any, vascular and other dementias outcomes compared to lipophilic statins (Figure 5.11). Additionally, in the AD outcomes, hydrophilic statins were associated with a small reduction in risk, compared to the weak evidence for an effect for lipophilic statins.




Figure 5.11: Analysis stratified by statin properties (hydrophilic vs lipophilic)

Impact of dementia code lists

When using the Smeeth *et al.* code lists to define dementia outcomes, effect estimates of HR: 1.19 (95%CI: 1.07-1.32) and HR: 1.33 (95%CI: 1.26-1.42) were obtained for the Alzheimer's disease and non-Alzheimer's ("other") dementia outcomes, respectively. While direct mapping to the outcomes used in this analysis was

not possible, the most comparable are probable Alzheimer's disease (HR:0.98, 95%CI:0.94-1.01) and other dementia (HR:1.19, 95%CI:1.15-1.24).

5.4 Discussion

5.4.1 Summary of findings

Lipid-regulating agents showed little evidence of an association with probable and possible Alzheimer's disease when compared to no treatment, but were associated with increased risk of an all-cause dementia, vascular dementia and other dementias diagnoses. The estimate observed in each case was driven by the effects observed statin subgroup, as a substantial majority of participants prescribed an LRA were prescribed a statin. For the other drug classes, no association was found with any outcome, with two exceptions. Ezetimibe was associated with increased risk of vascular and other dementias, while fibrates were associated with increase risk of all-cause dementia and probable Alzheimer's disease.

The effect estimates were robust to the exclusion of potentially pregnant participants, and for all outcomes except probable AD, no variation across grouped year of entry was observed. When looking at the statin subgroup alone, statin properties appeared to have a modifying effect, with hydrophilic statins being less harmful in the any, vascular and other dementias outcomes compared to lipophilic statins.

5.4.2 Interpretation of results

This section will expand on a potential explanation for the observed results detailed above. However, as the comparison of evidence across different sources is the aim of the triangulation exercise presented in later chapters, a detailed comparison with other published literature will not be provided here, except where needed to illustrate

a methodological point. For a detailed comparison of the result presented above with the existing evidence base identified by the systematic review, see Chapter 7.

Confounding by indication

A likely explanation for the observed increased risk of dementia outcomes with a vascular component (e.g. vascular and other dementias) with lipid-regulating agent use, and an important limitation of this analysis, is residual “confounding by indication”. While the term has been used to describe different sources of bias in epidemiological analyses,³¹² it is used here to described the role of risk factors that both prompt treatment and increase the risk of the outcome, thus causing a distorted positive association between the treatment and outcome (see Figure 5.12). In causal inference nomenclature, statins and dementia are said to be *d*-connected, as there is an open “backdoor” path between them via the uncontrolled confounders.³¹³ In the context of this analysis, this means a confounding variable (or, more likely, variables) both prompts prescription of statins and also represents a risk factor for the development of the vascular dementia. A similar confounding structure likely exists for ezetimibe, another hypercholesterolemia treatment, providing an explanation for the association of vascular/other dementia but not Alzheimer’s disease with this drug.




Figure 5.12: Causal diagram (directed acyclic graph) illustrating how confounding by indication could induce a positive association between statins and vascular dementia.

Conditioning entry into the study on being either “at-risk” or already diagnosed with hypercholesterolemia was employed in a pre-emptive attempt to mitigate confounding by indication, but evidence from the control outcomes suggests this was unsuccessful. The slight harmful effect for the backpain outcome is substantially smaller than that observed for the ischemic heart disease outcome, indicating that the majority of the uncontrolled confounding is likely related to vascular factors. The slight effect observed for the negative control of backpain could be due to incomplete control for socioeconomic status, as deprivation data was provided in quintiles to preserve patient privacy.^{314,315}

In line with this, an increasingly harmful effect is observed when moving from the probable and possible Alzheimer’s disease outcomes to the other dementia outcome, and finally to the vascular dementia outcomes. This pattern suggests that the strength of the residual confounding by indication increases as the proportion of cases with a vascular component in an outcome definition increases. Given confounding related to vascular factors, this pattern is also expected given the decision tree for assigning outcomes in the presence of greater than one dementia code. Under this system, the Alzheimer’s disease outcomes require a “pure” condition and the presence of any vascular or other dementias codes excludes participants from this group (Figure 5.1).

A review of other available literature suggests that this observation (a harmful effect of lipid-regulating agents on vascular-related outcomes) is not unusual. Using a conventional epidemiological technique, a previous analysis also found an increased risk of coronary heart disease (analogous to the ischemic heart disease outcome used in this analysis) in those taking statins (HR: 1.31, 95%CI: 1.04-1.66).³¹⁶ Controlling for confounding by indication in that study through the use of a trial emulation analysis gave an estimate of 0.89 (95%CI: 0.73-1.09), a comparable though less precise estimate to that observed in RCTs of statin use (0.73, 95%CI: 0.67-0.80).³¹⁷

Given the absence of vascular dementia in the published literature, as highlighted in the previous chapter, the unexpected increase in vascular dementia risk with statin use is particularly interesting. It is possible that this finding reflects publication bias if previous research encountered similar methodological issues to this analysis, and these results did not make it into the evidence base as unexpected or assumedly incorrect results are less likely to be submitted or accepted for publication.

Statin properties

This analysis found that hydrophilic statins were less harmful in the any, vascular and other dementias outcomes compared to lipophilic statins, and were associated with a small reduction in the risk of the probable and possible AD outcomes. The relative precision of the estimates in each group is expected, as the two most commonly prescribed statins are lipophilic (simvastatin and atorvastatin).³¹⁸

A widely discussed concept in the literature surrounding statin use and cognitive outcomes is the fact that lipophilic statins are more likely to be able to cross the blood brain barrier, and so have a more potent protective effect by directly lowering brain cholesterol.³¹⁹ My findings that hydrophilic statins appear to be more protective/less harmful than their lipophilic counterparts runs counter to this assertion.

An initial interpretation of the different associations observed in the two groups was that the lipophilic statins may be more potent, and so are prescribed to patients with

a higher underlying vascular load, leading to increased confounding by indication in this group. However, the statin with the strongest lipid lowering effect that is available via the NHS, rosuvastatin, is hydrophilic.

Impact of code lists

As part of a sensitivity analysis exploring the impact of outcome code-lists, I used definitions for Alzheimer's disease and other dementias obtained from a previously published paper (Smeeth *et al.*).²¹³ Using these lists in my analytical set-up, I found a harmful association of statin use with both outcomes.

This finding disagrees with the results of the original analysis, which found evidence for a protective effect of statin use on all-cause dementia (HR: 0.81, 95%CI: 0.69-0.96) and non-AD dementia (HR: 0.82, 95%CI: 0.69-0.97), but little evidence of an effect on AD (HR: 0.81, 95%CI: 0.49-1.35).

However, comparison of the results obtained using the two sets of code lists was deemed less useful following a detailed comparison of the codes used. While all of the codes used to define Alzheimer's in the Smeeth *et al.* paper are included in the probable Alzheimer's code-list (see Figure 5.13), I included several additional codes used to define this outcome (including, for example, "Eu00013: [X]AD disease type 2"). Additionally, several of the codes used to define "Possible Alzheimer's" in this analysis are included in the "Other dementia" code list used by Smeeth.




Figure 5.13: Sankey diagram comparing the codes used in this analysis with those used in the Smeeth *et al* paper.²¹³ The outcomes and number of codes contributing to each are presented (the Smeeth *et al.* outcomes are on the right hand side of the figure). The joining lines showing the overlap between the categories in the two analyses.

This analysis serves to illustrate the importance of the code lists chosen to define the outcomes of interest, particularly if they are used to define competing outcomes (e.g. AD vs non-AD dementia). This different codes used by Smeeth *et al.*, in addition to an analytical approach that adjusted for covariates defined after the index date, may go some way to explaining why my analysis obtained different results despite the substantial overlap in the data sources used.

5.4.3 Strengths and limitations

The primary strength of this analysis is the relative size of the CPRD and length of follow-up. Having reviewed the existing literature, as identified by the systematic review element of this thesis, this analysis of 1,684,564 participants is one of the largest available studies of this research question. Additionally, this analysis followed LRA users and non-users from a common index date, using a time-updating

5.4 - Discussion

treatment indicator to correctly assign time-at-risk to the exposed and unexposed groups. This approach has been less commonly used in the literature and allows for the mitigation of potential immortal time bias. Finally, it is one of the few that provide evidence on the association of lipid-regulating agent use and vascular and other dementias, and used negative and positive controls to assess the potential for residual confounding.

However, the findings of this analysis are subject to several limitations in addition to the confounding by indication discussed above. There is a strong possibility of differential misclassification of dementia-related conditions based on the exposure.³²⁰ As an illustrative example, those with memory complaints may be more likely to be classified as vascular dementia than Alzheimer's disease if their medical records contain prescriptions for lipid-regulating agents. Further, there is potential for general non-differential misclassification of the outcome due to the varying positive predictive value of electronic health record code lists to identify dementia cases.^{278,279}

Misclassification of outcomes is not the only issue introduced by the use of EHR codes to define outcomes. Comparing and contrasting between different studies is particularly difficult because of the impact that the use of different code lists can have on the analysis. This problem is illustrated by the discrepancy between the results when using the code lists defined for this study and those used by Smeeth *et al.* This presents a particular challenge in comparing research across different time-periods and coding systems, particularly the unexpected results I obtained for the vascular and other dementias outcomes.

A further limitation stems from the possibility of uncontrolled confounding due to genetic factors. The number of *ApoE4* alleles represents the strongest genetic risk factor for Alzheimer's disease, but also substantially increases LDL cholesterol levels,³²¹ potentially prompting treatment with a statin or other lipid regulating agent. I was unable to control for *ApoE* genotype in this analysis as I did not have access to genetic data on participants. As a result, any protective association

between LRA use and the Alzheimer's disease outcomes may be masked by residual negative confounding by *ApoE*.

Finally, as with many studies of dementia, there is a risk of reverse causation in my analysis. Dementia and associated conditions have a long prodromal period, during which preclinical disease could cause indications for the prescription of a lipid-regulating agent. Enforcing a minimum period of follow-up would address this limitation, but the use of a time-varying treatment indicator in this analysis prevents the use of this approach (see Appendix A.5.2 for a fuller discussion of this topic).

5.4.4 Enabling easy synthesis of this analysis

The raw data supporting this analysis is not publicly available, as access to the CPRD data is controlled by a data monitoring committee. However, when data are not readily available, sharing the analysis code and summary statistics represents a way for readers to validate the findings.¹³⁶

In light of this and my own experiences in attempting to extract information for papers assessing preventative treatments, as documented in Section 4.5.5, the outputs from this analysis have been made readily available. All code, Read code lists and summary statistics (namely the tables presented in this chapter plus summary tables of effect estimates) can be downloaded in a machine readable format from the archived repository for this project (Zotero DOI: **TBC**). This open approach should enable easy inclusion of this analysis in future evidence synthesis exercises, allowing new work to build on that presented here.

5.4.5 Summary

- In this chapter, I produced new evidence on the association of lipid-regulating agents with incidence of all-cause dementia, Alzheimer's disease, vascular

dementia, and other dementia.

- I found little evidence for an effect of lipid-regulating agents on probable or possible Alzheimer's disease. However, lipid-regulating agent use was associated with an increased risk of all-cause, vascular and other dementias. In all cases, the estimated associations for the any LRA analysis were largely due to those observed in the large statin subgroup.
- I attempted to account for important sources of bias through use of time-varying treatment indicator, though the control outcomes included in the analysis provided evidence for only partially controlled confounding by indication related to vascular factors. Additionally, there was the potential for differential misclassification of dementia subtype on the basis of the exposure. Combined, these biases reduce the confidence in my findings, in particular the unexpected increase in risk of vascular dementia associated with statin use.
- Findings from this analysis are used as an additional source of evidence in the triangulation exercise presented in Chapter 8.

These particular systematic reviews [individual participant data reviews] remain yardsticks against which the quality of other reviews continues to be judged.

— Ian Chalmers, 1993³²²

6

Individual participant data meta-analysis of blood lipid levels and dementia outcomes

Contents

Lay Summary	150
6.1 Introduction	150
6.2 Methods	151
6.3 Results	156
6.4 Discussion	168
6.5 Summary	175

Lay Summary

Individual participant data (IPD) meta-analyses are a special type of meta-analysis where those undertaking the study contact the authors of relevant studies to request access to the raw data on each participant. This is different to the meta-analyses presented in the previous chapter, which used published summary results.

To examine the relationship between lipid levels and dementia risk, I applied for access to 37 unique data sources containing information on these variables. However, only a small proportion of these data sources ($n = 3, 8.1\%$) provided the requested access.

The resulting analysis of these data sources did not suggest a relationship between any blood lipid and any dementia outcome. The sole exception was an increased risk of vascular dementia in those with higher triglyceride levels. In addition, the participants age and sex did not appear to influence the relationship of lipid fractions and dementia outcomes.

The reasons for the low response rate to requests for data are explored. Finally, I suggest future studies to investigate how data access rates could be improved.

6.1 Introduction

Individual participant data (IPD) meta-analyses are considered to be the gold standard form of evidence synthesis, allowing for the application of a common selection model and analytical approach across all identified cohorts.⁹² They are particularly useful when investigating the impact of participant-level characteristics, something that is not possible with aggregate data unless the results are stratified by the characteristic of interest.^{92,323} Knowledge of which groups a treatment will benefit most (or harm least) is a core aim of the move towards personalised medicine.^{98,324} IPD analysis also offer a mechanism by which previously unanalysed datasets can be incorporated into an analysis, thus expanding the evidence base for a particular research question.

Previous work has suggested a difference in the effect of lipids on dementia risk based on participant age and sex,^{59,220} though the systematic review presented in Chapters 3 & 4 did not identify strong evidence for an effect of sex or age at lipid measurement on the relationship between lipids and dementia. However, the number of included studies in the lipid fraction meta-analyses was small, due to both the relatively small number of studies reporting on the lipids exposure and the poor reporting of summary statistics of participant characteristics in those that did. Additionally, best practice guidance recommends against basing the decision to perform an IPD meta-analysis on the between-study heterogeneity in a summary data meta-analysis, as similar distributions of participant covariates across studies may mask a true effect.⁹⁸

As such, the aims of this analysis are two fold: firstly, to perform an IPD meta-analysis across identified cohorts to examine the impact of participant age-at-measurement and sex on the effect of lipids on dementia risk; and secondly, to expand the evidence base for the effect of lipids on dementia outcomes by obtaining estimates from previously unanalysed cohorts available via the Dementia Platform UK, a large consortia of dementia cohorts.

6.2 Methods

6.2.1 Eligibility criteria

Study design

Eligible data sources for this analysis were prospective cohort studies. Data sources which were cross-sectional, either by design or due to the available data (e.g. a study conducted across multiple waves, but only data from a single wave could be accessed) were excluded. Similarly, studies making use of population-level electronic health records, which often require an extensive project proposals in order to gain access to the data, were ineligible due to the time and cost involved in

applying. No restrictions were put on the number of participants or the length of follow-up, though participants must be free of dementia (or assumed to be free, based on age at entry) at baseline.

Exposures/outcome definition

I considered four blood lipid fractions as part of this analysis, namely: total cholesterol (TC), low-density lipoprotein cholesterol (LDL-c), high-density lipoprotein cholesterol (HDL-c) and triglycerides (TG). Cohorts were eligible for inclusion if they contained data on at least one of these in a continuous format (i.e. studies with binary “hypercholesterolemia” would be excluded).

In line with the analyses presented in other chapters, eligible cohorts were those containing data on at least one outcome of interest, namely: all-cause dementia, Alzheimer’s disease, or vascular dementia.

6.2.2 Applying for data access

Potentially eligible data sources as defined above were identified via two mechanisms, each with a distinct focus. For both approaches, the number of cohorts responding to the request for data access, and where applicable, the reasons given for a refusal were recorded.

Cohorts identified by the systematic review

The first approach focused on previously analysed observational prospective cohort studies examining the effect of blood lipid levels on dementia outcomes, as identified by the systematic review presented in Chapters 3 & 4. The data sources used in each of these cohort analyses were screened against the criteria listed in the previous

section, and eligible cohorts were approached for data access. In the first instance, the first/corresponding author of the publication was emailed in Autumn 2020. If this approach did not elicit a response within two months, the last author was contacted, on the basis that the first/corresponding author may have been a more junior member of the research group who had moved to a different institution.

Cohorts contained in Dementia Platform UK

The second approach focused on incorporating relevant, previously-unanalysed data into the analysis, thus providing additional evidence on the relationship between blood lipids and dementia risk. This was achieved through the Dementia Platform UK (DPUK), a collaborative grouping of existing dementia cohorts established by the MRC which works with data owners to make their data readily accessible for secondary analysis.¹⁰⁸ It provides access to 42 cohorts with over 3 million participants, and makes use of a central streamlined application processs for all cohorts with the intent of making it easier to access data from existing data sources.

Cohorts included in the DPUK were assessed against the eligibility criteria, and in Autumn 2020, an application for access to a subset of 17 cohorts was made via the common-access procedure.

6.2.3 Primary analysis

Data cleaning and harmonisation

Where data on one of these exposures (TC, LDL-c, HDL-c, TG) was missing, it was inferred from the other three fractions (where available) using the Friedwald formula (Equation (1.1)). Lipids levels were retained as continuous variables rather than dichotomising into a binary hypercholesterolemia exposure, given the additional

6.2 - Methods

statistical power this adds to the meta-analysis.^{98,325} Where lipid measurements were reported in *mg/dL*, these were converted to *mmol/L*.

Across all cohorts, data cleaning was performed in a similar manner, standardizing to commonly named variables so that a single model could be applied using functional programming.³²⁶ The advantage of this approach is that it reduces the likelihood of errors in model mis-specification if changes are required in variables names from cohort to cohort. Following data cleaning, summary statistics for each data source were calculated and compared with publicly available statistics to ensure no errors were introduced in the data cleaning process, in line with best practice.³²⁷

Covariate definition

A range of additional variables were included in the analysis, intended to address the potential for confounding. With an awareness that discrepancies in available covariates is a common issue in IPD studies, I defined an idealised set of covariate domains to be age, sex, education, BMI, *ApoE4* status, smoking/alcohol status, ethnicity, and prevalent diabetes or cardiovascular disease. This set covers key risk factors for dementia/Alzheimer's disease in addition to general cardiovascular risk factors. Details on how these variables were coded, given the available data, are presented in Section 6.3.3.

Missing data

Missing data in this analysis was classified as either relating to missing values (a cohort collected data on a variable of interest, but some values are missing) or missing variables (a cohort did not collect data on a variable of interest). Variables with missing values were identified, and 20 imputed datasets were created.²⁹⁷ Imputation was performed using the MICE (Multiple Imputation by Chained Equations) using the `mice` package in R.

Missing variables were originally intended to be addressed using a previously described method.³²⁸ Here, the correlation between the fully-adjusted and partially-adjusted estimate in cohorts with the full set of covariates is used to estimate the fully-adjusted effect in those cohorts missing covariates. However, this method requires several large cohorts to contain the full set of covariates, a condition this analysis failed to meet given the very low response rate. As such, two other common approaches were employed.³²⁸ In the primary analysis, all cohorts were analysed adjusted for the set of common covariates across cohorts (Model 1: age, sex, smoking, alcohol, diabetes). As a sensitivity analysis, cohorts with the full complement of covariates (Whitehall II and EPIC) were analysed using a maximally-adjusted model (Model 2: Model 1, further adjusted for education, ethnicity, and prevalent ischemic heart disease and BMI). Results between the common-set-adjusted (Model 1) and maximally-adjusted (Model 2) models were then compared.

IPD analysis

In this analysis, a two-stage IPD analysis was planned. Under a two-stage approach, estimates for the effect of each lipid fraction on incident dementia were first calculated for each data source. More specifically, a logistic regression model adjusted for relevant covariates as detailed in the above section was used to quantify the effect of a 1 mmol/L increase in each lipid fraction on each dementia outcome. Results were expressed as odds ratios (OR). Examination of the data available via the DPUK indicated that the common time-to-event approach used in studies of dementia outcomes would be precluded by the absence of detailed time-to-event data. An overall effect estimate was then produced by combining data-source-specific estimates in a random-effects meta-analysis (see Section 3.2.9 for a broader discussion of meta-analysis methods).

A two-stage IPD approach was employed for a number of reasons. Firstly, and most importantly, a two-stage approach allows for siloed data, enabling researchers

who are unwilling/unable to provide their data to analyse themselves following a specified analysis plan.⁹² Secondly, it allows for the production of forest plots of within-cohort estimates, something which is useful for the triangulation exercise in Chapter 7. Finally, a two-stage approach is simpler to model, as it uses standard well-documented summary effect estimate meta-analysis techniques, and automatically accounts for methodological issues such as clustering within cohorts³²⁹ and the potential for ecological bias.³³⁰

Investigating the effect of participant-level covariates

In order to investigate the interaction of participant-level characteristics (age and sex) with lipid levels, interaction terms for lipid-covariate terms were extracted and synthesised using a fixed effects meta-analysis.³³¹ To avoid ecological bias, where the between-study association does not reflect the within-study associations, cohorts where there was no within-study variation in the covariate of interest were excluded from the interaction analysis for that covariate.³³⁰ As an example, it is impossible to estimate the impact of sex on the lipid~dementia relationship in a study which contains only female participants. All interaction analysis were performed using the common-set-adjusted model (Model 1).

6.3 Results

6.3.1 Data access

Of the 37 studies to which I applied for data access, only three (8.1%) were included in the final analysis. Figure 6.1 details whether the cohorts eventually included in the review were identified by the systematic review or via DPUK portal. In addition, the reasons for cohorts not being included in the analysis are presented, where available, stratified by application approach.

In summary, the requests for data from cohorts identified by the systematic review were characterised by a very low response rate ($N = 5$, 25%). For the minority who did respond, common reasons given by authors for not sharing the data included that they: no longer worked with the same group and did not know if the data was available or how to obtain it; no longer had access to the data; or were currently performing, or intended to perform, a similar analysis as the one proposed.

With respect to the application to DPUK cohorts, where a dedicated project manager liaises with data owners on the applicants' behalf, the overall response rate was higher. However, even using this streamlined approach, a positive response was obtained for approximately half ($N = 9$, 53%) of the approached cohorts.




Figure 6.1: Flowchart of included cohorts, stratified by identification method (systematic review vs DPUK).

6.3 - Results

As highlighted in Figure 6.1, there was little overlap between cohorts identified by the systematic review and those contained in the DPUK (1), indicating that the DPUK is a useful source of unanalysed data with respect to this question. A single cohort (Whitehall II) was identified by the systematic review and was also present in the DPUK.

Nine cohorts (8 DPUK only, 1 Systematic review + DPUK) responded positively to requests for data access. However, on inspection of data provided, six of these cohorts were excluded and the reason for exclusion in each case is summarised in Table 6.1. In summary, two cohorts contained were memory-complaint cohorts, containing participants unlikely to be free of dementia at baseline (BRACE, MEMENTO). Two cohorts, despite being designed as multi-wave studies, only shared data related to a single wave and so represented cross-sectional data (Generation Scotland, NICOLA). Finally TRACK HD is a genetic cohort where cohort owners advised me that dementia will inevitably develop in mid-life driven by the Huntington's (HTT gene) mutation and that this effect would likely far outweigh the effect of lipids. Finally, the ELSA study was excluded based on the dichotomous definition of the exposure variable.

Table 6.1: Exclusion reasons for cohorts providing data

Cohort	Reason
BRACE	Memory complaint cohort - unlikely to be dementia free at baseline
ELSA	Lipid exposure reported as a binary variable
Generation Scotland	Cross-sectional - only one wave of data available
MEMENTO	Memory complaint cohort - unlikely to be dementia free at baseline
NICOLA	Cross-sectional - only one wave of data available
TRACK HD	HTT gene carriers (i.e. premanifest Huntington's Disease)

6.3.2 Included data sources

The three data sources used in this analysis are described in detail in the following sections. Of note, all included data sources were based in the United Kingdom. This is due to the majority of included datasets being identified via the Dementia Platform UK route (Figure 6.1), which as implied by the name, has a narrow geographical focus.¹⁰⁸

Caerphilly Prospective Study

The Caerphilly Prospective Study (CaPS) is a longitudinal study of men in South Wales, UK.^{332,333} Blood lipids (TC, LDL-c, HDL-c and TG) were measured at baseline in 1979-1983, and from Phase III (1989-1993) onwards, a battery of cognitive tests were introduced. A dementia was derived using data obtained during Phase V (2002-2004), giving between 19-25 years of follow-up. The cohort contains data on

dementia outcomes sub-classified as vascular and non-vascular dementia. Data was available for all covariates of interest except for education, BMI and *ApoE4* status.

Epic Norfolk

The European Prospective Investigation of Cancer (EPIC) - Norfolk is a population-based cohort, containing men and women recruited from 35 general practices in Norfolk between 1993 and 1998.^{334,335} Data provided by this cohort for the analysis, dementia was ascertained at the 5th Health Checkup (2016-2018), giving between 18-25 years of follow-up. The added evidential value of the EPIC cohort is small, given the fact that the data obtained contains only 8 dementia events. The cohort contains no information on dementia subtype, while all covariates of interest bar *ApoE4* were available.

Whitehall II

The Whitehall II study is a prospective cohort study of men and women recruited between 1985 and 1989 from the civil service in London.³³⁶ Lipids measurements were available from the third wave of the study, conducted between 1991-1993. The cohort is linked with the Hospital Episode Statistics (HES) database, a database containing details of participant events at NHS hospitals in England, which was used to define covariates at baseline and dementia outcomes.³³⁷ HES data for dementia outcomes was available up to March 2015, providing between 22-24 years of follow-up. Information was available on dementia type, which was classified as all-cause dementia, Alzheimer's disease or vascular dementia. This data source contained details on all covariates of interest except for *ApoE4* status.

Of note, the Whitehall II cohort was analysed in one of the included studies identified by the systematic review presented in Chapters 3 & 4,²⁴⁵ meaning that a comparison between the published result and the analysis reported here was possible.

6.3.3 Covariate definition & missing data

Based on available data across cohorts, smoking was classified as never/ever/current, while alcohol consumption was classified as never/ever. Education was categorised into 4 levels (None, O-levels, A-levels, Degree). BMI was treated as a continuous variable while presence of vascular co-morbidities was treated as dichotomous.

A key consideration in the definition of covariates across cohorts was the classification of age. The Whitehall II study, the largest cohort to which I had access, only shared age data in five-year age band (e.g. 40-44, 45-49, etc.). To ensure comparability across the cohorts, I created identical categories in the CaPS and EPIC data. This grouped age variable was then used in all subsequent analyses.

Missing values in collected variables was common across the cohorts. A matrix of covariates, and whether they were not available or contained missing values for each included cohort, can be seen Appendix A.6.1. The Whitehall II and EPIC cohorts contained data on all but one covariates of interest, while three missing covariates were identified in the CaPS cohort, namely education, prevalent ischemic heart disease, BMI and ethnicity. No included cohort provided information on the *ApoE4* status of participants, and so this variable could not be adjusted for in the analysis.

6.3.4 Analytical results

Descriptive statistics

Across the three cohorts, 11,835 participants were included in this analysis (Whitehall II = 8208, EPIC = 1115, CaPS = 2512). All cohorts contained data on the four exposure variables of interest (or sufficient data from which to calculate them) and on all-cause dementia outcomes. The only other dementia outcome examined across cohorts was vascular dementia, which was reported in the CaPS

6.3 - Results

and Whitehall II studies. The definitions of dementia outcomes used across cohorts can be seen in Appendix A.6.2. Cumulatively, there were 542 cases of all-cause dementia, with 114 further classified as vascular dementia. Summary statistics for each cohort are provided in Table 6.2.

Table 6.2: Summary of characteristics for cohorts included in the IPD analysis

	Whitehall II	Epic	CaPS
N	8208	1115	2512
Age group, Median	45-49	50-54	50-54
Male, N (%)	5679 (69.2)	495 (44)%	2512 (100)
BMI, Mean (SD)	25.3 (3.75)	25.8 (3.58)	-
Education >18 yrs, N (%)	2576 (31.4)	252 (22.6)	-
Smoke ever, N (%)	2678 (32)	490 (43)	2110 (83.9)
Alcohol ever, N (%)	6612 (80.6)	37 (3.3)	2235 (89)
Ethnicity (white), N (%)	7387 (90)	1107 (99.3)	-
IHD, N (%)	7 (0.1)	63 (5.7)	-
TC, Mean (SD)	6.49 (1.16)	5.93 (1.08)	5.93 (1.15)
LDL-c, Mean (SD)	4.40 (1.04)	3.77 (0.96)	3.74 (1.12)
HDL-c, Mean (SD)	1.43 (0.414)	1.43 (0.41)	1.37 (0.41)
TG, Mean (SD)	1.49 (1.14)	1.67 (1.04)	1.83 (1.21)
Dementia, N (%)	287 (3.5)	8 (0.7)	247 (9.8)
Alzheimer's, N (%)	92 (1.1)	-	-
Vascular dementia, N (%)	37 (0.5)	-	77 (3.1)
Non-vascular dementia, N (%)	-	-	170 (6.8)

Abbreviations: BMI - Body mass index; IHD - Ischemic heart disease; HDL - Low-density lipoprotein cholesterol; LDL - Low-density lipoprotein cholesterol; TG - Triglycerides.

Main effects

The results from the main effect analysis across the varying lipid fractions on each dementia, considered can be seen in Figures 6.2 & 6.3, respectively. There was weak evidence for an association of any lipid level with either all-cause or vascular dementia, with the exception of a harmful association between raised triglycerides and vascular dementia (OR: 1.22, 95%CI: 1.11-1.32).




Figure 6.2: IPD meta-analysis (Model 1) of all-cause dementia outcomes, stratified by lipid fraction

6.3 - Results




Figure 6.3: IPD meta-analysis (Model 1) of vascular dementia outcomes, stratified by lipid fraction. Note that the vascular dementia outcome was only available in the CaPS and Whitehall II cohorts.

Estimates from the common-set-adjusted model (Model 1: adjusted for age, sex, smoking, alcohol, and diabetes) were comparable to the fully-adjusted model (Model 2: Model 1 further adjusted for ethnicity, prevalent ischemic heart disease, education and BMI) for the effect of lipids on all-cause dementia in cohorts reporting all covariates of interest (Figure 6.4).




Figure 6.4: Comparison of results from the common-set-adjusted (Model 1) and fully-adjusted (Model 2) analyses for the all-cause dementia outcome.

Interaction effects

Given minimal effect of further adjustment for education and BMI on the effect estimates, the results from the interaction analysis presented here were obtained using the common-set-adjusted model. Further, while the maximally-adjusted model (Model 2) could have been employed in the analysis of the effect of sex, as the CaPS cohort contains only men and had to be excluded from this analysis, employing the same underlying model helped to ensure comparability between the age and sex interaction estimates.

6.3 - Results

For all-cause dementia, there was no evidence of an interaction of lipid levels with either age group or sex (Figures 6.5 & 6.6). Note that for the sex interaction analysis, the CaPS cohort was excluded as it contains only a single sex and therefore provides no information on exposure-sex interaction.




Figure 6.5: Meta-analysis of interaction terms representing the effect of a one-step increase in age group on the lipid/all-cause dementia association

6.3 - Results




Figure 6.6: Meta-analysis of interaction terms representing the effect of male gender on the lipid/all-cause dementia association

For the consideration of vascular dementia, I was only able to explore the effect of age, as only the CaPS and Whitehall cohorts contained details on vascular dementia as an outcome. As discussed above, the CaPS data contained a single sex excluding it from a exposure-sex analysis, leaving Whitehall II as the sole eligible study.




Figure 6.7: Meta-analysis of interaction terms representing the effect of a one-step increase in age group on the lipid/vascular dementia association

6.4 Discussion

6.4.1 Summary of findings

This analysis requested data from 37 data sources, but only obtained data from three, all of which were based in the United Kingdom. No evidence for an effect of lipids on the risk of dementia or related outcomes was identified, except for a harmful association of raised triglycerides with risk of vascular dementia. Similarly, there was weak evidence for an interaction of the effect of lipids levels on dementia outcomes with participants' age, grouped into 5 year bands, or sex.

A detailed comparison of the findings presented above with the existing evidence

base identified by the systematic review (Chapters 3 & 4), is presented as part of the triangulation exercise in Chapter 7. As such, this discussion will not provide a detailed comparison of the results of this analysis with other published literature, except to compare between this analysis and previously published results using the same data source.

6.4.2 Limitations

Low response rate to request for data

The obvious key limitation of this analysis is the very low response rate to requests for data access, which may bias the results if there are systematic differences in the association of lipids with dementia outcomes between cohorts that share data and those that do not.³³⁸ Whether or not to press ahead with an IPD analysis in the absence of all (or even most) data is a personal decision, and some previous analyses have highlighted where they decided not to pursue an IPD analysis.³³⁹ For the purposes of this thesis, the decision was made to conduct the IPD analysis, as it provided training in application of the methods in addition to providing additional evidence sources that could be incorporated into the triangulation exercise detailed in Chapter 7.

A low response rate is not unexpected, given that a review of IPD studies published between 1987 and 2015 found that fewer than half managed to obtain data from greater than 80% of studies, and that in many cases, the exact percentage of studies for which data was obtained was not accurately reported.¹⁰¹ However, it is assumed that a ~10% response rate is at the lower end of the scale.

There are likely several likely reasons for this low response rate to requests for data access. In general terms, there are several well-documented barriers that prevent data from being made readily available, including concerns regarding participant privacy, fear of “scooping” or “parasitic” behaviour, and a lack of trust between

6.4 - Discussion

primary and secondary researchers.¹⁰⁶ More specific to this analysis, IPD meta-analysis including studies other than randomised controlled trials have less success in obtaining IPD from studies.¹⁰¹ Additionally, while no evidence is available on whether the characteristics of the researcher requesting data access influences the response rate and eventual decision, there is the possibility that my position as a PhD student meant I was less likely to elicit a positive response than a well-known senior academic might. The timing of the requests for data access, coinciding with a global pandemic, may also have affected the response rate as researchers prioritised COVID-related work.

Finally, the method of contact used (email) has been shown to be less successful in eliciting responses from authors when compared with telephoning.³⁴⁰ One of the reasons for this may be that the email addresses reported on publications are more likely to be out of date for older publications. Anecdotally, post-hoc investigation of a subset of cohorts revealed that several corresponding/first authors were no longer at the same institution as when the study was reported, and as a result, were unlikely to have access to the institutional email address listed on the study publication.

The obstacles to data access described above are in theory what the DPUK was built to address. However, even with the help of the streamlined application process afforded by the DPUK, accessing sufficient data was a challenge. The response rate among DPUK cohorts a year after application was just 50%. In addition, some cohorts responded that that the proposed study question was already under investigation by another group, and that they would not share the data on this basis. In light of this, a centralised database of ongoing analysis being performed in DPUK would be of enormous help. Finally, the DPUK process would be aided by a clearer distinction between those cohorts that are “DPUK native” (i.e. where a copy of the data is already held on DPUK servers) versus externally hosted, as the response time for externally hosted cohorts is likely to be much longer.

Uncontrolled confounding

A key limitation of this analysis is the potential for uncontrolled confounding. Across all cohorts, adjustment for *ApoE4* was not possible as I did not have access to genetic data on participants. *ApoE4* is a strong risk factor for both increased LDL-c levels and Alzheimer's disease,^{321,341} and failing to adjust for this factor means residual confounding in the LDL-c fraction results is likely.

More generally, systematically missing variables meant that a trade-off between inclusion of cohort data and appropriate control for confounding was required, resulting in the potential for uncontrolled confounding due to education and BMI. However, sensitivity analysis comparing the common-set-adjusted (Model 1) and fully-adjusted (Model 2) models in cohorts with a full complement of covariates indicated that further adjustment for BMI and education had minimal impact on the effect estimates.

Comparison with previous analysis

For the single cohort (Whitehall II), a previously published analysis was available. Tynkkynen *et al.*²⁴⁵ analysed the association of blood lipids and risk of all-cause dementia and Alzheimer's disease across several cohorts, including the Whitehall II cohort.

In an attempt to validate my approach, I compared the results from the maximally-adjusted model (Model 2) in this analysis with those reported by Tynkkynen *et al.* for both all-cause dementia and Alzheimer's disease.

The results for the Alzheimer's disease outcome were comparable (Appendix A.6.3). However, for the vascular dementia outcome (Figure 6.8), I identified a discrepancy in the association of triglycerides with all-cause dementia estimated by the two analyses. Tynkkynen *et al.*²⁴⁵ found a protective effect of triglycerides on this

6.4 - Discussion

outcome (HR: 0.69, 95%CI: 0.56-0.85) while this analysis found evidence for a harmful effect (OR: 1.26, 95%CI: 1.12-1.41).




Figure 6.8: Comparison of results across the two analysis of the Whitehall II cohort for the all-cause dementia outcome.

Investigation revealed several possible reasons for this discrepancy. The first is that it was likely to be due to the Whitehall snapshot that each analysis had access to, as comparison of summary statistics indicated that this analysis had substantially more dementia events ($N = 287$ in this analysis vs. $n=114$ in Tynkkynen *et al.*). Additionally, this analysis uses multiple imputation to address missing data, while the Tynkkynen *et al.* does not mention how missing data was addressed, though the reduced precision in the estimates suggest a complete case analysis. Finally, the

covariates adjusted for in each analysis were similar, though the previous analysis had access to genetic data, allowing it to adjust for *ApoE4* status. However, given that *ApoE4* is a risk factor for increased LDL-c rather than triglycerides,³²¹ it seems unlikely that additional adjustment for this variable is responsible for the discrepancy in findings observed.

6.4.3 Strengths

While this analysis did not manage to systematically obtain and analyse a large proportion of identified data, a central strength of this analysis is the use of a systematic approach to identify and attempt to contact relevant cohorts. Further, it also enabled incorporation of two previously unanalysed datasets - the CaPS and EPIC Norfolk cohorts - thus providing additional evidence that is used in the triangulation exercise reported in Chapter 7.

A further strength of this analysis was the ability to investigate the effect of participant characteristics (namely age and sex) on the association of blood lipids and dementia outcomes. To the best of my knowledge, no previous review has examined the interaction of these factors with the observed associations.

Finally, this analysis provides new evidence on a previously unexplored outcome (vascular dementia), adding to the extremely small evidence base for this outcome identified by the systematic review presented in Chapters 3 & 4. This is relevant even in the previously analysed WhiteHall II cohort, as the previous analysis by Tynkkynen *et al.* did not report on this outcome.

6.4.4 Reflections on the process

In hindsight, attempting to undertake a large-scale IPD meta-analysis as part of a larger PhD project may have been overly ambitious. Data harmonization between

cohorts in an IPD analysis is an often under-appreciated challenge,³²⁷ and in line with this, data cleaning for this analysis took substantially longer than expected. While the cohort response rate was substantially lower than expected, given the time and resource of data cleaning and harmonisation for just three cohorts, a situation in which all 37 cohorts responded positively would have been logistically challenging within the scope of a PhD.

6.4.5 Future work

While it is tempting to suggest that an IPD analysis of lipid levels be reattempted, without empirically guided approaches to increase the response rate, this may just result in a similarly small set of studies as described here. In line with the limitations considered earlier in this chapter, future methodological work could formally consider the effect of requester characteristics (sex, location, career stage) on response rates in IPD analyses.

Additionally, the production of detailed guidance for handling cases where covariates are systematically missing. Much of the literature around IPD analysis is focused on the synthesis of RCTs, where additional covariate information is needed primarily for the assessment of treatment-covariate interactions rather than the adjustment of the effect estimate for confounding. Given the wide availability of non-randomised cohorts, improved guidance on this challenge would support future work.

Finally, a movement towards increased use of unique persistent identification of researchers, through schemes such as the ORCID programme,³⁴² would help with contact issues. Researchers move institutions regularly as their contracts come to an end, and so the institutional contact details provided on publications are frequently out of date.

6.5 Summary

- In this chapter, I performed an IPD meta-analysis to investigate the effect of blood lipid levels on the risk of incident dementia. There was a very low response rate to requests for data access, resulting in the analysis of three relevant cohorts.
- I found weak evidence for an association of any lipid with either all-cause or vascular dementia, except for an increased risk of vascular dementia associated with raised triglycerides. Similarly, there was weak evidence that the association of blood lipids and dementia outcomes varied by participant age or sex.
- I discussed potential reasons for the low response rate, and explored other limitations of this analysis. I highlighted the contribution of this work to the wider topic via the analysis of previously unexplored cohorts (CaPS & EPIC) and outcomes (vascular dementia). Finally, I recommended that future research formally investigate the impact of requestor characteristics on requests for data.
- The new evidence produced by this analysis will be incorporated, along with evidence identified or produced in the previous chapters, into the triangulation analysis presented in the following chapter.

Next on my list of features to be specially considered I would place the consistency of the observed association. Has it been repeatedly observed by different persons, in different places, circumstances and times?

— Austin Bradford Hill, 1965³⁴³

7

Aetiological triangulation across new and existing evidence sources

Contents

Lay summary	178
7.1 Introduction	178
7.2 Data sources	179
7.3 Qualitative (narrative) triangulation	182
7.4 Quantitative triangulation	185
7.5 Discussion	201
7.6 Summary	208

Lay summary

TBC

7.1 Introduction

Aetiological triangulation, or simply triangulation, is the process of comparing and contrasting across different sources of evidence. Triangulation is broadly comparable to Bradford-Hill's criteria of "consistency", that is the replication of an observed relationship across several different contexts,³⁴³ where contexts are assumed to have different underlying bias structures. More formally it can be defined as:

The practice of strengthening causal inferences by integrating results from several different approaches, where each approach has different (and assumed to be largely unrelated) key sources of potential bias.¹⁰⁹

This approach represents a significant step forward from the current practice of synthesising evidence from only one type of study (e.g. randomised controlled trials (RCTs), non-randomised studies of exposures (NRSE) or interventions (NRSI)). The most common implementation of this method to date has been in the form of *qualitative triangulation* - that is the identification and narrative comparison of diverse evidence sources with respect to their underlying bias structures.¹⁰⁹

However, qualitative triangulation faces issues at scale. There is a need to include all evidence to avoid potential confirmation bias,³⁴⁴ yet as the number of evidence sources increases, it is not possible to narratively compare and contrast across multiple results in any meaningful way. This fact is illustrated by previous exemplars of triangulation only considering a small number of individual results.^{109,345} To overcome this limitation, some attempts have considered the output of a meta-analysis of similarly-designed studies to be a single source of evidence, and have assessed bias in relation to this summary result. While this approach keeps the

number of individual sources of evidence manageable, it loses useful information on the specific biases inherent to each result contributing to the summary estimate. For example, while it may be true that all non-randomised studies share some minimal level of bias, competing extents and directions of bias in individual results may be masked by the use of summary estimates. In addition, previous attempts at qualitative triangulation have not systematically assessed the indirectness of the results, that is the differences between the question of interest and the question addressed by a study.

Taken together, these limitations to qualitative triangulation detail a need for a more systematic way to integrate across multiple evidence sources as the number of individual results contributing to the exercise increases. One such approach is *quantitative triangulation*, where results from different evidence sources are combined numerically while accounting for the key sources of bias and indirectness in each. This chapter, in addition to providing a narrative comparison of the existing and new evidence identified by this thesis, builds on recent developments in risk-of-bias assessment and methods for bias-/indirectness-adjusted meta-analysis to propose a generalised framework for quantitative triangulation. The framework, along with the methodological challenges to its implementation, is illustrated via two case studies examining the causal effect of lipids on dementia outcomes.

7.2 Data sources

The triangulation exercises presented in this chapter draw on the research produced in each of the preceding chapters. More specifically, this chapter builds on the comprehensive systematic review of existing evidence presented in Chapters 3 & 4. This evidence base is then supplemented by new evidence on the association of statin use with dementia outcomes in the CPRD (Chapter 5) and the association of lipids with dementia outcomes in previously unanalysed datasets accessed via the DPUK (Chapter 6).

7.2 - Data sources

Table 7.1 summarises the research question each data source has attempted to address, the exposures and outcomes considered in each case, and the contribution of each chapter to the triangulation exercise presented here.

Table 7.1: Summary of studies included in this thesis and used as evidence sources in this triangulation exercise. Note, Chapter 2 is intentionally not included in this table, as it describes a research tool rather than a research study.

Chapter	Research Question	Exposure/ Intervention	Outcome	Contribution to evidence synthesis framework
Chapter 3/4	Based on the available evidence; (i) are lipid fractions associated with subsequent dementia risk, stratified by subtype? (ii) Are lipid regulating agents associated with subsequent dementia risk, stratified by subtype?	Lipids (HDL-c, LDL-c, TC, TG),	Dementia, stratified by subtype	Provides overview of existing evidence, including detailed risk-of-bias assessments for each result
Chapter 5	Are lipid regulating agents associated with dementia risk in a large scale electronic health record database?	Seven classes of lipid regulating agents	Dementia, stratified by subtype	Provides additional observational data on vascular dementia (under-represented in the literature)
Chapter 6	Are lipid levels associated with dementia risk in previously unanalysed prospective cohort studies?	Lipids (HDL-c, LDL-c, TC, TG)	All-cause dementia VaD	Provides additional observational data on vascular dementia (under-represented in the literature)

7.3 - Qualitative (narrative) triangulation

Additionally, both the qualitative and quantitative triangulation exercises used the detailed risk-of-bias assessments performed and presented as part of the systematic review (Chapters 3 & 4). Similarly, risk-of-bias assessments were performed for each new source of evidence presented in this thesis. The risk of bias tools used to assess each result are described in detail in Section 3.2.8, but in summary, RCTs were assessed using the RoB 2 tool;¹⁶² NRSI using the ROBINS-I tool;¹⁵⁹ NRSE using an early adapted version of the ROBINS-E tool (see Section 3.2.8);^{163,164} and MR studies using the Mamluk et al. tool.¹⁶⁶

7.3 Qualitative (narrative) triangulation

As part of a qualitative triangulation of the new and existing evidence identified by this thesis, all information sources were initially grouped by outcome, and the findings from each source were narratively compared and contrasted. Potential reasons for heterogeneity between study designs were examined with specific reference to the risk-of-bias in each.

All-cause dementia

Analysis of lipid levels across both the systematic review (Chapters 3 & 4) and the IPD analysis (Chapter 6) both found extremely weak evidence for an association of any lipid fraction with all-cause dementia (TC HR: 0.97, 95%CI: 0.88-1.07; LDL-c HR: 0.97, 95%CI: 0.86-1.08; HDL-c HR: 1.05, 95%CI: 0.96-1.14; triglycerides HR: 0.90, 95%CI: 0.74-1.09). Similarly, meta-analysis of studies identified by the review which examined a binary hypercholesterolemia provided weak evidence for an association of this risk factor with Alzheimer's disease (HR: 1.11, 95%CI: 0.98-1.25). Finally, a meta-analysis of RCTs of statin use (OR: 1.07, 95%CI: 0.70-1.66) and of MR analyses of genetic instruments that mimic statin use (RR: 0.90, 95%CI:

7.3 - Qualitative (narrative) triangulation

0.29-2.81) both found weak evidence of an association, providing indirect evidence on the effect of lipid lowering on this outcome.

In contrast, the meta-analysis of NRSI of statin use found a slight protective effect (HR: 0.77, 95%CI: 0.69-0.87), though many of the studies in this analysis were found to be at risk of immortal time bias, which would be expected to favour the intervention. Finally, analysis of the association of statin use with all-cause dementia in the CPRD (Chapter 5) found evidence of a harmful association (HR:1.16, 95%CI:1.14-1.19), though this is likely to be driven by the substantial confounding by indication identified in the vascular dementia subgroup.

For fibrates, an alternative lipid-regulating agent, there was some evidence of an increased risk of all-cause dementia in the CPRD (HR:1.28, 95%CI:1.08-1.52), though this did not replicate in the meta-analysis of previous studies examining the association of fibrates and all-cause dementia (HR: 0.89, 95%CI: 0.75-1.07).

Alzheimer's disease

Meta-analysis of lipid levels in the systematic review found extremely weak evidence for an association of any lipid fraction with Alzheimer's disease (TC HR: 1.00, 95%CI: 0.94-1.06; LDL-c HR: 1.06, 95%CI: 0.98-1.16; HDL-c HR: 0.99, 95%CI: 0.91-1.07; triglycerides HR: 1.00, 95%CI: 0.84-1.18). Similarly, meta-analysis of studies examining a binary hypercholesterolemia provided weak evidence for an association of this risk factor with Alzheimer's disease (HR: 0.99, 95%CI: 0.78-1.25). A meta-analysis of MR analyses of genetic instruments that mimic statin use found weak evidence for an association of lipid lowering with Alzheimer's disease (RR: 0.76, 95%CI: 0.51-1.14).

In contrast, the meta-analysis of NRSI of statin use found a slight protective effect on this outcome (HR: 0.83, 95%CI: 0.69-0.99), though similar to the all-cause dementia outcome, many of the studies in this analysis were found to be at risk of immortal time bias. However, the CPRD analysis, which attempted to account

7.3 - Qualitative (narrative) triangulation

for this potential bias through the use of time-varying treatment indicator, found weak evidence for an effect (HR:0.98, 95%CI:0.94-1.01). This validity of this finding was limited by the potential for differential misclassification on the basis of the exposure in the CPRD analysis.

Finally, the IPD analysis provided no additional information on the Alzheimer's disease outcome, as a previously published analysis examined the association of lipid levels with this outcome in the Whitehall II cohort. The published analysis was included in the meta-analysis of NRSE, the results of which are presented above.

Vascular dementia

There was a general absence of existing evidence on vascular dementia outcomes. Meta-analysis of published results was only possible for the total cholesterol lipid fraction (HR: 1.05, 95%CI: 0.79-1.41). A meta-analysis of studies examining hypercholesterolemia also found weak evidence for an association with vascular dementia (HR: 1.20, 95%CI: 1.00-1.44). Similarly, a meta-analysis of statin use found weak evidence for an effect on this outcome (HR: 0.99, 95%CI: 0.79-1.25). The CPRD analysis found evidence for a harmful association of statin use (HR:1.81, 95%CI:1.73-1.9), though this finding is likely to be the result of severe confounding by indication related to vascular factors, as identified through the use of control outcomes.

The IPD analysis provided supporting evidence for the absence of an effect of any fraction on vascular dementia, with the sole exception of triglycerides. Raised triglycerides were associated with an increased incidence of vascular dementia across two previously unanalysed cohorts (OR: 1.25, 95%CI: 1.01-1.54), though there is the potential for uncontrolled confounding due to the limited information on covariates available. With respect to evidence on treatments for hypertriglyceridemia, examination of fibrates in the CPRD provided weak evidence of an association with vascular dementia (HR:1.29, 95%CI:0.83-2.02).

7.4 Quantitative triangulation

The qualitative comparison presented above identified no consistent effect of any lipid fraction on any dementia outcome across the evidence sources presented in this thesis. However, it is clear that this qualitative approach faces difficulties in interpretation when the number of individual results available is large, as discussed in the introduction to this chapter. This is true even when using summary estimates from meta-analyses of primary studies, which sacrifice valuable information on the biases inherent to each individual result.

In the following section, I propose and apply a novel quantitative triangulation framework to address these limitations. This approach incorporates recent advancements in the way that bias in results is assessed (namely the move to domain-based assessment tools)¹⁶² and existing methods for bias-/indirectness-adjusted meta-analysis³⁴⁶ to integrate the numerical results of the multiple approaches. The absence of any clear signals across the evidence base, as detailed in the previous section, does not make for particularly interesting case studies but I will nonetheless use the identified evidence to illustrate the novel methodological approach.

7.4.1 Methods

The proposed framework involves several steps, described in detail in the following sections. In summary, these steps are:

1. Define the causal question of interest
2. Identify of relevant evidence sources and standardise effect directions
3. Specify idealised version of each study
4. Assess the extent and direction of bias/indirectness in each result
5. Define modifying terms for bias and indirectness in each result
6. Calculate bias-/indirectness-adjusted results and perform meta-analysis

In order to illustrate the bias-/indirectness adjustment process detailed in the subsequent sections, the process for calculating the adjusted estimate is described in detail for a single result. Following this, the framework is applied to two case studies, as defined below.

Definition of the causal questions of interest (case-studies)

Using the ROBINS-E framework,¹⁶³ I defined the parameters of my causal questions of interest:

Case study #1: Effect of mid-life LDL-c on Alzheimer's disease

- *Population of interest:* General population
- *Exposure of interest:* Low density lipoprotein cholesterol
- *Exposure window of interest:* Mid-life (45-60)
- *Summary measure of exposure over time:* Average exposure
- *Outcome of interest:* Alzheimer's disease

Case study #2: Effect of mid-life triglycerides on vascular dementia

- *Population of interest:* General population
- *Exposure of interest:* Triglycerides
- *Exposure window of interest:* Mid-life (45-60)
- *Summary measure of exposure over time:* Average exposure
- *Outcome of interest:* Vascular dementia

These causal questions were chosen for several reasons. Firstly, work presented in previous chapters had identified some evidence for an association in these exposure/outcome pairs. Secondly, given the long prodromal period between the onset of physiological changes in the brain and presentation of dementia symptoms,

it seems likely that conditions during the mid-life period are particularly important. Finally, examining the causal impact of a lipid fraction on a specific outcome (e.g. Alzheimer's disease), rather than a composite outcome (e.g. all-cause dementia), focuses on a single causal pathway rather than including several conditions in the outcome definition that likely have very different mechanisms of disease. In addition, previous work presented in this thesis illustrated that the component conditions (in this case Alzheimer's disease and vascular dementia) are likely to be subject to very different confounding structures (see Section 5.4.2), and use of a composite all-cause outcome in the triangulation framework may mask this.

Identify relevant results

Once the causal question of interest had been defined, relevant individual results were obtained from the data sources described in Section 7.2. In a broad sense, this meant extracting results that examined the relationship between the exposure/outcome pair, either directly (e.g. non-randomised studies of LDL-c levels) or indirectly (e.g. non-randomised studies of statins, or Mendelian randomisation studies using genetic instruments for LDL-c levels).

Once the set of relevant results were identified, effects were standardised to refer to the risk resulting from a “reduction” in the lipid fraction. For both non-randomised studies and Mendelian randomisation analysis of lipid levels, the association is usually reported per 1-SD increase in the lipid fraction. In this case, the effect estimates were inverted to ensure consistency across study designs.

Specify idealised version of each study

Once the set of relevant results have been identified, an idealised study for each is described. An idealised study can be viewed as a replicate of the study producing the result of interest but with all sources of bias removed. The risk of bias is

7.4 - Quantitative triangulation

then assessed against this idealised study, while the causal question addressed by each idealised study is compared with the target causal question to define the indirectness of each result.

Note that all of the risk-of-bias tools used to assess non-randomised studies required specification of an idealised version of the study, against which bias was assessed. If using a risk-of-bias assessment tool that does not require this step, the idealised version of the study should be defined in advance of risk-of-bias assessment.

Assess risk and direction of bias in each result

The assessment of bias in each result is discussed in Section 7.2 above. For the sake of consistency and ease of computing, I mapped the different acceptable risk-of-bias judgements in each tool to a harmonised set: “Low”, “Moderate”, and “High”. The exact mapping can be seen in Table 7.2. Of note, no mapping was performed for the critical risk-of-bias levels present in the tools for non-randomised studies. This is because current best practice in evidence synthesis is to exclude all studies at critical risk of bias for further quantitative synthesis.¹⁵⁹

Table 7.2: Mapping of the different acceptable levels of bias across the assessment tools to a harmonised set. Note that no mapping for “Critical” risk of bias was performed, as best practice recommends that studies at critical risk of bias are excluded from quantitative syntheses.

Harmonised	RoB2	ROBINS-I / ROBINS-E	MR
Low	Low	Low	Low
Moderate	Some concerns	Moderate	Moderate
High	High	Serious	High
-	-	Critical	-

In addition to assessing the extent of bias, as defined by the three levels detailed above, I attempted to record the predicted direction and type of bias in each domain. This was achieved via an additional question at the end of each bias domain, which asked assessors to predict the expected direction of bias in that domain. Note that this question is currently only formally included in two existing tools (ROB2/ROBINS-I), but I employed an identical approach during assessment of non-randomised studies of exposure and Mendelian randomisation studies. Acceptable responses to this question included:

- “Favours experimental”/“Favours comparator” (defines an additive bias)
- “Towards null”/“Away from null” (defines a proportional bias)
- “Unpredictable”

As indicated in the options above, the response to this question is also used to determine the predicted type of bias as either additive or proportional. Proportional biases are dependent on the magnitude of the effect, while additive biases are not.³⁴⁶

For additive biases, determining the absolute direction of bias was simply a case of whether the bias is expected to shift the effect estimate to the left or right on an imaginary forest plot. For example, the estimate would be shifted to the right if a bias was predicted to favour the comparator (Figure 7.1).

In contrast, for proportional biases, whether the estimate should be increased or decreased proportionally depends on the current position of the point estimate relative to the null. For example, if the effect estimate represents a protective effect (below the null), then bias towards the null would be adjusted for by moving the effect estimate proportionally to the right. In contrast, if the effect of the intervention is harmful (effect estimate above the null), bias towards the null would be adjusted for by moving the effect estimate proportionally to the left. Both scenarios are illustrated using example data in Figure 7.1.




Figure 7.1: Illustration of calculating the absolute direction of biases (indicated by arrows), based on bias type. For additive biases, the direction is consistent regardless of the position of the effect estimate relative to the null - bias “favouring the comparator” will always pull the effect estimate to the right, and can induce a change from a protective to harmful effect. In contrast, the position of the effect estimate is accounted for when determining the absolute direction of proportional biases, defined as towards or away from the null.

The predicted direction and type of bias were not recorded for domains at “Low” risk of bias. Finally, for some domains, only one type of bias is allowed. For example, in the confounding domains in the ROBINS-I/ROBINS-E tools, only the “Favours experimental”/“Favours comparator” and “Unpredictable” options are available, meaning that bias in this domain will always be considered additive.

Visualisation of extent and direction of bias

To aid with the quantitative triangulation exercise, I created a new method to visualise the extent, direction and type of bias in each result, enabling detailed comparison across different studies contributing to a synthesis. Similar to the standard paired forest plots introduced in Chapter 3, these “bias direction” plots show the level (or extent) of bias across the domains for each result reported using coloured blocks. However, in contrast to the previous forest plots, symbols are

used to illustrate the predicted absolute direction of bias in that domain. Distinct symbol pairs are used to denote additive and proportional biases. Illustrations of these are provided in the Section 7.4.2.

Assign modifying values to risk/direction of bias

Following definition of the extent, type and direction of bias in each domain, I assigned a prior distribution on the log scale to each combination. For example, a “high” additive bias in j th domain of the i th study is defined as $\delta_{i,j}^{\text{High}}$ and uncertainty around this estimate is given by a distribution:

$$\delta_{i,j}^{\text{High}} = f(\mu_{i,j}^{\text{High}}, \sigma_{i,j}^{\text{High}}) \quad (7.1)$$

The sign of $\mu_{i,j}^{\text{High}}$ is defined by the absolute direction of the bias. If the bias is expected to pull the effect to the left, $\mu_{i,j}^{\text{High}}$ is given a negative sign, and if it is expected to pull the effect to the right, it is given a positive sign.

One key feature of the domain-based risk-of-bias tools is that the domains are considered interchangeable - i.e. a “high” judgement in one domain is equivalent to a “high” judgement in any other. As such, for this analysis, I assumed that the distributions assigned to each extent of bias were identical across all values of j , that is, were identical for all domains in the risk-of-bias assessment tool. To illustrate for additive biases:

$$\delta_{i,1}^{\text{High}} = \delta_{i,2}^{\text{High}} = \dots = \delta_{i,j}^{\text{High}}$$

and similarly

$$\delta_{i,1}^{\text{Moderate}} = \delta_{i,2}^{\text{Moderate}} = \dots = \delta_{i,j}^{\text{Moderate}}$$

Reasonable values for the position and spread of the prior distributions were informed by a previously-reported expert bias elicitation exercise.³⁴⁶ Using open data from that study, I calculated the mean values for the mean and variance of the adjustment distributions across all biases considered by that exercise (see Appendix A.7.1 for more details on how these were calculated). For the purposes of this illustrative analysis and to highlight the generalised nature of the code used to perform the analysis (see Section 7.5.5), I used the calculated values to inform two scenarios of modifying distributions (Table 7.3) for additive biases, and a single set for proportional biases. In the first scenario, the relationship between the position assigned to each additive judgement is linear. In the second scenario, the step from “Moderate” to “High” is twice that from “Low” to “Moderate”. Across both additive and proportional biases, the variance of the “High” judgement distribution was defined to be greater than that of the “Moderate” judgement.

Table 7.3: Prior distributions for extent and type of bias under two scenarios, on the log scale. Note that the distributions for proportional biases were kept consistent across the two scenarios.

Bias Level	Additive bias		Proportional bias
	Scenario 1	Scenario 2	
Low	-	-	-
Moderate	N(.09, .05)	N(.09, .05)	N(.03, .016)
High	N(.18, .1)	N(.27, .1)	N(.06, .032)

Where the direction of bias in a domain was unpredictable for a given result, the mean of the prior distribution for that domain was set to 0 but the appropriate variance for the recorded extent of bias was retained (e.g. for an unpredictable “High” additive bias, the distribution would be $N(0, 0.1)$ under Scenario 1). In other words, adjusting for bias in a domain with an unpredictable direction of bias has no impact on the position of the effect estimate but does increase the uncertainty around it.

Assessing and assigning prior values to indirectness

The indirectness of a result (also termed external bias, external validity, relevance, generalisability, or applicability) is defined here as the discrepancy between the research question addressed by the “idealised” study and the causal question of interest.

An identical approach was employed to assess, and assign prior distributions to, the indirectness in each result. The target question for each result was compared against the causal question of interest with respect to three domains, defined as important by the GRADE framework for assessing the certainty of evidence:³⁴⁷ population, intervention/exposure, and outcome. I again used the scale of “Low/”Moderate“/”High” to quantify the extent of indirectness in each domain. All indirectness was defined *a priori* as being proportional in nature (i.e. depending on the magnitude of the effect), in line with previous work on this topic.^{346,348} As above, data from the previous elicitation exercise were used to inform reasonable prior distributions for each level of indirectness (Table 7.4).

Table 7.4: Prior distributions for extent and type of bias under two scenarios, on the log scale. Note that the distributions for proportional biases were kept consistent across the two scenarios.

Bias Level	Proportional bias
Low	-
Moderate	N(.06, .016)
High	N(.12, .032)

As an illustrative example, consider the first causal question of the interest in this exercise, the effect of LDL-c in mid-life on Alzheimer’s disease. Here, studies of lipids levels in this time period would require minimal adjustment, whereas

studies examining statin use (indirect exposure/intervention) in late-life (indirect population) would be adjusted and down-weighted due to reduced relevance to the causal question.

Combine in a bias-adjusted meta-analysis

Using the method reported in Turner *et al.*,³⁴⁶ total additive and proportional bias and indirectness were calculated and used to define an adjusted estimate for each result included in the synthesis.

The adjusted estimate for each result, $\hat{\theta}$, is defined as:

$$\hat{\theta} = \frac{y_i - \mu_{i\beta}^{\text{In}}\mu_{i\delta}^{\text{In}} - \mu_{i\delta}^{\text{B}}}{\mu_{i\beta}^{\text{B}}\mu_{i\beta}^{\text{In}}} \quad (7.2)$$

where $\mu_{i\delta}^{\text{B}}$ and $\mu_{i\beta}^{\text{B}}$ refer to the total additive and proportional bias, and $\mu_{i\delta}^{\text{In}}$ and $\mu_{i\beta}^{\text{In}}$ refer to the total additive and proportional indirectness. Note that in this exercise, the total additive indirectness $\mu_{i\beta}^{\text{In}} = 1$, as all indirectness was defined *a priori* as being proportional.

The standard error of this estimate is then calculated as:

$$\text{SE}(\hat{\theta}) = \left(\frac{1}{\mu_{i\beta}^{\text{B}}\mu_{i\beta}^{\text{In}}} \right)^2 \left(s_i^2 + (\mu_{i\beta}^{\text{B}}2 + \sigma_{i\beta}^{\text{B}})(\hat{\theta}^2\sigma_{i\beta}^{\text{In}2} + \sigma_{i\delta}^{\text{In}2}) + \sigma_{i\beta}^{\text{B}}(\hat{\theta}\mu_{i\beta}^{\text{In}} + \mu_{i\delta}^{\text{In}})^2 + \sigma_{i\delta}^{\text{B}2} \right) \quad (7.3)$$

Once each individual result had been adjusted using this approach, I performed a random effects meta-analysis using the unadjusted and adjusted results. I also compared the adjusted results under the two scenarios of additive bias defined. results. I extracted the overall effect estimates, along with measures of heterogeneity in each case (τ^2 , I^2).^{349,350}

7.4.2 Results

Single study

As an illustrative example of the process using a single result, the bias and indirectness inherent to the estimate of the effect of statin use on Alzheimer's disease in the CPRD (as analysed in Chapter 5) are considered here. For this result, there were three domains at greater than low risk of bias: Bias due to confounding, Bias due to definition of the outcome, and Bias in the selection of the reported result (Figure 7.2). In Domain 1 (bias due to confounding), the study did not adjust for *ApoE4* status, which is predicted to mask any true association between statins and Alzheimer's disease (see Section 5.4.3 for a fuller discussion). As such, the bias is predicted to *favour the comparator* and so the absolute direction of bias is to the right. In Domain 7 (bias in measurement of the outcome), there was a high risk of differential misclassification on the basis of the exposure. Here, history of statin use may make a diagnosis of vascular dementia more likely than Alzheimer's disease, and so for this outcome, this bias is predicted to *favour the experimental* (in this case, an intervention). As such, the absolute direction of bias is to the left. In Domain 7 (bias in selection of the reported result), for consistency with the other NRSI, in the absence of a protocol the bias is assumed to be *away from the null*. Given the position of the effect estimate below the null, the absolute direction of bias in this domain is to the left.




Figure 7.2: Bias direction plot for a single study showing the predicted direction of bias in each domain. Additive biases are characterised using arrows, while the proportional biases are denoted using $<$ and $>$. A question mark indicates that the direction of bias was unpredictable.

Indirectness was intentionally not displayed in the plot above, as these figures are created using `robvis` which is focused on creating visualisations of risk-of-bias data. In addition, I found it more useful to articulate the indirectness of a result in tabular format. With respect to this example result, I judged it to be a low risk in the population and outcome domains (Table 7.5). However, the result refers to the effect of statins on Alzheimer's disease risk, rather than LDL-c levels directly, putting it at high risk of indirectness in the intervention/exposure domain. Assessing the predicted direction of indirectness was more challenging than assessing the predicted direction of bias. In this case, treatment with statins result in temporary lowering of lipid levels which could thus underestimate the true effect of LDL-c lowering (indirectness towards the null). In contrast, statins may have an effect on Alzheimer's disease through a mechanism other than LDL-c lowering effect of statins and so the effect is overestimated (indirectness away from null). As such, I judged the direction of indirectness in this domain to be unpredictable.

Table 7.5: Assessment of indirectness for a single example study.

Domain	Target Setting	Idealised version of single study	Level	Direction
Population	General population, dementia free at baseline	UK general population, >45 years of age, dementia free at baseline	Low	-
Intervention/ exposure	Average low density lipoprotein cholesterol levels at midlife (45-60)	Initiation of statin use	High	Unpredictable
Outcome	Alzheimer's disease, assessed using recognised clinical criteria	Alzheimer's disease cases extracted from EHR	Low	-

The unadjusted estimate was 0.98 (95%CI: 0.95-1.02). After accounting for the biases and indirectness described, the adjusted result was 1.08 (95%CI: 0.48-2.44).

Case study #1: effect of LDL-c at midlife on Alzheimer's disease

From the data sources identified in Section 7.2, I identified 18 results relevant to my first causal question of interest. Most ($N = 17$) were identified via the systematic review,^{54,72,189–191,196,200,204,205,210,212,213,219,245,248} while Chapter 5 provided an additional estimate from the CPRD analysis. The extent, type and direction of predicted bias for each result can be seen in the bias-direction plot presented in Figure 7.3, stratified by study design.

7.4 - Quantitative triangulation




Figure 7.3: Bias direction plot, summarising the internal biases related to effect of LDL-c on Alzheimer's disease, stratified by study design.

The results of the unadjusted and the bias-/indirectness-adjusted (under Scenario 1) are shown in Figure 7.4. Following adjustment for bias and indirectness, the observed effect attenuates to the null (unadjusted 0.94, 95%CI: 0.89-0.99; adjusted 0.95, 95%CI: 0.87-1.04). Additionally, adjustment for bias and indirectness substantially reduced the heterogeneity between studies (unadjusted $\tau^2 = 0.0017$, $I^2 = 17$; adjusted $\tau^2 = 0.0013569$, $I^2 = 1.3037272$). However, as can be seen from the right panel in Figure 7.4, this reduction in heterogeneity between studies is largely achieved via a reduction in the precision of each individual result.




Figure 7.4: Random effects meta-analysis of the association of midlife LDL-c with Alzheimer's disease, using unadjusted and bias-/indirectness-adjusted results.

No substantial differences were observed between the summary estimates obtained under the two scenarios of additive bias (Appendix A.7).

Case study #2: effect of triglycerides at midlife on vascular dementia

From the data sources identified in Section 7.2, I identified 7 results relevant to the causal effect of triglycerides at midlife on vascular dementia. Four were identified via the systematic review,^{225,237,248} while Chapter 5 provided two additional estimates from unanalysed datasets (CaPS and Whitehall II), and Chapter 6 provides an estimate for the effect of fibrates, a treatment for hypertriglyceridaemia. The extent, type and direction of predicted bias for each result can be seen in the bias-direction plot presented in Figure 7.5, stratified by study design.

7.4 - Quantitative triangulation




Figure 7.5: Bias direction plot, summarising the internal biases related to effect of triglycerides on vascular dementia, stratified by study design.

Comparison of the unadjusted and bias-/indirectness-adjusted results for the effect of triglycerides on vascular dementia did not demonstrate a substantial difference (unadjusted 0.82, 95%CI: 0.64-1.05; adjusted 0.79, 95%CI: 0.58-1.07; Figure 7.6), though again, heterogeneity between the adjusted results was greatly reduced (unadjusted $\tau^2 = 0.054$, $I^2 = 65$; adjusted $\tau^2 = 0$, $I^2 = 0$). Similarly there was minimal difference between the results under the two scenarios of bias (Figure A.7 in Appendix A.7).




Figure 7.6: Random effects meta-analysis of the association of midlife triglycerides and vascular dementia, using unadjusted and bias-/indirectness-adjusted (Scenario 1) results.

7.5 Discussion

7.5.1 Summary of findings

This chapter has narratively synthesised the evidence identified and produced by the previous chapters, highlighting the absence of any consistent association between any blood lipid and any dementia outcome.

In addition, it has sought to provide a generalised framework for quantitative triangulation, building on existing methods for systematic domain-based risk of bias assessment and bias-/indirectness-adjusted meta-analysis. To illustrate the method, I considered two causal questions: the effect of LDL-c at mid-life with Alzheimer's disease and the effect of triglycerides on vascular dementia. In the quantitative triangulation framework, there was weak evidence for an effect in relation to either of the causal questions considered. The heterogeneity between results produced by different study designs contributing to the analysis was substantially reduced

using this method, though this was largely due to a reduction in the precision of each individual result.

7.5.2 Limitations

The quantitative analysis presented in this chapter is subject to some strong methodological limitation, which are discussed in detail below.

Defining reasonable prior distributions for bias and indirectness

The key limitation of the quantitative synthesis presented in this chapter is the lack of empirical evidence available for the prior distributions for bias and indirectness. While I attempted to address this by informing the prior distributions using data from a previous expert elicitation exercise, the extent of bias/indirectness in that analysis may not generalise to the one presented here.

Ideally, these prior distributions would be based on empirical data on the effect of different domains of bias/indirectness on a result. Most meta-epidemiological studies examining the effect of different methodological issues studied only randomised controlled trials,^{351,352} and the estimates of bias they produce may not hold for non-randomised study designs. However, the prior distributions defined here are broadly comparable to previously reported estimates of the impact of bias in non-randomised trials. For example, a previous simulation study estimated that, in the absence of true effect of LDL-c, *ApoE4* would induce a RR of 1.09 with risk of dementia per 1-SD increase in LDL-c.³⁵³ This is comparable to the mean assigned to the prior log-normal distribution mapped to “Moderate” additive bias ($N(0.09, 0.05)$, Table 7.3).

A further limitation is the assumption that the distributions of bias are identical across domains of bias. Previous evidence from meta-epidemiological studies of randomised controlled trials that investigated the effects of different biases indicates

that this is unlikely to be the case.³⁵⁴ The generalised framework for quantitative triangulation, presented here and available via the `triangulate` package (see Section 7.5.5), allows for domain-specific distributions of bias/indirectness. As more information on the effect of specific sources of bias becomes available, such domain-specific distributions should be used.

Finally, it may in fact be more reasonable for domains such as bias due to confounding to adjust on a per-confounder basis rather than mapping to a “Moderate” or “High” extent of bias. In this approach, prior distributions are assigned to each of the confounders pre-specified as part of the ROBINS-I/ROBINS-E tools.¹⁵⁹ Then, rather than assigning a domain level distribution based on an arbitrary judgement of how many important confounders are missing, the result is adjusted for each confounder not accounted for in the original analysis.

Accuracy of bias-/indirectness-assessment

A key issue in this analysis is the accuracy of the assessment of bias and indirectness in each result. It has been widely demonstrated that the inter-rater agreement when using risk-of-bias tools is low with regards to the extent of bias assigned to each domain^{355–357} Though not previously studied, given the difficulty in assessing the predicted direction of bias, agreement on this aspect of the tools is likely to be lower still.

A further issue in this regard is that the rigour of the tools may not be equivalent. This issue can be illustrated by the Ostergaard *et al.* study⁷² having low risk of bias across all domains and thus receiving a substantial proportion of the weight in the first case study (Figure 7.4). While the RoB suite of tools (RoB 2.0, ROBINS-I, ROBINS-E) were developed by an expert working group and have been widely used, the Mamluk *et al.* tool¹⁶⁶ used to assess risk of bias in Mendelian randomisation studies was designed by the authors for use in their own review. If the Mamluk tool is failing to adequately assess bias in Mendelian randomisation studies, then any

analysis based on its assessments is likely to be subject to bias. This observation is particularly problematic given the default position of the ROBINS-I and ROBINS-E tool to require a “Moderate” judgement in the confounding domain. Under these conditions, it is possible that a well-performed cohort study (low risk of bias across everything other than bias due to confounding) will be down-weighted versus a potentially poor MR study.

Low and critical risk of bias

As illustrated in Table 7.2, studies at critical risk of bias were not included in the analysis. While this is in line with best practice, there is theoretically no reason why studies with this extent of bias could not be included in the proposed framework. However, the estimation of an appropriate prior distribution for the effect of “Critical” bias would be substantially more challenging, and so was avoided here.

A related issue is the “Low” risk of bias judgement. In this analysis, I assumed that domains at “Low” risk of bias did not require any adjustment. However, “Low” risk of bias is not equivalent to the absence of bias, and potentially should still be adjusted for, if only minimally. In this case, defining the predicted direction of bias would be particularly challenging in the absence of obvious sources of bias.

Combination of different effect measures

A final limitation to this analysis is the synthesis of different effect estimates. As discussed in Section 3.2.9, hazard ratios are not directly comparable to odds ratios, as they inherently account for person time-at-risk.¹⁶⁹ As non-randomised studies (NRSI/NRSI) of dementia outcomes report hazard ratios and Mendelian randomisation studies report odds ratios, the resulting synthesis may be biased by the integration of these two different effect measures. This is primarily a concern

for common outcomes, as when the outcome is rare, the odds, risk and hazard ratios approximate each other.

7.5.3 Strengths

The core strength of this analysis is that it is based on a comprehensive systematic review, supplemented by two additional primary studies to fill in the identified evidence gaps. In general, triangulation of any sort should be considered a natural extension of evidence synthesis, and therefore should follow best practice in relation to finding and critically assessing all relevant information. Additionally, the approach presented here also builds on recent developments in bias assessment to “explode out” the component results of a meta-analysis, and consider the effects of bias/indirectness in each separately.

A further advantage of this method comes from the ability to specify the prior distributions for each level of bias/indirectness in advance of performing the assessments. Expert elicitation of the extent of bias/indirectness using a panel of methodologists and topic experts is a powerful technique but the point at which it is deployed in the framework is important. In previous attempts at bias-/indirectness-adjusted meta-analysis, the extent of bias in each study was assessed via a elicitation process,^{346,348,358} during which experts were aware of the results of each analysis. This approach is potentially subject to differential misclassification of the impact of bias/indirectness on the basis of the results, as there is no way to ensure that results at a similar level of bias for confounding (for example) are being adjusted by the same amount across studies. As an illustrative example, if experts are influenced by the knowledge of the effect estimate in a study, then a result at “Moderate” risk of confounding and demonstrating a stronger protective effect may receive greater adjustment than a study at the same risk of confounding but with a more modest effect. This is likely to be particularly problematic where experts have preconceived ideas about the true effectiveness of the intervention.

Separating the assessment of bias/indirectness from the assignment of modifying values to each judgement will limit the potential for this misclassification. Using the framework proposed, reasonable modifying distributions for each level of bias can be defined *a priori* by the study team using the elicitation procedure detailed previously,³⁴⁶ similar to how important confounders and co-interventions are defined in advance when performing ROBINS-I/ROBINS-E assessments.¹⁵⁹ As an example, expert elicitation could be used to *a priori* define an additive bias distribution of $N(.1,0.7)$ for “Moderate” risk of bias in Domain 1 of the ROBINS-I tool (bias due to confounding). Risk-of-bias assessments are then performed, and each result at “Moderate” risk of bias in Domain 1 is adjusted using this pre-specified distribution.

Finally, accounting for bias using an incorrect prior distribution is no less problematic than synthesising and drawing conclusions from effect estimates as if they were unbiased, a common occurrence in systematic reviews.¹⁸⁵ While this analysis may be limited by the absence of empirically-derived adjustment distributions, it at the very least acknowledges the uncertainty of evidence due to bias/indirectness via a reduction in precision.

7.5.4 Future research

An obvious avenue for future work in relation to quantitative triangulation is the identification of empirical prior distributions for the effect of bias/indirectness in non-randomised studies using meta-epidemiological approaches. This will be substantially more challenging than examining the effect of bias in RCTs,³⁵⁴ given the absence of an underlying database of meta-analysis of non-randomised studies.

Future development of this framework should also account for bias at the analysis level in terms of missing evidence.¹⁶⁸ For example, there is at least one known study relevant to the vascular dementia case study that reported a non-significant result but provided insufficient details to be included in the analysis.²²³ Ideally, a quantitative triangulation framework would further account for this proportional

meta-bias (as bias due to missing evidence is most likely to bias away from the null due to publication bias mechanisms), in addition to result-specific biases.

Appreciation that both the risk-of-bias and indirectness of a result should be assessed is growing. Indeed, some existing for the assessment of diagnostic test accuracy³⁵⁹ and prediction models³⁶⁰ consider indirectness (termed applicability in these tools) alongside the assessment of bias. Future iterations of the core risk-of-bias tools (RoB 2, ROBINS-I, ROBINS-E) could take this into account, though there is an competing argument that indirectness is already handled via tools such as the GRADE framework.³⁴⁷ In addition, simple steps like harmonising the risk-of-bias judgements across tools and making the direction of bias question mandatory (even if users default to “Unpredictable”) would represent an improvement in the tools and encourage users to think about how the biases assessed affect the corresponding result. Similarly, software that implements a risk-of-bias tool should allow for direction/extent of bias question and carefully consider how this data will be exported.

7.5.5 Outputs

There are two key outputs from this project. The first is the `triangulate` R package. Personal communication with the authors of the original method resulted in the STATA code to implement the bias-/indirectness-adjusted model.³⁴⁶ This has since been generalised as part of the `triangulate` R package to enable other users to apply the approach detailed here. The package allows for preprocessing of domain-based risk-of-bias data to correctly identify the direction of the bias relative to the effect estimate, use of domain-specific prior distributions, production of bias direction plots (via the tool described in Appendix B.2, calculation of bias/indirectness-adjusted estimates for each result, and synthesis of these adjusted results in a standard random-effects meta-analysis.

The second key output is an annotated example dataset for the LDL-c/Alzheimer's disease causal question addressed in this chapter. Available via the `triangulate` package, this data will provide developers of new triangulation methods with an example dataset on which to test new tools, something which seriously hindered development of this work.

7.6 Summary

- In this chapter, I narratively described the evidence identified or produced in previous chapters, and highlighted the issues with a qualitative approach to triangulation when there are several sources of evidence.
- I then illustrated a proposed generalised framework for quantitative triangulation, exploiting existing methods in evidence synthesis and recent developments in the assessment of different risks of bias. I illustrated this method using two case studies: the effect of mid-life LDL-c on Alzheimer's disease, and the effect of mid-life triglycerides on vascular dementia.
- I discussed the limitations of this proposed method, given the absence of informative priors for the impact of bias/indirectness, and proposed future research (i.e. meta-epidemiological studies) that would address them.
- In summary, quantitative triangulation is a promising field that should be considered a natural extension of evidence synthesis. The implications of the synthesised evidence presented here to clinical practice, public health and future research is considered in the following chapter.

Hold

— Hold

8

Discussion

Contents

8.1	Lay summary	210
8.2	Introduction	210
8.3	Summary of clinical findings	211
8.4	Summary of methodological contributions	214
8.5	Overall strengths and limitations	216
8.6	Lessons learned	218
8.7	Future work	218
8.8	Overall conclusions	222

8.1 Lay summary

TBC

8.2 Introduction

The central aim of this thesis was to infer the causal effect of blood lipid levels on dementia outcomes via evidence synthesis methods. In this discussion, I will review the contribution of each chapter to this aim, summarise the principal findings and methodological innovations of the thesis, and discuss the implications of my findings for both clinical and public health practice. I consider the overall strengths and limitations of this thesis as a coherent body of work and suggest avenues for impactful future research that builds on the research presented.

8.2.1 Chapter summary

This thesis is comprised of a mix of methodological and applied research chapters.

In **Chapter 2**, I introduced the core concepts considered in this thesis and framed the research presented in relation to a theoretical framework of evidence synthesis.

In **Chapter 2**, I presented a new tool for the systematic searching of health-related preprints, and detailed two use cases for preprint metadata beyond the systematic searching of this evidence source.

In **Chapter 3**, I described the methods of a comprehensive systematic review of all available evidence on the association of both lipids and lipid-regulating agents with dementia outcomes. The results of this analysis are then presented in **Chapter 4**.

In **Chapter 5**, I described an analysis of the association of lipid-regulating agent use and dementia outcomes in the CPRD, a large electronic health record database. I make use of control outcomes to illustrate insufficiently controlled confounding.

In **Chapter 6**, I presented an individual participant data meta-analysis of the association of lipid levels and dementia outcomes, using previously unanalysed data accessed via the Dementia Platform UK.

Finally, in **Chapter 7**, I proposed a new method for the systematic integration of multiple evidence sources as part of quantitative triangulation framework. I illustrated the method using two case studies (the effect of LDL-c on Alzheimer's disease and the effect of triglycerides on vascular dementia), drawing on the evidence identified or produced by previous chapters.

8.3 Summary of clinical findings

8.3.1 Effect of blood lipids on dementia outcomes

As the results of the different components of this thesis are discussed in detail as part of the triangulation analysis (see Section 7.3), here I briefly summarise the key findings.

Overall, I did not identify a consistent effect of any blood lipid fraction (total cholesterol, LDL cholesterol, HDL cholesterol or triglycerides) on any dementia outcome (all-cause dementia, Alzheimer's disease or vascular dementia). While published non-randomised studies provided evidence for a protective effect of lipid-lowering via statin use on Alzheimer's disease and the IPD analysis suggested a harmful effect of raised triglycerides on vascular dementia, these effects were not maintained when incorporated along with other sources of evidence in a quantitative triangulation framework. Similarly, the IPD analysis did not provide evidence for an interaction of patient characteristics with the association of blood lipids on dementia outcomes, though the results are limited by low number of cohorts analysed.

8.3 - Summary of clinical findings

In the published literature, statins were by far the most studied lipid-regulating agent in relation to dementia outcomes. Given the large proportion of patients taking statins as indicated by my analysis of the CPRD data, this finding is unsurprising. Finally, there was a substantial absence of evidence for vascular dementia.

8.3.2 Comparison with new evidence

Since the searches underpinning the systematic review described in this thesis were performed (July 2019), further studies on this topic have been performed. Two of these, notable for the analysis of large UK cohorts, are discussed here.

The first is a large-scale analysis of 953,635 patients in the CPRD, followed from their first LDL-c measurement.³⁵³ The study found a slight increased risk of dementia was associated higher LDL-c measured at mid-life, driven by the finding for the Alzheimer's disease subgroup. These results are consistent with the findings presented in this thesis, particularly given the lack of adjustment for confounding by *ApoE4*, which was estimated by the study to induce a RR of 1.09 per 1-SD increase in LDL-c in the absence of a true causal effect.

The second analysis examined 502,226 participants in the UK Biobank, a large population-based prospective cohort study.^{361,362} The analysis found no association between lipids measured at mid-life and dementia outcomes, comparable to the results of this thesis. However, similar to the study discussed above, this analysis did not have access to genetic data and so may be subject to residual confounding by *ApoE4* genotype.

8.3.3 Implications for clinical practice

Given the absence of any consistent signals across the evidence sources analysed, there are no clear indications for clinical practice arising from this thesis. While

my research did not provide any strong evidence for an effect of lipid-lowering on dementia outcomes, the cumulative strength of evidence is also insufficient to rule out an effect.

Given this ambiguity, the use of statins should be restricted to their primary and well-understood purpose of lipid-lowering to reduce the risk of cardiovascular events. However, it is important to recognise that even under the scenario that a strong harmful effect of lipid-lowering on dementia outcomes was identified, whether or not to prescribe a statin at mid-life will always be guided by factors other than dementia prevention. Patients must live to a certain age in order to be at risk of dementia (with the exception of familial early-onset dementias), which is substantially less likely if hypercholesterolemia at mid-life is left unchecked.

In summary, clinicians should be aware of the uncertainty of evidence surrounding the effect of lipid-lowering on dementia outcomes and should be prepared to convey the same to patients.

8.3.4 Implications for public health

[Note: Yoav, I am particularly interested in your thoughts on this section]

Similarly, given the ambiguity of evidence, there are no clear implications for public health practice. However, the role of public health, and particularly continuous population surveillance, may be important in providing a further source of evidence in relation to the effect of lipid lowering on dementia. Public health practitioners could examine summary-level statistics of GP prescription data to attempt to relate lipid-lowering treatment to temporal or geographical trends in dementia incidence. It will be difficult to assess if any observed trends are due to a true effect or to confounding or ecological biases. However, as dementia outcomes will be monitored anyway for incidence/prevalence estimation and planning of services/care, this approach would provide a ready source of further evidence.

8.4 Summary of methodological contributions

In addition to addressing the causal impact of lipids on dementia outcomes, the research described in this thesis represents a number of novel contributions to the field of evidence synthesis methodology.

8.4.1 Inclusion of preprinted evidence

Preprints are an important source of evidence and searching of preprint repositories should become an accepted part of the evidence synthesis process. To support this, I developed a new tool (as described in Chapter 2) to enable researchers to readily search health-related preprints.

8.4.2 Bias

One of the central methodological contributions of this thesis has been on the conduct, visualisation, and incorporation of bias assessments as part of an evidence synthesis exercise.

In terms of assessment, I applied domain-based tools to evidence relevant to the effect of lipids on dementia outcomes and highlighted the limitation of the available tools for Mendelian randomisation studies. I also piloted and provided feedback on a forthcoming tool for assessing the risk of bias due to missing evidence. In the primary analysis of data from the CPRD, I used negative control outcomes to assess the potential for bias due to insufficiently controlled confounding by indication.

In terms of visualisation of bias, I have made a number of methodological contributions. I created a well-received R package to create paired forest plots (see

Appendix B.2), and built on this to enable creation of the bias direction plots illustrated in Chapter 7.

Finally, I made several novel contributions around the incorporation of bias into analyses. These were intended to encourage authors of reviews to actively consider the impact of bias in their synthesis, something that empirical evidence suggests happens infrequently at present.¹⁸⁵ The new methods introduced by this thesis range from the forest plots stratified by overall risk of bias level presented in Chapter 4 to the more advanced triangulation framework presented in Chapter 7.

8.4.3 Triangulation framework

The proposed quantitative triangulation framework represents the final novel contribution of this thesis to evidence synthesis methodology. Using pre-specified adjustment distributions mapped to the results of the domain-based risk-of-bias and indirectness assessments, this approach provides a systematic way to account for biases/indirectness across studies, reducing the potential for differential adjustment based on knowledge of the results.

Ideally, this framework should be based on the results of a comprehensive systematic review, which identifies all evidence (both direct and indirect) related to a causal question of interest. This would represent a break from the current practice of limiting review to a single type of evidence, and would result in fewer, more comprehensive reviews. However, in the current context of research waste, a concentration of effort in fewer reviews may be advantageous. Section 4.5.2 illustrates the duplication of reviews in relation to the topic of this thesis, while an assessment of the COVID-19 literature found that, for a specific clinical question, there were substantially more reviews (25) than available primary studies (17).³⁶³

Finally, while the framework proposed here could (fairly) be criticised over the validity of the prior distributions of bias/indirectness chosen, the assumptions made about the impact of bias in this thesis are no stronger than those made when

8.5 - Overall strengths and limitations

synthesising effect estimates as though they were unbiased. The proposed framework at the very least recognises the uncertainty introduced by bias/indirectness and decreases the precision of the overall effect estimate on this basis.

In summary, the proposed triangulation framework will enable better use of all available evidence to address causal questions.

8.5 Overall strengths and limitations

The strengths and limitations of each aspect of this thesis have been discussed in the respective chapter. Here, I highlight the strengths and weaknesses of this thesis as a body of work.

8.5.1 Strengths

There are several strengths to this thesis as a whole. Specifically, the identification and triangulation of evidence across study designs and publication status sets this thesis apart from previous synthesis of the evidence on this topic. Additionally, this thesis has also produced new evidence on the association of blood lipids with dementia outcomes in previously unanalysed cohorts and provided additional information on a previously under-studied association (lipids/LRA and vascular dementia).

A further strength of this thesis is the production of software to support the novel evidence synthesis techniques used. Extensive documentation has been written to guide users through usage of the tools and example data is provided for use. Finally, substantial effort has been invested to make the research documented in this thesis as reproducible as possible - this is documented more fully in Appendix A.8.1.

8.5.2 Weaknesses

However, there are also several strong limitations to this thesis. In the first instance, as with many studies of dementia outcomes, a limiting factor in the interpretation of the results is the absence of a detailed pathological mechanism. It is possible that statins have a true effect via some other pathway , but without knowledge of the mechanism, the plausibility of this relationship cannot be assessed.

Secondly, each analysis presented here makes use of secondary data sources, be it published literature, electronic health records, or existing cohorts. Secondary data can limit the type of analyses performed (for example, in the IPD analysis presented in Chapter 6, absence of time-to-event data prevented the use of hazard ratios to quantify dementia risk) and the validity of the data if collected for purposes other than research (for example, the accuracy of dementia diagnoses in electronic health records is known to be variable).²⁷⁹

Thirdly, missing evidence was a common limitation across this thesis. In Chapters 3 & 4, several studies stated that they did not report the results of an analysis because the results were not significant. This was particularly common among non-randomised studies of exposure examining blood lipids directly. Similarly, an absence of evidence on vascular dementia, potentially due to a publication bias mechanism, limited the analysis of this outcome. Missing evidence was also as an issue in the IPD meta-analysis reported in Chapter 6, in the form of a poor response to data access requests, while the absence of empirically-based distributions of bias/indirectness for use with the triangulation framework limited the credibility of the results produced.

A final weakness stems from the geographical focus of the data analysed in this thesis. All primary analysis presented drew on data from the UK (CPRD, CaPS, EPIC Norfolk, Whitehall II), while the majority of studies identified by the review were based in the Western world (Figure 4.2). This may limit the generalisability of the results presented to different populations.

8.6 Lessons learned

As part of a reflective learning process throughout my PhD, I maintained a catalogue of failures (available in Appendix A.8.2) describing analytical mistakes, failed experiments, and unsuccessful grant applications. However, I found there was one central learning point from this thesis, namely to be slightly less ambitious when planning future research projects.

The proposal for the program of research underpinning this thesis was created in advance of starting my PhD, as a detailed plan was required in order to secure the funding that supported me through my studies. In hindsight, the decision to take a broad approach to the inclusion of dementia outcomes resulted in a larger workload than anticipated, particularly when also considering evidence from multiple different study designs. Additionally, as dementia subtypes likely have very different aetiological pathways, it may have been better to focus on a single subtype (e.g. Alzheimer's disease) when considering the causal effects of lipids. Similarly, attempting to undertake a full individual participant data meta-analysis as a single part of a larger program of research was overly ambitious, as data cleaning and harmonisation for just three cohorts was a substantial undertaking.

In the future, armed with the experience gained through conduct of the research projects presented here, I will be better place to scope and design research projects.

8.7 Future work

Several avenues of future research could be pursued, building on the novel work presented here. These are grouped by topic in the following sections.

[Question: is it better to use “should” or “could” when listing ideas for future research? I.e. “future work could/should look at X”?]

8.7.1 Generation of new RCT evidence

It is customary at this point in the discussion of results based (primarily) on observational data to recommend that a large-scale randomised controlled trial be performed to assess the effect of the proposed intervention. Given the costs and logistical challenges associated with trials examining outcomes with long prodromal periods such as dementia, a more efficient approach is offered by the opportunistic post-trial follow-up of existing RCTs of lipid-regulating agents.

This approach has already been used to assess long-term (11-20 years) safety outcomes of statin use, using participant recontact³⁶⁴ and linkage with electronic health records to identify events.³⁶⁵ A similar approach could be used to assess the impact of randomisation to statins at midlife on subsequent risk of dementia outcomes. Use of data linkage would represent the most cost-efficient approach,³⁶⁶ though as noted in Chapter 5, there is the potential for non-differential misclassification when defining dementia outcomes using EHRs.

8.7.2 Evidence on vascular dementia

As highlighted by the results of the systematic review presented in Chapter 4, there is an absence of evidence on vascular dementia across the existing evidence base. While the primary studies presented here go some way towards supplementing this evidence base, future work is needed both to investigate the reasons for this evidence gap (e.g. due to an absence of primary data, challenges in the analysis of this outcome as documented in Chapter 5, or a publication bias mechanism) and to address it.

However, it seems unlikely that further analysis of observational data alone will be sufficient to investigate this outcome, as indicated by the presence of strong confounding by indication in the analysis of statins and vascular dementia (Chapter 5). Even using new methods, such as a target trial emulation approach, previous analyses have not replicated the known protective effect of statins on coronary heart

disease identified by RCTs (see Section 5.4.2).³¹⁶ Similarly, Mendelian randomisation studies are limited to a single one-sample analysis of HMGCR variants as a proxy for statin treatment. This study suggested a protective effect of lipid-lowering on vascular dementia, though there was a moderate risk of bias in this analysis and the effect was attenuated to the null when including a wider range of lipid-lowering variants.

Taken together, these points reinforce the need for large-scale GWAS of vascular dementia, similar to those current available for Alzheimer’s disease, to identify associated loci that future two-sample Mendelian randomisation studies could exploit. While GWAS of this outcome are likely to be methodologically challenging, given the difficult in diagnosing “pure” vascular dementia, it would be a worthwhile endeavour and allow for the assessment of genetically-driven risk factors beyond those considered in this thesis.

8.7.3 Preprinted evidence

In terms of evidence synthesis methodology, future work on preprinted evidence should address how to handle discrepancies between the preprinted and published versions of a paper when identifying results for inclusion in a synthesis. This will be particularly important in cases where the results are substantially different between the two versions or if a result available in the preprint manuscript is not presented in the published version.

The `medrxivr` tool enables programmatic searching of health-related preprints, and future work could incorporate the tool, along with software to search other literature sources, into an automated searching pipeline as part of a living systematic review.³⁶⁷ In addition, the ready access to preprint data afforded by the tool will enable future meta-epidemiological studies to examine factors which may influence eventual publication (e.g. significant result, geographical location/gender/career stage of first author, or some other factor). Finally, the tool will allow future research to assess the

impact of peer review and editorial guidelines on manuscripts through comparison of the same paper at two stages (preprinted versus published). This approach was used to explore the impact of editorial policies on open data sharing, as discussed in Chapter 2, but could be applied to other aspects of the publication process.

8.7.4 Reviewing Mendelian randomisation studies

As noted in Chapter 4, methods for the inclusion of Mendelian randomisation studies in evidence synthesis exercises are not yet sufficiently developed, particularly in comparison to other study designs. Future work should aim to validate search strategies for this study design, paying particular attention to the range of terms used to define the analytical approach (e.g. genetic instrumental variable analysis). In addition, as discussed in Section 7.5.2, the lack of an established risk-of-bias tool for Mendelian randomisation studies limits subsequent analyses that require systematic assessment of bias, such as the quantitative triangulation framework proposed here. Creation of a domain-based tool for this study design, which accounts for the differences in potential biases between one- and two-sample approaches, should be considered a priority. Finally, the inclusion of multiple two-sample Mendelian randomisation studies using identical summary statistics can falsely increase the precision of the summary effect estimate if treated as independent results. Development of best practice guidance for addressing this problem would represent a useful contribution to the field.

8.7.5 Systematic reviews and quantitative triangulation

Quantitative triangulation is an area ripe for future work. The approach presented here will benefit from future meta-epidemiological studies to better define the impact of bias/indirectness, as discussed in detail in Section 7.5.2. Similarly, future work should expand the framework proposed here to account for meta-biases, such as

8.8 - Overall conclusions

those introduced by missing evidence. Tools such as ROB-ME will enable this, but how to adjust for this analysis-level - as opposed to result-level - bias will need to be determined. Finally, the application of the framework to other causal questions will help to identify sticking points and may lead to refinement of the process.

8.8 Overall conclusions

To conclude, this thesis has provided new evidence concerning the role of blood lipids as a modifiable risk factor for dementia and highlighted the considerable uncertainty that still remains in relation to this causal question. In addition, it has developed new methods and tools, specifically around the inclusion of preprints in systematic reviews and the quantitative triangulation of evidence sources, which will enable future evidence synthesists to better address important causal questions.

Lasciate ogne speranza, voi ch'intrate. . .

9

Bibliography

1. Alzheimer, A. Über eine eigenartige Erkrankung der Hirnrinde. *Zentralbl. Nervenheil. Psych.* **18**, 177–179 (1907).
2. Stelzmann, R. A., Norman Schnitzlein, H. & Reed Murtagh, F. An english translation of alzheimer's 1907 paper, 'über eine eigenartige erkankung der hirnrinde'. *Clinical Anatomy* **8**, 429–431 (1995).
3. Edition, F. & others. Diagnostic and statistical manual of mental disorders. *Am Psychiatric Assoc* **21**, (2013).
4. Cerejeira, J., Lagarto, L. & Mukaetova-Ladinska, E. B. Behavioral and Psychological Symptoms of Dementia. *Frontiers in Neurology* **3**, (2012).
5. Kumar, C. S. & Kuriakose, J. R. End-of-life care issues in advanced dementia. *Mental Health in Family Medicine* **10**, 129–132 (2013).
6. Burns, A. & Iliffe, S. Dementia. *BMJ* **338**, b75 (2009).
7. Robinson, L., Tang, E. & Taylor, J.-P. Dementia: Timely diagnosis and early intervention. *BMJ* **350**, h3029 (2015).
8. Iadecola, C. The pathobiology of vascular dementia. *Neuron* **80**, (2013).
9. Venkat, P., Chopp, M. & Chen, J. Models and Mechanisms of Vascular Dementia. *Experimental neurology* **272**, 97–108 (2015).
10. Custodio, N. *et al.* Mixed dementia: A review of the evidence. *Dementia & Neuropsychologia* **11**, 364–370 (2017).
11. Sheehan, B. Assessment scales in dementia. *Therapeutic Advances in Neurological Disorders* **5**, 349–358 (2012).

12. Dubois, B. *et al.* Research criteria for the diagnosis of Alzheimer's disease: Revising the NINCDSADRDA criteria. *The Lancet Neurology* **6**, 734–746 (2007).
13. Román, G. C. *et al.* Vascular dementia: Diagnostic criteria for research studies: Report of the NINDS-AIREN International Workshop. *Neurology* **43**, 250–250 (1993).
14. Prince, M. *et al.* Recent global trends in the prevalence and incidence of dementia, and survival with dementia. *Alzheimer's Research & Therapy* **8**, 23 (2016).
15. Flier, W. M. van der & Scheltens, P. Epidemiology and risk factors of dementia. *Journal of Neurology, Neurosurgery & Psychiatry* **76**, v2–v7 (2005).
16. Baker, C., Jarrett, T. & Powell, T. *Dementia: Policy, services and statistics overview*. (2019).
17. Leading causes of death, UK - Office for National Statistics.
18. Wittenberg, R. *et al.* The costs of dementia in England. *International Journal of Geriatric Psychiatry* **34**, 1095–1103 (2019).
19. Prince, M., Albanese, E., Guerchet, M., Prina, M. & others. Dementia and risk reduction: An analysis of protective and modifiable factors. *World Alzheimer Report* 66–83 (2014).
20. Cummings, J., Lee, G., Ritter, A., Sabbagh, M. & Zhong, K. Alzheimer's disease drug development pipeline: 2020. *Alzheimer's & Dementia: Translational Research & Clinical Interventions* **6**, e12050 (2020).
21. Hampel, H. *et al.* The cholinergic system in the pathophysiology and treatment of Alzheimer's disease. *Brain* **141**, 1917–1933 (2018).
22. Pariente, A. *et al.* Prevalence of cholinesterase inhibitors in subjects with dementia in Europe. *Pharmacoepidemiology and Drug Safety* **17**, 655–660 (2008).
23. Marucci, G. *et al.* Efficacy of acetylcholinesterase inhibitors in Alzheimer's disease. *Neuropharmacology* 108352 (2020). doi:10.1016/j.neuropharm.2020.108352
24. Francis, P. T., Ramírez, M. J. & Lai, M. K. Neurochemical basis for symptomatic treatment of Alzheimer's disease. *Neuropharmacology* **59**, 221–229 (2010).
25. Winblad, B. *et al.* Defeating Alzheimer's disease and other dementias: A priority for European science and society. *The Lancet Neurology* **15**, 455–532 (2016).
26. Feingold, K. R. Introduction to Lipids and Lipoproteins. in *Endotext* (eds. Feingold, K. R. et al.) (MDText.com, Inc., 2000).
27. Peters, R. *et al.* Combining modifiable risk factors and risk of dementia: A systematic review and meta-analysis. *BMJ Open* **9**, e022846 (2019).

28. Anstey, K. J., Ee, N., Eramudugolla, R., Jagger, C. & Peters, R. A Systematic Review of Meta-Analyses that Evaluate Risk Factors for Dementia to Evaluate the Quantity, Quality, and Global Representativeness of Evidence. *Journal of Alzheimer's Disease* **70**, S165–S186 (2019).
29. Hughes, D. *et al.* Association of blood pressure lowering with incident dementia or cognitive impairment: A systematic review and meta-analysis. *Jama* **323**, 1934–1944 (2020).
30. Norton, S., Matthews, F. E., Barnes, D. E., Yaffe, K. & Brayne, C. Potential for primary prevention of Alzheimer's disease: An analysis of population-based data. *The Lancet Neurology* **13**, 788–794 (2014).
31. Pushpakom, S. *et al.* Drug repurposing: Progress, challenges and recommendations. *Nature Reviews Drug Discovery* **18**, 41–58 (2019).
32. Laufs, U., Parhofer, K. G., Ginsberg, H. N. & Hegele, R. A. Clinical review on triglycerides. *European Heart Journal* **41**, 99–109c (2020).
33. Zampelas, A. & Magriplis, E. New Insights into Cholesterol Functions: A Friend or an Enemy? *Nutrients* **11**, (2019).
34. Friedewald, W. T., Levy, R. I. & Fredrickson, D. S. Estimation of the Concentration of Low-Density Lipoprotein Cholesterol in Plasma, Without Use of the Preparative Ultracentrifuge. *Clinical Chemistry* **18**, 499–502 (1972).
35. Detection, N. C. E. P. (US). E. P. on. *Third report of the National Cholesterol Education Program (NCEP) Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III)*. (The Program, 2002).
36. Nelson, R. H. Hyperlipidemia as a Risk Factor for Cardiovascular Disease. *Primary care* **40**, 195–211 (2013).
37. Libby, P. *et al.* Atherosclerosis. *Nature Reviews Disease Primers* **5**, 1–18 (2019).
38. Collins, R. *et al.* Interpretation of the evidence for the efficacy and safety of statin therapy. *The Lancet* **388**, 2532–2561 (2016).
39. Law, M. R., Wald, N. J. & Rudnicka, A. R. Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: Systematic review and meta-analysis. *BMJ* **326**, 1423 (2003).
40. Schachter, M. Chemical, pharmacokinetic and pharmacodynamic properties of statins: An update. *Fundamental & Clinical Pharmacology* **19**, 117–125 (2005).
41. Sierra, S. *et al.* Statins as neuroprotectants: A comparative in vitro study of lipophilicity, blood-brain-barrier penetration, lowering of brain cholesterol, and decrease of neuron cell death. *Journal of Alzheimer's disease: JAD* **23**, 307–318 (2011).
42. Kosoglou, T. *et al.* Ezetimibe: A review of its metabolism, pharmacokinetics and drug interactions. *Clinical Pharmacokinetics* **44**, 467–494 (2005).

43. Genest, J. Combination of statin and ezetimibe for the treatment of dyslipidemias and the prevention of coronary artery disease. *The Canadian Journal of Cardiology* **22**, 863–867 (2006).
44. Chaudhary, R., Garg, J., Shah, N. & Sumner, A. PCSK9 inhibitors: A new era of lipid lowering therapy. *World Journal of Cardiology* **9**, 76–91 (2017).
45. McKenney, J. New perspectives on the use of niacin in the treatment of lipid disorders. *Archives of internal medicine* **164**, 697–705 (2004).
46. Skulas-Ray Ann C. *et al.* Omega-3 Fatty Acids for the Management of Hypertriglyceridemia: A Science Advisory From the American Heart Association. *Circulation* **140**, e673–e691 (2019).
47. Burns, M. P. *et al.* Co-localization of cholesterol, apolipoprotein E and fibrillar A β in amyloid plaques. *Molecular Brain Research* **110**, 119–125 (2003).
48. Mizuno, T. *et al.* Cholesterol-dependent Generation of a Seeding Amyloid β -Protein in Cell Culture *. *Journal of Biological Chemistry* **274**, 15110–15114 (1999).
49. Beecham, G. W. *et al.* Genome-Wide Association Meta-analysis of Neuropathologic Features of Alzheimer’s Disease and Related Dementias. *PLoS Genetics* **10**, (2014).
50. Harold, D. *et al.* Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer’s disease, and shows evidence for additional susceptibility genes. *Nature genetics* **41**, 1088–1093 (2009).
51. Meng, Y. *et al.* Association between SORL1 and Alzheimer disease in a genome-wide study. *Neuroreport* **18**, 1761–1764 (2007).
52. Kivipelto, M. *et al.* Apolipoprotein E E4 Allele, Elevated Midlife Total Cholesterol Level, and High Midlife Systolic Blood Pressure Are Independent Risk Factors for Late-Life Alzheimer Disease. *Annals of Internal Medicine* **137**, 149–155 (2002).
53. Kivipelto, M. *et al.* Obesity and vascular risk factors at midlife and the risk of dementia and Alzheimer disease. *Archives of Neurology* **62**, 1556–1560 (2005).
54. Schilling, S. *et al.* Differential associations of plasma lipids with incident dementia and dementia subtypes in the 3C Study: A longitudinal, population-based prospective cohort study. *PLoS medicine* **14**, e1002265 (2017).
55. Solomon, A., Kivipelto, M., Wolozin, B., Zhou, J. & Whitmer, R. A. Midlife Serum Cholesterol and Increased Risk of Alzheimer’s and Vascular Dementia Three Decades Later. *Dementia and Geriatric Cognitive Disorders* **28**, 75–80 (2009).
56. Whitmer, R. A., Sidney, S., Selby, J., Johnston, S. C. & Yaffe, K. Midlife cardiovascular risk factors and risk of dementia in late life. *Neurology* **64**, 277–281 (2005).
57. Li, G. *et al.* Serum cholesterol and risk of Alzheimer disease: A community-based cohort study. *Neurology* **65**, 1045–1050 (2005).

58. Mainous, A. G., Eschenbach, S. L., Wells, B. J., Everett, C. J. & Gill, J. M. Cholesterol, transferrin saturation, and the development of dementia and Alzheimer's disease: Results from an 18-year population-based cohort. *Family Medicine* **37**, 36–42 (2005).
59. Mielke, M. M. *et al.* The 32-year relationship between cholesterol and dementia from midlife to late life. *Neurology* **75**, 1888–1895 (2010).
60. Tan, Z. S. *et al.* Plasma Total Cholesterol Level as a Risk Factor for Alzheimer Disease: The Framingham Study. *Archives of Internal Medicine* **163**, 1053 (2003).
61. Mielke, M. M. *et al.* High total cholesterol levels in late life associated with a reduced risk of dementia. *Neurology* **64**, 1689–1695 (2005).
62. Reitz, C., Tang, M.-X., Luchsinger, J. & Mayeux, R. Relation of plasma lipids to Alzheimer disease and vascular dementia. *Archives of Neurology* **61**, 705–714 (2004).
63. Moroney, J. T. Low-Density Lipoprotein Cholesterol and the Risk of Dementia With Stroke. *JAMA* **282**, 254 (1999).
64. Ritchie, C. W., Terrera, G. M. & Quinn, T. J. Dementia trials and dementia tribulations: Methodological and analytical challenges in dementia research. *Alzheimer's Research & Therapy* **7**, 31 (2015).
65. McGuinness, B., Craig, D., Bullock, R. & Passmore, P. Statins for the prevention of dementia. *The Cochrane Database of Systematic Reviews* CD003160 (2016). doi:10.1002/14651858.CD003160.pub3
66. Trompet, S. *et al.* Pravastatin and cognitive function in the elderly. Results of the PROSPER study. *Journal of Neurology* **257**, 85–90 (2010).
67. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20 536 high-risk individuals: A randomised placebocontrolled trial. *The Lancet* **360**, 7–22 (2002).
68. Davey Smith, G. & Hemani, G. Mendelian randomization: Genetic anchors for causal inference in epidemiological studies. *Human Molecular Genetics* **23**, R89–R98 (2014).
69. Greenland, S. An introduction to instrumental variables for epidemiologists. *International Journal of Epidemiology* **29**, 722–729 (2000).
70. Davies, N. M., Holmes, M. V. & Smith, G. D. Reading Mendelian randomisation studies: A guide, glossary, and checklist for clinicians. *BMJ* **362**, k601 (2018).
71. Larsson, S. C. *et al.* Modifiable pathways in Alzheimer's disease: Mendelian randomisation analysis. *BMJ* **359**, (2017).
72. Østergaard, S. D. *et al.* Associations between Potentially Modifiable Risk Factors and Alzheimer Disease: A Mendelian Randomization Study. *PLOS Medicine* **12**, e1001841 (2015).
73. Kim, J., Basak, J. M. & Holtzman, D. M. The Role of Apolipoprotein E in Alzheimer's Disease. *Neuron* **63**, 287–303 (2009).

74. Benn, M., Nordestgaard, B. G., Frikke-Schmidt, R. & Tybjærg-Hansen, A. Low LDL cholesterol, PCSK9 and HMGCR genetic variation, and risk of Alzheimer's disease and Parkinson's disease: Mendelian randomisation study. *BMJ* **357**, j1648 (2017).
75. Donnelly, C. A. *et al.* Four principles to make evidence synthesis more useful for policy. *Nature* **558**, 361–364 (2018).
76. Chandler, J., Higgins, J., Deeks, J., Davenport, C. & Clarke, M. J. Chapter 1: introduction. in *Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021)* **5**, (Cochrane, 2020).
77. Paez, A. Grey literature: An important resource in systematic reviews. *Journal of Evidence-Based Medicine* (2017). doi:10.1111/jebm.12265
78. Conn, V. S., Valentine, J. C., Cooper, H. M. & Rantz, M. J. Grey literature in meta-analyses. *Nursing Research* **52**, 256–261 (2003 Jul-Aug).
79. McAuley, L., Pham, B., Tugwell, P. & Moher, D. Does the inclusion of grey literature influence estimates of intervention effectiveness reported in meta-analyses? *Lancet (London, England)* **356**, 1228–1231 (2000).
80. Hopewell, S., McDonald, S., Clarke, M. & Egger, M. Grey literature in meta-analyses of randomized trials of health care interventions. *The Cochrane Database of Systematic Reviews* MR000010 (2007). doi:10.1002/14651858.MR000010.pub3
81. Lefebvre, C. *et al.* Chapter 4: Searching for and selecting studies. in *Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021)* 67–107 (Cochrane, 2020).
82. Committee on Publication Ethics (COPE). *Discussion Document: preprints*. (2018).
83. Vale, R. D. & Hyman, A. A. Priority of discovery in the life sciences. *eLife* **5**, e16931 (2016).
84. Fraser, N. *et al.* Preprinting a pandemic: The role of preprints in the COVID-19 pandemic. *bioRxiv* 2020.05.22.111294 (2020). doi:10.1101/2020.05.22.111294
85. Rosenthal, R. The file drawer problem and tolerance for null results. *Psychological Bulletin* **86**, 638–641 (1979).
86. Maslove, D. M. Medical PreprintsA Debate Worth Having. *JAMA* **319**, 443–444 (2018).
87. Schalkwyk, M. C. I. van, Hird, T. R., Maani, N., Petticrew, M. & Gilmore, A. B. The perils of preprints. *BMJ* **370**, m3111 (2020).
88. Mahood, Q., Eerd, D. V. & Irvin, E. Searching for grey literature for systematic reviews: Challenges and benefits. *Research Synthesis Methods* **5**, 221–234 (2014).
89. Shi, X. *et al.* Assessment of Concordance and Discordance Among Clinical Studies Posted as Preprints and Subsequently Published in High-Impact Journals. *JAMA Network Open* **4**, e212110 (2021).

90. Klein, M., Broadwell, P., Farb, S. E. & Grappone, T. Comparing published scientific journal articles to their pre-print versions. *International Journal on Digital Libraries* **20**, 335–350 (2019).
91. Nicholson, D. N. *et al.* Linguistic Analysis of the bioRxiv Preprint Landscape. *bioRxiv* 2021.03.04.433874 (2021). doi:10.1101/2021.03.04.433874
92. Riley, R. D., Lambert, P. C. & Abo-Zaid, G. Meta-analysis of individual participant data: Rationale, conduct, and reporting. *BMJ* **340**, c221 (2010).
93. Stewart, L. A. & Parmar, M. K. B. Meta-analysis of the literature or of individual patient data: Is there a difference? *The Lancet* **341**, 418–422 (1993).
94. Arain, F. A., Kuniyoshi, F. H., Abdalrhim, A. D. & Miller, V. M. Sex/gender medicine. The biological basis for personalized care in cardiovascular medicine. *Circulation Journal: Official Journal of the Japanese Circulation Society* **73**, 1774–1782 (2009).
95. Clayton, J. A. Applying the new SABV (sex as a biological variable) policy to research and clinical care. *Physiology & Behavior* **187**, 2–5 (2018).
96. McCartney, M., Treadwell, J., Maskrey, N. & Lehman, R. Making evidence based medicine work for individual patients. *BMJ* **353**, i2452 (2016).
97. Letenneur, L. *et al.* Are sex and educational level independent predictors of dementia and Alzheimer's disease? Incidence data from the PAQUID project. *Journal of Neurology, Neurosurgery & Psychiatry* **66**, 177–183 (1999).
98. Riley, R. D. *et al.* Individual participant data meta-analysis to examine interactions between treatment effect and participant-level covariates: Statistical recommendations for conduct and planning. *Statistics in Medicine* **39**, 2115–2137 (2020).
99. Stewart, L. A. & Tierney, J. F. To IPD or not to IPD?: Advantages and Disadvantages of Systematic Reviews Using Individual Patient Data. *Evaluation & the Health Professions* **25**, 76–97 (2002).
100. Tugwell, P. & Knottnerus, J. A. Advantages of individual patient data analysis in systematic reviews. *Journal of Clinical Epidemiology* **63**, 233–234 (2010).
101. Nevitt, S. J. *et al.* Exploring changes over time and characteristics associated with data retrieval across individual participant data meta-analyses: Systematic review. *BMJ* **357**, j1390 (2017).
102. Ventresca, M. *et al.* Obtaining and managing data sets for individual participant data meta-analysis: Scoping review and practical guide. *BMC Medical Research Methodology* **20**, 113 (2020).
103. Alsheikh-Ali, A. A., Qureshi, W., Al-Mallah, M. H. & Ioannidis, J. P. A. Public Availability of Published Research Data in High-Impact Journals. *PLoS ONE* **6**, e24357 (2011).
104. Federer, L. M. *et al.* Data sharing in PLOS ONE: An analysis of Data Availability Statements. *PLOS ONE* **13**, e0194768 (2018).

105. Vines, T. H. *et al.* The Availability of Research Data Declines Rapidly with Article Age. *Current Biology* **24**, 94–97 (2014).
106. van Panhuis, W. G. *et al.* A systematic review of barriers to data sharing in public health. *BMC Public Health* **14**, 1144 (2014).
107. Wartenberg, D. & Thompson, W. D. Privacy Versus Public Health: The Impact of Current Confidentiality Rules. *American Journal of Public Health* **100**, 407–412 (2010).
108. Bauermeister, S. *et al.* The Dementias Platform UK (DPUK) Data Portal. *European Journal of Epidemiology* **35**, 601–611 (2020).
109. Lawlor, D. A., Tilling, K. & Davey Smith, G. Triangulation in aetiological epidemiology. *International Journal of Epidemiology* **45**, 1866–1886 (2016).
110. Munafò, M. R. & Smith, G. D. Robust research needs many lines of evidence. *Nature* **553**, 399–401 (2018).
111. Rawlinson, C. & Bloom, T. New preprint server for medical research. *BMJ* **365**, (2019).
112. Sever, R. *et al.* *bioRxiv: The preprint server for biology.* (Scientific Communication and Education, 2019). doi:10.1101/833400
113. Bealy, C. R Programming Humour. *Stack Exchange* (2013).
114. Bramer, W. M., de Jonge, G. B., Rethlefsen, M. L., Mast, F. & Kleijnen, J. A systematic approach to searching: An efficient and complete method to develop literature searches. *Journal of the Medical Library Association : JMLA* **106**, 531–541 (2018).
115. Gusenbauer, M. & Haddaway, N. R. Which academic search systems are suitable for systematic reviews or meta-analyses? Evaluating retrieval qualities of Google Scholar, PubMed, and 26 other resources. *Research Synthesis Methods* **11**, 181–217 (2020).
116. Ropensci/medrxivr: Access and search medRxiv and bioRxiv preprint data.
117. Wateridge, J. IT projects: A basis for success. *International Journal of Project Management* **13**, 169–172 (1995).
118. Abdill, R. J. & Blekhman, R. Rxivist.org: Sorting biology preprints using social media and readership metrics. *PLOS Biology* **17**, e3000269 (2019).
119. Rxivist: Explore popular biology preprints.
120. Iwema, C. L., LaDue, J., Zack, A. & Chattopadhyay, A. Search.bioPreprint: A discovery tool for cutting edge, preprint biomedical research articles. *F1000Research* **5**, 1396 (2016).
121. Shaw, M. "Self-healing": Softening precision to avoid brittleness: Position paper for WOSS '02: Workshop on self-healing systems. *Proceedings of the first workshop on Self-healing systems* 111–114 (2002). doi:10.1145/582128.582152

122. Laprie, J. C. Dependability: Basic Concepts and Terminology. in *Dependability: Basic Concepts and Terminology: In English, French, German, Italian and Japanese* (ed. Laprie, J. C.) 3–245 (Springer, 1992).
123. R Core Team. *R: A language and environment for statistical computing*. (R Foundation for Statistical Computing, 2019).
124. Git scraping: Track changes over time by scraping to a Git repository.
125. Rethlefsen, M. L. *et al.* PRISMA-S: An extension to the PRISMA statement for reporting literature searches in systematic reviews. *Systematic reviews* **10**, 1–19 (2021).
126. Kodvanj, I., Homolak, J., Virag, D. & Trkulja, V. Publishing of COVID-19 Preprints in Peer-reviewed Journals, Preprinting Trends, Public Discussion and Quality Issues. *bioRxiv* 2020.11.23.394577 (2020). doi:10.1101/2020.11.23.394577
127. Noone, C. *et al.* Investigating and evaluating evidence of the behavioural determinants of adherence to social distancing measures A protocol for a scoping review of COVID-19 research. *HRB Open Research* **3**, 46 (2020).
128. Grassly, N. C. *et al.* Comparison of molecular testing strategies for COVID-19 control: A mathematical modelling study. *The Lancet Infectious Diseases* **20**, 1381–1389 (2020).
129. McGuinness, L. A. & Sheppard, A. L. A descriptive analysis of the data availability statements accompanying medRxiv preprints and a comparison with their published counterparts. *MetaArXiv* (2020). doi:10.31222/osf.io/p75xe
130. Boettiger, C., Chamberlain, S., Hart, E. & Ram, K. Building Software, Building Community: Lessons from the rOpenSci Project. *Journal of Open Research Software* **3**, e8 (2015).
131. McGuinness, L. & Schmidt, L. Medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R. *Journal of Open Source Software* **5**, 2651 (2020).
132. Issue #380 - medrxivr: Accessing and searching medRxiv preprint data in R · · ropensci/software-review · GitHub. *GitHub · ropensci/software-review*
133. Abdill, R. J. & Blekhman, R. Tracking the popularity and outcomes of all bioRxiv preprints. *eLife* **8**, e45133 (2019).
134. Bong, S. M., McKay, J. L., Factor, S. A. & Ting, L. H. *Perception of whole-body motion during balance perturbations is impaired in Parkinson's disease and is associated with balance impairment*. (Neurology, 2019). doi:10.1101/19000265
135. Song, F. *et al.* Dissemination and publication of research findings: An updated review of related biases. *Health Technology Assessment* **14**, (2010).
136. Goldacre, B., Morton, C. E. & DeVito, N. J. Why researchers should share their analytic code. *BMJ* l6365 (2019). doi:10.1136/bmj.l6365
137. McKiernan, E. C. *et al.* How open science helps researchers succeed. *eLife* **5**, e16800 (2016).

138. Fraser, N. *Rbiorxiv: Client for the 'bioRxiv' API*. (2020).
139. Vuorre, M. & Crump, M. J. C. Sharing and organizing research products as R packages. *Behavior Research Methods* (2020). doi:10.3758/s13428-020-01436-x
140. Cochrane, A. L. 1931-1971: A critical review, with particular reference to the medical profession. *Medicines for the year 2000* (1979).
141. McGuinness, L., Higgins, J., Ben-Shlomo, Yoav, Coulthard, L. & Smith, G. The relationship between blood lipid levels and dementia: Protocol for a systematic review and triangulation of all available evidence. *OSF* (2020). doi:10.17605/OSF.IO/VTW5Y
142. Organization(WHO), W. H. *The ICD-10 classification of mental and behavioural disorders: Diagnostic criteria for research*. (World Health Organization, 1993).
143. Tennant, P. W. G., Arnold, K. F., Ellison, G. T. H. & Gilthorpe, M. S. Analyses of “change scores” do not estimate causal effects in observational data. *International Journal of Epidemiology* (2021). doi:10.1093/ije/dyab050
144. Greenhalgh, T. & Peacock, R. Effectiveness and efficiency of search methods in systematic reviews of complex evidence: Audit of primary sources. *BMJ : British Medical Journal* **331**, 1064–1065 (2005).
145. Wohlin, C. Guidelines for snowballing in systematic literature studies and a replication in software engineering. in *Proceedings of the 18th International Conference on Evaluation and Assessment in Software Engineering - EASE '14* 1–10 (ACM Press, 2014). doi:10.1145/2601248.2601268
146. Bramer, W. M., Giustini, D., de Jonge, G. B., Holland, L. & Bekhuis, T. De-duplication of database search results for systematic reviews in EndNote. *Journal of the Medical Library Association : JMLA* **104**, 240–243 (2016).
147. Hupe, M. EndNote X9. *Journal of Electronic Resources in Medical Libraries* **16**, 117–119 (2019).
148. Ouzzani, M., Hammady, H., Fedorowicz, Z. & Elmagarmid, A. Rayyan-a web and mobile app for systematic reviews. *Systematic Reviews* **5**, 210 (2016).
149. Page, M. J. *et al.* The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ* **372**, n71 (2021).
150. Gwet, K. L. Computing inter-rater reliability and its variance in the presence of high agreement. *British Journal of Mathematical and Statistical Psychology* **61**, 29–48 (2008).
151. Cohen, J. A Coefficient of Agreement for Nominal Scales. *Educational and Psychological Measurement* **20**, 37–46 (1960).
152. Wongpakaran, N., Wongpakaran, T., Wedding, D. & Gwet, K. L. A comparison of Cohen’s Kappa and Gwet’s AC1 when calculating inter-rater reliability coefficients: A study conducted with personality disorder samples. *BMC Medical Research Methodology* **13**, 61 (2013).

153. Brennan, P. & Silman, A. Statistical methods for assessing observer variability in clinical measures. *BMJ : British Medical Journal* **304**, 1491–1494 (1992).
154. McHugh, M. L. Interrater reliability: The kappa statistic. *Biochemia Medica* **22**, 276–282 (2012).
155. Higgins, J. P., Li, T. & Deeks, J. J. Section 6.5.2.10 : Combining groups. in *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated August 2019)* (eds. Higgins, J. P. T. et al.) (Cochrane, 2019).
156. Rugge, B. *et al.* Appendix A: Lipid Conversion Factors. in *Screening and Treatment of Subclinical Hypothyroidism or Hyperthyroidism* (Agency for Healthcare Research and Quality (US), 2011).
157. Reynders, R. M., Ladu, L. & Girolamo, N. D. Contacting of authors modified crucial outcomes of systematic reviews but was poorly reported, not systematic, and produced conflicting results. *Journal of Clinical Epidemiology* **115**, 64–76 (2019).
158. McGuinness, L. A., Higgins, J. P. T. & Sterne, J. A. C. Assessing the Credibility of Findings From Nonrandomized Studies of Interventions. *JAMA Cardiology* **3**, 905–906 (2018).
159. Sterne, J. A. *et al.* ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* **355**, i4919 (2016).
160. Campbell, D. T. Factors relevant to the validity of experiments in social settings. *Psychological Bulletin* **54**, 297–312 (1957).
161. Jüni, P., Altman, D. G. & Egger, M. Assessing the quality of controlled clinical trials. *BMJ* **323**, 42–46 (2001).
162. Sterne, J. A. C. *et al.* RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ* **366**, (2019).
163. Morgan R *et al.* A tool to assess Risk Of Bias In Non-randomised Studies - of Exposures (ROBINS-E). in *Advances in Evidence Synthesis: Special issue, Cochrane Database of Systematic Reviews* **9 Supp 1**, 320 (2020).
164. French, C. E. *et al.* Cannabis use and the risk of tuberculosis: A systematic review. *BMC Public Health* **19**, 1006 (2019).
165. Wells, G. A. *et al.* The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. (2000).
166. Mamluk, L. *et al.* Evidence of detrimental effects of prenatal alcohol exposure on offspring birthweight and neurodevelopment from a systematic review of quasi-experimental studies. *International Journal of Epidemiology* **49**, 1972–1995 (2020).
167. Spiga, F. *et al.* Tools for the assessment of quality and risk of bias in Mendelian randomization studies: A systematic review. *medRxiv* 2021.10.21.21265126 (2021). doi:10.1101/2021.10.21.21265126
168. Matthew J Page *et al.* *Risk Of Bias due to Missing Evidence (ROB-ME): A new tool for assessing risk of non-reporting biases in evidence syntheses - Detailed guidance.*

169. McKenzie, J. E. & Brennan, S. E. Section 12.1: Why a meta-analysis of effect estimates may not be possible. in *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated August 2019)* (eds. Higgins, J. P. T. et al.) (Cochrane, 2019).
170. VanderWeele, T. J. Optimal approximate conversions of odds ratios and hazard ratios to risk ratios. *Biometrics* **76**, 746–752 (2020).
171. Hedges, L. V. & Vevea, J. L. Fixed-and random-effects models in meta-analysis. *Psychological methods* **3**, 486 (1998).
172. DerSimonian, R. & Laird, N. Meta-analysis in clinical trials. *Controlled Clinical Trials* **7**, 177–188 (1986).
173. Deeks, J. J., Altman, D. G. & Higgins, J. P. T. Section 10.11.4: Meta-regression. in *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated August 2019)* (eds. Higgins, J. P. T. et al.) (Cochrane, 2019).
174. Viechtbauer, W. *Metafor: Meta-analysis package for r.* (2021).
175. Greenland, S. & Longnecker, M. P. Methods for Trend Estimation from Summarized Dose-Response Data, with Applications to Meta-Analysis. *American Journal of Epidemiology* **135**, 1301–1309 (1992).
176. Durrleman, S. & Simon, R. Flexible regression models with cubic splines. *Statistics in Medicine* **8**, 551–561 (1989).
177. Liu, Q., Cook, N. R., Bergström, A. & Hsieh, C.-C. A two-stage hierarchical regression model for meta-analysis of epidemiologic nonlinear doseresponse data. *Computational Statistics & Data Analysis* **53**, 4157–4167 (2009).
178. Gauthier, J., Wu, Q. V. & Gooley, T. A. Cubic splines to model relationships between continuous variables and outcomes: A guide for clinicians. *Bone Marrow Transplantation* **55**, 675–680 (2020).
179. White, I. R. Multivariate random-effects meta-analysis. *The Stata Journal* **9**, 40–56 (2009).
180. Gasparrini, A., Armstrong, B. & Kenward, M. G. Multivariate meta-analysis for non-linear and other multi-parameter associations. *Statistics in Medicine* **31**, 3821–3839 (2012).
181. Crippa, A. & Orsini, N. Multivariate dose-response meta-analysis: The dosresmeta R package. *Journal of Statistical Software, Code Snippets* **72**, 1–15 (2016).
182. Sterne, J. A. C. *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* **343**, d4002 (2011).
183. Saran, A. & White, H. Evidence and gap maps: A comparison of different approaches. *Campbell Systematic Reviews* **14**, 1–38 (2018).

184. Marušić, M. F. *et al.* Methodological tools and sensitivity analysis for assessing quality or risk of bias used in systematic reviews published in the high-impact anesthesiology journals. *BMC Medical Research Methodology* **20**, 121 (2020).
185. Katikireddi, S. V., Egan, M. & Petticrew, M. How do systematic reviews incorporate risk of bias assessments into the synthesis of evidence? A methodological study. *Journal of Epidemiology and Community Health* **69**, 189–195 (2015).
186. McGuinness, L. A. & Higgins, J. P. T. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Research Synthesis Methods* **12**, 55–61 (2021).
187. Feinstein, A. R. & Cicchetti, D. V. High agreement but low kappa: I. The problems of two paradoxes. *Journal of Clinical Epidemiology* **43**, 543–549 (1990).
188. Ridker, P. M. *et al.* Rosuvastatin to Prevent Vascular Events in Men and Women with Elevated C-Reactive Protein. *New England Journal of Medicine* **359**, 2195–2207 (2008).
189. Ancelin, M.-L. *et al.* Lipid lowering agents, cognitive decline, and dementia: The three-city study. *Journal of Alzheimer's Disease* **30**, 629–637 (2012).
190. Arvanitakis, Z. *et al.* Statins, incident Alzheimer disease, change in cognitive function, and neuropathology. *Neurology* **70**, 1795–1802 (2008).
191. Bettermann, K. *et al.* Statins, Risk of Dementia, and Cognitive Function: Secondary Analysis of the Ginkgo Evaluation of Memory Study. *Journal of Stroke and Cerebrovascular Diseases* **21**, 436–444 (2012).
192. Beydoun, M. A. *et al.* Statins and serum cholesterol's associations with incident dementia and mild cognitive impairment. *Journal of Epidemiology & Community Health* **65**, 949–957 (2011).
193. Chao, T.-F. *et al.* Statins and the risk of dementia in patients with atrial fibrillation: A nationwide population-based cohort study. *International Journal of Cardiology* **196**, 91–97 (2015).
194. Chen, J. M. *et al.* Effects of statins on incident dementia in patients with type 2 DM: A population-based retrospective cohort study in Taiwan. *PLoS ONE [Electronic Resource]* **9**, e88434 (2014).
195. Chitnis, A. S. *et al.* Use of statins and risk of dementia in heart failure: A retrospective cohort study. *Drugs & Aging* **32**, 743–754 (2015).
196. Chou, C.-Y., Chou, Y.-C., Chou, Y.-J., Yang, Y.-F. & Huang, N. Statin use and incident dementia: A nationwide cohort study of Taiwan. *International Journal of Cardiology* **173**, 305–310 (2014).
197. Chuang, C. S., Lin, C. L., Lin, M. C., Sung, F. C. & Kao, C. H. Decreased prevalence of dementia associated with statins: A national population-based study. *European Journal of Neurology* **22**, 912–918 (2015).

198. Cramer, C., Haan, M. N., Galea, S., Langa, K. M. & Kalbfleisch, J. D. Use of statins and incidence of dementia and cognitive impairment without dementia in a cohort study. *Neurology* **71**, 344–350 (2008).
199. Gnjidic, D. *et al.* STATIN THERAPY AND DEMENTIA IN OLDER ADULTS: ROLE OF DISEASE SEVERITY AND MULTIMORBIDITY. *Journal of the American Geriatrics Society* **64**, 223–224 (2016).
200. Haag, M. D., Hofman, A., Koudstaal, P. J., Stricker, B. H. & Breteler, M. M. Statins are associated with a reduced risk of Alzheimer disease regardless of lipophilicity. The Rotterdam Study. *Journal of Neurology, Neurosurgery & Psychiatry* **80**, 13–17 (2009).
201. Hendrie, H. C. *et al.* Statin use, incident dementia and alzheimer disease in elderly african americans. *Ethnicity & Disease* **25**, 345–354 (2015).
202. Hippisley-Cox, J. & Coupland, C. Unintended effects of statins in men and women in England and Wales: Population based cohort study using the QResearch database. *BMJ* **340**, (2010).
203. Jick, H., Zornberg, G. L., Jick, S. S., Seshadri, S. & Drachman, D. A. Statins and the risk of dementia. *Lancet (London, England)* **356**, 1627–1631 (2000).
204. Li, G. *et al.* Statin therapy and risk of dementia in the elderly: A community-based prospective cohort study. *Neurology* **63**, 1624–1628 (2004).
205. Li, G. *et al.* Age-varying association between statin use and incident Alzheimer's disease. *Journal of the American Geriatrics Society* **58**, 1311–1317 (2010).
206. Liao, M. T., Tsai, C. T. & Lin, J. L. Statins reduce the incidence of dementia in patients with atrial fibrillation: A nationwide cohort study. *European Heart Journal* **34**, 740–741 (2013).
207. Liu, D. *et al.* Incentive schemes to increase dementia diagnoses in primary care in England: A retrospective cohort study of unintended consequences. *British Journal of General Practice* **69**, e154–e163 (2019).
208. Pan, M. L., Hsu, C. C., Chen, Y. M., Yu, H. K. & Hu, G. C. Statin use and the risk of dementia in patients with stroke: A nationwide population-based cohort study. *Journal of Stroke & Cerebrovascular Diseases* **27**, 3001–3007 (2018).
209. Parikh, N. M. *et al.* Risk factors for dementia in patients over 65 with diabetes. *International Journal of Geriatric Psychiatry* **26**, 749–757 (2011).
210. Rea, T. D. *et al.* Statin Use and the Risk of Incident Dementia: The Cardiovascular Health Study. *Archives of Neurology* **62**, 1047–1051 (2005).
211. Redelmeier, D. A., Manzoor, F. & Thiruchelvam, D. Association between statin use and risk of dementia after a concussion. *JAMA Neurology* **20**, 20 (2019).
212. Reitz, C. *et al.* Association of higher levels of high-density lipoprotein cholesterol in elderly individuals and lower risk of late-onset Alzheimer disease. *Archives of Neurology* **67**, 1491–1497 (2010).

213. Smeeth, L., Douglas, I., Hall, A. J., Hubbard, R. & Evans, S. Effect of statins on a wide range of health outcomes: A cohort study validated by comparison with randomized trials. *British Journal of Clinical Pharmacology* **67**, 99–109 (2009).
214. Solomon, A. *et al.* Serum cholesterol changes after midlife and late-life cognition: Twenty-one-year follow-up study. *Neurology* **68**, 751–756 (2007).
215. Sparks, D. L. *et al.* Reduced risk of incident AD with elective statin use in a clinical trial cohort. *Current alzheimer research* **5**, 416–421 (2008).
216. Szwest, S. J. *et al.* Association of statin use with cognitive decline in elderly African Americans. *Neurology* **69**, 1873–1880 (2007).
217. Yang, Y. H. *et al.* Statins reduces the risk of dementia in patients with late-onset depression: A retrospective cohort study. *PLoS ONE [Electronic Resource]* **10**, e0137914 (2015).
218. Zamrini, E., McGwin, G. & Roseman, J. M. Association between statin use and Alzheimer's disease. *Neuroepidemiology* **23**, 94–98 (2004).
219. Zandi, P. P. *et al.* Do statins reduce risk of incident dementia and Alzheimer disease? The Cache County Study. *Archives of General Psychiatry* **62**, 217–224 (2005).
220. Ancelin, M. L. *et al.* Sex differences in the associations between lipid levels and incident dementia. *Journal of Alzheimer's Disease* **34**, 519–528 (2013).
221. Batty, G. D., Russ, T. C., Starr, J. M., Stamatakis, E. & Kivimaki, M. Modifiable cardiovascular disease risk factors as predictors of dementia death: Pooling of ten general population-based cohort studies. *Journal of Negative Results in Biomedicine* **13**, 8 (2014).
222. Bruce, D. G., Davis, W. A. & Davis, T. M. E. Low serum HDL-cholesterol concentrations in mid-life predict late-life cognitive impairment in type 2 diabetes: The Fremantle diabetes study. *Journal of Diabetes & its Complications* **31**, 945–947 (2017).
223. Chiang, C. J. *et al.* Midlife risk factors for subtypes of dementia: A nested case-control study in Taiwan. *American Journal of Geriatric Psychiatry* **15**, 762–771 (2007).
224. Dodge, H. H., Chang, C. C. H., Kamboh, I. M. & Ganguli, M. Risk of Alzheimer's disease incidence attributable to vascular disease in the population. *Alzheimer's and Dementia* **7**, 356–360 (2011).
225. Forti, P. *et al.* Metabolic syndrome and risk of dementia in older adults. *Journal of the American Geriatrics Society* **58**, 487–492 (2010).
226. Gottesman, R. F. *et al.* Associations between midlife vascular risk factors and 25-Year incident dementia in the atherosclerosis risk in communities (ARIC) cohort. *JAMA Neurology* **74**, 1246–1254 (2017).
227. Gustafson, D. R. *et al.* Epidemiological evidence for lipid-based dementia prevention: The Lipididiet approach. *Journal of Molecular Neuroscience* **1**), S45 (2012).

228. Hayden, K. M. *et al.* Vascular risk factors for incident Alzheimer disease and vascular dementia - The Cache County study. *Alzheimer Disease & Associated Disorders* **20**, 93–100 (2006).
229. Kimm, H. *et al.* Mid-life and late-life vascular risk factors and dementia in Korean men and women. *Archives of Gerontology & Geriatrics* **52**, e117–122 (2011).
230. Kivipelto, M. *et al.* Midlife vascular risk factors and Alzheimer's disease in later life: Longitudinal, population based study. *BMJ (Clinical research ed.)* **322**, 1447–1451 (2001).
231. Kuo, S. C. *et al.* Association between comorbidities and dementia in diabetes mellitus patients: Population-based retrospective cohort study. *Journal of Diabetes & its Complications* **29**, 1071–1076 (2015).
232. Mielke, M. *et al.* Serum ceramides predict incident alzheimer's disease: The women's health and aging study (whas) II. *Alzheimer's and Dementia* **1**, S607 (2011).
233. Muller, M. *et al.* Metabolic syndrome and dementia risk in a multiethnic elderly cohort. *Dementia & Geriatric Cognitive Disorders* **24**, 185–192 (2007).
234. Noale, M. *et al.* Incidence of dementia: Evidence for an effect modification by gender. The ILSA Study. *International Psychogeriatrics* **25**, 1867–1876 (2013).
235. Notkola, I. L. *et al.* Serum total cholesterol, apolipoprotein E epsilon 4 allele, and Alzheimer's disease. *Neuroepidemiology* **17**, 14–20 (1998).
236. Peters, R. *et al.* Cardiovascular and biochemical risk factors for incident dementia in the Hypertension in the Very Elderly Trial. *Journal of Hypertension* **27**, 2055–2062 (2009).
237. Raffaitin, C. *et al.* Metabolic syndrome and risk for incident Alzheimer's disease or vascular dementia: The Three-City Study. *Diabetes Care* **32**, 169–174 (2009).
238. Rantanen, K. *et al.* Cardiovascular risk factors and glucose tolerance in midlife and risk of cognitive disorders in old age up to a 49-year follow-up of the Helsinki businessmen study. *Annals of Medicine* **49**, 462–469 (2017).
239. Ronnemaa, E., Zethelius, B., Lannfelt, L. & Kilander, L. Vascular risk factors and dementia: 40-year follow-up of a population-based cohort. *Dementia & Geriatric Cognitive Disorders* **31**, 460–466 (2011).
240. Solomon, A. *et al.* Lipid-lowering treatment is related to decreased risk of dementia: A population-based study (FINRISK). *Neurodegenerative Diseases* **7**, 180–182 (2010).
241. Strand, B. H. *et al.* Midlife vascular risk factors and their association with dementia deaths: Results from a Norwegian prospective study followed up for 35 years. *Journal of the Neurological Sciences* **324**, 124–130 (2013).

242. Su, B. *et al.* [P3]: EXPLORING LATE-LIFE RISK FACTORS OF ALZHEIMER's DISEASE AND OTHER AGE-RELATED DEMENTIAS IN CPRD. *Alzheimer's & Dementia* **13**, P1178–P1179 (2017).
243. Svensson, T. *et al.* The association between midlife serum high-density lipoprotein and mild cognitive impairment and dementia after 19 years of follow-up. *Transl Psychiatry Psychiatry* **9**, 26 (2019).
244. Tynkkynen, J. *et al.* Apolipoproteins and HDL cholesterol do not associate with the risk of future dementia and Alzheimer's disease: The National Finnish population study (FINRISK). *Age (Melbourne, Vic.)* **38**, 465–473 (2016).
245. Tynkkynen, J. *et al.* Association of branched-chain amino acids and other circulating metabolites with risk of incident dementia and Alzheimer's disease: A prospective study in eight cohorts. *Alzheimer's & Dementia* **14**, 723–733 (2018).
246. Wang, K.-C. *et al.* Risk of Alzheimer's disease in relation to diabetes: A population-based cohort study. *Neuroepidemiology* **38**, 237–244 (2012).
247. Yamada, M. *et al.* Incidence and risks of dementia in Japanese women: Radiation Effects Research Foundation Adult Health Study. *Journal of the Neurological Sciences* **283**, 57–61 (2009).
248. Yoshitake, T. *et al.* Incidence and risk factors of vascular dementia and Alzheimer's disease in a defined elderly Japanese population: The Hisayama Study. *Neurology* **45**, 1161–1168 (1995).
249. Zimetbaum, P. *et al.* Plasma lipids and lipoproteins and the incidence of cardiovascular disease in the very elderly. The Bronx Aging Study. *Arteriosclerosis & Thrombosis* **12**, 416–423 (1992).
250. Andrews, S. J. *et al.* Causal Associations Between Modifiable Risk Factors and the Alzheimer's Phenome. *Annals of Neurology* **89**, 54–65 (2021).
251. Burgess, S. & Davey Smith, G. Mendelian randomization implicates high-density lipoprotein cholesterol-associated mechanisms in etiology of age-related macular degeneration. *Ophthalmology* **124**, 1165–1174 (2017).
252. Mukherjee, S. *et al.* Vascular disease, vascular risk factors and risk of late-onset Alzheimer's disease: Mendelian randomization analyses in the combined adgc dataset. *Alzheimer's and Dementia* **1**, P694 (2013).
253. So, H.-C., Chau, C. K. & Zhao, K. Exploring repositioning opportunities and side-effects of statins: A Mendelian randomization study of HMG-CoA reductase inhibition with 55 complex traits. 170241 (2017). doi:10.1101/170241
254. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nature Communications* **9**, 224 (2018).
255. Andrews, S. J. *et al.* Causal associations between potentially modifiable risk factors and the Alzheimer's disease phenotype: A Mendelian randomization study. *bioRxiv* (2019). doi:10.1101/689752
256. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from GWAS summary data. *bioRxiv* (2017). doi:10.1101/168674

257. Zhu, H. *kableExtra: Construct complex table with 'kable' and pipe syntax.* (2020).
258. Shepherd, J. *et al.* Pravastatin in elderly individuals at risk of vascular disease (PROSPER): A randomised controlled trial. *Lancet (London, England)* **360**, 1623–1630 (2002).
259. Yamada, M. *et al.* Incidence and risks of dementia in Japanese women: The adult health study. *Journal of the Neurological Sciences* **283**, 306 (2009).
260. Chu, C.-S. *et al.* Use of statins and the risk of dementia and mild cognitive impairment: A systematic review and meta-analysis. *Scientific Reports* **8**, 5804 (2018).
261. Yang, Z. *et al.* Association of blood lipids, atherosclerosis and statin use with dementia and cognitive impairment after stroke: A systematic review and meta-analysis. *Ageing Research Reviews* **57**, 100962 (2020).
262. Muangpaisan, W., Brayne, C. & the Alzheimer's Society Vascular Dementia Systematic Review Group. Systematic review of statins for the prevention of vascular dementia or dementia. *Geriatrics & Gerontology International* (2010). doi:10.1111/j.1447-0594.2009.00579.x
263. Poly, T. N. *et al.* Association between Use of Statin and Risk of Dementia: A Meta-Analysis of Observational Studies. *Neuroepidemiology* **54**, 214–226 (2020).
264. Kuźma, E. *et al.* Stroke and dementia risk: A systematic review and meta-analysis. *Alzheimer's & dementia : the journal of the Alzheimer's Association* **14**, 1416–1426 (2018).
265. Kuźma, E. *et al.* Which Risk Factors Causally Influence Dementia? A Systematic Review of Mendelian Randomization Studies. *Journal of Alzheimer's disease: JAD* **64**, 181–193 (2018).
266. Power, M. C., Weuve, J., Sharrett, A. R., Blacker, D. & Gottesman, R. F. Statins, cognition, and dementia systematic review and methodological commentary. *Nature reviews. Neurology* **11**, 220–229 (2015).
267. Aromataris, E. *et al.* Summarizing systematic reviews: Methodological development, conduct and reporting of an umbrella review approach. *International Journal of Evidence-Based Healthcare* **13**, 132–140 (2015).
268. Smith, V., Devane, D., Begley, C. M. & Clarke, M. Methodology in conducting a systematic review of systematic reviews of healthcare interventions. *BMC Medical Research Methodology* **11**, 15 (2011).
269. Skrivankova, V. W. *et al.* Strengthening the reporting of observational studies in epidemiology using mendelian randomisation (STROBE-MR): Explanation and elaboration. *BMJ* **375**, n2233 (2021).
270. Sohani, Z. N. *et al.* Assessing the quality of published genetic association studies in meta-analyses: The quality of genetic studies (Q-Genie) tool. *BMC Genetics* **16**, 50 (2015).

271. Waffenschmidt, S. *et al.* Development and validation of study filters for identifying controlled non-randomized studies in PubMed and Ovid MEDLINE. *Research Synthesis Methods* **11**, 617–626 (2020).
272. Wagner, M., Rosumeck, S., Küffmeier, C., Döring, K. & Euler, U. A validation study revealed differences in design and performance of MEDLINE search filters for qualitative research. *Journal of Clinical Epidemiology* **120**, 17–24 (2020).
273. Benn, M. Re: Low LDL cholesterol, PCSK9 and HMGCR genetic variation, and risk of Alzheimer's disease and Parkinson's disease: Mendelian randomisation study. *BMJ* (2017).
274. Meursinge Reynders, R., Ladu, L. & Di Girolamo, N. Contacting of authors modified crucial outcomes of systematic reviews but was poorly reported, not systematic, and produced conflicting results. *Journal of Clinical Epidemiology* **115**, 64–76 (2019).
275. Cooper, C., Bou, J. T. & Varley-Campbell, J. Evaluating the effectiveness, efficiency, cost and value of contacting study authors in a systematic review: A case study and worked example. *BMC Medical Research Methodology* **19**, 45 (2019).
276. Hennessy, E. A. *et al.* Ensuring Prevention Science Research is Synthesis-Ready for Immediate and Lasting Scientific Impact. *Prevention Science* (2021). doi:10.1007/s11121-021-01279-8
277. Hsieh, C.-Y. *et al.* Taiwan's National Health Insurance Research Database: Past and future. *Clinical Epidemiology* **11**, 349–358 (2019).
278. McGuinness, L. A., Warren-Gash, C., Moorhouse, L. R. & Thomas, S. L. The validity of dementia diagnoses in routinely collected electronic health records in the United Kingdom: A systematic review. *Pharmacoepidemiology and Drug Safety* **28**, 244–255 (2019).
279. Wilkinson, T. *et al.* Identifying dementia cases with routinely collected health data: A systematic review. *Alzheimer's & Dementia* **14**, 1038–1051 (2018).
280. Jeyaraman, M. M. *et al.* Methodologically rigorous risk of bias tools for nonrandomized studies had low reliability and high evaluator burden. *Journal of Clinical Epidemiology* **128**, 140–147 (2020).
281. Dorn, H. F. Philosophy of Inferences from Retrospective Studies. *American Journal of Public Health and the Nations Health* **43**, 677–683 (1953).
282. Suissa, S. Immortal Time Bias in Pharmacoepidemiology. *American Journal of Epidemiology* **167**, 492–499 (2008).
283. Walker, V. M., Davies, N. M., Jones, T., Kehoe, P. G. & Martin, R. M. Can commonly prescribed drugs be repurposed for the prevention or treatment of Alzheimer's and other neurodegenerative diseases? Protocol for an observational cohort study in the UK Clinical Practice Research Datalink. *BMJ Open* **6**, (2016).

284. Herrett, E. *et al.* Data Resource Profile: Clinical Practice Research Datalink (CPRD). *International Journal of Epidemiology* **44**, 827–836 (2015).
285. Williams, T., van Staa, T., Puri, S. & Eaton, S. Recent advances in the utility and use of the General Practice Research Database as an example of a UK Primary Care Data resource. *Therapeutic Advances in Drug Safety* **3**, 89–99 (2012).
286. Wood, L. & Coulson, R. Revitalizing the general practice research database: Plans, challenges, and opportunities. *Pharmacoepidemiology and drug safety* **10**, 379–383 (2001).
287. Mathur, R. *et al.* Completeness and usability of ethnicity data in UK-based primary care and hospital databases. *Journal of Public Health (Oxford, England)* **36**, 684–692 (2014).
288. Booth, N. What are the Read Codes? *Health Libraries Review* **11**, 177–182 (1994).
289. Wolf, A. *et al.* Data resource profile: Clinical Practice Research Datalink (CPRD) Aurum. *International Journal of Epidemiology* **48**, 1740–1740g (2019).
290. Wishart, D. S. *et al.* DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic Acids Research* **46**, D1074–D1082 (2017).
291. Walker, V. M., Davies, N. M., Martin, R. M. & Kehoe, P. G. Comparison of Antihypertensive Drug Classes for Dementia Prevention. *Epidemiology* **31**, 852–859 (2020).
292. Khan, N. F., Perera, R., Harper, S. & Rose, P. W. Adaptation and validation of the Charlson Index for Read/OXMIS coded databases. *BMC Family Practice* **11**, 1 (2010).
293. Taylor, G. M. J. *et al.* Effectiveness of varenicline versus nicotine replacement therapy on long-term smoking cessation in primary care: A prospective, cohort study of electronic medical records. *The Lancet* **388**, S107 (2016).
294. Wright, A. K. *et al.* Life Expectancy and Cause-Specific Mortality in Type 2 Diabetes: A Population-Based Cohort Study Quantifying Relationships in Ethnic Subgroups. *Diabetes Care* **40**, 338–345 (2017).
295. Charlson, M. E., Pompei, P., Ales, K. L. & MacKenzie, C. R. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of chronic diseases* **40**, 373–383 (1987).
296. Wells, B. J., Chagin, K. M., Nowacki, A. S. & Kattan, M. W. Strategies for handling missing data in electronic health record derived data. *Egems* **1**, (2013).
297. Sterne, J. A. C. *et al.* Multiple imputation for missing data in epidemiological and clinical research: Potential and pitfalls. *BMJ* **338**, b2393 (2009).
298. Moons, K. G., Donders, R. A., Stijnen, T. & Harrell Jr, F. E. Using the outcome for imputation of missing predictor values was preferred. *Journal of clinical epidemiology* **59**, 1092–1101 (2006).

299. Lévesque, L. E., Hanley, J. A., Kezouh, A. & Suissa, S. Problem of immortal time bias in cohort studies: Example using statins for preventing progression of diabetes. *BMJ* **340**, b5087 (2010).
300. Lamarca, R., Alonso, J., Gomez, G. & Munoz, A. Left-truncated Data With Age as Time Scale: An Alternative for Survival Analysis in the Elderly Population. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* **53A**, M337–M343 (1998).
301. Gail, M. H., Graubard, B., Williamson, D. F. & Flegal, K. M. Comments on “Choice of time scale and its effect on significance of predictors in longitudinal studies” by Michael J. Pencina, Martin G. Larson and Ralph B. D’Agostino, *Statistics in Medicine* 2007; **26** :1343-1359. *Statistics in Medicine* **28**, 1315–1317 (2009).
302. Pencina, M. J., Larson, M. G. & D’Agostino, R. B. Choice of time scale and its effect on significance of predictors in longitudinal studies. *Statistics in Medicine* **26**, 1343–1359 (2007).
303. Pigott, T. D. A review of methods for missing data. *Educational research and evaluation* **7**, 353–383 (2001).
304. Hughes, R. A., Heron, J., Sterne, J. A. C. & Tilling, K. Accounting for missing data in statistical analyses: Multiple imputation is not always the answer. *International Journal of Epidemiology* **48**, 1294–1304 (2019).
305. Lipsitch, M., Tchetgen, E. T. & Cohen, T. Negative Controls: A Tool for Detecting Confounding and Bias in Observational Studies. *Epidemiology (Cambridge, Mass.)* **21**, 383–388 (2010).
306. Selva-O’Callaghan, A. *et al.* Statin-induced myalgia and myositis: An update on pathogenesis and clinical recommendations. *Expert review of clinical immunology* **14**, 215–224 (2018).
307. Herrett, E. *et al.* Statin treatment and muscle symptoms: Series of randomised, placebo controlled n-of-1 trials. *BMJ* **372**, n135 (2021).
308. Macedo, A. F., Douglas, I., Smeeth, L., Forbes, H. & Ebrahim, S. Statins and the risk of type 2 diabetes mellitus: Cohort study using the UK clinical practice pesearch datalink. *BMC Cardiovascular Disorders* **14**, 85 (2014).
309. Smit, R. A. J. *et al.* Statin-induced LDL cholesterol response and type 2 diabetes: A bidirectional two-sample Mendelian randomization study. *The Pharmacogenomics Journal* **20**, 462–470 (2020).
310. Karalis, D. G., Hill, A. N., Clifton, S. & Wild, R. A. The risks of statin use in pregnancy: A systematic review. *Journal of Clinical Lipidology* **10**, 1081–1090 (2016 Sep-Oct).
311. Carbonari, D. M. *et al.* Use of demographic and pharmacy data to identify patients included within both the Clinical Practice Research Datalink (CPRD) and The Health Improvement Network (THIN). *Pharmacoepidemiology and Drug Safety* **24**, 999–1003 (2015).

312. Salas, M., Hotman, A. & Stricker, B. H. Confounding by Indication: An Example of Variation in the Use of Epidemiologic Terminology. *American Journal of Epidemiology* **149**, 981–983 (1999).
313. Suttorp, M. M., Siegerink, B., Jager, K. J., Zoccali, C. & Dekker, F. W. Graphical presentation of confounding in directed acyclic graphs. *Nephrology Dialysis Transplantation* **30**, 1418–1423 (2015).
314. Boruzs, K., Juhász, A., Nagy, C., Ádány, R. & Bíró, K. Relationship between Statin Utilization and Socioeconomic Deprivation in Hungary. *Frontiers in Pharmacology* **7**, 66 (2016).
315. Ikeda, T. *et al.* Socioeconomic inequalities in low back pain among older people: The JAGES cross-sectional study. *International Journal for Equity in Health* **18**, 15 (2019).
316. Danaei, G., Rodríguez, L. A. G., Cantero, O. F., Logan, R. & Hernán, M. A. Observational data for comparative effectiveness research: An emulation of randomised trials of statins and primary prevention of coronary heart disease. *Statistical Methods in Medical Research* **22**, 70–96 (2013).
317. Taylor, F. *et al.* Statins for the primary prevention of cardiovascular disease. *The Cochrane Database of Systematic Reviews* **2013**, CD004816 (2013).
318. Newman, C. B. *et al.* Statin Safety and Associated Adverse Events: A Scientific Statement From the American Heart Association. *Arteriosclerosis, Thrombosis, and Vascular Biology* **39**, e38–e81 (2019).
319. Shepardson, N. E. Cholesterol Level and Statin Use in Alzheimer Disease: II. Review of Human Trials and Recommendations. *Archives of Neurology* **68**, 1385 (2011).
320. Porta, M. *A dictionary of epidemiology*. (Oxford University Press, 2014).
321. Bennet, A. M. *et al.* Association of apolipoprotein E genotypes with lipid levels and coronary risk. *JAMA* **298**, 1300–1311 (2007).
322. Chalmers, I. The Cochrane collaboration: Preparing, maintaining, and disseminating systematic reviews of the effects of health care. *Annals of the New York Academy of Sciences* **703**, 156-163; discussion 163-165 (1993).
323. Thompson, S. G. & Higgins, J. P. T. Treating individuals 4: Can meta-analysis help target interventions at individuals most likely to benefit? *Lancet (London, England)* **365**, 341–346 (2005).
324. Hingorani, A. D. *et al.* Prognosis research strategy (PROGRESS) 4: Stratified medicine research. *The BMJ* **346**, e5793 (2013).
325. Ensor, J., Burke, D. L., Snell, K. I. E., Hemming, K. & Riley, R. D. Simulation-based power calculations for planning a two-stage individual participant data meta-analysis. *BMC Medical Research Methodology* **18**, 41 (2018).
326. Wickham, H. & Grolemund, G. *Chapter 21: Iteration / R for data science: Import, tidy, transform, visualize, and model data*. (O'Reilly, 2016).

327. Levis, B., Hattle, M. & Riley, R. D. PRIME-IPD SERIES Part 2. Retrieving, checking, and harmonizing data are underappreciated challenges in individual participant data meta-analyses. *Journal of Clinical Epidemiology* **136**, 221–223 (2021).
328. Fibrinogen Studies Collaboration. Systematically missing confounders in individual participant data meta-analysis of observational cohort studies. *Statistics in Medicine* **28**, 1218–1237 (2009).
329. Abo-Zaid, G. *et al.* Individual participant data meta-analyses should not ignore clustering. *Journal of Clinical Epidemiology* **66**, 865–873.e4 (2013).
330. Burke, D. L., Ensor, J. & Riley, R. D. Meta-analysis using individual participant data: One-stage and two-stage approaches, and why they may differ. *Statistics in Medicine* **36**, 855–875 (2017).
331. Fisher, D. J., Carpenter, J. R., Morris, T. P., Freeman, S. C. & Tierney, J. F. Meta-analytical methods to identify who benefits most from treatments: Daft, deluded, or deft approach? *BMJ* **356**, j573 (2017).
332. Caerphilly Prospective Study - About.
333. Elwood, P. *et al.* Healthy Lifestyles Reduce the Incidence of Chronic Diseases and Dementia: Evidence from the Caerphilly Cohort Study. *PLOS ONE* **8**, e81877 (2013).
334. Riboli, E. The EPIC Project: Rationale and study design. European Prospective Investigation into Cancer and Nutrition. *International Journal of Epidemiology* **26**, 6S–14 (1997).
335. Riboli, E. *et al.* European Prospective Investigation into Cancer and Nutrition (EPIC): Study populations and data collection. *Public Health Nutrition* **5**, 1113–1124 (2002).
336. Marmot, M. & Brunner, E. Cohort Profile: The Whitehall II study. *International Journal of Epidemiology* **34**, 251–256 (2005).
337. Hospital Episode Statistics (HES). *NHS Digital*
338. Ahmed, I., Sutton, A. J. & Riley, R. D. Assessment of publication bias, selection bias, and unavailable data in meta-analyses using individual participant data: A database survey. *BMJ* **344**, d7762 (2012).
339. Jaspers, G. J. & Degraeuwe, P. L. A failed attempt to conduct an individual patient data meta-analysis. *Systematic Reviews* **3**, 97 (2014).
340. Danko, K. J., Dahabreh, I. J., Ivers, N. M., Moher, D. & Grimshaw, J. M. Contacting authors by telephone increased response proportions compared with emailing: Results of a randomized study. *Journal of Clinical Epidemiology* **115**, 150–159 (2019).
341. Safieh, M., Korczyn, A. D. & Michaelson, D. M. ApoE4: An emerging therapeutic target for Alzheimer's disease. *BMC Medicine* **17**, 64 (2019).
342. Credit where credit is due. *Nature* **462**, 825–825 (2009).

343. Hill, A. B. The Environment and Disease: Association or Causation? *Proceedings of the Royal Society of Medicine* **58**, 295–300 (1965).
344. DuBroff, R. Confirmation bias, conflicts of interest and cholesterol guidance: Can we trust expert opinions? *QJM: An International Journal of Medicine* **111**, 687–689 (2018).
345. Ference, B. A. *et al.* Clinical effect of naturally random allocation to lower systolic blood pressure beginning before the development of hypertension. *Hypertension (Dallas, Tex.: 1979)* **63**, 1182–1188 (2014).
346. Turner, R. M., Spiegelhalter, D. J., Smith, G. C. S. & Thompson, S. G. Bias modelling in evidence synthesis. *Journal of the Royal Statistical Society. Series A, (Statistics in Society)* **172**, 21–47 (2009).
347. Guyatt, G. H. *et al.* GRADE guidelines: 9. Rating up the quality of evidence - indirectness. *Journal of Clinical Epidemiology* **64**, 1311–1316 (2011).
348. Thompson, S. *et al.* A proposed method of bias adjustment for meta-analyses of published observational studies. *International Journal of Epidemiology* **40**, 765–777 (2011).
349. Higgins, J. P. T., Thompson, S. G., Deeks, J. J. & Altman, D. G. Measuring inconsistency in meta-analyses. *BMJ : British Medical Journal* **327**, 557–560 (2003).
350. Higgins, J. P. T. Commentary: Heterogeneity in meta-analysis should be expected and appropriately quantified. *International Journal of Epidemiology* **37**, 1158–1160 (2008).
351. Amer, M. A. *et al.* A meta-epidemiological study of bias in randomized clinical trials of open and laparoscopic surgery. *British Journal of Surgery* **108**, 477–483 (2021).
352. Page, M. J. *et al.* Empirical Evidence of Study Design Biases in Randomized Trials: Systematic Review of Meta-Epidemiological Studies. *PLOS ONE* **11**, e0159267 (2016).
353. Iwagami, M. *et al.* Blood cholesterol and risk of dementia in more than 1·8 million people over two decades: A retrospective cohort study. *The Lancet Healthy Longevity* **2**, e498–e506 (2021).
354. Savović, J. *et al.* Association Between Risk-of-Bias Assessments and Results of Randomized Trials in Cochrane Reviews: The ROBES Meta-Epidemiologic Study. *American Journal of Epidemiology* **187**, 1113–1122 (2018).
355. Hartling, L. *et al.* Applying the Risk of Bias Tool in a Systematic Review of Combination Long-Acting Beta-Agonists and Inhaled Corticosteroids for Persistent Asthma. *PLOS ONE* **6**, e17242 (2011).
356. Minozzi, S. *et al.* Risk of bias in nonrandomized studies of interventions showed low inter-rater reliability and challenges in its application. *Journal of Clinical Epidemiology* **112**, 28–35 (2019).

357. Minozzi, S., Cinquini, M., Gianola, S., Gonzalez-Lorenzo, M. & Banzi, R. The revised Cochrane risk of bias tool for randomized trials (RoB 2) showed low interrater reliability and challenges in its application. *Journal of Clinical Epidemiology* **126**, 37–44 (2020).
358. Wilks, D. C. *et al.* Objectively Measured Physical Activity and Fat Mass in Children: A Bias-Adjusted Meta-Analysis of Prospective Studies. *PLOS ONE* **6**, e17205 (2011).
359. Whiting, P. F. *et al.* QUADAS-2: A revised tool for the quality assessment of diagnostic accuracy studies. *Annals of Internal Medicine* **155**, 529–536 (2011).
360. Moons, K. G. M. *et al.* PROBAST: A Tool to Assess Risk of Bias and Applicability of Prediction Model Studies: Explanation and Elaboration. *Annals of Internal Medicine* **170**, W1–W33 (2019).
361. Gong, J., Harris, K., Peters, S. A. E. & Woodward, M. Sex differences in the association between major cardiovascular risk factors in midlife and dementia: A cohort study using data from the UK Biobank. *BMC Medicine* **19**, 110 (2021).
362. Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLOS Medicine* **12**, e1001779 (2015).
363. Pérez-Gaxiola, G., Verdugo-Paiva, F., Rada, G. & Flórez, I. D. Assessment of Duplicate Evidence in Systematic Reviews of Imaging Findings of Children With COVID-19. *JAMA Network Open* **4**, e2032690 (2021).
364. Group, H. P. S. C. Effects on 11-year mortality and morbidity of lowering LDL cholesterol with simvastatin for about 5 years in 20 536 high-risk individuals: A randomised controlled trial. *The Lancet* **378**, 2013–2020 (2011).
365. Ford, I., Murray, H., McCowan, C. & Packard, C. J. Long-Term Safety and Efficacy of Lowering Low-Density Lipoprotein Cholesterol With Statin Therapy. *Circulation* **133**, 1073–1080 (2016).
366. Llewellyn-Bennett, R. *et al.* Post-trial follow-up methodology in large randomised controlled trials: A systematic review. *Trials* **19**, 298 (2018).
367. Elliott, J. H. *et al.* Living Systematic Reviews: An Emerging Opportunity to Narrow the Evidence-Practice Gap. *PLOS Medicine* **11**, e1001603 (2014).
368. Sterne, J. A. Why the Cochrane Risk of Bias Tool Should not Include Funding Source as a Standard Item. *Cochrane Database of Systematic Reviews* (2013). doi:10.1002/14651858.ED000076
369. Russ, T. C., Kivimäki, M., Starr, J. M., Stamatakis, E. & Batty, G. D. Height in relation to dementia death: Individual participant meta-analysis of 18 UK prospective cohort studies. *The British Journal of Psychiatry* **205**, 348–354 (2014).
370. Wilson, G. *et al.* Best Practices for Scientific Computing. *PLOS Biology* **12**, e1001745 (2014).

371. Wilson, G. *et al.* Good enough practices in scientific computing. *PLOS Computational Biology* **13**, e1005510 (2017).
372. R Core Team. *R: A language and environment for statistical computing*. (R Foundation for Statistical Computing, 2019).
373. Xie, Y. *Bookdown: Authoring books and technical documents with r markdown*. (2021).
374. Textror, J., der Zander, B. van & Ankan, A. *Dagitty: Graphical analysis of structural causal models*. (2021).
375. Dowle, M. & Srinivasan, A. *Data.table: Extension of ‘data.frame’*. (2021).
376. Iannone, R. *DiagrammeR: Graph/Network visualization*. (2020).
377. Wickham, H., François, R., Henry, L. & Müller, K. *Dplyr: A grammar of data manipulation*. (2021).
378. Gohel, D. *Flextable: Functions for tabular reporting*. (2021).
379. Barrett, M. *Ggdag: Analyze and create elegant directed acyclic graphs*. (2021).
380. Wickham, H. *et al.* *Ggplot2: Create elegant data visualisations using the grammar of graphics*. (2021).
381. Zhu, H. *kableExtra: Construct complex table with kable and pipe syntax*. (2020).
382. Xie, Y. *Knitr: A general-purpose package for dynamic report generation in r*. (2021).
383. Bache, S. M. & Wickham, H. *Magrittr: A forward-pipe operator for r*. (2020).
384. Bates, D. & Maechler, M. *Matrix: Sparse and dense matrix classes and methods*. (2019).
385. McGuinness, L. & Schmidt, L. *Medrxivr: Access and search MedRxiv and BioRxiv preprint data*. (2021).
386. Allaire, J. J., Gandrud, C., Russell, K. & Yetman, C. *networkD3: D3 JavaScript network graphs from r*. (2017).
387. Pedersen, T. L. *Patchwork: The composer of plots*. (2020).
388. Urbanek, S. *Png: Read and write PNG images*. (2013).
389. Wickham, H. & Bryan, J. *Readxl: Read excel files*. (2019).
390. McGuinness, L. *Robvis: Visualize the results of risk-of-bias (ROB) assessments*. (2021).
391. Pebesma, E. *Sf: Simple features for r*. (2021).

392. Xie, Y. *Bookdown: Authoring books and technical documents with R markdown.* (Chapman and Hall/CRC, 2016).
393. Textor, J., der Zander, B. van, Gilthorpe, M. S., Liśkiewicz, M. & Ellison, G. T. Robust causal inference using directed acyclic graphs: The R package 'dagitty'. *International Journal of Epidemiology* **45**, 1887–1894 (2016).
394. Wickham, H. *Ggplot2: Elegant graphics for data analysis.* (Springer-Verlag New York, 2016).
395. Xie, Y. *Dynamic documents with R and knitr.* (Chapman and Hall/CRC, 2015).
396. Xie, Y. Knitr: A comprehensive tool for reproducible research in R. in *Implementing reproducible computational research* (eds. Stodden, V., Leisch, F. & Peng, R. D.) (Chapman and Hall/CRC, 2014).
397. McGuinness, L. A. & Schmidt, L. Medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R. *Journal of Open Source Software* **5**, 2651 (2020).
398. Viechtbauer, W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* **36**, 1–48 (2010).
399. McGuinness, L. A. & Higgins, J. P. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Research Synthesis Methods* (2020). doi:10.1002/jrsm.1411
400. Pebesma, E. Simple features for r: Standardized support for spatial vector data. *The R Journal* **10**, 439–446 (2018).

Appendices

A

Appendices by Chapter

Contents

A.1	Chapter 1	254
A.2	Chapter 2	256
A.3	Chapter 3	258
A.4	Chapter 4	277
A.5	Chapter 5	278
A.6	Chapter 6	281
A.7	Chapter 7	283
A.8	Chapter 8	288

A.1 Chapter 1

A.1.1 Publications beyond the scope of this thesis

In addition to publications arising directly from this thesis (described in Section 1.7.1 & included in Appendix B.3), I have been involved in a number of external research projects. Taken together, these additional projects provided extensive experience in evidence synthesis methods and in collaborating across large teams.

During 2020, I was involved in the updating of the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement. My contributions were focused on the data sharing aspects of the guidelines and on building an interactive web application that would allow users to complete the PRISMA checklist.

Publications arising from this work are listed below:

- *Page, M.J., Moher, D., Bossuyt, P., Boutron, I., Hoffmann, T., Mulrow, C., Shamseer, L., Tetzlaff, J., Akl, E., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L., Thomas, J., Tricco, A., Welch, V.A., Whiting, P., McKenzie, J. (2021). PRISMA 2020 statement: updated guidelines for reporting systematic reviews and meta-analyses. BMJ 372:n71. DOI: 10.1136/bmj.n71*
- *Page, M.J., Moher, D., Bossuyt, P., Boutron, I., Hoffmann, T., Mulrow, C., Shamseer, L., Tetzlaff, J., Akl, E., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L., Thomas, J., Tricco, A., Welch, V.A., Whiting, P., McKenzie, J. (2021). PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. BMJ 372:n160. DOI: 10.1136/bmj.n160*

- Haddaway, N. R., Page, M. J., Pritchard, C. C., & McGuinness, L. A. (2021). PRISMA2020: An R package and Shiny app for producing PRISMA 2020-compliant flow diagrams, with interactivity for optimised digital transparency and Open Synthesis. *MedRxiv*, DOI: 10.1101/2021.07.14.21260492

In response to the COVID-19 pandemic, I worked with a team of evidence synthesists to develop a literature searching/screening pipeline. This work underpinned a living systematic review on impact of the COVID-19 pandemic on self-harm and suicidal behaviour. Subsequently, the pipeline enabled rapid reviews of related topics, such as an analysis of the available evidence by population.

- John, A., Okolie, C., Eyles, E., Webb, R.T., Schmidt, L., **McGuinness, L.A.**, Olorisade, B.K., Arensman, E., Hawton, K., Kapur, N. and Moran, P., 2020. *The impact of the COVID-19 pandemic on self-harm and suicidal behaviour: a living systematic review*. *F1000Research*, 9(1097). DOI: 10.12688/f1000research.25522.2
- John, A., Eyles, E., **McGuinness, L.A.**, Okolie, C., Olorisade, B.K., Schmidt, L., Webb, R.T., Arensman, E., Hawton, K., Kapur, N. and Moran, P., 2020. *The impact of the COVID-19 pandemic on self-harm and suicidal behaviour: protocol for a living systematic review*. *F1000Research*, 9(644), p.644. DOI: 10.12688/f1000research.24274.1
- Eyles, E., Moran, P., Okolie, C., Dekel, D., Macleod-Hall, C., Webb, R. T., Schmidt, L., Knipe, D., Sinyor, M., **McGuinness, L. A.**, Arensman, E., Hawton, K., O'Connor, R. C., Kapur, N., O'Neill, S., Olorisade, B., Cheng, H.-Y., Higgins, J. P. T., John, A., & Gunnell, D. (2021). Systematic review of the impact of the COVID-19 pandemic on suicidal behaviour amongst health and social care workers across the world. *Journal of Affective Disorders Reports*, 6, 100271. DOI: 10.1016/j.jadr.2021.100271

Finally, I have acted as part of a large review team for a number of other evidence synthesis exercises:

- *Grant, M. C., Geoghegan, L., Arbyn, M., Mohammed, Z., McGuinness, L.A., Clarke, E. L., & Wade, R. (2020). The Prevalence of Symptoms in 24,410 Adults Infected by the Novel Coronavirus (SARS-CoV-2; COVID-19): A Systematic Review and Meta-Analysis of 148 Studies from 9 Countries. PLOS ONE. DOI: 10.1371/journal.pone.0234765*
- *Cheng, H.Y., McGuinness, L.A., Elbers, R.G., MacArthur, G.J., Taylor, A., McAleenan, A., Dawson, S., López-López, J.A., Higgins, J.P., Cowlishaw, S. and Lingford-Hughes, A. (2020). Treatment interventions to maintain abstinence from alcohol in primary care: systematic review and network meta-analysis. BMJ, 371. DOI: 10.1136/bmj.m3934*
- *Schmidt, L., Olorisade, B.K., McGuinness, L.A., Thomas, J. and Higgins, J.P. (2020). Data extraction methods for systematic review (semi) automation. F1000Research, 10:401. DOI: 10.12688/f1000research.51117.1*

A.2 Chapter 2

A.2.1 Code for publication rate analysis

medRxiv automatically links preprints to their published counterparts, and provides the DOI of the published article as part of the metadata on a given preprint. Using `medrxivr` to access this data, I could assess the two-year publication rate of preprints originally posted to medRxiv in July 2019. The code used for this analysis is presented below:

```

# Load relevant packages

library(medrxivr)
library(dplyr)

# Use snapshot of published status from July 2021

mx_search(
  mx_snapshot("ccedfb8a44304b9fba4e3ba518a8ce4ed2294770"),
  query = "*",
  # Limit to those 129 records published in July 2019
  from_date = "2019-07-01",
  to_date = "2019-08-01"
) %>%

# Create indicator to show which records have been published

mutate(Status = ifelse(published == "NA",
                      "Not published",
                      "Published")) %>%

# Group by indicator variable ("Published"/"Not published")

group_by(Status) %>%

# Get total number per group, and percentage of total (N=129)

summarise(N = n(),
          Percent = comma(n() / 129 * 100))

```

As shown in Table A.1, two-thirds of preprints were published within two years of their submission to medRxiv.

Table A.1: Publication status of medRxiv preprints deposited in July 2019, allowing for a two-year lag

Status	N	Percent
Not published	42	33%
Published	87	67%

A.3 Chapter 3

A.3.1 Deviations from protocol

There were a number of deviations from the pre-registered protocol for the systematic review reported in Chapters 3 & 4:¹⁴¹

- I had originally intended to examine both dementia and mild cognitive impairment (MCI) as outcomes of interest. However, given the large number of relevant records for the dementia outcomes, I did not include MCI as an outcome.
- For Mendelian randomisation analyses, I had originally intended to use a panel of experts to assess the risk of bias in this study design. However, through private correspondence with the authors of a review of risk-of-bias assessment tools for Mendelian randomisation studies (since published¹⁶⁷), I became aware of the Mamluk *et al* tool. Given this tool was considered to be the best available for assessing Mendelian randomisation studies and that it followed a domain-based approach, enabling comparison with tools for other study designs, I decided to use it in the review.
- I performed an additional assessment of the risk of bias due to missing evidence using the ROB-ME tool. This analysis was not planned because the tool did not exist when the protocol was written.

- I had originally intended to assess sources of funding alongside bias. However, on further reading,³⁶⁸ I learned that any bias introduced by sources of funding would operate through one of the existing domains of the risk-of-bias tools. As such, I did not consider funding sources separately.
- Finally, I performed a dose response meta-analysis to assess the evidence for a non-linear relationship between lipids and dementia outcomes. This marked a departure from the published protocol, as I had not expected the number of studies reporting dementia risk across lipid sub-categories to be sufficient for a dose response analysis.

A.3.2 Web of Science databases searched

Table A.2: Summary of Web of Science databases searched. The contents of the Web of Science core collection can vary depending on an institution's subscription, and so detailing the exact databases and ranges searched is important.

Database	Abbreviation	Years
Science Citation Index Expanded	SCI-EXPANDED	1900-present
Social Sciences Citation Index	SSCI	1956-present
Arts & Humanities Citation Index	A&HCI	1975-present
Conference Proceedings Citation Index - Science	CPCI-S	1990-present
Conference Proceedings Citation Index - Social Science & Humanities	CPCI-SSH	1990-present
Emerging Sources Citation Index	ESCI	2015-present

A.3.3 Search strategy

An overview of the number of hits per database, in addition to the full line-by-line search results for each database, are presented below.

Table A.3: Summary of number of records identified in each database searched

Step	Hits
Medline	6045
EMBASE	10255
PyscINFO	800
CENTRAL	1473
Web of Science	4874
Total (pre-deduplication)	23447
Duplicates	-7338
Total (post-deduplication)	16109

Table A.4: Overview of the full Medline search strategy.

#	Search term	Hits
1	dement*.ti,ab.	103404
2	alzheimer*.ti,ab.	132832
3	exp Dementia/	154234
4	exp Alzheimer Disease/	87346
5	Pick* disease.ti,ab	2794
6	globular glial tauopathy.ti,ab	24
7	primary progressive aphasia.ti,ab	1051
8	logopaenic aphasia.ti,ab	0
9	posterior cortical atrophy.ti,ab	381
10	(age-associated) adj2 (memory decline).ti,ab	11
11	((mild or slight) adj2 (cognitive or cognition) adj2 (disorder* or defect* or deficit* or disabilit* or dysfunction or impair*)).ti,ab	14883
12	((cognit\$ or memory or cerebr\$ or mental\$) adj3 (declin\$ or impair\$ or los\$ or deteriorat\$ or degenerat\$ or complain\$ or disturb\$ or disorder\$)).ti,ab	182141
13	(MCI or aMCI or CIND or non-aMCI).ti,ab	16893
14	(cognitive impair*).ti,ab	56411
15	Cognition Disorders/	62602
16	Cognitive Dysfunction/	11999
17	Mild Cognitive Impairment/	11999
18	or/1-17	407352
19	lipid*.ti,ab.	462968
20	lipoprotein*.ti,ab.	140438
21	cholesterol.ti,ab.	227679
22	hypercholesterol*.ti,ab.	33093
23	hypcholesterol*.ti,ab.	3347
24	triacylglycerol.ti,ab.	11077
25	lipemia*.ti,ab.	1836
26	dyslipid?emia.ti,ab.	29128
27	hyperlipid?emia*.ti,ab.	25134
28	hypolipid?emia.ti,ab.	271
29	HDL.ti,ab.	61231
30	LDL.ti,ab.	71176
31	VLDL.ti,ab.	12485
32	triglyceride*.ti,ab.	104904
33	exp Dyslipidemias/	76480

Table A.4: Overview of the full Medline search strategy. (*continued*)

#	Search term	Hits
34	exp Cholesterol/	155339
35	exp Lipoproteins/	141558
36	or/19-35	777210
37	statin*.ti,ab.	39998
38	atorvastatin.ti,ab.	7994
39	cerivastatin.ti,ab	646
40	fluvastatin.ti,ab.	1795
41	pravastatin.ti,ab.	3940
42	rosuvastatin.ti,ab.	3175
43	simvastatin.ti,ab.	8933
44	pitavastatin.ti,ab	816
45	lovastatin.ti,ab.	3667
46	fibrat*.ti,ab.	3135
47	("fribic acid" adj3 derivat*).ti,ab.	341
48	bezafibrate.ti,ab	1523
49	fenofibrate.ti,ab	3109
50	gemfibrozil.ti,ab	1802
51	clofenapate.ti,ab	39
52	clofibrate.ti,ab	3035
53	ciprofibrate.ti,ab	481
54	(bile adj3 sequest*).ti,ab.	816
55	colestyramine.ti,ab	60
56	colestipol hydrochloride.ti,ab	52
57	colesevelam hydrochloride.ti,ab	71
58	nicotinic acid*.ti,ab.	5854
59	inositol nicotinate.ti,ab	30
60	niacin.ti,ab	4631
61	ezetimibe.ti,ab.	2766
62	acipimox.ti,ab	292
63	evolocumab.ti,ab	394
64	alirocumab.ti,ab	350
65	lomitapide.ti,ab	150
66	(omega-3-acid adj2 ethyl ester*).ti,ab	85
67	meglutol.ti,ab	2
68	Meglutol/	134
69	exp Anticholesteremic Agents/	71609
70	exp Fibric Acids/	9523
71	exp Ezetimibe/	1954
72	exp Nicotinic Acids/	36409
73	or/37-72	138108
74	18 and 36	19659
75	18 and 73	2287
76	74 or 75	21029
77	Animals/ not (Animals/ and Humans/)	4552498
78	76 not 77	18226
79	epidemiologic studies/ or case-control studies/ or cross-sectional studies/ or cohort studies/ or follow-up studies/ or longitudinal studies/ or prospective studies/ or retrospective studies/	2299133
80	((epidemiologic or prospective or retrospective or cross-sectional or case control* or cohort or longitudinal or followup or follow-up) adj3 (study or studies)).ti,ab,kf.	1043484
81	(case control* or cross-sectional or cohort? or follow-up or followup or longitudinal or prospective or retrospective or observational or population).ti.	656500
82	(cohort? adj2 (analys* or compar* or data or study or studies)).ab.	184866
83	(population adj2 (based or data* or study or studies or register? or survey? or surveillance)).ab.	200506
84	or/79-83	2933516

Table A.4: Overview of the full Medline search strategy. (*continued*)

#	Search term	Hits
85	controlled clinical trial.pt.	93095
86	randomized controlled trial.pt.	483099
87	clinical trials as topic/	187183
88	(randomi#ed or randomi#ation or randomi#ing).ti,ab,kf.	585795
89	(RCT or "at random" or (random* adj3 (administ* or allocat* or assign* or class* or cluster or crossover or cross-over or control* or determine* or divide* or division or distribut* or expose* or fashion or number* or place* or pragmatic or quasi or recruit* or split or subsitut* or treat*)).ti,ab,kf.	512675
90	placebo.ab,ti,kf.	203773
91	trial.ti.	199586
92	(control* adj3 group*).ab.	498141
93	(control* and (trial or study or group*) and (waitlist* or wait* list* or ((treatment or care) adj2 usual))).ti,ab,kf.	19035
94	((single or double or triple or treble) adj2 (blind* or mask* or dummy)).ti,ab,kf.	165010
95	double-blind method/ or random allocation/ or single-blind method/	266392
96	or/85-95	1616814
97	84 or 96	4175140
98	MENDELIAN RANDOMIZATION ANALYSIS/	736
99	Mendelian randomi*.ti,ab,kf.	1647
100	98 or 99	1738
101	RANDOMIZED CONTROLLED TRIALS AS TOPIC/	124147
102	(RCT? or (randomi#ed adj2 (control* or intervention* or experiment* or trial* or study or studies))).ti,ab,kf.	405207
103	((random* or comparative or intervention? or treatment?) adj3 (efficacy or effect*).ti,ab,kf.	435773
104	(clinical adj (intervention? or trial?)).ti,ab,kf.	346211
105	CLINICAL TRIALS AS TOPIC/ or CONTROLLED CLINICAL TRIALS AS TOPIC/	192430
106	TREATMENT EFFECT/	904484
107	or/101-106	1894420
108	100 AND 107	313
109	instrument* variab*.ti,ab,kf.	2380
110	((causal* or causative) adj3 (associat* or infer* or implicat* or effect* or predict* or factor? or risk? or relat*).ti,ab,kf.	54710
111	((gene* adj2 (associat* or risk? or varia* or determinant?)) or risk variant?).ti,ab,kf.	234808
112	(disease* adj3 (expos* or associat* or etiolog* or pathogenesis or risk?)).ti,ab,kf.	304605
113	risk factor?.mp.	1045594
114	exp CAUSALITY/	782487
115	"confounding factors (epidemiology)"/	9873
116	(confound* or nonconfound* or non-confound*).ti,ab,kf.	113902
117	(statistics or epidemiolog* or ((genetic* or molecular) and medicine)).jw.	205082
118	or/109 -117	1768577
119	108 and 118	273
120	98 and 101	27
121	119 or 120	277
122	97 or 121	4175143
123	78 and 122	6045

Table A.5: Overview of the full Embase search strategy.

#	Search term	Hits
1	dement*.ti,ab.	150297
2	alzheimer*.ti,ab.	182305
3	exp Dementia/	332818
4	exp Alzheimer Disease/	184019
5	Pick* disease.ti,ab	3399
6	globular glial tauopathy.ti,ab	52
7	primary progressive aphasia.ti,ab	1877
8	logopaenic aphasia.ti,ab	0
9	posterior cortical atrophy.ti,ab	714
10	(age-associated) adj2 (memory decline).ti,ab	17
11	((mild or slight) adj2 (cognitive or cognition) adj2 (disorder* or defect* or deficit* or disabilit* or dysfunction or impair*)).ti,ab	24705
12	((cognit\$ or memory or cerebr\$ or mental\$) adj3 (declin\$ or impair\$ or los\$ or deteriorat\$ or degenerat\$ or complain\$ or disturb\$ or disorder\$)).ti,ab.	259650
13	(MCI or aMCI or CIND or non-aMCI).ti,ab	31158
14	(cognitive impair*).ti,ab	88915
15	Cognition Disorders/	35208
16	Cognitive Dysfunction/	80947
17	Mild Cognitive Impairment/	23295
18	or/1-17	613530
19	lipid*.ti,ab.	568453
20	lipoprotein*.ti,ab.	168172
21	cholesterol.ti,ab.	295493
22	hypercholesterol*.ti,ab.	45908
23	hypcholesterol*.ti,ab.	3985
24	triacylglycerol.ti,ab.	12534
25	lipemia*.ti,ab.	2013
26	dyslipid?emia.ti,ab.	51477
27	hyperlipid?emia*.ti,ab.	40531
28	hypolipid?emia.ti,ab.	318
29	HDL.ti,ab.	95530
30	LDL.ti,ab.	108193
31	VLDL.ti,ab.	16654
32	triglyceride*.ti,ab.	147716
33	exp Dyslipidemias/	64190
34	exp Cholesterol/	286092
35	exp Lipoproteins/	242167
36	or/19-35	1045396
37	statin*.ti,ab.	65212
38	atorvastatin.ti,ab.	13401
39	cerivastatin.ti,ab	848
40	fluvastatin.ti,ab.	2551
41	pravastatin.ti,ab.	5573
42	rosuvastatin.ti,ab.	5706
43	simvastatin.ti,ab.	13858
44	pitavastatin.ti,ab	1417
45	lovastatin.ti,ab.	4742
46	fibrat*.ti,ab.	4434
47	("fibric acid" adj3 derivat*).ti,ab.	407
48	bezafibrate.ti,ab	1996
49	fenofibrate.ti,ab	4540
50	gemfibrozil.ti,ab	2309
51	clofibrate.ti,ab	32
52	clofibrate.ti,ab	3346
53	ciprofibrate.ti,ab	545
54	(bile adj3 sequest*).ti,ab.	1092

Table A.5: Overview of the full Embase search strategy. (*continued*)

#	Search term	Hits
55	colestyramine.ti,ab	133
56	colestipol hydrochloride.ti,ab	59
57	colesevelam hydrochloride.ti,ab	80
58	nicotinic acid*.ti,ab.	5159
59	inositol nicotinate.ti,ab	46
60	niacin.ti,ab	5622
61	ezetimibe.ti,ab.	4636
62	acipimox.ti,ab	394
63	evolocumab.ti,ab	707
64	alirocumab.ti,ab	648
65	lomitapide.ti,ab	239
66	(omega-3-acid adj2 ethyl ester*).ti,ab	141
67	meglutol.ti,ab	3
68	Meglutol/	219
69	exp Anticholesteremic Agents/	178271
70	exp Fibric Acids/	33072
71	exp Ezetimibe/	9146
72	exp Nicotinic Acids/	1946
73	or/37-72	204136
74	18 and 36	36463
75	18 and 73	6013
76	74 or 75	40002
77	Animals/ not (Animals/ and Humans/)	952788
78	76 not 77	39778
79	Clinical study/ OR Case control study/ OR Longitudinal study/ OR Retrospective study/ OR Prospective study/ OR Cohort analysis/ OR cross-sectional studies/	2034044
80	((epidemiologic or prospective or retrospective or cross-sectional or case control* or cohort or longitudinal or followup or follow-up) adj3 (study or studies)).ti,ab,kw.	1493506
81	(case control* or cross-sectional or cohort? or follow-up or followup or longitudinal or prospective or retrospective or observational or population).ti.	860479
82	(cohort? adj2 (analys* or compar* or data or study or studies)).ab.	280187
83	(population adj2 (based or data* or study or studies or register? or survey? or surveillance)).ab.	286930
84	or/79-83	3106288
85	randomized controlled trial/	551164
86	randomization.de.	82502
87	controlled clinical trial/	462715
88	*clinical trial/	17542
89	placebo.de.	334642
90	placebo.ti,ab.	287949
91	trial.ti.	270940
92	(randomi#ed or randomi#ation or randomi#ing).ti,ab,kw.	836677
93	(RCT or "at random" or (random* adj3 (administ* or allocat* or assign* or class* or control* or determine* or divide* or division or distribut* or expose* or fashion or number* or place* or recruit* or split or subsitut* or treat*)).ti,ab,kw.	667211
94	((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (blind\$ or mask\$ or dummy)).mp.	290312
95	(control* and (trial or study or group*)) and (waitlist* or wait* list* or ((treatment or care) adj2 usual))).ti,ab,kw.	25934
96	or/85-95	1682845
97	84 or 96	4464235
98	MENDELIAN RANDOMIZATION ANALYSIS/	1856
99	Mendelian randomi*.ti,ab.	2066
100	98 or 99	2447
101	RANDOMIZED CONTROLLED TRIALS AS TOPIC/	95688

Table A.5: Overview of the full Embase search strategy. (*continued*)

#	Search term	Hits
102	(RCT? or (randomi#ed adj2 (control* or intervention* or experiment* or trial* or study or studies))).ti,ab.	557079
103	((random* or comparative or intervention? or treatment?) adj3 (efficacy or effect*)).ti,ab.	611431
104	(clinical adj (intervention? or trial?)).ti,ab.	479774
105	CLINICAL TRIALS AS TOPIC/ or CONTROLLED CLINICAL TRIALS AS TOPIC/	79666
106	TREATMENT EFFECT/	1167457
107	or/101-106	2561978
108	100 AND 107	499
109	instrument* variab*.ti,ab.	2809
110	((causal* or causative) adj3 (associat* or infer* or implicat* or effect* or predict* or factor? or risk? or relat*)).ti,ab.	69566
111	((gene* adj2 (associat* or risk? or varia* or determinant?)) or risk variant?).ti,ab.	302507
112	(disease* adj3 (expos* or associat* or etiolog* or pathogenesis or risk?)).ti,ab.	420445
113	risk factor?.mp.	1243497
114	exp CAUSALITY/	2038
115	"confounding factors (epidemiology)" /	203764
116	(confound* or nonconfound* or non-confound*).ti,ab.	157400
117	(statistics or epidemiolog* or ((genetic* or molecular) and medicine)).jx.	210731
118	or/108-117	2325782
119	108 and 118	499
120	98 and 101	147
121	119 or 120	499
122	97 or 121	4464264
123	78 and 122	10255

Table A.6: Overview of the full PsycINFO search strategy.

#	Search term	Hits
1	dement*.ti,ab.	62001
2	alzheimer*.ti,ab.	56133
3	exp Dementia/	72554
4	exp Alzheimer Disease/	44097
5	Pick* disease.ti,ab	595
6	globular glial tauopathy.ti,ab	13
7	primary progressive aphasia.ti,ab	791
8	logopaenic aphasia.ti,ab	0
9	posterior cortical atrophy.ti,ab	270
10	(age-associated) adj2 (memory decline).ti,ab	8
11	((mild or slight) adj2 (cognitive or cognition) adj2 (disorder* or defect* or deficit* or disabilit* or dysfunction or impair*)).ti,ab.	9577
12	((cognit\$ or memory or cerebr\$ or mental\$) adj3 (declin\$ or impair\$ or los\$ or deteriorat\$ or degenerat\$ or complain\$ or disturb\$ or disorder\$)).ti,ab.	139004
13	(MCI or aMCI or CIND or non-aMCI).ti,ab	7092
14	(cognitive impair*).ti,ab	33109
15	Cognitive Dysfunction/	34042
16	or/1-15	215471
17	lipid*.ti,ab.	9083
18	lipoprotein*.ti,ab.	3102
19	cholesterol.ti,ab.	7650
20	hypercholesterol*.ti,ab.	826

Table A.6: Overview of the full PyscINFO search strategy. (*continued*)

#	Search term	Hits
21	hypcholesterol*.ti,ab.	30
22	triacylglycerol.ti,ab.	70
23	lipemia*.ti,ab.	9
24	dyslipid?emia.ti,ab.	1224
25	hyperlipid?emia*.ti,ab.	1027
26	hypolipid?emia.ti,ab.	2
27	HDL.ti,ab.	1628
28	LDL.ti,ab.	1364
29	VLDL.ti,ab.	100
30	triglyceride*.ti,ab.	2874
31	exp Cholesterol/	2081
32	exp Lipoproteins/	971
33	or/17-32	18467
34	statin*.ti,ab.	5057
35	atorvastatin.ti,ab.	222
36	cerivastatin.ti,ab	11
37	fluvastatin.ti,ab.	17
38	pravastatin.ti,ab.	90
39	rosuvastatin.ti,ab.	55
40	simvastatin.ti,ab.	280
41	lovastatin.ti,ab.	85
42	pitavastatin.ti,ab	20
43	fibrat*.ti,ab.	45
44	("fibric acid" adj3 derivat*).ti,ab.	1
45	bezafibrate.ti,ab	22
46	fenofibrate.ti,ab	40
47	gemfibrozil.ti,ab	22
48	clofenapate.ti,ab	0
49	clofibrate.ti,ab	12
50	ciprofibrate.ti,ab	0
51	(bile adj3 sequest*).ti,ab.	2
52	colestyramine.ti,ab	0
53	colestipol hydrochloride.ti,ab	0
54	colesevelam hydrochloride.ti,ab	0
55	nicotinic acid*.ti,ab.	154
56	inositol nicotinate.ti,ab	1
57	niacin.ti,ab	173
58	ezetimibe.ti,ab.	19
59	acipimox.ti,ab	1
60	evolocumab.ti,ab	3
61	alirocumab.ti,ab	1
62	lomitapide.ti,ab	1
63	(omega-3-acid adj2 ethyl ester*).ti,ab	1
64	meglutol.ti,ab	0
65	or/34-64	5782
66	16 and 33	3318
67	16 and 65	585
68	66 or 67	3682
69	Animals/ not (Animals/ and Humans/)	7175
70	68 not 69	3680
71	Clinical study/ OR Case control study/ OR Longitudinal study/ OR Retrospective study/ OR Prospective study/ OR Cohort analysis/ OR cross-sectional studies/	1317
72	((epidemiologic or prospective or retrospective or cross-sectional or case control* or cohort or longitudinal or followup or follow-up) adj3 (study or studies)).ti,ab,hw.	173681

Table A.6: Overview of the full PyscINFO search strategy. (*continued*)

#	Search term	Hits
73	(case control* or cross-sectional or cohort? or follow-up or followup or longitudinal or prospective or retrospective or observational or population).ti.	104744
74	(cohort? adj2 (analys* or compar* or data or study or studies)).ti,ab.	23369
75	(population adj2 (based or data* or study or studies or register? or survey? or surveillance)).ab.	48659
76	or/71-75	261952
77	randomization.de.	0
78	*clinical trial/	7000
79	placebo.de.	5253
80	placebo.ti,ab.	38285
81	trial.ti.	27967
82	(randomi#ed or randomi#ation or randomi#ing).ti,ab,hw.	78840
83	(RCT or "at random" or (random* adj3 (administ* or allocat* or assign* or class* or control* or determine* or divide* or division or distribut* or expose* or fashion or number* or place* or recruit* or split or substit* or treat*)).ti,ab,hw.	93430
84	((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (blind\$ or mask\$ or dummy)).mp.	25258
85	(control* and (trial or study or group*)) and (waitlist* or wait* list* or ((treatment or care) adj2 usual))).ti,ab,hw.	9382
86	or/77-85	159116
87	76 or 86	408627
88	Mendelian randomi*.ti,ab.	106
89	(RCT? or (randomi#ed adj2 (control* or intervention* or experiment* or trial* or study or studies))).ti,ab.	57324
90	((random* or comparative or intervention? or treatment?) adj3 (efficacy or effect*).ti,ab.	104711
91	(clinical adj (intervention? or trial?)).ti,ab.	34390
92	TREATMENT EFFECT/	22971
93	or/89-92	181514
94	88 AND 93	12
95	instrument* variab*.ti,ab.	1088
96	((causal* or causative) adj3 (associat* or infer* or implicat* or effect* or predict* or factor? or risk? or relat*).ti,ab.	20424
97	((gene* adj2 (associat* or risk? or varia* or determinant?)) or risk variant?).ti,ab.	28699
98	(disease* adj3 (expos* or associat* or etiolog* or pathogenesis or risk?)).ti,ab.	23709
99	risk factor?.mp.	118075
100	exp CAUSALITY/	4201
101	"confounding factors (epidemiology)"/	0
102	(confound* or nonconfound* or non-confound*).ti,ab.	28253
103	(statistics or epidemiolog* or ((genetic* or molecular) and medicine)).jw.	9677
104	or/95-103	211022
105	94 and 104	11
106	87 or 105	408627
107	70 and 106	800

Table A.7: Overview of the full CENTRAL search strategy.

#	Search term	Hits
1	dement*	21445
2	alzheimer*	11448
3	[mh Dementia]	5224
4	[mh "Alzheimer Disease"]	3015
5	Pick* disease	1059

Table A.7: Overview of the full CENTRAL search strategy. (*continued*)

#	Search term	Hits
6	globular glial tauopathy	0
7	primary progressive aphasia	115
8	logopaenic aphasia	0
9	posterior cortical atrophy	68
10	(age-associated) NEAR2 (memory decline)	2696
11	((mild or slight) NEAR2 (cognitive or cognition) NEAR2 (disorder* or defect* or deficit* or disabilit* or dysfunction or impair*))	136
12	((cognit\$ or memory or cerebr\$ or mental\$) NEAR3 (declin\$ or impair\$ or los\$ or deteriorat\$ or degenerat\$ or complain\$ or disturb\$ or disorder\$))	45
13	(MCI or aMCI or CIND or non-aMCI)	2288
14	cognitive impair*	14935
15	[mh "Mild Cognitive Impairment"]	922
16	[mh "cognition disorders"]	4475
17	[mh "cognitive dysfunction"]	922
18	#1 OR #2 OR #3 OR #4 OR #5 OR #6 or #7 or #8 or #9 or #10 or #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17	38448
19	lipid*	41298
20	lipoprotein*	22336
21	cholesterol	34213
22	hypercholesterol*	7724
23	hypcholesterol*	445
24	triacylglycerol	8029
25	lipemia*	375
26	dyslipid?emia	4929
27	hyperlipid?emia*	4667
28	hypolipid?emia	10
29	HDL	15420
30	LDL	18892
31	VLDL	1944
32	triglyceride*	20436
33	[mh Dyslipidemias]	6753
34	[mh Cholesterol]	9862
35	[mh Lipoproteins]	9416
36	#19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35	69284
37	statin*	9615
38	atorvastatin	4999
39	cerivastatin	196
40	fluvastatin	725
41	pravastatin	1954
42	rosuvastatin	2260
43	simvastatin	3698
44	lovastatin	950
45	pitavastatin	470
46	fibrat*	503
47	("fibric acid" NEAR/3 derivat*)	161
48	bezafibrate	458
49	fenofibrate	993
50	gemfibrozil	552
51	clofенапате	5
52	clofibrate	370
53	ciprofibrate	48
54	(bile NEAR/3 sequest*)	172
55	colestyramine	88
56	colestipol hydrochloride	29
57	colesevelam hydrochloride	104

Table A.7: Overview of the full CENTRAL search strategy. (*continued*)

#	Search term	Hits
58	nicotinic acid*	1183
59	inositol nicotinate	22
60	niacin	1084
61	ezetimibe	1541
62	acipimox	176
63	evolocumab	227
64	alirocumab	254
65	lomitapide	23
66	("omega-3-acid" NEAR/2 "ethyl ester")	95
67	meglutol	21
68	[mh meglutol]	2
69	[mh "Anticholesteremic Agents"]	5124
70	[mh "Fibric Acids"]	1251
71	[mh Ezetimibe]	635
72	[mh "Nicotinic Acids"]	1951
73	#37 OR #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56 OR #57 OR #58 OR #59 OR #60 OR #61 OR #62 OR #63 OR #64 OR #65 OR #66 OR #67 OR #68 OR #69 OR #70 OR #71 OR #72	21694
74	#18 and #36	1605
75	#18 and #73	727
76	#74 and #75	2025
	Limiting to trials rather than reviews/protocols	1473

Table A.8: Overview of the full Web of Science search strategy.

#	Search term	Hits
1	TS=dement*	154301
2	TS=alzheimer*	230302
3	TS=Pick* disease	8119
4	TS=globular glial tauopathy	39
5	TS=primary progressive aphasia	1992
6	TS=logopaenic aphasia	4
7	TS=posterior cortical atrophy	1271
8	TS=(age-associated NEAR/2 memory decline)	16
9	TS = ((mild or slight) NEAR/2 (cognitive or cognition) NEAR/2 (disorder* or defect* or deficit* or disabilit* or dysfunction or impair*))	
10	TS = ((cognit\$ or memory or cerebr\$ or mental\$) NEAR/3 (declin\$ or impair\$ or los\$ or deteriorat\$ or degenerat\$ or complain\$ or disturb\$ or disorder\$))	
11	TS = (MCI or aMCI or CIND or non-aMCI)	
12	TS = "cognitive impair"	
13	Y	313381
14	TS=lipid*	623475
15	TS=lipoprotein*	188675
16	TS=cholesterol	268455
17	TS=hypercholesterol*	46269
18	TS=hypocholesterol*	4146
19	TS=triacylglycerol	15348
20	TS=lipemia*	2882
21	TS=dyslipid?emia	4881
22	TS=hyperlipid?emia*	3324
23	TS=hypolipid?emia	45

Table A.8: Overview of the full Web of Science search strategy. (*continued*)

#	Search term	Hits
24	TS=HDL	63203
25	TS=LDL	76399
26	TS=VLDL	10962
27	TS=triglyceride*	108266
28	#14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27	920847
29	TS=statin*	54504
30	TS=atorvastatin	14422
31	TS=fluvastatin	2547
32	TS=pravastatin	8327
33	TS=rosuvastatin	5171
34	TS=simvastatin	15753
35	TS=lovastatin	6299
36	TS=fibrat*	6972
37	TS="fibrac acid" NEAR/3 derivat*)	313
38	TS=bezafibrate	1926
39	TS=fenofibrate	4194
40	TS=gemfibrozil	2606
41	TS=clofenapate	31
42	TS=clofibrate	3037
43	TS=ciprofibrate	608
44	TS=(bile NEAR/3 sequest*)	790
45	TS=colestyramine	64
46	TS=colestipol hydrochloride	73
47	TS=colesevelam hydrochloride	190
48	TS=nicotinic acid*	11925
49	TS=inositol nicotinate	49
50	TS=niacin	6199
51	TS=ezetimibe	3875
52	TS=acipimox	372
53	TS=evolocumab	723
54	TS=alirocumab	551
55	TS=lomitapide	179
56	TS=(omega-3-acid NEAR/2 "ethyl ester")	112
57	#29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56	108179
58	#13 and #28	18165
59	#13 and #57	2038
60	#58 or #59	19123
61	TS=((epidemiologic or prospective or retrospective or cross-sectional or "case control*" or cohort or longitudinal or followup or follow-up) NEAR/3 (study or studies))	1065411
62	TS=(case control* or cross-sectional or cohort? or follow-up or followup or longitudinal or prospective or retrospective or observational or population)	4896813
63	TS=(cohort* NEAR/2 (analys* or compar* or data or study or studies))	235118
64	TS=(population NEAR/2 (based or data* or study or studies or register? or survey? or surveillance))	326505
65	#61 OR #62 OR #63 OR #64	4948637
66	TS=controlled clinical trial	267207
67	TS=randomized controlled trial	412461
68	TS=(randomi?ed or randomi?ation or randomi?ing)	843224

Table A.8: Overview of the full Web of Science search strategy. (*continued*)

#	Search term	Hits
69	TS=(RCT or "at random" or (random* NEAR/3 (administ* or allocat* or assign* or class* or cluster or crossover or cross-over or control* or determine* or divide* or division or distribut* or expose* or fashion or number* or place* or pragmatic or quasi or recruit* or split or subsitut* or treat*)))	706888
70	TS=placebo	240446
71	TS=trial	1533550
72	TS=(control* NEAR/3 group*)	462747
73	TS=(control* and (trial or study or group*)) and (waitlist* or wait* list* or ((treatment or care) NEAR/2 usual)))	20886
74	TS=((single or double or triple or treble) NEAR/2 (blind* or mask* or dummy))	268471
75	#66 OR #67 OR #68 OR #69 OR #70 OR #71 OR #72 OR #73 OR #74	2393217
76	#65 or #75	6616705
77	TS=Mendelian randomisat*	273
78	TS=(RCT? or (randomi?ed NEAR/2 (control* or intervention* or experiment* or trial* or study or studies)))	654068
79	TS=((random* or comparative or intervention? or treatment?) NEAR/3 (efficacy or effect*))	196151
80	TS=(clinical NEAR/0 (intervention? or trial?))	242109
81	#78 OR #79 OR #80	975661
82	#77 AND #81	40
83	TS=instrument* variab*	55089
84	TS=((causal* or causative) NEAR/3 (associat* or infer* or implicat* or effect* or predict* or factor? or risk? or relat*))	67811
85	TS=((gene* NEAR/2 (associat* or risk? or varia* or determinant?)) or risk variant?)	394086
86	TS=(disease* NEAR/3 (expos* or associat* or etiolog* or pathogenesis or risk?))	257979
87	TS=risk factor?	841646
88	TS=(confound* or nonconfound* or non-confound*)	116352
89	TS=(statistics or epidemiolog* or ((genetic* or molecular) and medicine))	898995
90	#83 OR #84 OR #85 OR #86 OR #87 OR #88 OR #89	2342704
91	#82 and #90	36
92	#76 or #91	6616705
93	#60 and #92	4874

A.3.4 Code to search preprints

As detailed extensively in the main text, the inclusion of preprinted literature was a key aspect of the review presented in Chapters 3 & 4. Using the `medrxivr` tool presented in Chapter 2.

Given the relatively small number of records contained in the preprint server, no study design filters were employed. For the same reason, the number of search terms is smaller compared with the main database searches, as terms returning no records have been removed.

```

library(medrxivr)

# Define exposure search terms
topic1 <- c(mx_caps("statin"),
            mx_caps("ldl"),
            mx_caps("hdl"),
            mx_caps("TG"),
            mx_caps("triglycer"),
            paste0("\b",mx_caps("TC"),"\b"),
            mx_caps("ezetim"),
            mx_caps("fibrate"),
            mx_caps("bile acid"),
            mx_caps("lipoprotein"),
            mx_caps("lipid"),
            mx_caps("cholesterol"))

# Define outcome search terms
topic2 <- c(mx_caps("dementia"),
            mx_caps("alzheim"),
            mx_caps("MCI"),
            mx_caps("mild cognitive"))

# Combine search topics using Boolean AND
query <- list(topic1, topic2)

# Run search in bioRxiv
bx_data <- mx_api_content(server = "biorxiv",
                           to_date = "2019-08-01")
bx_results <- mx_search(bx_data, query)

```

```
# Run search in medRxiv
mx_data <- mx_api_content(server = "medrxiv",
                            to_date = "2019-08-01")
mx_results <- mx_search(mx_data, query)
```

A.3.5 Calculating Gwet's AC1

Gwet's agreement coefficient, $AC1$ is defined as:

$$AC1 = \frac{\text{observed agreement} - \text{chance agreement}}{1 - \text{chance agreement}} \quad (\text{A.1})$$

In reference to a two-by-two table with cells A, B, C and D, it is calculated using the following:

$$AC1 = \frac{\frac{A+D}{N} - e(\gamma)}{1 - e(\gamma)} \quad (\text{A.2})$$

where $e(\gamma)$ is the chance agreement between raters, given as $2q(1 - q)$, where

$$q = \frac{(A + C) + (A + B)}{2N} \quad (\text{A.3})$$

A.3.6 Risk-of-bias assessment tool for Mendelian randomisation studies

A copy of the risk of bias tool used to assess Mendelian randomisation studies is presented in Table ??, below.

Table A.9: Tool used to assess risk of bias in Mendelian randomisation studies, adapted from that developed by Mamluk et al.¹⁶⁶

Bias domain	Question	Risk of bias judgement		
		High	Moderate	Low
Weak instrument bias	Strength of association between instrument and exposure F statistic < 10 in the same sample (F < 10 indicating a weak instrument)	F < 10 F ~ 10	F missing or F > 10	

Table A.9: Tool used to assess risk of bias in Mendelian randomisation studies, adapted from that developed by Mamluk et al.¹⁶⁶ (*continued*)

Bias domain	Question	Risk of bias judgement		
		High	Moderate	Low
Genetic confounding bias	Reported test on association between confounders and IV (testing the assumption that the instrument is associated with the outcome only via the exposure)	Yes AND there is an obvious association	Not presented or Yes presented AND there is some degree of association	Presented and no obvious association
?Other? confounding bias	Included confounders in the IV analysis	Yes	No	

Table A.9: Tool used to assess risk of bias in Mendelian randomisation studies, adapted from that developed by Mamluk et al.¹⁶⁶ (*continued*)

Bias domain	Question	Risk of bias judgement		
		High	Moderate	Low
Additional direct effects between IV and outcome (exclusion restriction assumption)	Presence of pleiotropy for genetic IVs mechanism for genetic variant-lipid association (e.g. GWAS hit, could be acting through any pathway?)	Genetic IVs with no knowledge of mechanism for genetic variant-lipid association (e.g. GWAS hit, could be acting through any pathway?)	Biologically plausible lipid-specific mechanism of association for genetic variant-lipid (e.g. lipid metabolising genetic variants on outcome or its risk-factors variants)	Same as moderate AND checks that there is no other known effect of genetic variants on outcome or its risk-factors variants
Bias due to selection of participants	Homogenous population or similar ancestry? If no: Stratified by ethnicity or etc.) adjusted for population stratification (yes/no)?	Non-homogenous population (e.g. black and white together, etc.)	Population described as homogenous (e.g. whites only) BUT not corrected for ancestry informative markers like principal components derived from GWAS	?Population described as homogenous (e.g. whites only) AND corrected for ancestry informative markers like principal components derived from GWAS

A.4 Chapter 4

A.4.1 Dose response plots by fraction

In addition to examining a non-linear relationship between LDL-c and Alzheimer's disease (see Figure 4.16), I performed a dose response analysis for two other exposure-outcome pairs. The results of these analyses are presented below.




Figure A.1: Dose-response meta-analysis of total cholesterol on all-cause dementia




Figure A.2: Dose-response meta-analysis of low-density lipoprotein cholesterol on Alzheimer's disease

A.5 Chapter 5

A.5.1 Amendments to protocol

Any changes made to the approved protocol are “Minor amendments” as per the ISAC criteria, specifically falling under the category of “Additional methods to further control for confounding or sensitivity analysis provided these are to be reported as secondary to the main findings”.

Changes include:

- Use of a time-varying treatment indicator, to correctly classify time-at-risk.
- Inclusion of additional covariates that are adjusted for in the main model.

A.5.2 Minimum follow-up period

Enforcing a minimum period of follow-up is normally used to reduce the risk of reverse causation in an analysis by excluding events occurring within a certain time-frame post-exposure. With respect to dementia outcomes, which have a long prodromal period, enforcing a minimum period of follow-up can add biological validity to an analysis. Incident cases of dementia that occur within a few years of starting a drug may already be beyond effective preventive treatment at the time the drug was started, and so restricting the analysis to those with longer exposure to treatment may provide a more reliable estimate of the effect.

Panel A of Figure A.3 describes a standard analysis of an exposure with a minimum enforced follow-up period (in this case, 10 years). Participants enter the cohort and are classified at that point as either exposed or unexposed, for example, on the basis of a total cholesterol measurement. Participants are followed-up over time, but events occurring before the minimum period of follow-up are discarded.




Figure A.3: shortcap:

However, applying a similar approach to the analysis performed in Chapter 5, illustrated in Panel B of Figure A.3, is challenging. The use of a time updating treatment indicator results in two potential starting points for the 10-year lag: entry into the cohort or initiation of statin therapy. As the purpose of this analysis is to investigate the impact of statin treatment, starting the minimum period of follow-up from cohort entry is not helpful as it does not ensure that participants have been exposed to the drug for 10 years. Alternatively, defining the minimum period of follow-up from statin initiation cause issues around the treatment of time-at-risk before statin initiation and around the start point for the follow-up period in those never prescribed a statin. In summary, either approach is likely to be substantially biased.

With these caveats in mind, below I present a sensitivity analysis of the association of statins with dementia outcomes, enforcing a minimum 10-year period of follow-up from statin initiation in those prescribed a statin.

Hold for figure

Figure A.4: Estimates from the

A.6 Chapter 6

A.6.1 Covariate matrix

Table A.10: Overview of missing data across covariates (described as N (%)) included in IPD analysis. Note that LDL is missing completely for the Whitehall and CaPS datasets as this measure was derived from the measures for TC, HDL and TG following multiple imputation of missing values in these variables.

Variable	Whitehall II	Epic	CaPS
Age group	182 (2.2%)	Complete	Complete
Male	Complete	Complete	Complete
BMI	30 (0.4%)	60 (5.4%)	X
Education	Complete	1 (0.1%)	X
Smoke ever	1454 (17.7%)	7 (0.6%)	10 (0.4%)
Alcohol ever	1344 (16.4%)	26 (2.3%)	129 (5.1%)
Ethnicity	30 (0.4%)	6 (0.5%)	X
IHD	Complete	1 (0.1%)	X
Diabetes	372 (3.3%)	1 (0.1%)	9 (0.4%)
TC	82 (1%)	114 (10.2%)	55 (2.2%)
LDL-c	*	140 (12.6%)	*
HDL-c	108 (1.3%)	141 (12.6%)	98 (3.9%)
TG	82 (1%)	114 (10.2%)	115 (4.6%)

Values shown are N (

A.6.2 Dementia definitions across cohorts

Table A.11: Definition of dementia outcomes across cohorts included in the individual participant data meta-analysis. Note the codes used to define dementia and related outcomes in the HES data as part of the Whitehall II cohort were taken from a previously published analysis.³⁶⁹

Cohort	Primary detection methods	Identification of dementia	Identification of AD	Identification of VaD
CaPS	DPUK derived variable from Phase V (2002-2004) data	Assessment at clinic by a combination of interviewers and clinicians.	NA	Not reported
EPIC Norfolk	Self-reported during 5th Health Checkup (2016-2018)	Has your doctor ever told you that you had any of the following: Dementia (Alzheimer's, Vascular Dementia, Lewy Body or other)? YES/NO	NA	NA
Whitehall II	Electronic health records, via the Hospital Episodes Statistics database	ICD-9: 290.0-290.4, 331.0-331.2, 331.82, 331.9 ICD-10: F00, G30 F00, F01, F03, G30, G31	ICD-9: 331.0 ICD-10: F00, G30	ICD-9: 290.4 ICD-10: F01

A.6.3 Comparison with previous analysis of the Whitehall II cohort




Figure A.5: Comparison of results across the two analysis of the Whitehall II cohort for the Alzheimer's disease outcome.

A.7 Chapter 7

A.7.1 Calculation of average adjustment values from previous expert elicitation exercise

The paper describing the bias/indirectness-adjusted meta-analysis method used expert elicitation to define a distribution for bias and indirectness across eight

studies examining the effectiveness of routine anti-D prophylaxis in Rhesus negative women.³⁴⁶ The authors consider five sources of bias (selection bias, performance bias, attrition bias, detection bias and other bias) and four sources of indirectness (population, intervention, control, outcome). For each domain of bias/indirectness, four experts were asked to define the 67% interval around the effect expected due to bias/indirectness in the absence of a true effect. Additive or multiplicative scales were used to capture the intervals for additive and proportional bias/indirectness, respectively. The authors then use assumed log-normality of these distributions to derive a mean and variance for each on the log scale.

The resulting data for two domains of bias (attrition and detection) in one study are shown in Table A.12.

Table A.12: Distributions of bias, defined by position and variance terms, assigned by four independent experts to attrition and detection bias in a single study. This data was obtained from a previous publication,³⁴⁶ and represents an illustrative subset of the larger dataset.

Study	Domain	Assessor	Position	Variance
1	Attrition bias	1	-0.11	0.01
		2	-0.08	0.01
		3	-0.11	0.01
		4	-0.05	0.00
1	Detection bias	1	-0.09	0.02
		2	-0.32	0.14
		3	-0.23	0.02
		4	0.00	0.01

In an attempt to inform the prior distributions used in the triangulation exercise presented in Chapter 7, I first averaged the position and variance values across the four experts for each source of bias/indirectness within each study. I took the average of the absolute of the position value, as I wanted to obtain the average

adjustment regardless of direction. The results of this process for the example data presented below are shown in Table A.13.

Table A.13: Position and variance values for two example bias domains (attritiona and detection), obtained by averaging the absolute of the values contained in A.14 across the four reviewers.

Study	Domain	Position	Variance
1	Attrition bias	0.0875	0.0075
1	Detection bias	0.1600	0.0475

I then took the average and maximum of the expert-averaged position and variance values. The results of this for the example data are presented in Table A.14.

Table A.14: Mean and maximum of position and variance values described in Table A.14

Mean position	Mean variance	Max position	Max variance
0.12375	0.0275	0.16	0.0475

This two step approach (averaging across assessors before taking the average and maximum across domains/studies) was important. While the final average values will be the same whichever approach is taken, in a one-step process, the maximum value can be artificially inflated by a single expert's estimate. For example, in the illustrative data subset above, Assessor 2 assigns a distribution of $N(-0.32, 0.14)$, which would double the maximum position (0.16) and triple the maximum variance (0.0475) obtained when first averaging across experts.

The distributions arising from this applying this approach to the entire dataset, stratified by bias/indirectness and additive/proportional effects, are presented in Table A.15.

Table A.15: Mean and maximum distributions for bias and indirectness obtained from data resulting from the elicitation exercise reported in *Turner et al*³⁴⁶. Note I did not calculate values for additive indirectness, as in my analysis, all indirectness was assumed to be proportional.

	Mean	Max
Bias (Additive)	N(0.09, 0.06)	N(0.42, 0.29)
Bias (Proportional)	N(0.03, 0.02)	N(0.31, 0.21)
Indirectness (Proportional)	N(0.07, 0.02)	N(0.26, 0.16)

The mean values shown were used to define the distributions assigned to “Moderate” bias/indirectness judgements in my analysis (see Tables 7.3 & 7.4).

A.7.2 Comparison of two scenarios of bias

As discussed in the main text, I defined two sets of additive bias distributions as a sensitivity analysis. The results for the two case studies under the two scenarios are presented in the figures below. Under the first, shown on the left of the subsequent forest plot, the difference in the position value between the distributions assigned to “High” and “Moderate” bias levels is the same as between the “Moderate” and “Low” levels. Under the second scenario, the difference in the position value between the distributions assigned to “High” and “Moderate” is twice that between “Moderate” and “Low”.




Figure A.6: Comparison of bias-/indirectness-adjusted meta-analyses of LDL-c/Alzheimer's disease, using two different scenarios for the relationship of levels of bias.




Figure A.7: Comparison of bias-/indirectness-adjusted meta-analyses of triglycerides/vascular dementia, using two different scenarios for the relationship of levels of bias. Scenario 1: position value between the distributions assigned to "High" and "Moderate" bias levels is the same as between the "Moderate" and "Low" levels. Scenario 2: position value between the distributions assigned to "High" and "Moderate" bias levels is twice that between "Moderate" and "Low" levels

A.8 Chapter 8

A.8.1 Reproducible research

Reproducible science has been a key theme running through this thesis, as reflected by the development of open source tools to search preprinted literature, create visualisations of risk-of-bias data and to perform quantitative triangulation. In line with this, an open source copy of the code used to produce this thesis is available on GitHub, as is the code used to perform the analysis contained within it.

Unfortunately, given the nature of the data underpinning the CPRD (Chapter 5) and IPD (Chapter 6) analyses, it is not possible to make it publicly available. Access to the CPRD data is dependent on an application to its managing body, while access to the cohorts within the DPUK requires approval from both the DPUK and individual cohort owners. In the absence of data sharing, the best option to enable review and increase transparency is the open sharing of the code used to perform the analysis.¹³⁶

Finally, all software projects in these thesis attempt to conform to minimal best practices for research computing.^{370,371}

A.8.2 Catalogue of failures

Over the course of my PhD, I kept a record of attempted

Short section detailing the things I tried to do but which did not work:

- Failed to properly scope the literature and correctly estimate the number of studies returned by the search
- Failed IV analysis of anti-HC drugs because of low numbers of non-statins.
- Failed to realise that the initial code list used for the positive control outcome of IHD in the CPRD analysis (Chapter 5)

- Unsuccessful grant application with Andy
- Unsuccessful grant application with Chris Penfold
- Unsuccessful grant application with Matt
- Instrumental variable analysis of lipid regulation use using physicians prescribing preference.
- 2 x unsuccessful conference abstracts

B

Other Appendices

Contents

B.1	Software used to create this thesis	292
B.2	Producing risk-of-bias visualisations with <code>robvis</code>	295
B.3	Copies of papers arising from this thesis	298

B.1 Software used to create this thesis

This thesis was written in RMarkdown using the RStudio integrated development environment. Several R packages were used as part of this project.^{174,372–400} The version number of these packages is shown below:

```
## R version 3.6.1 (2019-07-05)
## Platform: x86_64-w64-mingw32/x64 (64-bit)
## Running under: Windows 10 x64 (build 19043)
##
## Matrix products: default
##
## locale:
## [1] LC_COLLATE=English_United Kingdom.1252
## [2] LC_CTYPE=English_United Kingdom.1252
## [3] LC_MONETARY=English_United Kingdom.1252
## [4] LC_NUMERIC=C
## [5] LC_TIME=English_United Kingdom.1252
##
## attached base packages:
## [1] stats      graphics    utils       datasets   grDevices methods   base
##
## other attached packages:
## [1] DiagrammeR_1.0.5     readxl_1.3.1      networkD3_0.4
## [4] ggdag_0.2.3          dagitty_0.3-1    sf_0.9-8
## [7] flextable_0.6.4       data.table_1.14.0 kableExtra_1.3.1
## [10] medrxivr_0.0.5.9000  robvis_0.3.0.900  png_0.1-7
## [13] metafor_3.0-2        Matrix_1.2-17    patchwork_1.1.1
## [16] ggplot2_3.3.5        dplyr_1.0.6      magrittr_2.0.1
```

B.1 - Software used to create this thesis

```
##  
## loaded via a namespace (and not attached):  
## [1] uuid_0.1-4                  backports_1.1.5          Hmisc_4.3-0  
## [4] systemfonts_1.0.1            plyr_1.8.5                igraph_1.2.6  
## [7] rbiorkxiv_0.2.0              sp_1.3-2                 splines_3.6.1  
## [10] TH.data_1.0-10              digest_0.6.27             htmltools_0.5.1.1  
## [13] magick_2.2                  viridis_0.5.1             fansi_0.4.2  
## [16] checkmate_2.0.0              cluster_2.1.0             openxlsx_4.1.4  
## [19] readr_1.3.1                 graphlayouts_0.5.0        extrafont_0.17  
## [22] officer_0.3.17              sandwich_3.0-1            extrafontdb_1.0  
## [25] colorspace_1.4-1            ggrepel_0.9.1             rvest_0.3.5  
## [28] haven_2.3.1                 xfun_0.22                 callr_3.7.0  
## [31] crayon_1.4.1                jsonlite_1.7.2            survival_3.2-3  
## [34] zoo_1.8-6                  glue_1.4.2                polyclip_1.10-0  
## [37] gtable_0.3.0                webshot_0.5.2             MatrixModels_0.4-1  
## [40] V8_3.0.1                   Rttf2pt1_1.3.7            rms_6.2-0  
## [43] SparseM_1.77                scales_1.1.0              mvtnorm_1.0-11  
## [46] DBI_1.1.1                  Rcpp_1.0.6                viridisLite_0.3.0  
## [49] htmlTable_1.13.2            units_0.6-6               irrCAC_1.0  
## [52] foreign_0.8-71              Formula_1.2-3            rsvg_1.3  
## [55] htmlwidgets_1.5.1            httr_1.4.1                RColorBrewer_1.1-2  
## [58] acepack_1.4.1                ellipsis_0.3.1            pkgconfig_2.0.3  
## [61] farver_2.0.1                nnet_7.3-12               utf8_1.2.1  
## [64] janitor_2.0.1              here_1.0.0                tidyselect_1.1.0  
## [67] labeling_0.3                 rlang_0.4.10              munsell_0.5.0  
## [70] cellranger_1.1.0            tools_3.6.1                visNetwork_2.0.9  
## [73] cachem_1.0.3                cli_2.5.0                 generics_0.1.0  
## [76] mathjaxr_1.4-0              evaluate_0.14             stringr_1.4.0  
## [79] mvmeta_1.0.3                 fastmap_1.0.1             yaml_2.2.1  
## [82] processx_3.5.2              utilities_0.3.0           knitr_1.36
```

B.1 - Software used to create this thesis

```
## [85] tidygraph_1.1.2           zip_2.1.1          dosresmeta_2.0.1
## [88] purrr_0.3.4              ggraph_2.0.0        nlme_3.1-140
## [91] quantreg_5.52            xml2_1.3.2         debugme_1.1.0
## [94] compiler_3.6.1           rstudioapi_0.13    curl_4.3
## [97] cranlogs_2.1.1           e1071_1.7-3        DiagrammeRsvg_0.1
## [100] tibble_3.1.0             tweenr_1.0.1       stringi_1.5.3
## [103] highr_0.8                ps_1.3.0          forcats_0.5.1
## [106] gdtools_0.2.3            rgeos_0.5-5        lattice_0.20-38
## [109] classInt_0.4-2           conflicted_1.0.4   vctrs_0.3.6
## [112] pillar_1.5.1             lifecycle_1.0.0    mixmeta_1.1.2
## [115] R6_2.5.0                 latticeExtra_0.6-28 bookdown_0.21.6
## [118] PRISMA2020_0.0.1        KernSmooth_2.23-15 gridExtra_2.3
## [121] rio_0.5.16               codetools_0.2-16    polspline_1.1.19
## [124] boot_1.3-22              MASS_7.3-51.6       rprojroot_2.0.2
## [127] withr_2.3.0              rnaturalearth_0.1.0 multcomp_1.4-17
## [130] hms_0.5.2                grid_3.6.1          rpart_4.1-15
## [133] tidyrr_1.1.3              rnaturalearthdata_0.1.0 class_7.3-15
## [136] rmarkdown_2.7.4            snakecase_0.11.0    ggforce_0.3.1
## [139] lubridate_1.7.9           base64enc_0.1-3
```

B.2 Producing risk-of-bias visualisations with **robvis**

The robvis package provides functions to convert a risk-of-bias assessment summary table into a summary plot or a traffic-light plot, formatted based on the specific risk-of-bias assessment tool used.

robvis currently contains templates for the following tools:

- ROB2
- ROBINS-I
- QUADAS-2
- ROB1

For the direction bias plots, the data expected by the tools is more comprehensive.

B.2 - Producing risk-of-bias visualisations with *robvis*

Table B.1: First five assessments contained in `data_rob2`, an example dataset for the Risk of Bias 2 tool for randomised controlled trials available via the `robvis` package.

Study	D1	D2	D3	D4	D5	Overall
Study 1	Low	Low	Low	Low	Low	Low
Study 2	Some concerns	Low	Low	Low	Low	Low
Study 3	No information	Low	Some concerns	Low	Low	Some concerns
Study 4	Low	Low	High	Low	Some concerns	High
Study 5	High	High	Low	Low	Some concerns	Low

Table B.2: Example data illustrating the structure required for bias direction plots. d_{ij} , d_{id} , and d_{it} define the extent (Serious, Moderate, Low), direction (right or left), and type (additive or proportional) of bias in the i th domain. Note that, because of the wide nature of the data, I have abbreviated the data in order to display it. A key for interpreting the data can be found below the table.

ID	yi	sei	d1j	d1d	d1t	d2j	d2d	d2t	d3j	d3d	d3t	d4j	d4d	d4t	d5j	d5d	d5t	d6j	d6d	d6t	d7j	d7d	d7t	overall
1	0.11	0.05	H	R	A	M	U	A	L	-	-	L	-	-	L	-	-	L	-	-	M	R	P	H
2	0.02	0.15	M	L	A	M	U	A	L	-	-	L	-	-	L	-	-	L	-	-	M	R	P	M
3	-0.06	0.13	M	L	A	M	U	A	L	-	-	L	-	-	L	-	-	L	-	-	M	L	P	M
4	-0.11	0.18	M	L	A	M	U	A	L	-	-	L	-	-	L	-	-	L	-	-	M	L	P	M
5	-0.12	0.18	H	L	A	L	-	-	L	-	-	L	-	-	L	-	-	L	-	-	M	L	P	H

Abbreviations: yi - Effect estimate; sei - Standard error of the effect estimate; H - High risk of bias; L - Low risk of bias; R - Right; L - Left; U - Unpredictable; A - Additive; P - Proportional

B.2.1 Simple visualisations

```
rob_summary(data = data_rob2, tool = "ROB2")
```




Figure B.1: shortcap:

```
rob_traffic_light(data = data_rob2,
                  tool = "ROB2",
                  psizes = 10)
```




Figure B.2: shortcap:

B.3 Copies of papers arising from this thesis

The following pages contain copies of the following publications:

- **McGuinness L. A., Higgins J. P. T., Walker, V. M., Davies, N. M., Martin, R. M., Coulthard, E., Davey-Smith, G., Kehoe, P. G., and Ben-Shlomo, Y.** (2021) *Association of lipid-regulating drugs with dementia and related conditions: an observational study of data from the Clinical Practice Research Datalink*. medRxiv. DOI: 10.1101/2021.10.21.21265131
- **McGuinness, L. A., and L Schmidt.** (2020) *medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R*. Journal of Open Source Software 5.54 2651. DOI: 10.21105/joss.02651
- **McGuinness, L. A., and Higgins J. P. T.** (2020) “*Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments.*” *Research Synthesis Method*). DOI: 10.1002/jrsm.1411
- **McGuinness, L. A., and Sheppard A. L.** 2020. “*A Descriptive Analysis of the Data Availability Statements Accompanying Medrxiv Preprints and a Comparison with Their Published Counterparts.*” *PLOS ONE* 16(5): e0250887. DOI: 10.1371/journal.pone.0250887
- *Hennessy, E. A., Acabchuk, R., Arnold, P. A., Dunn, A. G., Foo, Y. Z., Johnson, B. T., Geange, S. R., Haddaway, N. R., Nakagawa, S., Mapanga, W., Mengersen, K., Page, M., Sánchez-Tójar, A. Welch, V., and McGuinness L. A.* (2021). *Ensuring Prevention Science Research is Synthesis-Ready for Immediate and Lasting Scientific Impact.* *Prevention Science* . DOI: 10.1007/s11121-021-01279-8

These manuscripts arose from work performed as part of this thesis, as discussed more fully in Section 1.7.1.

1 Front matter

Type of manuscript:

Original Research Article

Title:

Association of lipid-regulating drugs with dementia and related conditions: an observational study of data from the Clinical Practice Research Datalink

Authors and affiliations:

Luke A McGuinness^{1,2,*}, Julian PT Higgins^{1,2,3}, Venexia M Walker^{1,2,4}, Neil M Davies^{1,2}, Richard M Martin^{1,2,3}, Elizabeth Coulthard⁵, George Davey-Smith^{1,2,3}, Patrick G Kehoe⁵ and Yoav Ben-Shlomo²

- (1) MRC University of Bristol Integrative Epidemiology Unit, Bristol, UK
- (2) Bristol Medical School: Population Health Sciences, University of Bristol, Bristol, UK
- (3) NIHR Biomedical Research Centre at University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol.
- (4) Department of Surgery, University of Pennsylvania Perelman School of Medicine, Philadelphia, USA
- (5) Bristol Medical School: Translational Health Sciences, University of Bristol, Bristol, UK

Corresponding author:

Luke McGuinness; Bristol Medical School, University of Bristol, Canyng Hall, 39 Whatley Road, Bristol, BS8 2PS, United Kingdom.; luke.mcguinness@bristol.ac.uk

Word counts

- Main text word count (max 4000): 3104
- Abstract word count (max 250): 250

2 Abstract

Background: There is some evidence that circulating blood lipids play a role in the development of Alzheimer's disease (AD) and dementia. These modifiable risk factors could be targeted by existing lipid-regulating agents, including statins, for dementia prevention. Here, we test the association between lipid-regulating agents and incidence of dementia and related conditions in the Clinical Practice Research Datalink (CPRD), an United Kingdom-based electronic health record database.

Methods: A retrospective cohort study was performed using routinely collected CPRD data (January 1995 and March 2016). Multivariable Cox proportional hazard models, allowing for a time-varying treatment indicator, were used to estimate the association between seven lipid-regulating drug classes (vs. no drug) and five dementia outcomes (all-cause, vascular and other dementias, and probable and possible Alzheimer's disease).

Results: We analyzed 1,684,564 participants with a total follow-up of 10,835,685 patient-years (median: 5.9 years (IQR:2.7-9.7)). We found little evidence that lipid-regulating agents were associated with incidence of Alzheimer's disease (probable HR:0.98, 95%CI:0.94-1.01; possible HR:0.97, 95%CI:0.93-1.01), but there was evidence of an increased risk of all-cause (HR:1.17, 95%CI:1.14-1.19), vascular (HR:1.81, 95%CI:1.73-1.89) and other dementias (HR:1.19, 95%CI:1.15-1.24). Evidence from a number of control outcomes indicated the presence of substantial residual confounding by indication (ischaemic heart disease HR: 1.62, 95%CI: 1.59-1.64; backpain HR: 1.04, 95%CI: 1.03-1.05; and Type 2 diabetes HR: 1.50, 95%CI: 1.48-1.51).

Conclusion: Lipid-regulating agents were not associated with reduced Alzheimer's disease risk. There was some evidence of an increased the risk of all-cause, vascular and other dementias, likely due to residual confounding by indication.

Keywords: Dementia; Alzheimer's disease; Lipids; Statins; Cohort study; Observational study; Electronic health records

3 Key messages

- A large cohort of patients from the Clinical Practice Research Datalink (CPRD) electronic health record database was assembled to examine the association of lipid-regulating agents, such as statins, with dementia outcomes.
- There was little evidence that lipid-regulating agents were associated with Alzheimer's disease, but there was some evidence for a harmful association with all-cause, vascular and other dementias. In all cases, the estimated associations were driven by the any statin subgroup, which comprised most participants in our cohort.
- Evidence from the control outcome analyses indicated strong residual confounding by indication, mostly likely related to vascular factors.

4 Introduction

Dementia is a major progressive neurocognitive disorder, the most common types of which are Alzheimer's disease, vascular dementia and Lewy Body dementia.(1) Despite an increasing number of cases globally and decades of research, there remains much unknown about the pathogenesis and progression of the disease, and, at present, no effective treatment exists to arrest, slow or reverse the cognitive decline associated with the condition.(2) Drug repurposing, the identification of new applications for previously approved drugs, may provide an efficient mechanism to discover new effective preventative and therapeutic treatments for dementia.(3,4)

Several cardiovascular factors have been identified as potential risk factors for dementia,(5) and of these, circulating lipid levels represent a promising target for intervention due to the ready availability of lipid-modifying treatments. In this context, determining whether lipid-regulating agents (LRA) could be repurposed for the prevention of dementia and related diseases would be helpful in the development of evidence-based prevention policy. Several existing prospective studies have examined the association of LRA use with dementia.(6–10) However, many of these studies are small, record few outcomes, and had limited follow-up.

The use of electronic health data for epidemiological research has several advantages.(11) As the data are collected through the routine care of a large cohort, they allow for nested cohort studies using sample sizes and time-scales which would be infeasible using traditional methods. In addition, data are collected for care provision and without a specific research question in mind, providing a holistic picture of a patient and their health experience. While they may not contain information on all potentially important confounders,(12) electronic health records provide routine data on common covariates for millions of participants.

We therefore aimed to test the association between several major classes of LRA and all-cause dementia, Alzheimer's disease, vascular dementia and other dementias, in the Clinical Practice Research Datalink (CPRD), a large, population-based electronic health record (EHR) database from the United Kingdom.(13)

5 Methods

5.1 Study design and protocol

We defined a longitudinal cohort using data from the CPRD. Our initial sample included all participants registered at a participating practice between 1 January 1995 and 29 February 2016 who had a flag for “research quality” data. Records pre-dating the 1995 cut-off were excluded from the analysis as data quality and reliability are thought to be higher after this date.(14) All events of interest were identified using predetermined code lists, which are available for inspection (see [Data/code availability](#)).

An *a priori* protocol for this study was published,(15) and amendments to this are recorded in Supplementary Materials 1. This study was reported in line with the RECORD guidelines (Supplementary Materials 2).(16)

5.2 Study Cohort

Participants were included in our study cohort if their record contained any of the following index events: (a) a code for a diagnosis of hypercholesterolemia or related condition; (b) a code for prescription of a LRA (such as statins); (c) a total cholesterol test result of >4mmol/L; (d) or a low-density lipoprotein cholesterol (LDL-c) test result of >2mmol/L.

These index events allowed us to define a population of participants who were either at risk of hypercholesterolemia, as indicated by the elevated total or LDL-c test results, or had already been diagnosed with it, as indicated by a diagnostic code or related prescription. This approach, conditioning entry into the study on being either “at-risk” or already diagnosed with hypercholesterolemia, was employed to reduce the “confounding by indication” that we would expect to observe if we had used a general population cohort.

An index date for each participant was defined as the date when the first relevant code or test result (as detailed above) was recorded on their clinical record. Participants were followed up until the earliest of (a) an outcome of interest; (b) death; (c) end of follow-up (29 February 2016); (d) last registration date with their GP practice; or (e) the last CPRD collection date for their practice. Participants were removed from our sample if they were less than 40 years of age at entry (as these patients are less likely to be prescribed a LRA), had less than 12 months of “research quality” data prior to their index date, were simultaneously prescribed more than one LRA (due to the difficulty of assigning these patients to a single exposure group), or were diagnosed with dementia before or on the date of the index event.

5.3 Exposures

We considered seven lipid-regulating drug classes based on groupings in the British National Formulary (BNF)(17), namely: statins, fibrates, bile acid sequestrants, ezetimibe, nicotinic acid groups, ezetimibe and statin (representing one treatment containing both drugs, rather than the two classes being prescribed concurrently), and omega-3 fatty acid groups.

To address the potential for immortal time bias, we employed a time-varying indicator of treatment status to correctly allocate time-at-risk to the exposed and unexposed groups.(18) Under this model, all participants entered the unexposed group on their index date and were moved into the exposed group on the date they were first prescribed an eligible LRA. Participants whose index event was a LRA prescription entered the study and the exposed group on the same day, and so contributed no time-at-risk to the unexposed group.

A participant's drug class was assigned based on their first recorded prescription, and any drug switching was ignored to mimic an intention-to-treat approach. We did however tabulate how often the initial drug class was stopped (defined as last prescription of the primary class being followed by at least six months of observation), added to (defined as a second drug class being prescribed before the last prescription of the initial class), or switched (defined as a second drug class being prescribed after the last prescription of the initial class).

5.4 Outcomes

We considered five outcomes as part of this analysis: probable Alzheimer's disease, possible Alzheimer's disease, vascular dementia, other dementias, and a composite all-cause dementia outcome. When two or more outcomes were coded in a participant's clinical record, a decision tree was used to differentiate between them (Supplementary Figure 1). The diagnosis date of the outcome was determined by the first record of a relevant code.

5.5 Covariates

The analysis was adjusted for a range of baseline covariates including sex, grouped year of entry into the cohort (<=2000, 2001-2005, 2006-2010, >2010), Charlson co-morbidity index, Index of Multiple Deprivation (IMD), consultation rate, alcohol use (current, former, never), smoking (current, former, never), BMI, baseline total cholesterol, and history of cardiovascular disease, coronary bypass surgery, coronary artery disease, peripheral arterial disease, hypertension, chronic kidney disease, and Type 1 and Type 2 diabetes. These variables were selected as potential confounders between dementia outcomes and use of a LRA. All covariates were determined at the index date and definitions for each can be found in Supplementary Table 1.

5.6 Analysis plan

All analyses were performed in STATA (V16, StataCorp LLC, College Station, TX). Cox proportional hazard models with a time-varying treatment indicator were used to estimate the hazard ratio and corresponding 95% confidence intervals, allowing for potential clustering of outcomes by practice. Participant's age was used as the time axis for all models.(19–21) To observe the effect of adjusting for additional covariates, we compared models adjusted for age only and age and sex with the fully adjusted model. Additional analyses stratified by outcome and drug class were also performed.

In the case of missing data, we used multiple imputation by chained equations (MICE) in STATA to create 20 imputed datasets.(22) All covariates included in the analytic model were also included in the imputation model.(23) The full imputation model is available for inspection (See [Data/Code availability](#) section).

5.7 Sensitivity analyses

We performed several sensitivity analyses. As statins are contraindicated in pregnancy,(24) we ran the models described above but excluding participants below the age of 55. Given the different ability of lipophilic statins to cross the blood-brain barrier,(25) we further stratified the statin exposure group into lipophilic (Atorvastatin, Lovastatin, Simvastatin, Cerivastatin) and hydrophilic (Pravastatin, Rosuvastatin, Fluvastatin) statins. Finally, we included three alternative outcomes with known associations with statin use as positive or negative controls using the fully adjusted model: back pain (negative control), ischaemic heart disease (positive protective control), and Type 2 diabetes (positive harmful control).(26,27)

6 Results

6.1 Patient characteristics

A total of 1,684,564 participants met the inclusion criteria for our cohort (see Supplementary Figure 2 for the attrition flowchart), with a total follow-up of 10,835,685 patient years at risk. Most participants were included in the cohort due to an elevated test result (elevated cholesterol test result: 93%, prescription of LRA: 5.6%, code for hypercholesterolemia: 1%). The median age at the index date was 57 years (inter-quartile range (IQR):48-67) and participants were followed up for a median of 5.9 years (IQR:2.7-9.7). During follow-up, an all-cause dementia diagnosis was recorded for 41,830 patients (12,647 probable Alzheimer's disease, 9,954 possible Alzheimer's disease, 8,466 vascular dementia, 10,763 other dementia).

The number of events, time-at-risk and crude rates for each drug class, tabulated by dementia outcome, are shown in Table 1. Most participants (98.1%) prescribed a lipid-regulating agent were prescribed a statin. We excluded the "Ezetimibe and statins" and "Nicotinic acid groups" drug classes from subsequent subgroup analyses based on the extremely small number of participants in these groups (Table 1). The distribution of baseline characteristics across the remaining seven drug classes can be seen in Table 2. The stopping, addition and switching of drug classes was common across all exposure groups (Supplementary Table 2).

6.2 Missing data

Full covariate information was available for 450,234 participants (26.7%). Five key variables had some missing data: IMD 2010 score, a proxy for socioeconomic position that is measured as twentiles with 1 indicating the least deprived and 20 indicating the most deprived, was missing for 625,788 participants (37.1%), because it is only recorded for selected English practices; alcohol status was missing for 269,526 participants (16%); smoking status was missing for 84,424 participants (5%); BMI, or a calculated BMI from height and weight measurements, was missing for 266,672 participants (15.8%); baseline total cholesterol was missing for 119,675 participants (7.1%); and baseline LDL cholesterol was missing for 787,289 participants (46.7%).

6.3 Primary analysis

Alzheimer's disease

As shown in Figure 1, our results show little evidence of an association between lipid-regulating agents and probable (HR:0.98, 95%CI:0.94-1.01) or possible (HR:0.97, 95%CI:0.93-1.01) Alzheimer's disease when compared with no treatment, except for an adverse association between fibrates and probable Alzheimer's disease (HR:1.52, 95%CI:1.13-2.03).

Non-Alzheimer's disease dementias

In contrast to the findings for Alzheimer's disease outcomes, lipid-regulating agents were associated with an increased risk of a subsequent diagnosis of vascular dementia (HR:1.81, 95%CI:1.73-1.89) or other dementias (HR:1.19, 95%CI:1.15-1.24). Again, the observed estimate was driven mainly by the any statin subgroup, but there was some evidence that ezetimibe was associated with an increased risk of vascular (HR:2.33, 95%CI:1.11-4.89) and other (HR:1.88, 95%CI:1.01-3.5) dementias.

All-cause dementia

For the composite all-cause dementia outcome, we found treatment with a lipid-regulating agent was associated with a slightly increased risk (HR:1.17, 95%CI:1.14-1.19), which lies between the associations for the Alzheimer and non-Alzheimer dementia outcomes as would be expected. There was also some evidence that fibrates were associated with increased risk of all-cause dementia (HR:1.28, 95%CI:1.08-1.52).

6.4 Sensitivity analyses

Adjustment for additional covariates beyond age and sex had a limited impact (Supplementary Figure 3), except for the probable Alzheimer's disease outcome, where the full adjustment attenuated to the null the protective association observed when adjusting only for age and sex.

Removing participants aged 55 and under at the index date from our analysis had minimal effect on our estimates (Supplementary Figure 4). When stratifying by statin properties, hydrophilic statins were less harmful in relation to the any, vascular and other dementias outcomes compared with lipophilic statins (Supplementary Figure 5). Similarly, hydrophilic statins were associated with a reduced incidence of Alzheimer's disease, compared with the absence of evidence for an association with lipophilic statins.

For our control outcomes (Supplementary Figure 6), there was some evidence that patients prescribed a lipid-regulating agent had an increased risk of back pain (HR: 1.04, 95%CI: 1.03-1.05), ischaemic heart disease (HR: 1.62, 95%CI: 1.59-1.64) and Type 2 diabetes (HR: 1.50, 95%CI: 1.48-1.51).

7 Discussion

7.1 Main findings

There was little evidence for an association between lipid-regulating agents and probable and possible Alzheimer's when compared with no treatment, but some evidence they were associated with an increased risk of an all-cause dementia, secondary to their association with vascular and other dementias diagnoses. The association observed in each case was driven by the any statin subgroup, which included a substantial majority of participants. For the other drug classes, there was limited evidence of an association with any outcome, with two exceptions. Ezetimibe was associated with increased risk of vascular and other dementias, while fibrates were associated with increased risk of all-cause dementia and probable Alzheimer's disease.

7.2 Comparison with other literature

Much of the existing literature focuses on the association of statins alone with neurodegenerative outcomes, with other lipid-regulating agents being grouped as "non-statin cholesterol-lowering drugs,"(8) echoing the distribution of participants in our analysis.

Statins and all-cause dementia

A recent Cochrane Review identified two randomized trials comparing treatment with statins versus non-treatment for the prevention of dementia, only one of which presented information on the incidence of dementia.(28) This study (Heart Protection Study) showed no effect of treatment with simvastatin on all-cause dementia risk (OR: 1.00, 95%CI:0.61-1.65),(29) but concerns were raised over the diagnostic criteria used. A meta-analysis of 30 observational studies found a reduced risk of all-cause dementia was associated with statin treatment (RR 0.83, 95%CI: 0.79–0.87).(30) This conflicts with the findings of our analysis, where statin use was associated with an increased risk of all-cause dementia (HR:1.17, 95%CI:1.14-1.19). Some of the included studies in the meta-analysis specifically exclude vascular dementia from the definition of all-cause dementia,(31) which may limit the ability for comparison with our findings for the all-cause dementia outcome.

Additionally, a previous analysis of the THIN EHR database using a propensity-score matched analysis found a protective effect of statins on all-cause dementia (HR:0.81, 95%CI:0.69-0.96)(32). Differences in the code-lists used to define dementia outcomes, in addition an analytical approach that adjusted for covariates defined after the index date, may go some way to explaining the discrepancy.

Statins and Alzheimer's disease

Our results are broadly in line with the findings of two distinct approaches examining the effect of statin treatment on subsequent Alzheimer's disease. No randomized trials of statins for the prevention of Alzheimer's disease have been reported, but a recent meta-analysis of 20 observational studies found statins were associated with a reduced risk of Alzheimer's disease (RR 0.69, 95% CI 0.60–0.80) with stronger evidence than observed in our analysis.(30) This review included case-control studies and analyses likely to be at risk of immortal time bias, which may account for the discrepancy with our findings.

Additionally, a recent Mendelian randomization study examining the effect of genetic inhibition of HMGCR on Alzheimer's disease (a genetic proxy for statin treatment) provided equivocal evidence (OR: 0.91, 95%CI: 0.63-1.31) but was consistent with our results.(33)

Our additional analyses stratified by statin properties found little evidence of differences in associations of lipophilic and hydrophilic statins and incidence of Alzheimer's disease, consistent with a recent meta-analysis of observational studies.(6)

Statins and vascular/other dementia

Far fewer studies have tested the association between lipid-regulating agents and vascular dementia or other dementias. A recent review found four observational studies examining the association of statins and vascular dementia found limited evidence for an effect (RR:0.93, 95% CI 0.74–1.16).(30) This contrasts with the harmful association found in our analysis (HR:1.81, 95%CI:1.73-1.89). When stratifying by lipid properties, lipophilic statins were more harmful than hydrophilic statins in vascular dementia, potentially due to their ability to cross the blood brain barrier.

Other drug classes

Apart from statins, few studies examining a lipid-regulating agent have been reported. One of the few classes for which a previous analysis was available were fibrates, which found little evidence of an association with all-cause dementia was identified,(8) inconsistent with our finding that patients prescribed fibrates had higher all-cause dementia risk than those prescribed other lipid lowering agents.

A previous Mendelian randomization study found little evidence that genetic variants that proxy for ezetimibe affect risk of Alzheimer's disease (OR: 1.17, 95%CI: 0.73-1.87),(33) consistent with our findings. To our knowledge, there is no previous study of the effect of preventative treatment with ezetimibe on risk of vascular dementia.

7.3 Strengths and limitations

A major strength of our analysis is the size of the included cohort and the length of follow-up that the use of electronic health records allowed. In addition, we followed users and non-users from a common index date, using a time-updating treatment indicator to correctly assign time-at-risk to the exposed and unexposed groups.

However, the findings of our analysis are subject to several limitations. Confounding by indication is a major bias in pharmacoepidemiological studies and could provide a potential explanation for the observed increased risk of vascular and other dementias with lipid regulating agent use. Patients with vascular risk factors are more likely to receive a statin prescription and also to be diagnosed with vascular dementia. Supporting evidence for this interpretation comes from a variety of sources, including the results of the control outcome analyses. This would explain the increased risk for the ischaemic heart disease outcome, for which statins are known to be protective, whilst almost no association was observed with the backpain outcome, indicating that most of the uncontrolled confounding is likely to be related to vascular factors. Additionally, we obtained the expected harmful result for Type 2 diabetes, where statins' mechanism of action on this outcome is unlikely to be vascular.(26,34,35) Further supporting evidence comes from the increasingly harmful association observed when moving from the probable and possible Alzheimer's disease outcome to the other dementias outcome, and finally to the vascular dementia outcomes. This pattern suggests that the strength of the residual confounding by indication increases as the proportion of cases with a vascular component in an outcome definition increases. A review of other available literature suggests that this observation (a harmful effect of lipid regulating agents on vascular-related outcome due to confounding by indication) is not unusual. Using a conventional epidemiological technique, a previous analysis also found an increased risk of coronary heart disease (analogous to the ischaemic heart disease outcome used in our analysis) in those taking statins (HR: 1.31, 95%CI: 1.04-1.66).(36) Controlling for confounding by indication in that study through the use of a trial emulation analysis gave an estimate of 0.89 (95%CI: 0.73-1.09), a comparable though less precise estimate to that observed in RCTs of statin use (0.73, 95%CI: 0.67-0.80).(37)

A secondary limitation is the potential for differential outcome misclassification based on the exposure, as we cannot exclude the possibility that for people with memory complaints, a diagnosis of vascular dementia might be made more frequently than Alzheimer's disease if their medical records contain prescriptions for lipid-regulating agents. Further, there is also the potential for non-differential misclassification of the outcome, based on the use of electronic health records to identify dementia cases.(38,39)

A further limitation stems from uncontrolled confounding due to genetic factors. Number of ApoE $\epsilon 4$ alleles represents the strongest genetic risk factor for Alzheimer's disease, but also substantially increases LDL-c levels,(40) potentially prompting treatment with a statin or other lipid regulating agent. We were unable to control for ApoE genotype in this analysis as we did not have access to genetic data on participants. As a result, any protective association between LRA use and the Alzheimer's disease outcomes may be masked by residual negative confounding by ApoE.

Finally, as with many studies of dementia, there is a risk of reverse causation in our analysis. Dementia and associated conditions have a long prodromal period, during which preclinical disease could cause indications for the prescription of a lipid-regulating agent.

8 Conclusions

We have provided new evidence on the association of lipid-regulating agent prescription, predominantly statins, with all-cause dementia, Alzheimer's disease, vascular dementia, and other dementias. We found limited evidence that use of lipid-regulating agents was associated with incidence of probable or possible Alzheimer's disease, but they were associated with an increased risk of all-cause, vascular and other dementias. Despite our attempts to account for bias in our analysis, there is a strong potential for residual confounding, misclassification and reverse causation.

This is likely to explain the unexpected increase in risk of vascular dementia associated with statin use. Future research should aim to address these biases using newer methods such as a trial emulation analysis.

9 Declarations

Ethics approval

The protocol for this study was approved by the CPRD's Independent Scientific Advisory Committee (ISAC 15_246R). This study did not directly involve patients, and so further ethical approval was not required.

Data/code availability

This analysis used the CPRD-GOLD primary care dataset March 2016 snapshot (ISAC 15_246R), which is available upon application to the CPRD Independent Scientific Advisory Committee. The code lists used to define the outcomes and covariates for this study are available as a supplementary file. The code lists, as well as the scripts used to create the study cohort and perform the analyses, are also available from the GitHub repository for this project (<https://github.com/mcguinlu/CPRD-LRA>), which was archived at the time of publication on Zenodo (DOI: TBC).

Supplementary data

Supplementary data are available at IJE online.

Sources of funding

LAM is supported by an NIHR Doctoral Research Fellowship (DRF-2018-11-ST2-048). JPTH, RMM and GDS are supported by the NIHR Biomedical Research Centre at University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. YBS and JPTH are supported by the NIHR Applied Research Collaboration West (ARC West) at University Hospitals Bristol NHS Foundation Trust. LAM, VMW, NMD, RMM, JPTH and GDS are members of the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_00011/1, MC_UU_00011/4). JPTH is a National Institute for Health Research (NIHR) Senior Investigator (NF-SI-0617-10145). The views expressed in this article are those of the authors and do not necessarily represent those of the NHS, the NIHR, MRC, or the Department of Health and Social Care. NMD is supported by a Norwegian Research Council Grant number 295989.

Conflict of interest statement

None declared.

10 Tables

Table 1: Summary of number of events, participant-time-at-risk and crude rates by drug class and dementia outcome.

Exposure group	Any dementia			Probable AD			Possible AD			Vascular dementia			Other dementia		
	Events	PYAR	Rate ^a	Events	PYAR	Rate ^a	Events	PYAR	Rate ^a	Events	PYAR	Rate ^a	Events	PYAR	Rate ^a
No LRA (unexposed)	18,608	5,872,717	317	6,368	5,818,047	109	2,637	5,800,964	45	4,813	5,811,594	83	4,790	5,808,285	82
By drug class															
Statins	22,920	4,871,568	470	6,190	4,758,526	130	5,773	4,753,437	121	5,871	4,755,258	123	5,086	4,747,237	107
Omega-3 FAGs	19	8,034	236	4	7,927	50	7	7,950	88	4	7,938	50	4	7,925	50
Fibrates	141	38,003	371	49	37,102	132	21	36,835	57	36	37,001	97	35	36,983	95
Ezetimibe	32	6,604	485	8	6,429	124	7	6,425	109	12	6,444	186	5	6,393	78
BAS	106	36,370	291	28	35,808	78	19	35,726	53	26	35,768	73	33	35,808	92
Ezetimibe + Statins ^b	0	986	-	0	986	-	0	986	-	0	986	-	0	986	-
NAG	4	1,403	-	0	1,379	-	2	1,391	-	1	1,389	-	1	1,382	-
Total	41,830	10,835,686	386	12,647	10,666,205	119	8,466	10,643,714	80	10,763	10,656,378	101	9,954	10,644,999	94

^a Crude rate per 100,000 participant-years-at-risk

^b One treatment containing both drugs, rather than the two classes being prescribed concurrently

Abbreviations: AD - Alzheimer's disease; BAS - Bile acid sequestrants; LRA - Lipid regulating agent; NAG - Nicotinic acid groups; Omega-3 FAGs - Omega-3 fatty acid groups; PYAR - Participant-years-at-risk.

Table 2: Patient characteristics by drug class. Summary statistics are presented as "% (N)" unless otherwise specified in the variable name.

	Whole Sample	None	Statins	Bile acid sequestrants	Ezetimibe	Fibrates	Omega-3 Fatty Acid Groups
Sample size (N)	1,684,564	1,087,704	585,528	5,396	763	3,889	992
Year of cohort entry (median)	2006	2007	2004	2005	2004	2001	2005
Female	53.0% (893174)	56.2% (610950)	47.1% (276043)	66.4% (3585)	54.5% (416)	38.6% (1500)	52.6% (522)
Age at cohort entry (median)	57	54	62	57	60	58	56
CAD	0.4% (7133)	0.1% (589)	1.1% (6465)	0.1% (6)	0.9% (7)	1.4% (53)	1.3% (13)
CBS	0.3% (5699)	0.1% (682)	0.8% (4926)	0.1% (4)	0.4% (3)	2.0% (78)	0.6% (6)
CVD	2.1% (34899)	1.1% (11619)	3.9% (22977)	1.6% (86)	2.6% (20)	4.4% (170)	1.7% (17)
Charlson (ever > 0)	30.6% (516135)	25.1% (272642)	40.7% (238403)	42.5% (2292)	41.7% (318)	50.8% (1976)	40.4% (401)
IMD-2010 (median)	9	8	9	8	9	10	10
Consultation rate (mean/SD)	5.4 (5.4)	5.0 (5.0)	6.2 (6.1)	8.6 (7.4)	7.4 (6.6)	7.1 (6.2)	8.0 (8.0)
Alcohol (ever)	85.9% (1447151)	86.6% (941648)	84.7% (496110)	82.8% (4468)	84.0% (641)	82.9% (3223)	82.0% (813)
Smoking (ever)	51.1% (861355)	47.1% (511826)	58.6% (343074)	55.2% (2978)	57.5% (439)	60.2% (2341)	53.7% (533)
BMI (mean/SD)	27.0 (5.3)	26.7 (5.2)	27.7 (5.3)	26.8 (5.8)	28.1 (5.7)	29.0 (5.2)	26.9 (5.5)
PAD	0.7% (12613)	0.4% (4039)	1.4% (8424)	0.9% (47)	0.9% (7)	1.9% (75)	1.0% (10)
Hypertension	16.0% (269804)	11.5% (124604)	24.4% (143101)	12.8% (692)	23.9% (182)	25.8% (1002)	15.7% (156)
Total cholesterol (mean/SD)	5.7 (10.1)	5.5 (6.4)	6.2 (15.3)	5.3 (1.3)	7.1 (26.5)	6.4 (5.6)	5.6 (1.6)
LDL cholesterol (mean/SD)	3.6 (4.9)	3.4 (5.3)	4.0 (3.7)	3.1 (1.0)	3.9 (1.1)	3.3 (1.8)	3.2 (1.0)
CKD	0.1% (1295)	0.1% (740)	0.1% (545)	0.1% (6)	0.1% (1)	0.0% (0)	0.3% (3)
Type 1 Diabetes	0.2% (4037)	0.1% (785)	0.5% (3196)	0.3% (14)	1.0% (8)	0.8% (31)	0.1% (1)
Type 2 Diabetes	2.9% (48557)	1.1% (11797)	6.1% (35941)	2.3% (123)	5.4% (41)	15.8% (614)	2.8% (28)

Note: The 'Nicotinic acid groups' (n=165) and 'Ezetimibe and Statins' (n=127) subgroups are not shown, but are included in the whole sample column
Abbreviations: BMI - Body mass index; CAD - Coronary arterial disease; CBS - Coronary bypass surgery; CKD - Chronic kidney disease; CVD - Cardiovascular disease; IMD - Index of multiple deprivation; LRA - Lipid regulating agent; PAD - Peripheral arterial disease; SD - Standard deviation.

11 Figures




Figure 1: Results from the primary analyses of CPRD data comparing prescription of an lipid-regulating drug with no prescription, stratified by dementia outcome and drug class. All results were obtained using the fully adjusted model and participant age as the time scale.

12 Bibliography

1. Prince M, Ali G-C, Guerchet M, Prina AM, Albanese E, Wu Y-T. Recent global trends in the prevalence and incidence of dementia, and survival with dementia. *Alzheimer's Research & Therapy*. 2016 Jul;8(1):23.
2. Jobke B, McBride T, Nevin L, Peiperl L, Ross A, Stone C, et al. Setbacks in Alzheimer research demand new strategies, not surrender. *PLoS medicine*. 2018 Feb;15(2):e1002518.
3. Ashburn TT, Thor KB. Drug repositioning: Identifying and developing new uses for existing drugs. *Nature Reviews Drug Discovery*. 2004;3(8):673.
4. Pushpakom S, Iorio F, Eyers PA, Escott KJ, Hopper S, Wells A, et al. Drug repurposing: Progress, challenges and recommendations. *Nature Reviews Drug Discovery*. 2019 Jan;18(1):41–58.
5. Walker VM, Davies NM, Martin RM, Kehoe PG. Comparison of Antihypertensive Drug Classes for Dementia Prevention. *Epidemiology*. 2020 Nov;31(6):852–9.
6. Chu C-S, Tseng P-T, Stubbs B, Chen T-Y, Tang C-H, Li D-J, et al. Use of statins and the risk of dementia and mild cognitive impairment: A systematic review and meta-analysis. *Scientific Reports*. 2018 Apr;8(1):5804.
7. Larsson SC, Markus HS. Does Treating Vascular Risk Factors Prevent Dementia and Alzheimer's Disease? A Systematic Review and Meta-Analysis. *Journal of Alzheimer's Disease*. 2018 Jan;64(2):657–68.
8. Ancelin M-L, Carrière I, Barberger-Gateau P, Auriacombe S, Rouaud O, Fourlanos S, et al. Lipid lowering agents, cognitive decline, and dementia: The three-city study. *Journal of Alzheimer's Disease*. 2012 Jan;30(3):629–37.
9. Bettermann K, Arnold AM, Williamson J, Rapp S, Sink K, Toole JF, et al. Statins, Risk of Dementia, and Cognitive Function: Secondary Analysis of the Ginkgo Evaluation of Memory Study. *Journal of Stroke and Cerebrovascular Diseases*. 2012 Aug;21(6):436–44.
10. Rea TD, Breitner JC, Psaty BM, Fitzpatrick AL, Lopez OL, Newman AB, et al. Statin Use and the Risk of Incident Dementia: The Cardiovascular Health Study. *Archives of Neurology*. 2005 Jul;62(7):1047–51.
11. Casey JA, Schwartz BS, Stewart WF, Adler NE. Using Electronic Health Records for Population Health Research: A Review of Methods and Applications. *Annual Review of Public Health*. 2016 Mar;37(1):61–81.
12. Farmer R, Mathur R, Bhaskaran K, Eastwood SV, Chaturvedi N, Smeeth L. Promises and pitfalls of electronic health record analysis. *Diabetologia*. 2018 Jun;61(6):1241–8.
13. Herrett E, Gallagher AM, Bhaskaran K, Forbes H, Mathur R, van Staa T, et al. Data Resource Profile: Clinical Practice Research Datalink (CPRD). *International Journal of Epidemiology*. 2015 Jun;44(3):827–36.

14. Wolf A, Dedman D, Campbell J, Booth H, Lunn D, Chapman J, et al. Data resource profile: Clinical Practice Research Datalink (CPRD) Aurum. *International Journal of Epidemiology*. 2019 Dec;48(6):1740–1740g.
15. Walker VM, Davies NM, Jones T, Kehoe PG, Martin RM. Can commonly prescribed drugs be repurposed for the prevention or treatment of Alzheimer's and other neurodegenerative diseases? Protocol for an observational cohort study in the UK Clinical Practice Research Datalink. *BMJ Open*. 2016;6(12).
16. Nicholls SG, Langan SM, Sørensen HT, Petersen I, Benchimol EI. The RECORD reporting guidelines: Meeting the methodological and ethical demands of transparency in research using routinely-collected health data. *Clinical epidemiology*. 2016;8:389.
17. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: A major update to the DrugBank database for 2018. *Nucleic Acids Research*. 2017;46(D1):D1074–82.
18. Lévesque LE, Hanley JA, Kezouh A, Suissa S. Problem of immortal time bias in cohort studies: Example using statins for preventing progression of diabetes. *BMJ*. 2010 Mar;340:b5087.
19. Lamarca R, Alonso J, Gomez G, Munoz A. Left-truncated Data With Age as Time Scale: An Alternative for Survival Analysis in the Elderly Population. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 1998 Sep;53A(5):M337–43.
20. Gail MH, Graubard B, Williamson DF, Flegal KM. Comments on 'Choice of time scale and its effect on significance of predictors in longitudinal studies' by Michael J. Pencina, Martin G. Larson and Ralph B. D'Agostino, *Statistics in Medicine* 2007; **26** :1343-1359. *Statistics in Medicine*. 2009 Apr;28(8):1315–7.
21. Pencina MJ, Larson MG, D'Agostino RB. Choice of time scale and its effect on significance of predictors in longitudinal studies. *Statistics in Medicine*. 2007 Mar;26(6):1343–59.
22. Sterne JAC, White IR, Carlin JB, Spratt M, Royston P, Kenward MG, et al. Multiple imputation for missing data in epidemiological and clinical research: Potential and pitfalls. *BMJ*. 2009 Jun;338:b2393.
23. Moons KG, Donders RA, Stijnen T, Harrell Jr FE. Using the outcome for imputation of missing predictor values was preferred. *Journal of clinical epidemiology*. 2006;59(10):1092–101.
24. National Institute for Health and Care Excellence. Cardiovascular disease: Risk assessment and reduction, including lipid modification. <https://www.nice.org.uk/guidance/cg181/ifp/chapter/Drug-treatments-to-lower-your-cholesterol>; NICE;
25. Sierra S, Ramos MC, Molina P, Esteo C, Vázquez JA, Burgos JS. Statins as neuroprotectants: A comparative in vitro study of lipophilicity, blood-brain-barrier

penetration, lowering of brain cholesterol, and decrease of neuron cell death. *Journal of Alzheimer's disease: JAD*. 2011;23(2):307–18.

26. Collins R, Reith C, Emberson J, Armitage J, Baigent C, Blackwell L, et al. Interpretation of the evidence for the efficacy and safety of statin therapy. *The Lancet*. 2016 Nov;388(10059):2532–61.
27. Herrett E, Williamson E, Brack K, Beaumont D, Perkins A, Thayne A, et al. Statin treatment and muscle symptoms: Series of randomised, placebo controlled n-of-1 trials. *BMJ*. 2021 Feb;372:n135.
28. McGuinness B, Craig D, Bullock R, Passmore P. Statins for the prevention of dementia. *The Cochrane Database of Systematic Reviews*. 2016 Jan;(1):CD003160.
29. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20 536 high-risk individuals: A randomised placebocontrolled trial. *The Lancet*. 2002 Jul;360(9326):7–22.
30. Poly TN, Islam MM, Walther BA, Yang H-C, Wu C-C, Lin M-C, et al. Association between Use of Statin and Risk of Dementia: A Meta-Analysis of Observational Studies. *Neuroepidemiology*. 2020;54(3):214–26.
31. Chao T-F, Liu C-J, Chen S-J, Wang K-L, Lin Y-J, Chang S-L, et al. Statins and the risk of dementia in patients with atrial fibrillation: A nationwide population-based cohort study. *International Journal of Cardiology*. 2015 Oct;196:91–7.
32. Smeeth L, Douglas I, Hall AJ, Hubbard R, Evans S. Effect of statins on a wide range of health outcomes: A cohort study validated by comparison with randomized trials. *British Journal of Clinical Pharmacology*. 2009 Jan;67(1):99–109.
33. Williams DM, Finan C, Schmidt AF, Burgess S, Hingorani AD. Lipid lowering and Alzheimer disease risk: A mendelian randomization study. *Annals of Neurology*. 2020 Jan;87(1):30–9.
34. Macedo AF, Douglas I, Smeeth L, Forbes H, Ebrahim S. Statins and the risk of type 2 diabetes mellitus: Cohort study using the UK clinical practice pesearch datalink. *BMC Cardiovascular Disorders*. 2014 Jul;14(1):85.
35. Smit RAJ, Trompet S, Leong A, Goodarzi MO, Postmus I, Warren H, et al. Statin-induced LDL cholesterol response and type 2 diabetes: A bidirectional two-sample Mendelian randomization study. *The Pharmacogenomics Journal*. 2020 Jun;20(3):462–70.
36. Danaei G, García Rodríguez LA, Cantero OF, Logan R, Hernán MA. Observational data for comparative effectiveness research: An emulation of randomised trials to estimate the effect of statins on primary prevention of coronary heart disease. *Statistical methods in medical research*. 2013 Feb;22(1):70–96.

37. Taylor F, Huffman MD, Macedo AF, Moore TH, Burke M, Davey Smith G, et al. Statins for the primary prevention of cardiovascular disease. *The Cochrane Database of Systematic Reviews*. 2013 Jan;2013(1):CD004816.
38. Wilkinson T, Ly A, Schnier C, Rannikmäe K, Bush K, Brayne C, et al. Identifying dementia cases with routinely collected health data: A systematic review. *Alzheimer's & Dementia*. 2018 Aug;14(8):1038–51.
39. McGuinness LA, Warren-Gash C, Moorhouse LR, Thomas SL. The validity of dementia diagnoses in routinely collected electronic health records in the United Kingdom: A systematic review. *Pharmacoepidemiology and Drug Safety*. 2019;28(2):244–55.
40. Bennet AM, Di Angelantonio E, Ye Z, Wensley F, Dahlin A, Ahlbom A, et al. Association of apolipoprotein E genotypes with lipid levels and coronary risk. *JAMA*. 2007 Sep;298(11):1300–11.

medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R

Luke A McGuinness^{1, 2} and Lena Schmidt¹

¹ Department of Population Health Science, University of Bristol ² MRC Intergrative Epidemiology Unit, University of Bristol

DOI: [10.21105/joss.02651](https://doi.org/10.21105/joss.02651)

Software

- [Review ↗](#)
- [Repository ↗](#)
- [Archive ↗](#)

Editor: Daniel S. Katz ↗

Reviewers:

- [@danielskatz](#)

Submitted: 08 September 2020

Published: 09 October 2020

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

An increasingly important source of health-related bibliographic content are preprints: preliminary versions of research articles that have yet to undergo peer review. The two preprint repositories most relevant to health-related sciences are [medRxiv](#) and [bioRxiv](#), both of which are operated by the Cold Spring Harbor Laboratory, a not-for-profit research and educational institution (Rawlinson & Bloom, 2019).

The goal of the `medrxivr` R package is two-fold. In the first instance, it provides programmatic access to the Cold Spring Harbour Laboratory (CSHL) API, allowing users to download medRxiv and bioRxiv preprint metadata (e.g., title, abstract, author list.) This functionality will be of interest to anyone who wishes to import medRxiv and/or bioRxiv preprint metadata into R, for example to explore the distribution of preprints by subject area or by publication year. Examples of this type of usage have already been reported (e.g., by Brierley, 2020).

In the second instance, the package provides functions that allow users to search the downloaded preprint metadata for relevant preprints using complex search strings, including functionality such as search term truncation, Boolean operators (AND, OR, NOT), and term proximity. Helper functions are provided that allow users to export the results of their search to a .bib file for import into a reference manager (e.g., Zotero) and to download the full-text PDFs of preprints matching their search. This aspect of the package will be more relevant to systematic reviewers, health librarians and others performing literature searches, allowing them to perform and document transparent and reproducible searches in these important evidence sources.

Acknowledgements

We acknowledge funding from NIHR (LAM through NIHR Doctoral Research Fellowship (DRF-2018-11-ST2-048), and LS through NIHR Systematic Reviews Fellowship (RM-SR-2017-09-028)). LAM is a member of the MRC Integrative Epidemiology Unit at the University of Bristol. The views expressed in this article are those of the authors and do not necessarily represent those of the NHS, the NIHR, MRC, or the Department of Health and Social Care.

References

- Brierley, L. (2020). The role of research preprints in the academic response to the COVID-19 epidemic. *Authorea*. doi:[10.22541/au.158516578.89167184](https://doi.org/10.22541/au.158516578.89167184)
- Rawlinson, C., & Bloom, T. (2019). New preprint server for medical research. *BMJ*, 365. doi:[10.1136/bmj.l2301](https://doi.org/10.1136/bmj.l2301)

Risk-of-bias VISualization (**robvis**): An R package and Shiny web app for visualizing risk-of-bias assessments

Luke A. McGuinness^{1,2}  | Julian P. T. Higgins^{1,2}

¹MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK

²Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

Correspondence

Luke A. McGuinness, Bristol Medical School, University of Bristol, Canyng Hall, 39 Whatley Road, Bristol, BS8 2PS, United Kingdom.

Email: luke.mcguinness@bristol.ac.uk

Funding information

National Institute for Health Research, Grant/Award Number: NF-SI-0617-10145Research Trainees Coordinating Centre, Grant/Award Number: DRF-2018-11-ST2-048

Despite a major increase in the range and number of software offerings now available to help researchers produce evidence syntheses, there is currently no generic tool for producing figures to display and explore the risk-of-bias assessments that routinely take place as part of systematic review. However, tools such as the R programming environment and **Shiny** (an R package for building interactive web apps) have made it straightforward to produce new tools to help in producing evidence syntheses. We present a new tool, **robvis** (Risk-Of-Bias VISualization), available as an R package and web app, which facilitates rapid production of publication-quality risk-of-bias assessment figures. We present a timeline of the tool's development and its key functionality.

KEY WORDS

data visualization, evidence synthesis, R, risk of bias

1 | INTRODUCTION

Synthesis of evidence from the totality of relevant research is becoming more important than ever in informing policy across an increasingly wide range of fields.¹ Risk of bias assessment—evaluation of the internal validity of studies included in a systematic review—often forms a key part of the evidence synthesis process, particularly in the health sciences.² A well-developed family of tools is widely used, which have in common the characteristic that they evaluate specific domains of bias rather than being constructed as a checklist or a quantitative score.² These tools include the RoB (Risk of Bias) 2 tool for randomized trials,³ the ROBINS-I (Risk Of Bias In Non-randomized Studies—of Interventions) tool,⁴ the QUADAS 2 (Quality and Applicability of Diagnostic Accuracy Studies) tool,⁵ and the ROBIS (Risk Of Bias in Systematic Reviews) tool.⁶ Within each bias domain a judgement is reached about the strength of the study in

that regard: for example, the first domain in the Cochrane RoB 2 tool deals with bias arising from the randomization process.³ Accessible graphics summarizing the results of these domain-based risk-of-bias assessments are included in reports of systematic reviews. A convenient plot in many reviews is a “traffic light” plot, which tabulates the judgement for each study in each domain. For larger numbers of studies, when such a table becomes unmanageable, a popular alternative is a (possibly weighted) bar plot, which shows the proportion of information with each judgement for each domain.⁷

Researchers can face a number of barriers in creating these plots. While some evidence synthesis platforms, such as Cochrane's Review Manager,⁸ can produce these visualizations, not all researchers use these systems to conduct their systematic reviews, and copying the risk-of-bias data into these systems solely to produce the plots is inefficient and error prone. On the other hand, creating the figures “by hand,” through software such as MS

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Research Synthesis Methods* published by John Wiley & Sons Ltd.

PowerPoint or Adobe Illustrator, may lead to unintentional errors and require the plots to be redrawn when a review is updated. Additionally, while the field of evidence synthesis software has grown rapidly in recent years,⁹ this growth has not been equally distributed across the different aspects of the systematic review process. For example, a recent review found several software offerings aimed specifically at the abstract screening stage of the review process,¹⁰ but no similar time- and error-reducing tool has been proposed for visualizing the results of risk-of-bias assessments.

It is now straightforward to produce such a tool, thanks to the availability of powerful computing offerings including R, RStudio, and **Shiny** (an R package for building interactive web apps).¹¹⁻¹³ Here, we present **robvis** (Risk Of Bias VISeualization),¹⁴ an R package and **Shiny** web-app that allows users to create publication-ready risk-of-bias plots quickly and easily. While primarily designed for use with the major risk-of-bias assessment tools used in health research (ROB2, ROBINS-I, and QUADAS-2), the tool allows users to visualize the results from any domain-based risk-of-bias assessment or quality appraisal tool.

The tool is open-source and available to use free of charge. Users can download a stable version of the R package from CRAN (<https://cran.r-project.org/package=robvis>); or access and contribute to the open-source code that powers the package via GitHub (<https://github.com/mcguinlu/robvis>). Extended guidance for the tool, including a step-by-step walk-through for those new to the R programming environment, is also available via the “Doing Meta-Analysis in R” online guide.¹⁵ Below, we discuss the tool’s development and key functionality.

2 | DEVELOPMENT

Development of **robvis** began in April 2019 at the Evidence Synthesis Hackathon (ESH), an event which brings together interested researchers, practitioners, and coders to discuss and develop new open-source evidence synthesis technologies. Test versions of both the R package and the web app were made available in early June 2019, with attendees of the ESH and members of the Bristol Appraisal and Review of Research (BARR) group at the University of Bristol being invited to test the tool and provide feedback. This feedback, along with other feature suggestions from the wider evidence synthesis community captured via GitHub issues, was incorporated and the first release version of the package was uploaded to CRAN in November 2019. The tool has been well received and is beginning to be cited in the evidence synthesis literature.¹⁶⁻²⁰

Highlights

- Risk-of-bias assessment is a key element of the systematic review workflow.
- No other tool exists explicitly for the purpose of visualizing risk-of-bias results.
- Here, we present **robvis**, an open-source R package and Shiny web app for creating publication-ready risk-of-bias assessment figures.
- **robvis** forms part of the metaverse, a collection of R packages designed to provide an evidence synthesis workflow in R.

3 | FUNCTIONALITY

3.1 | R package

3.1.1 | Tool templates and example data sets

At the time of writing, the tool includes templates for three major tools: the Cochrane RoB 2 tool for assessing randomized trials,³ the ROBINS-I tool for assessing non-randomized studies of interventions,⁴ and the QUADAS-2 tool for assessing diagnostic accuracy studies.⁵ These templates automatically apply the correct risk-of-bias domain names to the figures, and label the judgement levels appropriately (eg, “Low,” “Some Concerns,” and “High” in the case of the RoB 2 tool). In addition, **robvis** contains a general template that can be used to visualize the result of any domain-based assessment tool. The generic template has greater flexibility than the tool-specific templates, allowing a user-specified number of domains and custom domain titles to be used. It is suitable for use with the original version of the Cochrane risk-of-bias tool for randomized trials,²¹ in which flexibility in the specification of domains was permitted. Since users of more recent tools such as RoB 2 and ROBINS-I are not permitted to modify the domains, we strongly encourage use of the in-built templates for these.

In order to help users familiarize themselves with the package and its functionality, **robvis** contains built-in example data-sets for each template. We illustrate the example data for the RoB 2 tool for assessing risk of bias in randomized trials in Table 1.

3.1.2 | Data import

robvis expects the risk-of-bias data file to be arranged in a specific way (see Table 1 for an example). The first

TABLE 1 Example dataset for the ROB 2 tool contained within **robvis**. Data can be imported to the tool from with an Excel spreadsheet or a CSV file

Study	D1	D2	D3	D4	D5	Overall	Weight
Study 1	Low	Low	Low	Low	Low	Low	33.33
Study 2	Some concerns	Low	Low	Low	Low	Low	33.33
Study 3	Some concerns	Low	Some concerns	Low	Low	Some concerns	0.14
Study 4	Low	Low	High	Low	Some concerns	High	9.09
Study 5	High	High	Low	Low	Some concerns	Low	12.5
Study 6	Low	High	Some concerns	Low	Low	Some concerns	25
Study 7	Low	Some concerns	Some concerns	High	Low	Some concerns	200
Study 8	Low	Some concerns	Low	Low	Low	Low	11.11
Study 9	Low	Low	High	Low	Low	High	1.11

TABLE 2 Description of the arguments available in the two main functions of the **robvis** R package. “X” indicates that the option is available for the respective function

Argument	rob_traffic_light()	rob_summary()	Description
data	X	X	Defines the data frame containing the summary (domain) level risk-of-bias assessments. See the text and Table 1 for the format expected by robvis .
tool	X	X	Defines the risk of bias assessment tool used. The RoB2 (<code>tool = "ROB2"</code>), ROBINS-I (<code>tool = "ROBINS-I"</code>), and QUADAS-2 (<code>tool = "QUADAS-2"</code>) assessments tools are currently supported. Other tools can be visualized using the generic template (<code>tool = "Generic"</code>) [Note 1].
color	X	X	Defines the color scheme for the plot. The default is <code>colour = "cochrane"</code> which uses the “Cochrane” (red, yellow, green) colors, while a preset option for a color-blind friendly palette is also available (<code>colour = "colourblind"</code>). Alternatively, users can specify their own color scheme for example, <code>colour = c ("#f442c8," "#bef441," "#000000")</code> .
overall		X	Defines whether to include an additional bar showing the distribution of overall risk of bias judgments in the summary barplot figure. Default is <code>overall = FALSE</code> .
weighted		X	Defines whether weights should be used to produce the summary barplot figure. Default is <code>weighted = TRUE</code> , in line with current Cochrane Collaboration guidance.
psize	X		Defines the size of the points in the traffic light plot. Default is <code>psize = 20</code> .

Note: This option (`tool = "Generic"`) reflects the general template name used in the current development version of **robvis**, which will become the standard for all future iterations of the package. However, in the current CRAN version, the generic template is accessed using `tool = "ROB1"`.

column should contain the unique study/result identifier. The second-to-last column should contain the overall risk-of-bias judgments. The final column contains some measure of the result's precision (eg, the weight assigned to that result in a meta-analysis, or the sample size of the analysis that produced the result). This weight column is

used to create the summary bar plot, as current guidance recommends dividing the bars to show the proportion of information at each level of risk of bias, determined by the cumulative weight at that level, rather than simply showing the number of studies/results in each category.² If a measure of precision is not available, or to reproduce

“equally” weighted bar charts as have traditionally been presented in Cochrane Reviews to date, these weights may all be specified to be 1.

3.1.3 | Functions

robvis contains two main functions. The first, **rob_traffic_light()**, creates a *traffic light plot*. This displays every risk-of-bias judgement in a matrix, with domains along the horizontal and results/studies down the vertical, similar to the data set. The second function, **rob_summary()**, creates a *weighted bar plot*. This shows the proportion of information with each

risk-of-bias judgement separately for each domain in the assessment tool specified.

A worked example using these functions is outlined below, illustrating the simple steps involved in creating risk-of-bias plots using **robvis**. A detailed description of the additional options that can be used with each function is presented in Table 2. All examples produced in this article are created using the stable version available from CRAN.

To install and load the package in R, enter the following into the console:

```
install.packages("robvis")
library(robvis)
```




FIGURE 1 Example risk of bias traffic light plot of ROB2 assessments created using **robvis** and the colourblind palette [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 2 Example risk of bias summary plot of ROB2 assessments created using **robvis** and the standard Cochrane palette [Colour figure can be viewed at wileyonlinelibrary.com]

Using the example data set (`data_rob2`) that is built into the package and is presented in Table 1 for reference, the traffic light plot shown in Figure 1 is created using:

```
rob_traffic_light(data = data_rob2,
                  tool = "ROB2",
                  colour = "colourblind",
                  psize = 15)
```

Similarly, using the same data set, the summary barplot shown in Figure 2 is created using:

```
rob_summary(data = data_rob2,
            tool = "ROB2",
            overall = TRUE)
```

3.1.4 | Further customization

The `ggplot2` package in R, based on the “The Grammar of Graphics,” allows users to create detailed graphics and was used to create the templates found in **robvis**.^{22,23} As a result, both **robvis** functions return a `ggplot` object, meaning they can be easily customized further using the `ggplot2` framework. For example, to add a title to the summary bar plot:

```
library(ggplot2)

plot <- rob_summary(data_rob2,
                     tool = "ROB2")

plot +
  ggtitle("Summary Bar Plot")
```

For a full discussion of the range of post-production modifications that can be made to the plots via the `ggplot2` package, we refer the reader to the extensive guidance available.²³

3.2 | Shiny web app

Although **robvis** was originally designed for use in the command-line-based R programming environment, we developed a web app to make the tool accessible to those without knowledge of R. This is available via www.riskofbias.info.²⁴ The app was built using **Shiny**,¹³ an R package which makes it easy to produce interactive web-apps, and provides a graphical user interface (GUI) for the **robvis** package, allowing users to interact with the functions presented above without the need to download R or type any commands. Users can upload their data as either an Excel spreadsheet (recommended) or a comma-separate values (CSV) file, or can manually enter it directly into the app. Uploaded data are passed through a number of quality control checks that ensure the app will work correctly, and users are prevented from producing the plots until any issues identified have been addressed. Finally, users can customize their plots by defining a color scheme and other parameters, all through the online app.

4 | DISCUSSION

robvis facilitates the rapid production of two common risk-of-bias assessment figures at publication quality. By implementing **robvis** both as an R package and a **Shiny** web app, its functionality is available to evidence synthesists with varying levels of ability in R. **robvis** serves as an example of the advantages of “packaging” the R scripts that evidence synthesists often create for personal use.²⁵ It is likely that several other evidence synthesists have written scripts to produce similar risk-of-bias plots to those presented here—in fact, we personally know of at least one other research group that has done so. This duplication of time and effort is inefficient, and creating and sharing well-documented R packages represents one way to reduce this inefficiency. Taking this approach one step further, **Shiny** apps represent a straightforward way to provide a user-friendly GUI for a

newly created R package within a very short time-frame, expanding the potential pool of users of the package to anyone with an internet connection.

Creating a package using R has a number of particular advantages. R provides access to a range of powerful tools including the **ggplot** infrastructure as demonstrated above, and RMarkdown, which enables creation of documents that can be rendered in a range of formats such as PDF, HTML, or Word.²⁶ Furthermore, and focusing specifically on evidence synthesis, building new tools as packages in R allows for easy integration with the range of existing evidence synthesis packages. Recently, the **metaverse** project,²⁷ of which **robvis** is a part, has begun to curate a collection of R packages that cover different aspects of the systematic review and meta-analysis process which, when taken together, form a coherent end-to-end open-source alternative to commercial offerings such as Covidence or Review Manager. Key offerings in this suite of packages include **litsearcher**, which facilitates systematic search strategy development, **revtools**, a package for managing the review process and performing title and abstract screening, **metaDigitise**, a package for automatic extraction of data from figures in research papers, and **metafor**, a package for conducting meta-analyses in R.²⁸⁻³¹

While **robvis** is a stable package, a range of additional functionality could be added. At present, the number of tools with a specific template included in **robvis** is limited—adding additional templates is a priority. For example, a template for ROBIS, the risk-of-bias assessment tool for systematic reviews,⁶ is in development. Additionally, the **robvis** tool does not yet allow for the production of paired forest plots, where the risk-of-bias judgement is presented alongside each specific result included in the meta-analysis.² This was initially considered to be beyond the scope of the tool, as it involves the visualization of something other than risk-of-bias assessments. However, following user-driven demand, this functionality is in development and will be available in the near future. Finally, we would like to add similar functionality to that provided by the **metafor::reporter()** function, which generates a brief paragraph of text describing the results of a meta-analysis. The future **robvis::reporter()** function would provide a boilerplate description of the assessment tool used and the key domains at risk of bias.

ACKNOWLEDGMENTS

The following people provided valuable feedback that contributed to the development of the **robvis** tool: Emily Kothe, Eliza Grames, Matthew Page, Alexandra Bannach-Brown, Kyle Hamilton, Charles Gray, Vincent

Cheng, Wouter van Amsterdamm, Neal Haddaway, and Martin Westgate.

LAM is supported by an NIHR Doctoral Research Fellowship (DRF-2018-11-ST2-048). Attendance at the Evidence Synthesis Hackathon in April 2019 was supported by a grant from the Evidence Synthesis Hackathon itself and the University of Bristol Alumni Foundation. JPTH is a National Institute for Health Research (NIHR) Senior Investigator (NF-SI-0617-10145). JPTH is supported by NIHR Bristol Biomedical Research Centre at University Hospitals Bristol NHS Foundation Trust and the University of Bristol. JPTH is supported by the NIHR Applied Research Collaboration West (ARC West) at University Hospitals Bristol NHS Foundation Trust. JPTH is a member of the MRC Integrative Epidemiology Unit at the University of Bristol. The views expressed in this article are those of the authors and do not necessarily represent those of the NHS, the NIHR, MRC, or the Department of Health and Social Care.

CONFLICT OF INTEREST

The author reported no conflict of interest.

DATA AVAILABILITY STATEMENT

The software and data presented in this paper are freely available on GitHub: <https://github.com/mcguinlu/robvis>

ORCID

Luke A. McGuinness  <https://orcid.org/0000-0001-8730-9761>

REFERENCES

1. Donnelly CA, Boyd I, Campbell P, et al. Four principles to make evidence synthesis more useful for policy. *Nature*. 2018; 558(7710):361-364. <https://doi.org/10.1038/d41586-018-05414-4>.
2. Cochrane Handbook for Systematic Reviews of Interventions Version 6.0 (Updated August 2019)Boutron I, Higgins JP, Altman DG, et al. Chapter 7: considering bias and conflicts of interest among the included studies. In: JPT H, Chandler J, Cumpston M, et al., eds. Cochrane; 2019. Retrieved from. www.training.cochrane.org/handbook.
3. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:l4898. <https://doi.org/10.1136/bmj.l4898>.
4. Sterne JAC, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. <https://doi.org/10.1136/bmj.i4919>.
5. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*. 2011;155(8):529-536. <https://doi.org/10.7326/0003-4819-155-8-201110180-00009>.
6. Whiting P, Savović J, Higgins JPT, et al. ROBIS: a new tool to assess risk of bias in systematic reviews was developed. *J Clin*

- Epidemiol.* 2016;69:225-234. <https://doi.org/10.1016/j.jclinepi.2015.06.005>.
7. Higgins JPT, Altman DG. Chapter 8: Assessing risk of bias in included studies. In: JPT H, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Hoboken, New Jersey, United States: Wiley Online Library; 2008:187-241. <https://doi.org/10.1002/9780470712184.ch8>.
 8. Cochrane Collaboration. Review Manager (RevMan) [Computer Program]. 2014. <https://community.cochrane.org/help/tools-and-software/revman-5>.
 9. Marshall C, Brereton P. Systematic review toolbox: a catalogue of tools to support systematic reviews. Paper presented at: Proceedings of the 19th International Conference on Evaluation and Assessment in Software Engineering. EASE'15. Association for Computing Machinery; 2015: Article 23; 1–6. <https://doi.org/10.1145/2745802.2745824>.
 10. Harrison H, Griffin SJ, Kuhn I, Usher-Smith JA. Software tools to support title and abstract screening for systematic reviews in healthcare: an evaluation. *BMC Med Res Methodol.* 2020;20(1):7. <https://doi.org/10.1186/s12874-020-0897-3>.
 11. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2019. <https://www.R-project.org/>.
 12. RStudio Team. *RStudio: Integrated Development Environment for R*. Boston, MA: RStudio, Inc.; 2015.
 13. Chang W, Cheng J, Allaire J, Xie Y, McPherson J. Shiny: *Web Application Framework for R*. Boston, MA: RStudio; 2019. Retrieved from <https://CRAN.R-project.org/package=shiny>.
 14. McGuinness LA. Robvis: An R package for visualising risk-of-bias assessments. [Computer Software - v0.3.0]. November 2019. <https://doi.org/10.5281/zenodo.3552342>.
 15. Harrer M, Cuijpers P, Ebert D. *Doing Meta-Analysis in R: A Hands-on Guide*. Erlangen, Germany: Zenodo; 2019.
 16. Gibb K, Seeley A, Quinn T, et al. The consistent burden in published estimates of delirium occurrence in medical inpatients over four decades: a systematic review and meta-analysis study. *medRxiv*. 2019; 19005165. <https://doi.org/10.1101/19005165>.
 17. Habadi MI, Asiri AA, Aloufi SA, et al. Prevalence of panic disorders in the primary health care setting: a systematic review and meta-analysis. *EC Micro*. 2019;16(1):1-9.
 18. Veloso A, Vicente SG, Filipe MG. Effectiveness of cognitive training for school-aged children and adolescents with attention deficit/hyperactivity disorder: a systematic review. *Front Psych*. 2020;10:2983. <https://doi.org/10.3389/fpsyg.2019.02983>.
 19. Simillis C, Singh HKSI, Afentiotou T, et al. Postoperative chemotherapy improves survival in patients with resected high-risk stage II colorectal cancer: results of a systematic review and meta-analysis. *Colorectal Dis*. 2020; Published online 30th January. <https://doi.org/10.1111/codi.14994>.
 20. Tanneru K, Gautam S, Norez D, Kumar J., Alam M.U., Koocheckpour S., Balaji K.C., Joseph C. Meta-analysis and systematic review of intermediate-term follow-up of prostatic urethral lift for benign prostatic hyperplasia. *Int Urol Nephrol* February 2020. <https://doi.org/10.1007/s11255-020-02408-y>. Retrieved on 24 February 2020.
 21. Higgins JPT, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. <https://doi.org/10.1136/bmj.d5928>.
 22. Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. New York, NY: Springer-Verlag; 2016.
 23. Wilkinson L. *The Grammar of Graphics*. 2nd ed. New York, NY: Springer-Verlag; 2012.
 24. McGuinness L. Robvis web application. 2019. Retrieved on 20 January 2020 from <https://www.riskofbias.info/welcome/robvis-visualization-tool>
 25. Wickham H. R Packages: Organize, Test, Document, and Share Your Code. First. O'Reilly Media, Inc.; 2015. Retrieved on 20 February 2020 from <http://r-pkgs.had.co.nz/>
 26. Xie Y, Allaire JJ, Grolemund G. R Markdown: the Definitive Guide. Chapman and Hall/CRC; 2018. Retrieved on 1 February 2020 from <https://bookdown.org/yihui/rmarkdown/>
 27. Various Authors. Metaverse: an R Ecosystem for Meta-Research. 2020. Retrieved on 24 February 2020 from <https://rmetaverse.github.io/>
 28. Grames EM, Stillman AN, Tingley MW, Elphick CS. An automated approach to identifying search terms for systematic reviews using keyword co-occurrence networks. *Methods Ecol Evol*. 2019; 10:1645-1654. <https://doi.org/10.1111/2041-210X.13268>.
 29. Viechtbauer W. Conducting meta-analyses in R with the metanfor package. *J Stat Software*. 2010;36(3):1-48. <https://doi.org/10.18637/jss.v036.i03>.
 30. Pick JL, Nakagawa S, Noble DWA. Reproducible, flexible and high-throughput data extraction from primary literature: the metaDigitise R package. *Methods Ecol Evol*. 2018;10:426-431. <https://doi.org/10.1101/247775>.
 31. Westgate MJ. Revtools: an R package to support article screening for evidence synthesis. *Res Syn Meth*. 2019;10:606-614. <https://doi.org/10.1002/jrsm.1374>.

How to cite this article: McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Res Syn Meth*. 2021;12: 55–61. <https://doi.org/10.1002/jrsm.1411>

RESEARCH ARTICLE

A descriptive analysis of the data availability statements accompanying medRxiv preprints and a comparison with their published counterparts

Luke A. McGuinness^{1,2*}, Athena L. Sheppard^{1,3}

1 Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom, **2** MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, United Kingdom, **3** Department of Health Sciences, University of Leicester, Leicester, United Kingdom

* luke.mcguinness@bristol.ac.uk



Abstract

Objective

To determine whether medRxiv data availability statements describe open or closed data—that is, whether the data used in the study is openly available without restriction—and to examine if this changes on publication based on journal data-sharing policy. Additionally, to examine whether data availability statements are sufficient to capture code availability declarations.

Design

Observational study, following a pre-registered protocol, of preprints posted on the medRxiv repository between 25th June 2019 and 1st May 2020 and their published counterparts.

Main outcome measures

Distribution of preprinted data availability statements across nine categories, determined by a prespecified classification system. Change in the percentage of data availability statements describing open data between the preprinted and published versions of the same record, stratified by journal sharing policy. Number of code availability declarations reported in the full-text preprint which were not captured in the corresponding data availability statement.

Results

3938 medRxiv preprints with an applicable data availability statement were included in our sample, of which 911 (23.1%) were categorized as describing open data. 379 (9.6%) preprints were subsequently published, and of these published articles, only 155 contained an applicable data availability statement. Similar to the preprint stage, a minority (59 (38.1%)) of these published data availability statements described open data. Of the 151 records eligible for the comparison between preprinted and published stages, 57 (37.7%) were

OPEN ACCESS

Citation: McGuinness LA, Sheppard AL (2021) A descriptive analysis of the data availability statements accompanying medRxiv preprints and a comparison with their published counterparts. PLoS ONE 16(5): e0250887. <https://doi.org/10.1371/journal.pone.0250887>

Editor: Rafael Sarkis-Onofre, Meridional Faculty IMED, BRAZIL

Received: September 21, 2020

Accepted: April 15, 2021

Published: May 13, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0250887>

Copyright: © 2021 McGuinness, Sheppard. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All materials (data, code and [supporting information](#)) are available from GitHub (<https://github.com/mcguinlu/data->

[availability-impact](#)), archived at time of submission on Zenodo (DOI: [10.5281/zenodo.3968301](https://doi.org/10.5281/zenodo.3968301)).

Funding: LAM is supported by an National Institute for Health Research (NIHR; <https://www.nihr.ac.uk/>) Doctoral Research Fellowship (DRF-2018-11-ST2-048). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The views expressed in this article are those of the authors and do not necessarily represent those of the NHS, the NIHR, MRC, or the Department of Health and Social Care.

Competing interests: The authors have declared that no competing interests exist.

published in journals which mandated open data sharing. Data availability statements more frequently described open data on publication when the journal mandated data sharing (open at preprint: 33.3%, open at publication: 61.4%) compared to when the journal did not mandate data sharing (open at preprint: 20.2%, open at publication: 22.3%).

Conclusion

Requiring that authors submit a data availability statement is a good first step, but is insufficient to ensure data availability. Strict editorial policies that mandate data sharing (where appropriate) as a condition of publication appear to be effective in making research data available. We would strongly encourage all journal editors to examine whether their data availability policies are sufficiently stringent and consistently enforced.

1 Introduction

The sharing of data generated by a study is becoming an increasingly important aspect of scientific research [1, 2]. Without access to the data, it is harder for other researchers to examine, verify and build on the results of that study [3]. As a result, many journals now mandate data availability statements. These are dedicated sections of research articles, which are intended to provide readers with important information about whether the data described by the study are available and if so, where they can be obtained [4].

While requiring data availability statements is an admirable first step for journals to take, and as such is viewed favorably by journal evaluation rubrics such as the Transparency and Openness Promotion [TOP] Guidelines [5], a lack of review of the contents of these statements often leads to issues. Many authors claim that their data can be made “available on request”, despite previous work establishing that these statements are demonstrably untrue in the majority of cases—that when data is requested, it is not actually made available [6–8]. Additionally, previous work found that the availability of data “available on request” declines with article age, indicating that this approach is not a valid long term option for data sharing [9]. This suggests that requiring data availability statements without a corresponding editorial or peer review of their contents, in line with a strictly enforced data-sharing policy, does not achieve the intended aim of making research data more openly available. However, few journals actually mandate data sharing as a condition of publication. Of a sample of 318 biomedical journals, only ~20% had a data-sharing policy that mandated data sharing [10].

Several previous studies have examined the data availability statements of published articles [4, 11–13], but to date, none have examined the statements accompanying preprinted manuscripts, including those hosted on medRxiv, the preprint repository for manuscripts in the medical, clinical, and related health sciences [14]. Given that preprints, particularly those on medRxiv, have impacted the academic discourse around the recent (and ongoing) COVID-19 pandemic to a similar, if not greater, extent than published manuscripts [15], assessing whether these studies make their underlying data available without restriction (i.e. “open”), and adequately describe how to access it in their data availability statements, is worthwhile. In addition, by comparing the preprint and published versions of the data availability statements for the same paper, the potential impact of different journal data-sharing policies on data availability can be examined. This study aimed to explore the distribution of data availability statements’ description of the underlying data across a number of categories of “openness” and to

assess the change between preprint and journal-published data availability statements, stratified by journal data-sharing policy. We also intended to examine whether authors planning to make the data available upon publication actually do so, and whether data availability statements are sufficient to capture code availability declarations.

2 Methods

2.1 Protocol and ethics

A protocol for this analysis was registered in advance and followed at all stages of the study [16]. Any deviations from the protocol are described. Ethical approval was not required for this study.

2.2 Data extraction

The data availability statements of preprints posted on the medRxiv preprint repository between 25th June 2019 (the date of first publication of a preprint on medRxiv) and 1st May 2020 were extracted using the medrxivr and rvest R packages [17, 18]. Completing a data availability statement is required as part of the medRxiv submission process, and so a statement was available for all eligible preprints. Information on the journal in which preprints were subsequently published was extracted using the published DOI provided by medRxiv and rcross-ref [19]. Several other R packages were used for data cleaning and analysis [20–33].

To extract the data availability statements for published articles and the journals data-sharing policies, we browsed to the article or publication website and manually copied the relevant material (where available) into an Excel file. The extracted data are available for inspection (see Material availability section).

2.3 Categorization

A pre-specified classification system was developed to categorize each data availability statement as describing either open or closed data, with additional ordered sub-categories indicating the degree of openness (see Table 1). The system was based on the “Findability” and “Accessibility” elements of the FAIR framework [34], the categories used by previous effort to categorize published data availability statements [4, 11], our own experience of medRxiv data availability statements, and discussion with colleagues. Illustrative examples of each category were taken from preprints included in our sample [35–43].

The data availability statement for each preprinted record were categorized by two independent researchers, using the groups presented in Table 1, while the statements for published articles were categorized using all groups barring Category 3 and 4 (“Available in the future”). Records for which the data availability statement was categorized as “Not applicable” (Category 1 from Table 1) at either the preprint or published stage were excluded from further analyses. Researchers were provided only with the data availability statement, and as a result, were blind to the associated preprint metadata (e.g. title, authors, corresponding author institution) in case this could affect their assessments. Any disagreements were resolved through discussion.

Due to our large sample, if authors claimed that all data were available in the manuscript or as a S1 File, or that their study did not make use of any data, we took them at their word. Where a data availability statement met multiple categories or contained multiple data sources with varying levels of openness, we took a conservative approach and categorized it on the basis of the most restrictive aspect (see S1 File for some illustrative examples). We plotted the

Table 1. Categories used to classify the data availability statements.

Key	Main category	Sub-category	Example
0	Not applicable (protocol for a review, commentary, etc)		"Data sharing not applicable to this article as no datasets were generated or analysed during the current study." [35]
1	"Closed"	Data not made available	"Not available for public" [36]
2	"Closed"	Data available on request to authors	"Data can be available upon reasonable request to the corresponding author." [37]
3	"Closed"	Data will be made available in the future (link provided)	"The protocol and full dataset will be available at Open Science Framework upon peer review publication (https://osf.io/rvbuyl/)." [38]
4	"Closed"	Data will be made available in the future (no link provided)	"Data will be deposited in Dryad upon publication" [39]
5	"Closed"	Data available from central repository (access-controlled or open access), but insufficient detail available to find specific dataset	"Data were obtained from the international MSBase cohort study. Information regarding data availability can be obtained at https://www.msbase.org/ ." OR Daily diagnosis number of countries outside China is download from WHO situation reports (https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports). https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports [40]
6	"Closed"	Data available from central access-controlled repository, and sufficient details included to identify specific dataset e.g. via extract or accession ID or date stamp	"This research has been conducted using the UK Biobank Resource under application number 24494. All bona fide researchers can apply to use the UK Biobank resource for health related research that is in the public interest." [41]
7	"Open"	Data available in the manuscript/ S1 File	"All data related to this study are present in the paper or the S1 File ." [42]
8	"Open"	Data available via a online repository that is not access-controlled e.g. Dryad, Zenodo	"Extracted data used in this meta-analysis and analysis code are available at www.doi.org/10.5281/zenodo.3149365 ." [43]

Illustrative examples of each category were taken from preprints included in our sample (see "Data extraction").

<https://doi.org/10.1371/journal.pone.0250887.t001>

distribution of preprint and published data availability statements across the nine categories presented in [Table 1](#).

Similarly, the extracted data-sharing policies were classified by two independent reviewers according to whether the journal mandated data sharing (1) or not (0). Where the journal had no obvious data sharing policy, these were classified as not mandating data sharing.

2.4 Changes between preprinted and published statements

To assess if data availability statements change between preprint and published articles, we examined whether a discrepancy existed between the categories assigned to the preprinted and published statements, and the direction of the discrepancy ("more closed" or "more open"). Records were deemed to become "more open" if their data availability statement was categorized as "closed" at the preprint stage and "open" at the published stage. Conversely, records described as "more closed" were those moving from "open" at preprint to "closed" on publication.

We declare a minor deviation from our protocol for this analysis [16]. Rather than investigating the data-sharing policy only for journals with the largest change in openness as intended, which involved setting an arbitrary cut-off when defining "largest change", we systematically extracted and categorized the data-sharing policies for all journals in which preprints had subsequently been published using two categories (1: "requiring/mandating data sharing" and, 2: "not requiring/mandating data sharing"), and compared the change in openness between these two categories. Note that Category 2 includes journals that encourage data sharing, but do not make it a condition of publication.

To assess claims that data will be provided on publication, the data availability statements accompanying the published articles for all records in Category 3 (“Data available on publication (link provided)”) or Category 4 (“Data available on publication (no link provided)”) from [Table 1](#) were assessed, and any difference between the two categories examined.

2.5 Code availability

Finally, to assess whether data availability statements also capture the availability of programming code, such as STATA do files or R scripts, the data availability statement and full text PDF for a random sample of 400 preprinted records were assessed for code availability (1: “code availability described” and 2: “code availability not described”).

3 Results

The data availability statements accompanying 4101 preprints registered between 25th June 2019 and 1st May 2020 were extracted from the medRxiv preprint repository on the 26th May 2020 and were coded by two independent researchers according to the categories in [Table 1](#). During this process, agreement between the raters was high (Cohen’s Kappa = 0.98; “almost perfect agreement”) [44].

Of the 4101 preprints, 163 (4.0%) in Category 0 (“Not applicable”) were excluded following coding, leaving 3938 remaining records. Of these, 911 (23.1%) had made their data open as per the criteria in [Table 1](#). The distribution of data availability statements across the categories can be seen in [Fig 1](#). A total of 379 (9.6%) preprints had been subsequently published, and of these, only 159 (42.0%) had data availability statements that we could categorize. 4 (2.5%) records in




Fig 1. Distribution of the data availability statements of preprinted (Panel A) and published (Panel B) records by category from [Table 1](#).

<https://doi.org/10.1371/journal.pone.0250887.g001>

Table 2. Change in openness of data availability statements from preprint to published article, grouped by journal data-sharing policy.

Journal data sharing policy	Preprinted records subsequently published (N)	Open DAS in preprinted version % (N)	Open DAS in published version % (N)	Change in DAS from preprint to publication		
				More open (N)	More closed (N)	No change (N)
Does not mandate open data	94	20.2% (19)	22.3% (21)	10	8	76
Mandates open data	57	33.3% (19)	61.4% (35)	16	0	41

<https://doi.org/10.1371/journal.pone.0250887.t002>

Category 0 (“Not applicable”) were excluded, and of the 155 remaining, 59 (38.1%) had made their data open as per our criteria.

For the comparison of preprinted data availability statements with their published counterparts, we excluded records that were not published, that did not have a published data availability statement or that were labeled as “Not applicable” at either the preprint or published stage, leaving 151 records (3.7% of the total sample of 4101 records) records.

Data availability statements more frequently described open data on publication compared to the preprinted record when the journal mandated data sharing (Table 2). Moreover, the data availability statements for 8 articles published in journals that did not mandate open data sharing became less open on publication. The change in openness for preprints grouped by category and stratified by journal policy is shown in S1 Table in [S1 File](#), while the change for each individual journal included in our analysis is shown in S2 Table in [S1 File](#).

Interestingly, 22 records published in a journal mandating open data sharing did not have an open data availability statement. The majority of these records described data that was available from a central access-controlled repository (Category 5 or 6), while in others, legal restrictions were cited as the reason for lack of data sharing. However, in some cases, data was either insufficiently described or was only available on request (S3 Table in [S1 File](#)), indicating that journal policies which mandate data sharing may not always be consistently applied allowing some records to slip through the gaps.

161 (4.1%) preprints stated that data would be available on publication, but only 10 of these had subsequently been published (Table 3) and the number describing open data on publication did not seem to vary based on whether the preprinted data availability statements include a link to an embargoed repository or not, though the sample size is small.

Of the 400 records for which code availability was assessed, 75 mentioned code availability in the preprinted full-text manuscript. However, only 22 (29.3%) of these also described code availability in the corresponding data availability statement (S4 Table in [S1 File](#)).

Table 3. Assessment of whether researchers promising to make data available on publication actually do so, and whether this differs if researchers included a link to an embargoed repository or not.

Preprint Category	Number of preprints	Published Category	Number of published studies
Data available in the future, with a link to an embargoed repository provided	3	1. Data not made available 5. Data available from central repository (access-controlled or open access), but insufficient detail available to find specific dataset 8. Data available via a online repository that is not access-controlled e.g. Dryad, Zenodo	1 (33.3%) 1 (33.3%) 1 (33.3%)
Data available in the future, with no details of embargoed repository given	7	1. Data not made available 2. Data available on request to authors 7. Data available in the manuscript/ S1 File 8. Data available via a online repository that is not access-controlled e.g. Dryad, Zenodo	1 (14.3%) 1 (14.3%) 1 (14.3%) 4 (57.1%)

<https://doi.org/10.1371/journal.pone.0250887.t003>

4 Discussion

4.1 Principal findings and comparison with other studies

We have reviewed 4101 preprinted and 159 published data availability statements, coding them as “open” or “closed” according to a predefined classification system. During this labor-intensive process, we appreciated statements that reflected the authors’ enthusiasm for data sharing (“YES”) [45], their bluntness (“Data is not available on request.”) [46], and their efforts to endear themselves to the reader (“I promise all data referred to in the manuscript are available.”) [47]. Of the preprinted statements, almost three-quarters were categorized as “closed”, with the largest individual category being “available on request”. In light of the substantial impact that studies published as preprints on medRxiv have had on real-time decision making during the current COVID-19 pandemic [15], it is concerning that data for these preprints is so infrequently readily available for inspection.

A minority of published records we examined contained a data availability statement ($n = 159$ (42.0%)). This lack of availability statement at publication results in a loss of useful information. For at least one published article, we identified relevant information in the pre-printed statement that did not appear anywhere in the published article, due to it not containing a data availability statement [48, 49].

We provide initial descriptive evidence that strict data-sharing policies, which mandate that data be made openly available (where appropriate) as a condition of publication, appear to succeed in making research data more open than those that do not. Our findings, though based on a relatively small number of observations, agree with other studies on the effect of journal policies on author behavior. Recent work has shown that “requiring” a data availability statement was effective in ensuring that this element was completed [4], while “encouraging” authors to follow a reporting checklist (the ARRIVE checklist) had no effect on compliance [50, 51].

Finally, we also provide evidence that data availability statements alone are insufficient to capture code availability declarations. Even when researchers wish to share their code, as evidenced by a description of code availability in the main paper, they frequently do not include this information in the data availability statement. Code sharing has been advocated strongly elsewhere [52–54], as it provides an insight into the analytic decisions made by the research team, and there are few, if any, circumstances in which it is not possible to share the analytic code underpinning an analysis. Similar to data availability statements, a dedicated code availability statement which is critically assessed against a clear code-sharing policy as part of the editorial and peer review processes will help researchers to appraise published results.

4.2 Strengths and limitations

A particular strength of this analysis is that the design allows us to compare what is essentially the same paper (same design, findings and authorship team) under two different data-sharing policies, and assess the change in the openness of the statement between them. To our knowledge this is the first study to use this approach to examine the potential impact of journal editorial policies. This approach also allows us to address the issue of self-selection. When looking at published articles alone, it is not possible to tell whether authors always intended to make their data available and chose a given journal due to its reputation for data sharing. In addition, we have examined all available preprints within our study period and all corresponding published articles, rather than taking a sub-sample. Finally, categorization of the statements was carried out by two independent researchers using predefined categories, reducing the risk of misclassification.

However, our analysis is subject to a number of potential limitations. The primary one is that manuscripts (at both the preprint and published stages) may have included links to the data, or more information that uniquely identifies the dataset from a data portal, within the text (for example, in the Methods section). While this might be the case, if readers are expected to piece together the relevant information from different locations in the manuscript, it throws into question what having a dedicated data availability statement adds. A second limitation is that we do not assess the veracity of any data availability statements, which may introduce some misclassification bias into our categorization. For example, we do not check whether all relevant data can actually be found in the manuscript/S1 File (Category 7) or the linked repository (Category 8), meaning our results provide a conservative estimate of the scale of the issue, as previous work has suggested that this is unlikely to be the case [12]. A further consideration is that for Categories 1 (“No data available”) and 2 (“Available on request”), there will be situations where making research data available is not feasible, for example, due to cost or concerns about patient re-identifiability [55, 56]. This situation is perfectly reasonable, as long as statements are explicit in justifying the lack of open data.

4.3 Implications for policy

Data availability statements are an important tool in the fight to make studies more reproducible. However, without critical review of these statements in line with strict data-sharing policies, authors default to not sharing their data or making it “available on request”. Based on our analysis, there is a greater change towards describing open data between preprinted and published data availability statements in journals that mandate data sharing as a condition of publication. This would suggest that data sharing could be immediately improved by journals becoming more stringent in their data availability policies. Similarly, introduction of a related code availability section (or composite “material” availability section) will aid in reproducibility by capturing whether analytic code is available in a standardized manuscript section.

It would be unfair to expect all editors and reviewers to be able to effectively review the code and data provided with a submission. As proposed elsewhere [57], a possible solution is to assign an editor or reviewer whose sole responsibility in the review process is to examine the data and code provided. They would also be responsible for judging, when data and code are absent, whether the argument presented by the authors for not sharing these materials is valid.

However, while this study focuses primarily on the role of journals, some responsibility for enacting change rests with the research community at large. If researchers regularly shared our data, strict journal data-sharing policies would not be needed. As such, we would encourage authors to consider sharing the data underlying future publications, regardless of whether the journal actually mandates it.

5 Conclusion

Requiring that authors submit a data availability statement is a good first step, but is insufficient to ensure data availability, as our work shows that authors most commonly use them to state that data is only available on request. However, strict editorial policies that mandate data sharing (where appropriate) as a condition of publication appear to be effective in making research data available. In addition to the introduction of a dedicated code availability statement, a move towards mandated data sharing will help to ensure that future research is readily reproducible. We would strongly encourage all journal editors to examine whether their data availability policies are sufficiently stringent and consistently enforced.

Supporting information

S1 File.

(DOCX)

Acknowledgments

We must acknowledge the input of several people, without whom the quality of this work would have been diminished: Matthew Grainger and Neal Haddaway for their insightful comments on the subject of data availability statements; Phil Gooch and Sarah Nevitt for their skill in identifying missing published papers based on the vaguest of descriptions; Antica Culina, Julian Higgins and Alfredo Sánchez-Tójar for their comments on the preprinted version of this article; and Ciara Gardiner, for proof-reading this manuscript.

Author Contributions

Conceptualization: Luke A. McGuinness.

Data curation: Luke A. McGuinness.

Formal analysis: Luke A. McGuinness, Athena L. Sheppard.

Investigation: Luke A. McGuinness, Athena L. Sheppard.

Methodology: Luke A. McGuinness, Athena L. Sheppard.

Project administration: Luke A. McGuinness.

Software: Luke A. McGuinness.

Supervision: Luke A. McGuinness.

Validation: Luke A. McGuinness, Athena L. Sheppard.

Visualization: Luke A. McGuinness.

Writing – original draft: Luke A. McGuinness.

Writing – review & editing: Luke A. McGuinness, Athena L. Sheppard.

References

1. Packer M. Data sharing in medical research. *BMJ* 2018; k510. <https://doi.org/10.1136/bmj.k510> PMID: 29444885
2. Taichman DB, Backus J, Baethge C, et al. Sharing clinical trial data. *BMJ* 2016; i255. <https://doi.org/10.1136/bmj.i255> PMID: 26790902
3. Krumholz HM. Why data sharing should be the expected norm. *BMJ (Clinical research ed)* 2015; 350: h599. <https://doi.org/10.1136/bmj.h599> PMID: 25655137
4. Federer LM, Belter CW, Joubert DJ, et al. Data sharing in PLOS ONE: An analysis of Data Availability Statements. *PLOS ONE* 2018; 13:e0194768. <https://doi.org/10.1371/journal.pone.0194768> PMID: 29719004
5. Nosek BA, Alter G, Banks GC, et al. Promoting an open research culture. *Science* 2015; 348:1422–5. <https://doi.org/10.1126/science.aab2374> PMID: 26113702
6. Naudet F, Sakarovitch C, Janiaud P, et al. Data sharing and reanalysis of randomized controlled trials in leading biomedical journals with a full data sharing policy: Survey of studies published in *The BMJ* and *PLOS Medicine*. *BMJ* 2018; k400. <https://doi.org/10.1136/bmj.k400> PMID: 29440066
7. Miyakawa T. No raw data, no science: Another possible source of the reproducibility crisis. *Molecular Brain* 2020; 13:24. <https://doi.org/10.1186/s13041-020-0552-2> PMID: 32079532

8. Krawczyk M, Reuben E. (Un)Available upon Request: Field Experiment on Researchers' Willingness to Share Supplementary Materials. *Accountability in Research* 2012; 19:175–86. <https://doi.org/10.1080/08989621.2012.678688> PMID: 22686633
9. Vines TH, Albert AYK, Andrew RL, et al. The Availability of Research Data Declines Rapidly with Article Age. *Current Biology* 2014; 24:94–7. <https://doi.org/10.1016/j.cub.2013.11.014> PMID: 24361065
10. Vasilevsky NA, Minnier J, Haendel MA, et al. Reproducible and reusable research: Are journal data sharing policies meeting the mark? *PeerJ* 2017; 5. <https://doi.org/10.7717/peerj.3208> PMID: 28462024
11. Colavizza G, Hrynaszkiewicz I, Staden I, et al. The citation advantage of linking publications to research data. *PLOS ONE* 2020; 15:e0230416. <https://doi.org/10.1371/journal.pone.0230416> PMID: 32320428
12. Roche DG, Kruuk LEB, Lanfear R, et al. Public Data Archiving in Ecology and Evolution: How Well Are We Doing? *PLOS Biology* 2015; 13:e1002295. <https://doi.org/10.1371/journal.pbio.1002295> PMID: 26556502
13. Tan SC, Flanagan D, Morris E, et al. Research data repositories chosen by researchers across broad range of disciplines, from an analysis of 145,000 data availability statements. *Authorea Published Online First*: July 2020. <https://doi.org/10.22541/au.159422974.49069472>
14. Rawlinson C, Bloom T. New preprint server for medical research. *BMJ* 2019; 365. <https://doi.org/10.1136/bmj.l2301> PMID: 31167753
15. Fraser N, Brierley L, Dey G, et al. Preprinting a pandemic: The role of preprints in the COVID-19 pandemic. *bioRxiv* 2020;2020.05.22.111294. <https://doi.org/10.1101/2020.05.22.111294>
16. McGuinness LA, Sheppard AL. Protocol for a descriptive analysis of the data availability statements accompanying medRxiv preprints. 2020.
17. McGuinness L, Schmidt L. Medrxivr: Accessing and searching medRxiv and bioRxiv preprint data in R. *Journal of Open Source Software* 2020; 5:2651. <https://doi.org/10.21105/joss.02651>
18. Wickham H. *Rvest: Easily harvest (scrape) web pages*. 2019. <https://CRAN.R-project.org/package=rvest>
19. Chamberlain S, Zhu H, Jahn N, et al. *Rcrossref: Client for various 'CrossRef' APIs*. 2020. <https://CRAN.R-project.org/package=rcrossref>
20. R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria:: R Foundation for Statistical Computing 2019. <https://www.R-project.org/>
21. Wickham H, Hester J, Chang W. *Devtools: Tools to make developing r packages easier*. 2019. <https://CRAN.R-project.org/package=devtools>
22. Wickham H, François R, Henry L, et al. *Dplyr: A grammar of data manipulation*. 2019. <https://CRAN.R-project.org/package=dplyr>
23. Gohel D. *Flextable: Functions for tabular reporting*. 2020. <https://CRAN.R-project.org/package=flextable>
24. Wickham H. *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York 2016. <https://ggplot2.tidyverse.org>
25. Rodriguez-Sanchez F. *Grateful: Facilitate citation of r packages*. 2018. <https://github.com/Pakillo/grateful>
26. Müller K. *Here: A simpler way to find your files*. 2017. <https://CRAN.R-project.org/package=here>
27. Gamer M, Lemon J, Fellows I, et al. *Irr: Various coefficients of interrater reliability and agreement*. 2019. <https://CRAN.R-project.org/package=irr>
28. Gohel D. *Officer: Manipulation of microsoft word and PowerPoint documents*. 2020. <https://CRAN.R-project.org/package=officer>
29. Pedersen TL. *Patchwork: The composer of plots*. 2019. <https://CRAN.R-project.org/package=patchwork>
30. Neuwirth E. *RColorBrewer: ColorBrewer palettes*. 2014. <https://CRAN.R-project.org/package=RColorBrewer>
31. Chan C, Chan GC, Leeper TJ, et al. *Rio: A swiss-army knife for data file i/o*. 2018.
32. Wickham H. *Stringr: Simple, consistent wrappers for common string operations*. 2019. <https://CRAN.R-project.org/package=stringr>
33. Müller K, Wickham H. *Tibble: Simple data frames*. 2019. <https://CRAN.R-project.org/package=tibble>
34. Wilkinson MD, Dumontier M, Aalbersberg IJ, et al. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* 2016; 3:160018. <https://doi.org/10.1038/sdata.2016.18> PMID: 26978244

35. Ehrlich OG, Testaverde J, Heller C, et al. Crohns disease and ulcerative colitis patient perspectives on clinical trials and participation. *medRxiv* 2019;19000273. <https://doi.org/10.1101/19000273>
36. Septiandi AA, Aditiawarman A, Tjiong R, et al. Cost-Sensitive Machine Learning Classification for Mass Tuberculosis Screening. *medRxiv* 2019;19000190. <https://doi.org/10.1101/19000190>
37. Solis JCA, Storvoll I, Vanbelle S, et al. Impact of spectrograms on the classification of wheezes and crackles in an educational setting. An interrater study. *medRxiv* 2019;19005504. <https://doi.org/10.1101/19005504>
38. Ebbeling CB, Bielak L, Lakin PR, et al. Higher energy requirement during weight-loss maintenance on a low- versus high-carbohydrate diet: Secondary analyses from a randomized controlled feeding study. *medRxiv* Published Online First: July 2019. <https://doi.org/10.1101/19001248>
39. Barry A, Bradley J, Stone W, et al. Increased gametocyte production and mosquito infectivity in chronic versus incident Plasmodium falciparum infections. *medRxiv* 2020;2020.04.08.20057927. <https://doi.org/10.1101/2020.04.08.20057927>
40. Malpas CB, Ali Manouchehrinia A, Sharmin S, et al. Early clinical markers of aggressive multiple sclerosis. *medRxiv* Published Online First: July 2019. <https://doi.org/10.1101/19002063>
41. Knuppel A, Papier K, Fensom GK, et al. Meat intake and cancer risk: Prospective analyses in UK Biobank. *medRxiv* 2019;19003822. <https://doi.org/10.1101/19003822>
42. Thompson ER, Bates L, Ibrahim IK, et al. Novel delivery of cellular therapy to reduce ischaemia reperfusion injury in kidney transplantation. *medRxiv* 2019;19005546. <https://doi.org/10.1101/19005546>
43. Moriarty F, Ebell MH. A comparison of contemporary versus older studies of aspirin for primary prevention. *medRxiv* 2019;19004267. <https://doi.org/10.1101/19004267>
44. McHugh ML. Interrater reliability: The kappa statistic. *Biochemia Medica* 2012; 22:276–82. PMID: [23092060](#)
45. Chen L, Du X, Liu Y, et al. Comparison of the Clinical Implications among Two Different Nutritional Indic平es in Hospitalized Patients with COVID-19. *medRxiv* Published Online First: May 2020. <https://doi.org/10.1101/2020.04.28.20082644>
46. Hashmi M, Taqi A, Memon MI, et al. A national landscaping survey of critical care services in hospitals accredited for training in a lower-middle income country: Pakistan. *medRxiv* Published Online First: April 2020. <https://doi.org/10.1101/2020.04.22.20071555>
47. Peng L, Liu J, Xu W, et al. 2019 Novel Coronavirus can be detected in urine, blood, anal swabs and oropharyngeal swabs samples. *medRxiv* Published Online First: February 2020. <https://doi.org/10.1101/2020.02.21.20026179>
48. Martin J, Hosking G, Wadon M, et al. A brief report: De novo copy number variants in children with attention deficit hyperactivity disorder. *medRxiv* Published Online First: December 2019. <https://doi.org/10.1101/12.12.19014555>
49. Martin J, Hosking G, Wadon M, et al. A brief report: De novo copy number variants in children with attention deficit hyperactivity disorder. *Translational Psychiatry* 2020; 10:135. <https://doi.org/10.1038/s41398-020-0821-y> PMID: [32398668](#)
50. Hair K, Macleod MR, Sena ES, et al. A randomised controlled trial of an Intervention to Improve Compliance with the ARRIVE guidelines (IICARus). *Research Integrity and Peer Review* 2019; 4:12. <https://doi.org/10.1186/s41073-019-0069-3> PMID: [31205756](#)
51. Kilkenny C, Browne WJ, Cuthill IC, et al. Improving Bioscience Research Reporting: The ARRIVE Guidelines for Reporting Animal Research. *PLOS Biology* 2010; 8:e1000412. <https://doi.org/10.1371/journal.pbio.1000412> PMID: [20613859](#)
52. Goldacre B, Morton CE, DeVito NJ. Why researchers should share their analytic code. *BMJ* 2019; I6365. <https://doi.org/10.1136/bmj.I6365> PMID: [31753846](#)
53. Eglen SJ, Marwick B, Halchenko YO, et al. Towards standard practices for sharing computer code and programs in neuroscience. *Nature neuroscience* 2017; 20:770–3. <https://doi.org/10.1038/nn.4550> PMID: [28542156](#)
54. Culina A, van den Berg I, Evans S, et al. Low availability of code in ecology: A call for urgent action. *PLOS Biology* 2020; 18:e3000763. <https://doi.org/10.1371/journal.pbio.3000763> PMID: [32722681](#)
55. Goodhill GJ. Practical costs of data sharing. *Nature* 2014; 509:33–3. <https://doi.org/10.1038/509033b> PMID: [24784208](#)
56. Courbier S, Dimond R, Bros-Facer V. Share and protect our health data: An evidence based approach to rare disease patients' perspectives on data sharing and data protection—quantitative survey and recommendations. *Orphanet Journal of Rare Diseases* 2019; 14:175. <https://doi.org/10.1186/s13023-019-1123-4> PMID: [31300010](#)
57. Sánchez-Tójar A, Lagisz M, Moran NP, et al. The jury is still out regarding the generality of adaptive “transgenerational” effects. *Ecology Letters*;n/a. <https://doi.org/10.1111/ele.13479> PMID: [32844521](#)



Ensuring Prevention Science Research is Synthesis-Ready for Immediate and Lasting Scientific Impact

Emily A. Hennessy¹ · Rebecca L. Acabchuk² · Pieter A. Arnold³ · Adam G. Dunn⁴ · Yong Zhi Foo⁵ ·
Blair T. Johnson⁶ · Sonya R. Geange^{3,7} · Neal R. Haddaway^{8,9,10} · Shinichi Nakagawa⁵ · Witness Mapanga¹¹ ·
Kerrie Mengersen¹² · Matthew J. Page¹³ · Alfredo Sánchez-Tójar¹⁴ · Vivian Welch¹⁵ · Luke A. McGuinness¹⁶

Accepted: 23 June 2021
© Society for Prevention Research 2021

Abstract

When seeking to inform and improve prevention efforts and policy, it is important to be able to robustly synthesize all available evidence. But evidence sources are often large and heterogeneous, so understanding what works, for whom, and in what contexts can only be achieved through a systematic and comprehensive synthesis of evidence. Many barriers impede comprehensive evidence synthesis, which leads to uncertainty about the generalizability of intervention effectiveness, including inaccurate titles/abstracts/keywords terminology (hampering literature search efforts), ambiguous reporting of study methods (resulting in inaccurate assessments of study rigor), and poorly reported participant characteristics, outcomes, and key variables (obstructing the calculation of an overall effect or the examination of effect modifiers). To address these issues and improve the reach of primary studies through their inclusion in evidence syntheses, we provide a set of practical guidelines to help prevention scientists prepare synthesis-ready research. We use a recent mindfulness trial as an empirical example to ground the discussion and demonstrate ways to ensure the following: (1) primary studies are discoverable; (2) the types of data needed for synthesis are present; and (3) these data are readily synthesizable. We highlight several tools and practices that can aid authors in these efforts, such as using a data-driven approach for crafting titles, abstracts, and keywords or by creating a repository for each project to host all study-related data files. We also provide step-by-step guidance and software suggestions for standardizing data design and public archiving to facilitate synthesis-ready research.

Keywords Evidence synthesis · Meta-data · Reporting · Reproducibility · Synthesis-ready research · Transparency

Systematic reviews are an important tool for informing policy and practice. When done well, they provide a robust synthesis of available evidence, account for risks of bias in individual studies, and address problems associated with the volume of primary research being published (Johnson & Hennessy, 2019; Wilson & Tanner-Smith, 2014).

To conduct a comprehensive evidence synthesis, such as a systematic review and meta-analysis, primary studies related to the research aim must be identifiable and report relevant study and outcome data. Those who conduct evidence syntheses — hereafter referred to as *evidence synthesists* — often need information about study processes and measured variables aside from a manuscripts' primary outcome of interest to explore why there are differences (i.e.,

statistical heterogeneity) between studies examining the same intervention. This information is often missing, which leads to uncertainty about the generalizability of intervention effectiveness across different groups (e.g., race/ethnicity, gender, age) and can make these efforts slower and less accurate (Borah et al., 2017; Haddaway & Westgate, 2019; Nakagawa et al., 2020; Nuijten et al., 2016). Even when studies report their statistical data and results in detail, there is often still important information missing about other factors that require further attention before they can be accurately synthesized (e.g., description of the intervention, population, or outcome measures). Data and meta-data may be missing from a primary study for many reasons, including article length restrictions, a lack of detailed reporting standards or awareness of the importance of open data, or a desire to retain control of analyses, amongst others.

Given the potential impact of their work on human health and behaviors, prevention scientists should be at the

✉ Emily A. Hennessy
ehennessy@mgh.harvard.edu

Extended author information available on the last page of the article

forefront of ensuring primary research is transparent, reproducible, and furthermore, we argue, synthesis-ready. *Synthesis-ready research* is our term to describe a study (e.g., randomized controlled trial (RCT), quasi-experimental study (QED), or observational study), its *meta-data* (i.e., descriptive information about the study's data, including protocols, tools, algorithms, so that one could replicate the study), and *data* (information collected while conducting the study) that can be readily and accurately used in evidence syntheses.

Resources are available to enable authors to share detailed data and meta-data associated with a study; yet, these resources are often not directly connected to scientific journal practices. Some of these options make it much easier to discover, use, and cite authors' research, substantially increasing the impact and legacy of their work. Thus, our aim was to draw on resources from across disciplines (Grant et al., 2018; Montgomery et al., 2018; Schulz et al., 2010; Wilkinson et al., 2016) to produce a guide to reporting primary research that is open, accessible, and synthesis-ready. To illustrate synthesis-ready research principles and streamline this process for prevention scientists, we briefly present an empirical example from a mindfulness meditation RCT to improve college student well-being ("Introduction to Empirical Example"), followed by a demonstration of the application of best practices for synthesis-ready research to this example ("Evidence Synthesis Process Applied to Our Empirical Example" and "How to Make Research Synthesis-Ready: Steps from Study Design to Completion").

From Primary Study Research to Evidence Synthesis

Primary study conduct and reporting guidelines now advocate for reporting enough information for research to be transparent and reproducible and also provide specific information on how to do so (Chan et al., 2013; Grant et al., 2018; Schulz et al., 2010). Similarly, efforts have focused on improving the degree of FAIRness of primary studies, by making research Findable, Accessible, Interoperable, and Reusable (Wilkinson et al., 2016).¹

Findable indicates research is identifiable to anyone searching for it; data and meta-data and their locations clearly described. Data and meta-data that are retrievable by their identifier (e.g., a digital object identifier, DOI) and are freely obtainable in full without barriers to access (e.g.,

without needing to register with an external website or contacting study authors) are considered *Accessible*. Data and meta-data that are understandable across humans or machines without the need for specialized tools or translation and include linkage to their relevant data/meta-data are *Interoperable*. Finally, *Reusable* refers to the degree to which information can be appropriately repurposed (e.g., integrated/built upon). Thus, research data and meta-data should have a data usage license, detailed records of data origins, and be clearly and richly described in their accompanying materials. By following Open Science recommendations and addressing FAIR principles (Frankenhuis & Nettle, 2018; Vicente-Saez & Martinez-Fuentes, 2018), prevention scientists can provide synthesis-ready research that improves the translation of research into practice.

Evidence syntheses rely on the data reported in research studies and by the researchers themselves. Similar to the conduct of primary research, there are best practices for conducting and reporting of evidence syntheses, including the following: problem formulation, systematically identifying and selecting studies for inclusion, coding studies for key features, calculating effect sizes (if synthesizing quantitative data, such as by conducting a meta-analysis), analyzing the data (assess quality and risk of bias, synthesize coded information from studies), and interpreting and communicating results (Online, Fig. 1). The content and the comprehensiveness of the review (if the review authors follow best practice guidelines) are a direct result of the available information reported in primary studies.

Potential Advantages of Synthesis-Ready Research

Synthesis-ready research can increase the transparency, integrity, and reproducibility of research, ultimately leading to improved scientific evidence (Beugelsdijk et al., 2020). Pre-registration and subsequent data sharing practices should reduce the prevalence of post hoc hypothesis (HARKing; Forstmeier et al., 2017; Kerr, 1998) and falsifying or misrepresenting data in results (Banks et al., 2019; Miyakawa, 2020). Doing so can reduce duplication of effort and introduce findings to the research community sooner and ensure that all findings (beneficial and adverse) are reported, results that can prevent costly mistakes and accelerate the impact of prevention scientists (Gupta et al., 2015; Pasquetto et al., 2019). Indeed, synthesis-ready data can be used repeatedly and for multiple purposes, ultimately informing more research questions: thus it is more cost-effective when compared to data that is only used by the team that collected it.

Primary researchers who make their work synthesis-ready should also benefit from these efforts. Synthesis-ready research is more likely to be identified and used in evidence synthesis (Christensen et al., 2019; Gerstner et al., 2017).

¹ Although FAIR principles primarily refer to enhancing the ability of machines to find and use data, these recommendations also enhance data reuse by individuals. Where appropriate, we have adapted the FAIR components to more directly apply to human-readability efforts for the purposes of this manuscript.

For example, a recent study showed that hosting data in a repository resulted in 25% more citations than merely stating data were available by request (Colavizza et al., 2020). Additionally, depending on the organization of the metadata and the data repository used, the dataset may be cited directly, as may other study materials such as the analytical code, further enhancing the profile of the work and those who conducted it (Crosas, 2013).

Because synthesis-ready research is also more identifiable, there will be increased opportunities for future collaborations (Popkin, 2019), which potentially improves future resource-sharing and reduces costs. Producing synthesis-ready research can also increase the recognition of an individual study team to an international audience. Many funding sources require data sharing of their grantees and necessitate the inclusion of such plans in grant proposals (e.g., Wellcome Trust, the UK Medical Research Council, and the US National Institutes of Health: Gaba et al., 2020; NIH Data Sharing Policy & Guidance, 2003; Wellcome Trust Public Health Research Data Forum, 2012). Thus, researchers who prepare and deliver grant-funded research findings with synthesis-ready considerations are considered responsive to funders while potentially attaining these other discussed benefits.

Introduction to Empirical Example

Our empirical example is a RCT ($N=140$ participants; registered in Clinical Trials, NCT03402009) conducted across two university semesters in the 2018 academic year (Acabchuk et al., 2021). The study evaluated two active treatment conditions to determine what tools best assist university students in developing a personal meditation practice to self-manage symptoms of depression, anxiety, and stress. One group was assigned to the *10% Happier* meditation phone app (labeled the “App Group”; Ten Percent Happier: Mindfulness Meditation Courses, 2020), which provides guided meditations, and the other was assigned to the *10% Happier* meditation phone app in combination with a Muse EEG neurofeedback device (labeled the “Muse Group”; Choose Muse, 2018). The study included a baseline assessment, randomization to one of two groups, an orientation session for both groups, collection of salivary biomarkers, and follow-up survey assessments on a variety of well-being and behavior outcomes throughout the study (Study Timeline: Online, Fig. 2).

Evidence Synthesis Process Applied to Our Empirical Example

Drawing from our empirical example, if one sought to conduct a systematic review of all types of mindfulness interventions for college students on health-related outcomes

(*Problem Formulation*), they would first need to be able to identify all primary research matching their inclusion criteria (*Finding and Selecting Studies*). Clearly delineating the research problem is a foundational step in a review, as it drives the search strategy and ability to accurately identify and synthesize primary study research. Research questions in an evidence synthesis should address all the main elements of the studies to be included; many draw on the PICOS framework with its focus on the population(s), intervention(s), comparator(s), outcome(s), and study design(s) elements of a research problem (Stern et al., 2014). Once the relevant literature is identified, data are extracted from the studies (*Code Studies for Relevant Features*) so that the analysis can commence by synthesizing coded data (*Analyze the Systematic Review Database*).

Within this broader process, the types of data and metadata that need to be reported in a primary study depend on the task and/or type of evidence synthesis. For example, research methods need to be fully reported to assess risk of bias. If the review aimed to measure the effectiveness of mindfulness interventions on depression symptoms using a meta-analysis, then primary studies should report the data necessary to calculate an effect size from their depressive symptom assessment for all participants (e.g., means, standard deviations, and sample sizes for each group). To address relevant potential effect modifiers (e.g., age, year in school, gender, intervention fidelity), any data collected during the primary study on participant characteristics and the study methods (e.g., description of groups, duration of intervention) needs to be transparently reported. To examine geographic and temporal effect modifiers, key information about the study location and timing should be provided (Johnson et al., 2017). Additionally, if the synthesis focuses on participants’ perspectives on why interventions are more or less effective, they would need access to any qualitative data collected as part of the study (e.g., interviews, focus groups). Given this variability in the type of questions that could be addressed in a synthesis, it is important for investigators to transparently and comprehensively report a variety of information about their research.

How to Make Research Synthesis-Ready: Steps from Study Design to Completion

Approaching every step of a research project with data sharing in mind is useful and often necessary to ensure that data sharing is possible. This approach has the added benefit of encouraging better overall project management, including clearer workflows and comprehensive documentation of the analytical process. We now illustrate each step of these suggestions from study design to sharing with examples from our mindfulness trial (Fig. 1).




Fig. 1 Best practices in generating synthesis-ready research applied to mindfulness trial. DAS, data availability statement

Study Design

Preparing research for sharing can be a resource-intensive process, and so planning and provisioning for this task are vital to consider when submitting funding applications and considering funders' reporting requirements. When conducting research with human participants, advance planning for data sharing is essential because it ensures appropriate documentation is prepared. Any potential ethical concerns with data sharing should be addressed during the review of study materials by the governing Institutional Review Board, which ensures that participant consent documents include the correct language to ensure participants can consent to anonymized data sharing (Ohmann et al., 2017; Thorogood & Knoppers, 2017). The Inter-university Consortium for Political and Social Research (ICPSR) recommends carefully constructing these sections and provides sample text to use to ensure that participants are informed about the potential use of their anonymized data in future research (Inter-university Consortium for Political and Social Research, 2010).

The study protocol is the foundation of the study, detailing all the processes, procedures, and materials planned for a research project. Best practice for primary studies and secondary data analysis is to register a study protocol that includes primary questions and/or hypotheses, methodology, and planned analyses. With careful attention to developing a comprehensive protocol using a standard structure, this document can increase the synthesis-readiness of the research. The information provided in a study protocol includes important meta-data that may not be available in a single published manuscript but that can be used by evidence synthesists to address a variety of research questions, including using it to assess study quality and risk of bias due to selection of reported results or "outcome switching" (Falk Delgado & Falk Delgado, 2017; Kahan & Jairath, 2018). We recommend that authors review published guidelines for the best tool for the study design in question: The EQUATOR network provides an online resource to compare these tools (<https://www.equator-network.org/>). Online Tables 1 and 2 present a mapping of critical elements for a study protocol

(using the Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT); Chan et al., 2013) and for a study *manuscript* (using the Consolidated Standards of Reporting Trials for social and psychological interventions (CONSORT); Grant et al., 2018)) and indicate how study information provided following these guidelines could be used in an evidence synthesis. Table 1

The protocol for our mindfulness trial uses the SPIRIT guidelines (Chan et al., 2013) and is archived in a Zenodo repository (DOI: <https://doi.org/10.5281/zenodo.4011716>).

Study Conduct

To facilitate data sharing and to distribute the workload across the projects' lifecycle, transparent and reproducible data management workflows should be employed throughout the research process, including (1) defining data file contents; (2) clearly describing variable names, definitions, units, and formats; (3) implementing consistent data organization and file structure; (4) performing basic quality assurance to ensure that values fall within expected ranges; and (5) providing documentation including the analytical steps taken (crosswalks), data dictionaries (i.e., codebooks), and metadata (Barton et al., 2010; Strasser, 2012; Yenni et al., 2019). Building in these meta-data collection steps into the analysis process makes it easier to create items like variable crosswalks and data dictionaries to accompany shared research.

Study Dissemination

The most labor-intensive stage of making data synthesis-ready within a FAIR perspective occurs during preparation for study dissemination as it involves several steps. To ensure their research is *Findable* and *Accessible*, authors should (1) ensure all authors have a unique permanent identifier such as an ORCID iD (Open Researcher and Contributor ID, 2009) and include this identifier with manuscript and data submission (because institutional emails are often impermanent); (2) include a detailed data availability statement with manuscript submission; (3) assign data and meta-data a globally unique and persistent identifier (e.g., DOI), which ensures long-term discoverability; (4) use available data ‘tags’ in the archiving repository; (5) use consistent names/usernames across the different journal and archiving platforms; and (6) craft the title and abstract and select keywords that will optimize the chances of research being identified via the most typical evidence synthesis method of searching for research (i.e., electronic database searches). To ensure their data is also *Interoperable* and *Reusable*, researchers need to attend to licensing, data file structures and formats, and detailed meta-data when preparing their data for sharing.

Data Availability Statements to Ensure Findability A data availability statement (DAS; also known as “data access statements” or “availability of materials”) provides information about how and where other researchers can access the study data and is often required when submitting a manuscript for publication (Federer et al., 2018; Graf et al., 2020). In some closed-access journals, DAS are treated as article meta-data, so they are accessible even if the article is behind a paywall, ensuring the data are findable. Thus, the DAS should be as explicit as possible, even if some information that is in the manuscript is repeated in the DAS.

The DAS for our mindfulness trial example states: “Both the full raw dataset and the finalized analytical dataset used in this analysis are available from Zenodo (DOI: <https://doi.org/10.5281/zenodo.4011716>), along with comprehensive data dictionaries and crosswalks.”

Creating Titles, Abstracts, and Keywords to Optimize Findability Carefully crafting titles and abstracts, in addition to choosing appropriate keywords, is essential to help other researchers find your research. As much as possible given word limits, these should describe the condition, exposure, and settings of a study (Grant et al., 2018; Montgomery et al., 2018). These pieces of the submission are often the last to be created but are the main pieces of information indexed and displayed in electronic databases and are used to identify relevant research for an evidence synthesis. As a result of their ability to increase the findability of research, we suggest a data-driven approach to crafting these sections. The `litsearchr` package for the R programming environment offers a systematic approach to generate a list of words likely to improve the identifiability of the study manuscript (Grames et al., 2019; R Core Team, 2019).

Using our mindfulness trial example, we began with a set of keywords we thought likely to capture a set of relevant published articles from which we could draw the most highly used keywords. We conducted a search using these keywords in PubMed on August 24, 2020 and used `litsearchr` to systematically generate a list of important terms. The results suggest that several terms could be strategically incorporated in the title, abstract, and keywords to increase the findability of the trial (Online, Table 3 provides further details using our example).

Data Licenses Assigning a license to the data being shared is important to govern how the data can be reused and, more importantly, to help ensure that the study team is credited.²

For our mindfulness trial example, we chose the CC-BY-4.0 license, which allows the reuse of the data provided the original researchers are credited with the creation of the dataset.

² Comparing the full suite of licenses available is beyond the scope of this article; online resources such as ChooseALicense (<https://choosealicense.com/non-software/>) and Creative Commons (<https://creativecommons.org/licenses/>) provide detailed descriptions of the range of licenses available for data sharing.

Table 1 Comparison of the options for sharing data

Location	Examples	Advantages	Disadvantages	Examples
Non-affiliated repositories	Zenodo; Dryad; Figshare; Open Science Framework	Free (except Dryad) Self archiving. DOI assigned. Long-term storage. Can blind repositories for double-blind peer- review (Anonymous Github; OSF “Anonymous” Links)	Some have a Data Publishing Charge (e.g., Dryad charges \$120). No review of files uploaded Available from https://doi.org/10.5281/zenodo.3529382	Repository for Abbott et al., 2019, contains raw dataset, analysis data- set, cleaning code, analysis code.
Institutional repositories	data.bris; Harvard Dataverse; Inter-university Consortium for Political and Social Research; UK Data Service	Supported by institution DOI assigned. Often can get help from staff when uploading data	Not as well known as other repositories—limits findability	Pescosolido et al., 2020, contains scripts and data to replicate all fig- ures/analyses found in the original manuscript
Supplemental files attached to the online version of the published article	Many file types (.txt,.xlsx,.doc,.pdf) with data or metadata	Likely relatively less work to prepare compared to the other options	No DOI (for supplemental files) Can be closed-access. Potential for data to be lost. Least findable of the other options	Slep et al., 2020, contains mPLUS code and data Tomiczyk et al., 2020, open access, includes data in manuscript and sup- plemental files

Data File Structures and Formats Sharing individual participant data, i.e., data on each variable collected for each participant, affords those wishing to reuse or synthesize these data a range of advantages. Access to data on all variables collected, rather than just those used in the primary study's analysis, allows synthesists to ensure that an estimate from a primary study is adjusted for a common set of variables when it is included in a meta-analysis, even if the published effect estimate for that study was not (Riley et al., 2010). Even among individual participant data, different levels of granularity exist.

It may not be possible to share each individual participant data point, especially when participant confidentiality is at risk, for example, when studying a rare disease or among participants of specialized programs. In these cases, there are several options for sharing individual participant data. The first is to generate a "synthetic" dataset that mimics the individual-level data by preserving statistical properties and the relationships between variables. Software, for example, the synthpop R package, can help users generate a synthetic dataset (Quintana, 2020, provides a tutorial on using this package). A second option is to remove some individual characteristics that might give away participants' identities, such as age, sex, and location of recruitment source, but provide the rest of the variables as individual data points. Finally, if it is not possible to share any individual participant data, then at the very minimum, summary-level data (e.g., means and standard deviations or percentages which describe the distribution of all variables between intervention/exposure groups) should be provided for all variables included in the paper and, if possible, the dataset, not solely those thought by the primary study team to be relevant. Of course, for investigators conducting years of longitudinal studies examining numerous questions for which they are not ready to share data, we advise authors to explicitly state in their DAS the reasons for not sharing data. By beginning to capture the true reasons for not sharing data, data availability statements can then begin to be used as support for introducing better protection for and citation of data producers.

To illustrate the distinction in the level of data shared, for our mindfulness trial, we shared the raw data exported from the survey instrument, which contains each participant's answer to each individual question, and the processed individual participant data, which include scale score summaries (i.e., total DASS-21 score, comprised of 21 individual questions). To explore these different datasets, see the "REDCAP_data.xlsx" and "combined_data.xlsx" files in our Github repository which is archived by Zenodo (DOI: <https://doi.org/10.5281/zenodo.4011716>).

Study data can be shared as machine-readable (e.g., unformatted Excel/CSV files and raw DNA microarray

sequences) or human-readable formats (e.g., formatted PDF tables, figures, and data contained in text such as "There was evidence of an association (Odds ratio: 0.91, 95%CI: 0.85–0.96)"). Although tools are being developed to extract data automatically from human-readable sources for evidence syntheses (Marshall et al., 2020), it is often challenging to import these data into statistical analysis software, reducing interoperability and synthesis-readiness.

For our mindfulness trial example, all study data are shared via Excel files in our Github repository which is archived by Zenodo (DOI: <https://doi.org/10.5281/zenodo.4011716>). This includes the mindfulness trial's quantitative survey data. It also includes qualitative data, which were generated from in-person, semi-structured interviews and coded to examine adherence and problems with using the tools. To demonstrate synthesis-readiness, the relevant responses for qualitative coding are included in the data file rather than the full interview notes.

Meta-data Providing a README, a plain text file that contains useful information about data or software, can help ensure that data are correctly accessed, interpreted, and reused by other researchers. The README file should include a (1) definition of the primary research aims using a relevant alternative framework that addresses the main components of the research which will help others identify and use the research (e.g., PICOTS, TOPICS + M; Haynes, 2006; Johnson & Hennessy, 2019; Stern et al., 2014); (2) a short description of all the data files included in the project repository; (3) signposts to the detailed data dictionaries for each data file which contain definitions of all variables, including coded (e.g., "CRP_t2") and interpretable (e.g., "C-Reactive Protein levels at 2 months post-baseline") variable names, the values that each variable is allowed to take, and include missing data indicators and units of measurement; (4) a summary of data processing steps that were used to produce the final analytic datasets from the raw data files; (5) a description of each data file's relationship to other data files in the repository, and relevant linkage variables, such as participant ID; (6) permanent author contact details; and (7) details of the license, including the full name of the license holder, the year, and where the full text of the license can be found.

The README for our mindfulness trial contains the seven main elements necessary for a comprehensive metadata file; see Fig. 2 for a snapshot of these pieces and visit Zenodo to view and download all files: <https://doi.org/10.5281/zenodo.4011716>.

Archiving and Sharing Research Data

Research data sharing can be done via institutional and non-institutional repositories or alongside published articles

<p>Study description</p> <p>Taken from the Study protocol.docx document, which describes the study approach in detail:</p> <p>"Young adults in college experience high levels of stress, anxiety, and depression, which can lead to maladaptive behaviors and chronic health issues including inflammation and HPA-axis dysregulation. University mental health centers are seeking effective programs that teach coping skills to self-manage stress, with low stigma, that are easy to implement in large scale. The aim of this protocol is to describe the design of a study aiming to evaluate the effectiveness of a one-month self-guided meditation program for university students comparing various tools to improve mental health, physical health and health behaviors, and explore moderators. Secondary aims include reaching a diverse audience and documenting adverse experiences."</p>	<p>PICOTS</p> <p>Population - 140 university undergraduate or postgraduate students studying at an American university.</p> <p>Intervention - 10 minutes of self-guided meditation daily for four weeks, facilitated by the 10% Happier App, in combination with neurofeedback from the Muse Neurofeedback Tool.</p> <p>Comparison - 10 minutes of self-guided meditation daily for four weeks, facilitated by the 10% Happier App.</p> <p>Outcomes - Outcome measures included those related to Personality, Stress, Distress, Resilience (coping), Mindfulness, Emotion regulation, Sleep, Self-Regulation of Eating, Motivation Survey, and Student Adherence (for more details, see Section 2.4 of the Study protocol.docx document and the Instrument List.xlsx document).</p> <p>Time - 4 week intervention followed by post-intervention surveys (1-, 6- and 12-months). See Timeline.xlsx for more details on the tasks performed and the data collected at each timepoint.</p> <p>Study Design - randomized controlled trial with two intervention groups</p>
<p>(a)</p> <p>Data source files</p> <p>All data files are available as Excel spreadsheets in the <code>data/</code> subdirectory. The relationships between each data file, and between data files and other information contained in this repository, are described below:</p> <ul style="list-style-type: none"> • <code>RECAP_data.xlsx</code> (Data dictionary: Data Dictionary for REDCAP Data.pdf) contains the raw survey data exported from the REDCAP software. • <code>External_data.xlsx</code> (Data dictionary: Data Dictionary for External Data.xlsx) contains additional variables on each participant which were not captured by the REDCAP survey. Note some variables, such as C-Reactive protein or cortisol levels, are only available for a subset of participants, based on the availability of saliva samples for that participant. See Section 2.2.1 in Study protocol.docx for more details. • <code>Combined_data.xlsx</code> was created by merging the <code>RECAP_data.xlsx</code> and <code>External_data.xlsx</code>, and is used as the complete dataset from which composite variables are created using the <code>create_composite_variables.do</code> STATA do file. The crosswalk between the individual measures and the composite variables are described in more detail in <code>Stata Variable Crosswalk.xlsx</code> • <code>Debriefing Interview - Wave 1.xlsx</code> : contains qualitative responses from "Wave 1" participants to the debriefing interview questions. Note: the interview was only done for Wave 1 participants. <p>(c)</p>	<p>(b)</p> <p>Permanent contact about this data</p> <p>Rebeca L. Acabchuk, University of Connecticut, Institute for Collaboration on Health, Intervention, & Policy, Department of Psychological Sciences, United States. ORCID: 0000-0001-8254-2396</p> <p>› Licence</p> <p>The materials in this repository are licensed under the Creative Commons Attribution 4.0 (CC-BY-4.0) International. The full license file can be found here.</p> <p>(d)</p>

Fig. 2 Mindfulness trial metadata included in the README file. Panels **a** and **b** provide the study description according to the PICOTS framework; **c** provides data source files, locations, and rela-

tionships between the files; **d** provides permanent author information and licensing information

(Table 1). We have ordered these sharing options from best practice (most likely to be synthesis-ready) to good practice (still synthesis-ready but may require additional effort to prepare for synthesis).

Non-affiliated Repositories Non-affiliated repositories are commercial or charity repositories that allow researchers to self-archive, and assign a DOI to, their research data. Examples include Dryad, Open Science Framework, and Zenodo (Dryad, 2020; Open Science Framework, 2020; Zenodo—Research Shared, 2020). The main limitation of these repositories is that there are often conditions on how much data can be stored; some charge fees for storage.

For our mindfulness trial example, the repository was created in GitHub and archived using Zenodo, as GitHub deposition is non-permanent and does not assign a DOI to the uploaded data. Repositories of this type can be embargoed prior to publication to prevent “scooping,” and some repositories also provide services to help prevent scooping. For example, projects hosted in the Open Science Framework can have “anonymous” links to remove identifying meta-data as part of double-blind peer-review requirements.

Institutional Repositories An institutional repository is one hosted by an academic institution, such as the ICPSR (ICSPR, 2021). An advantage of affiliated repositories is that some provide staff to guide researchers who are new to data sharing. These repositories are less well-known, which may limit the findability of data when compared to other options. The terms of deposition should be reviewed to ensure that authors retain data ownership and copyright over the deposited materials.

Supplemental Files A common option for sharing research data is as a supplementary file to the published manuscript. However, this method has several major limitations. This approach reduces data findability because datasets shared as supplementary material are grouped under the publication’s DOI and are not uniquely identifiable. There is also the potential for supplementary files to become disassociated with their respective publications, for example, during a website redesign. If the published article is behind a paywall, then data attached to it as supplementary files may also be blocked and so data shared as supplemental files may not be fully accessible. Sharing data as supplementary files to an open-access publication should be considered the minimum level of data sharing needed to make research synthesis-ready.

Potential Challenges and Future Considerations

In this article, we have outlined the benefits of synthesis-ready research and provided guidelines on how to follow best practice and produce synthesis-ready research. Yet, we appreciate that there are many perceived and actual challenges to doing so, including costs to data sharing, lack of incentives, lack of awareness or training, not wanting to “give away” data or get “scooped” on findings, and the necessity of ensuring participant confidentiality (Evans, 2016; Tennant et al., 2019; Tenopir et al., 2011, 2015; van Panhuis et al., 2014; Walters, 2020).

To address the lack of knowledge and training, we have provided information, workflows, and a full demonstration using an empirical example, in addition to signposting to further literature, to help researchers plan for the resources needed for making research synthesis-ready at project conceptualization. Many funders have recognized the value of data sharing and will provide funds to do so based on investigator budgets which can offset the tangible (platform sharing fees) and intangible (time spent preparing data) costs of this process.

Researchers have valid concerns about having their research scooped, and yet, several measures can help prevent scooping including pre-registering study analysis plans and depositing preprints while awaiting publication, all core practices of open science. An additional solution is to license data at the point of publication or archival (“[How to Make Research Synthesis-Ready: Steps from Study Design to Completion](#)”). Finally, some journals have implemented “scoop-protection” policies for research, a system-level solution we hope more journals will implement to address this challenge (Transpose, 2020). Ensuring that all types of research data are reused appropriately is a warranted concern (Bishop, 2009; Yardley et al., 2014); yet, evidence suggests that individuals reusing data seek to collaborate with the original data creators to harness their knowledge about the data, rather than using it to scoop findings or to disprove the original publication’s findings, resulting in improved and more accurate data reuse and increased opportunities for publication (Pasquetto et al., 2019). As well, providing clearly detailed documentation in the data and meta-data files (“[How to Make Research Synthesis-Ready: Steps from Study Design to Completion](#)” and Fig. 2) should reduce inappropriate use of shared data. Finally, we have provided several different options for maintaining the confidentiality of participants while attending to synthesis-readiness (“[How to Make Research Synthesis-Ready: Steps from Study Design to Completion](#)”).

Conclusions

Despite the importance of making research available and synthesis-ready, not all scientific disciplines have embraced data/meta-data sharing and transparency at the same rate (Graf et al., 2020; Hardwicke et al., 2020; Sholler et al., 2019; Vasilevsky et al., 2017). If prevention scientists share research in synthesis-ready forms, the field would see improvements, especially in terms of impact on policy and practice. We presented methods for managing data sharing processes that should make it easier to produce synthesis-ready research. Changes in research practice that focus on synthesizability will reduce the delay between implementing effective primary study approaches (Nakagawa et al., 2020) and should have a positive impact on translation of prevention science into policy and practice.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11121-021-01279-8>.

Acknowledgements The paper was first developed during the 2019 Evidence Synthesis Hackathon.

Author Contribution Project administration (EAH; LAM), roles/writing — original draft (EAH; LAM), visualization (EAH; LAM); data curation (RLA; LAM); resources (LAM); funding acquisition (NRH); investigation (EAH; LAM; RLA), methodology (EAH; LAM); software (EAH; LAM); conceptualization (All authors), writing — review and editing (all authors).

Funding The Fenner School of Environment & Society (ANU), the University of New South Wales, and the Faculty of Humanities at the University of Johannesburg funded the Evidence Synthesis Hackathon 2019. EAH has support from NIAAA (K01 AA028536-01). MJP has support from an Australian Research Council Discovery Early Career Researcher Award (DE200101618). LAM has support by an NIH Doctoral Research Fellowship (DRF-2018-11-ST2-048).

Declarations

Conflict of Interest The authors declare no competing interests.

Disclaimer The views expressed in this article are those of the authors and do not necessarily represent those of the NIAAA, NHS, the NIHR, or the Department of Health and Social Care.

References

- Abbott, S., Christensen, H., Welton, N. J., & Brooks-Pollock, E. (2019). Estimating the effect of the 2005 change in BCG policy in England: A retrospective cohort study, 2000 to 2015. *Eurosurveillance*, 24(49), 1,900,220. <https://doi.org/10/gg9nqk>
- Acabchuk, R. L., Simon, M. A., Low, S., Brisson, J. M., & Johnson, B. T. (2021). Measuring meditation progress with a consumer-grade EEG device: Caution from a randomized controlled trial. *Mindfulness*, 12, 68–81. <https://doi.org/10.1007/s12671-020-01497-1>

- Banks, G. C., Field, J. G., Oswald, F. L., O'Boyle, E. H., Landis, R. S., Rupp, D. E., & Rogelberg, S. G. (2019). Answers to 18 questions about open science practices. *Journal of Business and Psychology*, 34, 257–270. <https://doi.org/10.1007/s10869-018-9547-8>
- Barton, C., Smith, R., & Weaver, R. (2010). Data practices, policy, and rewards in the information era demand a new paradigm. *Data Science Journal*, 9, IGY95–IGY99. https://doi.org/10.2481/dsj-SS_IGY-003
- Beugelsdijk, S., van Witteloostuijn, A., & Meyer, K. E. (2020). A new approach to data access and research transparency (DART). *Journal of International Business Studies*, 51, 887–905. <https://doi.org/10.1057/s41267-020-00323-z>
- Bishop, L. (2009). Ethical sharing and reuse of qualitative data. *Australian Journal of Social Issues*, 44(3), 255–272.
- Borah, R., Brown, A. W., Capers, P. L., & Kaiser, K. A. (2017). Analysis of the time and workers needed to conduct systematic reviews of medical interventions using data from the PROSPERO registry. *BMJ Open*, 7, e012545. <https://doi.org/10.1136/bmjopen-2016-012545>
- Chan, A. W., Tetzlaff, J. M., Altman, D. G., Laupacis, A., Gøtzsche, P. C., Krleža-Jerić, K., & Doré, C. J. (2013). *SPIRIT 2013 statement: Defining standard protocol items for clinical trials*. Annals of Internal Medicine; Ann Intern Med. <https://doi.org/10.7326/0003-4819-158-3-201302050-00583>
- Choose Muse. (2018, June 25). *A deep dive into brainwaves: Brainwave frequencies explained*. Muse. <https://choosemuse.com/blog/a-deep-dive-into-brainwaves-brainwave-frequencies-explained-2/>
- Christensen, G., Dafoe, A., Miguel, E., Moore, D. A., & Rose, A. K. (2019). A study of the impact of data sharing on article citations using journal policies as a natural experiment. *PLoS One*, 14, e0225883. <https://doi.org/10.1371/journal.pone.0225883>
- Colavizza, G., Hrynaszkiewicz, I., Staden, I., Whitaker, K., & McGillivray, B. (2020). The citation advantage of linking publications to research data. *PLoS One*, 15, 023–0416. <https://doi.org/10.1371/journal.pone.0230416>
- Crosas, M. (2013, October 30). *Joint Declaration of Data Citation Principles—FINAL*. FORCE11. <https://www.force11.org/datacitationprinciples>
- Dryad. (2020). https://datadryad.org/stash/our_mission
- Evans, S. R. (2016). Gauging the purported costs of public data archiving for long-term population studies. *PLoS Biology*, 14, e1002432. <https://doi.org/10.1371/journal.pbio.1002432>
- Falk Delgado, A., & Falk Delgado, A. (2017). Outcome switching in randomized controlled oncology trials reporting on surrogate endpoints: A cross-sectional analysis. *Scientific Reports*, 7. <https://doi.org/10.1038/s41598-017-09553-y>
- Federer, L. M., Belter, C. W., Joubert, D. J., Livinski, A., Lu, Y. L., Snyders, L. N., & Thompson, H. (2018). Data sharing in PLOS ONE: An analysis of Data Availability Statements. *PLoS One*, 13, e0194768. <https://doi.org/10/gdhb7j>
- Forstmeier, W., Wagenmakers, E. J., & Parker, T. H. (2017). Detecting and avoiding likely false-positive findings – a practical guide. *Biological Reviews*, 92, 1941–1968. <https://doi.org/10.1111/brv.12315>
- Frankenhuis, W. E., & Nettle, D. (2018). Open science is liberating and can foster creativity. *Perspectives on Psychological Science*, 13, 439–447. <https://doi.org/10.1177/1745691618767878>
- Gaba, J. F., Siebert, M., Dupuy, A., Moher, D., & Naudet, F. (2020). Funders' data-sharing policies in therapeutic research: A survey of commercial and non-commercial funders. *PLoS One*, 15, e0237464. <https://doi.org/10.1371/journal.pone.0237464>
- Gerstner, K., Moreno-Mateos, D., Gurevitch, J., Beckmann, M., Kambach, S., Jones, H. P., & Seppelt, R. (2017). Will your paper be used in a meta-analysis? Make the reach of your research broader and longer lasting. *Methods in Ecology and Evolution*, 8, 777–784. <https://doi.org/10.1111/2041-210X.12758>
- Graf, C., Flanagan, D., Wylie, L., & Silver, D. (2020). The open data challenge: An analysis of 124,000 data availability statements and an ironic lesson about data management plans. *Data Intelligence*, 1–15. https://doi.org/10.1162/dint_a_00061
- Grames, E. M., Stillman, A. N., Tingley, M. W., & Elphick, C. S. (2019). An automated approach to identifying search terms for systematic reviews using keyword co-occurrence networks. *Methods in Ecology and Evolution*, 10, 1645–1654.
- Grant, S., Mayo-Wilson, E., Montgomery, P., Macdonald, G., Michie, S., Hopewell, S., Moher, D., Aber, J. L., Altman, D., Bhui, K., Booth, A., Clark, D., Craig, P., Eisner, M., Fraser, M. W., Gardner, F., Grant, S., Hedges, L., Hollon, S., & CONSORT-SPI Group. (2018). CONSORT-SPI 2018 explanation and elaboration: Guidance for reporting social and psychological intervention trials. *Trials*, 19, 406. <https://doi.org/10.1186/s13063-018-2735-z>
- Gupta, Y. K., Meenu, M., & Mohan, P. (2015). The Tamiflu fiasco and lessons learnt. *Indian Journal of Pharmacology*, 47, 11–16. <https://doi.org/10.4103/0253-7613.150308>
- Haddaway, N. R., & Westgate, M. J. (2019). Predicting the time needed for environmental systematic reviews and systematic maps. *Conservation Biology*, 33(2), 434–443. <https://doi.org/10.1111/cobi.13231>
- Hardwicke, T. E., Wallach, J. D., Kidwell, M. C., Bendixen, T., Crüwell, S., & Ioannidis, J. P. A. (2020). An empirical assessment of transparency and reproducibility-related research practices in the social sciences (2014–2017). *Royal Society Open Science*, 7, 1908706. <https://doi.org/10.1098/rsos.190806>
- Haynes, R. B. (2006). Forming research questions. *Journal of Clinical Epidemiology*, 59, 881–886. <https://doi.org/10.1016/j.jclinepi.2006.06.006>
- ICPSR. (2021). Retrieved February 24, 2021, from <https://www.icpsr.umich.edu/web/pages/>
- Inter-university Consortium for Political and Social Research (2010). Guide to social science data preparation and archiving best practice throughout the data life cycle. *ICPSR Institute for Social Research University of Michigan*.
- Johnson, B. T., & Hennessy, E. A. (2019). Systematic reviews and meta-analyses in the health sciences: Best practice methods for research syntheses. *Social Science & Medicine*, 233, 237–251. <https://doi.org/10.1016/j.socscimed.2019.05.035>
- Johnson, B. T., Cromley, E. K., & Marrouch, N. (2017). Spatiotemporal meta-analysis: Reviewing health psychology phenomena over space and time. *Health Psychology Review*, 11, 280–291.
- Kahan, B. C., & Jairath, V. (2018). Outcome pre-specification requires sufficient detail to guard against outcome switching in clinical trials: A case study. *Trials*, 19, 265. <https://doi.org/10.1186/s13063-018-2654-z>
- Kerr, N. L. (1998). HARKing: Hypothesizing after the results are known. *Personality and Social Psychology Review*, 2, 196–217.
- Marshall, I. J., Johnson, B. T., Wang, Z., Rajasekaran, S., & Wallace, B. C. (2020). Semi-Automated evidence synthesis in health psychology: Current methods and future prospects. *Health Psychology Review*, 14, 145–158. <https://doi.org/10.1080/17437199.2020.1716198>
- Miyakawa, T. (2020). No raw data, no science: Another possible source of the reproducibility crisis. *Molecular Brain*, 13, 24. <https://doi.org/10.1186/s13041-020-0552-2>
- Montgomery, P., Grant, S., Mayo-Wilson, E., Macdonald, G., Michie, S., Hopewell, S., Moher, D., Lawrence Aber, J., Altman, D., Bhui, K., Booth, A., Clark, D., Craig, P., Eisner, M., Fraser, M. W., Gardner, F., Grant, S., Hedges, L., Hollon, S., & CONSORT-SPI Group. (2018). Reporting randomised trials of social and psychological interventions: The CONSORT-SPI 2018 Extension. *Trials*, 19, 407–407. <https://doi.org/10.1186/s13063-018-2733-1>
- Nakagawa, S., Dunn, A. G., Lagisz, M., Bannach-Brown, A., Grames, E. M., Sánchez-Tójar, A., O'Dea, R. E., Noble, D. W. A., Westgate,

- M. J., Arnold, P. A., Barrow, S., Bethel, A., Cooper, E., Foo, Y. Z., Geange, S. R., Hennessy, E., Mapanga, W., Mengersen, K., Munera, C., & Hackathon, E. S. (2020). A new ecosystem for evidence synthesis. *Nature Ecology & Evolution*, 4, 498–501. <https://doi.org/10.1038/s41559-020-1153-2>
- NIH data sharing policy and guidance. (2003). National Institutes of Health. https://grants.nih.gov/grants/policy/data_sharing/data_sharing_guidance.htm
- Nuijten, M. B., Hartgerink, C. H. J., van Assen, M. A. L. M., Epskamp, S., & Wicherts, J. M. (2016). The prevalence of statistical reporting errors in psychology (1985–2013). *Behavior Research Methods*, 48, 1205–1226. <https://doi.org/10.3758/s13428-015-0664-2>
- Ohmann, C., Banzi, R., Canham, S., Battaglia, S., Matei, M., Ariyo, C., Beclen, L., Bierer, B., Bowers, S., Clivio, L., Dias, M., Druml, C., Faure, H., Fenner, M., Galvez, J., Ghersi, D., Gluud, C., Groves, T., Houston, P., & Demotes-Mainard, J. (2017). Sharing and reuse of individual participant data from clinical trials: Principles and recommendations. *BMJ Open*, 7, e018647. <https://doi.org/10.1136/bmjopen-2017-018647>
- Open Science Framework. (2020). <https://osf.io/>
- ORCID iD. (2009). "ORCID website". orcid.org. Retrieved 18 April 2021.
- Pasquetto, I. V., Borgman, C. L., & Wofford, M. F. (2019). Uses and reuses of scientific data: The data creators' advantage. *Harvard Data Science Review*, 1. <https://doi.org/10.1162/99608f92.fc14bf2d>
- Pescosolido, B., Lee, B., & Kafadar, K. (2020). Replication data for: Socio-demographic similarity at the contextual level alters individual risk for completed suicide [Data set]. Harvard Dataverse. <https://doi.org/10.7910/DVN/35IV23>
- Popkin, G. (2019). Data sharing and how it can benefit your scientific career. *Nature*, 569, 445–447.
- Quintana, D. S. (2020). A synthetic dataset primer for the biobehavioral sciences to promote reproducibility and hypothesis generation. *Elife*, 9, e53275.
- R Core Team. (2019). *R: A language and environment for statistical computing*. <https://www.R-project.org/>
- Riley, R. D., Lambert, P. C., & Abo-Zaid, G. (2010). Meta-analysis of individual participant data: Rationale, conduct, and reporting. *BMJ*, 340. <https://doi.org/10.1136/bmj.c221>
- Schulz, K. F., Altman, D. G., Moher, D., & CONSORT Group. (2010). CONSORT 2010 statement: Updated guidelines for reporting parallel group randomised trials. *BMJ (clinical Research Ed.)*, 340, c332. <https://doi.org/10.1136/bmj.c332>
- Sholler, D., Ram, K., Boettiger, C., & Katz, D. S. (2019). Enforcing public data archiving policies in academic publishing: A study of ecology journals. *Big Data & Society*, 6, epub. <https://doi.org/10.1177/2053951719836258>
- Slep, A. M. S., Heyman, R. E., Lorber, M. F., Baucom, K. J. W., & Linkh, D. J. (2020). Evaluating the Effectiveness of NORTH STAR: A Community-Based Framework to Reduce Adult Substance Misuse, Intimate Partner Violence, Child Abuse, Suicidality, and Cumulative Risk. *Prevention Science*. <https://doi.org/10.gg898m>
- Stern, C., Jordan, Z., & McArthur, A. (2014). Developing the review question and inclusion criteria. *AJN the American Journal of Nursing*, 114, 53–56.
- Strasser, C. (2012). *Primer on data management: What you always wanted to know*.
- Ten Percent Happier: Mindfulness Meditation Courses. (2020). Ten percent happier. <https://www.tenpercent.com>
- Tennant, J. P., Crane, H., Crick, T., Davila, J., Enkhbayar, A., Havemann, J., Kramer, B., Martin, R., Masuzzo, P., Nobes, A., Rice, C., Rivera-López, B., Ross-Hellauer, T., Sattler, S., Thacker, P. D., & Vanholsbeeck, M. (2019). Ten hot topics around scholarly publishing. *Publications*, 7, 34. <https://doi.org/10.3390/publications7020034>
- Tenopir, C., Allard, S., Douglass, K., Aydinoglu, A. U., Wu, L., Read, E., Manoff, M., & Frame, M. (2011). Data sharing by scientists: Practices and perceptions. *PLoS One*, 6, e21101. <https://doi.org/10.1371/journal.pone.0021101>
- Tenopir, C., Dalton, E. D., Allard, S., Frame, M., Pjesivac, I., Birch, B., Pollock, D., & Dorsett, K. (2015). Changes in data sharing and data reuse practices and perceptions among scientists worldwide. *PLoS One*, 10, e0134826. <https://doi.org/10.1371/journal.pone.0134826>
- Thorogood, A., & Knoppers, B. M. (2017). Can research ethics committees enable clinical trial data sharing? *Ethics, Medicine and Public Health*, 3, 56–63.
- Tomczyk, S., Schomerus, G., Stolzenburg, S., Muehlan, H., & Schmidt, S. (2020). Ready, willing and able? An investigation of the theory of planned behaviour in help-seeking for a community sample with current untreated depressive symptoms. *Prevention Science*, 21(6), 749–760. <https://doi.org/10.gg898n>
- Transpose. (2020). *Transpose Database: A database of journal policies on peer review, co-reviewing, and preprinting*. <https://transpose-publishing.github.io/#/>
- van Panhuis, W. G., Paul, P., Emerson, C., Grefenstette, J., Wilder, R., Herbst, A. J., Heymann, D., & Burke, D. S. (2014). A systematic review of barriers to data sharing in public health. *BMC Public Health*, 14, 1144. <https://doi.org/10.1186/1471-2458-14-1144>
- Vasilevsky, N. A., Minnier, J., Haendel, M. A., & Champieux, R. E. (2017). Reproducible and reusable research: Are journal data sharing policies meeting the mark? *PeerJ*, 5.
- Vicente-Saez, R., & Martinez-Fuentes, C. (2018). Open Science now: A systematic literature review for an integrated definition. *Journal of Business Research*, 88, 428–436. <https://doi.org/10.1016/j.jbusres.2017.12.043>
- Walters, W. H. (2020). Data journals: Incentivizing data access and documentation within the scholarly communication system. *Insights*, 33, 18. <https://doi.org/10.1629/uksg.510>
- Wellcome Trust Public Health Research Data Forum. (2012). *Summary of Funders' data sharing policies*. <https://wellcome.ac.uk/sites/default/files/summary-of-phrdrf-funder-data-sharing-policies.pdf>
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J. W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., & Mons, B. (2016). The FAIR guiding principles for scientific data management and stewardship. *Scientific Data*, 3, 160018. <https://doi.org/10.1038/sdata.2016.18>
- Wilson, S. J., & Tanner-Smith, E. E. (2014). Meta-analysis in prevention science. In Z. Sloboda & H. Petras (Eds.), *Defining prevention science* (pp. 431–452). Springer US. https://doi.org/10.1007/978-1-4614-7424-2_19
- Yardley, S. J., Watts, K. M., Pearson, J., & Richardson, J. C. (2014). Ethical issues in the reuse of qualitative data: Perspectives from literature, practice, and participants. *Qualitative Health Research*, 24(1), 102–113.
- Yenni, G. M., Christensen, E. M., Bledsoe, E. K., Supp, S. R., Diaz, R. M., White, E. P., & Ernest, S. K. M. (2019). Developing a modern data workflow for regularly updated data. *PLoS Biology*, 17, e3000125. <https://doi.org/10.1371/journal.pbio.3000125>
- Zenodo—Research Shared. (2020). <https://about.zenodo.org/>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Emily A. Hennessy¹  · Rebecca L. Acabchuk²  · Pieter A. Arnold³  · Adam G. Dunn⁴  · Yong Zhi Foo⁵  ·
Blair T. Johnson⁶  · Sonya R. Geange^{3,7}  · Neal R. Haddaway^{8,9,10}  · Shinichi Nakagawa⁵  · Witness Mapanga¹¹  ·
Kerrie Mengersen¹²  · Matthew J. Page¹³  · Alfredo Sánchez-Tójar¹⁴ · Vivian Welch¹⁵ · Luke A. McGuinness¹⁶

¹ Center for Addiction Medicine, Recovery Research Institute, Massachusetts General Hospital & Harvard Medical School, Boston, MA, USA

² Institute for Collaboration On Health, Intervention, Department of Psychological Sciences, University of Connecticut & Policy, Mansfield, USA

³ Research School of Biology, The Australian National University, Canberra, Australia

⁴ Biomedical Informatics and Digital Health, School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney, Sydney, Australia

⁵ Evolution & Ecology Research Centre and School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia

⁶ Department of Psychological Sciences, University of Connecticut, 406 Babbidge Road, Unit 1020, Storrs, CT 06269-1020, USA

⁷ Department of Biological Sciences, University of Bergen, Bergen, Norway

⁸ Stockholm Environment Institute, Stockholm, Sweden

⁹ Mercator Research Institute On Global Commons and Climate Change, Berlin, Germany

¹⁰ Africa Centre for Evidence, Johannesburg, South Africa

¹¹ Faculty of Health Sciences, Non-Communicable Diseases Research Division of the Wits Health Consortium, University of the Witwatersrand, Johannesburg, South Africa

¹² School of Mathematical Sciences, Queensland University of Technology, Brisbane, Australia

¹³ School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia

¹⁴ Department of Evolutionary Biology, Bielefeld University, Bielefeld, Germany

¹⁵ School of Epidemiology and Public Health, Bruyere Research Institute, University of Ottawa, Ottawa, Canada

¹⁶ Department of Population Health Sciences, University of Bristol, Bristol, UK