

FINITE ELEMENTS IN N-D

COMPUTING ELEMENT ARRAYS

Maurizio M. Chiaramonte

COMPUTING THE ELEMENT ARRAYS

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and

$$\begin{aligned} \mathbf{k}_{ab}^e &= \mathbf{a}^e(\phi_a^e, \phi_b^e) = \int_{\Omega^e} \nabla_{\mathbf{x}} \phi_a^e(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \phi_b^e d\Omega = \\ &= \int_{\hat{\Omega}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_a(\xi)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_b(\xi)] \hat{j}(\xi) d\hat{\Omega} \\ &\approx \sum_{(\tilde{\xi}_Q, \omega_Q) \in \mathcal{Q}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_Q))^{-\top} \nabla_{\xi} \hat{\phi}_a(\tilde{\xi}_Q)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_Q))^{-\top} \nabla_{\xi} \hat{\phi}_b(\tilde{\xi}_Q)] \hat{j}^e(\tilde{\xi}_Q) \omega_Q \end{aligned}$$

where $\hat{j}^e(\xi) = \det(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))$.

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and

$$\begin{aligned} \mathbf{k}_{ab}^e &= \mathbf{a}^e(\phi_a^e, \phi_b^e) = \int_{\Omega^e} \nabla_{\mathbf{x}} \phi_a^e(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \phi_b^e d\Omega = \\ &= \int_{\hat{\Omega}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_a(\xi)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_b(\xi)] \hat{j}(\xi) d\hat{\Omega} \\ &\approx \sum_{(\tilde{\xi}_0, \omega_0) \in \mathcal{Q}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_0))^{-\top} \nabla_{\xi} \hat{\phi}_a(\tilde{\xi}_0)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_0))^{-\top} \nabla_{\xi} \hat{\phi}_b(\tilde{\xi}_0)] \hat{j}^e(\tilde{\xi}_0) \omega_0 \end{aligned}$$

where $\hat{j}^e(\xi) = \det(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))$.

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and

$$\begin{aligned} \mathbf{k}_{ab}^e &= \mathbf{a}^e(\phi_a^e, \phi_b^e) = \int_{\Omega^e} \nabla_{\mathbf{x}} \phi_a^e(\mathbf{x}) \cdot \nabla_{\mathbf{x}} \phi_b^e d\Omega = \\ &= \int_{\hat{\Omega}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_a(\xi)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))^{-\top} \nabla_{\xi} \hat{\phi}_b(\xi)] \hat{j}(\xi) d\hat{\Omega} \\ &\approx \sum_{(\tilde{\xi}_Q, \omega_Q) \in \mathcal{Q}} [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_Q))^{-\top} \nabla_{\xi} \hat{\phi}_a(\tilde{\xi}_Q)] \cdot [(\nabla_{\xi} \hat{\mathbf{x}}^e(\tilde{\xi}_Q))^{-\top} \nabla_{\xi} \hat{\phi}_b(\tilde{\xi}_Q)] \hat{j}^e(\tilde{\xi}_Q) \omega_Q \end{aligned}$$

where $\hat{j}^e(\xi) = \det(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))$.

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and ¹

$$\begin{aligned} \mathbf{f}_a^e &= F^e(\phi_a^e) = \int_{\Omega^e} \phi_a^e(\mathbf{x}) f(\mathbf{x}) d\Omega \\ &= \int_{\hat{\Omega}} \hat{\phi}_a(\xi) f(\hat{\mathbf{x}}^e(\xi)) \hat{j}^e(\xi) d\hat{\Omega} = \\ &\approx \sum_{(\tilde{\xi}_Q, \omega_Q) \in \mathcal{Q}} \hat{\phi}_a(\tilde{\xi}_Q) f(\hat{\mathbf{x}}^e(\tilde{\xi}_Q)) \hat{j}^e(\tilde{\xi}_Q) \omega_Q \end{aligned}$$

where $\hat{j}(\xi) = \det(\nabla_{\xi} \hat{\mathbf{x}}^e(\xi))$.

¹for simplicity we are assuming homogeneous Neumann boundary conditions hence

$$h = 0 \Rightarrow \int_{\Gamma_N} h \phi_i d\Gamma = 0$$

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and ¹

$$\begin{aligned} \mathbf{f}_a^e &= F^e(\phi_a^e) = \int_{\Omega^e} \phi_a^e(\mathbf{x}) f(\mathbf{x}) d\Omega \\ &= \int_{\hat{\Omega}} \hat{\phi}_a(\boldsymbol{\xi}) f(\hat{\mathbf{x}}^e(\boldsymbol{\xi})) \hat{j}^e(\boldsymbol{\xi}) d\hat{\Omega} = \\ &\approx \sum_{(\tilde{\xi}_Q, \omega_Q) \in \mathcal{Q}} \hat{\phi}_a(\tilde{\xi}_Q) f(\hat{\mathbf{x}}^e(\tilde{\xi}_Q)) \hat{j}^e(\tilde{\xi}_Q) \omega_Q \end{aligned}$$

where $\hat{j}(\boldsymbol{\xi}) = \det(\nabla_{\boldsymbol{\xi}} \hat{\mathbf{x}}^e(\boldsymbol{\xi}))$.

¹for simplicity we are assuming homogeneous Neumann boundary conditions hence

$$h = 0 \Rightarrow \int_{\Gamma_N} h \phi_i d\Gamma = 0$$

Constructing the Element Arrays

Recall that

$$\mathbf{K} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{k}^e, \quad \mathbf{F} = \mathbb{A}_{e=1}^{n_{el}} \mathbf{f}^e$$

where \mathbb{A} is the assembly operator, and ¹

$$\begin{aligned} \mathbf{f}_a^e &= F^e(\phi_a^e) = \int_{\Omega^e} \phi_a^e(\mathbf{x}) f(\mathbf{x}) d\Omega \\ &= \int_{\hat{\Omega}} \hat{\phi}_a(\boldsymbol{\xi}) f(\hat{\mathbf{x}}^e(\boldsymbol{\xi})) \hat{j}^e(\boldsymbol{\xi}) d\hat{\Omega} = \\ &\approx \sum_{(\tilde{\boldsymbol{\xi}}_Q, \omega_Q) \in \mathcal{Q}} \hat{\phi}_a(\tilde{\boldsymbol{\xi}}_Q) f(\hat{\mathbf{x}}^e(\tilde{\boldsymbol{\xi}}_Q)) \hat{j}^e(\tilde{\boldsymbol{\xi}}_Q) \omega_Q \end{aligned}$$

where $\hat{j}(\boldsymbol{\xi}) = \det(\nabla_{\boldsymbol{\xi}} \hat{\mathbf{x}}^e(\boldsymbol{\xi}))$.

¹for simplicity we are assuming homogeneous Neumann boundary conditions hence

$$h = 0 \Rightarrow \int_{\Gamma_N} h \phi_i d\Gamma = 0$$