

Kern- und Teilchenphysik II
Spring Term 2015

Exercise Sheet 1

Lecturers: Prof. F. Canelli, Prof. N. Serra
Assistants: Mr. E. Bowen, Dr. A. de Cosa

1. Dirac Matrices

Show that the Dirac matrices have the following properties:

- a) $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$
- b) $\{\gamma^\mu, \gamma^5\} = 0$
- c) $\gamma^{\mu\dagger}\gamma^0 = \gamma^0\gamma^\mu$

3 pt

2. Dirac Equation

The Dirac equation for the particle spinor $u(p)$ is:

$$(\gamma^\mu p_\mu - m) u(p) = 0$$

- a) Defining $\not{p} \equiv \gamma^\mu p_\mu$, write the equivalent equation for the antiparticle spinor $v(p)$.

1 pt

- b) Write the Dirac equation for the adjoint of the particle spinor, $\bar{u}(p)$ (which is defined as $\bar{u}(p) = u^\dagger(p)\gamma^0$), and for the antiparticle spinor $\bar{v}(p)$.

[Hint : Use the third property of the Dirac matrices listed in the previous exercise]

2 pt

- c) Verify the orthogonality of particle and antiparticle spinors:

$$u^{\dagger(1)}u^{(2)} = 0, \quad v^{\dagger(1)}v^{(2)} = 0$$

2 pt

- d) Show that:

$$\bar{u}u = 2mc, \quad \bar{v}v = -2mc$$

2 pt

- e) Show that they are complete:

$$\sum_{s=1,2} u^{(s)}\bar{u}^{(s)} = (\gamma^\mu p_\mu + mc), \quad \sum_{s=1,2} v^{(s)}\bar{v}^{(s)} = (\gamma^\mu p_\mu - mc)$$

4 pt

Considering that $u^{(s)}$ and $v^{(s)}$ are the canonical solutions to the Dirac equation.

3. Conservation of charge

Show how the conservation of charge can be deduced from the continuity equation:

$$\partial_\mu J^\mu = 0$$

2 pt

4. Coulomb Scattering

Consider the case of Coulomb scattering of a charged spin-0 particle from an external field A , which is a static field of a point charge Ze located in the origin:

$$A_{\mu} = (V, \vec{A}) = (V, 0)$$

The potential is given by:

$$V(x) = \frac{Ze}{4\pi |\mathbf{x}|}$$

Compute:

a) The transition amplitude T_{fi} for this process:

$$T_{fi} = -i \int d^4x j_f^\mu A_\mu$$

Hint : Make use of Fourier Transformation

$$\frac{1}{|\vec{p}_f - \vec{p}_i|^2} = \int d^3x e^{i(\vec{p}_f - \vec{p}_i)\vec{x}} \frac{1}{4\pi |\vec{x}|}$$

3 pt

b) Its transition probability:

$$\omega_{fi} = \lim_{T \rightarrow \infty} \frac{|T_{fi}|^2}{T}$$

Hint : Assume that the interaction occurs during a time period T from $t = -T/2$ up to $t = +T/2$

3 pt

c) The cross section:

$$d\sigma = \frac{\omega_{fi}}{flux_i} dLIPS$$

3 pt

d) Derive the Rutherford scattering cross section (i.e. consider the non-relativistic limit of the cross section computed in the previous point)

1 pt