

# Lab 11

McKenzie Mai (A16646664)

2024-05-13

## Section 1. Proportion of G/G in a population

Downloaded a CSV file from Ensembl < [https://useast.ensembl.org/Homo\\_sapiens/Variation/Sample?db=core;r=17:39580100-40210101;v=rs8067378;vdb=variation;vf=959672880#373531\\_tablePanel](https://useast.ensembl.org/Homo_sapiens/Variation/Sample?db=core;r=17:39580100-40210101;v=rs8067378;vdb=variation;vf=959672880#373531_tablePanel)

Here we read this CSV file

```
mxl <- read.csv("373531-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")
head(mxl)
```

```
## Sample..Male.Female.Unknown. Genotype..forward.strand. Population.s. Father
## 1 NA19648 (F) A|A ALL, AMR, MXL -
## 2 NA19649 (M) G|G ALL, AMR, MXL -
## 3 NA19651 (F) A|A ALL, AMR, MXL -
## 4 NA19652 (M) G|G ALL, AMR, MXL -
## 5 NA19654 (F) G|G ALL, AMR, MXL -
## 6 NA19655 (M) A|G ALL, AMR, MXL -
## Mother
## 1 -
## 2 -
## 3 -
## 4 -
## 5 -
## 6 -
```

```
table(mxl$Genotype..forward.strand.)
```

```
##
## A|A A|G G|A G|G
## 22 21 12 9
```

```
table(mxl$Genotype..forward.strand.)/nrow(mxl) * 100
```

```
##
## A|A A|G G|A G|G
## 34.3750 32.8125 18.7500 14.0625
```

Now let's look at a different population. I picked the GBR

```
gbr <- read.csv("373522-SampleGenotypes-Homo_sapiens_Variation_Sample_rs8067378.csv")
```

Find proportion of G|G

```
round(table(gbr$Genotype..forward.strand.) / nrow(gbr) * 100, 2)
```

```
##
## A|A A|G G|A G|G
```

```
## 25.27 18.68 26.37 29.67
```

This variant that is associated with childhood asthma is more frequent in the GBR population than the MKL population.

Let's now dig into this further.

## Section 4: Population Scale Analysis

One sample is obviously not enough to know what is happening in a population. You are interested in assessing genetic differences on a population scale. So, you processed about ~230 samples and did the normalization on a genome level. Now, you want to find whether there is any association of the 4 asthma-associated SNPs (rs8067378...) on ORMDL3 expression.

How many samples do we have?

```
expr <- read.table("rs8067378_ENSG00000172057.6.txt")
head(expr)
```

```
##      sample geno      exp
## 1 HG00367   A/G 28.96038
## 2 NA20768   A/G 20.24449
## 3 HG00361   A/A 31.32628
## 4 HG00135   A/A 34.11169
## 5 NA18870   G/G 18.25141
## 6 NA11993   A/A 32.89721
```

```
nrow(expr)
```

```
## [1] 462
```

Q13: Read this file into R and determine the sample size for each genotype and their corresponding median expression levels for each of these genotypes.

```
# sample size:
table(expr$geno)
```

```
##
## A/A A/G G/G
## 108 233 121
```

```
# median expression levels:
library(dplyr)
```

```
##
## Attaching package: 'dplyr'

## The following objects are masked from 'package:stats':
##
##      filter, lag

## The following objects are masked from 'package:base':
##
##      intersect, setdiff, setequal, union
```

```
expr %>%
  group_by(geno)%>%
  summarize(median_expression=median(exp))
```


```
## # A tibble: 3 x 2
##   geno median_expression
```

```
##   <chr>          <dbl>
## 1 A/A           31.2
## 2 A/G           25.1
## 3 G/G           20.1
```

```
library(ggplot2)
```

Lets make a boxplot

```
ggplot(expr) + aes(geno, exp, fill=geno) + geom_boxplot(notch=TRUE)
```



Q14: Generate a boxplot with a box per genotype, what could you infer from the relative expression value between A/A and G/G displayed in this plot? Does the SNP effect the expression of ORMDL3?

boxplot shown above; A/A is more expressed in comparison to G/G because it has a higher median expression value. yes, the SNP affects the expression because expression is seen to decreases as each allele changes.