

# Resampling Techniques and their Application

## -Class 7-

Frank Konietschke

Institut für Biometrie und Klinische Epidemiologie


Charité - Universitätsmedizin Berlin, Berlin

[frank.konietschke@charite.de](mailto:frank.konietschke@charite.de)



## Motivation and Examples-III

Researchers produce a pain killer using poison from a snake. They investigate the effect of the treatment on  $n_1$  mice in the **control** group and  $n_2 = 10$  mice in the **active treatment**. The response variable is the reaction time of the mice to signal pain when a stitch is applied to their tail. Is the treatment effective? (all mice survived the dose)



```
x = c(  
  2.4, 3.0, 3.0, 2.2, 2.2,  
  2.2, 2.2, 2.8, 2.0, 3.0)
```

```
y = c(  
  2.8, 2.2, 3.8, 9.4, 8.4,  
  3.0, 3.2, 4.4, 3.2, 7.4)
```

- **Aim:** Test  $H_0 : \mu_1 = \mu_2$  and confidence interval for  $\delta = \mu_1 - \mu_2$

# Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$

## Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$

## Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$

## Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$
- Estimators

# Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$
- Estimators
  - $\bar{X}_{1\cdot}$  and  $\bar{X}_{2\cdot}$ : means per group with

$$\bar{X}_{i\cdot} = \frac{1}{n_i} \sum_{k=1}^{n_i} X_{ik}$$

# Statistical Model

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$
- Estimators
  - $\bar{X}_{1\cdot}$  and  $\bar{X}_{2\cdot}$ : means per group with

$$\bar{X}_{i\cdot} = \frac{1}{n_i} \sum_{k=1}^{n_i} X_{ik}$$

- $\hat{\sigma}_1^2$  and  $\hat{\sigma}_2^2$ : empirical variances per group with

$$\hat{\sigma}_i^2 = \frac{1}{n_i - 1} \sum_{k=1}^{n_i} (X_{ik} - \bar{X}_{i\cdot})^2$$



# Satterthwaite-Welch t-Test

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$

# Satterthwaite-Welch t-Test

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$

# Satterthwaite-Welch t-Test

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$

# Satterthwaite-Welch t-Test

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$
- Test statistic

$$T = \frac{\bar{X}_{1\cdot} - \bar{X}_{2\cdot}}{\sqrt{\hat{\sigma}_1^2/n_1 + \hat{\sigma}_2^2/n_2}}$$

# Satterthwaite-Welch t-Test

- $X_{ik} \sim F_i, i = 1, 2; k = 1, \dots, n_i; N = n_1 + n_2$ 
  - $E(X_{i1}) = \mu_i; \text{Var}(X_{i1}) = \sigma_i^2$
  - Asymptotics:  $N \rightarrow \infty : n_i/N \rightarrow \kappa_i \in (0, 1)$

- Test statistic


$$T = \frac{\bar{X}_{1\cdot} - \bar{X}_{2\cdot}}{\sqrt{\hat{\sigma}_1^2/n_1 + \hat{\sigma}_2^2/n_2}}$$

- Reject  $H_0$ , if  $|T| \geq t_{1-\alpha/2}(\nu)$ ,

$$\nu = \frac{\left(\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}\right)^2}{\frac{\hat{\sigma}_1^4}{n_1^2(n_1-1)} + \frac{\hat{\sigma}_2^4}{n_2^2(n_2-1)}}$$

degrees of freedom (**Satterthwaite's approximation**).

Researchers produce pain killer using poison from a cobra. They investigate the effect of the treatment on  $n_1$  mice in the **control** group and  $n_2 = 10$  mice in the **active treatment**. The response variable is the reaction time of the mice to signal pain when a stitch is applied to their tail. Is the treatment effective?



```
react <- data.frame(resp=c(x,y),  
  grp=factor(c(rep(1,10),rep(2,10))))
```

```
t.test(resp~grp,data=react,  
  var.equal=TRUE)
```

```
t.test(resp~grp,data=react,  
  var.equal=FALSE)
```

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling
  - Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$



# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling
  - Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
    - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling
  - Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
    - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
    - $X_{11}^*, \dots, X_{1n_1}^*$ : group 1

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling
  - Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
    - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
    - $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
    - $X_{21}^*, \dots, X_{2n_2}^*$ : group 2

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling
  - Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
    - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
    - $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
    - $X_{21}^*, \dots, X_{2n_2}^*$ : group 2
    - $\bar{X}_{1\cdot}^*$  and  $\bar{X}_{2\cdot}^*$ : means

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling

- Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
  - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
  - $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
  - $X_{21}^*, \dots, X_{2n_2}^*$ : group 2
  - $\bar{X}_{1.}^*$  and  $\bar{X}_{2.}^*$ : means
  - $\hat{\sigma}_1^{2*}$  and  $\hat{\sigma}_2^{2*}$ : empirical variances

$$T^* = \frac{\bar{X}_{1.}^* - \bar{X}_{2.}^* - E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})}{\sqrt{\hat{\sigma}_1^{2*}/n_1 + \hat{\sigma}_2^{2*}/n_2}}$$

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling

- Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$ 
  - Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
  - $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
  - $X_{21}^*, \dots, X_{2n_2}^*$ : group 2
  - $\bar{X}_{1\cdot}^*$  and  $\bar{X}_{2\cdot}^*$ : means
  - $\hat{\sigma}_1^{2*}$  and  $\hat{\sigma}_2^{2*}$ : empirical variances

$$T^* = \frac{\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* - E(\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* | \mathbf{X})}{\sqrt{\hat{\sigma}_1^{2*}/n_1 + \hat{\sigma}_2^{2*}/n_2}}$$

- Repeat these steps  $nboot$ -times

# Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling

- Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$

- Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
- $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
- $X_{21}^*, \dots, X_{2n_2}^*$ : group 2
- $\bar{X}_{1\cdot}^*$  and  $\bar{X}_{2\cdot}^*$ : means
- $\hat{\sigma}_1^{2*}$  and  $\hat{\sigma}_2^{2*}$ : empirical variances

$$T^* = \frac{\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* - E(\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* | \mathbf{X})}{\sqrt{\hat{\sigma}_1^{2*}/n_1 + \hat{\sigma}_2^{2*}/n_2}}$$

- Repeat these steps  $nboot$ -times
- Reject  $H_0$ , if  $T < c_{\alpha/2}^*$  or  $T > c_{1-\alpha/2}^*$

## Resampling the $t$ -Test

- Goal: estimate the distribution of  $T$  via resampling

- Data:  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})'$

- Resampling variables:  $\mathbf{X}^* = (X_{11}^*, \dots, X_{2n_2}^*)'$
- $X_{11}^*, \dots, X_{1n_1}^*$ : group 1
- $X_{21}^*, \dots, X_{2n_2}^*$ : group 2
- $\bar{X}_{1\cdot}^*$  and  $\bar{X}_{2\cdot}^*$ : means
- $\hat{\sigma}_1^{2*}$  and  $\hat{\sigma}_2^{2*}$ : empirical variances

$$T^* = \frac{\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* - E(\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* | \mathbf{X})}{\sqrt{\hat{\sigma}_1^{2*}/n_1 + \hat{\sigma}_2^{2*}/n_2}}$$

- Repeat these steps  $nboot$ -times
- Reject  $H_0$ , if  $T < c_{\alpha/2}^*$  or  $T > c_{1-\alpha/2}^*$
- $c_{\alpha}^*$ :  $\alpha$ -quantile from resampling distribution



# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (**fixed values**)

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (**fixed values**)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (**fixed values**)

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$ 
  - $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$
  - $\mathbf{X}_1^* = (1, 1, 2, 3, 3)$

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$ 
  - $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$
  - $\mathbf{X}_1^* = (1, 1, 2, 3, 3)$
  - $\mathbf{X}_1^* = (2, 5, 5, 3, 3)$

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$ 
  - $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$
  - $\mathbf{X}_1^* = (1, 1, 2, 3, 3)$
  - $\mathbf{X}_1^* = (2, 5, 5, 3, 3)$
  - ...



# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$ 
  - $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$
  - $\mathbf{X}_1^* = (1, 1, 2, 3, 3)$
  - $\mathbf{X}_1^* = (2, 5, 5, 3, 3)$
  - ...
- In R: `sample(x1,replace=TRUE)`

# Group wise Nonparametric Bootstrap

- $\mathbf{X}_1 = (X_{11}, \dots, X_{1n_1})$  (fixed values)
- $\mathbf{X}_2 = (X_{21}, \dots, X_{2n_2})$  (fixed values)
- **Drawing with Replacement:** randomly draw  $n_1$  and  $n_2$  observations from  $\mathbf{X}_1$  and  $\mathbf{X}_2$
- Example  $\mathbf{X}_1 = (1, 2, 3, 4, 5) \Rightarrow$ 
  - $\mathbf{X}_1^* = (2, 2, 4, 3, 2)$
  - $\mathbf{X}_1^* = (1, 1, 2, 3, 3)$
  - $\mathbf{X}_1^* = (2, 5, 5, 3, 3)$
  - ...
- In R: `sample(x1,replace=TRUE)`
- Also known as **Group wise Nonparametric Bootstrap**

# Nonparametric Bootstrap

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**

# Nonparametric Bootstrap

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing with Replacement:** randomly draw  $N$  observations  $X_k^*$  from  $\mathbf{X}$  with replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

# Nonparametric Bootstrap

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing with Replacement:** randomly draw  $N$  observations  $X_k^*$  from  $\mathbf{X}$  with replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- In R: `sample(x,replace=TRUE)`

# Nonparametric Bootstrap

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing with Replacement:** randomly draw  $N$  observations  $X_k^*$  from  $\mathbf{X}$  with replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- In R: `sample(x,replace=TRUE)`
- Also known as **Nonparametric Bootstrap**

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$



# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}^* = (4, 1, 3, 2, 5)$

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}^* = (4, 1, 3, 2, 5)$   
 $\mathbf{X}^* = (5, 1, 2, 3, 4)$

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}^* = (4, 1, 3, 2, 5)$   
 $\mathbf{X}^* = (5, 1, 2, 3, 4)$   
 $\mathbf{X}^* = (3, 1, 2, 5, 4)$

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}^* = (4, 1, 3, 2, 5)$   
 $\mathbf{X}^* = (5, 1, 2, 3, 4)$   
 $\mathbf{X}^* = (3, 1, 2, 5, 4)$   
...

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
     $\mathbf{X}^* = (4, 1, 3, 2, 5)$   
     $\mathbf{X}^* = (5, 1, 2, 3, 4)$   
     $\mathbf{X}^* = (3, 1, 2, 5, 4)$   
    ...  
• In R: `sample(x)`

# Permutation

- **Data  $\mathbf{X} = (X_{11}, \dots, X_{2n_2})$  (fixed values)**
- **Drawing without Replacement:** randomly draw  $N$  observations  $X_{ik}^*$  from  $\mathbf{X}$  without replacement such that

$$P(X_{11}^* = X_{11}) = \frac{1}{N}$$

- Example  $\mathbf{X} = (1, 2, 3, 4, 5) \Rightarrow$   
 $\mathbf{X}^* = (4, 1, 3, 2, 5)$   
 $\mathbf{X}^* = (5, 1, 2, 3, 4)$   
 $\mathbf{X}^* = (3, 1, 2, 5, 4)$   
...
- In R: `sample(x)`
- Also known as **Permutation**

# Parametric Bootstrap

- **Data  $\mathbf{X}_i = (X_{ik}, \dots, X_{in_i})$  (fixed values)**



# Parametric Bootstrap

- **Data**  $\mathbf{X}_i = (X_{ik}, \dots, X_{in_i})$  (**fixed values**)
- **Resampling** randomly draw  $n_i$  observations  $X_{ik}^*$  from

$$N(0, \hat{\sigma}_i^2)$$

# Parametric Bootstrap

- **Data**  $\mathbf{X}_i = (X_{ik}, \dots, X_{in_i})$  (**fixed values**)
- **Resampling** randomly draw  $n_i$  observations  $X_{ik}^*$  from

$$N(0, \hat{\sigma}_i^2)$$

- In R: `rnorm(n, 0, sd(x))`

# Parametric Bootstrap

- **Data**  $\mathbf{X}_i = (X_{ik}, \dots, X_{in_i})$  (**fixed values**)
- **Resampling** randomly draw  $n_i$  observations  $X_{ik}^*$  from

$$N(0, \hat{\sigma}_i^2)$$

- In R: `rnorm(n, 0, sd(x))`
- Also known as **Parametric Bootstrap** (Why is that not equivalent to the t-approximation?)

# Skewed Parametric Bootstrap

- Data  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (fixed values)

# Skewed Parametric Bootstrap

- **Data  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (fixed values)**
- Estimate the skewness of each sample by

$$\hat{\mu}_{i,3} = \frac{n_i}{(n_i - 1)(n_i - 2)} \sum_{k=1}^{n_i} \left( \frac{X_{ik} - \bar{X}_{i\cdot}}{\hat{\sigma}_i} \right)^3$$

# Skewed Parametric Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Estimate the skewness of each sample by

$$\hat{\mu}_{i,3} = \frac{n_i}{(n_i - 1)(n_i - 2)} \sum_{k=1}^{n_i} \left( \frac{X_{ik} - \bar{X}_{i\cdot}}{\hat{\sigma}_i} \right)^3$$

- **Resampling** randomly draw  $n_i$  observations  $X_{ik}^*$  from

$$\text{sign}(\hat{\mu}_{i,3}) \hat{\sigma}_i \frac{\chi_{f_i}^2 - f_i}{\sqrt{2f_i}}$$

# Skewed Parametric Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Estimate the skewness of each sample by

$$\hat{\mu}_{i,3} = \frac{n_i}{(n_i - 1)(n_i - 2)} \sum_{k=1}^{n_i} \left( \frac{X_{ik} - \bar{X}_{i\cdot}}{\hat{\sigma}_i} \right)^3$$

- **Resampling** randomly draw  $n_i$  observations  $X_{ik}^*$  from

$$\text{sign}(\hat{\mu}_{i,3}) \hat{\sigma}_i \frac{\chi_{f_i}^2 - f_i}{\sqrt{2f_i}}$$

- $f_i = 8/\hat{\mu}_{i,3}^2$

# Wild Bootstrap

- **Data  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (fixed values)**



# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .

# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .
- **Resampling** randomly generate iid weights  $W_{ik}$  with  $E(W_{ik}) = 0$  and  $Var(W_{ik}) = 1$ . Generate  $X_{ik}^*$  by

$$X_{ik}^* = W_{ik} * Z_{ik}$$

# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .
- **Resampling** randomly generate iid weights  $W_{ik}$  with  $E(W_{ik}) = 0$  and  $\text{Var}(W_{ik}) = 1$ . Generate  $X_{ik}^*$  by

$$X_{ik}^* = W_{ik} * Z_{ik}$$

- Examples:  $W_{ik} \sim N(0, 1)$

# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .
- **Resampling** randomly generate iid weights  $W_{ik}$  with  $E(W_{ik}) = 0$  and  $\text{Var}(W_{ik}) = 1$ . Generate  $X_{ik}^*$  by

$$X_{ik}^* = W_{ik} * Z_{ik}$$

- Examples:  $W_{ik} \sim N(0, 1)$
- Rademacher:  $P(W_{ik} = 1) = P(W_{ik} = -1) = 1/2$

# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .
- **Resampling** randomly generate iid weights  $W_{ik}$  with  $E(W_{ik}) = 0$  and  $\text{Var}(W_{ik}) = 1$ . Generate  $X_{ik}^*$  by

$$X_{ik}^* = W_{ik} * Z_{ik}$$

- Examples:  $W_{ik} \sim N(0, 1)$
- Rademacher:  $P(W_{ik} = 1) = P(W_{ik} = -1) = 1/2$
- ...

# Wild Bootstrap

- **Data**  $\mathbf{X}_i = (X_{i1}, \dots, X_{in_i})$  (**fixed values**)
- Fix the values  $Z_{ik} = X_{ik} - \bar{X}_i$ .
- **Resampling** randomly generate iid weights  $W_{ik}$  with  $E(W_{ik}) = 0$  and  $\text{Var}(W_{ik}) = 1$ . Generate  $X_{ik}^*$  by

$$X_{ik}^* = W_{ik} * Z_{ik}$$

- Examples:  $W_{ik} \sim N(0, 1)$
- Rademacher:  $P(W_{ik} = 1) = P(W_{ik} = -1) = 1/2$
- ...
- Also known as **Wild-Bootstrap**

## Computation of $E(\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1\cdot}^* - \bar{X}_{2\cdot}^* | \mathbf{X})$  in the following cases

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap



## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation
- Compute  $Var(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation
- Compute  $Var(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation
- Compute  $Var(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap

## Computation of $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$

- Compute  $E(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation
- Compute  $Var(\bar{X}_{1.}^* - \bar{X}_{2.}^* | \mathbf{X})$  in the following cases
  - Group-wise nonparametric Bootstrap
  - Nonparametric Bootstrap
  - Permutation

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit



# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words
    - Resampling dist. mimics the distribution of  $T$  under  $H_0$

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words
    - Resampling dist. mimics the distribution of  $T$  under  $H_0$
    - The dist. of  $T$  departs from the resampling dist. under  $H_1$

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words
    - Resampling dist. mimics the distribution of  $T$  under  $H_0$
    - The dist. of  $T$  departs from the resampling dist. under  $H_1$
- References

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words
    - Resampling dist. mimics the distribution of  $T$  under  $H_0$
    - The dist. of  $T$  departs from the resampling dist. under  $H_1$
- References
  - Janssen (1997, 2005), Janssen and Pauls (2003)

# When do Resampling Tests Work?

- Limit distribution of  $T$ :  $N(0, 1)$  (Note that the  $t_{\nu}$  distribution is the  $N(0, 1)$  for large  $n$ )
- Limit Distribution of  $T^*$  given  $\mathbf{X}$ :  $N(0, 1)$ 
  - Both distributions coincide and have the same limit
  - Therefore, the resampling test will work (at least for large sample sizes)
  - What means 'large'?  $\hookrightarrow$  simulations necessary
  - In words
    - Resampling dist. mimics the distribution of  $T$  under  $H_0$
    - The dist. of  $T$  departs from the resampling dist. under  $H_1$
- References
  - Janssen (1997, 2005), Janssen and Pauls (2003)
  - Konietzschke and Pauly (2012 a, b)



# Confidence Intervals

- $(1 - \alpha)$  confidence interval for  $\delta = \mu_1 - \mu_2$
- Confidence intervals require the distribution under the alternative hypothesis

# Confidence Intervals

- $(1 - \alpha)$  confidence interval for  $\delta = \mu_1 - \mu_2$
- Confidence intervals require the distribution under the alternative hypothesis
- Computation of confidence interval for  $\delta$  is based on inverting

$$T = \frac{\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)}{\sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}}$$

# Confidence Intervals

- $(1 - \alpha)$  confidence interval for  $\delta = \mu_1 - \mu_2$
- Confidence intervals require the distribution under the alternative hypothesis
- Computation of confidence interval for  $\delta$  is based on inverting

$$T = \frac{\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)}{\sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}}$$

- For any  $\mu_1 - \mu_2$ , we have  $T \sim t_\nu$  (or  $N(0, 1)$  for large  $n$ )

# Confidence Intervals

- $(1 - \alpha)$  confidence interval for  $\delta = \mu_1 - \mu_2$
- Confidence intervals require the distribution under the alternative hypothesis
- Computation of confidence interval for  $\delta$  is based on inverting

$$T = \frac{\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)}{\sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}}$$

- For any  $\mu_1 - \mu_2$ , we have  $T \sim t_\nu$  (or  $N(0, 1)$  for large  $n$ )

$$P(t_{\alpha/2} \leq T \leq t_{1-\alpha/2}) = 1 - \alpha$$

$$CI = \left[ \bar{X}_1 - \bar{X}_2 \pm t_{\nu, (1-\alpha/2)} \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}} \right]$$

# Confidence Intervals

- Interpretation
  - A confidence interval is an estimator of  $\mu_1 - \mu_2$
  - We estimate the difference with  $(1 - \alpha)$  confidence
  - They should be compatible with the test result
  - It is false to say that  $CI$  covers  $\delta$  with  $(1 - \alpha)100\%$  probability (only holds for random prior observation)

## Resampling Based Confidence Intervals

- Can we use the resampling distribution to compute confidence intervals?

## Resampling Based Confidence Intervals

- Can we use the resampling distribution to compute confidence intervals?
- Did we ever assume that  $H_0$  holds when we computed the resampling distribution? **NO**

## Resampling Based Confidence Intervals

- Can we use the resampling distribution to compute confidence intervals?
- Did we ever assume that  $H_0$  holds when we computed the resampling distribution? **NO**
- Illustrate with 2 samples with large effect



# Resampling Based Confidence Intervals



- Can we use the resampling distribution to compute confidence intervals?
- Did we ever assume that  $H_0$  holds when we computed the resampling distribution? **NO**
- Illustrate with 2 samples with large effect

```
set.seed(1)
n1<-15;n2<-15; N<-n1+n2
x<-rnorm(n1,0)
y<-rnorm(n2,10)
boxplot(x,y)
erg<-c()
for(i in 1:100000){
  xx<-sample(c(x,y))
  xx1<-xx[1:n1]
  xx2<-xx[(n1+1):(N)]
  mxx1<-mean(xx1); mxx2<-mean(xx2)
  vxx1<-var(xx1); vxx2<-var(xx2)
  Tstar<-(mxx1-mxx2)/sqrt(vxx1/n1 + vxx2/n2)
  erg[i]<-Tstar}
hist(erg,freq=F)
```

# Resampling Based Confidence Intervals

- Can we use the resampling distribution to compute confidence intervals?
- Did we ever assume that  $H_0$  holds when we computed the resampling distribution? **NO**
- Illustrate with 2 samples with large effect

```
set.seed(1)
n1<-15;n2<-15; N<-n1+n2
x<-rnorm(n1,0)
y<-rnorm(n2,10)
boxplot(x,y)
erg<-c()
for(i in 1:100000){
  xx<-sample(c(x,y))
  xx1<-xx[1:n1]
  xx2<-xx[(n1+1):(N)]
  mxx1<-mean(xx1); mxx2<-mean(xx2)
  vxx1<-var(xx1); vxx2<-var(xx2)
  Tstar<-(mxx1-mxx2)/sqrt(vxx1/n1 + vxx2/n2)
  erg[i]<-Tstar}
hist(erg,freq=F)
```



# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide

# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide
- We can use the distribution of  $T^*$  for the computation of  $(1 - \alpha)100\%$ - confidence intervals

# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide
- We can use the distribution of  $T^*$  for the computation of  $(1 - \alpha)100\%$ - confidence intervals

$$P(c_{\alpha/2}^* \leq T \leq c_{1-\alpha/2}^*) \approx 1 - \alpha$$
$$CI_p = \left[ \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{1-\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}; \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}} \right]$$

# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide
- We can use the distribution of  $T^*$  for the computation of  $(1 - \alpha)100\%$ - confidence intervals

$$P(c_{\alpha/2}^* \leq T \leq c_{1-\alpha/2}^*) \approx 1 - \alpha$$
$$CI_p = \left[ \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{1-\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}; \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}} \right]$$

- Studentization „deletes“ the shift; dist. is invariant

# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide
- We can use the distribution of  $T^*$  for the computation of  $(1 - \alpha)100\%$ - confidence intervals

$$P(c_{\alpha/2}^* \leq T \leq c_{1-\alpha/2}^*) \approx 1 - \alpha$$
$$CI_p = \left[ \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{1-\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}; \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}} \right]$$

- Studentization „deletes“ the shift; dist. is invariant
- References

# Resampling Based Confidence Intervals

- Both the distribution of  $T$  and its resampling distribution coincide
- We can use the distribution of  $T^*$  for the computation of  $(1 - \alpha)100\%$ - confidence intervals


$$P(c_{\alpha/2}^* \leq T \leq c_{1-\alpha/2}^*) \approx 1 - \alpha$$
$$CI_p = \left[ \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{1-\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}}; \bar{X}_{1\cdot} - \bar{X}_{2\cdot} - c_{\alpha/2}^* \cdot \sqrt{\frac{\hat{\sigma}_1^2}{n_1} + \frac{\hat{\sigma}_2^2}{n_2}} \right]$$

- Studentization „deletes“ the shift; dist. is invariant
- References
  - Pauly, Asendorf and Konietzschke (2016)



## Confidence Intervals - II


- $X_{11}, \dots, X_{1n_1} \sim \text{Exp}(1); X_{21}, \dots, X_{2n_2} \sim \text{Exp}(1) + \delta$
- $n_{\text{sim}} = n_{\text{perm}} = 10,000$



```

x=c(2.4, 3.0, 3.0, 2.2, 2.2,
2.2, 2.2, 2.8, 2.0, 3.0)
y=c(2.8, 2.2, 3.8, 9.4, 8.4,
3.0, 3.2, 4.4, 3.2, 7.4)
n1<-length(x)
n2<-length(y)
N<-n1+n2
mx<-mean(x);my<-mean(y)
vx<-var(x);vy<-var(y)
T<-(mx-my)/sqrt(vx/n1 + vy/n2)
erg<-c()
for(i in 1:100000){
xx<-sample(c(x,y))
xx1<-xx[1:n1]
xx2<-xx[(n1+1):(N)]
mxx1<-mean(xx1); mxx2<-mean(xx2)
vxx1<-var(xx1); vxx2<-var(xx2)
Tstar<-(mxx1-mxx2)/sqrt(vxx1/n1 + vxx2/n2)
erg[i]<-Tstar}
hist(erg)
c1star<-quantile(erg,0.025)
c2star<-quantile(erg,0.975)

```



# Implementation

- Either in the same as in the 1-sample case, or
- Writing

$$\bar{X}_{1.} - \bar{X}_{2.} = \sum_{\ell=1}^N c_{\ell} X_{\ell}$$

- Permutation version

$$\bar{X}_{1.}^* - \bar{X}_{2.}^* = \sum_{\ell=1}^N c_{\ell} X_{\ell}^* = \sum_{\ell=1}^N c_{\ell}^* X_{\ell}$$

- Permute the “coefficients”  $c_{\ell}$
- Same strategy for the variance estimator
- Example next slide

```

myPermuCI<-function(nsim,nperm,n1,n2,v1,v2,delta, Distribution){
  PermCI=c()
  N<-n1+n2
  #-----Data Generation-----#
  vvec = sqrt(c(rep(v1,n1),rep(v2,n2)))
  if (Distribution == "Normal"){
    x1=matrix(rnorm(n1*nsim,delta)*sqrt(v1),ncol=nsim)
    x2=matrix(rnorm(n2*nsim)*sqrt(v2),ncol=nsim)}
  xy = rbind(x1,x2)
  x12 = x1^2; x22=x2^2
  mx = colMeans(x1); my = colMeans(x2)
  vx = (colSums(x12)-n1*mx^2)/(n1-1)
  vy = (colSums(x22)-n2*my^2)/(n2-1)
  df=(vx/n1+vy/n2)^2/(vx^2/(n1^2*(n1-1))+vy^2/(n2^2*(n2-1)))
  T.L <-mx-my-qt(0.975,df)*sqrt(vx/n1+vy/n2)
  T.U <-mx-my+qt(0.975,df)*sqrt(vx/n1+vy/n2)
  #-----Permutation Matrices-----#
  P<-t(apply(matrix(1:N,nrow=nperm,ncol=N,byrow=TRUE),1,sample))
  #-----Helping Variables for Permutation Distribution---#
  i1<-c(rep(1/n1,n1),rep(0,n2))
  i2<-c(rep(0,n1),rep(1/n2,n2))
  i3<-c(rep(1/(n1*(n1-1)),n1), rep(0,n2))
  i4<-c(rep(0,n1), rep(1/(n2*(n2-1)),n2))
  Im1<-matrix(i1[P],nrow=nperm,ncol=N)
  Im2<-matrix(i2[P],nrow=nperm,ncol=N)
  Iv1<-matrix(i3[P],nrow=nperm,ncol=N)
  Iv2<-matrix(i4[P],nrow=nperm,ncol=N)

```

```

#-----Begin of Simulation-----#
for (i in 1:nsim){
X<-xy[,i]
#-----Permutations-----#
mxP <- Im1%%X
myP = Im2%%X
vxP <- Iv1%%X^2 - n1/(n1*(n1-1))*mxP^2
vyP <- Iv2%%X^2 - n2/(n2*(n2-1))*myP^2
TP = (mxP -myP )/sqrt(vxP +vyP )
c1<-quantile(TP,0.025); c2<-quantile(TP,0.975)
lower <-mx[i]-my[i]-c2*sqrt(vx[i]/n1+vy[i]/n2)
upper <- mx[i]-my[i]-c1*sqrt(vx[i]/n1+vy[i]/n2)
PermCI[i]<-(lower<delta& upper >delta)
#-----End of Simulation-----#
}
Result <- data.frame(nsim=nsim,nperm=nperm,delta=delta,
n1=n1,n2=n2,v1=v1,v2=v2, SW=mean(T.L <delta & T.U >delta),
PermCI=mean(PermCI),
distribution=Distribution)
print(Result)
#-----End of Function-----#
}
myPermuCI(1000,1000,10,20,1,3,1,"Normal")

```

# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$

# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$
- Investigate normal and exponential distributions

# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$
- Investigate normal and exponential distributions
- Use  $\sigma_i^2 = 1$  under all settings and varying  $n_i \in \{10, 20, 30\}$



# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$
- Investigate normal and exponential distributions
- Use  $\sigma_j^2 = 1$  under all settings and varying  $n_i \in \{10, 20, 30\}$
- Use  $n_{sim} = 10,000$  and  $n_{perm} = 10,000$  permutation runs

# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$
- Investigate normal and exponential distributions
- Use  $\sigma_j^2 = 1$  under all settings and varying  $n_i \in \{10, 20, 30\}$
- Use  $n_{sim} = 10,000$  and  $n_{perm} = 10,000$  permutation runs
- Instead of using permutations, would a wild-bootstrap approach also be possible?

# Project

- Simulate the coverage probabilities of the 95%-confidence intervals  $CI$  and  $CI_p$  for  $\delta \in \{0, 0.1, \dots, 2\}$
- Investigate normal and exponential distributions
- Use  $\sigma_i^2 = 1$  under all settings and varying  $n_i \in \{10, 20, 30\}$
- Use  $n_{sim} = 10,000$  and  $n_{perm} = 10,000$  permutation runs
- Instead of using permutations, would a wild-bootstrap approach also be possible?
  - Compute  $(1 - \alpha)$ - confidence intervals for  $\mu$  (one-sample problem) and  $\delta = \mu_1 - \mu_2$  using a wild-bootstrap approach. Provide a detailed derivation (formula, no simulation)