

# Resampling Techniques and their Application

## -Class 13-

Frank Konietschke

Institut für Biometrie und Klinische Epidemiologie

Charité - Universitätsmedizin Berlin, Berlin

[frank.konietschke@charite.de](mailto:frank.konietschke@charite.de)



## Example: Yeast Study

- $n = 10$  medical yeast samples

## Example: Yeast Study


- $n = 10$  medical yeast samples
- Each sample grown on different growth mediums (increasing fertility)

## Example: Yeast Study

- $n = 10$  medical yeast samples
- Each sample grown on different growth mediums (increasing fertility)
- Response: protein concentration


## Example: Yeast Study

- $n = 10$  medical yeast samples
- Each sample grown on different growth mediums (increasing fertility)
- Response: protein concentration



## Example: Yeast Study

- $n = 10$  medical yeast samples
- Each sample grown on different growth mediums (increasing fertility)
- Response: protein concentration



ID	M1	M2	M3	M4	M5
1	6.16	6.04	7.21	7.23	6.22
2	4.28	5.42	7.44	6.23	6.03
3	5.26	5.72	7.40	7.02	5.87
4	6.11	6.65	7.44	7.09	6.62
5	6.15	6.67	7.79	5.62	5.80
6	5.33	7.50	8.23	7.38	6.42
7	5.47	6.82	7.94	7.01	6.57
8	6.10	6.57	8.73	7.90	6.90
9	7.39	5.44	7.50	7.32	5.36
10	7.07	7.51	8.88	7.70	7.85

# Statistical Model

- Statistical Model

# Statistical Model

- Statistical Model
  - $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$



# Statistical Model

- Statistical Model
  - $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$
  - $E(\mathbf{X}_k) = \boldsymbol{\mu} = (\mu_1, \dots, \mu_d)'$

# Statistical Model

- Statistical Model
  - $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$
  - $E(\mathbf{X}_k) = \boldsymbol{\mu} = (\mu_1, \dots, \mu_d)'$
  - Covariance matrix  $\mathbf{V} = \text{Cov}(\mathbf{X}_1)$

# Statistical Model

- Statistical Model
  - $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$
  - $E(\mathbf{X}_k) = \boldsymbol{\mu} = (\mu_1, \dots, \mu_d)'$
  - Covariance matrix  $\mathbf{V} = \text{Cov}(\mathbf{X}_1)$
  - No normal distribution

# Statistical Model

- Statistical Model

- $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$
- $E(\mathbf{X}_k) = \boldsymbol{\mu} = (\mu_1, \dots, \mu_d)'$
- Covariance matrix  $\mathbf{V} = \text{Cov}(\mathbf{X}_1)$
- No normal distribution

- Hypotheses  $H_0 : \mu_1 = \dots = \mu_d$

$$H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$$

# Statistical Model

- Statistical Model

- $\mathbf{X}_k = (X_{1k}, \dots, X_{dk})' \sim \mathbf{F}$
- $E(\mathbf{X}_k) = \boldsymbol{\mu} = (\mu_1, \dots, \mu_d)'$
- Covariance matrix  $\mathbf{V} = \text{Cov}(\mathbf{X}_1)$
- No normal distribution

- Hypotheses  $H_0 : \mu_1 = \dots = \mu_d$

$$H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$$

- $\mathbf{C}$ : Contrast matrix

# Point Estimators

- Means and covariance matrix

# Point Estimators

- Means and covariance matrix
  - $\bar{\mathbf{X}} = (\bar{X}_{1.}, \dots, \bar{X}_{d.})'$ : Means

# Point Estimators

- Means and covariance matrix
  - $\bar{\mathbf{X}} = (\bar{X}_1, \dots, \bar{X}_d)'$ : Means
  - Empirical covariance matrix

$$\hat{\mathbf{V}} = \frac{1}{n-1} \sum_{k=1}^n (\bar{\mathbf{X}}_k - \bar{\mathbf{X}})(\bar{\mathbf{X}}_k - \bar{\mathbf{X}})'$$

```
x=matrix(c(6.16, 6.04, 7.21, 7.23, 6.22,  
4.28, 5.42, 7.44, 6.23, 6.03,  
5.26, 5.72, 7.40, 7.02, 5.87,  
6.11, 6.65, 7.44, 7.09, 6.62,  
6.15, 6.67, 7.79, 5.62, 5.80,  
5.33, 7.50, 8.23, 7.38, 6.42,  
5.47, 6.82, 7.94, 7.01, 6.57,  
6.10, 6.57, 8.73, 7.90, 6.90,  
7.39, 5.44, 7.50, 7.32, 5.36,  
7.07, 7.51, 8.88, 7.70, 7.85),ncol=5,byrow=T)
```



# Point Estimators

- Means and covariance matrix
  - $\bar{\mathbf{X}} = (\bar{X}_{1.}, \dots, \bar{X}_{d.})'$ : Means
  - Empirical covariance matrix

$$\hat{\mathbf{V}} = \frac{1}{n-1} \sum_{k=1}^n (\bar{\mathbf{X}}_k - \bar{\mathbf{X}})(\bar{\mathbf{X}}_k - \bar{\mathbf{X}})'$$

```
x=matrix(c(6.16, 6.04, 7.21, 7.23, 6.22,  
4.28, 5.42, 7.44, 6.23, 6.03,  
5.26, 5.72, 7.40, 7.02, 5.87,  
6.11, 6.65, 7.44, 7.09, 6.62,  
6.15, 6.67, 7.79, 5.62, 5.80,  
5.33, 7.50, 8.23, 7.38, 6.42,  
5.47, 6.82, 7.94, 7.01, 6.57,  
6.10, 6.57, 8.73, 7.90, 6.90,  
7.39, 5.44, 7.50, 7.32, 5.36,  
7.07, 7.51, 8.88, 7.70, 7.85),ncol=5,byrow=T)
```

```
Xbar=colMeans(x)
```

```
Vhat=var(x)
```

# Test Statistics

- Multiple contrast test procedures (last classes)

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures
- Idea: If  $H_0 : \mu_1 = \mu_2 = \dots = \mu_a$  holds, then each  $\mu_\ell = \frac{1}{d} \sum_{j=1}^d \mu_j$

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures
- Idea: If  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  holds, then each  $\mu_\ell = \frac{1}{d} \sum_{j=1}^d \mu_j$
- So, contrast of interest is the *GrandMean* contrast

$$\mathbf{C} = \begin{pmatrix} \frac{d-1}{d} & -\frac{1}{d} & \dots & -\frac{1}{d} \\ -\frac{1}{d} & \frac{d-1}{d} & \dots & -\frac{1}{d} \\ \vdots & \vdots & \vdots & \vdots \\ -\frac{1}{d} & -\frac{1}{d} & \dots & \frac{d-1}{d} \end{pmatrix} = \mathbf{I} - \frac{1}{d} \mathbf{J}$$

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures
- Idea: If  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  holds, then each  $\mu_\ell = \frac{1}{d} \sum_{j=1}^d \mu_j$
- So, contrast of interest is the *GrandMean* contrast

$$\mathbf{C} = \begin{pmatrix} \frac{d-1}{d} & -\frac{1}{d} & \dots & -\frac{1}{d} \\ -\frac{1}{d} & \frac{d-1}{d} & \dots & -\frac{1}{d} \\ \vdots & \vdots & \vdots & \vdots \\ -\frac{1}{d} & -\frac{1}{d} & \dots & \frac{d-1}{d} \end{pmatrix} = \mathbf{I} - \frac{1}{d} \mathbf{J}$$

- $\mathbf{I} = \text{diag}(d)$  and  $\mathbf{J}$ :  $d \times d$  matrix of 1's

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures
- Idea: If  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  holds, then each  $\mu_\ell = \frac{1}{d} \sum_{j=1}^d \mu_j$
- So, contrast of interest is the *GrandMean* contrast

$$\mathbf{C} = \begin{pmatrix} \frac{d-1}{d} & -\frac{1}{d} & \dots & -\frac{1}{d} \\ -\frac{1}{d} & \frac{d-1}{d} & \dots & -\frac{1}{d} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{d} & -\frac{1}{d} & \dots & \frac{d-1}{d} \end{pmatrix} = \mathbf{I} - \frac{1}{d} \mathbf{J}$$

- $\mathbf{I} = \text{diag}(d)$  and  $\mathbf{J}$ :  $d \times d$  matrix of 1's
- Note that  $\mathbf{c}'_\ell \boldsymbol{\mu} = \mu_\ell - \frac{1}{d} \sum_{j=1}^d \mu_j$

# Test Statistics

- Multiple contrast test procedures (last classes)
- In addition today: global testing procedures
- Idea: If  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  holds, then each  $\mu_\ell = \frac{1}{d} \sum_{j=1}^d \mu_j$
- So, contrast of interest is the *GrandMean* contrast

$$\mathbf{C} = \begin{pmatrix} \frac{d-1}{d} & -\frac{1}{d} & \dots & -\frac{1}{d} \\ -\frac{1}{d} & \frac{d-1}{d} & \dots & -\frac{1}{d} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{d} & -\frac{1}{d} & \dots & \frac{d-1}{d} \end{pmatrix} = \mathbf{I} - \frac{1}{d} \mathbf{J}$$

- $\mathbf{I} = \text{diag}(d)$  and  $\mathbf{J}$ :  $d \times d$  matrix of 1's
- Note that  $\mathbf{c}'_\ell \boldsymbol{\mu} = \mu_\ell - \frac{1}{d} \sum_{j=1}^d \mu_j$
- $\mathbf{C}$  is also known as centering matrix



# Test Statistics: Wald-Type

- Wald-Type Statistics

# Test Statistics: Wald-Type

- Wald-Type Statistics

- $W = n(\mathbf{c}\bar{\mathbf{X}}.)' [\mathbf{c}\hat{\mathbf{V}}\mathbf{c}']^+ \mathbf{c}\bar{\mathbf{X}}.$

# Test Statistics: Wald-Type

- Wald-Type Statistics

- $W = n(\mathbf{c}\bar{\mathbf{X}}.)' [\mathbf{c}\hat{\mathbf{V}}\mathbf{c}']^+ \mathbf{c}\bar{\mathbf{X}}.$
- $[\mathbf{A}]^+$ : generalized inverse of a matrix

# Test Statistics: Wald-Type

- Wald-Type Statistics

- $W = n(\mathbf{c}\bar{\mathbf{X}}.)' [\mathbf{c}\hat{\mathbf{V}}\mathbf{c}']^+ \mathbf{c}\bar{\mathbf{X}}.$
- $[\mathbf{A}]^+$ : generalized inverse of a matrix
- Under  $H_0$ , WTS has a  $\chi^2_{rank(\mathbf{c})}$  distribution

# Test Statistics: Wald-Type

- Wald-Type Statistics

- $W = n(\mathbf{c}\bar{\mathbf{X}}.)' [\mathbf{c}\hat{\mathbf{V}}\mathbf{c}']^+ \mathbf{c}\bar{\mathbf{X}}.$
- $[\mathbf{A}]^+$ : generalized inverse of a matrix
- Under  $H_0$ , WTS has a  $\chi^2_{rank(\mathbf{c})}$  distribution
- When do we reject the hypothesis?

# Test Statistics: Wald-Type

- Wald-Type Statistics

- $W = n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\hat{\mathbf{V}}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}.$
- $[\mathbf{A}]^+$ : generalized inverse of a matrix
- Under  $H_0$ , WTS has a  $\chi^2_{rank(\mathbf{C})}$  distribution
- When do we reject the hypothesis?

```
library(multcomp)
library(MASS)
C=contrMat(n=rep(10,5),"GrandMean")
CX=C%*%Xbar
CVhat = C%*%Vhat%*%t(C)
W=n*t(CX)%*%ginv(CVhat)%*%CX
pvalue= 1-pchisq(W, d-1)
```

# Wald-Type Test: Properties

## Advantages

- 
- 
- 
- 

## Disadvantages

- 
- 
- 
-

## ANOVA-Type Statistic

- $\hat{V}$  causes issues in the WTS, let us remove it

$$\begin{aligned} A_1 &= n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\ &= n\bar{\mathbf{X}}.' \mathbf{C}' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\ &= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}. \end{aligned}$$



## ANOVA-Type Statistic

- $\hat{V}$  causes issues in the WTS, let us remove it

$$\begin{aligned}A_1 &= n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{C}' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}.\end{aligned}$$

- The final test is given by

$$\begin{aligned}A &= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}. / \text{Trace}(\mathbf{T}\hat{\mathbf{V}}), \\f &= \frac{\text{Trace}(\mathbf{T}\hat{\mathbf{V}})^2}{\text{Trace}(\mathbf{T}\hat{\mathbf{V}}\mathbf{T}\hat{\mathbf{V}})}\end{aligned}$$

## ANOVA-Type Statistic

- $\hat{\mathbf{V}}$  causes issues in the WTS, let us remove it

$$\begin{aligned}A_1 &= n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{C}' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}.\end{aligned}$$

- The final test is given by

$$\begin{aligned}A &= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}. / \text{Trace}(\mathbf{T}\hat{\mathbf{V}}), \\f &= \frac{\text{Trace}(\mathbf{T}\hat{\mathbf{V}})^2}{\text{Trace}(\mathbf{T}\hat{\mathbf{V}}\mathbf{T}\hat{\mathbf{V}})}\end{aligned}$$

- Under  $H_0$ ,  $ATS$  has a  $F(f, \infty)$  distribution

## ANOVA-Type Statistic

- $\hat{\mathbf{V}}$  causes issues in the WTS, let us remove it

$$\begin{aligned}A_1 &= n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{C}' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}.\end{aligned}$$

- The final test is given by

$$\begin{aligned}A &= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}. / \text{Trace}(\mathbf{T}\hat{\mathbf{V}}), \\f &= \frac{\text{Trace}(\mathbf{T}\hat{\mathbf{V}})^2}{\text{Trace}(\mathbf{T}\hat{\mathbf{V}}\mathbf{T}\hat{\mathbf{V}})}\end{aligned}$$

- Under  $H_0$ ,  $ATS$  has a  $F(f, \infty)$  distribution
- When do we reject the hypothesis?

## ANOVA-Type Statistic

- $\hat{V}$  causes issues in the WTS, let us remove it

$$\begin{aligned}A_1 &= n(\mathbf{C}\bar{\mathbf{X}}.)' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{C}' [\mathbf{C}\mathbf{C}']^+ \mathbf{C}\bar{\mathbf{X}}. \\&= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}.\end{aligned}$$

- The final test is given by

$$\begin{aligned}A &= n\bar{\mathbf{X}}.' \mathbf{T}\bar{\mathbf{X}}. / \text{Trace}(\mathbf{T}\hat{\mathbf{V}}), \\f &= \frac{\text{Trace}(\mathbf{T}\hat{\mathbf{V}})^2}{\text{Trace}(\mathbf{T}\hat{\mathbf{V}}\mathbf{T}\hat{\mathbf{V}})}\end{aligned}$$

- Under  $H_0$ ,  $ATS$  has a  $F(f, \infty)$  distribution
- When do we reject the hypothesis?

```
TT <- t(C)%*%ginv(C%*%t(C))%*%C
TrTV <-sum(c(diag(TT%*%Vhat)))
A <- n*t(Xbar)%*%TT%*%Xbar/TrTV
```

```
TVTV<-TT%*%Vhat%*%TT%*%Vhat
TrTVTV <-sum(c(diag(TVTV)))
f <- TrTV^2/TrTVTV
```

```
pvalue <- 1-pf(A,f,10^10)
```

```
#10^10=infty, arbitrary high nr#
```

# ANOVA-Type Test: Properties

## Advantages

- 
- 
- 
- 

## Disadvantages

- 
- 
- 
-

- R-package *MANOVA.RM*

```
data=data.frame(x=c(x),ID=rep(1:10,5),  
dose=sort(rep(1:5,10)))  
library(MANOVA.RM)
```

```
fit<-RM(x~dose, subject="ID", data=data)
```

```
summary(fit)
```

# MCTP: Motivation

- Is the global hypothesis always the question?

# MCTP: Motivation

- Is the global hypothesis always the question?
- Revise the example



# MCTP: Motivation

- Is the global hypothesis always the question?
- Revise the example
- Hypotheses

# MCTP: Motivation

- Is the global hypothesis always the question?
- Revise the example
- Hypotheses

- Contrast matrix  $\mathbf{C} = \begin{pmatrix} \mathbf{c}'_1 \\ \vdots \\ \mathbf{c}'_q \end{pmatrix} = \begin{pmatrix} c_{11} & \dots & c_{1d} \\ \vdots & \dots & \vdots \\ c_{1q} & \dots & c_{qd} \end{pmatrix}, (\mathbf{C}\mathbf{1} = \mathbf{0})$

# MCTP: Motivation

- Is the global hypothesis always the question?
- Revise the example
- Hypotheses

- Contrast matrix  $\mathbf{C} = \begin{pmatrix} \mathbf{c}'_1 \\ \vdots \\ \mathbf{c}'_q \end{pmatrix} = \begin{pmatrix} c_{11} & \dots & c_{1d} \\ \vdots & \dots & \vdots \\ c_{1q} & \dots & c_{qd} \end{pmatrix}, (\mathbf{C}\mathbf{1} = \mathbf{0})$
- $H_0^\ell : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  simultaneously

# MCTP: Motivation

- Is the global hypothesis always the question?
- Revise the example
- Hypotheses

- Contrast matrix  $\mathbf{C} = \begin{pmatrix} \mathbf{c}'_1 \\ \vdots \\ \mathbf{c}'_q \end{pmatrix} = \begin{pmatrix} c_{11} & \dots & c_{1d} \\ \vdots & \dots & \vdots \\ c_{1q} & \dots & c_{qd} \end{pmatrix}, (\mathbf{C}\mathbf{1} = \mathbf{0})$
- $H_0^\ell : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  simultaneously
- $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps
  1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$



# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$
- ...

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$
- ...

3. Confidence intervals for the effects

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$
- ...

3. Confidence intervals for the effects

- Regulatory authorities require confidence intervals (ICH E9)

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$
- ...

3. Confidence intervals for the effects

- Regulatory authorities require confidence intervals (ICH E9)

- Problems

# Why MCTP and SCI?

- ANOVA based evaluation: Three steps

1.  $H_0 : \mu_1 = \mu_2 = \dots = \mu_d$  (overall hypothesis)

(Using WTS or ATS)

2. Multiple Comparisons after rejecting the global hypothesis

- $H_0^{(1,2)} : \mu_1 = \mu_2$
- $H_0^{(1,3)} : \mu_1 = \mu_3$
- ...

3. Confidence intervals for the effects

- Regulatory authorities require confidence intervals (ICH E9)

- Problems

- Overall result and multiple comparisons may be incompatible

## Why MCTP and SCI?

- Better: Start with multiple comparisons (→ multiple contrast test)

## Why MCTP and SCI?

- Better: Start with multiple comparisons ( $\rightarrow$  multiple contrast test)
  - Control the multiple level  $\alpha$  in the strong sense

$$P(\text{reject at least one true } H_0^i)$$



# Why MCTP and SCI?

- Better: Start with multiple comparisons ( $\rightarrow$  multiple contrast test)
  - Control the multiple level  $\alpha$  in the strong sense

$$P(\text{reject at least one true } H_0^i)$$

- How?

## Why MCTP and SCI?

- Better: Start with multiple comparisons ( $\rightarrow$  multiple contrast test)
  - Control the multiple level  $\alpha$  in the strong sense

$$P(\text{reject at least one true } H_0^i)$$

- How?
  - Adapt the concept of MCTP from Bretz, Genz and Hothorn (2001)

# Why MCTP and SCI?

- Better: Start with multiple comparisons ( $\rightarrow$  multiple contrast test)
  - Control the multiple level  $\alpha$  in the strong sense

$$P(\text{reject at least one true } H_0^i)$$

- How?
  - Adapt the concept of MCTP from Bretz, Genz and Hothorn (2001)
  - Idea: For each partial hypothesis one t-Test

# Why MCTP and SCI?

- Better: Start with multiple comparisons ( $\rightarrow$  multiple contrast test)
  - Control the multiple level  $\alpha$  in the strong sense

$$P(\text{reject at least one true } H_0^i)$$

- How?
  - Adapt the concept of MCTP from Bretz, Genz and Hothorn (2001)
  - Idea: For each partial hypothesis one t-Test
  - Compare the t-values with one critical value (Gabriel, 1969)

# Multiple Contrast Tests

- Estimators

# Multiple Contrast Tests

- Estimators
  - $\bar{\mathbf{X}} = (\bar{X}_{1\cdot}, \dots, \bar{X}_{a\cdot})'$  (vector of means)

# Multiple Contrast Tests

- Estimators
  - $\bar{\mathbf{X}} = (\bar{X}_{1\cdot}, \dots, \bar{X}_{a\cdot})'$  (vector of means)
  - $\hat{\mathbf{V}}$ : empirical covariance matrix

# Multiple Contrast Tests

- Estimators
  - $\bar{\mathbf{X}} = (\bar{X}_{1\cdot}, \dots, \bar{X}_{a\cdot})'$  (vector of means)
  - $\hat{\mathbf{V}}$ : empirical covariance matrix
- Variance of a contrast

$$\sigma_\ell^2 = \text{Var}(\mathbf{c}_\ell' \bar{\mathbf{X}}) = \mathbf{c}_\ell' \mathbf{V} \mathbf{c}_\ell, \quad \hat{\sigma}_\ell^2 = \mathbf{c}_\ell' \hat{\mathbf{V}} \mathbf{c}_\ell$$



# Multiple Contrast Tests

- Estimators

- $\bar{\mathbf{X}}. = (\bar{X}_{1.}, \dots, \bar{X}_{a.})'$  (vector of means)

- $\hat{\mathbf{V}}$ : empirical covariance matrix

- Variance of a contrast

$$\sigma_\ell^2 = \text{Var}(\mathbf{c}_\ell' \bar{\mathbf{X}}.) = \mathbf{c}_\ell' \mathbf{V} \mathbf{c}_\ell, \quad \hat{\sigma}_\ell^2 = \mathbf{c}_\ell' \hat{\mathbf{V}} \mathbf{c}_\ell$$

- $\hat{\sigma}_\ell^2 = \mathbf{c}_\ell' \hat{\mathbf{V}} \mathbf{c}_\ell$

# Parametric MCTP

- Multiple Comparisons

# Parametric MCTP

- Multiple Comparisons
  - For  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$ :

$$T_\ell = \sqrt{n} \frac{\mathbf{c}'_\ell (\bar{\mathbf{X}} - \boldsymbol{\mu})}{\hat{\sigma}_\ell}, \ell = 1, \dots, q$$

# Parametric MCTP

- Multiple Comparisons
  - For  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$ :

$$T_\ell = \sqrt{n} \frac{\mathbf{c}'_\ell (\bar{\mathbf{X}} - \boldsymbol{\mu})}{\hat{\sigma}_\ell}, \ell = 1, \dots, q$$

- $T_\ell \sim t_{n-1}$  (under  $H_0$ )

# Parametric MCTP

- Multiple Comparisons

- For  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$ :

$$T_\ell = \sqrt{n} \frac{\mathbf{c}'_\ell (\bar{\mathbf{X}} - \boldsymbol{\mu})}{\hat{\sigma}_\ell}, \ell = 1, \dots, q$$

- $T_\ell \sim t_{n-1}$  (under  $H_0$ )
    - More than one test is performed, hence the type-1 error rate will increase

# Parametric MCTP

- Multiple Comparisons

- For  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$ :

$$T_\ell = \sqrt{n} \frac{\mathbf{c}'_\ell (\bar{\mathbf{X}} - \boldsymbol{\mu})}{\hat{\sigma}_\ell}, \ell = 1, \dots, q$$

- $T_\ell \sim t_{n-1}$  (under  $H_0$ )
    - More than one test is performed, hence the type-1 error rate will increase
    - Need to be adjusted

# Parametric MCTP

- Quality of the method?

# Parametric MCTP

- Quality of the method?
- Are the test statistics  $T_\ell$  and  $T_m$  independent?



# Parametric MCTP

- Quality of the method?
- Are the test statistics  $T_\ell$  and  $T_m$  independent?
- Covariance of  $\mathbf{c}'_\ell \bar{\mathbf{X}}$  and  $\mathbf{c}'_m \bar{\mathbf{X}}$ :

# Parametric MCTP

- Collect all test statistics in a vector  $\mathbf{T}$

# Parametric MCTP

- Collect all test statistics in a vector  $\mathbf{T}$
- $\mathbf{T} = (T_1, \dots, T_q)' \sim T(\mathbf{0}, n-1, \mathbf{R})$

# Parametric MCTP

- Collect all test statistics in a vector  $\mathbf{T}$
- $\mathbf{T} = (T_1, \dots, T_q)' \sim T(\mathbf{0}, n-1, \mathbf{R})$
- $\mathbf{R} =$

# Parametric MCTP

- Collect all test statistics in a vector  $\mathbf{T}$
- $\mathbf{T} = (T_1, \dots, T_q)' \sim T(\mathbf{0}, n-1, \mathbf{R})$
- $\mathbf{R} =$
- Empirical correlation  $\hat{\mathbf{R}} =$

# Parametric MCTP

- Test decisions

# Parametric MCTP

- Test decisions

- Reject  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  if  $|T_\ell| \geq t_{1-\alpha}(\hat{\mathbf{R}})$

$$CI_\ell = \mathbf{c}'_\ell \bar{\mathbf{X}}. \pm t_{1-\alpha}(\hat{\mathbf{R}})/\sqrt{n} \cdot \hat{\sigma}_\ell, \ell = 1, \dots, q$$

# Parametric MCTP

- Test decisions

- Reject  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  if  $|T_\ell| \geq t_{1-\alpha}(\hat{\mathbf{R}})$

$$CI_\ell = \mathbf{c}'_\ell \bar{\mathbf{X}}. \pm t_{1-\alpha}(\hat{\mathbf{R}}) / \sqrt{n} \cdot \hat{\sigma}_\ell, \ell = 1, \dots, q$$

- Reject  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$  if  $T_0 = \max(|T_1|, \dots, |T_q|) \geq t_{1-\alpha}(\hat{\mathbf{R}})$



# Parametric MCTP

- Test decisions

- Reject  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  if  $|T_\ell| \geq t_{1-\alpha}(\hat{\mathbf{R}})$

$$CI_\ell = \mathbf{c}'_\ell \bar{\mathbf{X}}. \pm t_{1-\alpha}(\hat{\mathbf{R}}) / \sqrt{n} \cdot \hat{\sigma}_\ell, \ell = 1, \dots, q$$

- Reject  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$  if  $T_0 = \max(|T_1|, \dots, |T_q|) \geq t_{1-\alpha}(\hat{\mathbf{R}})$
  - $t_{1-\alpha}(\mathbf{R}) : (1 - \alpha)$ -quantile of the multivariate  $T(\mathbf{0}, n - 1, \hat{\mathbf{R}})$  distribution

# Parametric MCTP

- Test decisions

- Reject  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  if  $|T_\ell| \geq t_{1-\alpha}(\hat{\mathbf{R}})$

$$CI_\ell = \mathbf{c}'_\ell \bar{\mathbf{X}}. \pm t_{1-\alpha}(\hat{\mathbf{R}}) / \sqrt{n} \cdot \hat{\sigma}_\ell, \ell = 1, \dots, q$$

- Reject  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$  if  $T_0 = \max(|T_1|, \dots, |T_q|) \geq t_{1-\alpha}(\hat{\mathbf{R}})$
  - $t_{1-\alpha}(\mathbf{R})$  :  $(1 - \alpha)$ -quantile of the multivariate  $T(\mathbf{0}, n - 1, \hat{\mathbf{R}})$  distribution
    - Different interpretation of the quantile?

# Parametric MCTP

- Test decisions

- Reject  $H_0 : \mathbf{c}'_\ell \boldsymbol{\mu} = 0$  if  $|T_\ell| \geq t_{1-\alpha}(\hat{\mathbf{R}})$

$$CI_\ell = \mathbf{c}'_\ell \bar{\mathbf{X}}. \pm t_{1-\alpha}(\hat{\mathbf{R}}) / \sqrt{n} \cdot \hat{\sigma}_\ell, \ell = 1, \dots, q$$

- Reject  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$  if  $T_0 = \max(|T_1|, \dots, |T_q|) \geq t_{1-\alpha}(\hat{\mathbf{R}})$
  - $t_{1-\alpha}(\mathbf{R})$  :  $(1 - \alpha)$ -quantile of the multivariate  $T(\mathbf{0}, n - 1, \hat{\mathbf{R}})$  distribution
    - Different interpretation of the quantile?
    - quantiles of the distribution of a maximum

# MCTP: Properties

## Advantages

- 
- 
- 
- 

## Disadvantages

- 
- 
- 
-

# MCTP vs ANOVA

- ANOVA

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\mu = \mathbf{0}$

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms
  - No information for particular dose levels available



# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms
  - No information for particular dose levels available
- MCTP

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\mu = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms
  - No information for particular dose levels available
- MCTP
  - Linear test statistics

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\mu = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms
  - No information for particular dose levels available
- MCTP
  - Linear test statistics
  - Test decisions via using multivariate critical values

# MCTP vs ANOVA

- ANOVA
  - Quadratic test procedure for global hypothesis  $H_0 : \mathbf{C}\mu = \mathbf{0}$
  - Derivation based on multivariate normality, but transformation in  $\mathbb{R}$  via quadratic forms
  - No information for particular dose levels available
- MCTP
  - Linear test statistics
  - Test decisions via using multivariate critical values
  - Informations and SCI for particular dose levels available

# Generalisations?

- Are there any drawbacks of the model assumptions?

# Generalisations?

- Are there any drawbacks of the model assumptions?
- Of the method?

# Generalisations?

- Are there any drawbacks of the model assumptions?
- Of the method?
- What to do in case of small sample sizes?

# Generalisations?

- Are there any drawbacks of the model assumptions?
- Of the method?
- What to do in case of small sample sizes?
- Resampling strategies?



# Resampling Strategies

- Goal: Estimate the quantiles from a distribution of a maximum of correlated  $t$ -statistics

# Resampling Strategies

- Goal: Estimate the quantiles from a distribution of a maximum of correlated  $t$ -statistics
- How?

# Resampling Strategies

- Goal: Estimate the quantiles from a distribution of a maximum of correlated  $t$ -statistics
- How?
- next slide

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}_k^*$

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^* - \bar{\mathbf{X}}.)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$



# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^* - \bar{\mathbf{X}}.)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and save in  $T_{0,s}^*$

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^* - \bar{\mathbf{X}})}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and save in  $T_{0,s}^*$
  - Repeat the above a large number of times ( $M = 10,000$ )

# Nonparametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  with replacement from  $\mathbf{X}$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^* - \bar{\mathbf{X}})}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and save in  $T_{0,s}^*$
  - Repeat the above a large number of times ( $M = 10,000$ )
- Estimate  $t_{1-\alpha}(\mathbf{R})$  by the  $(1 - \alpha)$ -quantile of

$$|T_{0,1}^*|, \dots, |T_{0,M}^*|$$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^*)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$



# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^*)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and safe in  $T_{0,s}^*$

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^*)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and save in  $T_{0,s}^*$
  - Repeat the above a large number of times ( $M = 10,000$ )

# Parametric Bootstrap

- Fix the data  $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_n)$ 
  - Draw  $n$  vectors  $\mathbf{X}_k^*$  from  $N(\mathbf{0}, \hat{\mathbf{V}})$
  - Compute their means:  $\bar{\mathbf{X}}^*$
  - Compute their empirical covariance matrix:  $\hat{\mathbf{V}}^*$
  - Compute  $\mathbf{T}^* = (T_1^*, \dots, T_q^*)'$ ;  $T_\ell^* = \sqrt{n} \frac{\mathbf{c}_\ell'(\bar{\mathbf{X}}^*)}{\sqrt{\mathbf{c}_\ell' \hat{\mathbf{V}}^* \mathbf{c}_\ell}}$
  - Compute  $T_0 = \max\{|T_1^*|, \dots, |T_q^*|\}$  and save in  $T_{0,s}^*$
  - Repeat the above a large number of times ( $M = 10,000$ )
- Estimate  $t_{1-\alpha}(\mathbf{R})$  by the  $(1 - \alpha)$ -quantile of

$$|T_{0,1}^*|, \dots, |T_{0,M}^*|$$