

# The Use of Real Time MRI in the Assessment of Articulatory Outcomes of Partial Glossectomy

Maury Lander-Portnoy

Saturday, November 11, 2017

## 1 Introduction

- I. Real-time MRI as exciting new imaging tool
  - A. Safe imaging protocol
    - i. Non-ionizing radiation
    - ii. Non-invasive measurements
  - B. Provides rich data sets
  - C. Allows task-specific customization
    - i. Can customize the trade-off between:
      - a. Spatial Resolution
      - b. # Slices Acquired
      - c. Temporal Resolution
  - D. Allows concurrent global imaging of vocal tract
- II. Current study details
  - A. Patient: Advanced stage basal tongue cancer
  - B. Operation: Partial glossectomy (cancerous tissue resection)
  - C. Provides relatively controlled manipulation of vocal tract
    - i. Non-operated articulators are unaffected
    - ii. Patient cognition remains largely unchanged
  - D. Provides good opportunity for controlled research questions
    - i. Here we explore how morphology (and its changes) influence articulator synergy
    - ii. Controls other factors to observe change in synergy due to change in morphology
- III. Current work's aims:
  - A. Pre-op and post-op data allow quantification of morphological changes
  - B. Vocal tract dynamics allow estimation of articulator synergies
  - C. Observation of relationship between morphology and synergies
- IV. Methods allow future examination of:
  - A. Individual variability in articulator synergies



- B. Relationship of morphological change and compensation
- C. Individual variability in degree of plasticity
- D. Relationship of accommodation to functional outcome
- E. Use of feedback to monitor articulatory task goals

## 2 Methods

- I. Participant: Male speaker of British English
- II. Scan format and time course:
  - A. Pre-op scan just prior to surgery
  - B. Post-op scan 6 months after surgery (when intelligibility plateaus)
  - C. Participant supine with motion minimized
  - D. Reading 30 second passages of scripted speech
- III. Stimuli:
  - A. Numerous repetitions collected during both preop and postop sessions
    - i. Monosyllabic (bVt) words
    - ii. Rainbow passage
    - iii. MOCHA-TIMIT sentences
- IV. Imaging:
  - A. Scanner: 1.5T GE Signa Excite HD
  - B. Max gradient amplitude: 4.0 G/cm
  - C. Max slew rate: 15.0 G/cm/ms
  - D. Transmission: body coil
  - E. Reception: 8-channel upper airway coil
- V. Data:
  - A. FOV: 200x200mm
  - B. Slice thickness: 6mm
  - C. Flip angle: 15°
  - D. Spatial resolution: 2.38x2.38mm
  - E. Temporal resolution: 11ms preop, 12ms postop

## VI. Pre-processing:

- A. Air-tissue boundary segmentation
  - i. Unsupervised algorithm
  - ii. Provided anatomic template
  - iii. Hierarchical gradient descent optimized fit
  - iv. Produces polyline vocal tract outline
  - v. Preserves different articulator identities:



- D. Relationship of weights to constrictions is mapped forming weight sequences
- E. Weight sequences represent articulatory strategy and can reveal articulatory differences
- F. Simulations of varying tasks
  - i. Utilizes aforementioned forward map
  - ii. Reveals differing articulatory strategies between forward maps

## VII. Analysis:


- A. Population sub-sampling:
  - i. Necessary due to data paucity
  - ii. Allows drawing meaningful conclusions from sample differences
  - iii. 5 iterations of random bisection and comparison
  - iv. Produced 20 sub-samples (5 iterations x 2 halves x 2 data-sets)
- B. Guided factor analysis:
  - i. Allowed isolation of articulator contributions
  - ii. Quantified levels of tongue and jaw activation in coordinated tasks
  - iii. Quantification allows synergy calculations
  - iv. Changes in synergies pre- vs. post-op show compensation in articulatory strategies

## VIII. Forward map estimation:


- A. Relies on task dynamics
  - i. Defines speech goal as constriction formation
  - ii. Constriction degree and location change for different speech tasks
- B. Calculates constrictions from segmentation data
- C. Each frame's constrictions can be made by weighting and summing articulator factors
  - i. Weights represent degree of utilization of that factor

## 3 Results


- I. Static vocal tract outline reveals tongue volume decrease basally in post-operative images



- II. We observe no systematic difference in articulatory synergy, therefore we find no evidence for compensation



III. We do see a difference in pre- vs. post-op factors, implying a difference in articulatory behavior



## References

- [1] E. Bresch, Y. C. Kim, K. Nayak, D. Byrd, and S. Narayanan. Seeing speech: Capturing vocal tract shaping using real-time magnetic resonance imaging [Exploratory DSP]. *IEEE Signal Processing Magazine*, 25(3):123–132, May 2008.
- [2] Tim Bressmann, Robert Sader, Tara L Whitehill, and Nabil Samman. Consonant intelligibility and tongue motility in patients with partial glossectomy. *Journal of Oral and Maxillofacial Surgery*, 62(3):298–303, March 2004.
- [3] Jun Cai, Yves Laprie, Julie Busset, and Fabrice Hirsch. Articulatory modeling based on semi-polar coordinates and guided PCA technique. In *10th Annual Conference of the International Speech Communication Association-INTERSPEECH 2009*, 2009.
- [4] Christina Hagedorn, Adam Lammert, Mary Bassily, Yihe Zu, Uttam Sinha, Louis Goldstein, and Shrikanth S. Narayanan. Characterizing postglossectomy speech using real-time MRI. In *International Seminar on Speech Production, Cologne, Germany*, 2014.
- [5] Y. Matsui, K. Ohno, Y. Yamashita, and K. Takahashi. Factors influencing postoperative speech function of tongue cancer patients following reconstruction with fasciocutaneous/myocutaneous flaps—a multicenter study. *International Journal of Oral and Maxillofacial Surgery*, 36(7):601–609, July 2007.
- [6] Y. Matsui, T. Shirota, Y. Yamashita, and K. Ohno. Analyses of speech intelligibility in patients after glossectomy and reconstruction with fasciocutaneous/myocutaneous flaps. *International Journal of Oral and Maxillofacial Surgery*, 38(4):339–345, April 2009.
- [7] Ken-ichi Michi, Satoko Imai, Yukari Yamashita, and Noriko Suzuki. Improvement of speech intelligibility by a secondary operation to mobilize the tongue after glossectomy. *Journal of Cranio-Maxillofacial Surgery*, 17(4):162–166, May 1989.
- [8] Shrikanth Narayanan, Krishna Nayak, Sungbok Lee, Abhinav Sethy, and Dani Byrd. An approach to real-time magnetic resonance imaging for speech production. *The Journal of the Acoustical Society of America*, 115(4):1771–1776, April 2004.
- [9] John E. Overall. Orthogonal Factors and Uncorrelated Factor Scores. *Psychological Reports*, 10(3):651–662, June 1962.
- [10] Vikram Ramanarayanan, Louis Goldstein, Dani Byrd, and Shrikanth S. Narayanan. An investigation of articulatory setting using real-time magnetic resonance imaging. *The Journal of the Acoustical Society of America*, 134(1):510–519, July 2013.
- [11] Elliot L. Saltzman and Kevin G. Munhall. A Dynamical Approach to Gestural Patterning in Speech Production. *Ecological Psychology*, 1(4):333–382, December 1989.
- [12] Satoko IMAI, Ken-ichi MICHI, Yukari YAMASHITA, Hiroshi YOSHIDA, Yukihiro MICHIWAKI, Kohsuke OHNO, and Noriko SUZUKI. Speech intelligibility after resection of the tongue and floor of the mouth. , 34(8):1567–1583, 1988.
- [13] Tanner Sorensen, Asterios Toutios, Louis Goldstein, and Shrikanth S. Narayanan. Characterizing Vocal Tract Dynamics Across Speakers Using Real-Time MRI. pages 465–469, September 2016.
- [14] Jian Sun, Yanqiu Weng, Jun Li, Guoming Wang, and Zhiyuan Zhang. Analysis of Determinants on Speech Function After Glossectomy. *Journal of Oral and Maxillofacial Surgery*, 65(10):1944–1950, October 2007.
- [15] Asterios Toutios and Shrikanth S. Narayanan. Factor analysis of vocal-tract outlines derived from real-time magnetic resonance imaging data. In *International Congress of Phonetic Sciences (ICPhS 2015)*. 2015.
- [16] Asterios Toutios and Shrikanth S. Narayanan. Advances in real-time magnetic resonance imaging of the vocal tract for speech science and technology research. *APSIPA Transactions on Signal and Information Processing*, 5, 2016.