

# Optimization notes

Ying-Jia Lin

National Cheng Kung University

May 3rd, 2021

# Directional derivative

From a starting point  $\underline{x}_0$  and a given direction  $\underline{u}$ :

- $\underline{x}(\lambda) = \underline{x}_0 + \lambda \underline{u}$ 
  - $\lambda$  is a scalar.
- $d\underline{x} = \underline{u}d\lambda$ 
  - For a small change in  $\lambda$ .
- $F(\lambda) = f(\underline{x}_0 + \lambda \underline{u})$

$$\begin{aligned}dF &= df = (\nabla f(\underline{x}))^\top d\underline{x} \\ &= (\nabla f(\underline{x}))^\top \underline{u}d\lambda = \nabla^\top f \underline{u} \lambda\end{aligned}$$

- $\frac{df}{d\lambda} = \nabla^\top f \underline{u}$ 
  - If  $f$  is minimized at  $\underline{x}^* = \underline{x}_0 + \lambda \underline{u}$ , then:
    - $\nabla f(\underline{x}^*)^\top \underline{u} = 0$
    - gradient  $f$  evaluated at the minimum point is orthogonal to  $\underline{u}$ .

# Weierstrass Theorem

If  $f(\underline{x})$  is continuous on a nonempty feasible set that is closed and bounded, then  $f(\underline{x})$  has a global minimum in this set.

- ▶ A set  $S$  is bounded if for any point  $\underline{x}$  in  $S$ , we have  $\underline{x}^T \underline{x} < c$ 
  - ▶  $c$  is a finite positive number.

# Single-variable unconstrained optimization

- Necessary condition
  - If a function  $f(x)$  has a local minimum at  $x = x^*$ , and  $f'(x)$  exists as a finite number at  $x = x^*$ , then  $f'(x^*) = 0$ .
  - $x^*$  at  $f'(x^*) = 0$  is called stationary point.
- Sufficient condition
  - Suppose  $f'(x^*) = f''(x^*) = \dots = f^{(m-1)}(x^*) = 0$ , but  $f^{(m)}(x^*) \neq 0$ , then  $f(x^*)$  is:
    - 1. a local minimum if  $f^{(m-1)}(x^*) > 0$  and  $m$  is even.
    - 2. a local maximum if  $f^{(m-1)}(x^*) < 0$  and  $m$  is even.
    - 3. neither a maximum nor a minimum if  $m$  is odd.

# Multi-variable unconstrained optimization (1)

Definition of  $r^{th}$  differential of function  $f$ :

$$d^r f(\underline{x}^*) = \sum_{i=1}^n \sum_{j=1}^n \cdots \sum_{k=1}^n h_i h_j \cdots h_k \frac{\partial^r f(\underline{x}^*)}{\partial x_i \partial x_j \cdots \partial x_k}$$

## Example

When (order)  $r = 2$  and (number of variables)  $n = 3$ , we have:

$$\begin{aligned} d^2 f(\underline{x}^*) &= d^2 f(x_1^*, x_2^*, x_3^*) = \sum_{i=1}^3 \sum_{j=1}^3 h_i h_j \frac{\partial^2 f(\underline{x}^*)}{\partial x_i \partial x_j} \\ &= h_1^2 \frac{\partial^2 f(\underline{x}^*)}{\partial x_1^2} + h_2^2 \frac{\partial^2 f(\underline{x}^*)}{\partial x_2^2} + h_3^2 \frac{\partial^2 f(\underline{x}^*)}{\partial x_3^2} \\ &\quad + 2h_1 h_2 \frac{\partial^2 f(\underline{x}^*)}{\partial x_1 \partial x_2} + 2h_2 h_3 \frac{\partial^2 f(\underline{x}^*)}{\partial x_2 \partial x_3} + 2h_1 h_3 \frac{\partial^2 f(\underline{x}^*)}{\partial x_1 \partial x_3} \end{aligned}$$

## Multi-variable unconstrained optimization (2)

- Necessary condition

$$\frac{\partial f(\underline{x}^*)}{\partial x_1} = \frac{\partial f(\underline{x}^*)}{\partial x_2} = \dots = \frac{\partial f(\underline{x}^*)}{\partial x_n} = 0$$

- In vector form,  $\nabla f(\underline{x}^*) = 0$ .
- $\underline{x}^*$  at  $\nabla f(\underline{x}^*) = 0$  is called stationary point.
- Sufficient condition
  - For a stationary point at  $\underline{x} = \underline{x}^*$ :
    - if the Hessian matrix of  $f(\underline{x})$  evaluated at  $\underline{x} = \underline{x}^*$  is **positive definite**, then  $\underline{x}^*$  is a local minimum.
    - if the Hessian matrix of  $f(\underline{x})$  evaluated at  $\underline{x} = \underline{x}^*$  is **negative definite**, then  $\underline{x}^*$  is a local maximum.

# Hessian Matrix

- $(H_f)_{i,j} = \frac{\partial^2 f}{\partial x_i \partial x_n}$

- In matrix form:  $H_f = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}$

# Definiteness

- A matrix is positive definite if all its **eigenvalues** are positive.
  - If some of the eigenvalues are positive and some are zero, then the matrix is positive semidefinite.
- **Checking the sign of the determinants** is an alternative way to determine the definiteness of a matrix.
- A twice differentiable function is convex if and only if its Hessian matrix is positive semi-definite.
  - The function is strictly convex if the Hessian matrix is positive definite.



# Multivariable optimization with equality constraints

## Lagrange Multiplier Theorem

- Suppose the point  $\underline{x}^*$  minimizes  $f(\underline{x})$  and satisfies the equality constraints:  $h_j(\underline{x}^*) = 0$ , for  $j = 1, 2, \dots, m$
- Assume that the constraint gradients  $\nabla h_j(\underline{x}^*)$  are linealy independent.
- Then there exists a unique set  $\lambda_j^*$  ( $j = 1, 2, \dots, m$ ) satisfying:

$$\frac{\partial L}{\partial x_i} = \frac{\partial f}{\partial x_i} + \sum_{j=1}^m \lambda_j^* \frac{\partial h_j}{\partial x_i} = 0$$

where  $i = 1, 2, \dots, n$ .

# Simplex method: Introduction

## Definition of Simplex

The geometric figure formed by a set of  $n + 1$  points in an  $n$ -dimensional space is called a simplex.

- When the points are equidistant, the simplex is said to be *regular*.
- In two dimensions the simplex is a triangle, and in three dimensions, it is a tetrahedron.
- The simplex method was originally given by Spendley et al. and was developed later by Nelder and Mead.

# Simplex method: Steps

Choose a reflection coefficient  $\alpha > 0$ , an expansion coefficient  $\gamma > 1$ , and a contraction coefficient  $0 < \beta < 1$ .

1. Identify  $\underline{x}_{\min}$  and  $\underline{x}_{\max}$  among  $\{\underline{x}_1, \underline{x}_2, \dots, \underline{x}_{n+1}\}$ 
  - such that  $f(\underline{x}_{\min})$  is the minimum and  $f(\underline{x}_{\max})$  is the maximum of all the  $f(\underline{x}_i)$ , for  $i = 1, 2, \dots, n + 1$ .
  - If  $|\underline{x}_{\max} - \underline{x}_{\min}| < \epsilon$ , stop. The minimum is at  $\underline{x}_{\min}$ .
  - Otherwise, let  $\underline{x}_a$  be the averaged position of  $\{\underline{x}_1, \underline{x}_2, \dots, \underline{x}_{n+1}\}$ , **excluding**  $\underline{x}_{\max}$ , and go to step 2.
2. Let the reflection point  $\underline{x}_r = \underline{x}_a + \alpha(\underline{x}_a - \underline{x}_{\max})$ .
  - If  $f(\underline{x}_{\min}) > f(\underline{x}_r)$ , let the expansion point  $\underline{x}_e = \underline{x}_a + \gamma(\underline{x}_r - \underline{x}_a)$ , and go to step 3.
  - Otherwise, go to step 4.

# Simplex method: Steps

3. If  $f(\underline{x}_r) > f(\underline{x}_e)$ , the point  $\underline{x}_{\max}$  is **replaced** by  $\underline{x}_e$ .
  - Otherwise,  $\underline{x}_{\max}$  is **replaced** by  $\underline{x}_r$ . A new set of  $n+1$  points is formed. Then go to step 1.
4. If the second largest  $f(\underline{x}_i) > f(\underline{x}_r)$ , then  $\underline{x}_{\max}$  is **replaced** by  $\underline{x}_r$  to form a new set of  $n+1$  points, and go to step 1.
  - Otherwise, go to step 5.
5. Let  $\underline{x}_p$  be defined such that  $f(\underline{x}_p) = \min \{f(\underline{x}_r), f(\underline{x}_{\max})\}$  and let the contraction point  $\underline{x}_c = \underline{x}_a + \beta(\underline{x}_p - \underline{x}_a)$ .
  - If  $f(\underline{x}_c) > f(\underline{x}_p)$ , **replace**  $\underline{x}_j$  by  $\underline{x}_j + (\underline{x}_{\min} - \underline{x}_j)/2$ , for  $j = 1, 2, \dots, n+1$ , and go to step 1.
  - Otherwise,  $\underline{x}_c$  **replaces**  $\underline{x}_{\max}$  to form a new set of  $n+1$  points, and go to step 1.