




Searching for ALPs from CCSNe with *Fermi* LAT's low-energy technique

Milena Crnogorčević^{1,2}, Regina Caputo², Manuel Meyer³, Nicola Omodei⁴, Michael Gustafsson⁵

AXIONLIKE PARTICLES

and where to find them




Sensitivity Analysis & Results

Analysis Steps

- Model backgrounds from the considered LLE-detected GRB sample
- Compute the extreme background levels
- Produce a grid of scaled ALP signals for two different progenitors.
- Produce 2000 realizations of the background + ALP spectrum and their corresponding (GRB) response functions (XSPEC fakeit)
- Fit the ALP and the "background-only" models
- Apply Wilks' Theorem and LLR test to find for which normalization ALP model is preferred

→ Find the coupling-distance parameter space for that normalization.



Distance limits for a LAT-LLE ALP 5-sigma detection for a 10-solar-mass progenitor for different background levels.

Conversion probability	Distance limit (Mpc)		
	Background level:		
	Low	Median	High
$P_y(g_0)$	4.4 (6.5)	4.9 (7.1)	6.6 (9.7)
0.1	3.1 (4.6)	3.5 (5.0)	4.7 (6.9)
0.05	1.4 (2.1)	1.5 (2.3)	2.1 (3.1)
0.01	0.4 (0.7)	0.5 (0.7)	0.7 (1.0)

Analogous to above, (including 18-solar-mass progenitor) for different background levels and conversion probabilities.


ALP Searches

Analysis Steps

- Data selection: 186 LAT-detected GRBs → final sample of 24 GRBs

Property	Selection Criterion
Distance	unassociated (no redshift)
Detection significance	$\geq 5\sigma$ in LAT-LLE ($\gtrsim 30$ MeV)
Observed time interval	\geq duration of the burst
Burst duration	long GRBs ($T_{95} \gtrsim 2$ seconds)

- Model comparison (null model: w/o ALP, alternative: w/ ALP)



An example of a spectral fit for GRB 101123A with and without the ALP component. The corresponding pre-trial LLR value is 5.8, or $\sim 2.4\sigma$. The corresponding post-trials value is $p \sim 0.3$.

- We report no statistically significant detections.

Conclusions & Current Work

- Novel method and results: using a transient data class as observed by Fermi to probe its sensitivity. Results are consistent with the analysis using the standard LAT data [Meyer et al. 2020]
- Sensitivity analysis pipeline: adaptable for calculating distance limits for the current and future gamma-ray instruments for the given ALP mass and coupling
- Good scientific case for the future instruments: they need more sensitivity in the MeV region in order to be able to increase the statistics of sources considered (see AMEGO-X posters!)

Current work: Comprehensive search for precursor emission in LGRBs, computing competitive limits on (m_a, g_{ay})

References

- M. Crnogorčević et al., Phys. Rev. D 104, 103001 (2021) • M. Meyer et al., Phys. Rev. Lett. 124, 231101 (2020); Erratum, Phys. Rev. Lett. 125, 119901 (2020). • A. Payez et al., J. Cosmol. Astropart. Phys. 02 (2015) 006. • M. Meyer et al., Phys. Rev. Lett. 118, 011103 (2017). • G. Raffelt et al., Phys. Rev. D 37, 1237 (1988). •

Read our paper!
arXiv:2109.05790


¹University of Maryland, College Park, ²NASA Goddard Space Flight Center, ³University of Hamburg, ⁴SLAC National Accelerator Laboratory, ⁵Stockholm University



Presenter's contact:
mcrnogor@umd.edu
[mcrnogor.github.io](https://github.com/mcrnogor)
[@mcrnogor](https://twitter.com/mcrnogor)
[milena-crnogorčević](https://orcid.org/0000-0002-1111-1111)



M. C. acknowledges support by NASA under Grant No. 80GSFC21M0002. M. M. acknowledges support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program Grant Agreement No. 948689 (AxionDM) and from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy—EXC 2121 "Quantum Universe"—39083306.



Take a photo to learn more about my research!