

DNA damage kinase signaling: checkpoint and repair at 30 years

Michael Charles Lanz, Diego Dibitetto & Marcus Bustamante Smolka^{*} 

Abstract

From bacteria to mammalian cells, damaged DNA is sensed and targeted by DNA repair pathways. In eukaryotes, kinases play a central role in coordinating the DNA damage response. DNA damage signaling kinases were identified over two decades ago and linked to the cell cycle checkpoint concept proposed by Weinert and Hartwell in 1988. Connections between the DNA damage signaling kinases and DNA repair were scant at first, and the initial perception was that the importance of these kinases for genome integrity was largely an indirect effect of their roles in checkpoints, DNA replication, and transcription. As more substrates of DNA damage signaling kinases were identified, it became clear that they directly regulate a wide range of DNA repair factors. Here, we review our current understanding of DNA damage signaling kinases, delineating the key substrates in budding yeast and humans. We trace the progress of the field in the last 30 years and discuss our current understanding of the major substrate regulatory mechanisms involved in checkpoint responses and DNA repair.

Keywords checkpoint; DNA damage response; DNA repair; kinase; mass spectrometry

DOI 10.15252/emboj.2019101801 | Received 14 February 2019 | Revised 3 July 2019 | Accepted 24 July 2019 | Published online 8 August 2019

The EMBO Journal (2019) 38: e101801

See the Glossary for abbreviations used in this article.

Introduction

From pathway to network

In eukaryotes, kinases play a central role in the DNA damage response, from sensing DNA damage to regulating cellular processes. Work in yeast and mammalian systems in the 1990s identified an evolutionarily conserved set of DNA damage signaling kinases, including phosphatidylinositol 3' kinase (PI3K)-related kinases (PIKKs) and PIKK-regulated downstream kinases. These kinases were found to be involved in cell cycle control (Allen *et al*, 1994; Carr, 1995; Morrow *et al*, 1995; Savitsky *et al*, 1995; Cimprich *et al*, 1996; Furnari *et al*, 1997; Peng *et al*, 1997; Sanchez *et al*, 1997) and were linked to the “cell cycle checkpoint” concept

proposed by Weinert and Hartwell (Weinert & Hartwell, 1988; Hartwell & Weinert, 1989). Subsequent work in the late 1990s and early 2000s revealed how these kinases establish the checkpoint and control processes beyond the cell cycle, such as apoptosis, transcription, and DNA replication (Santocanale & Diffley, 1998; Sun *et al*, 1998; Zhou & Elledge, 2000). In 2007, the use of mass spectrometry (MS)-based proteomics allowed a more systematic analysis of the network of phosphorylation events triggered by DNA damage signaling kinases (Matsuoka *et al*, 2007; Smolka *et al*, 2007). As a result, the perception that DNA damage signaling kinases operate within a simple signaling pathway (Fig 1; the classical “linear” depiction of DNA damage signaling) evolved to a more comprehensive view in which DNA damage signaling kinases function in an elaborate signaling network comprised of hundreds of substrates.

From checkpoint to DNA repair

After the discovery of DNA damage signaling kinases, mechanistic links between these kinases and the DNA repair machinery were virtually non-existent. It was not immediately appreciated that these kinases directly target and regulate the DNA repair machinery. Now that dozens of DNA repair proteins have been shown to be phosphorylated by DNA damage signaling kinases, there is little doubt that active and direct control of the DNA repair machinery is a core function of DNA damage signaling kinases. However, the precise mechanisms by which these kinases control the action of these substrates remain incompletely understood and represent a significant knowledge gap in the field. Here, we review our current understanding of the integrated action of DNA damage signaling kinases. We delineate the key substrates for checkpoint and DNA repair in budding yeast and highlight the potential parallels in humans. Based on the accumulated knowledge of the mechanisms of substrate regulation, we discuss how our understanding of the action of DNA damage signaling kinases in genome maintenance has evolved over the last 30 years.

DNA damage signaling kinases

DNA damage signaling kinases have been traditionally categorized as either apical or effector kinases. The apical PIKKs, or ATR, ATM, and DNA-PKcs in mammals and Mec1 and Tel1 in budding yeast (Fig 1; see Table 1 for gene name overview in model organisms),

Glossary

| | |
|-------------|--------------------------------------|
| BER | Base excision repair |
| BIR | Break-induced replication |
| CDK | Cyclin-dependent kinase |
| DDK | Dbf4-dependent kinase |
| dNTP | deoxyribonucleotide |
| GCR | Gross chromosomal rearrangement |
| HR | Homologous recombination |
| ICL | Interstrand crosslink |
| IR | Ionizing radiation |
| MMR | Mismatch repair |
| NER | Nucleotide excision repair |
| NHEJ | Non-homologous end joining |
| PIKK | PI3 kinase-related kinase |
| RNR | Ribonucleotide reductase |
| SAC | Spindle assembly checkpoint |
| SDSA | Synthesis-dependent strand annealing |
| SSA | Single-strand annealing |

associate with DNA structures that form as byproducts of DNA damage or replication stress, including single-strand DNA (ssDNA) and broken DNA ends (Fig 1). In the canonical mode of action, Mec1/ATR is recruited to ssDNA, whereas Tel1/ATM and DNA-PKcs associate with the ends of double-strand DNA (dsDNA) breaks. The mechanism of activation for each of these kinases is different, but recruitment to damaged DNA is often a requirement (for details on the mechanism of how kinases associate with DNA structures, as well as co-factors involved, refer to the following reviews: Blackford & Jackson, 2017; Di Domenico *et al*, 2014; Saldivar *et al*, 2017; Zou, 2013). Activated apical kinases transfer stimulatory phosphorylation to the downstream checkpoint kinases (Rad53, Chk1, and Dun1 in yeast; CHK2 and CHK1 in mammals), which catalyze phosphorylation events that mediate cellular responses to DNA damage as part of the canonical DNA damage checkpoint (Fig 1). Whereas Rad53 mediates nearly all checkpoint-related functions in budding yeast, with Chk1 playing only a minor role (Sanchez *et al*, 1999), a more balanced division of labor exists for CHK1 and CHK2 in humans. Human CHK2 shares sequence and structural similarities with yeast Rad53, including an FHA domain (Matsuoka *et al*, 1998), but the functional similarities are limited to the DSB signaling response. Human CHK1 and yeast Rad53 play a crucial role in the replication stress response, but they share little or no sequence/structural similarity.

Adaptor proteins as substrates for downstream signaling activation

In budding yeast, the transfer of phosphorylation from apical to downstream checkpoint kinases requires the checkpoint adaptor proteins Rad9 and Mrc1, which recruit the downstream checkpoint kinases in proximity to the apical kinase. Mec1 and Tel1 promote the recruitment of the Rad9 adaptor by phosphorylating lesion-proximal substrates, such as histone H2A (Downs *et al*, 2000) and the 9-1-1 complex (Paciotti *et al*, 1998), which are then recognized directly or indirectly by Rad9 (Fig 2; Toh *et al*, 2006; Hammet *et al*, 2007; Pfander & Diffley, 2011). Once recruited, Rad9 is phosphorylated by Mec1 or Tel1 (Emili, 1998; Vialard *et al*, 1998), which promotes its oligomerization (Soulier & Lowndes, 1999; Usui *et al*,

2009) and further stabilization on DNA (Naiki *et al*, 2004). Mec1- and Tel1-mediated phosphorylation of Rad9 creates docking sites for the recruitment of the downstream effector kinase Rad53 (Fig 2; Gilbert *et al*, 2001; Schwartz *et al*, 2002; Sun *et al*, 1998), which, upon recruitment to Rad9, is phosphorylated and activated by Mec1 or Tel1 (Sanchez *et al*, 1996; Sun *et al*, 1996). Chk1 also relies on Rad9 for its activation by Mec1 (Blankley & Lydall, 2004); however, unlike Rad53, the phosphorylation events that facilitate the recruitment of Chk1 to Rad9 are catalyzed by cyclin-dependent kinase (CDK; Fig 2; Abreu *et al*, 2013). Rad53 can also be activated via the Mrc1 adaptor (Alcasabas *et al*, 2001; Osborn & Elledge, 2003). Mrc1, being an intrinsic component of the replisome, is already “on-site” for mediating activation, obviating the need for a devoted recruitment mechanism, as is the case with Rad9. In fact, Mrc1 mediates a more rapid response to replication stress than Rad9-dependent DNA damage signaling (Pardo *et al*, 2017; Bacal *et al*, 2018). Similar to Rad9 and Mrc1, Sgs1 has been proposed to mediate Rad53 recruitment in a manner that depends upon its phosphorylation by Mec1 (Hegnauer *et al*, 2012). Mec1 and Tel1 also facilitate DNA damage signaling activation and propagation by recruiting chromatin modifiers and remodelers near sites of DNA damage (van Attikum *et al*, 2004; Downs *et al*, 2004; Morrison *et al*, 2004, 2007), which may help de-condense chromatin in a way that permits adaptor assembly.

Similar as in yeast, vertebrate ATM and ATR utilize checkpoint adaptor proteins to mediate their transfer of phosphorylation to the checkpoint effector kinases. ATR primarily relies on Claspin, the homolog of yeast Mrc1, to mediate the activation of CHK1 (Kumagai & Dunphy, 2000). Like Mrc1, Claspin associates with the replisome (Lee *et al*, 2003) and recruits CHK1 upon exposure to replication stress. Also similar to yeast, the Claspin–CHK1 interaction depends on ATR activity (Kumagai & Dunphy, 2003; Lindsey-Boltz *et al*, 2009). While it is currently unknown whether mammalian ATR directly phosphorylates Claspin, *Xenopus* ATR has been shown to directly phosphorylate Claspin at threonines 817 and 819, which are critical for CHK1 recruitment (Fig 2; the uncertainty of ATR’s direct phosphorylation of Claspin in humans is denoted by “?”; Yoo *et al*, 2006; in-depth discussion of the similarities and differences between Claspin orthologs can be found here: Smits *et al*, 2019). Once bound to CHK1, Claspin is thought to both stabilize CHK1 and tether it in proximity to ATR, allowing for extensive phosphorylation and full activation of the effector kinase (Liu *et al*, 2006). While ATR utilizes Claspin to facilitate its phosphorylation of CHK1, the phosphorylation of many other ATR substrates does not require Claspin, suggesting that, like in yeast, the adaptor proteins are primarily responsible for facilitating Mec1/ATR’s phosphorylation of the effector kinases and are not required for the phosphorylation of other substrates (like many of those depicted below in Fig 4).

Two mammalian adaptors have been linked to the ATM–CHK2 signaling axis: MDC1 and 53BP1. Despite extensive research on these proteins, it remains unclear precisely how they function in transducing ATM signaling toward CHK2 activation. While 53BP1 is the functional ortholog of yeast Rad9, MDC1 also shares functional similarities with Rad9 in the context of the DNA damage signaling response. Similar to Rad9, MDC1 possesses BRCT domains that directly bind to phosphorylated histone H2AX (Fig 2; γ H2AX; analogous to histone H2A phosphorylated at the C-terminal SQ site in yeast; Stucki *et al*, 2005). Once associated with γ H2AX at DNA




Figure 1. DNA damage signaling via PIKKs and checkpoint kinases in budding yeast and humans.

DNA damage signaling is initiated at DNA structures that form during DNA damage or replication stress, including single-strand DNA (ssDNA) and broken DNA ends. The apical PIKKs are recruited to these structures and become activated to initiate downstream signaling. Mec1/ATR is recruited to RPA-coated ssDNA, while Tel1/ATM and DNA-PKcs initially associate with DNA ends formed by double-strand breaks. Adaptor proteins are often required to mediate the transfer of phosphorylation from apical to downstream checkpoint kinases. Apical and downstream checkpoint kinases function coordinately to mediate cellular responses to DNA damage, either directly or through the regulation of additional downstream kinases. PIKKs also target an extensive network of substrates independently of downstream checkpoint kinases.

breaks, MDC1 is phosphorylated by ATM and contributes to CHK2 activation (Goldberg *et al*, 2003; Peng & Chen, 2003; Stewart *et al*, 2003; Wu *et al*, 2008). MDC1 has also been shown to be important for CHK1 activation through its interaction with the TOPBP1 scaffold (Wang *et al*, 2011; Leung *et al*, 2013), which may functionally resemble the Rad9–Dpb11 interaction in budding yeast (Fig 2).

ATM promotes the recruitment of 53BP1 through two distinct mechanisms. Similar to yeast Rad9, 53BP1 recognizes γ H2AX through its C-terminal pair of BRCT domains (Fig 2), an interaction which has been controversial, but has recently gained additional support (Baldock *et al*, 2015; Kleiner *et al*, 2015). Nonetheless, the prominent mechanism of 53BP1 recruitment to DNA breaks involves ATM- and γ H2AX-mediated recruitment of MDC1, which becomes phosphorylated by ATM and recruits the E3 ubiquitin ligases RNF8 and RNF168, leading to ubiquitylation of H2A that is recognized directly by 53BP1 (Fig 2; for a detailed review, Hustedt & Durocher, 2016). Notably, this ubiquitylation-dependent

recruiting mechanism is absent in budding yeast. Consistent with an adaptor function for 53BP1, it interacts with CHK2, and the loss of 53BP1 results in reduced ATM-mediated phosphorylation of CHK2 in response to low doses of ionizing radiation (IR; Wang *et al*, 2002). However, 53BP1 appears to regulate activation of the checkpoint through a more complex mechanism, as the physical interaction between CHK2 and 53BP1 rapidly decreases upon IR radiation rather than becoming stabilized (Wang *et al*, 2002), in contrast to the Rad9–Rad53 interaction, which increases after DNA damage in budding yeast. Both 53BP1 and Rad9 play a key role in the control of DNA end resection, highlighting the connection and coordination between checkpoint signaling and the regulation of DNA repair (discussed in detail later in this review; Lazzaro *et al*, 2008; Bunting *et al*, 2010; Chapman *et al*, 2013; Zimmermann *et al*, 2013; Ferrari *et al*, 2015; Liu *et al*, 2017). In both cases, phosphorylation of Rad9/53BP1 by Mec1 or Tel1 in yeast or ATM in humans is essential for suppressing DNA end resection (Bothmer *et al*, 2011; Ferrari

Table 1. Key protein names in budding yeast, fission yeast, and humans.

| | Budding yeast | Fission yeast | Humans |
|------------------------------|---------------|---------------|------------------------|
| Apical PIKK kinase | Mec1 | Rad3 | ATR |
| | Tel1 | Tel1 | ATM |
| | | | DNA-PKcs |
| Downstream checkpoint kinase | Rad53 | Cds1 | CHK2/CHK1 ^a |
| | Chk1 | Chk1 | CHK1? |
| MRN complex | Mre11 | Mre11 | MRE11 |
| | Rad50 | Rad50 | RAD50 |
| | Xrs2 | Nbs1 | NBS1 |
| ATR cofactor | Lcd1 (Ddc2) | Rad26 | ATRIP |
| 9-1-1 complex | Ddc1 | Rad9 | RAD9 |
| | Mec3 | Hus1 | HUS1 |
| | Rad17 | Rad1 | RAD1 |
| ATR activators | Dpb11 | Rad4/Cut5 | TOPBP1 |
| | Dna2 | | |
| | Ddc1 | | ETAA1 |
| Adaptors | Rad9 | Crb2 | MDC1/53BP1 |
| | Mrc1 | Mrc1 | CLASPIN |

^aHuman CHK1 is a functional analog of yeast Rad53.

et al, 2015), and it is possible that 53BP1's function in preventing resection contributes to the stabilization of ATM at breaks, which may indirectly promote ATM–CHK2 signaling.

Post-recruitment events in downstream checkpoint kinase activation

Recruitment to sites of DNA lesions via checkpoint adaptors enables the downstream checkpoint kinases to be directly phosphorylated by the upstream PIKKs, triggering initial kinase activation and subsequent autophosphorylation for further kinase activation. In the case of Rad53, for example, initial phosphorylation by Mec1 or Tel1 promotes its kinase activity by interfering with a kinase auto-inhibitory domain (Fiorani *et al*, 2008), which then stimulates Rad53 to phosphorylate other Rad9-bound Rad53 molecules (Jia-Lin Ma & Stern, 2008). Such trans-autophosphorylation events contribute to dissociate Rad53 from Rad9 and prevent further Rad9 oligomerization (Usui *et al*, 2009). Notably, overexpression of Rad53 in bacteria cells, which lack PIKKs and checkpoint adaptors, results in hyper-activated Rad53 (Gilbert *et al*, 2001). This finding supports a model whereby increased local concentration of Rad53 is enough for activation, with adaptors building increased local concentration at the site of lesions and PIKKs facilitating the initial trigger by reducing the minimal concentration threshold required for activation. In mammals, the apical kinases ATR and ATM drive the key events leading to activation of CHK1 and CHK2, respectively. ATR- and ATM-mediated phosphorylation not only recruits CHK1 and CHK2 to sites of DNA lesions, but also directly phosphorylates these downstream kinases to promote their activation. Like in yeast, these priming phosphorylation events are mainly required to relieve inhibitory domains or to drive monomer-to-oligomer kinase transition (reviewed in

Bartek & Lukas, 2003). ATM-mediated phosphorylation of CHK2 at threonine 68, an established marker of CHK2 activation, allows for the dimerization of two inactive CHK2 monomers and for their subsequent trans- and cis-phosphorylation (Ahn *et al*, 2000; Xu *et al*, 2002; Schwarz *et al*, 2003). CHK2 dimerization is a transient state, since the multiple trans- and cis-autophosphorylation events promote rapid dimer dissociation, leading to full active monomers (Ahn & Prives, 2002; Xu *et al*, 2002; Cai *et al*, 2009). Similar to CHK2, CHK1 activation is a multistep process that requires ATR phosphorylation at serine 317 and serine 345; however, unlike Rad53 or CHK2, CHK1 activation does not appear to involve dimerization or oligomerization (Liu *et al*, 2000; Zhao & Piwnicka-Worms, 2001).

Substrates mediating the core DNA damage signaling responses

Once activated, DNA damage signaling kinases mediate hallmark responses that include the arrest of the cell cycle, inhibition of origin firing, protection and restart of stalled replication forks, induction of a transcriptional response, initiation of apoptosis, and control of dNTP levels. More recent work has demonstrated that the DNA damage signaling kinases also regulate a range of other processes, such as autophagy, gene gating, chromosome mobility, transcription–replication conflicts, and many more whose mechanistic connections to DNA damage signaling and degrees of conservation across eukaryotes remain less clear. Here, we focus on a select set of core conserved functions of the DNA damage signaling kinases with defined substrates, delineating the parallels between budding yeast and humans (Fig 3).

Cell cycle control

The most classical and widely known function of DNA damage signaling kinases is the imposition of a cell cycle arrest that prevents entry into mitosis. Exactly how this arrest is imposed in budding yeast remains elusive. The paradigm for how cell cycle arrest occurs comes from a series of works conducted in fission yeast and metazoans in 1997 (Furnari *et al*, 1997; Peng *et al*, 1997; Sanchez *et al*, 1997; Weinert, 1997), which revealed that DNA damage signaling inhibits CDC25, a phosphatase that removes inhibitory phosphorylation at a key tyrosine residue in mitotic-CDK (M-CDK; Fig 3). In addition, DNA damage signaling kinases stimulate WEE1 (O'Connell *et al*, 1997; Boddy *et al*, 1998; Lee *et al*, 2001), a kinase responsible for phosphorylating the same inhibitory tyrosine site (Fig 3; Mueller *et al*, 1995). While DNA damage signaling in budding yeast does not appear to impinge upon the Cdc25 phosphatase homolog Mih1, the budding yeast kinase Swe1 (the WEE1 homolog) is regulated similar to its counterpart in higher eukaryotes and fission yeast. Swe1 is likely phosphorylated and activated by DNA damage signaling, which results in inhibition of M-CDK (Fig 3; Edenberg *et al*, 2015; Palou *et al*, 2015). DNA damage signaling in budding yeast is also able to suppress M-CDK activity through additional redundant mechanisms that remain unclear (Palou *et al*, 2015). In addition to inhibiting M-CDK-dependent activation of the anaphase-promoting complex (APC/C), DNA damage signaling in budding yeast more directly inhibits the onset of anaphase through Chk1, which




Figure 2. Recruitment of DNA damage signaling kinases and adaptor proteins to DNA lesions: conserved features between budding yeast and humans.

Phosphorylation and adaptor proteins play a key role in the recruitment of downstream checkpoint kinases. The colored ovals indicate phosphorylation events mediated by DNA damage signaling kinases (see kinase key). The orange lines indicate protein–protein interactions promoted by the indicated phosphorylation events (also methylation (me) or ubiquitylation (Ub)). Activation of the downstream checkpoint kinases by the apical PIKK kinases requires adaptor proteins (outlined in green). In most cases, these adaptor proteins act as scaffolds to directly bind to and recruit the downstream checkpoint kinase. The model, mostly based on extensive work in yeast, posits that the recruitment of the downstream checkpoint kinase to the proximity of the apical PIKK kinase enables the phosphorylation and activation of the downstream checkpoint kinase. In addition to activating the downstream checkpoint kinase, phosphorylation events mediated by the apical PIKK kinases are critical for scaffold assembly, often promoting protein–protein interactions. Accordingly, a conserved feature of several adaptor proteins in budding yeast and humans is the presence of protein domains responsible for binding phosphorylated proteins (FHA and BRCT domains). Notably, other kinases such as CDK and CK2 also catalyze phosphorylation events involved in adaptor recruitment, although these events are often not induced by DNA damage. For DNA-PKcs, while this kinase has been implicated in the phosphorylation of H2AX and 53BP1, it does not seem to be involved in CHK2 phosphorylation.

phosphorylates and stabilizes the anaphase-inhibiting Pds1/securin protein (Fig 3; Sanchez *et al*, 1999; Wang *et al*, 2001), preventing the separation of sister chromatids. Moreover, Rad53 can influence the mitotic spindle assembly checkpoint (SAC) through inhibition of the polo-like kinase Cdc5 (Fig 3; Sanchez *et al*, 1999; Valerio-Santiago *et al*, 2013; Zhang *et al*, 2009). Rad53-dependent inactivation of Cdc5, particularly in response to telomere damage, prevents Cdc5's phosphorylation of the spindle assembly checkpoint protein Bfa1 (Valerio-Santiago *et al*, 2013); however, it is unclear whether Rad53 phosphorylates Cdc5 directly (Fig 3). The inhibition of Cdc5 by DNA damage signaling also indirectly influences the resolution of joint molecules prior to M phase, as Cdc5 activity is important for promoting the activity of the Mus81 resolvase (Szakal & Branzel, 2013). Cdc5 has also been implicated in the down-regulation of the DNA damage response (Donnianni *et al*, 2010; Vidanes *et al*, 2010). A more in-depth

review exploring Cdc5's complex relationship with the DNA damage response in yeast can be found in Botchkarev and Haber (2018). Similar to its budding yeast counterpart, the mammalian polo-like kinase PLK1 is also inhibited by DNA damage signaling, albeit in an indirect manner (Fig 3; Bruinsma *et al*, 2017; Qin *et al*, 2013).

A critical mediator of cell cycle arrest in humans is the p53 transcription factor, whose classical function is to trigger the apoptotic program (reviewed in Chen, 2016). p53 is directly phosphorylated and stabilized by all the DNA damage signaling kinases, eliciting a p53-dependent transcriptional response that impacts the DNA damage response (reviewed in Kruse & Gu, 2009). p53-mediated expression of the CDK inhibitor protein p21 represents the primary mechanism by which p53 blocks progression through the cell cycle (Fig 3; Harper *et al*, 1993, 1995). Apical kinases in mammalian DNA damage signaling also phosphorylate the p53 inhibitor Mdm2, impairing its ability to promote




Figure 3. Yeast-to-human parallels in core checkpoint responses mediated by DNA damage signaling.

Substrate map highlighting the phosphorylation events involved in core DNA damage signaling responses in yeast and humans (see text for detailed discussion of each substrate). Conserved or functionally analogous phosphorylation events are positioned parallel to each other. The colored ovals indicate phosphorylation events mediated by DNA damage signaling kinases (see “kinase-dependency” key). The arrows or lines that emanate from the colored ovals represent the role phosphorylation plays in regulating that protein (see “role of phosphorylation” key). Question marks indicate uncertainty, either in the functionality of the phosphorylation event or in the identity of the kinase or substrate. Arrows that impinge on CDK demonstrate how DNA damage signaling can indirectly inhibit CDK activity.

p53 nuclear export and its subsequent degradation (Mayo *et al*, 1997; Maya *et al*, 2001; Shinozaki *et al*, 2003). Unlike most factors covered in Fig 3, *Saccharomyces cerevisiae* does not have clear p53 or Mdm2 homologs. That said, for reasons discussed in the section on dNTP regulation, the DNA damage signaling-mediated control of p53 might functionally resemble the control of the Crt1 transcription regulator by Rad53 and Dun1 (Huang *et al*, 1998).

Fork stability and protection

During DNA replication, stalled replication forks may be targeted and degraded by nucleases. Since fork degradation pathways impair replication fork restart after stalling, leading to persistent DNA lesions, they are considered a major driving force of genomic instability (recently reviewed in Pasero & Vindigni, 2017; Patel & Weiss, 2018). In *S. cerevisiae*, Rad53 is believed to play a major role in fork stability. It is worth mentioning that Rad53 is essential, and that the

lethality of *rad53* cells can be rescued by deletion of *SML1*, an inhibitor of the ribonucleotide reductase enzyme responsible for catalyzing the rate-limiting step in DNA precursor synthesis (Zhao *et al*, 1998). Nonetheless, even in the absence of *SML1*, *Rad53* mutants are exquisitely sensitive to chemical agents that damage DNA or stall DNA replication forks (Allen *et al*, 1994; Sanchez *et al*, 1999; Gunjan & Verreault, 2003). This sensitivity to DNA damaging agents has been primarily attributed to *Rad53*'s role in stabilizing and restarting stalled replication forks (reviewed in detail here: Segurado & Tercero, 2009). *Rad53* protects stalled replication forks from nucleolytic processing by phosphorylating and inhibiting the *Exo1* exonuclease (Fig 3; Cotta-Ramusino *et al*, 2005; Morin *et al*, 2008; Segurado & Diffley, 2008). In addition, *Rad53* regulates the *Pif1* and *Rrm3* helicases, potentially limiting replication fork reversal and subsequent fork degradation (Fig 3; Rossi *et al*, 2015). However, precisely how DNA damage signaling maintains replication fork integrity is unknown and represents a fundamental knowledge gap in the field. Recent work has revealed that *Rad53* phosphorylates the replicative helicase component *Cdc45*, which in turn recruits and stabilizes *Rad53* at replication complexes (Can *et al*, 2018). Identification of additional key substrates and recruiting mechanisms will be necessary to deconstruct *Rad53*'s functions in maintaining fork stability, which may involve a range of redundant phosphorylation events in more than one essential replisome protein.

Mammalian DNA damage signaling kinases also play key roles in protecting stalled replication forks (Fig 3). Human *EXO1* is phosphorylated by *ATR*, which promotes ubiquitylation-dependent degradation of *EXO1* and prevents chromosome fragmentation due to unrestrained *EXO1* processing activity (El-Shemerly *et al*, 2008; Tomimatsu *et al*, 2017). *CHK1* also phosphorylates *EXO1* directly on serine 746, creating a docking site for binding to 14-3-3 proteins, which prevent recruitment of *EXO1* to chromatin and limit *EXO1* action at stalled forks (Engels *et al*, 2011; Li *et al*, 2019). In addition, *ATR* governs the recruitment and/or stability of several helicases important for remodeling the stalled fork and promoting fork restart. For example, *ATR* phosphorylates the Werner syndrome helicase *WRN* at multiple S/T-Q sites, promoting *WRN*-RPA co-localization at replication stress sites (Ammazzalorso *et al*, 2010). Mutation of these *ATR* sites causes stalled fork breakage and severely impacts the resumption of DNA replication. Several studies have demonstrated that the Bloom syndrome helicase, *BLM*, is directly phosphorylated by *ATR* and that this phosphorylation is important for both promoting replication fork restart after HU-mediated arrest and suppressing dormant origin firing (Davies *et al*, 2004, 2007). *CHK1* may also constitutively phosphorylate *BLM* to prevent proteasome-dependent *BLM* degradation, suppress chromatin bridge formation, and promote an interaction with *53BP1* (Sengupta *et al*, 2004; Tripathi *et al*, 2007; Kaur *et al*, 2010; Petsalaki *et al*, 2014). Interestingly, in yeast, *Rad53* and *Sgs1* (a *BLM* ortholog) physically interact (Hegnauer *et al*, 2012), and *Sgs1* has been reported to display a strong genetic interaction with *Rad9*, *53BP1*'s yeast homolog (Nielsen *et al*, 2013). Finally, David Cortez's group described an additional mechanism of fork stability regulation through the *ATR*'s phosphorylation of the *SMARCAL1* helicase (Couch *et al*, 2013). *ATR*-dependent phosphorylation of *SMARCAL1* inhibits fork remodeling activity, preventing subsequent pathological fork degradation by active nucleases (Couch *et al*, 2013;

Kolinjivadi *et al*, 2017). Collectively, these examples highlight how yeast and human DNA damage signaling kinases protect replication fork integrity via the direct regulation of nucleases and helicases.

Inhibition of origin firing

One mechanism for preventing genome instability during replication stress is to inhibit the firing of late origins (reviewed in McIntosh & Blow, 2012; Yekezare *et al*, 2013). DNA damage signaling kinases control origin firing through two main pathways, which are conserved from yeast to higher eukaryotes (Fig 3). During genotoxic stress in S phase, *Rad53* phosphorylates *Dbf4*, a subunit of the *Dbf4*-dependent kinase (DDK) complex, and the *Sld3* component of pre-RCs (pre-replication complexes) to inhibit the firing of late-replicating origins (Lopez-Mosqueda *et al*, 2010; Zegerman & Diffley, 2010), thereby slowing the progression of DNA synthesis and preventing the exhaustion of RPA (Toledo *et al*, 2013, 2017). While the mechanism by which *Rad53* phosphorylation regulates DDK is unknown, the phosphorylation of *Sld3* by *Rad53* is thought to prevent the recruitment of *Dpb11* to primed replication origins (Lopez-Mosqueda *et al*, 2010; Zegerman & Diffley, 2010). In vertebrates, the *Dpb11* ortholog *TOPBP1* docks to CDK2-phosphorylated *Treslin* (the *Sld3* ortholog) to mediate origin firing (Kumagai *et al*, 2010; Boos *et al*, 2011). Conditions that perturb DNA replication and trigger *CHK1* activation have been shown to disrupt the *Treslin*-*TOPBP1* interaction (Boos *et al*, 2011) in a mechanism analogous to *Rad53*-dependent control of the *Sld3*-*Dpb11* interaction in *S. cerevisiae*. *CHK1* has also been shown to directly phosphorylate *Treslin* in *Xenopus* egg extracts, inhibiting DNA replication (Guo *et al*, 2015). Metazoan *ATR*/*ATM* and *CHK1* phosphorylate the *DBF4* subunit of DDK, inhibiting pre-RC assembly and origin firing (Costanzo *et al*, 2003; Heffernan *et al*, 2007; Lee *et al*, 2012). *CHK1* is also assumed to inhibit origin firing indirectly by suppressing CDK activity via the inhibition of *CDC25* (Shechter *et al*, 2004; Sorensen & Syljuasen, 2012). Consistent with this assumption, inhibition of CDK rescues replication stress sensitivity of cells lacking *ATR* or *CHK1* signaling, likely by reducing origin firing and preventing RPA exhaustion (Toledo *et al*, 2013; Dugrawala *et al*, 2015). Paradoxically, *Mec1* (and potentially *ATR*/*ATM*) also promotes DNA replication by phosphorylating core components of the replisome, such as the MCM helicase (Cortez *et al*, 2004; Yoo *et al*, 2004; Randell *et al*, 2010).

Transcriptional control

In yeast, *Rad53* plays a central role in the transcriptional response to DNA damage and replication stress. In G1, *Rad53* was proposed to influence the timing of START by phosphorylating *Swi6*, a component of the MBF/SBF transcription factor (analog of human E2F; Sidorova & Breen, 1997, 2003). *Rad53* can also regulate the transcription of MBF/SBF targets at the G1/S transition and during S phase through phosphorylation and inhibition of the *Nrm1* transcriptional repressor protein, promoting transcription of the largest set of co-regulated genes in the DNA damage response (Fig 3; Bastos de Oliveira *et al*, 2012; de Bruin *et al*, 2008; Travesa *et al*, 2012). In addition, *Dun1* up-regulates the transcription of a specific set of DNA damage-induced genes, including subunits of ribonucleotide reductase (RNR), by inducing the phosphorylation and inactivation of the transcriptional repressor protein, *Crt1* (Huang *et al*, 1998; Fig 3). In-depth transcriptome analyses performed in budding

yeast suggest that DNA damage signaling kinases may impinge upon other, as yet unknown, transcription factors (Jaehnig *et al*, 2013).

DNA damage signaling kinases profoundly impact multiple transcriptional programs in mammals (Fig 3). The E2F pathway (analogous to yeast SBF/MBF) represents a major transcriptional system regulated by DNA damage signaling kinases. ATM-, ATR-, and CHK2-dependent phosphorylation activates E2F1 (Lin *et al*, 2001; Stevens *et al*, 2003), whereas CHK1 inhibits the transcriptional repressor E2F6 through direct phosphorylation (Bertoli *et al*, 2013a). Interestingly, CHK1's inhibition of E2F6 resembles the Rad53-dependent inhibition of the *S. cerevisiae* transcriptional repressor Nrm1, as discussed above (more details can be found here: Bertoli *et al*, 2013b). Similar to yeast, E2F-dependent transcription in humans includes a large set of genes that play a role in DNA replication, cell cycle, and DNA repair (reviewed here: Bracken *et al*, 2004; Poppy Roworth *et al*, 2015). Not surprisingly, inhibition of the ATR–CHK1 pathway causes major remodeling of the proteome via control of the E2F transcriptional circuit and strongly impacts the DNA repair machinery (Kim *et al*, 2018).

Finally, recent work in mammalian cells has revealed a role for ATR in enforcing a checkpoint between the S and G2 phases of the cell cycle. ATR suppresses CDK-dependent phosphorylation and activation of the FOXM1 transcription factor by limiting CDK activity during S phase (Fig 3), thereby preventing a premature expression of G2-specific genes (Saldivar *et al*, 2018). While the function of the FOXM1 transcription factor is conserved in yeast (Pic-Taylor *et al*, 2004; Murakami *et al*, 2010), it is unclear whether the budding yeast checkpoint kinases also restrict the expression of G2/M-specific transcripts through an analogous mechanism (Fig 3).

Regulation of dNTP production

In budding yeast, the regulation of dNTP production is considered an essential function of DNA damage signaling. This is supported by the fact that the lethality caused by deletion of *MEC1* or *RAD53* can be suppressed by deletion of *SML1*, an inhibitor of the ribonucleotide reductase enzyme complex, which catalyzes the limiting step in dNTP production (Zhao *et al*, 1998). The Rad53–Dun1 signaling axis induces the activity of the RNR complex by controlling several of its subunits via distinct mechanisms (Allen *et al*, 1994; Bashkirov *et al*, 2003; Fig 3). Once activated by Rad53, Dun1 directly phosphorylates Sml1, resulting in its degradation and the activation of the Rnr1 subunit (Zhao *et al*, 2001; Zhao & Rothstein, 2002; Lee *et al*, 2008). In addition, Dun1-mediated phosphorylation of Dif1, an *SML1* paralog, results in Dif1's degradation and the subsequent export of the Rnr2–Rnr4 subunits from the nucleus to the cytosol, which enables the formation of the active RNR complex (Lee *et al*, 2008). Dun1 also up-regulates the transcription of *RNR3* by inducing the phosphorylation and inactivation of the transcriptional repressor protein, Crt1 (Huang *et al*, 1998; Fig 3). Rad53 also plays a Dun1-independent role in RNR control. For example, Rad53's phosphorylation of the transcriptional repressor protein Nrm1, upon exposure to replication stress, results in the MBF-mediated induction of *RNR1* transcription (Bastos de Oliveira *et al*, 2012; Huang *et al*, 1998; Travesa *et al*, 2012; Fig 3). Since cells lacking Dun1 are viable, in contrast to *rad53Δ* cells, it is possible that the essential function of Rad53 is due to both its Dun1-dependent and Dun1-independent roles in modulating RNR or, alternatively, a combination of

Rad53's function in dNTP regulation (via Dun1) and other roles in, for example, replication fork stability.

Despite the critical importance of dNTP regulation in yeast, no clear Sml1, Dif1, or Dun1 orthologs or functional analogs have been identified so far in mammalian cells. Nevertheless, it is intriguing that an extra genomic copy of the *RRM2* gene (RNR regulatory subunit) can extend the lifespan of ATR-deficient mice (Lopez-Contreras *et al*, 2015). Whether ATR directly phosphorylates *RRM2* is to date not clear, although ATR may promote *RRM2* stabilization by regulating the degradation of the Cyclin F subunit of the SCF ubiquitin ligase complex (Fig 3), which prevents SCF-mediated degradation of *RRM2* (D'Angiolella *et al*, 2012). *RRM2* transcription is also induced by E2F and thus can be indirectly up-regulated by the ATR–CHK1 kinases (as discussed in the section “transcriptional control”; Ishida *et al*, 2001). Interestingly, p53 promotes transcription of the *RRM2B* subunit of human RNR (Tanaka *et al*, 2000), and it is tempting to speculate on a potential analogy to the role of yeast Crt1 in inducing *RNR3* transcription (Huang *et al*, 1998). In response to hypoxia, this p53-dependent induction of *RRM2B* is important for maintaining the fidelity of DNA replication in low oxygen conditions (Foskolou *et al*, 2017). However, a direct link to the DNA damage signaling kinases has yet to be made.

Control of DNA repair by DNA damage signaling kinases

The control of DNA repair is a core function of DNA damage signaling kinases. Mounting genetic evidence from budding yeast illustrates the central role of DNA damage signaling to maintaining genome integrity. For example, using a genetic assay to assess chromosome instability in yeast, known as the gross chromosomal rearrangement (GCR) assay, the Kolodner Lab showed that cells lacking *Mec1* and *Tel1* exhibit extremely high rates of GCRs (Myung *et al*, 2001). In fact, *mec1Δ tel1Δ* cells are some of the most genetically unstable strains isolated to date. These cells undergo GCRs at a rate of five orders of magnitude higher than wild-type cells (Myung *et al*, 2001). As discussed below, current evidence supports a model whereby the ability of *Mec1* and *Tel1* to suppress GCRs and related forms of genetic instabilities is associated with the role these kinases play in directly targeting and regulating components of the DNA repair machinery. In mammals, ATM and ATR are also responsible for maintaining chromosome stability (Xu *et al*, 1996; Brown & Baltimore, 2000); as shown in Fig 4, ATM and ATR phosphorylate and regulate a large set of DNA repair factors. Importantly, the high levels of genetic instability in yeast or mammalian cells lacking *Mec1*/ATR and *Tel1*/ATM have not yet been recapitulated by any combination of mutations of phosphorylation sites, suggesting that the key substrates through which these kinases suppress genomic instabilities remain unclear.

Phosphorylation control of the homologous recombination machinery: substrates and mechanisms

In both yeast and humans, DNA damage signaling kinases phosphorylate proteins in homologous recombination (HR)-directed repair and have been shown to play an important regulatory role. For example, *Mec1* phosphorylates Rtt107, Slx4, and Sgs1, all of which are known to play multiple roles in HR-mediated repair, from the control of resection to the dissolution or resolution of joint




Figure 4.

Figure 4. Yeast-to-human parallels in the regulation of DNA repair proteins by DNA damage signaling: substrates and mechanisms.

Substrate map cataloging the DNA damage signaling events regulating DNA repair proteins (see text for detailed discussion of each substrate). Conserved or analogous substrates involved in related DNA repair pathways are positioned parallel to each other. The colored ovals indicate phosphorylation events mediated by DNA damage signaling kinases (see “kinase-dependency” key). The arrows or lines that emanate from the colored nodes represent the role phosphorylation plays in regulating that protein (see “role of phosphorylation” key). Question marks indicate uncertainty, either in the functionality of the phosphorylation event or in the identity of the kinase or substrate.

chromosome structures (Sarbjana & West, 2014; Hang & Zhao, 2016; Cussiol *et al*, 2017; Guervilly & Gaillard, 2018). Recent work from our laboratory has revealed that phosphorylation of these proteins correlates with the ability of cells to suppress GCRs, suggesting that proper control of HR repair is essential for preventing chromosomal rearrangements (Lanz *et al*, 2018). Phosphorylation of Slx4 by Mec1 promotes DNA end resection, the first step in the HR process (Dibitetto *et al*, 2016; Liu *et al*, 2017). Mechanistically, Mec1 phosphorylation mediates the interaction between Slx4 and Dpb11, a multi-BRCT domain scaffold that bridges Slx4 to the 9-1-1 clamp loaded at 5' recessed junctions (Ohouo *et al*, 2010; Cussiol *et al*, 2015; Fig 4). Such stabilization of Slx4 at DNA lesions, which is also dependent on Rtt107's interaction with phosphorylated H2A (Fig 4), is believed to counteract a resection block promoted by Rad9, thereby allowing long-range resection to occur (Dibitetto *et al*, 2016; Liu *et al*, 2017). Importantly, the role of Rad9 in counteracting resection relies on its oligomerization and Rad53 signaling (Clerici *et al*, 2014; Ferrari *et al*, 2015; Gobbin *et al*, 2015), which are both dependent on Mec1-mediated phosphorylation events. Therefore, Mec1 plays opposing roles in HR, inhibiting resection via the Rad9–Rad53 signaling axis or promoting resection by mediating the Slx4–Dpb11 interaction (Fig 4).

Mec1's phosphorylation of Slx4 has also been linked to a role in single-strand annealing (SSA), specifically in regulating cleavage of 3' non-homologous DNA tails by Rad1–Rad10 (Toh *et al*, 2010). In another example, which supports a key role for Mec1 in the control of HR, phosphorylation of multiple S/T-Q residues in Sae2, a pro-resection factor homologous to human CtIP, contributes to Mec1-mediated GCR suppression (Liang *et al*, 2015). Furthermore, Mec1 phosphorylates the strand exchange factor Rad55, and Rad53 promotes the phosphorylation and DNA binding of the Tid1/Rdh54 translocase (Fig 4; Bashkirov *et al*, 2006; Ferrari *et al*, 2013; Herzberg *et al*, 2006). Mec1 also phosphorylates Rad51 at serine 192, likely supporting its ATPase activity and capacity for strand invasion (Fig 4; Flott *et al*, 2011).

The ssDNA binding protein RPA, which plays a critical role in coating resected DNA ends during the homologous recombination process, is one of the most established targets of the PIKKs. In both yeast and humans, RPA, through an interaction with Rad52, is thought to promote the loading of Rad51 on ssDNA (Park *et al*, 1996; Sugiyama *et al*, 1998; Sugiyama & Kowalczykowski, 2002). How PIKK phosphorylation regulates RPA function is not well understood, although studies point to a role in homologous recombination. RPA phosphorylation was initially proposed to promote the interaction between RPA and Rad52 (Deng *et al*, 2009); however, recent work using single-molecule imaging reports that RPA phosphorylation inhibits DNA end resection via inhibition of the BLM helicase (Soniat *et al*, 2019). RPA phosphorylation in human cells has also been reported to impact DNA synthesis under stress (Vassin *et al*, 2009), the recruitment of factors to stalled replication forks (Murphy *et al*, 2014), and the

imposition of the cell cycle checkpoint (Vassin *et al*, 2009). In budding yeast, mutation of the residues in RPA targeted by Mec1 and Tel1 has yet to yield an observable phenotype. Overall, the accumulated evidence supports a model whereby DNA damage signaling plays a key role in the control of the HR machinery by targeting multiple components that participate in discrete steps of the HR process (Fig 4).

ATR is a key regulator of homologous recombination-mediated repair

ATR's primary action occurs in S phase, a period during the cell cycle where a sister DNA template is available for homology-directed repair, and it functions mainly as a pro-HR kinase. Depletion or inhibition of ATR impairs the ability of cells to utilize HR and leads to synergistic cell death with replication stress-inducing drugs (Wang *et al*, 2004a; Vriend *et al*, 2016; Yazinski *et al*, 2017; Kim *et al*, 2018). One model proposes that during HR-mediated repair, an ATM-to-ATR transition occurs, where ATM initiates resection and triggers the ATR activation that governs the later steps of homologous recombination (Cuadrado *et al*, 2006; Shiotani & Zou, 2009). Our laboratory proposed that ATR drives HR by promoting the stabilization of the pro-resection factor BRCA1 at DNA lesions via interaction with TOPBP1 (human ortholog of Dpb11), which may counteract recruitment of the anti-resection factor 53BP1 to sites of DNA lesions (Liu *et al*, 2017). It is tempting to speculate that this mechanism could be analogous to the Mec1-mediated Dpb11–Slx4 interaction in yeast, which is also important for DSB resection (Dibitetto *et al*, 2016; Liu *et al*, 2017; Ohouo *et al*, 2010; Fig 4). A recent study from the Zou Lab provided additional insights into how ATR promotes HR-mediated repair. They showed that while initial DSB resection requires CDK activity, later steps in HR require a “CDK-to-ATR switch” to promote proper recruitment of the key HR factors PALB2 and BRCA2, which are required for strand invasion. Mechanistically, ATR activated at resected ends recruits PALB2–BRCA2 to DNA damage sites by phosphorylating PALB2 on serine 59 to promote its interaction with BRCA1 (Buisson *et al*, 2017). At the same time, ATR inhibits CDK to prevent a CDK-dependent phosphorylation at PALB2's serine 64 that inhibits the BRCA1–PALB2 interaction (Buisson *et al*, 2017). Finally, the control of E2F transcription via ATR–CHK1 signaling strongly impacts the ability of cells to utilize HR-mediated repair by ensuring proper expression of key components of the HR machinery (Kim *et al*, 2018). Similar to yeast, RAD51 deposition and RAD51-mediated strand invasion are regulated by DNA damage signaling kinases. CHK1 has been shown to drive the formation of RAD51 foci at stalled replication forks, likely through direct phosphorylation (Sorensen *et al*, 2005). CHK1 and CHK2 may also be important to regulate the association of RAD51 with BRCA2 (Bahassi *et al*, 2008). Moreover, RAD51C, a putative homolog of yeast Rad55, associates with XRCC3, whose phosphorylation by ATM and ATR in response to IR is important for HR regulation (Somyajit *et al*, 2013).

ATM and DNA-PKcs in resection control

ATM signaling blocks DNA end resection via recruitment of an anti-resection complex formed by 53BP1-RIF1-REV7-SHIELDIN (Setiaputra & Durocher, 2019). This process is achieved through a defined order of events: phosphorylation of H2AX and MDC1, followed by recruitment of the RNF8 and RNF168 ubiquitin ligases, followed by ubiquitylation of H2A and recruitment of 53BP1, whose phosphorylation by ATM serves as a docking platform for RIF1 and, indirectly, the REV7-SHIELDIN complex (Fig 4). In addition, ATM limits resection by phosphorylating the proteasomal shuttle factor UBQLN4, which leads to degradation of MRE11, a component of the MRN complex required for initial steps in resection (Jachimowicz *et al*, 2019). Paradoxically, ATM may also promote resection by phosphorylating CtIP, a key factor required for resection initiation (Shibata *et al*, 2011). DNA-PKcs also plays opposing roles in resection control. Whereas binding of DNA-PKcs to DNA ends has a major anti-resection function by preventing recruitment of the EXO1 nuclease, DNA-PKcs' autophosphorylation is required to promote dissociation from DNA ends, thereby allowing EXO1 binding and resection (Zhou & Paull, 2013). More work is needed to better understand and define the extent to which the actions of ATM and DNA-PKcs impact overall resection in cells.

ATM and DNA-PKcs promote NHEJ

ATM and DNA-PKcs are key players in the control of non-homologous end joining (NHEJ), a pathway for DSB repair in which DNA ends are ligated without the need for a homologous template. ATM and DNA-PKcs play partially redundant roles in promoting NHEJ, as impairment of both kinases results in a stronger NHEJ defect compared with impairment of one of the kinases (Gapud & Sleckman, 2011; Gapud *et al*, 2011; Zha *et al*, 2011). In addition, combined ATM and DNA-PKcs deficiency leads to embryonic lethality in mice and a more severe DSB repair defect during immunoglobulin class-switch recombination than loss of only one of the kinases (Callen *et al*, 2009). The ability of ATM and DNA-PKcs to limit DNA end resection is a key part of their pro-NHEJ function; however, these kinases likely regulate additional processes to promote NHEJ, since chemical inhibition of both kinases severely inhibits NHEJ without inducing resection due to persistent DNA-PKcs at break ends. ATM and DNA-PKcs redundantly phosphorylate several NHEJ repair components, such as XLF, Artemis, and POL λ (Zhang *et al*, 2004; Goodarzi *et al*, 2006; Yu *et al*, 2008; Sastre-Moreno *et al*, 2017) (Fig 4). In addition, DNA-PKcs solely phosphorylates the XRCC4, KU70, KU80, and LIG-IV core components (Chan *et al*, 1999; Lee *et al*, 2004; Wang *et al*, 2004b; Douglas *et al*, 2005; Amiri Moghani *et al*, 2018). However, mutation of these phosphorylation sites, which are dependent on ATM and/or DNA-PKcs, does not impair NHEJ or phenocopy the loss or inhibition of ATM and DNA-PKcs function. Notably, phosphorylation of a non-conventional DNA-PKcs substrate, the transcription factor NR4A, was proposed to be important for NHEJ, as demonstrated by a phosphosite mutation, but the impact of the mutation in NHEJ remains unclear (Malewicz *et al*, 2011).

In *S. cerevisiae*, little is known about the crosstalk between DNA damage signaling kinases and NHEJ repair factors. Tel1 phosphorylates Pol4, stimulating its gap-filling activity and preventing the appearance of deleterious chromosome translocations (Ruiz *et al*, 2013). Another report showed that the Dun1 kinase is also

important for NHEJ by phosphorylating Nej1, a key activator of the NHEJ ligase Dnl4 (Ahnesorg & Jackson, 2007). DNA damage signaling kinases in budding yeast do not appear to impinge upon the NHEJ pathway as extensively as they do in humans and may reflect budding yeast's strong preference for homologous recombination-mediated DNA repair (Jasin & Rothstein, 2013).

Other DNA repair pathways

While DNA damage signaling kinases appear to extensively impinge upon multiple factors and steps of recombinational DNA repair, they also target and regulate other repair mechanisms (Fig 4), suggesting that their action is context-dependent and, in some cases, important in repair pathway choice. In budding yeast, Mec1 phosphorylates the nucleotide excision repair (NER) protein Rad26 and regulates the transcription-coupled NER mode of repair (Taschner *et al*, 2010). In humans, ATR also controls NER by phosphorylating XPA (Wu *et al*, 2006) and, in conjunction with ATM, the histone acetyltransferase HBO1. HBO1 phosphorylation stabilizes its interaction with DDB2, promoting its proteasome-mediated degradation (Matsunuma *et al*, 2016). In addition, ATR phosphorylates POL η at serine 601, facilitating post-replication repair and promoting checkpoint activation at UV damage sites (Gohler *et al*, 2011). In yeast, Mec1 regulates the translesion synthesis polymerase Rev1 potentially through direct phosphorylation (Pages *et al*, 2009). This complements earlier studies, where it was shown that Mec1 is essential for Rev1 binding to chromosomes (Hirano & Sugimoto, 2006; Sabbioneda *et al*, 2007). DNA damage signaling kinases are also important for the execution of mismatch repair (MMR) and base excision repair (BER). ATM phosphorylates the MMR protein MLH1, contributing to its stabilization after DNA damage (Romeo *et al*, 2011). Also, ATM directly phosphorylates the BER proteins TDP1 (Das *et al*, 2009) and MPG (Agnihotri *et al*, 2014), enhancing their DNA repair activity and promoting cell survival after exposure to alkylating agents. DNA-PKcs is also able to phosphorylate TDP1 at serine 81 (Das *et al*, 2009), allowing interaction with the DNA repair protein XRCC1, which is itself a CHK2 target for BER regulation (Chou *et al*, 2008). Finally, Rad53 was shown to be important in break-induced replication (BIR) through the phosphorylation of the Pif1 helicase (Vasianovich *et al*, 2014).

Fanconi anemia pathway

The Fanconi anemia pathway is a genetic network involved in the repair of DNA interstrand crosslinks (ICLs) in the genome (Niraj *et al*, 2019). It combines the action of nucleotide excision repair and homologous recombination factors, several of which are regulated by ATR (Fig 4). ATR plays a crucial role at an early step in the Fanconi anemia pathway by phosphorylating the FANCI protein and inducing ubiquitylation of the FANCI-FANCD2 complex (Ishiai *et al*, 2008; Shigechi *et al*, 2012). The ubiquitylated FANCI-FANCD2 complex serves as a key scaffold for the recruitment of several DNA repair proteins, including structure-specific nucleases and TLS polymerases involved in ICL repair (Knipscheer *et al*, 2009). In addition to promoting repair and replication fork start, phosphorylation of FANCI by ATR modulates DNA replication by inhibiting dormant origin firing (Chen *et al*, 2015). FANCD2 is also phosphorylated by ATM and ATR (Ho *et al*, 2006), whereas CHK1 phosphorylation is important for the FANCD2-BRCA2 interaction (Zhi *et al*, 2009). CHK1 phosphorylates FANCE at threonine 346 and serine 374 to

promote resistance to mitomycin C (Wang *et al.*, 2007). Additionally, ICL resistance has been linked to the ATR-mediated phosphorylation of FANCA at serine 1449 and FANCM at serine 1045 (Collins *et al.*, 2009; Singh *et al.*, 2013; Fig 4).

Coordination of checkpoint and DNA repair: where are we 30 years later?

Following the checkpoint paper by Weinert and Hartwell in 1988 (Weinert & Hartwell, 1988), it was not at all clear how eukaryotic cells sense DNA damage and couple lesion detection to the control of the cell cycle. At that time, Weinert and Hartwell defined checkpoint as “control mechanisms enforcing dependency in the cell cycle”, implying the existence of some undefined regulatory mechanism. While the 1988 paper marks the identification of the first checkpoint factor in budding yeast (Rad9), the checkpoint concept largely preceded the identification of most key components mediating lesion detection and signaling. It was not even clear whether kinases were involved; in fact, the word “kinase” is absent from both the checkpoint papers of 1988 and 1989 (Weinert & Hartwell, 1988; Hartwell & Weinert, 1989). Identification of the apical PIKK kinases and downstream checkpoint kinases, as well as their involvement in mediating DNA damage responses, would not become evident until the mid-1990s. Decades later, our understanding of both the mechanisms of lesion detection and the signaling circuitry connecting lesion detection to regulation of biological processes has radically changed. The field has defined many of the key proteins required for lesion detection and uncovered an extensive network of kinase substrates. The concept that checkpoint is a simple mechanism coupling lesion detection to cell cycle arrest also dramatically changed to incorporate both a range of biological outputs and a network of functional and physical connections. Nonetheless, at its essence, the central concept that DNA damage checkpoints coordinate cell cycle with DNA repair continues to hold true.

A unified model for the coordination of checkpoint and repair in the DSB response

When Weinert and Hartwell first proposed the checkpoint concept, it would be considered improbable that the circuitry responsible for promoting cell cycle arrest also directly and actively controls the DNA repair machinery. However, over the last 30 years, accumulated knowledge from work in yeast and mammals points to a key role for the apical PIKKs and downstream checkpoint kinases in the spatiotemporal coordination of checkpoint responses and DNA repair transactions. The response to DSBs epitomizes such intricate coordination. Mounting evidence supports a model whereby early response to DSBs involves rapid activation of apical PIKK kinases and the establishment of an emergency response that inhibits improper nuclease action (and resection) to prevent improper processing of DNA ends. These early response actions include cell cycle arrest, general inhibition of origin firing (if in S phase), and extensive transcriptional re-programming. Following establishment of a robust checkpoint response and early resection block, cells must decide when to initiate DNA repair, including which repair pathway to utilize and when to resume the cell cycle. This decision is simpler in G1 cells, as the engagement of NHEJ for the repair of

DSBs is highly preferred and can occur as a natural consequence of the resection block. However, for many DNA lesions in S phase and G2, the use of HR-mediated repair is the pathway of choice for error-free repair. Cells must, therefore, oppose the initial anti-resection effects of DNA damage signaling to initiate resection, the first step in HR. Given the importance of the decision to initiate (or not to initiate) resection for repair pathway choice, this step is under intense regulation by phosphorylation. In yeast, Mec1 promotes resection by phosphorylating Slx4 and likely Sae2, opposing its own anti-resection function via Rad53 activation. Interestingly, such dual and opposite functions for Mec1 could be key to providing a highly controllable system in which resection is spatiotemporally fine-tuned. This is particularly important, as the extent of resection impacts which HR-mediated mechanisms will follow [e.g., canonical HR, single-strand annealing, synthesis-dependent strand annealing (SDSA), and break-induced replication (BIR)]. In humans, current evidence points to ATR, ATM, and DNA-PKcs playing an important role in the regulation of resection, although much less is understood compared with yeast. In both yeast and humans, phosphatases are expected to play a key role in counter-balancing the phosphorylation events that impose the early resection block and influence the DNA repair process. The balance between DNA damage signaling kinases and the phosphatases that oppose them is better appreciated for the more canonical aspects of the checkpoint, but more work is needed to understand how this balance affects choices made during the DNA repair process.

In sum, apical kinases perform highly elaborate actions in the spatiotemporal control of checkpoint responses and DNA repair. Since checkpoint functions and early resection blockades can counteract HR-mediated DNA repair mechanisms, the ability of cells to modulate DNA damage signaling is essential for repair pathway control, and, in particular, proper execution of HR-mediated repair. More work is needed to better understand how DNA damage signaling kinases monitor and control subsequent steps in HR (including strand invasion, homology search, and resolution/dissolution), as well as how such regulation ensures the fidelity of HR and prevents the erroneous recombination that can give rise to genetic instability.

What's next? Toward a holistic view of the DNA damage signaling network

In budding yeast and humans, DNA damage signaling kinases target dozens, if not hundreds, of DNA repair proteins, thereby modulating repair pathways. As described above, our understanding of the mechanisms by which these kinases coordinate DNA repair machineries is incomplete. When considering a function associated with the action of a kinase, the gold standard is to demonstrate that the function is impaired by mutation of a discrete set of phosphorylatable residues within a substrate protein(s). In almost all the cases discussed above, the DNA repair-related phenotypes associated with the disruption of kinase function are far stronger than the phenotypes associated with the introduction of phosphosite mutations on the kinase substrates. Thus, either the proper combination of phosphosite mutations has not yet been uncovered, or there are still many more undiscovered phosphorylation events mediated by the DNA damage signaling kinases that are critical for the control of DNA repair. In some cases, fundamental phenotypes associated with

the loss of DNA damage signaling still lack defined substrates and underlying molecular mechanisms. For example, a mechanistic understanding of the genetic instability (mostly monitored as accumulation of GCR) observed upon loss of Mec1 and Tel1 is unknown and represents a significant gap in our understanding of DNA damage signaling in budding yeast. In humans, the lethality and chromosomal fragmentation observed upon inhibition of ATR also lack a clear mechanistic underpinning. Furthermore, while it is established that inhibition of ATM and DNA-PKcs strongly impair NHEJ, it is unclear which phosphorylation events are disrupted upon inhibition and in what ways they are necessary for NHEJ.

Mass spectrometry as a tool to study DNA damage signaling kinases: strengths, weaknesses, and future directions

Mass spectrometry has been instrumental for discovering *in vivo* substrates of the DNA damage signaling kinases (Matsuoka *et al*, 2007; Smolka *et al*, 2007; Bastos de Oliveira *et al*, 2015; Wagner *et al*, 2016; Lanz *et al*, 2018). The ability to quantitatively assess the phosphoproteome of an organism opened the door for unbiased identification of kinase substrates. Phosphoproteomic screens, performed in budding yeast and humans, have identified hundreds of phosphorylation events catalyzed by the DNA damage signaling kinases. Reassuringly, many of these phosphorylation events map to previously known substrate proteins. Excitingly, most of these events occur in substrate proteins not yet studied, many of which are associated with a broad range of nuclear processes, including DNA repair. However, the scope of the DNA damage signaling network revealed by phosphoproteomics raises the question of how many phosphorylation events have tangible biological significance. The ability to distinguish functional phosphorylation from kinase promiscuity represents the primary challenge of large-scale phosphoproteomic datasets. For such an inquiry, it is imperative to generate mutant strains that either lack or constitutively mimic the phosphorylated residues in the substrate protein, with the ultimate goal to phenocopy the effects of the kinase's action or inaction. However, this is equivalent to finding a needle in a haystack, and better strategies are needed to enhance our ability to efficiently and systematically predict functional sites and, when necessary, to dissect functional redundancy and combinatorial effects.

Identification of *in vivo* kinase substrates using mass spectrometry is often performed using quantitative MS-based approaches, primary SILAC, which provide a highly quantitative comparison between the phosphoproteomic profiles of two different cell populations (Bastos de Oliveira *et al*, 2015, 2018). Budding yeast represents an ideal system for mapping DNA damage signaling using mass spectrometry due to its relative simplicity compared with the mammalian system. A range of genetic tools and mutants enable powerful genetic-proteomic strategies to deconstruct the action of apical and downstream kinases. For example, loss of Mec1/ATR results in lethality in most eukaryotic systems, but suppressor mutations that rescue the lethality of *mec1Δ* cells (such as *SML1* deletion) are available in yeast and are vital for the identification of proteins containing Mec1/Tel1- and Rad53-dependent phosphorylation using phosphoproteomics (Smolka *et al*, 2007; Bastos de Oliveira *et al*, 2015; Lanz *et al*, 2018).

Phosphoproteomics has also been used to identify targets of the DNA damage signaling kinases in mammalian cell lines (Matsuoka

et al, 2007; Stokes *et al*, 2007; Bennetzen *et al*, 2010; Pines *et al*, 2011; Beli *et al*, 2012; Kirkpatrick *et al*, 2013; Wagner *et al*, 2016). The sheer complexity of the human phosphoproteome makes in-depth phosphoproteomic analyses more challenging than in yeast. Nonetheless, phosphoproteomic studies, such as the 2007 landmark paper by the Elledge group (Matsuoka *et al*, 2007), have revealed hundreds of potential substrates of DNA damage signaling kinases. One limitation of this work is that it relied on the use of phospho-specific antibodies for the SQ/TQ phosphosite motif, so it did not identify substrates of the downstream DDC kinases CHK1 and CHK2, nor did it identify potentially direct ATR or ATM substrates phosphorylated at non-canonical motifs (non-SQ/TQ). Unlike in yeast, matching a substrate to a specific DNA damage signaling kinase is still a challenge, although the recent development of specific chemical inhibitors of ATR, ATM, and CHK1 has allowed more efficient assignment of kinase dependency. In fact, recent studies have made use of these chemical inhibitors to find ATR-, ATM-, and CHK1-dependent phosphorylation events *in vivo* (Blasius *et al*, 2011; Wagner *et al*, 2016). Of note, the substrates of DNA-PKcs, a third, human-specific PI3 kinase-like kinase, have yet to be extensively profiled by MS, although low-throughput studies have uncovered substrate proteins associated with NHEJ (DNA-PKcs is reviewed in detail here: Blackford & Jackson, 2017).

An alternative approach to screening for substrates of a kinase-of-interest involves the mutation of gate-keeping residues in the kinase's ATP-binding pocket, which enables the accommodation and subsequent utilization of bulky ATP analogs (Hertz *et al*, 2010). The addition of an analog-compatible kinase to human cell extracts results in the *in vitro* modification of its substrate proteins, which are then isolated and identified based on the covalent attachment of analog phosphate groups. This chemical-genetics approach (commonly referred to as the "Shokat method") has been used to map the signaling network of several CDKs (Blethrow *et al*, 2008; Chi *et al*, 2008) and, more recently, to identify CHK1 substrate proteins in human cell extracts (Blasius *et al*, 2011). However, these chemical-genetic strategies require the generation of mutant kinase alleles that are able to utilize bulky ATP analogs, and the kinase labeling reactions normally take place in cell lysates rather than *in vivo*. Also, efforts to apply the Shokat method to the upstream PIKKs have thus far been unsuccessful, as ATM and ATR appear to be less amenable to mutations in their ATP-binding pockets.

Overall, phosphoproteomic studies have revealed that the human DNA damage signaling network is extensive, consisting of many nuclear proteins, but also non-nuclear proteins. In addition to the established DNA repair-related substrates previously described in low-throughput studies, phosphoproteomic analyses have uncovered a large proportion of substrates involved in transcription and RNA processing, although, like in yeast, the functional significance of these phosphorylation events remains unclear. While the work done thus far in mammalian systems has yielded many candidate substrates, the DNA damage signaling kinases may operate differently in different cell types. Thus, it will be important for future investigations to thoroughly assess how the DNA damage signaling network varies in different cell types and pathological statuses. Attaining full coverage of the phosphoproteome is also a concern, as available phosphoproteome datasets do not cover all phosphorylation events in a cell due to technical limitations with mass

spectrometry (Engholm-Keller & Larsen, 2013). Currently, however, instruments have become more sensitive, and the ability to use mass spectrometry to cover the full phosphoproteome may become a reality in the near future. Looking ahead, it will be crucial to develop more efficient approaches for identifying functional phosphorylation events within the large phosphoproteomic datasets. One possibility is to use structural prediction analysis to identify phosphorylation events that are likely to impair protein function (such as disrupting protein–protein or protein–DNA interactions) (preprint: Lanz *et al*, 2019). In conclusion, a holistic understanding of the role and action of DNA damage signaling kinases will require powerful technologies capable of quantitatively monitoring the full extent of the phosphoproteome in combination with systematic approaches for functional and mechanistic analyses.

Acknowledgements

We thank Ethan Sanford and Robert Weiss for careful reading of the article and suggestions. This work was supported by M.B.S. grants from the National Institutes of Health (R01GM097272 and R01HD095296).

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Abreu CM, Kumar R, Hamilton D, Dawdy AW, Creavin K, Eivers S, Finn K, Balsbaugh JL, O'Connor R, Kiely PA *et al* (2013) Site-specific phosphorylation of the DNA damage response mediator rad9 by cyclin-dependent kinases regulates activation of checkpoint kinase 1. *PLoS Genet* 9: e1003310
- Agnihotri S, Burrell K, Buczkowicz P, Remke M, Golbourn B, Chornenkyy Y, Gajadhar A, Fernandez NA, Clarke ID, Barszczyk MS *et al* (2014) ATM regulates 3-methylpurine-DNA glycosylase and promotes therapeutic resistance to alkylating agents. *Cancer Discov* 4: 1198–1213
- Ahn JY, Schwarz JK, Piwnicka-Worms H, Canman CE (2000) Threonine 68 phosphorylation by ataxia telangiectasia mutated is required for efficient activation of Chk2 in response to ionizing radiation. *Cancer Res* 60: 5934–5936
- Ahn J, Prives C (2002) Checkpoint kinase 2 (Chk2) monomers or dimers phosphorylate Cdc25C after DNA damage regardless of threonine 68 phosphorylation. *J Biol Chem* 277: 48418–48426
- Ahnesorg P, Jackson SP (2007) The non-homologous end-joining protein Nej1p is a target of the DNA damage checkpoint. *DNA Repair (Amst)* 6: 190–201
- Alcasabas AA, Osborn AJ, Bachant J, Hu F, Werler PJ, Bousset K, Furuya K, Diffley JF, Carr AM, Elledge SJ (2001) Mrc1 transduces signals of DNA replication stress to activate Rad53. *Nat Cell Biol* 3: 958–965
- Allen JB, Zhou Z, Siede W, Friedberg EC, Elledge SJ (1994) The SAD1/RAD53 protein kinase controls multiple checkpoints and DNA damage-induced transcription in yeast. *Genes Dev* 8: 2401–2415
- Amiri Moghani AR, Sharma MK, Matsumoto Y (2018) In cellulo phosphorylation of DNA double-strand break repair protein XRCC4 on Ser260 by DNA-PK. *J Radiat Res* 59: 700–708
- Ammazzalorso F, Pirzio LM, Bignami M, Franchitto A, Pichierri P (2010) ATR and ATM differently regulate WRN to prevent DSBs at stalled replication forks and promote replication fork recovery. *EMBO J* 29: 3156–3169
- van Attikum H, Fritsch O, Hohn B, Gasser SM (2004) Recruitment of the INO80 complex by H2A phosphorylation links ATP-dependent chromatin remodeling with DNA double-strand break repair. *Cell* 119: 777–788
- Bacal J, Moriel-Carretero M, Pardo B, Barthe A, Sharma S, Chabes A, Lengronne A, Pasero P (2018) Mrc1 and Rad9 cooperate to regulate initiation and elongation of DNA replication in response to DNA damage. *EMBO J* 37: e99319
- Bahassi EM, Ovesen JL, Riesenberger AL, Bernstein WZ, Hasty PE, Stambrook PJ (2008) The checkpoint kinases Chk1 and Chk2 regulate the functional associations between hBRCA2 and Rad51 in response to DNA damage. *Oncogene* 27: 3977–3985
- Baldock RA, Day M, Wilkinson OJ, Cloney R, Jeggo PA, Oliver AW, Watts FZ, Pearl LH (2015) ATM localization and heterochromatin repair depend on direct interaction of the 53BP1-BRCT2 domain with gammaH2AX. *Cell Rep* 13: 2081–2089
- Bartek J, Lukas J (2003) Chk1 and Chk2 kinases in checkpoint control and cancer. *Cancer Cell* 3: 421–429
- Bashkirov VI, Bashkirova EV, Haghnazari E, Heyer WD (2003) Direct kinase-to-kinase signaling mediated by the FHA phosphoprotein recognition domain of the Dun1 DNA damage checkpoint kinase. *Mol Cell Biol* 23: 1441–1452
- Bashkirov VI, Herzberg K, Haghnazari E, Vlasenko AS, Heyer WD (2006) DNA damage-induced phosphorylation of Rad55 protein as a sentinel for DNA damage checkpoint activation in *S. cerevisiae*. *Methods Enzymol* 409: 166–182
- Bastos de Oliveira FM, Harris MR, Brazauskas P, de Bruin RA, Smolka MB (2012) Linking DNA replication checkpoint to MBF cell-cycle transcription reveals a distinct class of G1/S genes. *EMBO J* 31: 1798–1810
- Bastos de Oliveira FM, Kim D, Cussiol JR, Das J, Jeong MC, Doerfler L, Schmidt KH, Yu H, Smolka MB (2015) Phosphoproteomics reveals distinct modes of Mec1/ATR signaling during DNA replication. *Mol Cell* 57: 1124–1132
- Bastos de Oliveira FM, Kim D, Lanz M, Smolka MB (2018) Quantitative analysis of DNA damage signaling responses to chemical and genetic perturbations. *Methods Mol Biol* 1672: 645–660
- Beli P, Lukashchuk N, Wagner SA, Weinert BT, Olsen JV, Baskcomb L, Mann M, Jackson SP, Choudhary C (2012) Proteomic investigations reveal a role for RNA processing factor THRAP3 in the DNA damage response. *Mol Cell* 46: 212–225
- Bennetzen MV, Larsen DH, Bunkenborg J, Bartek J, Lukas J, Andersen JS (2010) Site-specific phosphorylation dynamics of the nuclear proteome during the DNA damage response. *Mol Cell Proteomics* 9: 1314–1323
- Bertoli C, Klier S, McGowan C, Wittenberg C, de Bruin RA (2013a) Chk1 inhibits E2F6 repressor function in response to replication stress to maintain cell-cycle transcription. *Curr Biol* 23: 1629–1637
- Bertoli C, Skotheim JM, de Bruin RA (2013b) Control of cell cycle transcription during G1 and S phases. *Nat Rev Mol Cell Biol* 14: 518–528
- Blackford AN, Jackson SP (2017) ATM, ATR, and DNA-PK: the trinity at the heart of the DNA damage response. *Mol Cell* 66: 801–817
- Blankley RT, Lydall D (2004) A domain of Rad9 specifically required for activation of Chk1 in budding yeast. *J Cell Sci* 117: 601–608
- Blasius M, Forment JV, Thakkar N, Wagner SA, Choudhary C, Jackson SP (2011) A phospho-proteomic screen identifies substrates of the checkpoint kinase Chk1. *Genome Biol* 12: R78
- Blethrow JD, Glavy JS, Morgan DO, Shokat KM (2008) Covalent capture of kinase-specific phosphopeptides reveals Cdk1-cyclin B substrates. *Proc Natl Acad Sci USA* 105: 1442–1447
- Boddy MN, Furnari B, Mondesert O, Russell P (1998) Replication checkpoint enforced by kinases Cds1 and Chk1. *Science* 280: 909–912

- Boos D, Sanchez-Pulido L, Rappas M, Pearl LH, Oliver AW, Ponting CP, Diffley JF (2011) Regulation of DNA replication through Sld3-Dpb11 interaction is conserved from yeast to humans. *Curr Biol* 21: 1152–1157
- Botchkarev Jr VW, Haber JE (2018) Functions and regulation of the Polo-like kinase Cdc5 in the absence and presence of DNA damage. *Curr Genet* 64: 87–96
- Bothmer A, Robbiani DF, Di Virgilio M, Bunting SF, Klein IA, Feldhahn N, Barlow J, Chen HT, Bosque D, Callen E et al (2011) Regulation of DNA end joining, resection, and immunoglobulin class switch recombination by 53BP1. *Mol Cell* 42: 319–329
- Bracken AP, Ciro M, Cocito A, Helin K (2004) E2F target genes: unraveling the biology. *Trends Biochem Sci* 29: 409–417
- Brown EJ, Baltimore D (2000) ATR disruption leads to chromosomal fragmentation and early embryonic lethality. *Genes Dev* 14: 397–402
- de Bruin RA, Kalashnikova TI, Aslanian A, Wohlschlegel J, Chahwan C, Yates III JR, Russell P, Wittenberg C (2008) DNA replication checkpoint promotes G1-S transcription by inactivating the MBF repressor Nrm1. *Proc Natl Acad Sci USA* 105: 11230–11235
- Bruinsma W, Aprelia M, Garcia-Santisteban I, Kool J, Xu YJ, Medema RH (2017) Inhibition of Polo-like kinase 1 during the DNA damage response is mediated through loss of Aurora A recruitment by Bora. *Oncogene* 36: 1840–1848
- Buisson R, Niraj J, Rodrigue A, Ho CK, Kreuzer J, Foo TK, Hardy EJ, Dellaire G, Haas W, Xia B et al (2017) Coupling of homologous recombination and the checkpoint by ATR. *Mol Cell* 65: 336–346
- Bunting SF, Callen E, Wong N, Chen HT, Polato F, Gunn A, Bothmer A, Feldhahn N, Fernandez-Capetillo O, Cao L et al (2010) 53BP1 inhibits homologous recombination in Brca1-deficient cells by blocking resection of DNA breaks. *Cell* 141: 243–254
- Cai Z, Chehab NH, Pavletich NP (2009) Structure and activation mechanism of the CHK2 DNA damage checkpoint kinase. *Mol Cell* 35: 818–829
- Callen E, Jankovic M, Wong N, Zha S, Chen HT, Difilippantonio S, Di Virgilio M, Heidkamp G, Alt FW, Nussenzweig A et al (2009) Essential role for DNA-PKcs in DNA double-strand break repair and apoptosis in ATM-deficient lymphocytes. *Mol Cell* 34: 285–297
- Can G, Kauerhof AC, Macak D, Zegerman P (2018) Helicase subunit Cdc45 targets the checkpoint kinase Rad53 to both replication initiation and elongation complexes after fork stalling. *Mol Cell* 73: 562–573.e3
- Carr AM (1995) DNA structure checkpoints in fission yeast. *Semin Cell Biol* 6: 65–72
- Chan DW, Ye R, Veillette CJ, Lees-Miller SP (1999) DNA-dependent protein kinase phosphorylation sites in Ku 70/80 heterodimer. *Biochemistry* 38: 1819–1828
- Chapman JR, Barral P, Vannier JB, Borel V, Steger M, Tomas-Loba A, Sartori AA, Adams IR, Batista FD, Boulton SJ (2013) RIF1 is essential for 53BP1-dependent nonhomologous end joining and suppression of DNA double-strand break resection. *Mol Cell* 49: 858–871
- Chen YH, Jones MJ, Yin Y, Crist SB, Colnaghi L, Sims III RJ, Rothenberg E, Jallepalli PV, Huang TT (2015) ATR-mediated phosphorylation of FANCI regulates dormant origin firing in response to replication stress. *Mol Cell* 58: 323–338
- Chen J (2016) The cell-cycle arrest and apoptotic functions of p53 in tumor initiation and progression. *Cold Spring Harb Perspect Med* 6: a026104
- Chi Y, Welcker M, Hizli AA, Posakony JJ, Aebbersold R, Clurman BE (2008) Identification of CDK2 substrates in human cell lysates. *Genome Biol* 9: R149
- Chou WC, Wang HC, Wong FH, Ding SL, Wu PE, Shieh SY, Shen CY (2008) Chk2-dependent phosphorylation of XRCC1 in the DNA damage response promotes base excision repair. *EMBO J* 27: 3140–3150
- Cimprich KA, Shin TB, Keith CT, Schreiber SL (1996) cDNA cloning and gene mapping of a candidate human cell cycle checkpoint protein. *Proc Natl Acad Sci USA* 93: 2850–2855
- Clerici M, Trovesi C, Galbiati A, Lucchini G, Longhese MP (2014) Mec1/ATR regulates the generation of single-stranded DNA that attenuates Tel1/ATM signaling at DNA ends. *EMBO J* 33: 198–216
- Collins NB, Wilson JB, Bush T, Thomashevski A, Roberts KJ, Jones NJ, Kupfer GM (2009) ATR-dependent phosphorylation of FANCA on serine 1449 after DNA damage is important for FA pathway function. *Blood* 113: 2181–2190
- Cortez D, Glick C, Elledge SJ (2004) Minichromosome maintenance proteins are direct targets of the ATM and ATR checkpoint kinases. *Proc Natl Acad Sci USA* 101: 10078–10083
- Costanzo V, Shechter D, Lupardus PJ, Cimprich KA, Gottesman M, Gautier J (2003) An ATR- and Cdc7-dependent DNA damage checkpoint that inhibits initiation of DNA replication. *Mol Cell* 11: 203–213
- Cotta-Ramusino C, Fachinetti D, Lucca C, Doksan Y, Lopes M, Sogo J, Foiani M (2005) Exo1 processes stalled replication forks and counteracts fork reversal in checkpoint-defective cells. *Mol Cell* 17: 153–159
- Couch FB, Bansbach CE, Driscoll R, Luzwick JW, Glick GG, Betous R, Carroll CM, Jung SY, Qin J, Cimprich KA et al (2013) ATR phosphorylates SMARCA1 to prevent replication fork collapse. *Genes Dev* 27: 1610–1623
- Cuadrado M, Martinez-Pastor B, Murga M, Toledo LI, Gutierrez-Martinez P, Lopez E, Fernandez-Capetillo O (2006) ATM regulates ATR chromatin loading in response to DNA double-strand breaks. *J Exp Med* 203: 297–303
- Cussiol JR, Jablonowski CM, Yimit A, Brown GW, Smolka MB (2015) Dampening DNA damage checkpoint signalling via coordinated BRCT domain interactions. *EMBO J* 34: 1704–1717
- Cussiol JR, Dibitetto D, Pelliccioli A, Smolka MB (2017) Slx4 scaffolding in homologous recombination and checkpoint control: lessons from yeast. *Chromosoma* 126: 45–58
- D'Angiolella V, Donato V, Forrester FM, Jeong YT, Pellacani C, Kudo Y, Saraf A, Florens L, Washburn MP, Pagano M (2012) Cyclin F-mediated degradation of ribonucleotide reductase M2 controls genome integrity and DNA repair. *Cell* 149: 1023–1034
- Das BB, Antony S, Gupta S, Dexheimer TS, Redon CE, Garfield S, Shiloh Y, Pommier Y (2009) Optimal function of the DNA repair enzyme TDP1 requires its phosphorylation by ATM and/or DNA-PK. *EMBO J* 28: 3667–3680
- Davies SL, North PS, Dart A, Lakin ND, Hickson ID (2004) Phosphorylation of the Bloom's syndrome helicase and its role in recovery from S-phase arrest. *Mol Cell Biol* 24: 1279–1291
- Davies SL, North PS, Hickson ID (2007) Role for BLM in replication-fork restart and suppression of origin firing after replicative stress. *Nat Struct Mol Biol* 14: 677–679
- Deng X, Prakash A, Dhar K, Baia GS, Kolar C, Oakley GG, Borgstahl GE (2009) Human replication protein A-Rad52-single-stranded DNA complex: stoichiometry and evidence for strand transfer regulation by phosphorylation. *Biochemistry* 48: 6633–6643
- Di Domenico EG, Romano E, Del Porto P, Ascenzioni F (2014) Multifunctional role of ATM/Tel1 kinase in genome stability: from the DNA damage response to telomere maintenance. *Biomed Res Int* 2014: 787404
- Dibitetto D, Ferrari M, Rawal CC, Balint A, Kim T, Zhang Z, Smolka MB, Brown GW, Marini F, Pelliccioli A (2016) Slx4 and Rtt107 control checkpoint signalling and DNA resection at double-strand breaks. *Nucleic Acids Res* 44: 669–682
- Donnianni RA, Ferrari M, Lazzaro F, Clerici M, Tamilselvan Nachimuthu B, Plevani P, Muzi-Falconi M, Pelliccioli A (2010) Elevated levels of the polo

- kinase Cdc5 override the Mec1/ATR checkpoint in budding yeast by acting at different steps of the signaling pathway. *PLoS Genet* 6: e1000763
- Douglas P, Gupta S, Morrice N, Meek K, Lees-Miller SP (2005) DNA-PK-dependent phosphorylation of Ku70/80 is not required for non-homologous end joining. *DNA Repair (Amst)* 4: 1006–1018
- Downs JA, Lowndes NF, Jackson SP (2000) A role for *Saccharomyces cerevisiae* histone H2A in DNA repair. *Nature* 408: 1001–1004
- Downs JA, Allard S, Jobin-Robitaille O, Javaheri A, Auger A, Bouchard N, Kron SJ, Jackson SP, Cote J (2004) Binding of chromatin-modifying activities to phosphorylated histone H2A at DNA damage sites. *Mol Cell* 16: 979–990
- Dungrawala H, Rose KL, Bhat KP, Mohni KN, Glick GG, Couch FB, Cortez D (2015) The replication checkpoint prevents two types of fork collapse without regulating replisome stability. *Mol Cell* 59: 998–1010
- Edenberg ER, Mark KG, Toczyski DP (2015) Ndd1 turnover by SCF(Grr1) is inhibited by the DNA damage checkpoint in *Saccharomyces cerevisiae*. *PLoS Genet* 11: e1005162
- El-Shemerly M, Hess D, Pyakurel AK, Moselhy S, Ferrari S (2008) ATR-dependent pathways control hEXO1 stability in response to stalled forks. *Nucleic Acids Res* 36: 511–519
- Emili A (1998) MEC1-dependent phosphorylation of Rad9p in response to DNA damage. *Mol Cell* 2: 183–189
- Engels K, Giannattasio M, Muzi-Falconi M, Lopes M, Ferrari S (2011) 14-3-3 proteins regulate exonuclease 1-dependent processing of stalled replication forks. *PLoS Genet* 7: e1001367
- Engholm-Keller K, Larsen MR (2013) Technologies and challenges in large-scale phosphoproteomics. *Proteomics* 13: 910–931
- Ferrari M, Nachimuthu BT, Donnianni RA, Klein H, Pelliccioli A (2013) Tid1/Rdh54 translocase is phosphorylated through a Mec1- and Rad53-dependent manner in the presence of DSB lesions in budding yeast. *DNA Repair (Amst)* 12: 347–355
- Ferrari M, Dibitetto D, De Gregorio G, Eapen VV, Rawal CC, Lazzaro F, Tsabar M, Marini F, Haber JE, Pelliccioli A (2015) Functional interplay between the 53BP1-ortholog Rad9 and the Mre11 complex regulates resection, end-tethering and repair of a double-strand break. *PLoS Genet* 11: e1004928
- Fiorani S, Mimun G, Caleca L, Piccini D, Pelliccioli A (2008) Characterization of the activation domain of the Rad53 checkpoint kinase. *Cell Cycle* 7: 493–499
- Flott S, Kwon Y, Pigli YZ, Rice PA, Sung P, Jackson SP (2011) Regulation of Rad51 function by phosphorylation. *EMBO Rep* 12: 833–839
- Foskolou IP, Jorgensen C, Leszczynska KB, Olcina MM, Tarhonskaya H, Haisma B, D'Angiolella V, Myers WK, Domene C, Flashman E et al (2017) Ribonucleotide reductase requires subunit switching in hypoxia to maintain DNA replication. *Mol Cell* 66: 206–220 e9
- Furnari B, Rhind N, Russell P (1997) Cdc25 mitotic inducer targeted by chk1 DNA damage checkpoint kinase. *Science* 277: 1495–1497
- Gapud EJ, Dorsett Y, Yin B, Callen E, Bredemeyer A, Mahowald GK, Omi KQ, Walker LM, Bednarski JJ, McKinnon PJ et al (2011) Ataxia telangiectasia mutated (Atm) and DNA-PKcs kinases have overlapping activities during chromosomal signal joint formation. *Proc Natl Acad Sci USA* 108: 2022–2027
- Gapud EJ, Sleckman BP (2011) Unique and redundant functions of ATM and DNA-PKcs during V(D)J recombination. *Cell Cycle* 10: 1928–1935
- Gilbert CS, Green CM, Lowndes NF (2001) Budding yeast Rad9 is an ATP-dependent Rad53 activating machine. *Mol Cell* 8: 129–136
- Gobbini E, Villa M, Gnugnoli M, Menin L, Clerici M, Longhese MP (2015) Sae2 function at DNA double-strand breaks is bypassed by dampening Tel1 or Rad53 activity. *PLoS Genet* 11: e1005685
- Gohler T, Sabbioneda S, Green CM, Lehmann AR (2011) ATR-mediated phosphorylation of DNA polymerase η is needed for efficient recovery from UV damage. *J Cell Biol* 192: 219–227
- Goldberg M, Stucki M, Falck J, D'Amours D, Rahman D, Pappin D, Bartek J, Jackson SP (2003) MDC1 is required for the intra-S-phase DNA damage checkpoint. *Nature* 421: 952–956
- Goodarzi AA, Yu Y, Riballo E, Douglas P, Walker SA, Ye R, Harer C, Marchetti C, Morrice N, Jeggo PA et al (2006) DNA-PK autophosphorylation facilitates Artemis endonuclease activity. *EMBO J* 25: 3880–3889
- Guervilly JH, Gaillard PH (2018) SLX4: multitasking to maintain genome stability. *Crit Rev Biochem Mol Biol* 53: 475–514
- Gunjan A, Verreault A (2003) A Rad53 kinase-dependent surveillance mechanism that regulates histone protein levels in *S. cerevisiae*. *Cell* 115: 537–549
- Guo C, Kumagai A, Schlacher K, Shevchenko A, Shevchenko A, Dunphy WG (2015) Interaction of Chk1 with Treslin negatively regulates the initiation of chromosomal DNA replication. *Mol Cell* 57: 492–505
- Hammet A, Magill C, Heierhorst J, Jackson SP (2007) Rad9 BRCT domain interaction with phosphorylated H2AX regulates the G1 checkpoint in budding yeast. *EMBO Rep* 8: 851–857
- Hang L, Zhao X (2016) The Rtt107 BRCT scaffold and its partner modification enzymes collaborate to promote replication. *Nucleus* 7: 346–351
- Harper JW, Adami GR, Wei N, Keyomarsi K, Elledge SJ (1993) The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75: 805–816
- Harper JW, Elledge SJ, Keyomarsi K, Dynlacht B, Tsai LH, Zhang P, Dobrowolski S, Bai C, Connell-Crowley L, Swindell E et al (1995) Inhibition of cyclin-dependent kinases by p21. *Mol Biol Cell* 6: 387–400
- Hartwell LH, Weinert TA (1989) Checkpoints: controls that ensure the order of cell cycle events. *Science* 246: 629–634
- Heffernan TP, Unsal-Kacmaz K, Heinloth AN, Simpson DA, Paules RS, Sancar A, Cordeiro-Stone M, Kaufmann WK (2007) Cdc7-Dbp4 and the human S checkpoint response to UVC. *J Biol Chem* 282: 9458–9468
- Hegnauer AM, Hustedt N, Shimada K, Pike BL, Vogel M, Amsler P, Rubin SM, van Leeuwen F, Guenole A, van Attikum H et al (2012) An N-terminal acidic region of Sgs1 interacts with Rpa70 and recruits Rad53 kinase to stalled forks. *EMBO J* 31: 3768–3783
- Hertz NT, Wang BT, Allen JJ, Zhang C, Dar AC, Burlingame AL, Shokat KM (2010) Chemical genetic approach for kinase-substrate mapping by covalent capture of thiophosphopeptides and analysis by mass spectrometry. *Curr Protoc Chem Biol* 2: 15–36
- Herzberg K, Bashkurov VI, Rolfmeier M, Haghazari E, McDonald WH, Anderson S, Bashkurova EV, Yates III JR, Heyer WD (2006) Phosphorylation of Rad55 on serines 2, 8, and 14 is required for efficient homologous recombination in the recovery of stalled replication forks. *Mol Cell Biol* 26: 8396–8409
- Hirano Y, Sugimoto K (2006) ATR homolog Mec1 controls association of DNA polymerase ζ -Rev1 complex with regions near a double-strand break. *Curr Biol* 16: 586–590
- Ho GP, Margossian S, Taniguchi T, D'Andrea AD (2006) Phosphorylation of FANCD2 on two novel sites is required for mitomycin C resistance. *Mol Cell Biol* 26: 7005–7015
- Huang M, Zhou Z, Elledge SJ (1998) The DNA replication and damage checkpoint pathways induce transcription by inhibition of the Crt1 repressor. *Cell* 94: 595–605
- Hustedt N, Durocher D (2016) The control of DNA repair by the cell cycle. *Nat Cell Biol* 19: 1–9

- Ishiai M, Kitao H, Smogorzewska A, Tomida J, Kinomura A, Uchida E, Saberi A, Kinoshita E, Kinoshita-Kikuta E, Koike T *et al* (2008) FANCI phosphorylation functions as a molecular switch to turn on the Fanconi anemia pathway. *Nat Struct Mol Biol* 15: 1138–1146
- Ishida S, Huang E, Zuzan H, Spang R, Leone G, West M, Nevins JR (2001) Role for E2F in control of both DNA replication and mitotic functions as revealed from DNA microarray analysis. *Mol Cell Biol* 21: 4684–4699
- Jachimowicz RD, Beleggia F, Isensee J, Velpula BB, Goergens J, Bustos MA, Doll MA, Shenoy A, Checa-Rodriguez C, Wiederstein JL *et al* (2019) UBQLN4 represses homologous recombination and is overexpressed in aggressive tumors. *Cell* 176: 505–519 e22
- Jaehnjg EJ, Kuo D, Hombauer H, Ideker TG, Kolodner RD (2013) Checkpoint kinases regulate a global network of transcription factors in response to DNA damage. *Cell Rep* 4: 174–188
- jasin M, Rothstein R (2013) Repair of strand breaks by homologous recombination. *Cold Spring Harb Perspect Biol* 5: a012740
- Jia-Lin Ma N, Stern DF (2008) Regulation of the Rad53 protein kinase in signal amplification by oligomer assembly and disassembly. *Cell Cycle* 7: 808–817
- Kaur S, Modi P, Srivastava V, Mudgal R, Tikoo S, Arora P, Mohanty D, Sengupta S (2010) Chk1-dependent constitutive phosphorylation of BLM helicase at serine 646 decreases after DNA damage. *Mol Cancer Res* 8: 1234–1247
- Kim D, Liu Y, Oberly S, Freire R, Smolka MB (2018) ATR-mediated proteome remodeling is a major determinant of homologous recombination capacity in cancer cells. *Nucleic Acids Res* 46: 8311–8325
- Kirkpatrick DS, Bustos DJ, Dogan T, Chan J, Phu L, Young A, Friedman LS, Belvin M, Song Q, Bakalarski CE *et al* (2013) Phosphoproteomic characterization of DNA damage response in melanoma cells following MEK/PI3K dual inhibition. *Proc Natl Acad Sci USA* 110: 19426–19431
- Kleiner RE, Verma P, Molloy KR, Chait BT, Kapoor TM (2015) Chemical proteomics reveals a gammaH2AX-53BP1 interaction in the DNA damage response. *Nat Chem Biol* 11: 807–814
- Knipscheer P, Raschle M, Smogorzewska A, Enoiu M, Ho TV, Schärer OD, Elledge SJ, Walter JC (2009) The Fanconi anemia pathway promotes replication-dependent DNA interstrand cross-link repair. *Science* 326: 1698–1701
- Kolinjivadi AM, Sannino V, De Antoni A, Zadorozhny K, Kilkenny M, Techer H, Baldi G, Shen R, Ciccia A, Pellegrini L *et al* (2017) Smarcal1-mediated fork reversal triggers Mre11-dependent degradation of nascent DNA in the absence of Brca2 and Stable Rad51 nucleofilaments. *Mol Cell* 67: 867–881 e7
- Kruse JP, Gu W (2009) Modes of p53 regulation. *Cell* 137: 609–622
- Kumagai A, Dunphy WG (2000) Claspin, a novel protein required for the activation of Chk1 during a DNA replication checkpoint response in *Xenopus* egg extracts. *Mol Cell* 6: 839–849
- Kumagai A, Dunphy WG (2003) Repeated phosphopeptide motifs in Claspin mediate the regulated binding of Chk1. *Nat Cell Biol* 5: 161–165
- Kumagai A, Shevchenko A, Shevchenko A, Dunphy WG (2010) Treslin collaborates with TopBP1 in triggering the initiation of DNA replication. *Cell* 140: 349–359
- Lanz MC, Oberly S, Sanford EJ, Sharma S, Chabes A, Smolka MB (2018) Separable roles for Mec1/ATR in genome maintenance, DNA replication, and checkpoint signaling. *Genes Dev* 32: 822–835
- Lanz MC, Yugandhar K, Gupta S, Sanford E, Faça V, Vega S, Joiner A, Fromme C, Yu H (2019) In-depth and 3-dimensional exploration of the budding yeast phosphoproteome. *bioRxiv* <https://doi.org/10.1101/700070> [PREPRINT]
- Lazzaro F, Sapountzi V, Granata M, Pelliccioli A, Vaze M, Haber JE, Plevani P, Lydall D, Muzi-Falconi M (2008) Histone methyltransferase Dot1 and Rad9 inhibit single-stranded DNA accumulation at DSBs and uncapped telomeres. *EMBO J* 27: 1502–1512
- Lee J, Kumagai A, Dunphy WG (2001) Positive regulation of Wee1 by Chk1 and 14-3-3 proteins. *Mol Biol Cell* 12: 551–563
- Lee J, Kumagai A, Dunphy WG (2003) Claspin, a Chk1-regulatory protein, monitors DNA replication on chromatin independently of RPA, ATR, and Rad17. *Mol Cell* 11: 329–340
- Lee KJ, Jovanovic M, Udayakumar D, Bladen CL, Dynan WS (2004) Identification of DNA-PKcs phosphorylation sites in XRCC4 and effects of mutations at these sites on DNA end joining in a cell-free system. *DNA Repair (Amst)* 3: 267–276
- Lee YD, Wang J, Stubbe J, Elledge SJ (2008) Dif1 is a DNA-damage-regulated facilitator of nuclear import for ribonucleotide reductase. *Mol Cell* 32: 70–80
- Lee AY, Chiba T, Truong LN, Cheng AN, Do J, Cho MJ, Chen L, Wu X (2012) Dbf4 is direct downstream target of ataxia telangiectasia mutated (ATM) and ataxia telangiectasia and Rad3-related (ATR) protein to regulate intra-S-phase checkpoint. *J Biol Chem* 287: 2531–2543
- Leung CC, Sun L, Gong Z, Burkat M, Edwards R, Assmus M, Chen J, Glover JN (2013) Structural insights into recognition of MDC1 by TopBP1 in DNA replication checkpoint control. *Structure* 21: 1450–1459
- Li S, Lavagnino Z, Lemacon D, Kong L, Ustione A, Ng X, Zhang Y, Wang Y, Zheng B, Piwnicka-Worms H *et al* (2019) Ca(2+)-stimulated AMPK-dependent phosphorylation of Exo1 protects stressed replication forks from aberrant resection. *Mol Cell* 74: 1123–1137 e6
- Liang J, Suhandynata RT, Zhou H (2015) Phosphorylation of Sae2 mediates forkhead-associated (FHA) domain-specific interaction and regulates its DNA repair function. *J Biol Chem* 290: 10751–10763
- Lin WC, Lin FT, Nevins JR (2001) Selective induction of E2F1 in response to DNA damage, mediated by ATM-dependent phosphorylation. *Genes Dev* 15: 1833–1844
- Lindsey-Boltz LA, Sercin O, Choi JH, Sancar A (2009) Reconstitution of human claspin-mediated phosphorylation of Chk1 by the ATR (ataxia telangiectasia-mutated and rad3-related) checkpoint kinase. *J Biol Chem* 284: 33107–33114
- Liu Q, Guntuku S, Cui XS, Matsuoka S, Cortez D, Tamai K, Luo G, Carattini-Rivera S, DeMayo F, Bradley A *et al* (2000) Chk1 is an essential kinase that is regulated by Atr and required for the G(2)/M DNA damage checkpoint. *Genes Dev* 14: 1448–1459
- Liu S, Bekker-Jensen S, Mailand N, Lukas C, Bartek J, Lukas J (2006) Claspin operates downstream of TopBP1 to direct ATR signaling towards Chk1 activation. *Mol Cell Biol* 26: 6056–6064
- Liu Y, Cussiol JR, Dibitetto D, Sims JR, Twayana S, Weiss RS, Freire R, Marini F, Pelliccioli A, Smolka MB (2017) TOPBP1Dpb11 plays a conserved role in homologous recombination DNA repair through the coordinated recruitment of 53BP1Rad9. *J Cell Biol* 216: 623–639
- Lopez-Contreras AJ, Specks J, Barlow JH, Ambrogio C, Desler C, Vikingsson S, Rodrigo-Perez S, Green H, Rasmussen LJ, Murga M *et al* (2015) Increased Rrm2 gene dosage reduces fragile site breakage and prolongs survival of ATR mutant mice. *Genes Dev* 29: 690–695
- Lopez-Mosqueda J, Maas NL, Jonsson ZO, Defazio-Eli LG, Wohlschlegel J, Toczyski DP (2010) Damage-induced phosphorylation of Sld3 is important to block late origin firing. *Nature* 467: 479–483
- Malewicz M, Kadkhodaei B, Kee N, Volakakis N, Hellman U, Viktorsson K, Leung CY, Chen B, Lewensohn R, van Gent DC *et al* (2011) Essential role for DNA-PK-mediated phosphorylation of NR4A nuclear

- orphan receptors in DNA double-strand break repair. *Genes Dev* 25: 2031–2040
- Matsunuma R, Niida H, Ohhata T, Kitagawa K, Sakai S, Uchida C, Shiotani B, Matsumoto M, Nakayama KI, Ogura H *et al* (2016) UV damage-induced phosphorylation of HBO1 triggers CRL4DDB2-mediated degradation to regulate cell proliferation. *Mol Cell Biol* 36: 394–406
- Matsuoka S, Huang M, Elledge SJ (1998) Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 282: 1893–1897
- Matsuoka S, Ballif BA, Smogorzewska A, McDonald III ER, Hurov KE, Luo J, Bakalarski CE, Zhao Z, Solimini N, Lerenthal Y *et al* (2007) ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science* 316: 1160–1166
- Maya R, Balass M, Kim ST, Shkedy D, Leal JF, Shifman O, Moas M, Buschmann T, Ronai Z, Shiloh Y *et al* (2001) ATM-dependent phosphorylation of Mdm2 on serine 395: role in p53 activation by DNA damage. *Genes Dev* 15: 1067–1077
- Mayo MW, Wang CY, Cogswell PC, Rogers-Graham KS, Lowe SW, Der CJ, Baldwin Jr AS (1997) Requirement of NF-kappaB activation to suppress p53-independent apoptosis induced by oncogenic Ras. *Science* 278: 1812–1815
- McIntosh D, Blow JJ (2012) Dormant origins, the licensing checkpoint, and the response to replicative stresses. *Cold Spring Harb Perspect Biol* 4: a012955
- Morin I, Ngo HP, Greenall A, Zubko MK, Morrice N, Lydall D (2008) Checkpoint-dependent phosphorylation of Exo1 modulates the DNA damage response. *EMBO J* 27: 2400–2410
- Morrison AJ, Highland J, Krogan NJ, Arbel-Eden A, Greenblatt JF, Haber JE, Shen X (2004) INO80 and gamma-H2AX interaction links ATP-dependent chromatin remodeling to DNA damage repair. *Cell* 119: 767–775
- Morrison AJ, Kim JA, Person MD, Highland J, Xiao J, Wehr TS, Hensley S, Bao Y, Shen J, Collins SR *et al* (2007) Mec1/Tel1 phosphorylation of the INO80 chromatin remodeling complex influences DNA damage checkpoint responses. *Cell* 130: 499–511
- Morrow DM, Tagle DA, Shiloh Y, Collins FS, Hieter P (1995) TEL1, an *S. cerevisiae* homolog of the human gene mutated in ataxia telangiectasia, is functionally related to the yeast checkpoint gene MEC1. *Cell* 82: 831–840
- Mueller PR, Coleman TR, Dunphy WG (1995) Cell cycle regulation of a *Xenopus* Wee1-like kinase. *Mol Biol Cell* 6: 119–134
- Murakami H, Aiba H, Nakanishi M, Murakami-Tonami Y (2010) Regulation of yeast forkhead transcription factors and FoxM1 by cyclin-dependent and polo-like kinases. *Cell Cycle* 9: 3233–3242
- Murphy AK, Fitzgerald M, Ro T, Kim JH, Rabinowitsch AI, Chowdhury D, Schildkraut CL, Borowiec JA (2014) Phosphorylated RPA recruits PALB2 to stalled DNA replication forks to facilitate fork recovery. *J Cell Biol* 206: 493–507
- Myung K, Datta A, Kolodner RD (2001) Suppression of spontaneous chromosomal rearrangements by S phase checkpoint functions in *Saccharomyces cerevisiae*. *Cell* 104: 397–408
- Naiki T, Wakayama T, Nakada D, Matsumoto K, Sugimoto K (2004) Association of Rad9 with double-strand breaks through a Mec1-dependent mechanism. *Mol Cell Biol* 24: 3277–3285
- Nielsen I, Bentsen IB, Andersen AH, Gasser SM, Bjergbaek L (2013) A Rad53 independent function of Rad9 becomes crucial for genome maintenance in the absence of the Recq helicase Sgs1. *PLoS One* 8: e81015
- Niraj J, Farkkila A, D'Andrea AD (2019) The fanconi anemia pathway in cancer. *Annu Rev Cancer Biol* 3: 457–478
- O'Connell MJ, Raleigh JM, Verkade HM, Nurse P (1997) Chk1 is a wee1 kinase in the G2 DNA damage checkpoint inhibiting cdc2 by Y15 phosphorylation. *EMBO J* 16: 545–554
- Ohouo PY, Bastos de Oliveira FM, Almeida BS, Smolka MB (2010) DNA damage signaling recruits the Rtt107-Slx4 scaffolds via Dpb11 to mediate replication stress response. *Mol Cell* 39: 300–306
- Osborn AJ, Elledge SJ (2003) Mrc1 is a replication fork component whose phosphorylation in response to DNA replication stress activates Rad53. *Genes Dev* 17: 1755–1767
- Paciotti V, Lucchini G, Plevani P, Longhese MP (1998) Mec1p is essential for phosphorylation of the yeast DNA damage checkpoint protein Ddc1p, which physically interacts with Mec3p. *EMBO J* 17: 4199–4209
- Pages V, Santa Maria SR, Prakash L, Prakash S (2009) Role of DNA damage-induced replication checkpoint in promoting lesion bypass by translesion synthesis in yeast. *Genes Dev* 23: 1438–1449
- Palou G, Palou R, Zeng F, Vashisht AA, Wohlschlegel JA, Quintana DG (2015) Three different pathways prevent chromosome segregation in the presence of DNA damage or replication stress in budding yeast. *PLoS Genet* 11: e1005468
- Pardo B, Crabbe L, Pasero P (2017) Signaling pathways of replication stress in yeast. *FEMS Yeast Res* 17: fow101
- Park MS, Ludwig DL, Stigger E, Lee SH (1996) Physical interaction between human RAD52 and RPA is required for homologous recombination in mammalian cells. *J Biol Chem* 271: 18996–19000
- Pasero P, Vindigni A (2017) Nucleases acting at stalled forks: how to reboot the replication program with a few shortcuts. *Annu Rev Genet* 51: 477–499
- Patel DR, Weiss RS (2018) A tough row to hoe: when replication forks encounter DNA damage. *Biochem Soc Trans* 46: 1643–1651
- Peng CY, Graves PR, Thoma RS, Wu Z, Shaw AS, Piwnicka-Worms H (1997) Mitotic and G2 checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on serine-216. *Science* 277: 1501–1505
- Peng A, Chen PL (2003) NFB1, like 53BP1, is an early and redundant transducer mediating Chk2 phosphorylation in response to DNA damage. *J Biol Chem* 278: 8873–8876
- Petsalaki E, Dandoulaki M, Morrice N, Zachos G (2014) Chk1 protects against chromatin bridges by constitutively phosphorylating BLM serine 502 to inhibit BLM degradation. *J Cell Sci* 127: 3902–3908
- Pfander B, Diffley JF (2011) Dpb11 coordinates Mec1 kinase activation with cell cycle-regulated Rad9 recruitment. *EMBO J* 30: 4897–4907
- Pic-Taylor A, Darieva Z, Morgan BA, Sharrocks AD (2004) Regulation of cell cycle-specific gene expression through cyclin-dependent kinase-mediated phosphorylation of the forkhead transcription factor Fkh2p. *Mol Cell Biol* 24: 10036–10046
- Pines A, Kelstrup CD, Vrouwe MG, Puigvert JC, Typas D, Misovic B, de Groot A, von Stechow L, van de Water B, Danen EH *et al* (2011) Global phosphoproteome profiling reveals unanticipated networks responsive to cisplatin treatment of embryonic stem cells. *Mol Cell Biol* 31: 4964–4977.
- Poppy Roworth A, Ghari F, La Thangue NB (2015) To live or let die – complexity within the E2F1 pathway. *Mol Cell Oncol* 2: e970480
- Qin B, Gao B, Yu J, Yuan J, Lou Z (2013) Ataxia telangiectasia-mutated- and Rad3-related protein regulates the DNA damage-induced G2/M checkpoint through the Aurora A cofactor Bora protein. *J Biol Chem* 288: 16139–16144
- Randell JC, Fan A, Chan C, Francis LI, Heller RC, Galani K, Bell SP (2010) Mec1 is one of multiple kinases that prime the Mcm2-7 helicase for phosphorylation by Cdc7. *Mol Cell* 40: 353–363
- Romeo F, Falbo L, Di Sanzo M, Misaggi R, Faniello MC, Viglietto G, Cuda G, Costanzo F, Quaresima B (2011) BRCA1 is required for hMLH1 stabilization following doxorubicin-induced DNA damage. *Int J Biochem Cell Biol* 43: 1754–1763

- Rossi SE, Ajazi A, Carotenuto W, Foiani M, Giannattasio M (2015) Rad53-mediated regulation of Rrm3 and Pif1 DNA helicases contributes to prevention of aberrant fork transitions under replication stress. *Cell Rep* 13: 80–92
- Ruiz JF, Pardo B, Sastre-Moreno G, Aguilera A, Blanco L (2013) Yeast pol4 promotes tel1-regulated chromosomal translocations. *PLoS Genet* 9: e1003656
- Sabbioneda S, Bortolomai I, Giannattasio M, Plevani P, Muzi-Falconi M (2007) Yeast Rev1 is cell cycle regulated, phosphorylated in response to DNA damage and its binding to chromosomes is dependent upon MEC1. *DNA Repair (Amst)* 6: 121–127
- Saldivar JC, Cortez D, Cimprich KA (2017) The essential kinase ATR: ensuring faithful duplication of a challenging genome. *Nat Rev Mol Cell Biol* 18: 622–636
- Saldivar JC, Hamperl S, Bocek MJ, Chung M, Bass TE, Cisneros-Soberanis F, Samejima K, Xie L, Paulson JR, Earnshaw WC et al (2018) An intrinsic S/G2 checkpoint enforced by ATR. *Science* 361: 806–810
- Sanchez Y, Desany BA, Jones WJ, Liu Q, Wang B, Elledge SJ (1996) Regulation of RAD53 by the ATM-like kinases MEC1 and TEL1 in yeast cell cycle checkpoint pathways. *Science* 271: 357–360
- Sanchez Y, Wong C, Thoma RS, Richman R, Wu Z, Piwnicka-Worms H, Elledge SJ (1997) Conservation of the Chk1 checkpoint pathway in mammals: linkage of DNA damage to Cdk regulation through Cdc25. *Science* 277: 1497–1501
- Sanchez Y, Bachant J, Wang H, Hu F, Liu D, Tetzlaff M, Elledge SJ (1999) Control of the DNA damage checkpoint by chk1 and rad53 protein kinases through distinct mechanisms. *Science* 286: 1166–1171
- Santocanale C, Diffley JF (1998) A Mec1- and Rad53-dependent checkpoint controls late-firing origins of DNA replication. *Nature* 395: 615–618
- Sarbajna S, West SC (2014) Holliday junction processing enzymes as guardians of genome stability. *Trends Biochem Sci* 39: 409–419
- Sastre-Moreno G, Pryor JM, Moreno-Onate M, Herrero-Ruiz AM, Cortes-Ledesma F, Blanco L, Ramsden DA, Ruiz JF (2017) Regulation of human pollambda by ATM-mediated phosphorylation during non-homologous end joining. *DNA Repair (Amst)* 51: 31–45
- Savitsky K, Bar-Shira A, Gilad S, Rotman G, Ziv Y, Vanagaite L, Tagle DA, Smith S, Uziel T, Sfez S et al (1995) A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 268: 1749–1753
- Schwartz MF, Duong JK, Sun Z, Morrow JS, Pradhan D, Stern DF (2002) Rad9 phosphorylation sites couple Rad53 to the *Saccharomyces cerevisiae* DNA damage checkpoint. *Mol Cell* 9: 1055–1065
- Schwarz JK, Lovly CM, Piwnicka-Worms H (2003) Regulation of the Chk2 protein kinase by oligomerization-mediated cis- and trans-phosphorylation. *Mol Cancer Res* 1: 598–609
- Segurado M, Diffley JF (2008) Separate roles for the DNA damage checkpoint protein kinases in stabilizing DNA replication forks. *Genes Dev* 22: 1816–1827
- Segurado M, Tercero JA (2009) The S-phase checkpoint: targeting the replication fork. *Biol Cell* 101: 617–627
- Sengupta S, Robles AI, Linke SP, Sinogeeva NI, Zhang R, Pedoux R, Ward IM, Celeste A, Nussenzweig A, Chen J et al (2004) Functional interaction between BLM helicase and 53BP1 in a Chk1-mediated pathway during S-phase arrest. *J Cell Biol* 166: 801–813
- Setiawati D, Durocher D (2019) Shieldin – the protector of DNA ends. *EMBO Rep* 20: e47560
- Shechter D, Costanzo V, Gautier J (2004) ATR and ATM regulate the timing of DNA replication origin firing. *Nat Cell Biol* 6: 648–655
- Shibata A, Conrad S, Birraux J, Geuting V, Barton O, Ismail A, Kakarougkas A, Meek K, Taucher-Scholz G, Lobrich M et al (2011) Factors determining DNA double-strand break repair pathway choice in G2 phase. *EMBO J* 30: 1079–1092
- Shigechi T, Tomida J, Sato K, Kobayashi M, Eykelenboom JK, Pessina F, Zhang Y, Uchida E, Ishiai M, Lowndes NF et al (2012) ATR-ATRIP kinase complex triggers activation of the Fanconi anemia DNA repair pathway. *Cancer Res* 72: 1149–1156
- Shinozaki T, Nota A, Taya Y, Okamoto K (2003) Functional role of Mdm2 phosphorylation by ATR in attenuation of p53 nuclear export. *Oncogene* 22: 8870–8880
- Shiotani B, Zou L (2009) Single-stranded DNA orchestrates an ATM-to-ATR switch at DNA breaks. *Mol Cell* 33: 547–558
- Sidorova JM, Breeden LL (1997) Rad53-dependent phosphorylation of Swi6 and down-regulation of CLN1 and CLN2 transcription occur in response to DNA damage in *Saccharomyces cerevisiae*. *Genes Dev* 11: 3032–3045
- Sidorova JM, Breeden LL (2003) Rad53 checkpoint kinase phosphorylation site preference identified in the Swi6 protein of *Saccharomyces cerevisiae*. *Mol Cell Biol* 23: 3405–3416
- Singh TR, Ali AM, Paramasivam M, Pradhan A, Wahengbam K, Seidman MM, Meetei AR (2013) ATR-dependent phosphorylation of FANCM at serine 1045 is essential for FANCM functions. *Cancer Res* 73: 4300–4310
- Smits VAJ, Cabrera E, Freire R, Gillespie DA (2019) Clasp – checkpoint adaptor and DNA replication factor. *FEBS J* 286: 441–455
- Smolka MB, Albuquerque CP, Chen SH, Zhou H (2007) Proteome-wide identification of *in vivo* targets of DNA damage checkpoint kinases. *Proc Natl Acad Sci USA* 104: 10364–10369
- Somyajit K, Basavaraju S, Scully R, Nagaraju G (2013) ATM- and ATR-mediated phosphorylation of XRCC3 regulates DNA double-strand break-induced checkpoint activation and repair. *Mol Cell Biol* 33: 1830–1844
- Soniati MM, Myler LR, Kuo HC, Paull TT, Finkelstein IJ (2019) RPA phosphorylation inhibits DNA resection. *Mol Cell* 75: 145–153.e5
- Sorensen CS, Hansen LT, Dziegielewska J, Syljuasen RG, Lundin C, Bartek J, Helleday T (2005) The cell-cycle checkpoint kinase Chk1 is required for mammalian homologous recombination repair. *Nat Cell Biol* 7: 195–201
- Sorensen CS, Syljuasen RG (2012) Safeguarding genome integrity: the checkpoint kinases ATR, CHK1 and WEE1 restrain CDK activity during normal DNA replication. *Nucleic Acids Res* 40: 477–486
- Soulier J, Lowndes NF (1999) The BRCT domain of the *S. cerevisiae* checkpoint protein Rad9 mediates a Rad9-Rad9 interaction after DNA damage. *Curr Biol* 9: 551–554
- Stevens C, Smith L, La Thangue NB (2003) Chk2 activates E2F-1 in response to DNA damage. *Nat Cell Biol* 5: 401–409
- Stewart GS, Wang B, Bignell CR, Taylor AM, Elledge SJ (2003) MDC1 is a mediator of the mammalian DNA damage checkpoint. *Nature* 421: 961–966
- Stokes MP, Rush J, Macneill J, Ren JM, Sprott K, Nardone J, Yang V, Beausoleil SA, Gygi SP, Livingstone M et al (2007) Profiling of UV-induced ATM/ATR signaling pathways. *Proc Natl Acad Sci USA* 104: 19855–19860
- Stucki M, Clapperton JA, Mohammad D, Yaffe MB, Smerdon SJ, Jackson SP (2005) MDC1 directly binds phosphorylated histone H2AX to regulate cellular responses to DNA double-strand breaks. *Cell* 123: 1213–1226
- Sugiyama T, New JH, Kowalczykowski SC (1998) DNA annealing by RAD52 protein is stimulated by specific interaction with the complex of replication protein A and single-stranded DNA. *Proc Natl Acad Sci USA* 95: 6049–6054
- Sugiyama T, Kowalczykowski SC (2002) Rad52 protein associates with replication protein A (RPA)-single-stranded DNA to accelerate

- Rad51-mediated displacement of RPA and presynaptic complex formation. *J Biol Chem* 277: 31663–31672
- Sun Z, Fay DS, Marini F, Foiani M, Stern DF (1996) Spk1/Rad53 is regulated by Mec1-dependent protein phosphorylation in DNA replication and damage checkpoint pathways. *Genes Dev* 10: 395–406
- Sun Z, Hsiao J, Fay DS, Stern DF (1998) Rad53 FHA domain associated with phosphorylated Rad9 in the DNA damage checkpoint. *Science* 281: 272–274
- Szakai B, Brnzei D (2013) Premature Cdk1/Cdc5/Mus81 pathway activation induces aberrant replication and deleterious crossover. *EMBO J* 32: 1155–1167
- Tanaka H, Arakawa H, Yamaguchi T, Shiraishi K, Fukuda S, Matsui K, Takei Y, Nakamura Y (2000) A ribonucleotide reductase gene involved in a p53-dependent cell-cycle checkpoint for DNA damage. *Nature* 404: 42–49
- Taschner M, Harreman M, Teng Y, Gill H, Anindya R, Maslen SL, Skehel JM, Waters R, Svejstrup JQ (2010) A role for checkpoint kinase-dependent Rad26 phosphorylation in transcription-coupled DNA repair in *Saccharomyces cerevisiae*. *Mol Cell Biol* 30: 436–446
- Toh GW, O'Shaughnessy AM, Jimeno S, Dobbie IM, Grenon M, Maffini S, O'Rourke A, Lowndes NF (2006) Histone H2A phosphorylation and H3 methylation are required for a novel Rad9 DSB repair function following checkpoint activation. *DNA Repair (Amst)* 5: 693–703
- Toh GW, Sugawara N, Dong J, Toth R, Lee SE, Haber JE, Rouse J (2010) Mec1/Tel1-dependent phosphorylation of Slx4 stimulates Rad1-Rad10-dependent cleavage of non-homologous DNA tails. *DNA Repair (Amst)* 9: 718–726
- Toledo LI, Altmeyer M, Rask MB, Lukas C, Larsen DH, Povlsen LK, Bekker-Jensen S, Mailand N, Bartek J, Lukas J (2013) ATR prohibits replication catastrophe by preventing global exhaustion of RPA. *Cell* 155: 1088–1103
- Toledo L, Neelsen KJ, Lukas J (2017) Replication catastrophe: when a checkpoint fails because of exhaustion. *Mol Cell* 66: 735–749
- Tomimatsu N, Mukherjee B, Harris JL, Boffo FL, Hardebeck MC, Potts PR, Khanna KK, Burma S (2017) DNA-damage-induced degradation of EXO1 exonuclease limits DNA end resection to ensure accurate DNA repair. *J Biol Chem* 292: 10779–10790
- Travesa A, Kuo D, de Bruin RA, Kalashnikova TI, Guaderrama M, Thai K, Aslanian A, Smolka MB, Yates III JR, Ideker T et al (2012) DNA replication stress differentially regulates G1/S genes via Rad53-dependent inactivation of Nrm1. *EMBO J* 31: 1811–1822
- Tripathi V, Nagarjuna T, Sengupta S (2007) BLM helicase-dependent and -independent roles of 53BP1 during replication stress-mediated homologous recombination. *J Cell Biol* 178: 9–14
- Usui T, Foster SS, Petrini JH (2009) Maintenance of the DNA-damage checkpoint requires DNA-damage-induced mediator protein oligomerization. *Mol Cell* 33: 147–159
- Valerio-Santiago M, de Los Santos-Velazquez AI, Monje-Casas F (2013) Inhibition of the mitotic exit network in response to damaged telomeres. *PLoS Genet* 9: e1003859
- Vasianovich Y, Harrington LA, Makovets S (2014) Break-induced replication requires DNA damage-induced phosphorylation of Pif1 and leads to telomere lengthening. *PLoS Genet* 10: e1004679
- Vassin VM, Anantha RW, Sokolova E, Kanner S, Borowiec JA (2009) Human RPA phosphorylation by ATR stimulates DNA synthesis and prevents ssDNA accumulation during DNA-replication stress. *J Cell Sci* 122: 4070–4080
- Vialard JE, Gilbert CS, Green CM, Lowndes NF (1998) The budding yeast Rad9 checkpoint protein is subjected to Mec1/Tel1-dependent hyperphosphorylation and interacts with Rad53 after DNA damage. *EMBO J* 17: 5679–5688
- Vidanes GM, Sweeney FD, Galicia S, Cheung S, Doyle JP, Durocher D, Toczyski DP (2010) CDC5 inhibits the hyperphosphorylation of the checkpoint kinase Rad53, leading to checkpoint adaptation. *PLoS Biol* 8: e1000286
- Vriend LE, Prakash R, Chen CC, Vanoli F, Cavallo F, Zhang Y, Jasin M, Krawczyk PM (2016) Distinct genetic control of homologous recombination repair of Cas9-induced double-strand breaks, nicks and paired nicks. *Nucleic Acids Res* 44: 5204–5217
- Wagner SA, Oehler H, Voigt A, Dalic D, Freiwald A, Serve H, Beli P (2016) ATR inhibition rewires cellular signaling networks induced by replication stress. *Proteomics* 16: 402–416
- Wang H, Liu D, Wang Y, Qin J, Elledge SJ (2001) Pds1 phosphorylation in response to DNA damage is essential for its DNA damage checkpoint function. *Genes Dev* 15: 1361–1372
- Wang B, Matsuoka S, Carpenter PB, Elledge SJ (2002) 53BP1, a mediator of the DNA damage checkpoint. *Science* 298: 1435–1438
- Wang H, Powell SN, Iliakis G, Wang Y (2004a) ATR affecting cell radiosensitivity is dependent on homologous recombination repair but independent of nonhomologous end joining. *Cancer Res* 64: 7139–7143
- Wang YC, Nnakwe C, Lane WS, Modesti M, Frank KM (2004b) Phosphorylation and regulation of DNA ligase IV stability by DNA-dependent protein kinase. *J Biol Chem* 279: 37282–37290
- Wang X, Kennedy RD, Ray K, Stuckert P, Ellenberger T, D'Andrea AD (2007) Chk1-mediated phosphorylation of FANCD1 is required for the Fanconi anemia/BRCA pathway. *Mol Cell Biol* 27: 3098–3108
- Wang J, Gong Z, Chen J (2011) MDC1 collaborates with TopBP1 in DNA replication checkpoint control. *J Cell Biol* 193: 267–273
- Weinert TA, Hartwell LH (1988) The RAD9 gene controls the cell cycle response to DNA damage in *Saccharomyces cerevisiae*. *Science* 241: 317–322
- Weinert T (1997) A DNA damage checkpoint meets the cell cycle engine. *Science* 277: 1450–1451
- Wu X, Shell SM, Yang Z, Zou Y (2006) Phosphorylation of nucleotide excision repair factor xeroderma pigmentosum group A by ataxia telangiectasia mutated and Rad3-related-dependent checkpoint pathway promotes cell survival in response to UV irradiation. *Cancer Res* 66: 2997–3005
- Wu L, Luo K, Lou Z, Chen J (2008) MDC1 regulates intra-S-phase checkpoint by targeting NBS1 to DNA double-strand breaks. *Proc Natl Acad Sci USA* 105: 11200–11205
- Xu Y, Ashley T, Brainerd EE, Bronson RT, Meyn MS, Baltimore D (1996) Targeted disruption of ATM leads to growth retardation, chromosomal fragmentation during meiosis, immune defects, and thymic lymphoma. *Genes Dev* 10: 2411–2422
- Xu X, Tsvetkov LM, Stern DF (2002) Chk2 activation and phosphorylation-dependent oligomerization. *Mol Cell Biol* 22: 4419–4432
- Yazinski SA, Comaills V, Buisson R, Genois MM, Nguyen HD, Ho CK, Todorova Kwan T, Morris R, Lauffer S, Nussenzweig A et al (2017) ATR inhibition disrupts rewired homologous recombination and fork protection pathways in PARP inhibitor-resistant BRCA-deficient cancer cells. *Genes Dev* 31: 318–332
- Yekezare M, Gomez-Gonzalez B, Diffley JF (2013) Controlling DNA replication origins in response to DNA damage – inhibit globally, activate locally. *J Cell Sci* 126: 1297–1306
- Yoo HY, Shevchenko A, Dunphy WG (2004) Mcm2 is a direct substrate of ATM and ATR during DNA damage and DNA replication checkpoint responses. *J Biol Chem* 279: 53353–53364

- Yoo HY, Jeong SY, Dunphy WG (2006) Site-specific phosphorylation of a checkpoint mediator protein controls its responses to different DNA structures. *Genes Dev* 20: 772–783
- Yu Y, Mahaney BL, Yano K, Ye R, Fang S, Douglas P, Chen DJ, Lees-Miller SP (2008) DNA-PK and ATM phosphorylation sites in XLF/Cernunnos are not required for repair of DNA double strand breaks. *DNA Repair (Amst)* 7: 1680–1692
- Zegerman P, Diffley JF (2010) Checkpoint-dependent inhibition of DNA replication initiation by Sld3 and Dbf4 phosphorylation. *Nature* 467: 474–478
- Zha S, Jiang W, Fujiwara Y, Patel H, Goff PH, Brush JW, Dubois RL, Alt FW (2011) Ataxia telangiectasia-mutated protein and DNA-dependent protein kinase have complementary V(D)J recombination functions. *Proc Natl Acad Sci USA* 108: 2028–2033
- Zhang X, Succi J, Feng Z, Prithivirajasingh S, Story MD, Legerski RJ (2004) Artemis is a phosphorylation target of ATM and ATR and is involved in the G2/M DNA damage checkpoint response. *Mol Cell Biol* 24: 9207–9220
- Zhang T, Nirantar S, Lim HH, Sinha I, Surana U (2009) DNA damage checkpoint maintains CDH1 in an active state to inhibit anaphase progression. *Dev Cell* 17: 541–551
- Zhao X, Muller EG, Rothstein R (1998) A suppressor of two essential checkpoint genes identifies a novel protein that negatively affects dNTP pools. *Mol Cell* 2: 329–340
- Zhao H, Piwnicka-Worms H (2001) ATR-mediated checkpoint pathways regulate phosphorylation and activation of human Chk1. *Mol Cell Biol* 21: 4129–4139
- Zhao X, Chabes A, Domkin V, Thelander L, Rothstein R (2001) The ribonucleotide reductase inhibitor Sml1 is a new target of the Mec1/Rad53 kinase cascade during growth and in response to DNA damage. *EMBO J* 20: 3544–3553
- Zhao X, Rothstein R (2002) The Dun1 checkpoint kinase phosphorylates and regulates the ribonucleotide reductase inhibitor Sml1. *Proc Natl Acad Sci USA* 99: 3746–3751
- Zhi G, Wilson JB, Chen X, Krause DS, Xiao Y, Jones NJ, Kupfer GM (2009) Fanconi anemia complementation group FANCD2 protein serine 331 phosphorylation is important for fanconi anemia pathway function and BRCA2 interaction. *Cancer Res* 69: 8775–8783
- Zhou BB, Elledge SJ (2000) The DNA damage response: putting checkpoints in perspective. *Nature* 408: 433–439
- Zhou Y, Paull TT (2013) DNA-dependent protein kinase regulates DNA end resection in concert with Mre11-Rad50-Nbs1 (MRN) and ataxia telangiectasia-mutated (ATM). *J Biol Chem* 288: 37112–37125
- Zimmermann M, Lottersberger F, Buonomo SB, Sfeir A, de Lange T (2013) 53BP1 regulates DSB repair using Rif1 to control 5' end resection. *Science* 339: 700–704
- Zou L (2013) Four pillars of the S-phase checkpoint. *Genes Dev* 27: 227–233