

---

UNIVERSITY OF TEXAS AT AUSTINHomework assignment 10

Continuous probability. The normal distribution.

---

Please, provide your final answer only to the following questions:

**Problem 10.1.** (2 points) A linear combination of two jointly normally distributed random variables is always also normally distributed. Assume that a constant is also considered normally distributed with variance zero. *True or false?*

**Solution: TRUE**

**Problem 10.2.** (2 points) If the random variable  $X$  has the distribution function  $F_X$ , then the distribution function of the random variable  $Y = |X|$  equals

$$F_Y(y) = 2F_X(y).$$

*True or false?*

**Solution: FALSE**

**Problem 10.3.** (2 points) Let  $X_1, \dots, X_n$  be random variables with finite expectations and let  $\alpha_1, \dots, \alpha_n$  be constants. Then, we always have that

$$\mathbb{E}[\alpha_1 X_1 + \dots + \alpha_n X_n] = \sum_{i=1}^n \alpha_i \mathbb{E}[X_i].$$

*True or false?*

**Solution: TRUE**

---

Please, provide your complete solution to the following problems. Only the final answer without justification will receive zero credit.

**Problem 10.4.** (2 points) It is possible that a cumulative distribution function be even. *True or false? Why?*

**Solution: FALSE**

Any cumulative distribution function  $F$  must satisfy  $F(-\infty) = 0$  and  $F(+\infty) = 1$ . Therefore, it cannot be even.

**Problem 10.5.** (2 pts) If the random variable  $X$  is standard normal, then the distribution function of the random variable  $Y = |X|$  equals

$$F_Y(a) = 2\Phi(a) - 1 \text{ for every } a \geq 0.$$

*True or false? Why?*

**Solution: TRUE**

For every  $a \geq 0$ ,

$$\begin{aligned}
 F_Y(a) &= \mathbb{P}[Y \leq a] \\
 &= \mathbb{P}[|X| \leq a] \\
 &= \mathbb{P}[-a \leq X \leq a] \\
 &= \mathbb{P}[X \leq a] - \mathbb{P}[X < -a] \\
 &= \mathbb{P}[X \leq a] - (1 - \mathbb{P}[X \geq -a]) \\
 &= \mathbb{P}[X \leq a] - (1 - \mathbb{P}[X \leq a]) \\
 &= 2\Phi(a) - 1.
 \end{aligned}$$

**Problem 10.6.** (10 pts) Let  $Z$  be a standard normal random variable. Find the following probabilities:

- i.  $\mathbb{P}[-1.33 < Z \leq 0.24]$
- ii.  $\mathbb{P}[0.49 < |Z|]$
- iii.  $\mathbb{P}[Z^4 < 0.0256]$
- iv.  $\mathbb{P}[e^{2Z} < 2.25]$
- v.  $\mathbb{P}[\frac{1}{Z} < 2]$

**Solution:**

i.

$$\begin{aligned}
 \mathbb{P}[-1.33 < Z \leq 0.24] &= \mathbb{P}[Z \leq 0.24] - \mathbb{P}[Z \leq -1.33] = \mathbb{P}[Z \leq 0.24] - (1 - \mathbb{P}[Z \leq 1.33]) \\
 &= 0.5948 - 1 + 0.9082 = 0.503
 \end{aligned}$$

ii.

$$\begin{aligned}
 \mathbb{P}[0.49 < |Z|] &= \mathbb{P}[Z < -0.49] + \mathbb{P}[0.49 < Z] = 2\mathbb{P}[Z > 0.49] \\
 &= 2(1 - \mathbb{P}[Z \leq 0.49]) = 2(1 - 0.6879) = 0.6242
 \end{aligned}$$

iii.

$$\begin{aligned}
 \mathbb{P}[Z^4 < 0.0256] &= \mathbb{P}[|Z| < \sqrt[4]{0.0256}] = \mathbb{P}[|Z| < 0.4] = \mathbb{P}[Z < 0.4] - \mathbb{P}[Z < -0.4] \\
 &= 2\mathbb{P}[Z < 0.4] - 1 = 2(0.6554) - 1 = 0.3108
 \end{aligned}$$

iv.

$$\mathbb{P}[e^{2Z} < 2.25] = \mathbb{P}[2Z < \ln(2.25)] = \mathbb{P}[Z < 0.5 \ln(2.25)] \approx \mathbb{P}[Z \leq 0.41] = 0.6591$$

v.

$$\begin{aligned}
 \mathbb{P}[\frac{1}{Z} < 2] &= \mathbb{P}[\frac{1}{Z} < 0] + \mathbb{P}[0 < \frac{1}{Z} < 2] \\
 &= \mathbb{P}[Z < 0] + \mathbb{P}[Z > 0.5] = 0.5 + (1 - \mathbb{P}[Z \leq 0.5]) = 0.5 + (1 - 0.6915) = 0.8085.
 \end{aligned}$$

**Problem 10.7.** (5 points) Let the density function of a random variable  $X$  be given as

$$f_X(x) = cx, \quad x \in [0, 1],$$

for some constant  $c$ . Find  $\mathbb{E}[X^3]$ .

**Solution:** If you draw the graph of  $f_X$ , you will see that the integral of the density is the area of a triangle. You get  $c = 2$ . Then,

$$\mathbb{E}[X^3] = 2 \int_0^1 x^4 dx = 2/5.$$

**Problem 10.8.** (10 points) Two laser pointers are used to measure the length  $\ell$  of a building. The error made by the less accurate laser pointer is normally distributed with mean 0 and standard deviation  $0.0144\ell$ . The error made by the more accurate laser pointer is normally distributed with mean 0 and standard deviation  $0.0036\ell$ . The errors from the two laser pointers are independent of each other. Calculate the probability that the average value of the two measurements is within  $0.001\ell$  of the true length  $\ell$  of the building.

**Solution:** Let the error of the measurement from the first laser pointer be denoted by  $\varepsilon_1$  and let the error of the measurement from the second laser pointer be denoted by  $\varepsilon_2$ . According to our modelling assumptions

$$\varepsilon_1 \sim \text{Normal}(\text{mean} = 0, \text{sd} = 0.0144\ell),$$

$$\varepsilon_2 \sim \text{Normal}(\text{mean} = 0, \text{sd} = 0.0036\ell).$$

Let the average error be  $\varepsilon = \frac{1}{2}(\varepsilon_1 + \varepsilon_2)$ . Then,  $\varepsilon$  is normally distributed with mean 0 and variance

$$\text{Var}[\varepsilon] = \frac{1}{4}(\text{Var}[\varepsilon_1] + \text{Var}[\varepsilon_2]) = \frac{1}{4}((0.0144\ell)^2 + (0.0036\ell)^2).$$

So,

$$\varepsilon \sim \text{Normal}(\text{mean} = 0, \text{sd} = 0.0074\ell)$$

We need to find the probability  $\mathbb{P}[|\varepsilon| < 0.001\ell]$ . We have

$$\begin{aligned} \mathbb{P}[|\varepsilon| < 0.001\ell] &= \mathbb{P}[-0.001\ell < \varepsilon < 0.001\ell] \\ &= \mathbb{P}\left[\frac{-0.001\ell - 0}{0.0074\ell} < \frac{\varepsilon - 0}{0.0074\ell} < \frac{0.001\ell - 0}{0.0074\ell}\right] = \mathbb{P}[-0.1351 < Z < 0.1351] \end{aligned}$$

where  $Z \sim N(0, 1)$ . So,

$$\mathbb{P}[|\varepsilon| < 0.001\ell] = 2N(0.1351) - 1 \approx 2N(0.14) - 1 = 2(0.5557) - 1 = 0.1114.$$

**Problem 10.9.** (10 points) An astronomical instrument measures the distance  $d$  to a far-away planet. You know that the instrument is calibrated so that its measurement error is normally distributed, centered around zero, and with variance  $(0.0001d)^2$ . The different measurements using this same instrument are assumed to be independent. How many independent measurements would you have to perform so that your average is within  $10^{-6}d$  with probability 99%?

**Solution:** For every  $i \in \mathbb{N}$ , let  $\varepsilon_i$  denote the error of the  $i^{\text{th}}$  measurement. Then, with  $n$  being the number of measurements, the average error  $\varepsilon = \frac{1}{n} \sum_{i=1}^n \varepsilon_i$  is normally distributed with mean 0 and with variance

$$\text{Var}[\varepsilon] = \text{Var}\left[\frac{1}{n}(\varepsilon_1 + \varepsilon_2 + \dots + \varepsilon_n)\right] = \frac{1}{n^2} \text{Var}[\varepsilon_1 + \varepsilon_2 + \dots + \varepsilon_n].$$

Due to the fact that different measurements are independent, we get

$$\text{Var}[\varepsilon] = \frac{1}{n^2}(\text{Var}[\varepsilon_1] + \text{Var}[\varepsilon_2] + \dots + \text{Var}[\varepsilon_n]) = \frac{1}{n^2}(n)(0.0001d)^2 = \frac{(0.0001d)^2}{n}.$$

The probability we are given is

$$\begin{aligned} \mathbb{P}[|\varepsilon| < 10^{-6}d] = 0.99 &\Rightarrow \mathbb{P}[-10^{-6}d < \varepsilon < 10^{-6}d] = 0.99 \\ &\Rightarrow \mathbb{P}\left[-\frac{10^{-6}d - 0}{\frac{0.0001d}{\sqrt{n}}} < \frac{\varepsilon - 0}{\frac{0.0001d}{\sqrt{n}}} < \frac{10^{-6}d - 0}{\frac{0.0001d}{\sqrt{n}}}\right] = 0.99 \\ &\Rightarrow \mathbb{P}\left[-\frac{10^{-6}\sqrt{n}}{10^{-4}} < Z < \frac{10^{-6}\sqrt{n}}{10^{-4}}\right] = 0.99 \end{aligned}$$

where  $Z \sim N(0, 1)$ . So, we have to find a condition for  $n$  in

$$\mathbb{P}\left[-\frac{\sqrt{n}}{100} < Z < \frac{\sqrt{n}}{100}\right] = 0.99 \Rightarrow 2N\left(\frac{\sqrt{n}}{100}\right) - 1 = 0.99 \Rightarrow N\left(\frac{\sqrt{n}}{100}\right) = 0.995.$$

From the standard normal tables, we get

$$\frac{\sqrt{n}}{100} = 2.575 \Rightarrow n \geq (257.5)^2 = 66306.25 \Rightarrow n \geq 66307.$$

**Problem 10.10.** (5 points) The profit of a certain company are modelled using a normal distribution with mean 1,000,000 and standard deviation 400,000. Given that the profit is positive, what is the probability that it is below 1,200,000?

**Solution:** Let  $X$  denote the profit random variable. Then, we need to calculate

$$\mathbb{P}[X < 1200000 | X > 0] = \frac{\mathbb{P}[0 < X < 1200000]}{\mathbb{P}[X > 0]} = \frac{\mathbb{P}[X < 1200000] - \mathbb{P}[X \leq 0]}{\mathbb{P}[X > 0]}.$$

The given model for  $X$  is

$$X \sim \text{Normal}(\text{mean} = 1000000, \text{sd} = 400000).$$

First, let's calculate the probability that the profit is below zero

$$\mathbb{P}[X \leq 0] = \mathbb{P}\left[\frac{X - 1000000}{400000} \leq \frac{0 - 1000000}{400000}\right] = \mathbb{P}[Z \leq -2.5] = 0.0062.$$

Above, we denoted by  $Z$  a standard normal random variable. So,  $\mathbb{P}[X > 0] = 0.9938$ . Next, we calculate the probability that the profit is below 1,200,000.

$$\mathbb{P}[X \leq 1200000] = \mathbb{P}\left[\frac{X - 1000000}{400000} \leq \frac{1200000 - 1000000}{400000}\right] = \mathbb{P}[Z \leq 0.5] = 0.6915.$$

Finally, our answer is

$$\mathbb{P}[X < 1200000 | X > 0] = \frac{0.6915 - 0.0062}{0.9938} = 0.6895754.$$