


UNIVERSITY OF TEXAS AT AUSTIN

Homework Assignment 8Classification trees.

Please, provide your complete solutions to the following problems. Final answers only, even if correct will earn zero points for those problems.

**Problem 8.1.** (10 points) Solve Problem 8.4.3 (pp.361-362) in the textbook.

**Solution:** Here is the plot:



**Problem 8.2.** (10 points) A classification tree is constructed to predict whether a student will pass *Predictive Analytics*. There are two categorical predictors:

- $X_1$  indicating whether the student passed *Linear Algebra* prior to enrolling in *Predictive Analytics*, and

- $X_2$  indicating whether the student passed *Mathematical Statistics* prior to enrolling in *Predictive Analytics*.

Here is the table from a data set of students:

$X_1$	$X_2$	outcome
0	0	5 passed, 20 didn't
1	0	10 passed, 10 didn't
0	1	15 passed, 20 didn't
1	1	15 passed, 5 didn't

What's the first split made using the Gini index? Be careful to calculate the **weighted** average.

**Solution:** Since both  $X_1$  and  $X_2$  are binary, we can make the first split in a unique way with respect to  $X_1$  or with respect to  $X_2$ . If we split along  $X_1$ , we have that

$X_1 = 0$  20 passed and 40 didn't;

$X_1 = 1$  25 passed and 15 didn't.

So, the Gini index is

$$\frac{60}{100} \cdot 2 \cdot \left( \frac{20}{60} \left( 1 - \frac{20}{60} \right) \right) + \frac{40}{100} \cdot 2 \cdot \left( \frac{25}{40} \left( 1 - \frac{25}{40} \right) \right) = 0.4541667$$

If we split along  $X_2$ , we have that

$X_2 = 0$  15 passed and 30 didn't;

$X_2 = 1$  30 passed and 25 didn't.

So, the Gini index is

$$\frac{45}{100} \cdot 2 \cdot \left( \frac{15}{45} \left( 1 - \frac{15}{45} \right) \right) + \frac{55}{100} \cdot 2 \cdot \left( \frac{30}{55} \left( 1 - \frac{30}{55} \right) \right) = 0.4727273$$

Since the first Gini index is smaller, we should split along  $X_1$ .

**Problem 8.3.** (30 points) A classification tree is fitted to be used to classify drivers into one of three categories: *Good*, *Medium* and *Bad*. There are three terminal nodes in the classification tree designating the regions  $R_1$ ,  $R_2$  and  $R_3$ . Here is the breakdown of categories in each region:

Region	Good	Medium	Bad
$R_1$	70	20	10
$R_2$	30	50	10
$R_3$	20	25	35

Calculate the overall classification error rate, Gini index, and cross-entropy using **weighted averages**.

**Solution:** There are  $n_1 = 100$  observations in  $R_1$ ,  $n_2 = 90$  observations in  $R_2$ , and  $n_3 = 80$  observations in  $R_3$ . The total number of observations is  $n = 270$ . So, in each calculation, the weight for  $R_1$  is  $w_1 = \frac{100}{270} = \frac{10}{27}$ , the weight for  $R_2$  is  $w_2 = \frac{90}{270} = \frac{1}{3}$ , and the weight for  $R_3$  is  $w_3 = \frac{80}{270} = \frac{8}{27}$ .

- (i) In  $R_1$ , the most common category is *Good*, so the classification error rate equals

$$\frac{20 + 10}{100} = 0.3.$$

In  $R_2$ , the most common category is *Medium*, so the classification error rate equals

$$\frac{30 + 10}{90} = \frac{4}{9}.$$

In  $R_3$ , the most common category is *Bad*, so the classification error rate equals

$$\frac{20 + 25}{80} = 0.5625.$$

Overall, we obtain

$$\frac{10}{27} \cdot \frac{3}{10} + \frac{1}{3} \cdot \frac{4}{9} + \frac{8}{27} \cdot \frac{9}{16} = 0.4259259.$$

Equivalently, and way more easily, we could have calculated

$$\frac{20 + 10 + 30 + 10 + 20 + 25}{100 + 90 + 80} = \frac{115}{270} = 0.4259259.$$

(ii) In  $R_1$ , the Gini index is

$$\frac{70}{100} \cdot \frac{30}{100} + \frac{20}{100} \cdot \frac{80}{100} + \frac{10}{100} \cdot \frac{90}{100} = 0.46$$

In  $R_2$ , the Gini index is

$$\frac{30}{90} \cdot \frac{60}{90} + \frac{50}{90} \cdot \frac{40}{90} + \frac{10}{90} \cdot \frac{80}{90} = 0.5679012$$

In  $R_3$ , the Gini index is

$$\frac{20}{80} \cdot \frac{60}{80} + \frac{25}{80} \cdot \frac{55}{80} + \frac{35}{80} \cdot \frac{45}{80} = 0.6484375$$

Overall, we have

$$\frac{10}{27}(0.46) + \frac{1}{3}(0.5679012) + \frac{8}{27}(0.6484375) = 0.5518004$$

(iii) In  $R_1$ , the cross-entropy is

$$-(0.7 \ln(0.7) + 0.2 \ln(0.2) + 0.1 \ln(0.1)) = 0.8018186.$$

In  $R_2$ , the cross-entropy is

$$-\left(\frac{30}{90} \ln\left(\frac{30}{90}\right) + \frac{50}{90} \ln\left(\frac{50}{90}\right) + \frac{10}{90} \ln\left(\frac{10}{90}\right)\right) = 0.9368883.$$

In  $R_3$ , the cross-entropy is

$$-\left(\frac{20}{80} \ln\left(\frac{20}{80}\right) + \frac{25}{80} \ln\left(\frac{25}{80}\right) + \frac{35}{80} \ln\left(\frac{35}{80}\right)\right) = 1.07173$$

Overall, we have

$$\frac{10}{27}(0.8018186) + \frac{1}{3}(0.9368883) + \frac{8}{27}(1.07173) = 0.9268156.$$