

## Approximate Confidence Intervals for $p$ .

Consider a population in which a specific trait occurs w/ an unknown probability  $p$ .

Let  $(Y_1, Y_2, \dots, Y_n)$  be a random sample from the Bernoulli dist'n w/ the unknown  $p$ .

Goal: Designing a confidence interval for  $p$ .

Idea: Look @ the natural point estimator for  $p$ .

$$\rightarrow \bar{Y} = \frac{1}{n} (Y_1 + Y_2 + \dots + Y_n) \sim B(n, p)$$

Note:  $E[\bar{Y}] = p$  of course, the dist'n of  $\bar{Y}$  depends on  $p$ .

We need a pivotal quantity.

By de Moivre-Laplace Thm:

$$\text{Set } S_n = Y_1 + \dots + Y_n$$

$$\text{Then, } \frac{S_n - np}{\sqrt{np(1-p)}} \xrightarrow{D} N(0, 1)$$

$$\text{We note } \bar{Y} = \frac{1}{n} S_n$$

$$U = \frac{\bar{Y} - p}{\sqrt{\frac{p(1-p)}{n}}} \xrightarrow{D} N(0, 1) \text{ is still not a pivotal quantity}$$

But we can still create an approximate confidence interval based on it!

Say,  $C$  is a confidence level.

$$\text{Let } z^* = \Phi^{-1}\left(\frac{1+C}{2}\right) = q_{\text{norm}}((1+C)/2)$$

$$\mathbb{P}[-z^* \leq u \leq z^*] \approx C \quad \text{dM.L.}$$

$$\mathbb{P}\left[-z^* \leq \frac{\bar{Y} - p}{\sqrt{\frac{p(1-p)}{n}}} \leq z^*\right] \approx C$$

$$\mathbb{P}\left[\bar{Y} - z^* \sqrt{\frac{p(1-p)}{n}} \leq p \leq \bar{Y} + z^* \sqrt{\frac{p(1-p)}{n}}\right] \approx C$$

Usually, we write

$$\bar{Y} = \hat{p}$$

We construct the confidence interval for  $p$  as

$$p = \hat{p} \pm z^* \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

**Problem 16.2.** Gallup's inaugural measure of global loneliness shows over one in five people worldwide (23%) said they felt loneliness "a lot of the day yesterday."<sup>1</sup> However, there were considerable variations between countries. For instance, out of 1000 individuals polled in Taiwan, 11% reported having felt loneliness "a lot of the day" before. What 90% confidence would you report for the population proportion of Taiwanese who had felt lonely the day before?

→:  $\hat{p} = 0.11$

$z^* = \Phi^{-1}(0.95) = 1.645$

$p = \hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} = 0.11 \pm 1.645 \sqrt{\frac{0.11 \cdot 0.89}{1000}}$

□

Example. How do we figure out the necessary sample size w/ a required margin of error  $m$ ?

→:  $n = ?$

$z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq m$  /<sup>2</sup>

w/  $(z^*)$  the critical value for the confidence level  $C$


$(z^*)^2 \cdot \frac{\hat{p}(1-\hat{p})}{n} \leq m^2$

$(z^*)^2 \cdot \frac{\hat{p}(1-\hat{p})}{m^2} \leq n$

We cannot know  $\hat{p}$  before knowing  $n$ .

<sup>1</sup><https://news.gallup.com/poll/646718/people-worldwide-feel-lonely-lot.aspx>

Consider  $\hat{p}(1-\hat{p})$  as a function of  $\hat{p}$ .



The conservative choice for  $\hat{p}$  is  $\frac{1}{2}$ .

The conservative choice for the sample size

$$n \geq \left( \frac{z^*}{2m} \right)^2$$

