

# M378K Introduction to Mathematical Statistics

## Problem Set #2

### Discrete random variables.

---

**2.1. Probability mass function.** Recall the following definition from the last class:

**Definition 2.1.** Given a set  $B$ , we say that a random variable  $Y$  is  $B$ -valued if

$$\mathbb{P}[Y \in B] = 1.$$

We reserve special terminology for random variables  $Y$  depending on the cardinality of the set  $B$  from the above definition. In particular, we have the following definition:

**Definition 2.2.** A random variable  $Y$  is said to be discrete if there exists a set  $S$  such that :

- $Y$  is  $S$ -valued, and
- $S$  is either **finite** or **countable**.

**Problem 2.1.** Provide an example of a **discrete** random variable.

**Solution:** A roll of a fair die.

Our next task is to try to keep track of the probabilities that  $Y$  takes specific values from  $S$ . In order to be more "economical", we introduce the following concept:

**Definition 2.3.** The support  $S_Y$  of a random variable  $Y$  is the **smallest** set  $S$  such that  $Y$  is  $S$ -valued.

**Problem 2.2.** What is the **support** of the random variable you provided as an example in the above problem?

**Solution:**

$$S_Y = \{1, 2, 3, 4, 5, 6\}$$

**Problem 2.3.** Let  $y \in S_Y$  where  $Y$  is a discrete random variable. Is it possible to have  $\mathbb{P}[Y = y] = 0$ ?

**Solution:** No, it is not possible. If  $Y$  is a discrete random variable and there exists  $\tilde{y} \in S_Y$  such that  $\mathbb{P}[Y = \tilde{y}] = 0$ , then we can consider the set  $\tilde{S}_Y = S_Y \setminus \{\tilde{y}\}$ . Then,

$$\mathbb{P}[Y \in \tilde{S}_Y] = \mathbb{P}[Y \in S_Y] - \mathbb{P}[Y = \tilde{y}] = \mathbb{P}[Y \in S_Y] = 1 \quad (2.1)$$

and  $\tilde{S}_Y$  is a smaller set than  $S_Y$  such that  $Y$  is  $\tilde{S}_Y$ -valued. This contradicts the definition of support.

Usually, we are interested in calculating and modeling probabilities that look like this

$$\mathbb{P}[Y \in A] \quad \text{for some } A \subseteq S_Y.$$

Note that, if we know the probabilities of the form

$$\mathbb{P}[Y = y] \quad \text{for all } y \in S_Y,$$

then we can calculate any probability of the above form. *How?*

**Solution:**

$$\mathbb{P}[Y \in A] = \sum_{y \in A} \mathbb{P}[Y = y].$$

So, if we "tabulate" the probabilities of the form  $\mathbb{P}[Y = y]$  for all  $y \in S_Y$ , we have sufficient information to calculate any probability of interest to do with the random variable  $Y$ . This observation motivates the following definition:

**Definition 2.4.** *The probability mass function (pmf) of a **discrete** random variable  $Y$  is the function  $p_Y : S_Y \rightarrow \mathbb{R}$  defined as*

$$p_Y(y) = \mathbb{P}[Y = y] \quad \text{for all } y \in S_Y.$$

Can you think of different ways in which to display the pmf?

What is the pmf of the random variable which you provided as an example above?

**Solution:**

$$p_Y(y) = \begin{cases} \frac{1}{6} & \text{if } y \in \{1, 2, 3, 4, 5, 6\}, \\ 0 & \text{otherwise.} \end{cases}$$

What are the immediate properties of every pmf?

**Solution:** For every discrete random variable  $Y$ , its pmf  $p_Y$  satisfies the following two properties:

- $p_Y(y) > 0$  for all  $y \in S_Y$ , and
- $\sum_{y \in S_Y} p_Y(y) = 1$ .

Does the "reverse" hold, i.e., if a function  $p_Y$  satisfies you stated, is it always a pmf of **some** random variable?

**Solution:** Yes, it does. Given a function  $p_Y$  satisfying the two properties above, we can always construct an outcome space  $\Omega$ , a probability  $\mathbb{P}$  on it, and a discrete random variable  $Y$  on it such that  $p_Y$  is the pmf of  $Y$ .

For example, we can consider the outcome space  $\Omega = S_Y$ . Then, we can define the probability measure  $\mathbb{P}$  on  $\Omega$  as

$$\mathbb{P}[\{\omega\}] = p_Y(\omega) \quad \text{for all } \omega \in \Omega.$$

Finally, we can define the random variable  $Y$  as the identity function on  $\Omega$ , i.e.,  $Y(\omega) = \omega$  for all  $\omega \in \Omega$ .

**Problem 2.4.** *The number of pieces of gossip that break out in a particular high school in a week is modeled by a random variable  $Y$  with the following probability mass function:*

$$p_Y(n) = \frac{1}{(n+1)(n+2)} \quad \text{for all } n \in \mathbb{N}_0.$$

*Is the above a well-defined probability mass function?*

**Solution:** We need to verify that the two requirements are satisfied, namely, that

- $p_Y(n) > 0$  for all  $n \in \mathbb{N}_0$ , and
- $\sum_{n=0}^{\infty} p_Y(n) = 1$ .

The first condition is obviously satisfied. As for the second one, we have that for every  $N \in \mathbb{N}_0$

$$\sum_{n=0}^N p_Y(n) = \sum_{n=0}^N \frac{1}{(n+1)(n+2)} = \sum_{n=0}^N \left( \frac{1}{n+1} - \frac{1}{n+2} \right) = 1 - \frac{1}{N+2}.$$

As  $N \rightarrow \infty$ , the above sum goes to 1 (which is a definition of the sum of a series).

**2.2. Conditional probability.** In order to "build" more complicated (and useful!) random variables, it helps to review a bit more probability.

**Definition 2.5.** Let  $E$  and  $F$  be two events on the same  $\Omega$  such that  $\mathbb{P}[E] > 0$ . The conditional probability of  $F$  given  $E$  is defined as

$$\mathbb{P}[F | E] = \frac{\mathbb{P}[E \cap F]}{\mathbb{P}[E]}.$$

Let's spend a moment with the geometric/informational perspective on this definition.

By far, the most popular problems relying on the notion of **conditional probability** are those to do with **specificity** and **sensitivity**<sup>1</sup> of medical tests.

**Problem 2.5.** At any given time, 2% of the population actually has a particular disease.

A test indicates the presence of a particular disease 96% of the time in people who actually have the disease. The same test is positive 1% of the time when actually healthy people are tested.

Calculate the probability that a particular person actually has the disease **given** that they tested positive.

**Solution:** Let  $F$  denote the event that a person has the disease and let  $E$  denote the event that the person tested positive. We need to calculate the following conditional probability:

$$\mathbb{P}[F | E] = \frac{\mathbb{P}[E \cap F]}{\mathbb{P}[E]} = \frac{0.02(0.96)}{0.02(0.96) + 0.98(0.01)} = \frac{2(96)}{2(96) + 98} = \frac{96}{145}.$$

---

<sup>1</sup>[https://en.wikipedia.org/wiki/Sensitivity\\_and\\_specificity](https://en.wikipedia.org/wiki/Sensitivity_and_specificity)

Moreover, now that we remember the definition of **conditional probability**, we can solve interesting problems such as this one:

**Problem 2.6.** *The number of pieces of gossip that break out in a particular high school in a week is modeled by a random variable  $Y$  with the following probability mass function:*

$$p_Y(n) = \frac{1}{(n+1)(n+2)} \quad \text{for all } n \in \mathbb{N}_0.$$

*Calculate the probability that at least one piece of gossip occurred in a week **given** that at most four pieces of gossip occurred.*

**Solution:** Here, we calculate

$$\mathbb{P}[Y \geq 1 \mid Y \leq 4] = \frac{\mathbb{P}[1 \leq Y \leq 4]}{\mathbb{P}[Y \leq 4]} = \frac{p_Y(1) + p_Y(2) + p_Y(3) + p_Y(4)}{p_Y(0) + p_Y(1) + p_Y(2) + p_Y(3) + p_Y(4)}.$$

Using the same reasoning as in part (ii), we get

$$\mathbb{P}[Y \geq 1 \mid Y \leq 4] = \frac{\frac{1}{2} - \frac{1}{6}}{1 - \frac{1}{6}} = \frac{\frac{2}{6}}{\frac{5}{6}} = \frac{2}{5}.$$

### 3. INDEPENDENT EVENTS

What if knowing that an event happened in fact does **not** give any information about the probability of another event?

**Definition 3.1.** We say that events  $E$  and  $F$  on  $\Omega$  are independent if

$$\mathbb{P}[E \cap F] = \mathbb{P}[E]\mathbb{P}[F].$$

In the case when  $E$  or  $F$  have a positive probability, it's possible to rewrite the above condition in a different (illustrative!) way. *How?*

Now that we know the notion of **independence**, we can construct random variables in many creative ways.

**Example 3.2.** A fair coin is tossed repeatedly and **independently** until the first Heads. Let the random variable  $Y$  represent the total number of Tails observed by the end of the procedure.

*What is the support of the random variable  $Y$ ?*

*What is the **probability mass function** of the random variable  $Y$ ?*