
Name:

UTeid:

M378K Introduction to Mathematical Statistics

Fall 2024

University of Texas at Austin

In-Term Exam III

Instructor: Milica Čudina

Notes: This is a closed book and closed notes exam. The maximal score on the exam is 100 points.

Time: 50 minutes

All written work handed in by the student is considered to be
their own work, prepared without unauthorized assistance.

The University Code of Conduct

"The core values of The University of Texas at Austin are learning, discovery, freedom, leadership, individual opportunity, and responsibility. Each member of the university is expected to uphold these values through integrity, honesty, trust, fairness, and respect toward peers and community. As a student of The University of Texas at Austin, I shall abide by the core values of the University and uphold academic integrity."

"I agree that I have complied with the UT Honor Code during my completion of this exam."

Signature:

3.1. Formulas. If Y has the binomial distribution with parameters n and p , then $p_Y(k) = \mathbb{P}[Y = k] = \binom{n}{k} p^k (1-p)^{n-k}$, for $k = 0, \dots, n$, $\mathbb{E}[Y] = np$, $\text{Var}[Y] = np(1-p)$. The binomial coefficients are defined as follows for integers $0 \leq k \leq n$, $\binom{n}{k} = \frac{n!}{k!(n-k)!}$. The moment generating function of Y is given by $m_Y(t) = (pe^t + q)^n$.

If Y has a geometric distribution with parameter p , then $p_Y(k) = p(1-p)^k$ for $k = 0, 1, \dots$, $\mathbb{E}[Y] = \frac{1-p}{p}$, $\text{Var}[Y] = \frac{1-p}{p^2}$. Its mgf is $m_Y(t) = \frac{p}{1-qe^t}$ for t such that $qe^t < 1$.

If Y has a Poisson distribution with parameter λ , then $p_Y(k) = e^{-\lambda} \frac{\lambda^k}{k!}$ for $k = 0, 1, \dots$, $\mathbb{E}[Y] = \text{Var}[Y] = \lambda$. Its mgf is $m_Y(t) = e^{\lambda(e^t-1)}$.

If Y has a uniform distribution on $[l, r]$, its density is

$$f_Y(y) = \frac{1}{r-l} \mathbf{1}_{(l,r)}(y),$$

its mean is $\frac{l+r}{2}$, and its variance is $\frac{(r-l)^2}{12}$. Let $U \sim U(0, 1)$. The mgf of U is $m_U(t) = \frac{1}{t}(e^t - 1)$.

If Y has the standard normal distribution, then its mean is zero, its variance is one, and its density equals

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \quad x \in \mathbb{R}.$$

Its mgf is $m_Y(t) = e^{\frac{t^2}{2}}$.

If Y has the exponential distribution with parameter τ , then its cumulative distribution function is $F_Y(y) = 1 - e^{-\frac{y}{\tau}}$ for $y \geq 0$, its probability density function is $f_Y(y) = \frac{1}{\tau} e^{-y/\tau}$ for $y \geq 0$. Also, $\mathbb{E}[Y] = \text{SD}[Y] = \tau$. Its mgf is $m_Y(t) = \frac{1}{1-\tau t}$.

The mgf of $Y \sim \Gamma(k, \tau)$ is

$$m_Y(t) = \frac{1}{(1-\tau t)^k} \text{ for } t < 1/\tau.$$

Its expectation is $k\tau$ and its variance is $k\tau^2$. The χ^2 -distribution with n degrees of freedom is the special case $\Gamma(\frac{n}{2}, 2)$

3.2. DEFINITIONS.

Problem 3.1. (10 points) Write down the definition of the **bias** of an estimator $\hat{\theta}$ of a parameter θ .

Solution:

$$\text{bias}(\hat{\theta}) = \mathbb{E}[\hat{\theta}] - \theta$$

Problem 3.2. (10 points) Write down the definition of the **mean squared error** of an estimator $\hat{\theta}$ of a parameter θ .

Solution:

$$\text{MSE}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \theta)^2]$$

3.3. TRUE/FALSE QUESTIONS.

Problem 3.3. (5 points) Let Z be a standard normal random variable and let Q^2 have the χ^2 -distribution with $\nu \geq 2$ degrees of freedom. Assume that Z and Q^2 are independent. Set

$$T = \frac{Z}{\sqrt{\frac{Q^2}{\nu}}}.$$

Then, T has a t -distribution with ν degrees of freedom. *True or false? Why?*

Solution: TRUE

From the definition of the t -distribution, we know that

$$T = \frac{Z}{\sqrt{\frac{Q^2}{\nu}}}$$

has the t -distribution with ν degrees of freedom.

3.4. Free-response problems.

Please, explain carefully all your statements and assumptions. Numerical results or single-word answers without an explanation (even if they're correct) are worth 0 points.

Problem 3.4. (15 points) Let Y_1, Y_2, \dots, Y_n be a random sample from a population with a uniform distribution on $(\theta, \theta + 1)$. Then,

$$\hat{\theta}_n = \bar{Y}_n - \frac{1}{2}$$

is a **consistent estimator** for θ

Solution: We can use the same theorem we used in class, i.e., we can demonstrate that

- $\hat{\theta}_n$ is *unbiased*; and
- $\text{Var}[\hat{\theta}_n] \rightarrow 0$, as $n \rightarrow \infty$.

To show that $\hat{\theta}_n$ is unbiased, we must prove that

$$\mathbb{E}[\hat{\theta}_n] = \theta.$$

From the given definition of $\hat{\theta}_n$, we have

$$\mathbb{E}[\hat{\theta}_n] = \mathbb{E}\left[\bar{Y}_n - \frac{1}{2}\right] = \mathbb{E}[Y_1] - \frac{1}{2} = \theta + \frac{1}{2} - \frac{1}{2} = \theta.$$

Hence, $\hat{\theta}_n$ is, indeed, unbiased.

As for the other claim, we have that

$$\text{Var}[\hat{\theta}_n] = \text{Var}\left[\bar{Y}_n - \frac{1}{2}\right] = \text{Var}[\bar{Y}_n].$$

Since Y_1, \dots, Y_n is a random sample, the additive formula for the variance applies. So, we get

$$\text{Var}[\hat{\theta}_n] = \frac{\text{Var}[Y_1]}{n} = \frac{1}{12n}.$$

The above converges to 0 as $n \rightarrow \infty$ which concludes our proof.

Problem 3.5. (20 points) Let Y_1, Y_2, \dots, Y_n be a random sample from $E(\tau)$. Consider the following two estimators for τ :

$$\hat{\theta}_1 = \bar{Y} \quad \text{and} \quad \hat{\theta}_2 = nY_{(n)}$$

You know that $\hat{\theta}_2$ is unbiased and that $\text{MSE}(\hat{\theta}_2) = \tau^2$. Find the efficiency of $\hat{\theta}_1$ relative to $\hat{\theta}_2$.

Solution: We have shown in class (multiple times) that \bar{Y} is unbiased for the population mean. So, $\hat{\theta}_1$ is unbiased.

By definition, the relative efficiency we are looking for is

$$\text{eff}(\hat{\theta}_1, \hat{\theta}_2) = \frac{\text{Var}[\hat{\theta}_2]}{\text{Var}[\hat{\theta}_1]}.$$

We have shown in class that

$$\text{Var}[\hat{\theta}_1] = \text{Var}[\bar{Y}] = \frac{\tau^2}{n}.$$

From the given information, we know that

$$\text{Var}[\hat{\theta}_2] = \text{MSE}(\hat{\theta}_2) = \tau^2.$$

So, the relative efficiency is

$$\text{eff}(\hat{\theta}_1, \hat{\theta}_2) = \frac{\tau^2}{\frac{\tau^2}{n}} = n.$$

Problem 3.6. (20 points) Let Y_1, \dots, Y_n be a random sample from an exponential distribution with an unknown parameter τ . What is the maximum likelihood estimator for τ ? Make sure that you prove your claim!

Solution: Let y_1, \dots, y_n represent a set of observations of the above random sample. Then, the likelihood function is

$$L(\tau; y_1, \dots, y_n) = \prod_{i=1}^n \left(\frac{1}{\tau} e^{-\frac{y_i}{\tau}} \right) = \frac{1}{\tau^n} e^{-\frac{1}{\tau} \sum y_i}.$$

The log-likelihood function is

$$\ell(\tau; y_1, \dots, y_n) = -n \ln(\tau) - \frac{1}{\tau} \sum_{i=1}^n y_i.$$

Differentiating with respect to τ , we get

$$\ell'(\tau; y_1, \dots, y_n) = -\frac{n}{\tau} + \frac{\sum_{i=1}^n y_i}{\tau^2}.$$

Equating the above to 0 and solving for τ , we get

$$\hat{\tau}_{MLE} = \bar{Y}.$$

3.5. MULTIPLE CHOICE QUESTIONS.

Problem 3.7. (5 points) In a sample Y_1, \dots, Y_n from the uniform distribution on $(0, \theta)$ with parameter $\theta > 0$, $U = cY_{(n)}$ is a pivotal quantity if the value of the constant c is

- (a) $\frac{1}{2}$
- (b) $1/\theta$
- (c) θ
- (d) $\frac{n+1}{n}$
- (e) none of the above

Solution: The correct answer is **(b)**.

The pivotal quantity

$$U = \frac{Y_{(n)}}{\theta} = \max(Y_1, \dots, Y_n)/\theta$$

has the distribution with cdf y^n on $[0, 1]$.

Problem 3.8. (5 points) In a random sample of 100 voters 64 prefer candidate A and the rest prefer candidate B . The (approximate) $(1 - \alpha)$ -confidence interval for the parameter p (the population proportion of A voters) is of the form

$$[0.64 - z_{\alpha/2} \times c, 0.64 + z_{\alpha/2} \times c],$$

where $z_{\alpha/2} = \text{qnorm}(1 - \alpha/2, 0, 1)$.

The value of c is:

- (a) 0.016
- (b) 0.024
- (c) 0.036
- (d) 0.048
- (e) ≥ 0.05

Solution: The correct answer is **(d)** since $c = \sqrt{\hat{p}(1 - \hat{p})/n} = \sqrt{0.64 \times 0.36/100} = 0.048$.

Problem 3.9. (5 points) A sample of size $n = 2$ from normal distribution with unknown μ and σ is collected and the data are

$$y_1 = 1 \text{ and } y_2 = 5.$$

The right end-point of a symmetric 95% confidence interval for σ^2 is

- (a) $8/\text{qchisq}(0.975, 2)$
- (b) $16/\text{qchisq}(0.025, 1)$
- (c) $8/\text{qchisq}(0.025, 1)$
- (d) $16/\text{qchisq}(0.975, 2)$
- (e) **None of the above**

Solution: The correct answer is (c).

The confidence interval is based on the pivotal quantity $(n-1)S^2/\sigma^2$ whose distribution is $\chi^2(n-1)$. In this case, $n = 2$, $\bar{Y} = 3$ so that $(n-1)s^2 = (y_1 - \bar{y})^2 + (y_2 - \bar{y})^2 = 2^2 + 2^2 = 8$, which produces the interval $[8/\text{qchisq}(0.975, 1), 8/\text{qchisq}(0.025, 1)]$.

Problem 3.10. (5 points) Let Y_1, \dots, Y_5 be a random sample from the normal distribution $N(\mu, 2)$, with an unknown mean μ and the known standard deviation $\sigma = 2$. The collected data turn out to be

$$y_1 = 2, y_2 = 5, y_3 = 1, y_4 = 4, y_5 = 3.$$

The right end-point $\hat{\mu}_R$ of the symmetric 90%-confidence interval $[\hat{\mu}_L, \hat{\mu}_R]$ for μ is

- (a) $3 + \frac{2}{\sqrt{5}}\text{qnorm}(0.95, 0, 1)$.
- (b) $3 - \frac{2}{\sqrt{5}}\text{qnorm}(0.95, 0, 1)$.
- (c) $3 - \frac{1}{\sqrt{5}}\text{qt}(0.95, 4)$.
- (d) $3 + \frac{1}{5}\text{qnorm}(0.9, 5)$.
- (e) none of the above

Solution: The correct answer is (a).

The confidence interval in this case is based on the pivotal quantity $\sqrt{5} \left(\frac{\mu - \bar{Y}}{2} \right)$ which has the $N(0, 1)$ distribution. Therefore, for $a = \text{qnorm}(0.95, 0, 1)$ we have

$$\mathbb{P} \left[-a \leq \sqrt{5} \left(\frac{\mu - \bar{Y}}{2} \right) \leq a \right] = 0.90.$$

We solve for μ to obtain

$$\mathbb{P} \left[\bar{Y} - \frac{2}{\sqrt{5}}a \leq \mu \leq \bar{Y} + \frac{2}{\sqrt{5}}a \right] = 0.90.$$

For our data set $\bar{y} = 3$, so $\hat{\mu}_R = 3 + \frac{2}{\sqrt{5}}\text{qnorm}(0.95, 0, 1)$.