

UNIVERSITY OF TEXAS AT AUSTIN

Problem Set # 7

The Central Limit Theorem.

Let $\{X_n, n = 1, 2, 3, \dots\}$ be a sequence of independent, identically distributed random variables such that $\mu_X = \mathbb{E}[X_1] < \infty$ and $\text{Var}[X] = \sigma_X^2 < \infty$. For every $n = 1, 2, \dots$ define

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \quad \text{sample mean}$$

Problem 7.1. Find the expected value of \bar{X}_n for every n .

$$\begin{aligned} \mathbb{E}[\bar{X}_n] &= \mathbb{E}\left[\left(\frac{1}{n}\right)(X_1 + X_2 + \dots + X_n)\right] = \text{linearity} \\ &= \frac{1}{n} \mathbb{E}[X_1 + X_2 + \dots + X_n] = \text{linearity} \\ &= \frac{1}{n} (\mathbb{E}[X_1] + \mathbb{E}[X_2] + \dots + \mathbb{E}[X_n]) \quad \text{identically dist'd} \\ &= \frac{1}{n} \cdot n \cdot \mu_X = \mu_X \quad \text{accuracy} \end{aligned}$$

Problem 7.2. Find the variance and standard deviation of \bar{X}_n for every n .

$$\begin{aligned} \text{Var}[\bar{X}_n] &= \text{Var}\left[\left(\frac{1}{n}\right)(X_1 + X_2 + \dots + X_n)\right] \\ &= \frac{1}{n^2} \text{Var}[X_1 + X_2 + \dots + X_n] \quad \text{independence} \\ &= \frac{1}{n^2} (\text{Var}[X_1] + \text{Var}[X_2] + \dots + \text{Var}[X_n]) \quad \text{identically dist'd} \\ &= \frac{1}{n^2} \cdot n \cdot \text{Var}[X_1] = \frac{\sigma_X^2}{n} \\ \text{SD}[\bar{X}_n] &= \frac{\sigma_X}{\sqrt{n}} \quad \text{precision} \end{aligned}$$

Theorem 7.1. The Central Limit Theorem (CLT). If the above conditions are satisfied, we have that

$$\frac{\bar{X}_n - \mu_X}{\frac{\sigma_X}{\sqrt{n}}} \xrightarrow{D} N(0, 1) \quad \text{as } n \rightarrow \infty.$$

Practically, for "large enough" n , \bar{X}_n is approximately normal with mean μ_X and variance $\frac{\sigma_X^2}{n}$. The rule of thumb is that we use the theorem for $n \geq 30$. If that is the case, we have that for any real $a < b$,

$$\mathbb{P}[a < \bar{X}_n \leq b] = \mathbb{P}\left[\frac{a - \mu_X}{\frac{\sigma_X}{\sqrt{n}}} < \frac{\bar{X}_n - \mu_X}{\frac{\sigma_X}{\sqrt{n}}} \leq \frac{b - \mu_X}{\frac{\sigma_X}{\sqrt{n}}}\right] \approx \Phi\left(\frac{b - \mu_X}{\frac{\sigma_X}{\sqrt{n}}}\right) - \Phi\left(\frac{a - \mu_X}{\frac{\sigma_X}{\sqrt{n}}}\right).$$

Problem 7.3. Travel time by sled between Whoville and Whoburgh takes on average 36 minutes with a standard deviation of 6 minutes. Over a particular weekend, 64 sled trips take place. What is the (approximate) probability that the average sled trip took more than 38 minutes?

→: $n = 64 \geq 30 \checkmark$

$\bar{X}_n \approx \text{Normal}(\text{mean} = 36, \text{sd} = \frac{6}{\sqrt{64}} = \frac{3}{4})$

CLT

$\mathbb{P}[\bar{X}_n \geq 38] = 1 - \text{pnorm}(38, \text{mean} = 36, \text{sd} = 0.75) = 0.003830381$ \square

Problem 7.4. The amount of time your friendly taquero at *Torchy's Tacos* spends to assemble any one tasty taco is a random variable with mean 3 minutes and 15 seconds and standard deviation of thirty seconds. You and your 31 friends from *Applied Statistics* celebrate by ordering two tacos each. What is the probability that the average taco-assembly time is:

- less than 2 minutes and 30 seconds;
- more than 3 minutes and 15 seconds; *symmetry $\Rightarrow 0.5$*
- at least 3 minutes but at most 3 minutes and 30 seconds?

$n = 64 \geq 30 \checkmark : \bar{X}_n \approx \text{Normal}(\text{mean} = 3.25, \text{sd} = \frac{0.5}{\sqrt{64}} = 0.0625)$

$\mathbb{P}[\bar{X}_n \leq 2.5] = \text{pnorm}(2.5, \text{mean} = 3.25, \text{sd} = 0.0625) = 1.776482 \cdot 10^{-33}$

$\mathbb{P}[3 < \bar{X}_n < 3.5] = \text{pnorm}(3.5, 3.25, 0.0625) - \text{pnorm}(3, 3.25, 0.0625)$
 $= 0.9999367$