H33	AD:	Nov	embe	4 4th 6	2024.						
Stan	dard	N	oma	l Dis	tribut	ion.					
We	sa	y t	hat	a ra	ndom	vania	de Z	has	the		
						ormal					
4	its	Pro	bab	ility	densi	ty fun	chion	(pdf)) has	this	form
			£	"(之) =	φ(≈)	ty fun	e 2	J	for al	I zel	5
		•			•	symmet					
							φ(Z):	= 41-	E), i.	e., eve	?n
	/					•	mean/	medi	an/i	node= 1	0
		0'			7						
The	CION	0	tie	dist	huti	on func	tion (J +4	o std	namel
JILE.						, , , , , , , , , , , , , , , , , , ,			4 14	e siu	
			N	z) = (I)(z):	=P[Z	{ 2]	*		u 2	
					5	- η <u>L</u> L - ω <u>+</u>	1) du =	·	<u>1</u> 2₹7	- Z du	
						-∞ T₄	1 0.	- <u></u>	£	>	
-0							lo And	elync	rorm		
		_				tables	-				
he c	an c	u se	74	out	LN	Pnorm	and			in R	•
						N	wnite	/			
						10.					

The Normal Distributions We completely specify any normal distribution by providing its mean H_X and variance σ_X^2 (or its std deviation σ_X^2 . we write: $\times \sim Normal (mean = \mu_{\times}, variance = \sigma_{\times}^{2})$ X can be written as a linear transform of a standard normal Z: $X = H_X + \sigma_X \cdot Z$ can check:

• $E[X] = E[H_X + \sigma_X \cdot Z] = H_X + \sigma_X \cdot E[Z] = M_X$ We can check: · Var[x] = Var[Hx + ox Z] a deterministic shift which = $Var[O_X] = O_X^2$ $Var[Z] = O_X^2$ doesn't affect the variance I