

appropriate term.^{657,658} Poor glycemic control has been suggested, although not universally accepted, as a KLA risk factor, given impaired neutrophil phagocytosis of K1/K2 capsular serotypes.⁶⁵⁹ Strains isolated from patients with invasive syndrome have distinct virulence features compared with less invasive, comparator strains.⁶⁶⁰ These factors have genetic similarity with invasive disease isolates despite geographic diversity.⁶⁶¹ Capsular type K1 has emerged as an important factor in KLA and demonstrates a distinct link to a specific clonal complex, CC23, that involves a virulence plasmid; several of the factors discussed later portend higher community acquisition rates, KLA formation, and metastatic spread.^{627,661–663}

The K1/2 serotypes demonstrate high rates of hypermucoviscous phenotype,⁶⁵⁰ but hyperviscous phenotype is not limited to only the K1/2 serotypes. Presence of *magA* (mucoviscosity-associated gene), ubiquitous in K1 isolates, was initially proposed as a risk factor for both KLA and metastatic infection,⁶⁶⁴ but it appears to be associated with K1 capsule and not a driver of virulence.^{650,665,666} The regulator of mucoid phenotype (*rmpA*) gene increases mucoid phenotype for most isolates.^{634,650,667–670}

K. pneumoniae can produce a variety of fimbrial types, including types 1 and 3 pili, aiding host cell adherence.⁶⁷¹ Aerobactin, a siderophore, increases *K. pneumoniae* virulence,^{651,672–674} with increased production in hypervirulent phenotypes supporting its role in invasive disease.^{675,676} Much like *E. coli*, KLA isolates may use several iron acquisition systems, of which a majority are regulated by *tonB*.⁶⁷³

All strains of *K. pneumoniae* are resistant to ampicillin as a result of the presence of a chromosomal gene encoding a penicillin-specific β -lactamase.⁶⁷⁷ Preceding ampicillin use has been shown to increase risk of KLA⁶³³ and postneurosurgical infections.⁶²² In addition, nosocomial isolates are frequently resistant to numerous other antibiotics as a result of the acquisition of MDR plasmids. For example, *K. pneumoniae* is one of the most common organisms to carry plasmids encoding ESBLs and carbapenemases, and bacteremia with such strains is associated with higher rates of treatment failure and death.^{678,679} After the initial description of the novel carbapenemase, KPC-1, in a *Klebsiella* isolate in the United States, *Klebsiella* is an archetype of emerging resistance.⁶⁸⁰ Therapeutic options for infections caused by non-MDR strains include first-generation cephalosporins, penicillin/ β -lactamase inhibitor combinations, trimethoprim-sulfamethoxazole, fluoroquinolones, and aminoglycosides. For MDR strains, especially those expressing ESBLs, treatment options are often limited to fourth-generation cephalosporins or carbapenems. Carbapenemase-producing strains leave scarce options.⁷

The *K. pneumoniae* subspecies *rhinoscleromatis* is the causative agent of respiratory scleroma, also known as rhinoscleroma, a chronic granulomatous infection of the nasal passages and other parts of the respiratory tract.^{627,681} In contrast to *K. pneumoniae*, *K. rhinoscleromatis* is not found outside human hosts.⁶⁸² The disease it causes is characterized by chronic inflammation and remains difficult to diagnose because of reliance on biopsy, lack of early symptom specificity, and variation of routine biochemical properties between strains.⁶⁸³ Often, rhinoscleroma presents in advanced stages with disfigurement and respiratory compromise following a progressive clinical pathway from atrophic through granulomatous to fibrotic stages.⁶⁸⁴ The disease is found primarily in impoverished areas of Central and South America, Africa, and Asia or in immigrants from these areas^{681,682} and is transmitted by close contact and affected by genetic control of the host immune responses.⁶⁸² Respiratory scleroma is characterized by nodules and masses often involving the nasal passages and, less commonly, other areas of the respiratory tract. Occasionally, the infection can erode through bone and invade the CNS. The differential diagnosis includes tuberculosis, leprosy, fungal infections, granulomatosis with polyangiitis, malignancies, and sarcoidosis. Biopsy specimens show granulomatous inflammation with foamy macrophages (Mikulicz cells) containing intracellular organisms (Fig. 218.6).⁶⁸⁵ The bacteria can be grown on ordinary laboratory media. Prolonged antibiotic therapy, often for months, and, occasionally, surgical débridement are required for cure, but relapses are common.⁶⁸² Traditionally, streptomycin or tetracycline has been used, but more recent reports suggest that trimethoprim-sulfamethoxazole or fluoroquinolones could be alternatives.⁶⁸²

The *K. pneumoniae* subspecies *ozaenae* can colonize the nasopharynx of healthy individuals. This subspecies has been associated with chronic




FIG. 218.6 Photomicrograph of Mikulicz cells in a patient with rhinoscleroma. The periodic acid-Schiff–positive structures within the macrophage (arrows) are the causative organism, *Klebsiella rhinoscleromatis*. (From ID Images. © The Infectious Diseases Society of America. Case presented by Danielle Osterholzer, Tom E. Davis, and Stephen D. Allen.)

atrophic rhinitis (*ozena* means “stench”) marked by bone resorption and thick, foul-smelling discharge that often forms crusts, but its etiologic role is controversial. The organism has only rarely been isolated from patients with a variety of infections, including otitis media and mastoiditis, cystitis and pyelonephritis, soft tissue infections, bacteremia associated with neutropenia, pneumonia, and meningitis.^{686,687} It is usually susceptible to multiple antibiotics except when isolated from hospitalized patients who have received previous antimicrobial therapy.

K. oxytoca, like *K. pneumoniae*, can cause a variety of nosocomial infections and is distinguished from *K. pneumoniae* based on its ability to produce indole from tryptophan. *K. oxytoca* is rarely isolated in stool specimens of asymptomatic patients and is not believed to be a constituent of the normal bowel microbiota.^{688–690} There is evidence that *K. oxytoca* has increased cytotoxin production not found in other *Klebsiella* species, which may account for its ability to induce antibiotic-associated hemorrhagic colitis with histopathologic findings similar to those of STEC or *Shigella*.^{691,692} It may also cause outbreaks and a variety of nosocomial infections and may be resistant to multiple antibiotics.

K. granulomatis (formerly *Calymmatobacterium granulomatis*) is a fastidious member of the genus that causes granuloma inguinale or donovanosis, a chronic genital ulcerative disease (see Chapter 235).

Enterobacter Species, Pantoea agglomerans, and Chronobacter sakazakii

Enterobacter species have undergone reclassification. *Enterobacter cloacae* is responsible for most *Enterobacter* infections in humans. *Enterobacter aerogenes* was renamed *Klebsiella aerogenes* owing to genetic similarity.^{692a,692b} *Enterobacter sakazakii* was reclassified into the genus *Chronobacter*, which includes at least seven distinct species.⁶⁹³ Several less common *Enterobacter* spp. have been reorganized into the new genera of *Lelliottia*, *Pluralibacter*, and *Kosakonia*.⁶⁹⁴ *P. agglomerans* (formerly *Enterobacter agglomerans*) is also a common isolate and is grouped with the *Enterobacter* spp. here. These bacteria ferment lactose, are motile, and form mucoid colonies. *Enterobacter* strains are part of the intestinal microbiota and rose to notoriety with a large nationwide outbreak⁶⁹⁵ and continue to be implicated in clonal outbreaks.⁶⁹⁶ The normal colonization may become infection, especially in patients who have received antimicrobial therapy or who are admitted to the ICU.⁶⁹⁷ *Enterobacter* spp. may cause a wide variety of nosocomial infections,^{612,698,699} including pneumonia, UTIs, wound and burn infections, and infections of intravascular and other prosthetic devices. Other than neonates and postneurosurgical patients, *Enterobacter* rarely causes meningitis. There do not seem to be distinguishing characteristics among infections caused by *E. cloacae* and *E. aerogenes*, although *E. cloacae* may have higher

rates of antimicrobial resistance, and *E. aerogenes* may have poorer outcomes.⁷⁰⁰

Given the considerable difficulty in correctly identifying *Chronobacter* spp., not all reports of *E. sakazakii* prior to reclassification in 2007 can be directly attributed to *Chronobacter*.⁶⁹⁴ Given those limitations, we know that *Chronobacter* may cause bacteremia, necrotizing enterocolitis, and meningitis, primarily in neonates of low birth weight, and in such infants is associated with consumption of powdered milk formula, with mortality rates in excess of 40%.^{701–705}

P. agglomerans, a plant pathogen, is often associated with contaminated catheters or penetrating trauma.⁷⁰⁸

Enterobacter, *Pantoea*, and *Chronobacter* use virulence factors typical of other Enterobacteriaceae.^{709–712} *Enterobacter* strains associated with illness may deploy a variety of mechanisms, including heavy metal resistance and additional siderophore assembly kits to gain a fitness advantage against other microbes, that facilitate their survival in diverse environments.⁷¹¹ Efflux pumps are very active in the wide array of antimicrobial resistance in *Enterobacter* but may also help explain *E. cloacae* competitive virulence.⁷¹² *C. sakazakii* can adhere to and invade brain microvascular endothelial cells, similar to NMEC.^{713–715} The outer membrane of *C. sakazakii* is encoded by *ompA*, its best-described virulence marker.⁷¹⁶ *C. sakazakii* resists desiccation, which may explain its role in contaminating powdered milk, where it can survive for more than 12 months.^{703,717–721} *C. sakazakii* is the only *Chronobacter* species that can metabolize sialic acid, which is found in breast milk, infant formula, the gastrointestinal tract, and the brain, possibly explaining the specific association with these syndromes.⁷²²

Strains belonging to the genus *Enterobacter* often show antimicrobial resistance, earning them a place in the ESKAPE mnemonic.^{723,724} *E. cloacae*, *E. aerogenes*, and most strains of *Chronobacter* are intrinsically resistant to ampicillin and first- and second-generation cephalosporins as a result of an inducible AmpC chromosomal β -lactamase that is controlled by both positive and negative regulators.⁷²⁵ Furthermore, mutants that constitutively produce high levels of β -lactamase, conferring resistance to third-generation cephalosporins, arise at frequencies of 10^{-4} to 10^{-7} , often as a result of mutations in the regulatory loci, such that resistant mutants may already be constitutively present before initiation of therapy. As with other members of the Enterobacteriaceae, *Enterobacter* spp. may carry plasmids encoding MDR phenotypes.^{698,726,727} (see Chapter 18 on antimicrobial resistance). Therapy must therefore be tailored to local resistance patterns and results of individual isolate susceptibility testing. Clinicians must be aware that emergence of stably derepressed resistant mutants may lead to treatment failure when third-generation cephalosporins are used, even if isolates appear susceptible at initial testing.⁷²⁸ Therefore, fourth-generation cephalosporins and carbapenems may be empirically chosen for severe infections.

Serratia Species

Of the many species in the genus *Serratia*, *Serratia marcescens* is the one most commonly isolated from human infections, and *Serratia liquefaciens* is occasionally grown. *Serratia* strains are motile, rarely ferment lactose, and produce an extracellular DNase. The organism is widespread in the environment but not a commonly recognized component of the human fecal microbiota; thus most infections appear to be acquired exogenously. Many environmental and some clinical strains of *S. marcescens* produce a red pigment (Fig. 218.7), prodigiosin. Bartolemeo Bizio, an Italian pharmacist, first described the organism in 1819 as the cause of red discoloration of polenta, thereby discrediting the claim that the growth was due to the miraculous appearance of blood.⁷²⁹ He named the bacterial genus to honor Serafino Serrati, an Italian physicist, and its species name (*marcescens*) for the Latin word for “to decay” because of the tendency of the pigment to change color as the colonies age.⁷³⁰ The production of prodigiosin and the misconception that the bacterium was harmless led to its frequent use as a biologic marker to study, among other things, the transmission of bacteria through speech and contact, ascending bladder colonization from urinary catheters, and the dissemination of aerosolized bacteria after experimental release into the environment in models of biologic warfare.^{731,732} It is now appreciated that *S. marcescens* can cause a wide variety of infections, ranging from UTIs, bacteremia, pneumonia, and CNS infections, to



FIG. 218.7 *Serratia marcescens* colonies producing prodigiosin responsible for characteristic red pigmentation. (From Mahlen SD. *Serratia* infections: from military experiments to current practice. Clin Microbiol Rev. 2011;24:755–791.)

other less common infections, including ocular infections. The most common site of infection is the urinary tract, but the organism is frequently isolated from the respiratory tract and wounds.⁷³²

In large epidemiologic surveys, *Serratia* species account for only a small percentage of pneumonia and bacteremia. *Serratia* is typically considered a nosocomial pathogen, but in a population-based study in Australia conducted over 10 years, nearly half of the cases of *Serratia* bacteremia was found to be of community onset.⁷³³ A similar laboratory-based surveillance study from Canada found that nearly two-thirds of incident isolates were of community onset; however, the investigators were unable to distinguish community-acquired from health care-associated isolates.⁷³⁴ Such data suggest that community infections may play a larger role than previously thought, and that infections may arise from endogenous (microbiota) strains, in addition to being transmitted nosocomially.

There have been more than 200 reported *S. marcescens* outbreaks⁷³² implicating sources such as compounded drugs,⁷³⁵ prefilled syringes,⁷³⁶ contaminated soaps and water,^{737–739} medical devices,⁷⁴⁰ and total parenteral nutrition.^{741,742} *Serratia* outbreaks are of particular concern in pediatric ICUs.⁷³⁷ *S. liquefaciens* has been noted to be a cause of transfusion-related outbreaks.^{743,744}

In addition, *Serratia* species have a historical association with infections in injection drug users who are at particular risk for *S. marcescens* infections, including endocarditis^{731,745} and infections at other sites after hematogenous spread.⁷³¹ *Serratia* spp. have a predilection for ocular infection, accounting for up to 20% of nosocomial *Serratia* infections, second only to respiratory infections in a retrospective review.⁷⁴⁶ *Serratia* can cause keratitis, corneal ulceration, and endophthalmitis, which may follow hematogenous dissemination, and is second only to *Pseudomonas* spp. in the frequency of gram-negative ocular infections and rivals it with regard to infections associated with contact lens wear.^{747–749}

Potential virulence factors of *Serratia* have not received as much attention as those of *E. coli* or *Klebsiella*, but *Serratia* likely uses many of the common virulence factors present in other Enterobacteriaceae. The ability to degrade macrophages has been proposed as a possible virulence attribute for *S. liquefaciens*.⁷⁵⁰ This process is independent of internalization, but because proximity and contact are required, extracellular toxins or other bacterial effectors may be implicated. More recently, the RssAB-FlhDC-ShlA pathway has been considered as a complex host invasive pathogenic pathway involving numerous factors, including quorum sensing and swarming.^{752,754} Through this pathway, ShlA both exerts cytotoxic properties and contributes to urinary pathogenesis. ShlA, codependent on its activator, ShlB, requires cellular contact consistent with the effects on macrophages noted earlier.⁷⁵² RssAB is a regulatory system that decreases ShlA production and thereby swarming.⁷⁵³ FlhDC is important for quorum sensing and positively regulates

swarming, but negatively affects biofilm formation.^{754,755} *Serratia* colonization, invasiveness, and persistence may be aided by biofilm formation⁷⁵⁶ and are likely, in part, regulated by quorum sensing.^{732,757} However, biofilm formation was not evident in *Serratia* keratitis infections; accordingly, exoenzyme production may be important.⁷⁵⁸

Serratia isolates are resistant to ampicillin and first-generation cephalosporins because of an inducible, chromosomal AmpC β -lactamase similar to that of *Enterobacter*.^{759,760} *Serratia* have been associated with nearly all mechanisms of β -lactamase production seen in other gram-negative organisms,⁷⁶¹ including KPC enzymes conferring resistance to carbapenems (see Chapter 18 on antimicrobial resistance). Given the variety of resistance mechanisms found, treatment of infections caused by *S. marcescens* can be difficult, and every effort should be made to identify the sources of outbreaks to control the spread of the organism in susceptible populations.

Citrobacter Species

Members of the genus *Citrobacter* are named for their use of citrate as their sole carbon source. Of the more than 10 recognized species, *Citrobacter freundii* and *Citrobacter koseri* (formerly *Citrobacter diversus*) are the most important human pathogens. They are differentiated by their ability to convert tryptophan to indole, ferment lactose, and use malonate.⁷⁶² *C. freundii* produces hydrogen sulfide and therefore can be confused with *Salmonella*, with which it was classified at one time. The urinary tract is the most frequent site from which *Citrobacter* is cultured, often in association with an indwelling catheter, and *Citrobacter* may be a frequent cause of nosocomial UTIs⁷⁶³ and outbreaks.⁷⁶⁴ These bacteria may also be seen in intraabdominal and pulmonary infections.^{765,766} Bacteremia from *C. freundii* may have an intraabdominal source more commonly than from the urinary tract, especially in patients with polymicrobial bacteremia and those with underlying malignancy.^{765,768} Neonatal meningitis due to *C. koseri* often (approximately 75%) causes multiple brain abscesses associated with high mortality and severe neurologic sequelae in survivors.⁷⁶⁹ The predilection for CNS infection has been postulated to be due to a combination of a unique outer membrane in *C. koseri* and infiltration of macrophages infected with *C. koseri* resistant to phagolysosomal effects across the BBB.^{769,770} Nosocomial outbreaks have been accompanied by high rates of intestinal colonization by the organism in infants and by carriage of the bacteria on the hands of health care workers.^{764,771} A carbapenemase-expressing *C. freundii* outbreak was considered to have a potential foodborne source.⁷⁷²

C. freundii strains, like strains of *Enterobacter* and *Serratia*, have inducible AmpC genes encoding resistance to ampicillin and first-generation cephalosporins that can be produced constitutively at high levels after mutations selected by prior exposure. Furthermore, isolates of *Citrobacter* may be resistant to multiple other antibiotics as a result of plasmid-encoded resistance genes. The exposure of commensal *Citrobacter* spp. of low virulence to repeated antibiotic courses may select for accumulation of multiple resistance genes, leading to extensive resistance profiles.⁷⁷³

Hafnia alvei

H. alvei (formerly *Enterobacter hafniae*) was previously considered to be the sole species in the genus *Hafnia*. Through more recent 16S rRNA gene studies, a novel species, *H. paralvei*, now has been confirmed.⁷⁷⁴ There are not clear virulence differences between the species, although *H. alvei* may be more cytotoxic.⁷⁷⁵ As with most of the disease-implicated members of the Enterobacteriaceae, *Hafnia* reside in the gastrointestinal tract of humans and many animal species as part of the normal microbiota.⁷⁷⁶ *Hafnia* are motile but do not ferment lactose. Although *H. alvei* may be cultured from various sites, it is rarely implicated in monomicrobial infections⁷⁷⁷; this finding suggests that its pathogenicity may be quite low, with the exception of rare case reports of isolated bacteremia. Most infections with *H. alvei* occur in patients with severe underlying illness, including malignancy, trauma, and recent surgery. There have been numerous reports linking diarrhea to the isolation of *H. alvei* from stool specimens and several studies have implicated *H. alvei* in outbreaks, although the evidence is not strong.⁷⁷⁶ Furthermore, those strains initially reported to produce A/E lesions similar to EPEC (as described earlier) were reclassified as *E. albertii*.⁹⁴ However, in a

case-control study the prevalence in stools of *H. alvei* was significantly greater in those with diarrhea than without, consistent with a possible etiologic role.⁷⁷⁸ These strains, though, were all negative for the *eae* gene encoding intimin, found in EPEC and *E. albertii*.⁹⁴ A majority of *Hafnia* strains tested in vitro produced a cytolytic toxin that has been implicated in gastroenteritis.⁷⁷⁵ *H. alvei* are naturally resistant or of intermediate susceptibility to tetracyclines, amoxicillin, amoxicillin-clavulanate, ampicillin-sulbactam, narrow-spectrum cephalosporins, azithromycin, and fosfomycin.⁷⁷⁹ Treatment of serious *H. alvei* infections is guided by antimicrobial susceptibility testing. High rates of colistin resistance have been described, given the close phylogenetic relationship with other intrinsically colistin-resistant Enterobacteriaceae.⁷⁸⁰ Multiple resistance patterns have been noted.^{779,781}

Proteus, Providencia, and Morganella

The genera *Proteus*, *Providencia*, and *Morganella* are related members of the Enterobacteriaceae that are lactose negative and motile and produce phenylalanine deaminase. *Proteus mirabilis* and *Proteus vulgaris* account for most clinical *Proteus* isolates. Both produce urease and hydrogen sulfide, and the latter is indole positive. In contrast to most Enterobacteriaceae, these bacteria may express large numbers of fimbriae and flagella, transforming them into highly elongated rods that translocate rapidly across the surface of agar plates, resulting in a characteristic “swarming” motility. This motility produces a characteristic pattern on growth plates (Fig. 218.8) from sequential rounds of swarm cell differentiation and can overwhelm accompanying organisms. This swarming motility is regulated through a complex network acting on the flagellar transcription regulator, *flhDC*.⁷⁸² The name *Proteus* follows from the character in Homer’s *Odyssey* who is capable of changing form.

Providencia stuartii is the most common species of its genus isolated from clinical specimens, but *Providencia rettgeri* is occasionally grown. These bacteria can be differentiated from *Proteus* and *Morganella* based on their ability to use citrate and ferment D-mannitol.⁷⁸³ *Morganella morganii* is, at present, the only member of its genus and is citrate negative.

Proteus spp. are common causes of UTIs, occasionally in normal hosts⁷⁸⁴ but more often in those with indwelling catheters or anatomic or functional urinary tract abnormalities.^{785,786} *Proteus* are common among the gram-negative bloodstream isolates, with most secondary to UTI and often associated with urinary catheters.^{639,787–789}




FIG. 218.8 Characteristic bull's eye pattern of sequential rounds of swarming and consolidation frequently seen with *Proteus*. (From Schaffer JN, Pearson MM. *Proteus mirabilis* and urinary tract infections. Microbiol Spectr. 2015;3[5].)

Community-acquired *Proteus* UTIs in the presence of hydronephrosis or urolithiasis have increased risk of bacteremia.⁷⁹⁰ *Proteus* has a remarkable ability to persist in the urinary tract despite antibiotics and catheter exchange, likely due to immune evasion and the protective reservoir that urinary stones provide. Infections not related to the urinary tract or bloodstream have been reported with much less frequency.

The pathogenesis of UTIs caused by *P. mirabilis* has received considerable attention.^{782,791} These microorganisms have multiple features devoted to adherence and motility⁷⁹² and may produce several types of pili, the most important of which are mannose-resistant *Proteus*-like (MR/P) fimbriae,^{793,794} which have been shown to contribute to bladder colonization in a murine model.⁷⁹⁵ Ascending UTIs are common, and fimbriae and adhesins are necessary, but not sufficient, to produce infection.⁷⁸² Potent urease production contributes to colonization, stone formation, and obstruction,^{786,796–798} without which the infectivity of *Proteus* is diminished.^{799,800} By hydrolyzing urea to form CO₂ and ammonia, urease leads to the alkalization of urine, which leads to the precipitation of struvite, formation of calculi, and subsequent urinary catheter obstruction. Urinary pH ≥7 suggests infection with a urease-positive bacterium, most often a *Proteus* species. *P. mirabilis* can cause monomicrobial UTIs but is frequently found in polymicrobial catheter-associated urinary tract infections. In coinfection models with *P. stuartii*, urolithiasis and bacteremia are promoted in a urease-dependent fashion, and the presence of the other uropathogens regardless of their own urease production enhances the urease activity of *Proteus*.^{801–803}

Given the increased severity of *Proteus* UTIs and the propensity of this organism to cause bacteremia in community-acquired infections and patients with hydronephrosis or urolithiasis, radiographic studies such as renal sonography or computed tomography should be considered in patients with severe UTIs, especially those complicated by bacteremia.⁷⁹⁰ Obstruction due to calculi must be promptly relieved to avert severe clinical outcomes and to avoid persistent sources that can lead to recurrent infection.⁷⁸⁵ Most *Proteus* strains are susceptible to commonly used antibiotics, except for nitrofurantoin and tetracycline.⁷⁸³ Like other members of Enterobacteriaceae, MDR strains exist and are increasing; strains of *P. vulgaris* are generally more resistant.

P. stuartii and *P. rettgeri* are relatively uncommon clinical isolates, except in elderly patients with long-term indwelling urinary catheters,^{797,804,805} and are emerging in hospital settings.⁸⁰⁶ These infections are sometimes complicated by bacteremia and may cause death.^{804,807} *Providencia* uses many of the same virulence factors as *Proteus*, although they are less often studied. Biofilm production, weak urease production, and urease tolerance are proposed virulence mechanisms^{797,808} and have been implicated in the organism's ability to disseminate in a hospitalized setting.⁸⁰⁹ *P. stuartii* and *P. rettgeri* are often resistant to multiple antibiotics, including gentamicin, first-generation cephalosporins, and ampicillin,⁸¹⁰ although MDR phenotypes have been described.^{811–814} Therapy is guided by susceptibility testing.

M. morganii is an infrequent nosocomial isolate, usually isolated from urine or wounds.^{783,815} *M. morganii* bacteremia most often is due to a urinary or hepatobiliary port of entry and is often community acquired.^{816,817} *Morganella* spp. also may be present in polymicrobial infections and in wounds, which are the next most common source in bacteremic cases.⁸¹⁶ As with other members of this group, *M. morganii* may cause nosocomial outbreaks.⁸¹⁸ *M. morganii* strains possess inducible AmpC β-lactamases and therefore are intrinsically resistant to ampicillin

and first-generation cephalosporins; spontaneous derepressed mutants resistant to ESBLs similar to those described previously for *Enterobacter* spp. may arise.⁸¹⁹

Other Genera

The majority of clinically relevant Enterobacteriaceae are discussed in the previous sections, although as sequencing studies and in-depth biochemical analyses increase, the number of distinct genera within the Enterobacteriaceae continues to expand. Descriptions of clinical disease limited to case-level data include *Averyella*, *Buttiauxella*, *Cedecea*, *Kosakonia*, *Leclercia*, *Leminorella*, *Moellerella*, *Pluralibacter*, *Rahnella*, *Tatumella*, and *Yokenella*, in addition to those highlighted here.⁶⁹⁴

Edwardsiella tarda is found in freshwater environments, has been associated with diarrhea, and can cause wound infections, abscesses, and bacteremia, sometimes linked to marine exposure, including in patients "noodling" or hand-fishing of catfish.^{820,821} In patients with liver disease and iron overload, the mortality rates from *Edwardsiella* infections are high.⁸²² Genes related to iron scavenging have shown adaptive evolution, which may facilitate virulence in *Edwardsiella* to a broad range of hosts.⁸²³

Plesiomonas shigelloides is another organism that is found in water and has been associated with diarrhea and rarely with extraintestinal infections.⁸²⁴ *P. shigelloides* was previously grouped with vibrios, given their shared oxidase production and polar flagellae, but has been reassigned based on phylogenetic studies based on ribosomal RNA gene sequencing.^{825,826} *P. shigelloides* is an infrequent isolate from patients with gastroenteritis. Although the pathogenicity of *P. shigelloides* was questioned initially, it is now an accepted enteric pathogen based on case-control studies and numerous diarrheal illness outbreaks, with strong epidemiologic evidence.⁸²⁴ *P. shigelloides*-associated diarrhea manifests most commonly with secretory gastroenteritis but can also cause grossly bloody stool or chronic diarrheal illness. Risk factors include recently ingested raw shellfish or travel exposure.⁸²⁷

Ewingella americana, named after William Ewing, who made many contributions to our understanding of the microbiology of the Enterobacteriaceae, is a rare cause of nosocomial bacteremia, peritonitis associated with peritoneal dialysis, and conjunctivitis.^{828–830} Most isolates are highly sensitive to antibiotics, although multidrug resistance has been reported.⁸³¹

Infections caused by organisms belonging to the genus *Kluyvera*, which closely resembles *E. coli*, are rare. These bacteria have been recovered from urine, sputum, and wounds; and in many cases, the pathologic significance of their presence is unclear. However, pyelonephritis, bacteremia, and abdominal and soft tissue infections caused by *Kluyvera* spp. have occurred, particularly in immunosuppressed patients, with some infections being fatal.^{832–834} Chromosomal genes in *Kluyvera* spp. are likely the origin of CTX-M ESBLs, which have become an increasing cause of MDR and have demonstrated global spread.^{835–837} Although bacteria belonging to the genus *Photobacterium* are fascinating bioluminescent nematode symbionts, the nonluminescent species *Photobacterium asymbiotica* is the only one recovered in humans.^{694,838,839}

ACKNOWLEDGMENT

The authors thank Dr. Michael Donnenberg for his many contributions to the field, and his work on this chapter in the 6th, 7th, and 8th editions of this book.

Key References

The complete reference list is available online at Expert Consult.

- Doi Y, Bonomo RA, Hooper DC, et al. Gram-Negative Committee of the Antibacterial Resistance Leadership Group (ARLG).a. Gram-negative bacterial infections: research priorities, accomplishments, and future directions of the Antibacterial Resistance Leadership Group. *Clin Infect Dis*. 2017;64(suppl_1):S30–S35, Review.
- Peleg AY, Hooper DC. Hospital-acquired infections due to gram-negative bacteria. *N Engl J Med*. 2010;362:1804–1813.
- Hayashi F, Smith KD, Ozinsky A, et al. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature*. 2001;410:1099–1103.
- Welch RA, Burland V, Plunkett G 3rd, et al. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc Natl Acad Sci USA*. 2002;99:17020–17024.
- Kuehn MJ, Heuser J, Normark S, et al. Pili in uropathogenic *E. coli* are composite fibres with distinct fibrillar adhesive tips. *Nature*. 1992;356:252–255.
- Merz AJ, So M, Sheetz MP. Pilus retraction powers bacterial twitching motility. *Nature*. 2000;407:98–102.
- Connell H, Agace W, Klemm P, et al. Type 1 fimbrial expression enhances *Escherichia coli* virulence for the urinary tract. *Proc Natl Acad Sci USA*. 1996;93:9827–9832.
- Luo Y, Frey EA, Pfuetzner RA, et al. Crystal structure of enteropathogenic *Escherichia coli* intimin-receptor complex. *Nature*. 2000;405:1073–1077.
- Kenny B, DeVinney R, Stein M, et al. Enteropathogenic *E. coli* (EPEC) transfers its receptor for intimate adherence into mammalian cells. *Cell*. 1997;91:511–520.
- Koronakis V, Sharff A, Koronakis E, et al. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature*. 2000;405:914–919.
- Stebbins CE, Galán JE. Structural mimicry in bacterial virulence. *Nature*. 2001;412:701–705.
- Agnese DM, Calvano JE, Hahn SJ, et al. Human Toll-like receptor 4 mutations but not CD14 polymorphisms are associated with an increased risk of gram-negative infections. *J Infect Dis*. 2002;186:1522–1525.
- Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol*. 2004;2:123–140, Review.

96. Hyma KE, Lacher DW, Nelson AM, et al. Evolutionary genetics of a new pathogenic *Escherichia* species: *Escherichia albertii* and related *Shigella boydii* strains. *J Bacteriol*. 2005;187:619–628.
98. Rasko DA, Rosovitz MJ, Myers GS, et al. The pangenome structure of *Escherichia coli*: comparative genomic analysis of *E. coli* commensal and pathogenic isolates. *J Bacteriol*. 2008;190:6881–6893.
100. Croxen MA, Law RJ, Scholz R, et al. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev*. 2013;26:822–880. Review.
101. McDaniel TK, Jarvis KG, Donnenberg MS, et al. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc Natl Acad Sci USA*. 1995;92:1664–1668.
105. Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev*. 1998;11:142–201.
110. Rasko DA, Webster DR, Sahl JW, et al. Origins of the *E. coli* strain causing an outbreak of hemolytic-uremic syndrome in Germany. *N Engl J Med*. 2011;365:709–717.
149. Freedman SB, Xie J, Neufeld MS, et al; Alberta Provincial Pediatric Enteric Infection Team (APPETITE), Nettel-Aguirre A, Chuck A, Lee B, Johnson D, Currie G, Talbot J, Jiang J, Dickinson J, Kellner J, MacDonald J, Svenson L, Chui L, Louie M, Lavoie M, Eltorki M, Vanderkooi O, Tellier R, Ali S, Drews S, Graham T, Pang XL. Shiga toxin-producing *Escherichia coli* infection, antibiotics, and risk of developing hemolytic uremic syndrome: a meta-analysis. *Clin Infect Dis*. 2016;62:1251–1258. Review. [Epub 2016 Feb 24].
183. Levine MM, Nataro JP, Karch H, et al. The diarrheal response of humans to some classic serotypes of enteropathogenic *Escherichia coli* is dependent on a plasmid encoding an enteroadhesiveness factor. *J Infect Dis*. 1985;152:550–559.
184. Donnenberg MS, Tacket CO, James SP, et al. The role of the *eaeA* gene in experimental enteropathogenic *Escherichia coli* infection. *J Clin Invest*. 1993;92:1412–1417.
185. Kotloff KL, Nataro JP, Blackwelder WC, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet*. 2013;382:209–222.
187. Cohen MB, Nataro JP, Bernstein DI, et al. Prevalence of diarrheagenic *Escherichia coli* in acute childhood enteritis: a prospective controlled study. *J Pediatr*. 2005;146:54–61.
191. Bieber D, Ramer SW, Wu CY, et al. Type IV pili, transient bacterial aggregates, and virulence of enteropathogenic *Escherichia coli*. *Science*. 1998;280:2114–2118.
212. Deng W, Puente JL, Gruenheid S, et al. Dissecting virulence: systematic and functional analyses of a pathogenicity island. *Proc Natl Acad Sci USA*. 2004;101:3597–3602.
257. Garg AX, Suri RS, Barrowman N, et al. Long-term renal prognosis of diarrhea-associated hemolytic uremic syndrome—a systematic review, meta-analysis, and meta-regression. *JAMA*. 2003;290:1360–1370.
262. Wong CS, Mooney JC, Brandt JR, et al. Risk factors for the hemolytic uremic syndrome in children infected with *Escherichia coli* O157:H7: a multivariable analysis. *Clin Infect Dis*. 2012;55:33–41. [Epub 2012 Mar 19].
265. Frank C, Werber D, Cramer JP, et al. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. *N Engl J Med*. 2011;365:1771–1780.
266. Menne J, Nitschke M, Stingle R, et al; EHEC-HUS consortium. Validation of treatment strategies for enterohaemorrhagic *Escherichia coli* O104:H4 induced haemolytic uraemic syndrome: case-control study. *BMJ*. 2012;345:e4565.
289. O'Brien AD, Newland JW, Miller SE, et al. Shiga-like toxin-converting phages from *Escherichia coli* strains that cause hemorrhagic colitis or infantile diarrhea. *Science*. 1984;226:694–696.
348. Adachi JA, Jiang ZD, Mathewson JJ, et al. Enterotoxigenic *Escherichia coli* as a major etiologic agent in traveler's diarrhea in 3 regions of the world. *Clin Infect Dis*. 2001;32:1706–1709.
464. Johnson JR, Russo TA. Uropathogenic *Escherichia coli* as agents of diverse non-urinary tract extraintestinal infections. *J Infect Dis*. 2002;186:859–864.
473. Banerjee R, Johnson JR. A new clone sweeps clean: the enigmatic emergence of *Escherichia coli* sequence type 131. *Antimicrob Agents Chemother*. 2014;58:4997–5004. Review. [Epub 2014 May 27].
567. Kim KS, Itabashi H, Genski P, et al. The K1 capsule is the critical determinant in the development of *Escherichia coli* meningitis in the rat. *J Clin Invest*. 1992;90:897–905.
569. Basmaci R, Bonacorsi S, Bidet P, et al. *Escherichia coli* meningitis features in 325 children from 2001 to 2013 in France. *Clin Infect Dis*. 2015;61:779–786. [Epub 2015 May 5].
612. Weiner LM, Webb AK, Limbago B, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. *Infect Control Hosp Epidemiol*. 2016;37:1288–1301. [Epub 2016 Aug 30].
627. Brisse S, Fevre C, Passet V, et al. Virulent clones of *Klebsiella pneumoniae*: identification and evolutionary scenario based on genomic and phenotypic characterization. *PLoS ONE*. 2009;4:e4982. [Epub 2009 Mar 25].
657. Shon AS, Bajwa RP, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. *Virulence*. 2013;4:107–118. [Epub 2013 Jan 9].
679. Ben-David D, Kordevani R, Keller N, et al. Outcome of carbapenem resistant *Klebsiella pneumoniae* bloodstream infections. *Clin Microbiol Infect*. 2012;18:54–60.
694. Forsythe SJ, Abbott SL, Pitout J. *Klebsiella*, *Enterobacter*, *Citrobacter*, *Cronobacter*, *Serratia*, *Plesiomonas*, and other Enterobacteriaceae. In: *Manual of Clinical Microbiology*. 10th ed. Washington, DC: American Society of Microbiology; 2015:714–737.
723. Boucher HW, Talbot GH, Bradley JS, et al. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;48:1–12.
732. Mahlen SD. *Serratia* infections: from military experiments to current practice. *Clin Microbiol Rev*. 2011;24:755–791.
785. Jacobsen SM, Stickler DJ, Mobley HL, et al. Complicated catheter-associated urinary tract infections due to *Escherichia coli* and *Proteus mirabilis*. *Clin Microbiol Rev*. 2008;21:26–59.
782. Armbruster CE, Mobley HL. Merging mythology and morphology: the multifaceted lifestyle of *Proteus mirabilis*. *Nat Rev Microbiol*. 2012;10:743–754. [Epub 2012 Oct 8].
783. O'Hara CM, Brenner FW, Miller JM. Classification, identification, and clinical significance of *Proteus*, *Providencia*, and *Morganella*. *Clin Microbiol Rev*. 2000;13:534–546.

References

- Pupo GM, Karaolis DKR, Lan RT, et al. Evolutionary relationships among pathogenic and nonpathogenic *Escherichia coli* strains inferred from multilocus enzyme electrophoresis and *mdh* sequence studies. *Infect Immun*. 1997;65:2685–2692.
- Mackowiak PA, Martin RM, Jones SR, et al. Pharyngeal colonization by gram-negative bacilli in aspiration-prone persons. *Arch Intern Med*. 1978;138:1224–1227.
- Johanson WG, Pierce AK, Sanford JP. Changing pharyngeal bacterial flora of hospitalized patients: emergence of gram-negative bacilli. *N Engl J Med*. 1969;281:1137–1140.
- Raz R, Stamm WE. A controlled trial of intravaginal estriol in postmenopausal women with recurrent urinary tract infections. *N Engl J Med*. 1993;329:753–756.
- Gupta K, Hillier SL, Hooton TM, et al. Effects of contraceptive method on the vaginal microbial flora: a prospective evaluation. *J Infect Dis*. 2000;181:595–601.
- Doi Y, Bonomo RA, Hooper DC, et al. Gram-Negative Committee of the Antibacterial Resistance Leadership Group (ARLG). Gram-negative bacterial infections: research priorities, accomplishments, and future directions of the Antibacterial Resistance Leadership Group. *Clin Infect Dis*. 2017;64(suppl_1):S30–S35, Review.
- Peleg AY, Hooper DC. Hospital-acquired infections due to gram-negative bacteria. *N Engl J Med*. 2010;362:1804–1813.
- Sandora TJ, Goldmann DA. Preventing lethal hospital outbreaks of antibiotic-resistant bacteria. *N Engl J Med*. 2012;367:2168–2170.
- Moellering RC Jr. NDM-1—a cause for worldwide concern. *N Engl J Med*. 2010;363:2377–2379.
- Leverstein-van Hall MA, Blok HEM, Donders AR, et al. Multi-drug resistance among Enterobacteriaceae is strongly associated with the presence of integrons and is independent of species or isolate origin. *J Infect Dis*. 2003;187:251–259.
- Allos BM, Moore MR, Griffin PM, et al. Surveillance for sporadic foodborne disease in the 21st century: the FoodNet perspective. *Clin Infect Dis*. 2004;38(suppl 3):S115–S120.
- Schwaber MJ, Carmeli Y. The Impact of a Carbapenem-Resistant Enterobacteriaceae Outbreak on Facilitating Development of a National Infrastructure for Infection Control in Israel. *Clin Infect Dis*. 2017;65:2144–2149.
- Snitkin ES, Zelazny AM, Thomas PJ, et al. NISC Comparative Sequencing Program Group, Henderson DK, Palmore TN, Segre JA. Tracking a hospital outbreak of carbapenem-resistant *Klebsiella pneumoniae* with whole-genome sequencing. *Sci Transl Med*. 2012;4.
- Kadner RJ. Inner membrane. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:58–87.
- Blattner FR, Plunkett G 3rd, Bloch CA, et al. The complete genome sequence of *Escherichia coli* K-12. *Science*. 1997;277:1453–1462.
- Oliver DB. Periplasm. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:88–103.
- Park JT. The murein sacculus. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:48–57.
- Koebnik R, Locher KP, Van Gelder P. Structure and function of bacterial outer membrane proteins: barrels in a nutshell. *Mol Microbiol*. 2000;37:239–253.
- Nikaido H. Outer membrane. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:29–47.
- Rick PD, Silver RP. Enterobacterial common antigen and capsular polysaccharides. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:104–122.
- Ørskov F, Ørskov I. *Escherichia coli* serotyping and disease in man and animals. *Can J Microbiol*. 1992;38:699–704.
- Whitfield C, Roberts IS. Structure, assembly and regulation of expression of capsules in *Escherichia coli*. *Mol Microbiol*. 1999;31:1307–1319.
- Girón JA. Expression of flagella and motility by *Shigella*. *Mol Microbiol*. 1995;18:63–75.
- Hayashi F, Smith KD, Ozinsky A, et al. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature*. 2001;410:1099–1103.
- Gewirtz AT, Simon PO Jr, Schmitt CK, et al. *Salmonella typhimurium* translocates flagellin across intestinal epithelia, inducing a proinflammatory response. *J Clin Invest*. 2001;107:99–109.
- Steiner TS, Nataro JP, Potet-Smith CE, et al. Enterotoxigenic *Escherichia coli* expresses a novel flagellin that causes IL-8 release from intestinal epithelial cells. *J Clin Invest*. 2000;105:1769–1777.
- Wang L, Rothmund D, Curd H, et al. Species-wide variation in the *Escherichia coli* flagellin (H-antigen) gene. *J Bacteriol*. 2003;185:2936–2943.
- Ottow JC. Ecology, physiology, and genetics of fimbriae and pili. *Annu Rev Microbiol*. 1975;29:79–108.
- Lessl M, Lanka E. Common mechanisms in bacterial conjugation and Ti-mediated T-DNA transfer to plant cells. *Cell*. 1994;77:321–324.
- McClelland M, Sanderson KE, Spieth J, et al. Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature*. 2001;413:852–856.
- Parkhill J, Dougan G, James KD, et al. Complete genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. *Nature*. 2001;413:848–852.
- Welch RA, Burland V, Plunkett G 3rd, et al. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc Natl Acad Sci USA*. 2002;99:17020–17024.
- Thanassi DG, Saulino ET, Lombardo MJ, et al. The PapC usher forms an oligomeric channel: implications for pilus biogenesis across the outer membrane. *Proc Natl Acad Sci USA*. 1998;95:3146–3151.
- Dodson KW, Jacob-Dubuisson F, Striker RT, et al. Outer-membrane PapC molecular usher discriminately recognizes periplasmic chaperone-pilus subunit complexes. *Proc Natl Acad Sci USA*. 1993;90:3670–3674.
- Jones CH, Pinkner JS, Roth R, et al. FimH adhesin of type 1 pili is assembled into a fibrillar tip structure in the Enterobacteriaceae. *Proc Natl Acad Sci USA*. 1995;92:2081–2085.
- Kuehn MJ, Heuser J, Normark S, et al. P pili in uropathogenic *E. coli* are composite fibres with distinct fibrillar adhesive tips. *Nature*. 1992;356:252–255.
- Girón JA, Ho ASY, Schoolnik GK. An inducible bundle-forming pilus of enteropathogenic *Escherichia coli*. *Science*. 1991;254:710–713.
- Girón JA, Levine MM, Kaper JB. Longus: a long pilus ultrastructure produced by human enterotoxigenic *Escherichia coli*. *Mol Microbiol*. 1994;12:71–82.
- Xicotencatl-Cortes J, Monteiro-Neto V, Ledesma MA, et al. Intestinal adherence associated with type IV pili of enterohemorrhagic *Escherichia coli* O157:H7. *J Clin Invest*. 2007;117:3519–3529.
- Zhang XL, Tsui IS, Yip CM, et al. *Salmonella enterica* serovar Typhi uses type IVB pili to enter human intestinal epithelial cells. *Infect Immun*. 2000;68:3067–3073.
- Collyn F, Lety MA, Nair S, et al. *Yersinia pseudotuberculosis* harbors a type IV pilus gene cluster that contributes to pathogenicity. *Infect Immun*. 2002;70:6196–6205.
- Craig L, Pique ME, Tainer JA. Type IV pilus structure and bacterial pathogenicity. *Nat Rev Microbiol*. 2004;2:363–378.
- Merz AJ, So M, Sheetz MP. Pilus retraction powers bacterial twitching motility. *Nature*. 2000;407:98–102.
- Falkow S. Molecular Koch's postulates applied to microbial pathogenicity. *Rev Infect Dis*. 1988;10(suppl 2):S274–S276.
- Finlay BB, Falkow S. Common themes in microbial pathogenicity revisited. *Microbiol Rev*. 1997;61:136–169.
- Connell H, Agace W, Klemm P, et al. Type 1 fimbrial expression enhances *Escherichia coli* virulence for the urinary tract. *Proc Natl Acad Sci USA*. 1996;93:9827–9832.
- Hung CS, Bouckaert J, Hung D, et al. Structural basis of tropism of *Escherichia coli* to the bladder during urinary tract infection. *Mol Microbiol*. 2002;44:903–915.
- Ofek I, Mirelman D, Sharon N. Adherence of *Escherichia coli* to human mucosal cells mediated by mannose receptors. *Nature*. 1977;265:623–625.
- Abraham JM, Freitag CS, Clements JR, et al. An invertible element of DNA controls phase variation of type 1 fimbriae of *Escherichia coli*. *Proc Natl Acad Sci USA*. 1985;82:5724–5727.
- Thomas WE, Trintchina E, Forero M, et al. Bacterial adhesion to target cells enhanced by shear force. *Cell*. 2002;109:913–923.
- Bloch CA, Stocker BAD, Orndorff PE. A key role for type 1 pili in enterobacterial communicability. *Mol Microbiol*. 1992;6:697–701.
- Bloch CA, Orndorff PE. Impaired colonization by and full invasiveness of *Escherichia coli* K1 bearing a site-directed mutation in the type 1 pilin gene. *Infect Immun*. 1990;58:275–278.
- Tran Van Nhieu G, Isberg RR. The *Yersinia pseudotuberculosis* invasin protein and human fibronectin bind to mutually exclusive sites on the $\alpha_5\beta_1$ integrin receptor. *J Biol Chem*. 1991;266:24367–24375.
- Kenny B, DeVinney R, Stein M, et al. Enteropathogenic *E. coli* (EPEC) transfers its receptor for intimate adherence into mammalian cells. *Cell*. 1997;91:511–520.
- Hamburger ZA, Brown MS, Isberg RR, et al. Crystal structure of invasin: a bacterial integrin-binding protein. *Science*. 1999;286:291–295.
- Luo Y, Frey EA, Pfuetzner RA, et al. Crystal structure of enteropathogenic *Escherichia coli* intimin-receptor complex. *Nature*. 2000;405:1073–1077.
- Kelly G, Prasanna S, Daniell S, et al. Structure of the cell-adhesion fragment of intimin from enteropathogenic *Escherichia coli*. *Nat Struct Biol*. 1999;6:313–318.
- Economou A, Christie PJ, Fernandez RC, et al. Secretion by numbers: protein traffic in prokaryotes. *Mol Microbiol*. 2006;62:308–319.
- Henderson IR, Nataro JP. Virulence functions of autotransporter proteins. *Infect Immun*. 2001;69:1231–1243.
- Dutta PR, Cappello R, Navarro-Garcia F, et al. Functional comparison of serine protease autotransporters of Enterobacteriaceae. *Infect Immun*. 2002;70:7105–7113.
- Dautin N. Serine protease autotransporters of Enterobacteriaceae (SPATEs): biogenesis and function. *Toxins (Basel)*. 2010;2:1179–1206.
- Koronakis V, Sharff A, Koronakis E, et al. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature*. 2000;405:914–919.
- Thanabalu T, Koronakis E, Hughes C, et al. Substrate-induced assembly of a contiguous channel for protein export from *E. coli*: reversible bridging of an inner-membrane translocase to an outer membrane exit pore. *EMBO J*. 1998;17:6487–6496.
- Foreman DT, Martinez Y, Coombs G, et al. TolC and DsbA are needed for the secretion of ST₁₀, a heat-stable enterotoxin of *Escherichia coli*. *Mol Microbiol*. 1995;18:237–245.
- Soloaga A, Veiga MP, García-Segura LM, et al. Insertion of *Escherichia coli* α -haemolysin in lipid bilayers as a non-transmembrane integral protein: prediction and experiment. *Mol Microbiol*. 1999;31:1013–1024.
- Uhlén P, Laestadius Å, Jahnukainen T, et al. Alpha-haemolysin of uropathogenic *E. coli* induces Ca²⁺ oscillations in renal epithelial cells. *Nature*. 2000;405:694–697.
- Sandkvist M. Biology of type II secretion. *Mol Microbiol*. 2001;40:271–283.
- Latham WW, Grys TE, Witowski SE, et al. StcE, a metalloprotease secreted by *Escherichia coli* O157:H7, specifically cleaves Cl esterase inhibitor. *Mol Microbiol*. 2002;45:277–288.
- Galán JE, Collmer A. Type III secretion machines: bacterial devices for protein delivery into host cells. *Science*. 1999;284:1322–1328.
- Cornelis GR. The type III secretion injectisome. *Nat Rev Microbiol*. 2006;4:811–825.
- Stebbins CE, Galán JE. Structural mimicry in bacterial virulence. *Nature*. 2001;412:701–705.
- Reife RA, Shapiro RA, Bamber BA, et al. *Porphyromonas gingivalis* lipopolysaccharide is poorly recognized by molecular components of innate host defense in a mouse model of early inflammation. *Infect Immun*. 1995;63:4686–4694.
- Raetz CRH. Bacterial lipopolysaccharides: a remarkable family of bioactive macroamphiphiles. In: Neidhardt FC, ed. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: ASM Press; 1996:1035–1063.
- Stenutz R, Weintraub A, Widmalm G. The structures of *Escherichia coli* O-polysaccharide antigens. *FEMS Microbiol Rev*. 2006;30:382–403, Review.
- Agnese DM, Calvano JE, Hahn SJ, et al. Human Toll-like receptor 4 mutations but not CD14 polymorphisms are associated with an increased risk of gram-negative infections. *J Infect Dis*. 2002;186:1522–1525.
- Faber J, Henninger N, Finn A, et al. A Toll-like receptor 4 variant is associated with fatal outcome in children with invasive meningococcal disease. *Acta Paediatr*. 2009;98:448–452.
- Horwitz MA, Silverstein SC. Influence of the *Escherichia coli* capsule on complement fixation and on phagocytosis and killing by human phagocytes. *J Clin Invest*. 1980;65:82–94.
- Cross AS, Kim KS, Wright DC, et al. Role of lipopolysaccharide and capsule in the serum resistance of bacteremic strains of *Escherichia coli*. *J Infect Dis*. 1986;154:497–503.
- Russo TA, Moffitt MC, Hammer CH, et al. TnphoA-mediated disruption of K54 capsular polysaccharide genes in *Escherichia coli* confers serum sensitivity. *Infect Immun*. 1993;61:3578–3582.
- Hoffman JA, Wass C, Stins MF, et al. The capsule supports survival but not traversal of *Escherichia coli* K1 across the blood-brain barrier. *Infect Immun*. 1999;67:3566–3570.

81. Podschun R, Ullmann U. *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clin Microbiol Rev*. 1998;11:589–603.
82. Escolar L, Pérez-Martín J, De Lorenzo V. Opening the iron box: transcriptional metalloreulation by the Fur protein. *J Bacteriol*. 1999;181:6223–6229.
83. Troxell B, Hassan HM. Transcriptional regulation by Ferric Uptake Regulator (Fur) in pathogenic bacteria. *Front Cell Infect Microbiol*. 2013;3:59, eCollection 2013. Review.
84. Raymond KN, Dertz EA, Kim SS. Enterobactin: an archetype for microbial iron transport. *Proc Natl Acad Sci USA*. 2003;100:3584–3588.
85. Carbonetti NH, Boonchai S, Parry SH, et al. Aerobactin-mediated iron uptake by *Escherichia coli* isolates from human extraintestinal infections. *Infect Immun*. 1986;51:966–968.
86. Johnson JR, Moseley SL, Roberts PL, et al. Aerobactin and other virulence factor genes among strains of *Escherichia coli* causing urosepsis: association with patient characteristics. *Infect Immun*. 1988;56:405–412.
87. Torres AG, Redford P, Welch RA, et al. TonB-dependent systems of uropathogenic *Escherichia coli*: aerobactin and heme transport and TonB are required for virulence in the mouse. *Infect Immun*. 2001;69:6179–6185.
88. Jiang XQ, Payne MA, Cao ZH, et al. Ligand-specific opening of a gated-pore channel in the outer membrane of living bacteria. *Science*. 1997;276:1261–1264.
89. Shulman ST, Friedmann HC, Sims RH. Theodor Escherich: the first pediatric infectious diseases physician? *Clin Infect Dis*. 2007;45:1025–1029.
90. Escherich T. The intestinal bacteria of the neonate and breast-fed infant 1884. *Rev Infect Dis*. 1988;10:1220–1225.
91. Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol*. 2004;2:123–140.
92. Albert MJ, Alam K, Islam M, et al. *Hafnia alvei*, a probable cause of diarrhea in humans. *Infect Immun*. 1991;59:1507–1513.
93. Albert MJ, Faruque SM, Ansaruzzaman M, et al. Sharing of virulence-associated properties at the phenotypic and genetic levels between enteropathogenic *Escherichia coli* and *Hafnia alvei*. *J Med Microbiol*. 1992;37:310–314.
94. Huys G, Cnockaert M, Janda JM, et al. *Escherichia albertii* sp. nov., a diarrhoeagenic species isolated from stool specimens of Bangladeshi children. *Int J Syst Evol Microbiol*. 2003;53:807–810.
95. Janda JM, Abbott SL, Albert MJ. Prototypal diarrheagenic strains of *Hafnia alvei* are actually members of the genus *Escherichia*. *J Clin Microbiol*. 1999;37:2399–2401.
96. Hyma KE, Lacher DW, Nelson AM, et al. Evolutionary genetics of a new pathogenic *Escherichia* species: *Escherichia albertii* and related *Shigella boydii* strains. *J Bacteriol*. 2005;187:619–628.
97. Ooka T, Seto K, Kawano K, et al. Clinical significance of *Escherichia albertii*. *Emerg Infect Dis*. 2012;18:488–492.
98. Rasko DA, Rosovitz MJ, Myers GS, et al. The pangenome structure of *Escherichia coli*: comparative genomic analysis of *E. coli* commensal and pathogenic isolates. *J Bacteriol*. 2008;190:6881–6893.
99. Robins-Browne RM, Holt KE, Ingle DJ, et al. Are *Escherichia coli* pathotypes still relevant in the era of whole-genome sequencing? *Front Cell Infect Microbiol*. 2016;6:141, eCollection 2016. Review.
100. Croxson MA, Law RJ, Scholz R, et al. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev*. 2013;26:822–880.
101. McDaniel TK, Jarvis KG, Donnenberg MS, et al. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc Natl Acad Sci USA*. 1995;92:1664–1668.
102. Leimbach A, Hacker J, Dobrindt U. *E. coli* as an all-rounder: the thin line between commensalism and pathogenicity. *Curr Top Microbiol Immunol*. 2013;358:3–32. Review.
103. Dobrindt U, Hochhut B, Hentschel U, et al. Genomic islands in pathogenic and environmental microorganisms. *Nat Rev Microbiol*. 2004;2:414–424. Review.
104. Hacker J, Kaper JB. Pathogenicity islands and the evolution of microbes. *Annu Rev Microbiol*. 2000;54:641–679. Review.
105. Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev*. 1998;11:142–201.
106. Cravioto A, Gross RJ, Scotland SM, et al. An adhesive factor found in strains of *Escherichia coli* belonging to the traditional infantile enteropathogenic serotypes. *Curr Microbiol*. 1979;3:95–99.
107. Scaletsky ICA, Silva MLM, Trablusi LR. Distinctive patterns of adherence of enteropathogenic *Escherichia coli* to HeLa cells. *Infect Immun*. 1984;45:534–536.
108. Clements A, Young JC, Constantinou N, et al. Infection strategies of enteric pathogenic *Escherichia coli*. *Gut Microbes*. 2012;3:71–87. Review. [Epub 2012 Mar 1].
109. Karmali MA. Emerging Public Health Challenges of Shiga Toxin-Producing *Escherichia coli* Related to Changes in the Pathogen, the Population, and the Environment. *Clin Infect Dis*. 2017;64:371–376. Review. [Epub 2016 Oct 21].
110. Rasko DA, Webster DR, Sahl JW, et al. Origins of the *E. coli* strain causing an outbreak of hemolytic-uremic syndrome in Germany. *N Engl J Med*. 2011;365:709–717.
111. Rohde H, Qin J, Cui Y, et al. *E. coli* O104:H4 genome analysis Crowd-Sourcing consortium. Open-Source genomic analysis of Shiga-toxin-producing *E. coli* O104:H4. *N Engl J Med*. 2011;365:718–724. [Epub 2011 Jul 27].
112. Russo TA, Johnson JR. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J Infect Dis*. 2000;181:1753–1754.
113. Wang MC, Tseng CC, Chen CY, et al. The role of bacterial virulence and host factors in patients with *Escherichia coli* bacteremia who have acute cholangitis or upper urinary tract infection. *Clin Infect Dis*. 2002;35:1161–1166.
114. Bray J. Isolation of antigenically homogeneous strains of *Bact. coli neapolitanum* from summer diarrhoea of infants. *J Pathol Bacteriol*. 1945;57:239–247.
115. Afset JE, Bergh K, Bevanger L. High prevalence of atypical enteropathogenic *Escherichia coli* (EPEC) in Norwegian children with diarrhoea. *J Med Microbiol*. 2003;52(Pt 11):1015–1019.
116. Philippsborn R, Ahmed SM, Brosi BJ, et al. Climatic drivers of diarrheagenic *Escherichia coli* incidence: a systematic review and meta-analysis. *J Infect Dis*. 2016;214:6–15. Review. [Epub 2016 Feb 29].
117. Shane AL, Mody RK, Crump JA, et al. 2017 Infectious Diseases Society of America clinical practice guidelines for the diagnosis and management of infectious diarrhea. *Clin Infect Dis*. 2017;65:1963–1973.
118. Riddle MS, Connor P, Fraser J, et al. TrEAT TD Study Team. Trial Evaluating Ambulatory Therapy of Travelers' diarrhea (TrEAT TD) study: a randomized controlled trial comparing 3 single-dose antibiotic regimens with loperamide. *Clin Infect Dis*. 2017;65:2008–2017.
119. Riddle MS, DuPont HL, Connor BA. ACG clinical guideline: diagnosis, treatment, and prevention of acute diarrheal infections in adults. *Am J Gastroenterol*. 2016;111:602–622. [Epub 2016 Apr 12].
120. Fang FC, Patel R. 2017 Infectious Diseases Society of America infectious diarrhea guidelines: a view from the clinical laboratory. *Clin Infect Dis*. 2017;65:1974–1976.
121. Binnicker MJ. Multiplex molecular panels for diagnosis of gastrointestinal infection: performance, result interpretation, and cost-effectiveness. *J Clin Microbiol*. 2015;53:3723–3728. Review.
122. Lääveri T, Antikainen J, Pakkanen SH, et al. Prospective study of pathogens in asymptomatic travellers and those with diarrhoea: aetiological agents revisited. *Clin Microbiol Infect*. 2016;22:535–541. [Epub 2016 Mar 9].
123. Shah N, DuPont HL, Ramsey DJ. Global etiology of travelers' diarrhea: systematic review from 1973 to the present. *Am J Trop Med Hyg*. 2009;80:609–614. Review.
124. Qadri F, Svennerholm AM, Faruque AS, et al. Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin Microbiol Rev*. 2005;18:465–483. Review.
125. Adachi JA, Ostrosky-Zeichner L, DuPont HL, et al. Empirical antimicrobial therapy for traveler's diarrhea. *Clin Infect Dis*. 2000;31:1079–1083.
126. Adachi JA, Ericsson CD, Jiang ZD, et al. Azithromycin found to be comparable to levofloxacin for the treatment of US travelers with acute diarrhea acquired in Mexico. *Clin Infect Dis*. 2003;37:1165–1171.
127. DuPont HL, Jiang ZD, Ericsson CD, et al. Rifaximin versus ciprofloxacin for the treatment of traveler's diarrhea: a randomized, double-blind clinical trial. *Clin Infect Dis*. 2001;33:1807–1815.
128. De Bruyn G, Hahn S, Borwick A. Antibiotic treatment for travellers' diarrhoea. *Cochrane Database Syst Rev*. 2000;(3):CD002242.
129. Kantele A, Lääveri T, Mero S, et al. Antimicrobials increase travelers' risk of colonization by extended-spectrum betalactamase-producing Enterobacteriaceae. *Clin Infect Dis*. 2015;60:837–846. [Epub 2015 Jan 21].
130. Connor BA, Keystone JS. Editorial commentary: antibiotic self-treatment of travelers' diarrhea: helpful or harmful? *Clin Infect Dis*. 2015;60:847–848. [Epub 2015 Jan 21].
131. DuPont HL, Ericsson CD, Farthing MJ, et al. Expert review of the evidence base for self-therapy of travelers' diarrhea. *J Travel Med*. 2009;16:161–171. Review. [Epub 2009 Mar 19].
132. Ericsson CD, DuPont HL. Travelers' diarrhea: approaches to prevention and treatment. *Clin Infect Dis*. 1993;16:616–626.
133. Tafuri S, Guerra R, Gallone MS, et al. Effectiveness of pre-travel consultation in the prevention of travel-related diseases: a retrospective cohort study. *Travel Med Infect Dis*. 2014;12(6 Pt B):745–749.
134. Shlim DR. Looking for evidence that personal hygiene precautions prevent traveler's diarrhea. *Clin Infect Dis*. 2005;41(suppl 8):S531–S535. Review.
135. Kozicki M, Steffen R, Schär M. 'Boil it, cook it, peel it or forget it': does this rule prevent travellers' diarrhoea? *Int J Epidemiol*. 1985;14:169–172.
136. DuPont HL, Ericsson CD. Drug therapy: prevention and treatment of traveler's diarrhea. *N Engl J Med*. 1993;328:1821–1827.
137. Flores J, Dupont HL, Jiang ZD, et al. A randomized, double-blind, pilot study of rifaximin 550 mg versus placebo in the prevention of travelers' diarrhea in Mexico during the dry season. *J Travel Med*. 2011;18:333–336. [Epub 2011 Aug 1].
138. Gupta SK, Keck J, Ram PK, et al. Analysis of data gaps pertaining to enterotoxigenic *Escherichia coli* infections in low and medium human development index countries, 1984–2005. *Epidemiol Infect*. 2008;136:721–738. Review. [Epub 2007 Aug 9].
139. Beatty ME, Adcock PM, Smith SW, et al. Epidemic diarrhea due to enterotoxigenic *Escherichia coli*. *Clin Infect Dis*. 2006;42:329–334.
140. Devasia RA, Jones TF, Ward J, et al. Endemically acquired foodborne outbreak of enterotoxin-producing *Escherichia coli* serotype O169:H41. *Am J Med*. 2006;119:168, e7–10.
141. Beatty ME, Bopp CA, Wells JG, et al. Enterotoxin-producing *Escherichia coli* O169:H41, United States. *Emerg Infect Dis*. 2004;10:518–521.
142. Madhavan TP, Sakellaris H. Colonization factors of enterotoxigenic *Escherichia coli*. *Adv Appl Microbiol*. 2015;90:155–197. Review. [Epub 2014 Nov 12].
143. Elsinghorst EA. Enterotoxigenic *Escherichia coli*. In: Donnenberg MS, ed. *Escherichia coli: Virulence Mechanisms of a Versatile Pathogen*. San Diego: Academic Press; 2002:155–187.
144. Steinsland H, Valentiner-Branth P, Perch M, et al. Enterotoxigenic *Escherichia coli* infections and diarrhea in a cohort of young children in Guinea-Bissau. *J Infect Dis*. 2002;186:1740–1747.
145. von Mentzer A, Connor TR, Wieler LH, et al. Identification of enterotoxigenic *Escherichia coli* (ETEC) clades with long-term global distribution. *Nat Genet*. 2014;46:1321–1326. [Epub 2014 Nov 10].
146. Levine MM, Nalin DR, Hoover DL, et al. Immunity to enterotoxigenic *Escherichia coli*. *Infect Immun*. 1979;23:729–736.
147. Qadri F, Saha A, Ahmed T, et al. Disease burden due to enterotoxigenic *Escherichia coli* in the first 2 years of life in an urban community in Bangladesh. *Infect Immun*. 2007;75:3961–3968. [Epub 2007 Jun 4].
148. Steinsland H, Valentiner-Branth P, Gjessing HK, et al. Protection from natural infections with enterotoxigenic *Escherichia coli*: longitudinal study. *Lancet*. 2003;362:286–291.
149. Freedman SB, Xie J, Neufeld MS, et al. Alberta Provincial Pediatric Enteric Infection Team (APPETITE), Nettel-Aguirre A, Chuck A, Lee B, Johnson D, Currie G, Talbot J, Jiang J, Dickinson J, Kellner J, MacDonald J, Svenson L, Chui L, Louie M, Lavoie M, Eltorik M, Vanderkooi O, Tellier R, Ali S, Drews S, Graham T, Pang XL. Shiga toxin-producing *Escherichia coli* infection, antibiotics, and risk of developing hemolytic uremic syndrome: a meta-analysis. *Clin Infect Dis*. 2016;62:1251–1258. Review. [Epub 2016 Feb 24].
150. Roy K, Kansal R, Bartels SR, et al. Adhesin degradation accelerates delivery of heat-labile toxin by enterotoxigenic *Escherichia coli*. *J Biol Chem*. 2011;286:29771–29779, Aug 26. [Epub 2011 Jul 8].
151. Isidean SD, Riddle MS, Savarino SJ, et al. A systematic review of ETEC epidemiology focusing on colonization factor and toxin expression. *Vaccine*. 2011;29:6167–6178. Review. [Epub 2011 Jul 1].
152. Sixma TK, Pronk SE, Kalk KH, et al. Crystal structure of a cholera toxin-related heat-labile enterotoxin from *E. coli*. *Nature*. 1991;351:371–377.
153. Tauschek M, Gorrell RJ, Strugnell RA, et al. Identification of a protein secretory pathway for the secretion of heat-labile enterotoxin by an enterotoxigenic strain of *Escherichia coli*. *Proc Natl Acad Sci USA*. 2002;99:7066–7071.
154. Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clin Microbiol Rev*. 1998;11:142–201. Review. Erratum in: *Clin Microbiol Rev* 1998 Apr;11(2):403.
155. Sears CL, Kaper JB. Enteric bacterial toxins: mechanisms of action and linkage to intestinal secretion. *Microbiol Rev*. 1996;60:167–215. Review.
156. Joffré E, von Mentzer A, Abd El Ghany M, et al. Allele variants of enterotoxigenic *Escherichia coli* heat-labile toxin are globally transmitted and associated with

- colonization factors. *J Bacteriol.* 2015;197:392–403. [Epub 2014 Nov 17].
157. Nada RA, Shaheen HI, Khalil SB, et al. Discovery and phylogenetic analysis of novel members of class B enterotoxigenic *Escherichia coli* adhesive fimbriae. *J Clin Microbiol.* 2011;49:1403–1410. [Epub 2011 Feb 2].
 158. Rodrigues JF, Mathias-Santos C, Sbraggio-Almeida ME, et al. Functional diversity of heat-labile toxins (LT) produced by enterotoxigenic *Escherichia coli*: differential enzymatic and immunological activities of LT1 (hLT) AND LT4 (pLT). *J Biol Chem.* 2011;286:5222–5233. [Epub 2010 Dec 6].
 159. Lasaro MA, Rodrigues JF, Mathias-Santos C, et al. Genetic diversity of heat-labile toxin expressed by enterotoxigenic *Escherichia coli* strains isolated from humans. *J Bacteriol.* 2008;190:2400–2410. [Epub 2008 Jan 25].
 160. Forte LR, Eber SL, Turner JT, et al. Guanylin stimulation of Cl⁻ secretion in human intestinal T₈₄ cells via cyclic guanosine monophosphate. *J Clin Invest.* 1993;91:2423–2428.
 161. Field M. Intestinal ion transport and the pathophysiology of diarrhea. *J Clin Invest.* 2003;111:931–943.
 162. Lucas ML. Diarrhoeal disease through enterocyte secretion: a doctrine untroubled by proof. *Exp Physiol.* 2010;95:479–484. Review.
 163. Lucas ML, Thom MM, Bradley JM, et al. *Escherichia coli* heat stable (STa) enterotoxin and the upper small intestine: lack of evidence in vivo for net fluid secretion. *J Membr Biol.* 2005;206:29–42.
 164. Joffré E, von Mentzer A, Svennerholm AM, et al. Identification of new heat-stable (STa) enterotoxin allele variants produced by human enterotoxigenic *Escherichia coli* (ETEC). *Int J Med Microbiol.* 2016;306:586–594. [Epub 2016 Jun 15].
 165. Rodas C, Iniguez V, Qadri F, et al. Development of multiplex PCR assays for detection of enterotoxigenic *Escherichia coli* colonization factors and toxins. *J Clin Microbiol.* 2009;47:1218–1220. [Epub 2009 Feb 25].
 166. Aranda KR, Fabbriotti SH, Fagundes-Neto U, et al. Single multiplex assay to identify simultaneously enteropathogenic, enteroaggregative, enterotoxigenic, enteroinvasive and Shiga toxin-producing *Escherichia coli* strains in Brazilian children. *FEMS Microbiol Lett.* 2007;267:145–150.
 167. Toma C, Martínez Espinosa E, Song T, et al. Distribution of putative adhesins in different serotypes of Shiga toxin-producing *Escherichia coli*. *J Clin Microbiol.* 2004;42:4937–4946.
 168. Vidal R, Vidal M, Lagos R, et al. Multiplex PCR for diagnosis of enteric infections associated with diarrheagenic *Escherichia coli*. *J Clin Microbiol.* 2004;42:1787–1789.
 169. Ozaki CY, Silveira CR, Andrade FB, et al. Single chain variable fragments produced in *Escherichia coli* against heat-labile and heat-stable toxins from enterotoxigenic *E. coli*. *PLoS ONE.* 2015;10:e0131484. eCollection 2015.
 170. Begum YA, Talukder KA, Azmi IJ, et al. Resistance pattern and molecular characterization of enterotoxigenic *Escherichia coli* (ETEC) strains isolated in Bangladesh. *PLoS ONE.* 2016;11:e0157415. eCollection 2016.
 171. Svennerholm AM. From cholera to enterotoxigenic *Escherichia coli* (ETEC) vaccine development. *Indian J Med Res.* 2011;133:188–196. Review.
 172. Sack DA, Shimko J, Torres O, et al. Randomised, double-blind, safety and efficacy of a killed oral vaccine for enterotoxigenic *E. coli* diarrhoea of travellers to Guatemala and Mexico. *Vaccine.* 2007;25:4392–4400. [Epub 2007 Apr 4].
 173. Frech SA, DuPont HL, Bourgeois AL, et al. Use of a patch containing heat-labile toxin from *Escherichia coli* against travellers' diarrhoea: a phase II, randomised, double-blind, placebo-controlled field trial. *Lancet.* 2008;371:2019–2025.
 174. Behrens RH, Cramer JP, Jelinek T, et al. Efficacy and safety of a patch vaccine containing heat-labile toxin from *Escherichia coli* against travellers' diarrhoea: a phase 3, randomised, double-blind, placebo-controlled field trial in travellers from Europe to Mexico and Guatemala. *Lancet Infect Dis.* 2014;14:197–204. [Epub 2013 Nov 29].
 175. Lundgren A, Bourgeois L, Carlin N, et al. Safety and immunogenicity of an improved oral inactivated multivalent enterotoxigenic *Escherichia coli* (ETEC) vaccine administered alone and together with dmlT adjuvant in a double-blind, randomized, placebo-controlled Phase I study. *Vaccine.* 2014;32:7077–7084. [Epub 2014 Nov 5].
 176. Roy K, Hilliard GM, Hamilton DJ, et al. Enterotoxigenic *Escherichia coli* EtpA mediates adhesion between flagella and host cells. *Nature.* 2009;457:594–598.
 177. Roy K, Hamilton D, Allen KP, et al. The EtpA exoprotein of enterotoxigenic *Escherichia coli* promotes intestinal colonization and is a protective antigen in an experimental model of murine infection. *Infect Immun.* 2008;76:2106–2112.
 178. Luo Q, Vickers TJ, Fleckenstein JM. Immunogenicity and protective efficacy against enterotoxigenic *Escherichia coli* colonization following intradermal, sublingual, or oral vaccination with EtpA adhesin. *Clin Vaccine Immunol.* 2016;23:628–637. Print 2016 Jul.
 179. Roy K, Hamilton D, Ostmann MM, et al. Vaccination with EtpA glycoprotein or flagellin protects against colonization with enterotoxigenic *Escherichia coli* in a murine model. *Vaccine.* 2009;27:4601–4608. [Epub 2009 Jun 11].
 180. Robins-Browne RM. Traditional enteropathogenic *Escherichia coli* of infantile diarrhea. *Rev Infect Dis.* 1987;9:28–53. Review.
 181. Neter E, Westphal O, Luderitz O, et al. Demonstration of antibodies against enteropathogenic *Escherichia coli* in sera of children of various ages. *Pediatrics.* 1955;16:801–808.
 182. Girón JA, Donnenberg MS, Martin WC, et al. Distribution of the bundle-forming pilus structural gene (*bfpA*) among enteropathogenic *Escherichia coli*. *J Infect Dis.* 1993;168:1037–1041.
 183. Levine MM, Nataro JP, Karch H, et al. The diarrheal response of humans to some classic serotypes of enteropathogenic *Escherichia coli* is dependent on a plasmid encoding an enteroadhesiveness factor. *J Infect Dis.* 1985;152:550–559.
 184. Donnenberg MS, Tacket CO, James SP, et al. The role of the *eaeA* gene in experimental enteropathogenic *Escherichia coli* infection. *J Clin Invest.* 1993;92:1412–1417.
 185. Kotloff KL, Nataro JP, Blackwelder WC, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet.* 2013;382:209–222.
 186. Afset JE, Bevanger L, Romundstad P, et al. Association of atypical enteropathogenic *Escherichia coli* (EPEC) with prolonged diarrhoea. *J Med Microbiol.* 2004;53:1137–1144.
 187. Cohen MB, Nataro JP, Bernstein DI, et al. Prevalence of diarrheagenic *Escherichia coli* in acute childhood enteritis: a prospective controlled study. *J Pediatr.* 2005;146:54–61.
 188. Robins-Browne RM, Bordun AM, Tauschek M, et al. *Escherichia coli* and community-acquired gastroenteritis, Melbourne, Australia. *Emerg Infect Dis.* 2004;10:1797–1805.
 189. Gomes TAT, Vieira MAM, Wachsmuth IK, et al. Serotype-specific prevalence of *Escherichia coli* strains with EPEC adherence factor genes in infants with and without diarrhea in São Paulo, Brazil. *J Infect Dis.* 1989;160:131–135.
 190. Trabulsi LR, Keller R, Gomes TAT. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg Infect Dis.* 2002;8:508–513.
 191. Bieber D, Ramer SW, Wu CY, et al. Type IV pili, transient bacterial aggregates, and virulence of enteropathogenic *Escherichia coli*. *Science.* 1998;280:2114–2118.
 192. Foster MA, Iqbal J, Zhang C, et al. Enteropathogenic and enteroaggregative *E. coli* in stools of children with acute gastroenteritis in Davidson County, Tennessee. *Diagn Microbiol Infect Dis.* 2015;83:319–324. [Epub 2015 Jul 26].
 193. Tobias J, Kassem E, Rubinstein U, et al. Involvement of main diarrheagenic *Escherichia coli*, with emphasis on enteroaggregative *E. coli*, in severe non-epidemic pediatric diarrhea in a high-income country. *BMC Infect Dis.* 2015;15:79.
 194. Hu J, Torres AG. Enteropathogenic *Escherichia coli*: foe or innocent bystander? *Clin Microbiol Infect.* 2015;21:729–734. Aug. Review. [Epub 2015 Jan 28].
 195. Gomes TA, Irino K, Girão DM, et al. Emerging enteropathogenic *Escherichia coli* strains? *Emerg Infect Dis.* 2004;10:1851–1855. Oct.
 196. Paulozzi LJ, Johnson KE, Kamahele LM, et al. Diarrhea associated with adherent enteropathogenic *Escherichia coli* in an infant and toddler center, Seattle, Washington. *Pediatrics.* 1986;77:296–300.
 197. Wu S-X, Peng R-Q. Studies on an outbreak of neonatal diarrhea caused by EPEC O127:H6 with plasmid analysis restriction analysis and outer membrane protein determination. *Acta Paediatr Scand.* 1992;81:217–221.
 198. Blake PA, Ramos S, Macdonald KL, et al. Pathogen-specific risk factors and protective factors for acute diarrheal disease in urban Brazilian infants. *J Infect Dis.* 1993;167:627–632.
 199. Thorén A, Stintzing G, Tufvesson B, et al. Aetiology and clinical features of severe infantile diarrhoea in Addis Ababa, Ethiopia. *J Trop Pediatr.* 1982;28:127–131.
 200. Gomes TAT, Rassi V, Macdonald KL, et al. Enteropathogens associated with acute diarrheal disease in urban infants in São Paulo, Brazil. *J Infect Dis.* 1991;164:331–337.
 201. Bower JR, Congeni BL, Cleary TG, et al. *Escherichia coli* O114: nonmotile as a pathogen in an outbreak of severe diarrhea associated with a day care center. *J Infect Dis.* 1989;160:243–247.
 202. Clausen CR, Christie DL. Chronic diarrhea in infants caused by adherent enteropathogenic *Escherichia coli*. *J Pediatr.* 1982;100:358–361.
 203. Rothbaum R, McAdams AJ, Giannella R, et al. A clinicopathological study of enterocyte-adherent *Escherichia coli*: a cause of protracted diarrhea in infants. *Gastroenterology.* 1982;83:441–454.
 204. Nougayrède JP, Fernandes PJ, Donnenberg MS. Adhesion of enteropathogenic *Escherichia coli* to host cells. *Cell Microbiol.* 2003;5:359–372. Review.
 205. Aroeti B, Friedman G, Zlotkin-Rivkin E, et al. Retraction of enteropathogenic *E. coli* type IV pili promotes efficient host cell colonization, effector translocation and tight junction disruption. *Gut Microbes.* 2012;3:267–271.
 206. Zahavi EE, Lieberman JA, Donnenberg MS, et al. Bundle forming pilus retraction enhances enteropathogenic *Escherichia coli* infectivity. *Mol Biol Cell.* 2011;22:2436–2447.
 207. Moreira CG, Palmer K, Whiteley M, et al. Bundle-forming pili and EspA are involved in biofilm formation by enteropathogenic *Escherichia coli*. *J Bacteriol.* 2006;188:3952–3961.
 208. Ashokkumar B, Kumar JS, Hecht GA, et al. Enteropathogenic *Escherichia coli* inhibits intestinal vitamin B1 (thiamin) uptake: studies with human-derived intestinal epithelial Caco-2 cells. *Am J Physiol Gastrointest Liver Physiol.* 2009;297:G825–G833. [Epub 2009 Jul 23].
 209. Garcia BG, Ooka T, Gotoh Y, et al. Genetic relatedness and virulence properties of enteropathogenic *Escherichia coli* strains of serotype O119:H6 expressing localized adherence or localized and aggregate adherence-like patterns on HeLa cells. *Int J Med Microbiol.* 2016;306:152–164. [Epub 2016 Feb 27].
 210. Kaper JB. Defining EPEC. *Rev Microbiol Sao Paulo.* 1996;27(suppl 1):130–133.
 211. McDaniel TK, Kaper JB. A cloned pathogenicity island from enteropathogenic *Escherichia coli* confers the attaching and effacing phenotype on K-12 *E. coli*. *Mol Microbiol.* 1997;23:399–407.
 212. Deng W, Puente JL, Gruenheid S, et al. Dissecting virulence: systematic and functional analyses of a pathogenicity island. *Proc Natl Acad Sci USA.* 2004;101:3597–3602.
 213. Iguchi A, Thomson NR, Ogura Y, et al. Complete genome sequence and comparative genome analysis of enteropathogenic *Escherichia coli* O127:H6 strain E2348/69. *J Bacteriol.* 2009;191:347–354.
 214. Cepeda-Molero M, Berger CN, Walsham ADS, et al. Attaching and effacing (A/E) lesion formation by enteropathogenic *E. coli* on human intestinal mucosa is dependent on non-LEE effectors. *PLoS Pathog.* 2017;13:e1006706. eCollection 2017 Oct.
 215. Knutton S, Rosenshine I, Pallen MJ, et al. A novel EspA-associated surface organelle of enteropathogenic *Escherichia coli* involved in protein translocation into epithelial cells. *EMBO J.* 1998;17:2166–2176.
 216. Frankel G, Phillips AD. Attaching effacing *Escherichia coli* and paradigms of Tir-triggered actin polymerization: getting off the pedestal. *Cell Microbiol.* 2008;10:549–556.
 217. Staley TE, Jones EW, Corley LD. Attachment and penetration of *Escherichia coli* into intestinal epithelium of the ileum in newborn pigs. *Am J Pathol.* 1969;56:371–392.
 218. Moon HW, Whipp SC, Argenzio RA, et al. Attaching and effacing activities of rabbit and human enteropathogenic *Escherichia coli* in pig and rabbit intestines. *Infect Immun.* 1983;41:1340–1351.
 219. Schüller S, Chong Y, Lewin J, et al. Tir phosphorylation and Nck/N-WASP recruitment by enteropathogenic and enterohaemorrhagic *Escherichia coli* during ex vivo colonization of human intestinal mucosa is different to cell culture models. *Cell Microbiol.* 2007;9:1352–1364.
 220. Kalman D, Weiner OD, Goosney DL, et al. Enteropathogenic *E. coli* acts through WASP and Arp2/3 complex to form actin pedestals. *Nat Cell Biol.* 1999;1:389–391.
 221. Sanger JM, Chang R, Ashton F, et al. Novel form of actin-based motility transports bacteria on the surfaces of infected cells. *Cell Motil Cytoskeleton.* 1996;34:279–287.
 222. Gao X, Wang X, Pham TH, et al. NleB, a bacterial effector with glycosyltransferase activity, targets GAPDH function to inhibit NF- κ B activation. *Cell Host Microbe.* 2013;13:87–99.
 223. Newton HJ, Pearson JS, Badea L, et al. The type III effectors NleE and NleB from enteropathogenic *E. coli* and OspZ from *Shigella* block nuclear translocation of NF- κ B. *PLoS Pathog.* 2010;6.

224. Sham HP, Shames SR, Croxson MA, et al. Attaching and effacing bacterial effector NleC suppresses epithelial inflammatory responses by inhibiting NF-kappaB and p38 mitogen-activated protein kinase activation. *Infect Immun*. 2011;79:3552–3562.
225. Tacket CO, Szein MB, Losonsky G, et al. Role of EspB in experimental human enteropathogenic *Escherichia coli* infection. *Infect Immun*. 2000;68:3689–3695.
226. Cleary J, Lai L-C, Donnenberg MS, et al. Enteropathogenic *E. coli* (EPEC) adhesion to intestinal epithelial cells: role of bundle-forming pili (BFP), EspA filaments and intimin. *Microbiology*. 2004;150:527–538.
227. Wong AR, Pearson JS, Bright MD, et al. Enteropathogenic and enterohaemorrhagic *Escherichia coli*: even more subversive elements. *Mol Microbiol*. 2011;80:1420–1438. Review. [Epub 2011 May 5].
228. Weflen AW, Alto NM, Hecht GA. Tight junctions and enteropathogenic *E. coli*. *Ann N Y Acad Sci*. 2009;1165:169–174.
229. Crane JK, McNamara BP, Donnenberg MS. Role of EspF in host cell death induced by enteropathogenic *Escherichia coli*. *Cell Microbiol*. 2001;3:197–211.
230. McNamara BP, Koutsouris A, O'Connell CB, et al. Translocated EspF protein from enteropathogenic *Escherichia coli* disrupts host intestinal barrier function. *J Clin Invest*. 2001;107:621–629.
231. Gill RK, Borthakur A, Hodges K, et al. Mechanism underlying inhibition of intestinal apical Cl/OH exchange following infection with enteropathogenic *E. coli*. *J Clin Invest*. 2007;117:428–437. [Epub 2007 Jan 25].
232. Aranda KR, Fagundes-Neto U, Scaletsky IC. Evaluation of multiplex PCRs for diagnosis of infection with diarrheagenic *Escherichia coli* and *Shigella* spp. *J Clin Microbiol*. 2004;42:5849–5853.
233. Rocha LB, Santos AR, Munhoz DD, et al. Development of a rapid agglutination latex test for diagnosis of enteropathogenic and enterohemorrhagic *Escherichia coli* infection in developing world: defining the biomarker, antibody and method. *PLoS Negl Trop Dis*. 2014;8:e3150, eCollection 2014 Sep.
234. Senerwa D, Mutanda LN, Gathuma JM, et al. Antimicrobial resistance of enteropathogenic *Escherichia coli* strains from a nosocomial outbreak in Kenya. *APMIS*. 1991;99:728–734.
235. Thorén A, Wolde-Mariam T, Stintzing G, et al. Antibiotics in the treatment of gastroenteritis caused by enteropathogenic *Escherichia coli*. *J Infect Dis*. 1980;141:27–31.
236. Antai SP, Anozie SO. Incidence of infantile diarrhoea due to enteropathogenic *Escherichia coli* in Port Harcourt metropolis. *J Appl Bacteriol*. 1987;62:227–229.
237. Vila J, Vargas M, Casals C, et al. Antimicrobial resistance of diarrheagenic *Escherichia coli* isolated from children under the age of 5 years from Ifakara, Tanzania. *Antimicrob Agents Chemother*. 1999;43:3022–3024.
238. Lim YS, Ngan CCL, Tay L. Enteropathogenic *Escherichia coli* as a cause of diarrhoea among children in Singapore. *J Trop Med Hyg*. 1992;95:339–342.
239. Huang Z, Pan H, Zhang P, et al. Prevalence and Antimicrobial Resistance Patterns of Diarrheagenic *Escherichia coli* in Shanghai, China. *Pediatr Infect Dis J*. 2016;35:835–839.
240. Subramanian K, Selvakumar C, Vinaykumar KS, et al. Tackling multiple antibiotic resistance in enteropathogenic *Escherichia coli* (EPEC) clinical isolates: a diarylheptanoid from *Alpinia officinarum* shows promising antibacterial and immunomodulatory activity against EPEC and its lipopolysaccharide-induced inflammation. *Int J Antimicrob Agents*. 2009;33:244–250. [Epub 2008 Dec 17].
241. Giles C, Sangster G, Smith J. Epidemic gastroenteritis of infants in Aberdeen during 1947. *Arch Dis Child*. 1949;24:45–53.
242. Robins-Browne R, Still CS, Miliotis MD, et al. Summer diarrhoea in African infants and children. *Arch Dis Child*. 1980;55:923–928.
243. Taylor J, Powell BW, Wright J. Infantile diarrhea and vomiting: a clinical and bacteriological investigation. *Br Med J*. 1949;2:117–141.
244. Camara LM, Carbonare SB, Silva MLM, et al. Inhibition of enteropathogenic *Escherichia coli* (EPEC) adhesion to HeLa cells by human colostrum: detection of specific sIgA related to EPEC outer-membrane proteins. *Int Arch Allergy Immunol*. 1994;103:307–310.
245. Cravioto A, Tello A, Villafán H, et al. Inhibition of localized adhesion of enteropathogenic *Escherichia coli* to HEp-2 cells by immunoglobulin and oligosaccharide fractions of human colostrum and breast milk. *J Infect Dis*. 1991;163:1247–1255.
246. Loureiro I, Frankel G, Adu-Bobie J, et al. Human colostrum contains IgA antibodies reactive to enteropathogenic *Escherichia coli* virulence-associated proteins: intimin, BfpA, EspA, and EspB. *J Pediatr Gastroenterol Nutr*. 1998;27:166–171.
247. Noguera-Obenza M, Ochoa TJ, Gomez HF, et al. Human milk secretory antibodies against attaching and effacing *Escherichia coli* antigens. *Emerg Infect Dis*. 2003;9:545–551.
248. Parissi-Crivelli A, Parissi-Crivelli JM, Girón JA. Recognition of enteropathogenic *Escherichia coli* virulence determinants by human colostrum and serum antibodies. *J Clin Microbiol*. 2000;38:2696–2700.
249. Costa-Carvalho BT, Bertipaglia A, Solé D, et al. Detection of immunoglobulin (IgG and IgA) anti-outer-membrane proteins of enteropathogenic *Escherichia coli* (EPEC) in saliva, colostrum, breast milk, serum, cord blood and amniotic fluid. Study of inhibition of localized adherence of EPEC to HeLa cells. *Acta Paediatr*. 1994;83:870–873.
250. Liu J, Wang WD, Liu YJ, et al. Mice vaccinated with enteropathogenic *Escherichia coli* ghosts show significant protection against lethal challenges. *Lett Appl Microbiol*. 2012;54:255–262.
251. Stakenborg T, Vandekerckhove D, Mariën J, et al. Protection of rabbits against enteropathogenic *Escherichia coli* (EPEC) using an intimin null mutant. *BMC Vet Res*. 2006;2:22.
252. de Souza Campos Fernandes RC, Quintana Flores VM, Sousa de Macedo Z, et al. Coproantibodies to the enteropathogenic *Escherichia coli* vaccine candidates BfpA and EspB in breastfed and artificially fed children. *Vaccine*. 2003;21:1725–1731.
253. Quintana Flores VM, Campos de Souza Fernandes RC, Sousa de Macedo Z, et al. Expression and purification of the recombinant enteropathogenic *Escherichia coli* vaccine candidates BfpA and EspB. *Protein Expr Purif*. 2002;25:16–22.
254. Donnenberg MS, Tacket CO, Losonsky G, et al. Effect of prior experimental human enteropathogenic *Escherichia coli* infection on illness following homologous and heterologous challenge. *Infect Immun*. 1998;66:52–58.
255. Blanco JE, Blanco M, Alonso MP, et al. Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from human patients: prevalence in Lugo, Spain, from 1992 through 1999. *J Clin Microbiol*. 2004;42:311–319.
256. Griffin PM, Ostroff SM, Tauxe RV, et al. Illnesses associated with *Escherichia coli* O157:H7 infections: a broad clinical spectrum. *Ann Intern Med*. 1988;109:705–712.
257. Garg AX, Suri RS, Barrowman N, et al. Long-term renal prognosis of diarrhea-associated hemolytic uremic syndrome—a systematic review, meta-analysis, and meta-regression. *JAMA*. 2003;290:1360–1370.
258. Tzipori S, Gunzer F, Donnenberg MS, et al. The role of the *eaeA* gene in diarrhea and neurological complications in a gnotobiotic piglet model of enterohemorrhagic *Escherichia coli* infection. *Infect Immun*. 1995;63:3621–3627.
259. Loos S, Aubert W, Hoppe B, et al. Intermediate follow-up of pediatric patients with hemolytic uremic syndrome during the 2011 outbreak caused by *E. coli* O104:H4. *Clin Infect Dis*. 2017;64:1637–1643.
260. Spinale JM, Ruebner RL, Copelovitch L, et al. Long-term outcomes of Shiga toxin hemolytic uremic syndrome. *Pediatr Nephrol*. 2013;28:2097–2105. Review. [Epub 2013 Jan 4].
261. Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiol Rev*. 1991;13:60–98.
262. Wong CS, Mooney JC, Brandt JR, et al. Risk factors for the hemolytic uremic syndrome in children infected with *Escherichia coli* O157:H7: a multivariable analysis. *Clin Infect Dis*. 2012;55:33–41. [Epub 2012 Mar 19].
263. Alpers K, Werber D, Frank C, et al. Sorbitol-fermenting enterohaemorrhagic *Escherichia coli* O157:H- causes another outbreak of haemolytic uraemic syndrome in children. *Epidemiol Infect*. 2009;137:389–395. [Epub 2009 Feb 9].
264. Rangel JM, Sparling PH, Crowe C, et al. Epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982–2002. *Emerg Infect Dis*. 2005;11:603–609.
265. Frank C, Werber D, Cramer JP, et al. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. *N Engl J Med*. 2011;365:1771–1780.
266. Menne J, Nitschke M, Stingle R, et al. EHEC-HUS consortium. Validation of treatment strategies for enterohaemorrhagic *Escherichia coli* O104:H4 induced haemolytic uraemic syndrome: case-control study. *BMJ*. 2012;345:e4565.
267. Schimmer B, Nygard K, Eriksen HM, et al. Outbreak of hemolytic uraemic syndrome in Norway caused by stx2-positive *Escherichia coli* O103:H25 traced to cured mutton sausages. *BMC Infect Dis*. 2008;8:41.
268. Johnson KE, Thorpe CM, Sears CL. The emerging clinical importance of non-O157 Shiga toxin-producing *Escherichia coli*. *Clin Infect Dis*. 2006;43:1587–1595. [Epub 2006 Nov 9].
269. Gonzalez AG, Cerqueira AM, Guth BE, et al. Serotypes, virulence markers and cell invasion ability of Shiga toxin-producing *Escherichia coli* strains isolated from healthy dairy cattle. *J Appl Microbiol*. 2016;121:1130–1143. [Epub 2016 Aug 25].
270. Cordeiro F, da Silva RI, Vargas-Stampe TL, et al. Cell invasion and survival of Shiga toxin-producing *Escherichia coli* within cultured human intestinal epithelial cells. *Microbiology*. 2013;159(Pt 8):1683–1694. [Epub 2013 May 23].
271. Matheus-Guimarães C, Gonçalves EM, Cabilio Guth BE. Interactions of O157 and non-O157 Shiga toxin-producing *Escherichia coli* (STEC) recovered from bovine hide and carcass with human cells and abiotic surfaces. *Foodborne Pathog Dis*. 2014;11:248–255. [Epub 2013 Dec 6].
272. Biscola FT, Abe CM, Guth BE. Determination of adhesin gene sequences in, and biofilm formation by, O157 and non-O157 Shiga toxin-producing *Escherichia coli* strains isolated from different sources. *Appl Environ Microbiol*. 2011;77:2201–2208. [Epub 2011 Feb 11].
273. Crowe SJ, Bottichio L, Shade LN, et al. Shiga toxin-producing *E. coli* infections associated with flour. *N Engl J Med*. 2017;377:2036–2043.
274. Jay MT, Cooley M, Carychao D, et al. *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle, central California coast. *Emerg Infect Dis*. 2007;13:1908–1911.
275. Olsen SJ, Miller G, Breuer T, et al. A waterborne outbreak of *Escherichia coli* O157:H7 infections and hemolytic uremic syndrome: implications for rural water systems. *Emerg Infect Dis*. 2002;8:370–375.
276. Crump JA, Sulka AC, Langer AJ, et al. An outbreak of *Escherichia coli* O157:H7 infections among visitors to a dairy farm. *N Engl J Med*. 2002;347:555–560.
277. Michino H, Araki K, Minami S, et al. Massive outbreak of *Escherichia coli* O157:H7 infection in schoolchildren in Sakai City, Japan, associated with consumption of white radish sprouts. *Am J Epidemiol*. 1999;150:787–796.
278. Bell BP, Goldoft M, Griffin PM, et al. A multistate outbreak of *Escherichia coli* O157:H7-associated bloody diarrhea and hemolytic uremic syndrome from hamburgers: the Washington experience. *JAMA*. 1994;272:1349–1353.
279. Tilden J Jr, Young W, McNamara AM, et al. A new route of transmission for *Escherichia coli*: infection from dry fermented salami. *Am J Public Health*. 1996;86:1142–1145.
280. Hale CR, Scallan E, Cronquist AB, et al. Estimates of enteric illness attributable to contact with animals and their environments in the United States. *Clin Infect Dis*. 2012;54(suppl 5):S472–S479.
281. Snedeker KG, Shaw DJ, Locking ME, et al. Primary and secondary cases in *Escherichia coli* O157 outbreaks: a statistical analysis. *BMC Infect Dis*. 2009;9:144.
282. Centers for Disease Control and Prevention (CDC). Outbreak of Shiga toxin-producing *Escherichia coli* O157 infection associated with a day camp petting zoo—Pinellas County, Florida, May–June 2007. *MMWR Morb Mortal Wkly Rep*. 2009;58:426–428.
283. Chase-Topping M, Gally D, Low C, et al. Super-shedding and the link between human infection and livestock carriage of *Escherichia coli* O157. *Nat Rev Microbiol*. 2008;6:904–912.
284. Solomon EB, Yaron S, Matthews KR. Transmission of *Escherichia coli* O157:H7 from contaminated manure and irrigation water to lettuce plant tissue and its subsequent internalization. *Appl Environ Microbiol*. 2002;68:397–400.
285. Tarr PI, Gordon CA, Chandler WL. Shiga-toxin-producing *Escherichia coli* and haemolytic uraemic syndrome. *Lancet*. 2005;365:1073–1086. Review.
286. Frank C, Werber D, Cramer JP, et al; HUS Investigation Team. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. *N Engl J Med*. 2011;365:1771–1780. [Epub 2011 Jun 22].
287. Serna A 4th, Boedeker EC. Pathogenesis and treatment of Shiga toxin-producing *Escherichia coli* infections. *Curr Opin Gastroenterol*. 2008;24:38–47. Review.
288. Brown JA, Hite DS, Gillim-Ross LA, et al. Outbreak of shiga toxin-producing *Escherichia coli* serotype O26: H11 infection at a child care center in Colorado. *Pediatr Infect Dis J*. 2012;31:379–383.
289. O'Brien AD, Newland JW, Miller SF, et al. Shiga-like toxin-converting phages from *Escherichia coli* strains that cause hemorrhagic colitis or infantile diarrhea. *Science*. 1984;226:694–696.
290. Sandvig K, Garred O, Prydz K, et al. Retrograde transport of endocytosed Shiga toxin to the endoplasmic reticulum. *Nature*. 1992;358:510–512.

291. Melton-Celsa AR. Shiga toxin (Stx) classification, structure, and function. *Microbiol Spectr*. 2014;2:EHEC-0024-2013. Review.
292. Tesh VL. Activation of cell stress response pathways by Shiga toxins. *Cell Microbiol*. 2012;14:1-9. Review. [Epub 2011 Sep 22].
293. Jones NL, Islur A, Haq R, et al. *Escherichia coli* Shiga toxins induce apoptosis in epithelial cells that is regulated by the Bcl-2 family. *Am J Physiol Gastrointest Liver Physiol*. 2000;278:G811-G819.
294. Te Loo DM, van Hinsbergh VW, van den Heuvel LP, et al. Detection of verocytotoxin bound to circulating polymorphonuclear leukocytes of patients with hemolytic uremic syndrome. *J Am Soc Nephrol*. 2001;12:800-806.
295. Yoshida T, Fukada M, Koide N, et al. Primary cultures of human endothelial cells are susceptible to low doses of Shiga toxins and undergo apoptosis. *J Infect Dis*. 1999;180:2048-2052.
296. Chandler WL, Jelacic S, Boster DR, et al. Prothrombotic coagulation abnormalities preceding the hemolytic-uremic syndrome. *N Engl J Med*. 2002;346:23-32.
297. Perna NT, Mayhew GF, Pósfai G, et al. Molecular evolution of a pathogenicity island from enterohemorrhagic *Escherichia coli* O157:H7. *Infect Immun*. 1998;66:3810-3817.
298. Curtis MM, Sperandio V. A complex relationship: the interaction among symbiotic microbes, invading pathogens, and their mammalian host. *Mucosal Immunol*. 2011;4:133-138. Review. [Epub 2011 Jan 19].
299. Mellies JL, Barron AM, Carmona AM. Enteropathogenic and enterohemorrhagic *Escherichia coli* virulence gene regulation. *Infect Immun*. 2007;75:4199-4210. Review. [Epub 2007 Jun 18].
300. Tobe T, Beatson SA, Taniguchi H, et al. An extensive repertoire of type III secretion effectors in *Escherichia coli* O157 and the role of lambdoid phages in their dissemination. *Proc Natl Acad Sci USA*. 2006;103:14941-14946. [Epub 2006 Sep 21].
301. Clougher P, Hurd S, Hoefler D, et al. Assessment of physician knowledge and practices concerning Shiga toxin-producing *Escherichia coli* infection and enteric illness, 2009, Foodborne Diseases Active Surveillance Network (FoodNet). *Clin Infect Dis*. 2012;54(suppl 5):S446-S452.
302. Marcon MJ. Point: should all stools be screened for Shiga toxin-producing *Escherichia coli*? *J Clin Microbiol*. 2011;49:2390-2394. [Epub 2011 May 4].
303. Gould LH, Bopp C, Strockbine N, et al; Centers for Disease Control and Prevention (CDC). Recommendations for diagnosis of shiga toxin-producing *Escherichia coli* infections by clinical laboratories. *MMWR Recomm Rep*. 2009;58(RR-12):1-14.
304. Centers for Disease Control and Prevention (CDC). Surveillance for foodborne disease outbreaks - United States, 2006. *MMWR Morb Mortal Wkly Rep*. 2009;58:609-615.
305. Zhang XP, McDaniel AD, Wolf LE, et al. Quinolone antibiotics induce Shiga toxin-encoding bacteriophages, toxin production, and death in mice. *J Infect Dis*. 2000;181:664-670.
306. Wong CS, Jelacic S, Habeeb RL, et al. The risk of the hemolytic-uremic syndrome after antibiotic treatment of *Escherichia coli* O157:H7 infections. *N Engl J Med*. 2000;342:1930-1936.
307. McGannon CM, Fuller CA, Weiss AA. Different classes of antibiotics differentially influence shiga toxin production. *Antimicrob Agents Chemother*. 2010;54:3790-3798. [Epub 2010 Jun 28].
308. Holtz LR, Neill MA, Tarr PI. Acute bloody diarrhea: a medical emergency for patients of all ages. *Gastroenterology*. 2009;136:1887-1898. Review. [Epub 2009 May 7].
309. Michael M, Elliott EJ, Ridley GF, et al. Interventions for haemolytic uraemic syndrome and thrombotic thrombocytopenic purpura. *Cochrane Database Syst Rev*. 2009;(1):CD003595.
310. Bell BP, Griffin PM, Lozano P, et al. Predictors of hemolytic uremic syndrome in children during a large outbreak of *Escherichia coli* O157:H7 infections. *Pediatrics*. 1997;100:E12.
311. Loos S, Ahlenstiel T, Kranz B, et al. An outbreak of Shiga toxin-producing *Escherichia coli* O104:H4 hemolytic uremic syndrome in Germany: presentation and short-term outcome in children. *Clin Infect Dis*. 2012;55:753-759. [Epub 2012 Jun 5].
312. Hickey CA, Beattie TJ, Cowieson J, et al. Early volume expansion during diarrhea and relative nephroprotection during subsequent hemolytic uremic syndrome. *Arch Pediatr Adolesc Med*. 2011;165:884-889. [Epub 2011 Jul 22].
313. Pape L, Hartmann H, Bange FC, et al. Eculizumab in typical hemolytic uremic syndrome (HUS) with neurological involvement. *Medicine (Baltimore)*. 2015;94:e1000.
314. Delmas Y, Vendrely B, Clouzeau B, et al. Outbreak of *Escherichia coli* O104:H4 haemolytic uraemic syndrome in France: outcome with eculizumab. *Nephrol Dial Transplant*. 2014;29:565-572. [Epub 2013 Nov 28].
315. Lapeyroue AL, Malina M, Fremeaux-Bacchi V, et al. Eculizumab in severe Shiga-toxin-associated HUS. *N Engl J Med*. 2011;364:2561-2563. Jun 30. [Epub 2011 May 25].
316. Barton Behravesh C, Jones TF, Vugia DJ, et al; FoodNet Working Group. Deaths associated with bacterial pathogens transmitted commonly through food: foodborne diseases active surveillance network (FoodNet), 1996-2005. *J Infect Dis*. 2011;204:263-267. Jul 15.
317. Chauvet C. Survival and control of *Escherichia coli* O157:H7 in foods, beverages, soil and water. *Virulence*. 2011;2:593-601. [Epub 2011 Nov 1].
318. Mukherjee J, Chios K, Fishwild D, et al. Human Stx2-specific monoclonal antibodies prevent systemic complications of *Escherichia coli* O157:H7 infection. *Infect Immun*. 2002;70:612-619.
319. Nishikawa K, Matsuoka K, Kita E, et al. A therapeutic agent with oriented carbohydrates for treatment of infections by Shiga toxin-producing *Escherichia coli* O157:H7. *Proc Natl Acad Sci USA*. 2002;99:7669-7674.
320. Armstrong GD, Rowe PC, Goodyer P, et al. A phase I study of chemically synthesized verotoxin (Shiga-like toxin) Pk-trisaccharide receptors attached to chromosorb for preventing hemolytic-uremic syndrome. *J Infect Dis*. 1995;171:1042-1045.
321. Mulvey GL, Marcato P, Kitov PI, et al. Assessment in mice of the therapeutic potential of tailored, multivalent Shiga toxin carbohydrate ligands. *J Infect Dis*. 2003;187:640-649.
322. Jeong KI, Tzipori S, Sheoran AS. Shiga toxin 2-specific but not Shiga toxin 1-specific human monoclonal antibody protects piglets challenged with enterohemorrhagic *Escherichia coli* producing Shiga toxin 1 and Shiga toxin 2. *J Infect Dis*. 2010;201:1081-1083.
323. López EL, Contrini MM, Glatstein E, et al. Safety and pharmacokinetics of urtoxazumab, a humanized monoclonal antibody, against Shiga-like toxin 2 in healthy adults and in pediatric patients infected with Shiga-like toxin-producing *Escherichia coli*. *Antimicrob Agents Chemother*. 2010;54:239-243. [Epub 2009 Oct 12].
324. Bitzan M, Poole R, Mehran M, et al. Safety and pharmacokinetics of chimeric anti-Shiga toxin 1 and anti-Shiga toxin 2 monoclonal antibodies in healthy volunteers. *Antimicrob Agents Chemother*. 2009;53:3081-3087. [Epub 2009 May 4].
325. Dowling TC, Chavaille PA, Young DG, et al. Phase I safety and pharmacokinetic study of chimeric murine-human monoclonal antibody c alpha Stx2 administered intravenously to healthy adult volunteers. *Antimicrob Agents Chemother*. 2005;49:1808-1812.
326. Trachtman H, Cnaan A, Christen E, et al. Effect of an oral Shiga toxin-binding agent on diarrhea-associated hemolytic uremic syndrome in children—a randomized controlled trial. *JAMA*. 2003;290:1337-1344.
327. Ahmed A, Li J, Shiloach Y, et al. Safety and immunogenicity of *Escherichia coli* O157 O-specific polysaccharide conjugate vaccine in 2-5-year-old children. *J Infect Dis*. 2006;193:515-521.
328. Mayr UB, Haller C, Haidinger W, et al. Bacterial ghosts as an oral vaccine: a single dose of *Escherichia coli* O157:H7 bacterial ghosts protects mice against lethal challenge. *Infect Immun*. 2005;73:4810-4817.
329. Potter AA, Klashinsky S, Li Y, et al. Decreased shedding of *Escherichia coli* O157:H7 by cattle following vaccination with type III secreted proteins. *Vaccine*. 2004;22:362-369.
330. Dean-Nystrom EA, Gansheroff LJ, Mills M, et al. Vaccination of pregnant dams with intimin O157 protects suckling piglets from *Escherichia coli* O157:H7 infection. *Infect Immun*. 2002;70:2414-2418.
331. Konadu E, Donohue-Rolfe A, Calderwood SB, et al. Syntheses and immunologic properties of *Escherichia coli* O157 O-specific polysaccharide and Shiga toxin 1 B subunit conjugates in mice. *Infect Immun*. 1999;67:6191-6193.
332. Lu X, Skurnik D, Pozzi C, et al. A Poly-N-acetylglucosamine-Shiga toxin broad-spectrum conjugate vaccine for Shiga toxin-producing *Escherichia coli*. *MBio*. 2014;5:e00974-14.
333. Cai K, Tu W, Liu Y, et al. Novel fusion antigen displayed-bacterial ghosts vaccine candidate against infection of *Escherichia coli* O157:H7. *Sci Rep*. 2015;5:17479.
334. Paton AW, Chen AY, Wang H, et al. Protection against Shiga-toxicogenic *Escherichia coli* by non-genetically modified organism receptor mimic bacterial ghosts. *Infect Immun*. 2015;83:3526-3533. [Epub 2015 Jun 22].
335. Nataro JP, Kaper JB, Robins-Browne R, et al. Patterns of adherence of diarrheagenic *Escherichia coli* to HEp-2 cells. *Pediatr Infect Dis J*. 1987;6:829-831.
336. Cravioto A, Tello A, Navarro A, et al. Association of *Escherichia coli* HEp-2 adherence patterns with type and duration of diarrhoea. *Lancet*. 1991;337:262-264.
337. Yamamoto T, Koyama Y, Matsumoto M, et al. Localized, aggregative, and diffuse adherence to HeLa cells, plastic, and human small intestines by *Escherichia coli* isolated from patients with diarrhea. *J Infect Dis*. 1992;166:1295-1310.
338. Nataro JP, Hicks S, Phillips AD, et al. T84 cells in culture as a model for enteroregulative *Escherichia coli* pathogenesis. *Infect Immun*. 1996;64:4761-4768.
339. Huang DB, Nataro JP, DuPont HL, et al. Enteroregulative *Escherichia coli* is a cause of acute diarrheal illness: a meta-analysis. *Clin Infect Dis*. 2006;43:556-563.
340. Hebbelstrup Jensen B, Olsen KE, Struve C, et al. Epidemiology and clinical manifestations of enteroregulative *Escherichia coli*. *Clin Microbiol Rev*. 2014;27:614-630. Review.
341. Nataro JP, Mai V, Johnson J, et al. Diarrheagenic *Escherichia coli* infection in Baltimore, Maryland, and New Haven, Connecticut. *Clin Infect Dis*. 2006;43:402-407. [Epub 2006 Jul 11].
342. Nataro JP, Yikang D, Cookson S, et al. Heterogeneity of enteroregulative *Escherichia coli* virulence demonstrated in volunteers. *J Infect Dis*. 1995;171:465-468.
343. Okeke IN, Lamikanra A, Czeizulin J, et al. Heterogeneous virulence of enteroregulative *Escherichia coli* strains isolated from children in southwest Nigeria. *J Infect Dis*. 2000;181:252-260.
344. Czeizulin JR, Whittam TS, Henderson IR, et al. Phylogenetic analysis of enteroregulative and diffusely adherent *Escherichia coli*. *Infect Immun*. 1999;67:2692-2699.
345. Buchholz U, Bernard H, Werber D, et al. German outbreak of *Escherichia coli* O104:H4 associated with sprouts. *N Engl J Med*. 2011;365:1763-1770. [Epub 2011 Oct 26].
346. Gault G, Weill FX, Mariani-Kurkdjian P, et al. Outbreak of haemolytic uraemic syndrome and bloody diarrhoea due to *Escherichia coli* O104:H4, south-west France, June 2011. *Euro Surveill*. 2011;16: pii: 19905.
347. Adachi JA, Ericsson CD, Jiang ZD, et al. Natural history of enteroregulative and enterotoxigenic *Escherichia coli* infection among US travelers to Guadalajara, Mexico. *J Infect Dis*. 2002;185:1681-1683.
348. Adachi JA, Jiang ZD, Mathewson JJ, et al. Enteroregulative *Escherichia coli* as a major etiologic agent in traveler's diarrhea in 3 regions of the world. *Clin Infect Dis*. 2001;32:1706-1709.
349. Dallman TJ, Chattaway MA, Cowley LA, et al. An investigation of the diversity of strains of enteroregulative *Escherichia coli* isolated from cases associated with a large multi-pathogen foodborne outbreak in the UK. *PLoS ONE*. 2014;9:e98103. eCollection 2014.
350. Okhuysen PC, Jiang ZD, Carlin L, et al. Post-diarrhea chronic intestinal symptoms and irritable bowel syndrome in North American travelers to Mexico. *Am J Gastroenterol*. 2004;99:1774-1778.
351. Itoh Y, Nagano I, Kunishima M, et al. Laboratory investigation of enteroregulative *Escherichia coli* O untypeable:H10 associated with a massive outbreak of gastrointestinal illness. *J Clin Microbiol*. 1997;35:2546-2550.
352. Bhatnagar S, Bhan MK, Sommerfelt H, et al. Enteroregulative *Escherichia coli* may be a new pathogen causing acute and persistent diarrhea. *Scand J Infect Dis*. 1993;25:579-583.
353. Wanke CA, Schorling JB, Barrett LJ, et al. Potential role of adherence traits of *Escherichia coli* in persistent diarrhea in an urban Brazilian slum. *Pediatr Infect Dis J*. 1991;10:746-751.
354. Bhan MK, Raj P, Levine MM, et al. Enteroregulative *Escherichia coli* associated with persistent diarrhea in a cohort of rural children in India. *J Infect Dis*. 1989;159:1061-1064.
355. Gassama-Sow A, Sow PS, Guèye M, et al. Characterization of pathogenic *Escherichia coli* in human immunodeficiency virus-related diarrhea in Senegal. *J Infect Dis*. 2004;189:75-78.
356. Durrer P, Zbinden R, Fleisch E, et al. Intestinal infection due to enteroregulative *Escherichia coli* among human immunodeficiency virus-infected persons. *J Infect Dis*. 2000;182:1540-1544.
357. Wanke CA, Mayer H, Weber R, et al. Enteroregulative *Escherichia coli* as a potential cause of diarrheal disease in adults infected with human immunodeficiency virus. *J Infect Dis*. 1998;178:185-190.

358. Fang GD, Lima AA, Martins CV, et al. Etiology and epidemiology of persistent diarrhea in northeastern Brazil: a hospital-based, prospective, case-control study. *J Pediatr Gastroenterol Nutr.* 1995;21:137–144.
359. Steiner TS, Lima AAM, Nataro JP, et al. Enterotoxigenic *Escherichia coli* produce intestinal inflammation and growth impairment and cause interleukin-8 release from intestinal epithelial cells. *J Infect Dis.* 1998;177:88–96.
360. Rogawski ET, Guerrant RL, Havt A, et al. MAL-ED Network Investigators. Epidemiology of enterotoxigenic *Escherichia coli* infections and associated outcomes in the MAL-ED birth cohort. *PLoS Negl Trop Dis.* 2017;11:e0005798, eCollection 2017 Jul.
361. Khan K, Konar M, Goyal A, et al. Enterotoxigenic *Escherichia coli* infection induces IL-8 production via activation of mitogen-activated protein kinases and the transcription factors NF- κ B and AP-1 in INT-407 cells. *Mol Cell Biochem.* 2010;337:17–24. [Epub 2009 Oct 24].
362. Mohamed JA, DuPont HL, Flores J, et al. Single nucleotide polymorphisms in the promoter of the gene encoding the lipopolysaccharide receptor CD14 are associated with bacterial diarrhea in US and Canadian travelers to Mexico. *Clin Infect Dis.* 2011;52:1332–1341.
363. Mohamed JA, DuPont HL, Jiang ZD, et al. A single-nucleotide polymorphism in the gene encoding osteoprotegerin, an anti-inflammatory protein produced in response to infection with diarrheagenic *Escherichia coli*, is associated with an increased risk of nonsecretory bacterial diarrhea in North American travelers to Mexico. *J Infect Dis.* 2009;199:477–485.
364. Mohamed JA, DuPont HL, Jiang ZD, et al. A novel single-nucleotide polymorphism in the lactoferrin gene is associated with susceptibility to diarrhea in North American travelers to Mexico. *Clin Infect Dis.* 2007;44:945–952. [Epub 2007 Feb 20].
365. Jiang ZD, Okhuysen PC, Guo DC, et al. Genetic susceptibility to enterotoxigenic *Escherichia coli* diarrhea: polymorphism in the interleukin-8 promoter region. *J Infect Dis.* 2003;188:506–511. [Epub 2003 Jul 25].
366. Nataro JP, Deng Y, Maneval DR, et al. Aggregative adherence fimbriae I of enterotoxigenic *Escherichia coli* mediate adherence to HEp-2 cells and hemagglutination of human erythrocytes. *Infect Immun.* 1992;60:2297–2304.
367. Czeizulin JR, Balepur S, Hicks S, et al. Aggregative adherence fimbria II, a second fimbrial antigen mediating aggregative adherence in enterotoxigenic *Escherichia coli*. *Infect Immun.* 1997;65:4135–4145.
368. Boisen N, Struve C, Scheutz F, et al. New adhesin of enterotoxigenic *Escherichia coli* related to the Afa/Dtr/AAF family. *Infect Immun.* 2008;76:3281–3292.
369. Harrington SM, Strauman MC, Abe CM, et al. Aggregative adherence fimbriae contribute to the inflammatory response of epithelial cells infected with enterotoxigenic *Escherichia coli*. *Cell Microbiol.* 2005;7:1565–1578.
370. Sheikh J, Czeizulin JR, Harrington S, et al. A novel dispersin protein in enterotoxigenic *Escherichia coli*. *J Clin Invest.* 2002;110:1329–1337.
371. Morin N, Santiago AE, Ernst RK, et al. Characterization of the AggR regulon in enterotoxigenic *Escherichia coli*. *Infect Immun.* 2013;81:122–132. [Epub 2012 Oct 22].
372. Jiang ZD, Greenberg D, Nataro JP, et al. Rate of occurrence and adherence effect of enterotoxigenic *Escherichia coli* virulence factors in international travelers. *J Clin Microbiol.* 2002;40:4185–4190.
373. Huang DB, Mohamed JA, Nataro JP, et al. Virulence characteristics and the molecular epidemiology of enterotoxigenic *Escherichia coli* isolates from travellers to developing countries. *J Med Microbiol.* 2007;56(Pt 10):1386–1392.
374. Schiebel J, Böhm A, Nitschke J, et al. Genotypic and phenotypic characteristics associated with biofilm formation by human clinical *Escherichia coli* isolates of different pathotypes. *Appl Environ Microbiol.* 2017;83: pii: e01660-17.
375. Kong H, Hong X, Li X. Current perspectives in pathogenesis and antimicrobial resistance of enterotoxigenic *Escherichia coli*. *Microb Pathog.* 2015;85:44–49, Review. [Epub 2015 Jun 6].
376. Sheikh J, Hicks S, Dall'Agnol M, et al. Roles for Fis and YafK in biofilm formation by enterotoxigenic *Escherichia coli*. *Mol Microbiol.* 2001;41:983–997.
377. Navarro-García F, Gutierrez-Jimenez J, Garcia-Tovar C, et al. Pic, an autotransporter protein secreted by different pathogens in the Enterobacteriaceae family, is a potent mucus secretagogue. *Infect Immun.* 2010;78:4101–4109. [Epub 2010 Aug 9].
378. Savarino SJ, Fasano A, Robertson DC, et al. Enterotoxigenic *Escherichia coli* elaborate a heat-stable enterotoxin demonstrable in an in vitro rabbit intestinal model. *J Clin Invest.* 1991;87:1450–1455.
379. Savarino SJ, Fasano A, Watson J, et al. Enterotoxigenic *Escherichia coli* heat-stable enterotoxin 1 represents another subfamily of *E. coli* heat-stable toxin. *Proc Natl Acad Sci USA.* 1993;90:3093–3097.
380. Fleckenstein JM, Hardwidge PR, Munson GP, et al. Molecular mechanisms of enterotoxigenic *Escherichia coli* infection. *Microbes Infect.* 2010;12:89–98, Review. [Epub 2009 Oct 31].
381. Eslava C, Navarro-García F, Czeizulin JR, et al. Pet, an auto-transporter enterotoxin from enterotoxigenic *Escherichia coli*. *Infect Immun.* 1998;66:3155–3163.
382. Henderson IR, Hicks S, Navarro-García F, et al. Involvement of the enterotoxigenic *Escherichia coli* plasmid-encoded toxin in causing human intestinal damage. *Infect Immun.* 1999;67:5338–5344.
383. Iwanaga M, Song T, Higa N, et al. Enterotoxigenic *Escherichia coli*: incidence in Japan and usefulness of the clump-formation test. *J Infect Chemother.* 2002;8:345–348.
384. Albert MJ, Qadri F, Haque A, et al. Bacterial dump formation at the surface of liquid culture as a rapid test for identification of enterotoxigenic *Escherichia coli*. *J Clin Microbiol.* 1993;31:1397–1399.
385. Jenkins C, Tembo M, Chart H, et al. Detection of enterotoxigenic *Escherichia coli* in faecal samples from patients in the community with diarrhoea. *J Med Microbiol.* 2006;55(Pt 11):1493–1497.
386. Glandt M, Adachi JA, Mathewson JJ, et al. Enterotoxigenic *Escherichia coli* as a cause of traveler's diarrhea: clinical response to ciprofloxacin. *Clin Infect Dis.* 1999;29:335–338.
387. Wanke CA, Gerrior J, Blais V, et al. Successful treatment of diarrheal disease associated with enterotoxigenic *Escherichia coli* in adults infected with human immunodeficiency virus. *J Infect Dis.* 1998;178:1369–1372.
388. Nitschke M, Sayk F, Härtel C, et al. Association between azithromycin therapy and duration of bacterial shedding among patients with Shiga toxin-producing enterotoxigenic *Escherichia coli* O104:H4. *JAMA.* 2012;307:1046–1052.
389. Seifert ME, Tarr PI. Azithromycin decolonization of STEC—a new risk emerges. *Nat Rev Nephrol.* 2012;8:429.
390. Seifert ME, Tarr PI. Therapy: azithromycin and decolonization after HUS. *Nat Rev Nephrol.* 2012;8:317–318.
391. Chattaway MA, Jenkins C, Rajendram D, et al. Enterotoxigenic *Escherichia coli* have evolved independently as distinct complexes within the *E. coli* population with varying ability to cause disease. *PLoS ONE.* 2014;9:e112967, eCollection 2014.
392. Do Nascimento V, Day MR, Doumth M, et al. Comparison of phenotypic and WGS-derived antimicrobial resistance profiles of enterotoxigenic *Escherichia coli* isolated from cases of diarrhoeal disease in England, 2015–16. *J Antimicrob Chemother.* 2017;72:3288–3297.
393. Frech SA, Dupont HL, Bourgeois AL, et al. Use of a patch containing heat-labile toxin from *Escherichia coli* against travellers' diarrhoea: a phase II, randomised, double-blind, placebo-controlled field trial. *Lancet.* 2008;371:2019–2025.
394. Wirth T, Falush D, Lan R, et al. Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol.* 2006;60:1136–1151.
395. Martinez MB, Whittam TS, McGraw EA, et al. Clonal relationship among invasive and non-invasive strains of enteroinvasive *Escherichia coli* serogroups. *FEMS Microbiol Lett.* 1999;172:145–151.
396. Prosseda G, Di Martino ML, Campilongo R, et al. Shedding of genes that interfere with the pathogenic lifestyle: the Shigella model. *Res Microbiol.* 2012;163:399–406, Review. [Epub 2012 Jul 20].
397. Lan R, Reeves PR. *Escherichia coli* in disguise: molecular origins of Shigella. *Microbes Infect.* 2002;4:1125–1132, Review.
398. Maurelli AT, Fernandez RE, Bloch CA, et al. “Black holes” and bacterial pathogenicity: a large genomic deletion that enhances the virulence of *Shigella* spp. and enteroinvasive *Escherichia coli*. *Proc Natl Acad Sci USA.* 1998;95:3943–3948.
399. DuPont HL, Formal SB, Hornick RB, et al. Pathogenesis of *Escherichia coli* diarrhea. *N Engl J Med.* 1971;285:1–9.
400. Ewing WH, Gravatt JL. Shigella types encountered in the Mediterranean area. *J Bacteriol.* 1947;53:191–195.
401. Lozer DM, Souza TB, Monfardini MV, et al. Genotypic and phenotypic analysis of diarrheagenic *Escherichia coli* strains isolated from Brazilian children living in low socioeconomic level communities. *BMC Infect Dis.* 2013;13:418.
402. Moreno AC, Filho AF, Gomes Tdo A, et al. Etiology of childhood diarrhea in the northeast of Brazil: significant emergent diarrheal pathogens. *Diagn Microbiol Infect Dis.* 2010;66:50–57. [Epub 2008 May 27].
403. Formal SB, Hornick RB. Invasive *Escherichia coli*. *J Infect Dis.* 1978;137:641–644.
404. Sakazaki R, Tamura K, Saito M. Enteropathogenic *Escherichia coli* associated with diarrhea in children and adults. *Jpn J Med Sci Biol.* 1967;20:387–399.
405. Trabulsi LR, Manissadjian A, Penna HA, et al. [Infantile diarrhea caused by enteropathogenic colibacilli. Preliminary studies on the occurrence of certain groups and serological types in São Paulo]. *Rev Inst Med Trop São Paulo.* 1961;3:267–270, Portuguese.
406. Newitt S, MacGregor V, Robbins V, et al. Two linked enteroinvasive *Escherichia coli* outbreaks, Nottingham, UK, June 2014. *Emerg Infect Dis.* 2016;22:1178–1184.
407. Michelacci V, Prosseda G, Maugliani A, et al. Characterization of an emergent clone of enteroinvasive *Escherichia coli* circulating in Europe. *Clin Microbiol Infect.* 2016;22:287, e11–9. [Epub 2015 Nov 10].
408. Harris JR, Mariano J, Wells JG, et al. Person-to-person transmission in an outbreak of enteroinvasive *Escherichia coli*. *Am J Epidemiol.* 1985;122:245–252.
409. Day WA, Maurelli AT. Shigella and enteroinvasive *Escherichia coli*: paradigms for pathogen evolution and host-parasite interactions. In: Donnenberg MS, ed. *Escherichia coli: Virulence Mechanisms of a Versatile Pathogen*. San Diego: Academic Press; 2002:209–237.
410. Formal SB, Hale TL, Sansonetti PJ. Invasive enteric pathogens. *Rev Infect Dis.* 1983;5(suppl 4):S702–S707, Review.
411. Harris JR, Wachsmuth IK, Davis BR, et al. High-molecular-weight plasmid correlates with *Escherichia coli* enteroinvasiveness. *Infect Immun.* 1982;37:1295–1298.
412. Bando SY, Moreno AC, Albuquerque JA, et al. Expression of bacterial virulence factors and cytokines during in vitro macrophage infection by enteroinvasive *Escherichia coli* and Shigella flexneri: a comparative study. *Mem Inst Oswaldo Cruz.* 2010;105:786–791.
413. Moreno AC, Ferreira LG, Martinez MB. Enteroinvasive *Escherichia coli* vs. Shigella flexneri: how different patterns of gene expression affect virulence. *FEMS Microbiol Lett.* 2009;301:156–163. [Epub 2009 Oct 7].
414. Silva RM, Toledo MR, Trabulsi LR. Biochemical and cultural characteristics of invasive *Escherichia coli*. *J Clin Microbiol.* 1980;11:441–444.
415. van den Beld MJ, Reubsaet FA. Differentiation between Shigella, enteroinvasive *Escherichia coli* (EIEC) and noninvasive *Escherichia coli*. *Eur J Clin Microbiol Infect Dis.* 2012;31:899–904, Review. [Epub 2011 Sep 7].
416. DuPont HL. Clinical practice. Bacterial diarrhea. *N Engl J Med.* 2009;361:1560–1569, Review.
417. Ahmed AM, Miyoshi S, Shinoda S, et al. Molecular characterization of a multidrug-resistant strain of enteroinvasive *Escherichia coli* O164 isolated in Japan. *J Med Microbiol.* 2005;54(Pt 3):273–278.
418. Gomes TA, Vieira MA, Abe CM, et al. Adherence patterns and adherence-related DNA sequences in *Escherichia coli* isolates from children with and without diarrhea in São Paulo city, Brazil. *J Clin Microbiol.* 1998;36:3609–3613.
419. Tacket CO, Moseley SL, Kay B, et al. Challenge studies in volunteers using *Escherichia coli* strains with diffuse adherence to HEp-2 cells. *J Infect Dis.* 1990;162:550–552.
420. Germani Y, Begaud E, Duval P, et al. Prevalence of enteropathogenic, enterotoxigenic, and diffusely adherent *Escherichia coli* among isolates from children with diarrhea in New Caledonia. *J Infect Dis.* 1996;174:1124–1126.
421. Gunzburg ST, Chang BJ, Elliott SJ, et al. Diffuse and enterotoxigenic patterns of adherence of enteric *Escherichia coli* isolated from aboriginal children from the Kimberley region of Western Australia. *J Infect Dis.* 1993;167:755–758.
422. Lopes LM, Fabbriotti SH, Ferreira AJ, et al. Heterogeneity among strains of diffusely adherent *Escherichia coli* isolated in Brazil. *J Clin Microbiol.* 2005;43:1968–1972.
423. Scaletsky IC, Fabbriotti SH, Carvalho RL, et al. Diffusely adherent *Escherichia coli* as a cause of acute diarrhea in young children in Northeast Brazil: a case-control study. *J Clin Microbiol.* 2002;40:645–648.
424. Knutton S, Shaw R, Phillips AD, et al. Phenotypic and genetic analysis of diarrhea-associated *Escherichia coli* isolated from children in the United Kingdom. *J Pediatr Gastroenterol Nutr.* 2001;33:32–40.
425. Servin AL. Pathogenesis of human diffusely adhering *Escherichia coli* expressing Afa/Dr adhesins (Afa/Dr DAEC): current insights and future challenges. *Clin Microbiol Rev.* 2014;27:823–869, Review.

426. Tenaillon O, Skurnik D, Picard B, et al. The population genetics of commensal *Escherichia coli*. *Nat Rev Microbiol*. 2010;8:207–217. Mar. Review.
427. Snelling AM, Macfarlane-Smith LR, Fletcher JN, et al. The commonly-used DNA probe for diffusely-adherent *Escherichia coli* cross-reacts with a subset of enteroaggregative *E. coli*. *BMC Microbiol*. 2009;9:269.
428. Le Bouguéneq C, Servin AL. Diffusely adherent *Escherichia coli* strains expressing Afa/Dr adhesins (Afa/Dr DAEC): hitherto unrecognized pathogens. *FEMS Microbiol Lett*. 2006;256:185–194. Review.
429. Labigne-Roussel AF, Lark D, Schoolnik G, et al. Cloning and expression of an afimbrial adhesin (AFA-I) responsible for P blood group-independent, mannose-resistant hemagglutination from a pyelonephritic *Escherichia coli* strain. *Infect Immun*. 1984;46:251–259.
430. Le Bouguéneq C, Lalioui L, du Merle L, et al. Characterization of AfaE adhesins produced by extraintestinal and intestinal human *Escherichia coli* isolates: PCR assays for detection of Afa adhesins that do or do not recognize Dr blood group antigens. *J Clin Microbiol*. 2001;39:1738–1745.
431. Servin AL. Pathogenesis of Afa/Dr diffusely adhering *Escherichia coli*. *Clin Microbiol Rev*. 2005;18:264–292. Review.
432. Peiffer I, Guignot J, Barbat A, et al. Structural and functional lesions in brush border of human polarized intestinal Caco-2/TC7 cells infected by members of the Afa/Dr diffusely adhering family of *Escherichia coli*. *Infect Immun*. 2000;68:5979–5990.
433. Peiffer I, Blanc-Potard AB, Bernet-Camard MF, et al. Afa/Dr diffusely adhering *Escherichia coli* C1845 infection promotes selective injuries in the junctional domain of polarized human intestinal Caco-2/TC7 cells. *Infect Immun*. 2000;68:3431–3442.
434. Goluszko P, Selvarangan R, Popov V, et al. Decay-accelerating factor and cytoskeleton redistribution pattern in HeLa cells infected with recombinant *Escherichia coli* strains expressing Dr family of adhesins. *Infect Immun*. 1999;67:3989–3997.
435. Peiffer I, Servin AL, Bernet-Camard MF. Piracy of decay-accelerating factor (CD55) signal transduction by the diffusely adhering strain *Escherichia coli* C1845 promotes cytoskeletal F-actin rearrangements in cultured human intestinal INT407 cells. *Infect Immun*. 1998;66:4036–4042.
436. Bernet-Camard MF, Coconnier MH, Hudault S, et al. Pathogenicity of the diffusely adhering strain *Escherichia coli* C1845: F1845 adhesin-decay accelerating factor interaction, brush border microvilli injury, and actin disassembly in cultured human intestinal epithelial cells. *Infect Immun*. 1996;64:1918–1928.
437. Nowicki B, Moulds J, Hull R, et al. A hemagglutinin of uropathogenic *Escherichia coli* recognizes the Dr blood group antigen. *Infect Immun*. 1988;56:1057–1060.
438. Arikawa K, Nishikawa Y. Interleukin-8 induction due to diffusely adherent *Escherichia coli* possessing Afa/Dr genes depends on flagella and epithelial Toll-like receptor 5. *Microbiol Immunol*. 2010;54:491–501.
439. Bétis F, Brest P, Hoffman V, et al. Afa/Dr diffusely adhering *Escherichia coli* infection in T84 cell monolayers induces increased neutrophil transepithelial migration, which in turn promotes cytokine-dependent upregulation of decay-accelerating factor (CD55), the receptor for Afa/Dr adhesins. *Infect Immun*. 2003;71:1774–1783.
440. Bétis F, Brest P, Hoffman V, et al. The Afa/Dr adhesins of diffusely adhering *Escherichia coli* stimulate interleukin-8 secretion, activate mitogen-activated protein kinases, and promote polymorphonuclear transepithelial migration in T84 polarized epithelial cells. *Infect Immun*. 2003;71:1068–1074.
441. Darfeuille-Michaud A, Neut C, Barnich N, et al. Presence of adherent *Escherichia coli* strains in ileal mucosa of patients with Crohn's disease. *Gastroenterology*. 1998;115:1405–1413.
442. Berstad AE, Brandtzaeg P. Expression of cell membrane complement regulatory glycoproteins along the normal and diseased human gastrointestinal tract. *Gut*. 1998;42:522–529.
443. Boudeau J, Glasser AL, Masseret E, et al. Invasive ability of an *Escherichia coli* strain isolated from the ileal mucosa of a patient with Crohn's disease. *Infect Immun*. 1999;67:4499–4509.
444. Mann EA, Saeed SA. Gastrointestinal infection as a trigger for inflammatory bowel disease. *Curr Opin Gastroenterol*. 2012;28:24–29. Review.
445. Spehlmann ME, Begun AZ, Burghardt J, et al. Epidemiology of inflammatory bowel disease in a German twin cohort: results of a nationwide study. *Inflamm Bowel Dis*. 2008;14:968–976.
446. Smith EJ, Thompson AP, O'Driscoll A, et al. Pathogenesis of adherent-invasive *Escherichia coli*. *Future Microbiol*. 2013;8:1289–1300. Review.
447. Darfeuille-Michaud A, Boudeau J, Bulois P, et al. High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology*. 2004;127:412–421.
448. Martin HM, Campbell BJ, Hart CA, et al. Enhanced *Escherichia coli* adherence and invasion in Crohn's disease and colon cancer. *Gastroenterology*. 2004;127:80–93.
449. O'Brien CL, Bringer MA, Holt KE, et al. Comparative genomics of Crohn's disease-associated adherent-invasive *Escherichia coli*. *Gut*. 2017;66:1382–1389. [Epub 2016 Apr 20].
450. Kotlowski R, Bernstein CN, Sepehri S, et al. High prevalence of *Escherichia coli* belonging to the B2+D phylogenetic group in inflammatory bowel disease. *Gut*. 2007;56:669–675. [Epub 2006 Oct 6].
451. Johnson JR, Delavari P, Kuskowski M, et al. Phylogenetic distribution of extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis*. 2001;183:78–88. [Epub 2000 Nov 22]. Erratum in: *J Infect Dis* 2001 Jun 15;183(12):1834–5.
452. Bingen E, Picard B, Brahimi N, et al. Phylogenetic analysis of *Escherichia coli* strains causing neonatal meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J Infect Dis*. 1998;177:642–650.
453. Chassaing B, Rolhion N, de Vallée A, et al. Crohn disease-associated adherent-invasive *E. coli* bacteria target mouse and human Peyer's patches via long polar fimbriae. *J Clin Invest*. 2011;121:966–975. [Epub 2011 Feb 21].
454. Barnich N, Darfeuille-Michaud A. Abnormal CEACAM6 expression in Crohn disease patients favors gut colonization and inflammation by adherent-invasive *E. coli*. *Virulence*. 2010;1:281–282.
455. Rolhion N, Barnich N, Bringer MA, et al. Abnormally expressed ER stress response chaperone Gp96 in CD favours adherent-invasive *Escherichia coli* invasion. *Gut*. 2010;59:1355–1362. [Epub 2010 Jun 29].
456. Barnich N, Carvalho FA, Glasser AL, et al. CEACAM6 acts as a receptor for adherent-invasive *E. coli*, supporting ileal mucosa colonization in Crohn disease. *J Clin Invest*. 2007;117:1566–1574. [Epub 2007 May 24].
457. Bringer MA, Glasser AL, Tung CH, et al. The Crohn's disease-associated adherent-invasive *Escherichia coli* strain LF82 replicates in mature phagolysosomes within I774 macrophages. *Cell Microbiol*. 2006;8:471–484.
458. Glasser AL, Boudeau J, Barnich N, et al. Adherent invasive *Escherichia coli* strains from patients with Crohn's disease survive and replicate within macrophages without inducing host cell death. *Infect Immun*. 2001;69:5529–5537.
459. Pranter C, Scribano ML. Antibiotics and probiotics in inflammatory bowel disease: why, when, and how. *Curr Opin Gastroenterol*. 2009;25:329–333. Review.
460. Picard B, Garcia JS, Gouriou S, et al. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun*. 1999;67:546–553.
461. Hacker J, Blum-Oehler G, Mühlendorfer I, et al. Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Mol Microbiol*. 1997;23:1089–1097. Review.
462. Bidet P, Mahjoub-Messai F, Blanco J, et al. Combined multilocus sequence typing and O serogrouping distinguishes *Escherichia coli* subtypes associated with infant urosepsis and/or meningitis. *J Infect Dis*. 2007;196:297–303. [Epub 2007 Jun 4].
463. Kunin CM, Hua TH, Krishnan C, et al. Isolation of a nicotinamide-requiring clone of *Escherichia coli* O18:K1:H7 from women with acute cystitis: resemblance to strains found in neonatal meningitis. *Clin Infect Dis*. 1993;16:412–416.
464. Johnson JR, Russo TA. Uropathogenic *Escherichia coli* as agents of diverse non-urinary tract extraintestinal infections. *J Infect Dis*. 2002;186:859–864.
465. Johnson JR, Delavari P, O'Bryan TT. *Escherichia coli* O18:K1:H7 isolates from patients with acute cystitis and neonatal meningitis exhibit common phylogenetic origins and virulence factor profiles. *J Infect Dis*. 2001;183:425–434.
466. Benton J, Chawla J, Parry S, et al. Virulence factors in *Escherichia coli* from urinary tract infections in patients with spinal injuries. *J Hosp Infect*. 1992;22:117–127.
467. Johnson JR, Roberts PL, Stamm WE. P fimbriae and other virulence factors in *Escherichia coli* urosepsis: association with patients' characteristics. *J Infect Dis*. 1987;156:225–229.
468. Warren JW. Clinical presentations and epidemiology of urinary tract infections. In: Mobley HLT, Warren JW, eds. *Urinary Tract Infections: Molecular Pathogenesis and Clinical Management*. Washington, DC: ASM Press; 1996:3–27.
469. Starčič Erjavec M, Žgur-Bertok D. Virulence potential for extraintestinal infections among commensal *Escherichia coli* isolated from healthy humans—the Trojan horse within our gut. *FEMS Microbiol Lett*. 2015;362. [Epub 2015 Feb 5].
470. Peirano G, Mulvey GL, Armstrong GD, et al. Virulence potential and adherence properties of *Escherichia coli* that produce CTX-M and NDM β -lactamases. *J Med Microbiol*. 2013;62(Pt 4):525–530. [Epub 2013 Jan 14].
471. Dale AP, Woodford N. Extra-intestinal pathogenic *Escherichia coli* (ExPEC): disease, carriage and clones. *J Infect*. 2015;71:615–626. Review. [Epub 2015 Sep 26].
472. Russo TA, Johnson JR. Medical and economic impact of extraintestinal infections due to *Escherichia coli*: focus on an increasingly important endemic problem. *Microbes Infect*. 2003;5:449–456. Review.
473. Banerjee R, Johnson JR. A new clone sweeps clean: the enigmatic emergence of *Escherichia coli* sequence type 131. *Antimicrob Agents Chemother*. 2014;58:4997–5004. Review. [Epub 2014 May 27].
474. Peirano G, van der Bij AK, Gregson DB, et al. Molecular epidemiology over an 11-year period (2000 to 2010) of extended-spectrum β -lactamase-producing *Escherichia coli* causing bacteremia in a centralized Canadian region. *J Clin Microbiol*. 2012;50:294–299. [Epub 2011 Dec 7].
475. Peirano G, Schreckenberger PC, Pitout JD. Characteristics of NDM-1-producing *Escherichia coli* isolates that belong to the successful and virulent clone ST131. *Antimicrob Agents Chemother*. 2011;55:2986–2988. [Epub 2011 Mar 28].
476. Kumarasamy KK, Toleman MA, Walsh TR, et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis*. 2010;10:597–602. [Epub 2010 Aug 10].
477. Peirano G, Pitout JD. Molecular epidemiology of *Escherichia coli* producing CTX-M beta-lactamases: the worldwide emergence of clone ST131 O25:H4. *Int J Antimicrob Agents*. 2010;35:316–321. Review. [Epub 2010 Jan 13].
478. Queenan AM, Bush K. Carbapenemases: the versatile beta-lactamases. *Clin Microbiol Rev*. 2007;20:440–458. table of contents. Review.
479. Pitout JD, Gregson DB, Church DL, et al. Population-based laboratory surveillance for AmpC beta-lactamase-producing *Escherichia coli*, Calgary. *Emerg Infect Dis*. 2007;13:443–448.
480. Flores-Mireles AL, Walker JN, Caparon M, et al. Urinary tract infections: epidemiology, mechanisms of infection and treatment options. *Nat Rev Microbiol*. 2015;13:269–284. Review. [Epub 2015 Apr 8].
481. Moreno E, Andreu A, Pigrau C, et al. Relationship between *Escherichia coli* strains causing acute cystitis in women and the fecal *E. coli* population of the host. *J Clin Microbiol*. 2008;46:2529–2534. [Epub 2008 May 21].
482. Yamamoto S, Tsukamoto T, Terai A, et al. Genetic evidence supporting the fecal-perineal-urethral hypothesis in cystitis caused by *Escherichia coli*. *J Urol*. 1997;157:1127–1129.
483. Manges AR, Johnson JR, Foxman B, et al. Widespread distribution of urinary tract infections caused by a multidrug-resistant *Escherichia coli* clonal group. *N Engl J Med*. 2001;345:1007–1013.
484. Phillips I, Eykyn S, King A, et al. Epidemic multiresistant *Escherichia coli* infection in West Lambeth health district. *Lancet*. 1988;1:1038–1041.
485. Tullus K, Hörlin K, Svenson SB, et al. Epidemic outbreaks of acute pyelonephritis caused by nosocomial spread of P fimbriated *Escherichia coli* in children. *J Infect Dis*. 1984;150:728–736.
486. Johnson JR, Stapleton AE, Russo TA, et al. Characteristics and prevalence within serogroup O4 of a J96-like clonal group of uropathogenic *Escherichia coli* O4:H5 containing the class I and class III alleles of *papG*. *Infect Immun*. 1997;65:2153–2159.
487. Johnson JR, O'Bryan TT, Delavari P, et al. Clonal relationships and extended virulence genotypes among *Escherichia coli* isolates from women with a first or recurrent episode of cystitis. *J Infect Dis*. 2001;183:1508–1517.
488. Zhang LX, Foxman B, Tallman P, et al. Distribution of *drb* genes coding for Dr binding adhesins among uropathogenic and fecal *Escherichia coli* isolates and identification of new subtypes. *Infect Immun*. 1997;65:2011–2018.
489. Olesen B, Scheut F, Andersen RL, et al. Enterotoxigenic *Escherichia coli* O78:H10, the cause of an outbreak of urinary tract infection. *J Clin Microbiol*. 2012;50:3703–3711. [Epub 2012 Sep 12].
490. Mobley HLT, Green DM, Trifillis AL, et al. Pyelonephritogenic *Escherichia coli* and killing of cultured human renal proximal tubular epithelial cells: role of hemolysin in some strains. *Infect Immun*. 1990;58:1281–1289.

491. Johnson DE, Russell RG. Animal models of urinary tract infection. In: Mobley HLT, Warren JW, eds. *Urinary Tract Infections: Molecular Pathogenesis and Clinical Management*. Washington, DC: ASM Press; 1996:377–403.
492. Johnson DE, Locketell CV, Russell RG, et al. Comparison of *Escherichia coli* strains recovered from human cystitis and pyelonephritis infections in transurethrally challenged mice. *Infect Immun*. 1998;66:3059–3065.
493. Langermann S, Molloy R, Burlein JE, et al. Vaccination with FimH adhesin protects cynomolgus monkeys from colonization and infection by uropathogenic *Escherichia coli*. *J Infect Dis*. 2000;181:774–778.
494. Roberts JA, Kaack B, Källénius G, et al. Receptors for pyelonephritogenic *Escherichia coli* in primates. *J Urol*. 1984;131:163–168.
495. Buckles EL, Bahrani-Mougeot FK, Molina A, et al. Identification and characterization of a novel uropathogenic *Escherichia coli*-associated fimbrial gene cluster. *Infect Immun*. 2004;72:3890–3901.
496. Brzuszkiewicz E, Brüggemann H, Liesegang H, et al. How to become a uropathogen: comparative genomic analysis of extraintestinal pathogenic *Escherichia coli* strains. *Proc Natl Acad Sci USA*. 2006;103:12879–12884.
497. Rendón MA, Saldaña Z, Erdem AL, et al. Commensal and pathogenic *Escherichia coli* use a common pilus adherence factor for epithelial cell colonization. *Proc Natl Acad Sci USA*. 2007;104:10637–10642.
498. Min G, Zhou G, Schapira M, et al. Structural basis of urothelial permeability barrier function as revealed by Cryo-EM studies of the 16 nm uropilin particle. *J Cell Sci*. 2003;116(Pt 20):4087–4094.
499. Wu XR, Sun TT, Medina JJ. In vitro binding of type 1-fimbriated *Escherichia coli* to uropilins Ia and Ib: relation to urinary tract infections. *Proc Natl Acad Sci USA*. 1996;93:9630–9635.
500. Mulvey MA, Lopez-Boado YS, Wilson CL, et al. Induction and evasion of host defenses by type 1-piliated uropathogenic *Escherichia coli*. *Science*. 1998;282:1494–1497.
501. Duncan MJ, Li G, Shin JS, et al. Bacterial penetration of bladder epithelium through lipid rafts. *J Biol Chem*. 2004;279:18944–18951. [Epub 2004 Feb 19].
502. Gunther NW, Snyder JA, Locketell V, et al. Assessment of virulence of uropathogenic *Escherichia coli* type 1 fimbrial mutants in which the invertible element is phase-locked on or off. *Infect Immun*. 2002;70:3344–3354.
503. Gunther NW, Locketell V, Johnson DE, et al. In vivo dynamics of type 1 fimbria regulation in uropathogenic *Escherichia coli* during experimental urinary tract infection. *Infect Immun*. 2001;69:2838–2846.
504. Lim JK, Gunther NW, Zhao H, et al. In vivo phase variation of *Escherichia coli* type 1 fimbrial genes in women with urinary tract infection. *Infect Immun*. 1998;66:3303–3310.
505. Stärk K, Khandige S, Kolmos HJ, et al. Uropathogenic *Escherichia coli* express type 1 fimbriae only in surface adherent populations under physiological growth conditions. *J Infect Dis*. 2016;213:386–394. [Epub 2015 Aug 19].
506. Yakovenko O, Sharma S, Forero M, et al. FimH forms catch bonds that are enhanced by mechanical force due to allosteric regulation. *J Biol Chem*. 2008;283:11596–11605. [Epub 2008 Feb 21].
507. Thomas WE, Nilsson LM, Forero M, et al. Shear-dependent 'stick-and-roll' adhesion of type 1 fimbriated *Escherichia coli*. *Mol Microbiol*. 2004;53:1545–1557.
508. Floyd RV, Upton M, Hultgren SJ, et al. *Escherichia coli*-mediated impairment of ureteric contractility is uropathogenic *E. coli* specific. *J Infect Dis*. 2012;206:1589–1596. [Epub 2012 Sep 21].
509. Väisänen V, Elo J, Tallgren LG, et al. Mannose-resistant haemagglutination and P antigen recognition are characteristic of *Escherichia coli* causing primary pyelonephritis. *Lancet*. 1981;2:1366–1369.
510. Källénius G, Svenson S, Möllby R, et al. Structure of carbohydrate part of receptor on human uroepithelial cells for pyelonephritogenic *Escherichia coli*. *Lancet*. 1981;2:604–606.
511. Johnson JR. Virulence factors in *Escherichia coli* urinary tract infection. *Clin Microbiol Rev*. 1991;4:80–128.
512. Donnenberg MS, Welch RA. Virulence determinants of uropathogenic *Escherichia coli*. In: Mobley HLT, Warren JW, eds. *Urinary Tract Infections: Molecular Pathogenesis and Clinical Management*. Washington, DC: ASM Press; 1996:135–174.
513. Johanson I, Lindstedt R, Svanborg C. Roles of the *pap*- and *prs*-encoded adhesins in *Escherichia coli* adherence to human uroepithelial cells. *Infect Immun*. 1992;60:3416–3422.
514. Korhonen TK, Virkola R, Holthofer H. Localization of binding sites for purified *Escherichia coli* P fimbriae in the human kidney. *Infect Immun*. 1986;54:328–332.
515. Roberts JA, Marklund B-I, Ilver D, et al. The Gal(α 1-4) Gal-specific tip adhesin of *Escherichia coli* P-fimbriae is needed for pyelonephritis to occur in the normal urinary tract. *Proc Natl Acad Sci USA*. 1994;91:11889–11893.
516. Stapleton A, Nudelman E, Clausen H, et al. Binding of uropathogenic *Escherichia coli* R45 to glycolipids extracted from vaginal epithelial cells is dependent on histo-blood group secretor status. *J Clin Invest*. 1992;90:965–972.
517. Wroblewska-Seniuk K, Selvarangan R, Hart A, et al. Dra/ AfaE adhesin of uropathogenic Dr/Afa+ *Escherichia coli* mediates mortality in pregnant rats. *Infect Immun*. 2005;73:7597–7601.
518. Foxman B, Zhang L, Tallman P, et al. Virulence characteristics of *Escherichia coli* causing first urinary tract infection predict risk of second infection. *J Infect Dis*. 1995;172:1536–1541.
519. Stapleton A, Moseley S, Stamm WE. Urovirulence determinants in *Escherichia coli* isolates causing first-episode and recurrent cystitis in women. *J Infect Dis*. 1991;163:773–779.
520. Nowicki B, Svanborg-Edén C, Hull R, et al. Molecular analysis and epidemiology of the Dr hemagglutinin of uropathogenic *Escherichia coli*. *Infect Immun*. 1989;57:446–451.
521. Archambaud M, Courcoux P, Labigne-Roussel A. Detection by molecular hybridization of *pap*, *afa*, and *sfa* adherence systems in *Escherichia coli* strains associated with urinary and enteric infections. *Ann Inst Pasteur Microbiol*. 1988;139:575–588.
522. Selvarangan R, Goluszko P, Singhal J, et al. Interaction of Dr adhesin with collagen type IV is a critical step in *Escherichia coli* renal persistence. *Infect Immun*. 2004;72:4827–4835.
523. Goluszko P, Moseley SL, Truong LD, et al. Development of experimental model of chronic pyelonephritis with *Escherichia coli* O75:K5:H-bearing Dr fimbriae: mutation in the *dra* region prevented tubulointerstitial nephritis. *J Clin Invest*. 1997;99:1662–1672.
524. Regua-Mangia AH, Irino K, da Silva Pacheco R, et al. Molecular characterization of uropathogenic and diarrheagenic *Escherichia coli* pathotypes. *J Basic Microbiol*. 2010;50(suppl 1):S107–S115.
525. Wallace-Gadsden F, Johnson JR, Wain J, et al. Enterotoxigenic *Escherichia coli* related to uropathogenic clonal group A. *Emerg Infect Dis*. 2007;13:757–760.
526. Boll EJ, Struve C, Boisen N, et al. Role of enterotoxigenic *Escherichia coli* virulence factors in uropathogenesis. *Infect Immun*. 2013;81:1164–1171. [Epub 2013 Jan 28].
527. Lewis AJ, Richards AC, Mulvey MA. Invasion of Host Cells and Tissues by Uropathogenic Bacteria. *Microbiol Spectr*. 2016;4.
528. Robino L, Scavone P, Araujo L, et al. Intracellular bacteria in the pathogenesis of *Escherichia coli* urinary tract infection in children. *Clin Infect Dis*. 2014;59:e158–e164. [Epub 2014 Aug 4].
529. Bokil NJ, Totsika M, Carey AJ, et al. Intramacrophage survival of uropathogenic *Escherichia coli*: differences between diverse clinical isolates and between mouse and human macrophages. *Immunobiology*. 2011;216:1164–1171. [Epub 2011 May 24].
530. Rosen DA, Hooton TM, Stamm WE, et al. Detection of intracellular bacterial communities in human urinary tract infection. *PLoS Med*. 2007;4:e329.
531. Mysorekar IU, Hultgren SJ. Mechanisms of uropathogenic *Escherichia coli* persistence and eradication from the urinary tract. *Proc Natl Acad Sci USA*. 2006;103:14170–14175. [Epub 2006 Sep 12].
532. Wiles TJ, Dhakal BK, Eto DS, et al. Inactivation of host Akt/protein kinase B signaling by bacterial pore-forming toxins. *Mol Biol Cell*. 2008;19:1427–1438. [Epub 2008 Jan 30].
533. Smith YC, Rasmussen SB, Grande KK, et al. Hemolysin of uropathogenic *Escherichia coli* evokes extensive shedding of the uroepithelium and hemorrhage in bladder tissue within the first 24 hours after intraurethral inoculation of mice. *Infect Immun*. 2008;76:2978–2990.
534. Nagamatsu K, Hannan TJ, Guest RL, et al. Dysregulation of *Escherichia coli* α -hemolysin expression alters the course of acute and persistent urinary tract infection. *Proc Natl Acad Sci USA*. 2015;112:E871–E880, Feb 24. [Epub 2015 Feb 9].
535. Schmidt G, Sehr P, Wilm M, et al. Gln 63 of Rho is deamidated by *Escherichia coli* cytotoxic necrotizing factor-1. *Nature*. 1997;387:725–729.
536. Boquet P. The cytotoxic necrotizing factor 1 (CNF1) from *Escherichia coli*. *Toxicon*. 2001;39:1673–1680, Review.
537. Ripperre-Lampe KE, O'Brien AD, Conran R, et al. Mutation of the gene encoding cytotoxic necrotizing factor type 1 (*cnf1*) attenuates the virulence of uropathogenic *Escherichia coli*. *Infect Immun*. 2001;69:3954–3964.
538. Buckles EL, Wang X, Lane MC, et al. The role of K2 capsule in *Escherichia coli* urinary tract infection and serum resistance. *J Infect Dis*. 2009;199:1689–1697.
539. Russo TA, Brown JJ, Jodush ST, et al. The O4 specific antigen moiety of lipopolysaccharide but not the K54 group 2 capsule is important for urovirulence of an extraintestinal isolate of *Escherichia coli*. *Infect Immun*. 1996;64:2343–2348.
540. Bahrani-Mougeot FK, Buckles EL, Locketell CV, et al. Type-1 fimbriae and extracellular polysaccharides are preeminent uropathogenic *Escherichia coli* virulence determinants in the murine urinary tract. *Mol Microbiol*. 2002;45:1079–1093.
541. Nagy G, Dobrindt U, Schneider G, et al. Loss of regulatory protein RfaH attenuates virulence of uropathogenic *Escherichia coli*. *Infect Immun*. 2002;70:4406–4413.
542. Buckles EL, Wang X, Locketell CV, et al. PhoU enhances the ability of extraintestinal pathogenic *Escherichia coli* strain CF7073 to colonize the murine urinary tract. *Microbiology*. 2006;152:153–160.
543. Sarkar S, Ulett GC, Totsika M, et al. Role of capsule and O antigen in the virulence of uropathogenic *Escherichia coli*. *PLoS ONE*. 2014;9:e94786, eCollection 2014.
544. O'Brien VP, Hannan TJ, Nielsen HV, et al. Drug and Vaccine Development for the Treatment and Prevention of Urinary Tract Infections. *Microbiol Spectr*. 2016;4.
545. Schwenger EM, Tejani AM, Loewen PS. Probiotics for preventing urinary tract infections in adults and children. *Cochrane Database Syst Rev*. 2015;(12):CD008772.
546. Stapleton AE, Au-Yeung M, Hooton TM, et al. Randomized, placebo-controlled phase 2 trial of a *Lactobacillus crispatus* probiotic given intravaginally for prevention of recurrent urinary tract infection. *Clin Infect Dis*. 2011;52:1212–1217. [Epub 2011 Apr 14].
547. Baerheim A, Larsen E, Digraanes A. Vaginal application of lactobacilli in the prophylaxis of recurrent lower urinary tract infection in women. *Scand J Prim Health Care*. 1994;12:239–243.
548. Jepson RG, Williams G, Craig JC. Cranberries for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD001321.
549. Stapleton AE, Dziura J, Hooton TM, et al. Recurrent urinary tract infection and urinary *Escherichia coli* in women ingesting cranberry juice daily: a randomized controlled trial. *Mayo Clin Proc*. 2012;87:143–150.
550. Barbosa-Cesnik C, Brown MB, Buxton M, et al. Cranberry juice fails to prevent recurrent urinary tract infection: results from a randomized placebo-controlled trial. *Clin Infect Dis*. 2011;52:23–30, Jan 1.
551. Gupta K, Chou MY, Howell A, et al. Cranberry products inhibit adherence of p-fimbriated *Escherichia coli* to primary cultured bladder and vaginal epithelial cells. *J Urol*. 2007;177:2357–2360.
552. Howell AB, Foxman B. Cranberry juice and adhesion of antibiotic-resistant uropathogens. *JAMA*. 2002;287:3082–3083.
553. Lee BS, Bhuta T, Simpson JM, et al. Methenamine hippurate for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD003265.
554. Perrotta C, Aznar M, Mejia R, et al. Oestrogens for preventing recurrent urinary tract infection in postmenopausal women. *Cochrane Database Syst Rev*. 2008;(2):CD005131.
555. Eriksen B. A randomized, open, parallel-group study on the preventive effect of an estradiol-releasing vaginal ring (Estring) on recurrent urinary tract infections in postmenopausal women. *Am J Obstet Gynecol*. 1999;180:1072–1079.
556. Mydock-McGrane L, Cusumano Z, Han Z, et al. Antivirulence C-mannosides as antibiotic-sparing, oral therapeutics for urinary tract infections. *J Med Chem*. 2016;59:9390–9408. [Epub 2016 Oct 14].
557. Totsika M, Kostakioti M, Hannan TJ, et al. A FimH inhibitor prevents acute bladder infection and treats chronic cystitis caused by multidrug-resistant uropathogenic *Escherichia coli* ST131. *J Infect Dis*. 2013;208:921–928. [Epub 2013 Jun 3].
558. Terlizzi ME, Gribaudo G, Maffei ME. Uropathogenic *Escherichia coli* (UEPC) infections: virulence factors, bladder responses, antibiotic, and non-antibiotic antimicrobial strategies. *Front Microbiol*. 2017;8:1566, eCollection 2017. Review.
559. Brumbaugh AR, Mobley HL. Preventing urinary tract infection: progress toward an effective *Escherichia coli* vaccine. *Expert Rev Vaccines*. 2012;11:663–676, Review.
560. Brumbaugh AR, Smith SN, Subashchandrase S, et al. Blocking yersiniabactin import attenuates extraintestinal pathogenic *Escherichia coli* in cystitis and pyelonephritis and represents a novel target to prevent urinary tract infection. *Infect Immun*. 2015;83:1443–1450. [Epub 2015 Jan 26].

561. Brumbaugh AR, Smith SN, Mobley HL. Immunization with the yersiniabactin receptor, FyuA, protects against pyelonephritis in a murine model of urinary tract infection. *Infect Immun*. 2013;81:3309–3316. [Epub 2013 Jun 24].
562. Hopkins WJ, Elkhwaji J, Beierle LM, et al. Vaginal mucosal vaccine for recurrent urinary tract infections in women: results of a phase 2 clinical trial. *J Urol*. 2007;177:1349–1353, quiz 1591.
563. Huttner A, Hatz C, van den Dobbelen G, et al. Safety, immunogenicity, and preliminary clinical efficacy of a vaccine against extraintestinal pathogenic *Escherichia coli* in women with a history of recurrent urinary tract infection: a randomised, single-blind, placebo-controlled phase 1b trial. *Lancet Infect Dis*. 2017;17:528–537. [Epub 2017 Feb 24].
564. Taha Neto KA, Nogueira Castilho L, Reis LO. Oral vaccine (OM-89) in the recurrent urinary tract infection prophylaxis: a realistic systematic review with meta-analysis. *Actas Urol Esp*. 2016;40:203–208, Review. English, Spanish. [Epub 2015 Oct 24].
565. Beerepoot MA, Geerlings SE, van Haarst EP, et al. Nonantibiotic prophylaxis for recurrent urinary tract infections: a systematic review and meta-analysis of randomized controlled trials. *J Urol*. 2013;190:1981–1989, Review. [Epub 2013 Jul 15].
566. Naber KG, Cho YH, Matsumoto T, et al. Immunoactive prophylaxis of recurrent urinary tract infections: a meta-analysis. *Int J Antimicrob Agents*. 2009;33:111–119, Review. [Epub 2008 Oct 28].
567. Kim KS, Itabashi H, Gerns P, et al. The K1 capsule is the critical determinant in the development of *Escherichia coli* meningitis in the rat. *J Clin Invest*. 1992;90:897–905.
568. Siegel JD, McCracken GH Jr. Sepsis neonatorum. *N Engl J Med*. 1981;304:642–647, Review.
569. Basmaci R, Bonacorsi S, Bide P, et al. *Escherichia coli* meningitis features in 325 children from 2001 to 2013 in France. *Clin Infect Dis*. 2015;61:779–786. [Epub 2015 May 5].
570. Pabst HF, Kreth HW. Ontogeny of the immune response as a basis of childhood disease. *J Pediatr*. 1980;97:519–534, Review.
571. Robbins JB, McCracken GH Jr, Gotschlich EC, et al. *Escherichia coli* K1 capsular polysaccharide associated with neonatal meningitis. *N Engl J Med*. 1974;290:1216–1220.
572. Kasper DL, Winkelhake JL, Zollinger WD, et al. Immunochemical similarity between polysaccharide antigens of *Escherichia coli* 07: K1(L):NM and group B *Neisseria meningitidis*. *J Immunol*. 1973;110:262–268.
573. Kim KS, Itabashi H, Gerns P, et al. The K1 capsule is the critical determinant in the development of *Escherichia coli* meningitis in the rat. *J Clin Invest*. 1992;90:897–905.
574. Wooster DG, Maruvada R, Blom AM, et al. Logarithmic phase *Escherichia coli* K1 efficiently avoids serum killing by promoting C4bp-mediated C3b and C4b degradation. *Immunology*. 2006;117:482–493.
575. Siitonen A, Takala A, Ratiner YA, et al. Invasive *Escherichia coli* infections in children: bacterial characteristics in different age groups and clinical entities. *Pediatr Infect Dis J*. 1993;12:606–612.
576. Korhonen TK, Parkkinen J, Hacker J, et al. Binding of *Escherichia coli* S fimbriae to human kidney epithelium. *Infect Immun*. 1986;54:322–327.
577. Parkkinen J, Rogers GN, Korhonen T, et al. Identification of the O-linked sialyloligosaccharides of glycoprotein A as the erythrocyte receptors for S-fimbriated *Escherichia coli*. *Infect Immun*. 1986;54:37–42.
578. Parkkinen J, Korhonen TK, Pere A, et al. Binding sites in the rat brain for *Escherichia coli* S fimbriae associated with neonatal meningitis. *J Clin Invest*. 1988;81:860–865.
579. Krishnan S, Chen S, Turcatel G, et al. Regulation of Toll-like receptor 2 interaction with Ecp96 controls *Escherichia coli* K1 invasion of brain endothelial cells. *Cell Microbiol*. 2013;15:63–81. [Epub 2012 Oct 4].
580. Krishnan S, Fernandez GE, Sacks DB, et al. IQGAP1 mediates the disruption of adherens junctions to promote *Escherichia coli* K1 invasion of brain endothelial cells. *Cell Microbiol*. 2012;14:1415–1433. [Epub 2012 May 24].
581. Khan NA, Kim Y, Shin S, et al. FimH-mediated *Escherichia coli* K1 invasion of human brain microvascular endothelial cells. *Cell Microbiol*. 2007;9:169–178.
582. Prasadarao NV, Wass CA, Weiser JN, et al. Outer membrane protein A of *Escherichia coli* contributes to invasion of brain microvascular endothelial cells. *Infect Immun*. 1996;64:146–153.
583. Korhonen TK, Valtonen MV, Parkkinen J, et al. Serotypes, hemolysin production, and receptor recognition of *Escherichia coli* strains associated with neonatal sepsis and meningitis. *Infect Immun*. 1985;48:486–491.
584. Johnson JR, Oswald E, O'Bryan TT, et al. Phylogenetic distribution of virulence-associated genes among *Escherichia coli* isolates associated with neonatal bacterial meningitis in the Netherlands. *J Infect Dis*. 2002;185:774–784.
585. Blum G, Falbo V, Caprioli A, et al. Gene clusters encoding the cytotoxic necrotizing factor type 1, Prs-fimbriae and α -hemolysin form the pathogenicity island II of the uropathogenic *Escherichia coli* strain J96. *FEMS Microbiol Lett*. 1995;126:189–196.
586. Falbo V, Famiglietti M, Caprioli A. Gene block encoding production of cytotoxic necrotizing factor 1 and hemolysin in *Escherichia coli* isolates from extraintestinal infections. *Infect Immun*. 1992;60:2182–2187.
587. Badger JL, Wass CA, Weissman SJ, et al. Application of signature-tagged mutagenesis for identification of *Escherichia coli* K1 genes that contribute to invasion of human brain microvascular endothelial cells. *Infect Immun*. 2000;68:5056–5061.
588. Badger JL, Wass CA, Kim KS. Identification of *Escherichia coli* K1 genes contributing to human brain microvascular endothelial cell invasion by differential fluorescence induction. *Mol Microbiol*. 2000;36:174–182.
589. McCarthy AJ, Martin P, Cloup E, et al. The genotoxin colibactin is a determinant of virulence in *Escherichia coli* K1 experimental neonatal systemic infection. *Infect Immun*. 2015;83:3704–3711. [Epub 2015 Jul 6].
590. Burdet C, Clermont O, Bonacorsi S, et al; COLIBAFI Group. *Escherichia coli* bacteremia in children: age and portal of entry are the main predictors of severity. *Pediatr Infect Dis J*. 2014;33:872–879.
591. Berman H, Barberino MG, Moreira ED Jr, et al. Distribution of strain type and antimicrobial susceptibility of *Escherichia coli* isolates causing meningitis in a large urban setting in Brazil. *J Clin Microbiol*. 2014;52:1418–1422. [Epub 2014 Feb 12].
592. Diekema DJ, Pfaffner MA, Jones RN, et al. Survey of bloodstream infections due to gram-negative bacilli: frequency of occurrence and antimicrobial susceptibility of isolates collected in the United States, Canada, and Latin America for the SENTRY Antimicrobial Surveillance Program, 1997. *Clin Infect Dis*. 1999;29:595–607.
593. Sannes MR, Kuskowski MA, Owens K, et al. Virulence factor profiles and phylogenetic background of *Escherichia coli* isolates from veterans with bacteremia and uninfected control subjects. *J Infect Dis*. 2004;190:2121–2128. [Epub 2004 Nov 15].
594. Lefort A, Panhard X, Clermont O, et al; COLIBAFI Group. Host factors and portal of entry outweigh bacterial determinants to predict the severity of *Escherichia coli* bacteremia. *J Clin Microbiol*. 2011;49:777–783. [Epub 2010 Dec 22].
595. Macutkiewicz C, Carlson G, Clark E, et al. Characterisation of *Escherichia coli* strains involved in transcytosis across gut epithelial cells exposed to metabolic and inflammatory stress. *Microbes Infect*. 2008;10:424–431. [Epub 2008 Jan 12].
596. Wolochow H, Hildebrand GJ, Lamanna C. Translocation of microorganisms across the intestinal wall of the rat: effect of microbial size and concentration. *J Infect Dis*. 1966;116:523–528.
597. Berg RD. Bacterial translocation from the gastrointestinal tract. *Trends Microbiol*. 1995;3:149–154, Review.
598. Samet A, Sledzińska A, Krawczyk B, et al. Leukemia and risk of recurrent *Escherichia coli* bacteremia: genotyping implicates *E. coli* translocation from the colon to the bloodstream. *Eur J Clin Microbiol Infect Dis*. 2013;32:1393–1400. [Epub 2013 May 7].
599. Krawczyk B, Sledzińska A, Szemiako K, et al. Characterisation of *Escherichia coli* isolates from the blood of haematological adult patients with bacteraemia: translocation from gut to blood requires the cooperation of multiple virulence factors. *Eur J Clin Microbiol Infect Dis*. 2015;34:1135–1143. [Epub 2015 Feb 6].
600. Poole NM, Green SI, Rajan A, et al. Role for FimH in Extraintestinal Pathogenic *Escherichia coli* Invasion and Translocation through the Intestinal Epithelium. *Infect Immun*. 2017;85.
601. Mahjoub-Messai F, Bide P, Caro V, et al. *Escherichia coli* isolates causing bacteremia via gut translocation and urinary tract infection in young infants exhibit different virulence genotypes. *J Infect Dis*. 2011;203:1844–1849. [Epub 2011 May 5].
602. Poolman JT, Wacker M. Extraintestinal Pathogenic *Escherichia coli*, a Common Human Pathogen: challenges for Vaccine Development and Progress in the Field. *J Infect Dis*. 2016;213:6–13, Review. [Epub 2015 Sep 2].
603. Moriel DG, Bertoldi I, Spagnuolo A, et al. Identification of protective and broadly conserved vaccine antigens from the genome of extraintestinal pathogenic *Escherichia coli*. *Proc Natl Acad Sci USA*. 2010;107:9072–9077. [Epub 2010 May 3].
604. Kanki M, Yoda T, Tsukamoto T, et al. *Klebsiella pneumoniae* produces no histamine: *Raoultella planticola* and *Raoultella ornithinolytica* strains are histamine producers. *Appl Environ Microbiol*. 2002;68:3462–3466.
605. Drancourt M, Bollet C, Carta A, et al. Phylogenetic analyses of *Klebsiella* species delineate *Klebsiella* and *Raoultella* gen. nov., with description of *Raoultella ornithinolytica* comb. nov., *Raoultella terrigena* comb. nov. and *Raoultella planticola* comb. nov. *Int J Syst Evol Microbiol*. 2001;51(Pt 3):925–932.
606. Gorrie CL, Mirceta M, Wick RR, et al. Gastrointestinal carriage is a major reservoir of *Klebsiella pneumoniae* infection in intensive care patients. *Clin Infect Dis*. 2017;65:208–215.
607. Conlan S, Kong HH, Segre JA. Species-level analysis of DNA sequence data from the NIH Human Microbiome Project. *PLoS ONE*. 2012;7:e47075. [Epub 2012 Oct 10].
608. Siu LK, Yeh KM, Lin JC, et al. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. *Lancet Infect Dis*. 2012;12:881–887, Review.
609. Lin YT, Siu LK, Lin JC, et al. Seroepidemiology of *Klebsiella pneumoniae* colonizing the intestinal tract of healthy Chinese and overseas Chinese adults in Asian countries. *BMC Microbiol*. 2012;12:13.
610. Al-Hasan MN, Lahr BD, Eckel-Passow JE, et al. Seasonal variation in *Escherichia coli* bloodstream infection: a population-based study. *Clin Microbiol Infect*. 2009;15:947–950.
611. Anderson DJ, Richet H, Chen LF, et al. Seasonal variation in *Klebsiella pneumoniae* bloodstream infection on 4 continents. *J Infect Dis*. 2008;197:752–756.
612. Weiner LM, Webb AK, Limbago B, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. *Infect Control Hosp Epidemiol*. 2016;37:1288–1301. [Epub 2016 Aug 30].
613. Pitout JD, Nordmann P, Poirer L. Carbapenemase-producing *Klebsiella pneumoniae*, a key pathogen set for global nosocomial dominance. *Antimicrob Agents Chemother*. 2015;59:5873–5884, Review. [Epub 2015 Jul 13].
614. Munoz-Price LS, Poirer L, Bonomo RA, et al. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis*. 2013;13:785–796, Review.
615. Zhanel GG, Adam HJ, Baxter MR, et al; Canadian Antimicrobial Resistance Alliance. Antimicrobial susceptibility of 22746 pathogens from Canadian hospitals: results of the CANWARD 2007–11 study. *J Antimicrob Chemother*. 2013;68(suppl 1):i7–i22.
616. Vincent JL, Rello J, Marshall J, et al; EPIC Int Group of Investigators. International study of the prevalence and outcomes of infection in intensive care units. *JAMA*. 2009;302:2323–2329.
617. Pagano L, Caira M, Treacchi EM, et al. Carbapenemase-producing *Klebsiella pneumoniae* and hematologic malignancies. *Emerg Infect Dis*. 2014;20:1235–1236, Review.
618. Schwaber MJ, Klarfeld-Lidji S, Navon-Venezia S, et al. Predictors of carbapenem-resistant *Klebsiella pneumoniae* acquisition among hospitalized adults and effect of acquisition on mortality. *Antimicrob Agents Chemother*. 2008;52:1028–1033. [Epub 2007 Dec 17].
619. Tzouvelekis LS, Markogiannakis A, Psichogiou M, et al. Carbapenemases in *Klebsiella pneumoniae* and other Enterobacteriaceae: an evolving crisis of global dimensions. *Clin Microbiol Rev*. 2012;25:682–707, Review.
620. McClelland S 3rd, Hall WA. Postoperative central nervous system infection: incidence and associated factors in 2111 neurosurgical procedures. *Clin Infect Dis*. 2007;45:55–59. [Epub 2007 May 21].
621. Weisfelt M, van de Beek D, Spanjaard L, et al. Nosocomial bacterial meningitis in adults: a prospective series of 50 cases. *J Hosp Infect*. 2007;66:71–78. [Epub 2007 Apr 11].
622. Korinek AM, Baugnon T, Gormard JL, et al. Risk factors for adult nosocomial meningitis after craniotomy: role of antibiotic prophylaxis. *Neurosurgery*. 2006;59:126–133, discussion 126–33.
623. Tang LM, Chen ST. *Klebsiella pneumoniae* meningitis: prognostic factors. *Scand J Infect Dis*. 1994;26:95–102.
624. Durand ML, Calderwood SB, Weber DJ, et al. Acute bacterial meningitis in adults. A review of 493 episodes. *N Engl J Med*. 1993;328:21–28.
625. Song JY, Jung SJ, Park CW, et al. Prognostic significance of infection acquisition sites in spontaneous bacterial peritonitis: nosocomial versus community acquired. *J Korean Med Sci*. 2006;21:666–671.
626. Such J, Runyon BA. Spontaneous bacterial peritonitis. *Clin Infect Dis*. 1998;27:669–674, quiz 675–6. Review.
627. Brisse S, Fevre C, Passet V, et al. Virulent clones of *Klebsiella pneumoniae*: identification and evolutionary

- scenario based on genomic and phenotypic characterization. *PLoS ONE*. 2009;4:e4982. [Epub 2009 Mar 25].
628. Luderan ER, Crum NE. Pyogenic liver abscess with a focus on *Klebsiella pneumoniae* as a primary pathogen: an emerging disease with unique clinical characteristics. *Am J Gastroenterol*. 2005;100:322–331. Review.
 629. Rahimian J, Wilson T, Oram V, et al. Pyogenic liver abscess: recent trends in etiology and mortality. *Clin Infect Dis*. 2004;39:1654–1659. [Epub 2004 Nov 9].
 630. Yang CC, Yen CH, Ho MW, et al. Comparison of pyogenic liver abscess caused by non-*Klebsiella pneumoniae* and *Klebsiella pneumoniae*. *J Microbiol Immunol Infect*. 2004;37:176–184.
 631. Liu YC, Cheng DL, Lin CL. *Klebsiella pneumoniae* liver abscess associated with septic endophthalmitis. *Arch Intern Med*. 1986;146:1913–1916.
 632. Wang JH, Liu YC, Lee SS, et al. Primary liver abscess due to *Klebsiella pneumoniae* in Taiwan. *Clin Infect Dis*. 1998;26:1434–1438.
 633. Lin YT, Liu CJ, Yeh YC, et al. Ampicillin and amoxicillin use and the risk of *Klebsiella pneumoniae* liver abscess in Taiwan. *J Infect Dis*. 2013;208:211–217. [Epub 2013 Apr 8].
 634. Lee SS, Chen YS, Tsai HC, et al. Predictors of septic metastatic infection and mortality among patients with *Klebsiella pneumoniae* liver abscess. *Clin Infect Dis*. 2008;47:642–650.
 635. Fang CT, Lai SY, Yi WC, et al. *Klebsiella pneumoniae* genotype K1: an emerging pathogen that causes septic ocular or central nervous system complications from pyogenic liver abscess. *Clin Infect Dis*. 2007;45:284–293. Review. [Epub 2007 Jun 19].
 636. Fung CP, Chang FY, Lee SC, et al. A global emerging disease of *Klebsiella pneumoniae* liver abscess: is serotype K1 an important factor for complicated endophthalmitis? *Gut*. 2002;50:420–424.
 637. Cheng DL, Liu YC, Yen MY, et al. Septic metastatic lesions of pyogenic liver abscess. Their association with *Klebsiella pneumoniae* bacteremia in diabetic patients. *Arch Intern Med*. 1991;151:1557–1559.
 638. Garcia de la Torre M, Romero-Vivas J, Martinez-Beltrán J, et al. *Klebsiella* bacteremia: an analysis of 100 episodes. *Rev Infect Dis*. 1985;7:143–150.
 639. Al-Hasan MN, Eckel-Passow JE, Baddour LM. Bacteremia complicating gram-negative urinary tract infections: a population-based study. *J Infect*. 2010;60:278–285. [Epub 2010 Jan 28].
 640. Togawa A, Toh H, Onozawa K, et al. Influence of the bacterial phenotypes on the clinical manifestations in *Klebsiella pneumoniae* bacteremia patients: a retrospective cohort study. *J Infect Chemother*. 2015;21:531–537. [Epub 2015 Apr 22].
 641. Chetcuti Zammit S, Azzopardi N, Sant J. Mortality risk score for *Klebsiella pneumoniae* bacteraemia. *Eur J Intern Med*. 2014;25:571–576. [Epub 2014 May 9].
 642. Tseng CP, Wu HS, Wu TH, et al. Clinical characteristics and outcome of patients with community-onset *Klebsiella pneumoniae* bacteremia requiring intensive care. *J Microbiol Immunol Infect*. 2013;46:217–223. [Epub 2012 Jul 24].
 643. Meatherall BL, Gregson D, Ross T, et al. Incidence, risk factors, and outcomes of *Klebsiella pneumoniae* bacteremia. *Am J Med*. 2009;122:866–873.
 644. Carpenter JL. *Klebsiella* pulmonary infections: occurrence at one medical center and review. *Rev Infect Dis*. 1990;12:672–682.
 645. Mandell LA, Wunderink RG, Anzueto A, et al; Infectious Diseases Society of America; American Thoracic Society. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis*. 2007;44(suppl 2):S27–S72.
 646. File TM. Community-acquired pneumonia. *Lancet*. 2003;362:1991–2001. Review.
 647. Hu HC, Huang CC, Tsai YH, et al. Outcome analysis of patients requiring mechanical ventilation with severe community-acquired pneumonia and identified bacterial pathogens. *Chang Gung Med J*. 2005;28:229–236.
 648. Lin YT, Wang YP, Wang FD, et al. Community-onset *Klebsiella pneumoniae* pneumonia in Taiwan: clinical features of the disease and associated microbiological characteristics of isolates from pneumonia and nasopharynx. *Front Microbiol*. 2015;9:122. eCollection 2015.
 649. Janssens JP. Pneumonia in the elderly (geriatric) population. *Curr Opin Pulm Med*. 2005;11:226–230. Review.
 650. Yeh KM, Kurup A, Siu LK, et al. Capsular serotype K1 or K2, rather than magA and rmpA, is a major virulence determinant for *Klebsiella pneumoniae* liver abscess in Singapore and Taiwan. *J Clin Microbiol*. 2007;45:466–471. [Epub 2006 Dec 6].
 651. Yu VL, Hansen DS, Ko WC, et al; International *Klebsiella* Study Group. Virulence characteristics of *Klebsiella* and clinical manifestations of *K. pneumoniae* bloodstream infections. *Emerg Infect Dis*. 2007;13:986–993.
 652. Paczosa MK, Mecsas J. *Klebsiella pneumoniae*: going on the Offense with a Strong Defense. *Microbiol Mol Biol Rev*. 2016;80:629–661. Print 2016 Sep. Review.
 653. Pomakova DK, Hsiao CB, Beanan JM, et al. Clinical and phenotypic differences between classic and hypervirulent *Klebsiella pneumoniae*: an emerging and under-recognized pathogenic variant. *Eur J Clin Microbiol Infect Dis*. 2012;31:981–989. [Epub 2011 Sep 15].
 654. Lee CH, Liu JW, Su LH, et al. Hypermucoviscosity associated with *Klebsiella pneumoniae*-mediated invasive syndrome: a prospective cross-sectional study in Taiwan. *Int J Infect Dis*. 2010;14:e688–e692. [Epub 2010 May 23].
 655. Lee HC, Chuang YC, Yu WL, et al. Clinical implications of hypermucoviscosity phenotype in *Klebsiella pneumoniae* isolates: association with invasive syndrome in patients with community-acquired bacteraemia. *J Intern Med*. 2006;259:606–614.
 656. Kawai T. Hypermucoviscosity: an extremely sticky phenotype of *Klebsiella pneumoniae* associated with emerging destructive tissue abscess syndrome. *Clin Infect Dis*. 2006;42:1359–1361. [Epub 2006 Apr 11].
 657. Shon AS, Bajwa RP, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. *Virulence*. 2013;4:107–118. Review. [Epub 2013 Jan 9].
 658. Lin YC, Lu MC, Tang HL, et al. Assessment of hypermucoviscosity as a virulence factor for experimental *Klebsiella pneumoniae* infections: comparative virulence analysis with hypermucoviscosity-negative strain. *BMC Microbiol*. 2011;11:50.
 659. Lin JC, Siu LK, Fung CP, et al. Impaired phagocytosis of capsular serotypes K1 or K2 *Klebsiella pneumoniae* in type 2 diabetes mellitus patients with poor glycemic control. *J Clin Endocrinol Metab*. 2006;91:3084–3087. [Epub 2006 May 23].
 660. Ko WC, Paterson DL, Sagnimeni AJ, et al. Community-acquired *Klebsiella pneumoniae* bacteremia: global differences in clinical patterns. *Emerg Infect Dis*. 2002;8:160–166.
 661. Turton JB, Engleider H, Gabriel SN, et al. Genetically similar isolates of *Klebsiella pneumoniae* serotype K1 causing liver abscesses in three continents. *J Med Microbiol*. 2007;56(Pt 5):593–597.
 662. Struve C, Roe CC, Stegger M, et al. Mapping the Evolution of Hypervirulent *Klebsiella pneumoniae*. *MBio*. 2015;6:e00630.
 663. Luo Y, Wang Y, Ye L, et al. Molecular epidemiology and virulence factors of pyogenic liver abscess causing *Klebsiella pneumoniae* in China. *Clin Microbiol Infect*. 2014;20:O818–O824. [Epub 2014 Jun 14].
 664. Fang CT, Chuang YP, Shun CT, et al. A novel virulence gene in *Klebsiella pneumoniae* strains causing primary liver abscess and septic metastatic complications. *J Exp Med*. 2004;199:697–705.
 665. Yeh KM, Chang FY, Fung CP, et al. Serotype K1 capsule, rather than magA per se, is really the virulence factor in *Klebsiella pneumoniae* strains that cause primary pyogenic liver abscess. *J Infect Dis*. 2006;194:403–404. author reply 404–5.
 666. Yeh KM, Chang FY, Fung CP, et al. magA is not a specific virulence gene for *Klebsiella pneumoniae* strains causing liver abscess but is part of the capsular polysaccharide gene cluster of *K. pneumoniae* serotype K1. *J Med Microbiol*. 2006;55(Pt 6):803–804.
 667. Qu TT, Zhou JC, Jiang Y, et al. Clinical and microbiological characteristics of *Klebsiella pneumoniae* liver abscess in East China. *BMC Infect Dis*. 2015;15:161.
 668. Yu WL, Lee MF, Tang HJ, et al. Low prevalence of rmpA and high tendency of rmpA mutation correspond to low virulence of extended spectrum β -lactamase-producing *Klebsiella pneumoniae* isolates. *Virulence*. 2015;6:162–172.
 669. Yu WL, Ko WC, Cheng KC, et al. Association between rmpA and magA genes and clinical syndromes caused by *Klebsiella pneumoniae* in Taiwan. *Clin Infect Dis*. 2006;42:1351–1358. [Epub 2006 Apr 11].
 670. Ma LC, Fang CT, Lee CZ, et al. Genomic heterogeneity in *Klebsiella pneumoniae* strains is associated with primary pyogenic liver abscess and metastatic infection. *J Infect Dis*. 2005;192:117–128. [Epub 2005 May 25].
 671. Alcántar-Curiel MD, Blackburn D, Saldaña Z, et al. Multi-functional analysis of *Klebsiella pneumoniae* fimbrial types in adherence and biofilm formation. *Virulence*. 2013;4:129–138. [Epub 2013 Jan 9].
 672. Guo Y, Wang S, Zhan L, et al. Microbiological and Clinical Characteristics of Hypermucoviscous *Klebsiella pneumoniae* Isolates Associated with Invasive Infections in China. *Front Cell Infect Microbiol*. 2017;7:24. eCollection 2017.
 673. Hsieh PF, Lin TL, Lee CZ, et al. Serum-induced iron-acquisition systems and TonB contribute to virulence in *Klebsiella pneumoniae* causing primary pyogenic liver abscess. *J Infect Dis*. 2008;197:1717–1727.
 674. Nassif X, Sansonetti PJ. Correlation of the virulence of *Klebsiella pneumoniae* K1 and K2 with the presence of a plasmid encoding aerobactin. *Infect Immun*. 1986;54:603–608.
 675. Russo TA, Olson R, MacDonald U, et al. Aerobactin, but not yersiniabactin, salmochelin, or enterobactin, enables the growth/survival of hypervirulent (hypermucoviscous) *Klebsiella pneumoniae* ex vivo and in vivo. *Infect Immun*. 2015;83:3325–3333. [Epub 2015 Jun 8].
 676. Russo TA, Olson R, MacDonald U, et al. Aerobactin mediates virulence and accounts for increased siderophore production under iron-limiting conditions by hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*. *Infect Immun*. 2014;82:2356–2367. [Epub 2014 Mar 24].
 677. Hæggman S, Löfdahl S, Burman LG. An allelic variant of the chromosomal gene for class A beta-lactamase K2, specific for *Klebsiella pneumoniae*, is the ancestor of SHV-1. *Antimicrob Agents Chemother*. 1997;41:2705–2709.
 678. Tumbarello M, Spanu T, Sanguinetti M, et al. Bloodstream infections caused by extended-spectrum-beta-lactamase-producing *Klebsiella pneumoniae*: risk factors, molecular epidemiology, and clinical outcome. *Antimicrob Agents Chemother*. 2006;50:498–504.
 679. Ben-David D, Kordevani R, Keller N, et al. Outcome of carbapenem resistant *Klebsiella pneumoniae* bloodstream infections. *Clin Microbiol Infect*. 2012;18:54–60.
 680. Yigit H, Queenan AM, Anderson GJ, et al. Novel carbapenem-hydrolyzing beta-lactamase, KPC-1, from a carbapenem-resistant strain of *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2001;45:1151–1161. Erratum in: *Antimicrob Agents Chemother*. 2008 Feb;52(2):809.
 681. Andraca R, Edson RS, Kern EB. Rhinoscleroma: a growing concern in the United States? Mayo Clinic experience. *Mayo Clin Proc*. 1993;68:1151–1157.
 682. de Pontual L, Ovetchkine P, Rodriguez D, et al. Rhinoscleroma: a French national retrospective study of epidemiological and clinical features. *Clin Infect Dis*. 2008;47:1396–1402.
 683. Favre C, Passet V, Deletoile A, et al. PCR-based identification of *Klebsiella pneumoniae* subsp. *rhinoscleromatis*, the agent of rhinoscleroma. *PLoS Negl Trop Dis*. 2011;5:e1052. [Epub 2011 May 24].
 684. Hart CA, Rao SK. Rhinoscleroma. *J Med Microbiol*. 2000;49:395–396.
 685. Lenis A, Ruff T, Diaz JA, et al. Rhinoscleroma. *South Med J*. 1988;81:1580–1582.
 686. Goldstein EJ, Lewis RP, Martin WJ, et al. Infections caused by *Klebsiella ozaenae*: a changing disease spectrum. *J Clin Microbiol*. 1978;8:413–418.
 687. Tang LM, Chen ST. *Klebsiella ozaenae* meningitis: report of two cases and review of the literature. *Infection*. 1994;22:58–61.
 688. Hoffmann KM, Deutschmann A, Weitzer C, et al. Antibiotic-associated hemorrhagic colitis caused by cytotoxin-producing *Klebsiella oxytoca*. *Pediatrics*. 2010;125:e960–e963. [Epub 2010 Mar 1].
 689. Shinjoh M, Iwata S, Takahashi T. *Klebsiella oxytoca*-positive, penicillin-associated hemorrhagic enterocolitis in children. *Pediatr Int*. 2010;52:132–133.
 690. Beaugerie L, Metz M, Barbut F, et al; Infectious Colitis Study Group. *Klebsiella oxytoca* as an agent of antibiotic-associated hemorrhagic colitis. *Clin Gastroenterol Hepatol*. 2003;1:370–376.
 691. Högenauer C, Langner C, Beubler E, et al. *Klebsiella oxytoca* as a causative organism of antibiotic-associated hemorrhagic colitis. *N Engl J Med*. 2006;355:2418–2426.
 692. Joainig MM, Gorkiewicz G, Leitner E, et al. Cytotoxic effects of *Klebsiella oxytoca* strains isolated from patients with antibiotic-associated hemorrhagic colitis or other diseases caused by infections and from healthy subjects. *J Clin Microbiol*. 2010;48:817–824. [Epub 2010 Jan 6].
 - 692a. Tindall BJ, Sutton G, Garrity GM. Enterobacter aerogenes Hormaeche and Edwards 1960 (Approved Lists 1980) and *Klebsiella mobilis* Bascomb et al. 1971 (Approved Lists 1980) share the same nomenclatural type (ATCC 13048) on the Approved Lists and are homotypic synonyms, with consequences for the name *Klebsiella mobilis* Bascomb et al. 1971 (Approved Lists 1980). *Int J Syst Evol Microbiol*. 2017;67:502–504.
 - 692b. Munson E, Carroll KC. An Update on the Novel Genera and Species and Revised Taxonomic Status of Bacterial Organisms Described in 2016 and 2017. *J Clin Microbiol*. 2019;57:e01181-18.
 693. Iversen C, Lehner A, Mullane N, et al. The taxonomy of *Enterobacter sakazakii*: proposal of a new genus

- Cronobacter* gen. nov. and descriptions of *Cronobacter sakazakii* comb. nov. *Cronobacter sakazakii* subsp. *sakazakii*, comb. nov., *Cronobacter sakazakii* subsp. *malonaticus* subsp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov. and *Cronobacter* genomospecies 1. *BMC Evol Biol.* 2007;7:64.
694. Forsythe SJ, Abbott SL, Pitout J. *Klebsiella*, *Enterobacter*, *Citrobacter*, *Cronobacter*, *Serratia*, *Plesiomonas*, and other Enterobacteriaceae. In: *Manual of Clinical Microbiology*. 10th ed. Washington, DC: American Society of Microbiology; 2015:714–737.
 695. Maki DG, Rhame FS, Mackel DC, et al. Nationwide epidemic of septicemia caused by contaminated intravenous products. I. Epidemiologic and clinical features. *Am J Med.* 1976;60:471–485.
 696. Pestourie N, Garnier F, Barraud O, et al. Outbreak of AmpC β -lactamase-hyper-producing *Enterobacter cloacae* in a neonatal intensive care unit in a French teaching hospital. *Am J Infect Control.* 2014;42:456–458.
 697. Lockhart SR, Abramson MA, Beekmann SE, et al. Antimicrobial resistance among Gram-negative bacilli causing infections in intensive care unit patients in the United States between 1993 and 2004. *J Clin Microbiol.* 2007;45:3352–3359. [Epub 2007 Aug 22].
 698. Davin-Regli A, Pagès JM. *Enterobacter aerogenes* and *Enterobacter cloacae*: versatile bacterial pathogens confronting antibiotic treatment. *Front Microbiol.* 2015;6:392. eCollection 2015. Review.
 699. Carbone A, Arnaud I, Maugat S, et al; MDRB Surveillance National Steering Group (BMR-Raisin). National multidrug-resistant bacteria (MDRB) surveillance in France through the RAISIN network: a 9 year experience. *J Antimicrob Chemother.* 2013;68:954–959. [Epub 2012 Nov 29].
 700. Song EH, Park KH, Jang EY, et al. Comparison of the clinical and microbiologic characteristics of patients with *Enterobacter cloacae* and *Enterobacter aerogenes* bacteremia: a prospective observation study. *Diagn Microbiol Infect Dis.* 2010;66:436–440. [Epub 2010 Jan 13].
 701. Bowen AB, Braden CR. Invasive *Enterobacter sakazakii* disease in infants. *Emerg Infect Dis.* 2006;12:1185–1189.
 702. Centers for Disease Control and Prevention (CDC). *Cronobacter* species isolation in two infants - New Mexico, 2008. *MMWR Morb Mortal Wkly Rep.* 2009;58:1179–1183.
 703. Hunter CJ, Petrosyan M, Ford HR, et al. *Enterobacter sakazakii*: an emerging pathogen in infants and neonates. *Surg Infect (Larchmt).* 2008;9:533–539. Review.
 704. Drudy D, Mullane NR, Quinn T, et al. *Enterobacter sakazakii*: an emerging pathogen in powdered infant formula. *Clin Infect Dis.* 2006;42:996–1002. Review. [Epub 2006 Feb 22].
 705. Centers for Disease Control and Prevention (CDC). *Enterobacter sakazakii* infections associated with the use of powdered infant formula—Tennessee, 2001. *MMWR Morb Mortal Wkly Rep.* 2002;51:297–300.
 706. Kalyantanda G, Shumyak L, Archibald LK. *Cronobacter* Species Contamination of Powdered Infant Formula and the Implications for Neonatal Health. *Front Pediatr.* 2015;3:56. eCollection 2015. Review.
 707. Friedemann M. Epidemiology of invasive neonatal *Cronobacter* (*Enterobacter sakazakii*) infections. *Eur J Clin Microbiol Infect Dis.* 2009;28:1297–1304. [Epub 2009 Aug 7].
 708. Cruz AT, Cazacu AC, Allen CH. *Pantoea agglomerans*, a plant pathogen causing human disease. *J Clin Microbiol.* 2007;45:1989–1992.
 709. Enterobacteriaceae. In: Ryan KJ, eds. *Sherris Medical Microbiology*, 7th ed. New York, NY: McGraw-Hill.
 710. Chart H. In: Greenwood D, Slack RC, Barer MR, et al, eds. *Medical Microbiology: A Guide to Microbial Infections: Pathogenesis, Immunity, Laboratory Diagnosis and Control*. 18th ed. Edinburgh; New York: Churchill Livingstone/Elsevier; 2012. Chapter 27, *Klebsiella*, *Enterobacter*, *Proteus* and other Enterobacteriaceae: Pneumonia; urinary tract infection; opportunistic infection; p. 290–297.
 711. Liu WY, Wong CF, Chung KM, et al. Comparative genome analysis of *Enterobacter cloacae*. *PLoS ONE.* 2013;8:e74487. eCollection 2013.
 712. Pérez A, Poza M, Fernández A, et al. Involvement of the AcrAB-TolC efflux pump in the resistance, fitness, and virulence of *Enterobacter cloacae*. *Antimicrob Agents Chemother.* 2012;56:2084–2090. [Epub 2012 Jan 30].
 713. Townsend S, Hurrell E, Forsythe S. Virulence studies of *Enterobacter sakazakii* isolates associated with a neonatal intensive care unit outbreak. *BMC Microbiol.* 2008;8:64.
 714. Singh N, Goel G, Raghav M. Insights into virulence factors determining the pathogenicity of *Cronobacter sakazakii*. *Virulence.* 2015;6:433–440. Review. [Epub 2015 May 7].
 715. Nair MK, Venkitanarayanan K, Silbart LK, et al. Outer membrane protein A (OmpA) of *Cronobacter sakazakii* binds fibronectin and contributes to invasion of human brain microvascular endothelial cells. *Foodborne Pathog Dis.* 2009;6:495–501.
 716. Mohan Nair MK, Venkitanarayanan KS. Cloning and sequencing of the ompA gene of *Enterobacter sakazakii* and development of an ompA-targeted PCR for rapid detection of *Enterobacter sakazakii* in infant formula. *Appl Environ Microbiol.* 2006;72:2539–2546.
 717. Fei P, Jiang Y, Feng J, et al. Antibiotic and Desiccation Resistance of *Cronobacter sakazakii* and *C. malonaticus* Isolates from Powdered Infant Formula and Processing Environments. *Front Microbiol.* 2017;8:316. eCollection 2017.
 718. Yan QQ, Condell O, Power K, et al. *Cronobacter* species (formerly known as *Enterobacter sakazakii*) in powdered infant formula: a review of our current understanding of the biology of this bacterium. *J Appl Microbiol.* 2012;113:1–15. Review. [Epub 2012 Apr 11].
 719. Walsh D, Molloy C, Iversen C, et al. Survival characteristics of environmental and clinically derived strains of *Cronobacter sakazakii* in infant milk formula (IMF) and ingredients. *J Appl Microbiol.* 2011;110:697–703. [Epub 2011 Jan 24].
 720. Breeuwer P, Lardeau A, Peterz M, et al. Desiccation and heat tolerance of *Enterobacter sakazakii*. *J Appl Microbiol.* 2003;95:967–973.
 721. Nazarowec-White M, Farber JM. Thermal resistance of *Enterobacter sakazakii* in reconstituted dried-infant formula. *Lett Appl Microbiol.* 1997;24:9–13.
 722. Joseph S, Hariri S, Masood N, et al. Sialic acid utilization by *Cronobacter sakazakii*. *Microb Inform Exp.* 2013;3:3.
 723. Boucher HW, Talbot GH, Bradley JS, et al. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis.* 2009;48:1–12.
 724. Rice LB. Federal funding for the study of antimicrobial resistance in nosocomial pathogens: no ESKAPE. *J Infect Dis.* 2008;197:1079–1081.
 725. Korfmann G, Wiedemann B. Genetic control of beta-lactamase production in *Enterobacter cloacae*. *Rev Infect Dis.* 1988;10:793–799.
 726. Izdebski R, Baraniak A, Herda M, et al; MOSAR WP2, WP3 and WP5 Study Groups. MLST reveals potentially high-risk international clones of *Enterobacter cloacae*. *J Antimicrob Chemother.* 2015;70:48–56. [Epub 2014 Sep 12].
 727. Mezzatesta ML, Gona F, Stefani S. *Enterobacter cloacae* complex: clinical impact and emerging antibiotic resistance. *Future Microbiol.* 2012;7:887–902. Review.
 728. Choi SH, Lee JE, Park SJ, et al. Emergence of antibiotic resistance during therapy for infections caused by Enterobacteriaceae producing AmpC beta-lactamase: implications for antibiotic use. *Antimicrob Agents Chemother.* 2008;52:995–1000. [Epub 2007 Dec 17].
 729. Bizio B. Lettera di Bartolomeo Bizio al chiarissimo canonico Angelo Bellani sopra il fenomeno della polenta porporina. *Bibl Ital G Lett Sci Art (Anno VIII).* 1823;30:275–295. [in Italian].
 730. Sehdev PS, Donnenberg MS. Arcanum: the 19th-century Italian pharmacist pictured here was the first to characterize what are now known to be bacteria of the genus *Serratia*. *Clin Infect Dis.* 1999;29:925.
 731. Yu VL. *Serratia marcescens*: historical perspective and clinical review. *N Engl J Med.* 1979;300:887–893.
 732. Mahlen SD. *Serratia* infections: from military experiments to current practice. *Clin Microbiol Rev.* 2011;24:755–791. Review.
 733. Engel HJ, Collignon PJ, Whiting PT, et al. *Serratia* sp. bacteremia in Canberra, Australia: a population-based study over 10 years. *Eur J Clin Microbiol Infect Dis.* 2009;28:821–824. [Epub 2009 Feb 5].
 734. Laupland KB, Parkins MD, Gregson DB, et al. Population-based laboratory surveillance for *Serratia* species isolates in a large Canadian health region. *Eur J Clin Microbiol Infect Dis.* 2008;27:89–95. [Epub 2007 Oct 25].
 735. Sunenshine RH, Tan ET, Terashita DM, et al. A multistate outbreak of *Serratia marcescens* bloodstream infection associated with contaminated intravenous magnesium sulfate from a compounding pharmacy. *Clin Infect Dis.* 2007;45:527–533. [Epub 2007 Jul 24].
 736. Blossom D, Noble-Wang J, Su J, et al; *Serratia* in Prefilled Syringes Investigation Team Group. Multistate outbreak of *Serratia marcescens* bloodstream infections caused by contamination of prefilled heparin and isotonic sodium chloride solution syringes. *Arch Intern Med.* 2009;169:1705–1711.
 737. Buffet-Bataillon S, Rabier V, Bérémieux P, et al. Outbreak of *Serratia marcescens* in a neonatal intensive care unit: contaminated unmedicated liquid soap and risk factors. *J Hosp Infect.* 2009;72:17–22. [Epub 2009 Feb 25].
 738. Horcajada JP, Martínez JA, Alcón A, et al. Acquisition of multidrug-resistant *Serratia marcescens* by critically ill patients who consumed tap water during receipt of oral medication. *Infect Control Hosp Epidemiol.* 2006;27:774–777. [Epub 2006 Jun 9].
 739. Takahashi H, Kramer MH, Yasui Y, et al. Nosocomial *Serratia marcescens* outbreak in Osaka, Japan, from 1999 to 2000. *Infect Control Hosp Epidemiol.* 2004;25:156–161.
 740. Kirschke DL, Jones TF, Craig AS, et al. *Pseudomonas aeruginosa* and *Serratia marcescens* contamination associated with a manufacturing defect in bronchoscopes. *N Engl J Med.* 2003;348:214–220.
 741. Gupta N, Hovevar SN, Moulton-Meissner HA, et al. Outbreak of *Serratia marcescens* bloodstream infections in patients receiving parenteral nutrition prepared by a compounding pharmacy. *Clin Infect Dis.* 2014;59:1–8. [Epub 2014 Apr 11].
 742. Pan A, Dolcetti L, Barosi C, et al. An outbreak of *Serratia marcescens* bloodstream infections associated with misuse of drug vials in a surgical ward. *Infect Control Hosp Epidemiol.* 2006;27:79–82. [Epub 2006 Jan 6].
 743. Grohskopf LA, Roth VR, Feikin DR, et al. *Serratia liquefaciens* bloodstream infections from contamination of epoetin alfa at a hemodialysis center. *N Engl J Med.* 2001;344:1491–1497.
 744. Roth VR, Arduino MJ, Nobilotti J, et al. Transfusion-related sepsis due to *Serratia liquefaciens* in the United States. *Transfusion.* 2000;40:931–935.
 745. Mills J, Drew D. *Serratia marcescens* endocarditis: a regional illness associated with intravenous drug abuse. *Ann Intern Med.* 1976;84:29–35.
 746. Samonis G, Vouloumanou EK, Christofaki M, et al. *Serratia* infections in a general hospital: characteristics and outcomes. *Eur J Clin Microbiol Infect Dis.* 2011;30:653–660. [Epub 2011 Jan 10].
 747. Lakhundi S, Siddiqui R, Khan NA. Pathogenesis of microbial keratitis. *Microb Pathog.* 2017;104:97–109. Review. [Epub 2016 Dec 18].
 748. Das S, Shorey H, Taylor HR, et al. Association between cultures of contact lens and corneal scraping in contact lens related microbial keratitis. *Arch Ophthalmol.* 2007;125:1182–1185.
 749. Cheng KH, Leung SL, Hoekman HW, et al. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet.* 1999;354:181–185.
 750. Remuzgo-Martínez S, Aranzamendi-Zaldunbide M, Pilares-Ortega L, et al. Interaction of macrophages with a cytotoxic *Serratia liquefaciens* human isolate. *Microbes Infect.* 2013;15:480–490. [Epub 2013 Mar 22].
 751. Kurz CL, Chauvet S, Andrés E, et al. Virulence factors of the human opportunistic pathogen *Serratia marcescens* identified by in vivo screening. *EMBO J.* 2003;22:1451–1460.
 752. Hertle R. The family of *Serratia* type pore forming toxins. *Curr Protein Pept Sci.* 2005;6:313–325. Review.
 753. Lai HC, Soo PC, Wei JR, et al. The RssAB two-component signal transduction system in *Serratia marcescens* regulates swarming motility and cell envelope architecture in response to exogenous saturated fatty acids. *J Bacteriol.* 2005;187:3407–3414.
 754. Lin CS, Horng JT, Yang CH, et al. RssAB-FlhDC-ShlBA as a major pathogenesis pathway in *Serratia marcescens*. *Infect Immun.* 2010;78:4870–4881. [Epub 2010 Aug 16].
 755. Van Houdt R, Givskov M, Michiels CW. Quorum sensing in *Serratia*. *FEMS Microbiol Rev.* 2007;31:407–424. Review. [Epub 2007 Apr 25].
 756. Shanks RM, Stella NA, Kalivoda EJ, et al. A *Serratia marcescens* OxyR homolog mediates surface attachment and biofilm formation. *J Bacteriol.* 2007;189:7262–7272. [Epub 2007 Aug 3].
 757. Remuzgo-Martínez S, Lázaro-Díez M, Mayer C, et al. Biofilm formation and quorum-sensing-molecule production by clinical isolates of *Serratia liquefaciens*. *Appl Environ Microbiol.* 2015;81:3306–3315. [Epub 2015 Mar 6].
 758. Pinna A, Usai D, Sechi LA, et al. Detection of virulence factors in *Serratia* strains isolated from contact lens-associated corneal ulcers. *Acta Ophthalmol.* 2011;89:382–387. [Epub 2009 Oct 20].
 759. Mahlen SD, Morrow SS, Abdalhamid B, et al. Analyses of *ampC* gene expression in *Serratia marcescens* reveal new regulatory properties. *J Antimicrob Chemother.* 2003;51:791–802.
 760. Stock I, Grueger T, Wiedemann B. Natural antibiotic susceptibility of strains of *Serratia marcescens* and the *S. liquefaciens* complex: *S. liquefaciens* sensu stricto, *S. proteamaculans* and *S. grimesii*. *Int J Antimicrob Agents.* 2003;22:35–47.
 761. Jacoby GA. AmpC beta-lactamases. *Clin Microbiol Rev.* 2009;22:161–182. Table of Contents. Review.
 762. Lipsky BA, Hook EW III, Smith AA, et al. *Citrobacter* infections in humans: experience at the Seattle Veterans Administration Medical Center and a review of the literature. *Rev Infect Dis.* 1980;2:746–760.

763. Ranjan KP, Ranjan N. *Citrobacter*: an emerging health care associated urinary pathogen. *Urol Ann*. 2013;5:313–314.
764. Jiménez A, Castro JG, Munoz-Price LS, et al. Outbreak of *Klebsiella pneumoniae* Carbenapemase-Producing *Citrobacter freundii* at a Tertiary Acute Care Facility in Miami, Florida. *Infect Control Hosp Epidemiol*. 2017;38:320–326. [Epub 2016 Dec 7].
765. Shih CC, Chen YC, Chang SC, et al. Bacteremia due to *Citrobacter* species: significance of primary intraabdominal infection. *Clin Infect Dis*. 1996;23:543–549.
766. Samonis G, Karageorgopoulos DE, Kofteridis DP, et al. *Citrobacter* infections in a general hospital: characteristics and outcomes. *Eur J Clin Microbiol Infect Dis*. 2009;28:61–68.
767. Deleted in review.
768. Liu LH, Wang NY, Wu AY, et al. *Citrobacter freundii* bacteremia: risk factors of mortality and prevalence of resistance genes. *J Microbiol Immunol Infect*. 2018;51:565–572. [Epub 2017 Jun 22].
769. Townsend SM, Gonzalez-Gomez I, Badger JL. fliP influences *Citrobacter koseri* macrophage uptake, cytokine expression and brain abscess formation in the neonatal rat. *J Med Microbiol*. 2006;55(Pt 12):1631–1640.
770. Li J, Musser JM, Beltran P, et al. Genotypic heterogeneity of strains of *Citrobacter diversus* expressing a 32-kilodalton outer membrane protein associated with neonatal meningitis. *J Clin Microbiol*. 1990;28:1760–1765.
771. Williams WW, Mariano J, Spurrier M, et al. Nosocomial meningitis due to *Citrobacter diversus* in neonates: new aspects of the epidemiology. *J Infect Dis*. 1984;150:229–235.
772. Pletz MW, Wollny A, Dobermann UH, et al. A nosocomial foodborne outbreak of a VIM carbapenemase-expressing *Citrobacter freundii*. *Clin Infect Dis*. 2018;67:58–64.
773. Pepperell C, Kus JV, Gardam MA, et al. Low-virulence *Citrobacter* species encode resistance to multiple antimicrobials. *Antimicrob Agents Chemother*. 2002;46:3555–3560.
774. Huys G, Cnockaert M, Abbott SL, et al. *Hafnia paralvei* sp. nov., formerly known as *Hafnia alvei* hybridization group 2. *Int J Syst Evol Microbiol*. 2010;60(Pt 8):1725–1728. [Epub 2009 Sep 4].
775. Abbott SL, Moler S, Green N, et al. Clinical and laboratory diagnostic characteristics and cytotoxicigenic potential of *Hafnia alvei* and *Hafnia paralvei* strains. *J Clin Microbiol*. 2011;49:3122–3126. [Epub 2011 Jul 27].
776. Janda JM, Abbott SL. The genus *Hafnia*: from soup to nuts. *Clin Microbiol Rev*. 2006;19:12–18.
777. Günthard H, Pennekamp A. Clinical significance of extraintestinal *Hafnia alvei* isolates from 61 patients and review of the literature. *Clin Infect Dis*. 1996;22:1040–1045.
778. Ridell J, Siitonen A, Paulin L, et al. *Hafnia alvei* in stool specimens from patients with diarrhea and healthy controls. *J Clin Microbiol*. 1994;32:2335–2337.
779. Stock I, Rahman M, Sherwood KJ, et al. Natural antimicrobial susceptibility patterns and biochemical identification of *Escherichia albertii* and *Hafnia alvei* strains. *Diagn Microbiol Infect Dis*. 2005;51:151–163.
780. Jayal A, Saly M, Nordmann P, et al. *Hafnia*, an enterobacterial genus naturally resistant to colistin revealed by three susceptibility testing methods. *J Antimicrob Chemother*. 2017;72:2507–2511.
781. Skurnik D, Nucci A, Ruimy R, et al. Emergence of carbapenem-resistant *Hafnia*: the fall of the last soldier. *Clin Infect Dis*. 2010;50:1429–1431.
782. Armbruster CE, Mobley HL. Merging mythology and morphology: the multifaceted lifestyle of *Proteus mirabilis*. *Nat Rev Microbiol*. 2012;10:743–754. Review. [Epub 2012 Oct 8].
783. O'Hara CM, Brenner FW, Miller JM. Classification, identification, and clinical significance of *Proteus*, *Providencia*, and *Morganella*. *Clin Microbiol Rev*. 2000;13:534–546.
784. Karlowsky JA, Lagacé-Wiens PR, Simmer PJ, et al. Antimicrobial resistance in urinary tract pathogens in Canada from 2007 to 2009: CANWARD surveillance study. *Antimicrob Agents Chemother*. 2011;55:3169–3175. [Epub 2011 May 2].
785. Jacobsen SM, Stickler DJ, Mobley HL, et al. Complicated catheter-associated urinary tract infections due to *Escherichia coli* and *Proteus mirabilis*. *Clin Microbiol Rev*. 2008;21:26–59.
786. Stickler DJ, Feneley RC. The encrustation and blockage of long-term indwelling bladder catheters: a way forward in prevention and control. *Spinal Cord*. 2010;48:784–790. Review. [Epub 2010 Apr 6].
787. Sader HS, Flamm RK, Jones RN. Frequency of occurrence and antimicrobial susceptibility of Gram-negative bacteremia isolates in patients with urinary tract infection: results from United States and European hospitals (2009–2011). *J Chemother*. 2014;26:133–138. [Epub 2013 Dec 6].
788. Adams-Sapper S, Sergeeva-Selezneva J, Tartof S, et al. Globally dispersed mobile drug-resistance genes in gram-negative bacterial isolates from patients with bloodstream infections in a US urban general hospital. *J Med Microbiol*. 2012;61(Pt 7):968–974. [Epub 2012 Apr 5].
789. Mylotte JM. Nursing home-acquired bloodstream infection. *Infect Control Hosp Epidemiol*. 2005;26:833–837. Review.
790. Chen CY, Chen YH, Lu PL, et al. *Proteus mirabilis* urinary tract infection and bacteremia: risk factors, clinical presentation, and outcomes. *J Microbiol Immunol Infect*. 2012;45:228–236. [Epub 2012 May 7].
791. Schaffer JN, Pearson MM. *Proteus mirabilis* and Urinary Tract Infections. *Microbiol Spectr*. 2015;3.
792. Pearson MM, Sebahia M, Churcher C, et al. Complete genome sequence of uropathogenic *Proteus mirabilis*, a master of both adherence and motility. *J Bacteriol*. 2008;190:4027–4037. [Epub 2008 Mar 28].
793. Li X, Lockatell CV, Johnson DE, et al. Identification of MrpI as the sole recombinase that regulates the phase variation of MR/P fimbria, a bladder colonization factor of uropathogenic *Proteus mirabilis*. *Mol Microbiol*. 2002;45:865–874.
794. Zhao H, Li X, Johnson DE, et al. *In vivo* phase variation of MR/P fimbrial gene expression in *Proteus mirabilis* infecting the urinary tract. *Mol Microbiol*. 1997;23:1009–1019.
795. Bahrani FK, Massad G, Lockatell CV, et al. Construction of an MR/P fimbrial mutant of *Proteus mirabilis*: role in virulence in a mouse model of ascending urinary tract infection. *Infect Immun*. 1994;62:3363–3371.
796. Torzewska A, Budzyńska A, Białczak-Kokot M, et al. *In vitro* studies of epithelium-associated crystallization caused by uropathogens during urinary calculi development. *Microb Pathog*. 2014;71:72–25–31. [Epub 2014 May 4].
797. Stickler DJ. Bacterial biofilms in patients with indwelling urinary catheters. *Nat Clin Pract Urol*. 2008;5:598–608. Review. [Epub 2008 Oct 14].
798. Jones BV, Mahenthalingam E, Sabbuba NA, et al. Role of swarming in the formation of crystalline *Proteus mirabilis* biofilms on urinary catheters. *J Med Microbiol*. 2005;54(Pt 9):807–813.
799. Dattelbaum JD, Lockatell CV, Johnson DE, et al. UreR, the transcriptional activator of the *Proteus mirabilis* urease gene cluster, is required for urease activity and virulence in experimental urinary tract infections. *Infect Immun*. 2003;71:1026–1030.
800. Jones BD, Lockatell CV, Johnson DE, et al. Construction of a urease-negative mutant of *Proteus mirabilis*: analysis of virulence in a mouse model of ascending urinary tract infection. *Infect Immun*. 1990;58:1120–1123.
801. Armbruster CE, Forsyth-DeOrnellas V, Johnson AO, et al. Genome-wide transposon mutagenesis of *Proteus mirabilis*: essential genes, fitness factors for catheter-associated urinary tract infection, and the impact of polymicrobial infection on fitness requirements. *PLoS Pathog*. 2017;13:e1006434. eCollection 2017 Jun.
802. Armbruster CE, Smith SN, Johnson AO, et al. The pathogenic potential of *Proteus mirabilis* is enhanced by other uropathogens during polymicrobial urinary tract infection. *Infect Immun*. 2017;85.
803. Armbruster CE, Smith SN, Yep A, et al. Increased incidence of urolithiasis and bacteremia during *Proteus mirabilis* and *Providencia stuartii* coinfection due to synergistic induction of urease activity. *J Infect Dis*. 2014;209:1524–1532. [Epub 2013 Nov 26].
804. Wie SH. Clinical significance of *Providencia* bacteremia or bacteriuria. *Korean J Intern Med*. 2015;30:167–169. [Epub 2015 Feb 27].
805. De Vecchi E, Sitta S, Romanò CL, et al. Aetiology and antibiotic resistance patterns of urinary tract infections in the elderly: a 6-month study. *J Med Microbiol*. 2013;62(Pt 6):859–863. [Epub 2013 Mar 8].
806. Tumbarello M, Citton R, Spanu T, et al. ESBL-producing multidrug-resistant *Providencia stuartii* infections in a university hospital. *J Antimicrob Chemother*. 2004;53:277–282. [Epub 2003 Dec 19].
807. Muder RR, Brennen C, Wagener MM, et al. Bacteremia in a long-term-care facility: a five-year prospective study of 163 consecutive episodes. *Clin Infect Dis*. 1992;14:647–654.
808. El Khatib M, Tran QT, Nasrallah C, et al. *Providencia stuartii* form biofilms and floating communities of cells that display high resistance to environmental insults. *PLoS ONE*. 2017;12:e0174213. eCollection 2017.
809. Lahlaoui H, Poirer L, Moussa MB, et al. Nosocomial dissemination of extended-spectrum β -lactamase VEB-1a-producing *Providencia stuartii* isolates in a Tunisian hospital. *Eur J Clin Microbiol Infect Dis*. 2011;30:1267–1270. [Epub 2011 Mar 27].
810. O'Hara CM, Brenner FW, Miller JM. Classification, identification, and clinical significance of *Proteus*, *Providencia*, and *Morganella*. *Clin Microbiol Rev*. 2000;13:534–546. Review.
811. Oikonomou O, Liakopoulos A, Phee LM, et al. *Providencia stuartii* isolates from Greece: co-carriage of cephalosporin (blaSHV-5, blaVEB-1), carbapenem (blaVIM-1), and aminoglycoside (rmtB) resistance determinants by a multidrug-resistant outbreak clone. *Microb Drug Resist*. 2016;22:379–386. [Epub 2016 Jan 26].
812. Douka E, Perivolioti E, Kraniotaki E, et al. Emergence of a pandrug-resistant VIM-1-producing *Providencia stuartii* clonal strain causing an outbreak in a Greek intensive care unit. *Int J Antimicrob Agents*. 2015;45:533–536. [Epub 2015 Feb 16].
813. Mataseje LF, Boyd DA, Lefebvre B, et al; Canadian Nosocomial Infection Surveillance Program. Complete sequences of a novel blaNDM-1-harboring plasmid from *Providencia rettgeri* and an FII-type plasmid from *Klebsiella pneumoniae* identified in Canada. *J Antimicrob Chemother*. 2014;69:637–642. [Epub 2013 Nov 24].
814. Zavaski AP, Carvalhaes CG, da Silva GL, et al. Outbreak of carbapenem-resistant *Providencia stuartii* in an intensive care unit. *Infect Control Hosp Epidemiol*. 2012;33:627–630. [Epub 2012 Apr 24].
815. Falagas ME, Kavvadia PK, Mantadakis E, et al. *Morganella morganii* infections in a general tertiary hospital. *Infection*. 2006;34:315–321.
816. Lin TY, Chan MC, Yang YS, et al. Clinical manifestations and prognostic factors of *Morganella morganii* bacteremia. *Eur J Clin Microbiol Infect Dis*. 2015;34:231–236. [Epub 2014 Aug 9].
817. Lee IK, Liu JW. Clinical characteristics and risk factors for mortality in *Morganella morganii* bacteremia. *J Microbiol Immunol Infect*. 2006;39:328–334.
818. Cêtre JC, Nicolle MC, Salord H, et al. Outbreaks of contaminated broncho-alveolar lavage related to intrinsically defective bronchoscopes. *J Hosp Infect*. 2005;61:39–45.
819. Poirer L, Guibert M, Girlich D, et al. Cloning, sequence analyses, expression, and distribution of *ampC-ampR* from *Morganella morganii* clinical isolates. *Antimicrob Agents Chemother*. 1999;43:769–776.
820. Slaven EM, Lopez FA, Hart SM, et al. Myonecrosis caused by *Edwardsiella tarda*: a case report and case series of extraintestinal *E. tarda* infections. *Clin Infect Dis*. 2001;32:1430–1433.
821. Crosby SN, Snoddy MC, Atkinson CT, et al. Upper extremity myonecrosis caused by *Edwardsiella tarda* resulting in transhumeral amputation: case report. *J Hand Surg Am*. 2013;38:129–132. [Epub 2012 Nov 30].
822. Janda JM, Abbott SL. Infections associated with the genus *Edwardsiella*: the role of *Edwardsiella tarda* in human disease. *Clin Infect Dis*. 1993;17:742–748.
823. Yang M, Lv Y, Xiao J, et al. *Edwardsiella* comparative phylogenomics reveal the new intra/inter-species taxonomic relationships, virulence evolution and niche adaptation mechanisms. *PLoS ONE*. 2012;7:e36987. [Epub 2012 May 10].
824. Janda JM, Abbott SL, McIver CJ. *Plesiomonas shigelloides* Revisited. *Clin Microbiol Rev*. 2016;29:349–374. Review.
825. Ruimy R, Breittmayer V, Elbaze P, et al. Phylogenetic analysis and assessment of the genera *Vibrio*, *Photobacterium*, *Aeromonas*, and *Plesiomonas* deduced from small-subunit rRNA sequences. *Int J Syst Bacteriol*. 1994;44:416–426.
826. Salerno A, Delétoile A, Lefevre M, et al. Recombining population structure of *Plesiomonas shigelloides* (Enterobacteriaceae) revealed by multilocus sequence typing. *J Bacteriol*. 2007;189:7808–7818. [Epub 2007 Aug 10].
827. Holmberg SD, Wachsmuth IK, Hickman-Brenner FW, et al. *Plesiomonas* enteric infections in the United States. *Ann Intern Med*. 1986;105:690–694.
828. Devreese K, Claeys G, Verschraegen G. Septicemia with *Ewingella americana*. *J Clin Microbiol*. 1992;30:2746–2747.
829. Kati C, Bibashi E, Kokolina E, et al. Case of peritonitis caused by *Ewingella americana* in a patient undergoing continuous ambulatory peritoneal dialysis. *J Clin Microbiol*. 1999;37:3733–3734.
830. Pien FD, Bruce AE. Nosocomial *Ewingella americana* bacteremia in an intensive care unit. *Arch Intern Med*. 1986;146:111–112.
831. Pound MW, Tart SB, Okoye O. Multidrug-resistant *Ewingella americana*: a case report and review of the literature. *Ann Pharmacother*. 2007;41:2066–2070. [Epub 2007 Oct 23].
832. Carter JE, Evans TN. Clinically significant *Kluyvera* infections: a report of seven cases. *Am J Clin Pathol*. 2005;123:334–338.

833. Sarria JC, Vidal AM, Kimbrough RCIII. Infections caused by *Kluyvera* species in humans. *Clin Infect Dis*. 2001;33:E69–E74.
834. Cheruvattath R, Balan V, Stewart R, et al. *Kluyvera* co-infection in two solid organ transplant recipients: an emerging pathogen or a colonizer bystander? *Transpl Infect Dis*. 2007;9:83–86.
835. Zhao WH, Hu ZQ. Epidemiology and genetics of CTX-M extended-spectrum β -lactamases in Gram-negative bacteria. *Crit Rev Microbiol*. 2013;39:79–101, Review. [Epub 2012 Jun 15].
836. Poirel L, Kämpfer P, Nordmann P. Chromosome-encoded Ambler class A beta-lactamase of *Kluyvera georgiana*, a probable progenitor of a subgroup of CTX-M extended-spectrum beta-lactamases. *Antimicrob Agents Chemother*. 2002;46:4038–4040.
837. Humeniuk C, Arlet G, Gautier V, et al. Beta-lactamases of *Kluyvera ascorbata*, probable progenitors of some plasmid-encoded CTX-M types. *Antimicrob Agents Chemother*. 2002;46:3045–3049.
838. Akhurst RJ, Boemare NE, Janssen PH, et al. Taxonomy of Australian clinical isolates of the genus *Phototrhaddus* and proposal of *Phototrhaddus asymbiotica* subsp. *asymbiotica* subsp. nov. and *P. asymbiotica* subsp. *australis* subsp. nov. *Int J Syst Evol Microbiol*. 2004;54:1301–1310.
839. Gerrard JG, McNevin S, Alfredson D, et al. *Phototrhaddus* species: bioluminescent bacteria as emerging human pathogens? *Emerg Infect Dis*. 2003;9:251–254.

Pseudomonas aeruginosa and Other *Pseudomonas* Species

Rafael Araos and Erika D'Agata

SHORT VIEW SUMMARY

Epidemiology

- *Pseudomonas* species are gram-negative bacteria that inhabit diverse environments, including soil, water, plants, insects, and animals.
- *P. aeruginosa* is the most important *Pseudomonas* species affecting humans, often causing serious infections associated with substantial morbidity and mortality.
- Other *Pseudomonas* species, including *P. fluorescens*, *P. luteola*, *P. putida*, and *P. stutzeri*, are less virulent but are still implicated in a wide variety of infections, primarily occurring among immunocompromised patients.
- *P. aeruginosa* has many virulence factors, including pili, flagella, enzyme secretion systems, and quorum-sensing molecules.

- *P. aeruginosa* is one of the most frequent pathogens implicated in hospital-acquired infections, including ventilator-associated pneumonia and catheter-associated urinary tract infections. Long-term acute care hospitals have very high rates of infections caused by *P. aeruginosa*.
- Characteristic infections caused by *P. aeruginosa* include ecthyma gangrenosum, malignant otitis externa, hot hand-foot syndrome, and hot tub folliculitis.
- Resistance to single and multiple antimicrobial agents is rising rapidly.

Microbiology

- *P. aeruginosa* is an aerobic gram-negative, rod-shaped bacteria.
- Growth occurs in a variety of culture media, forming smooth round colonies with a

characteristic grapelike or “corn-taco” odor and a green-blue coloration.

- Identification is based on colony morphology, coloration, oxidase positivity, and growth at 42°C.

Therapy

- The majority of evidence suggests that empirical treatment with combination therapy is indicated if antimicrobial resistance rates are high.
- Streamlining to a single antimicrobial agent, once antimicrobial susceptibility profiles are available, should then be considered.
- Effective antimicrobials include quinolones, carbapenems, and broad-spectrum cephalosporins ± β -lactamase inhibitors (Table 219.3)

Pseudomonas species are ubiquitous gram-negative bacteria capable of inhabiting a wide variety of diverse environments, including soil, water, plants, insects, and animals. Among all *Pseudomonas* species, *P. aeruginosa* is the most important species affecting humans and is responsible for serious debilitating and life-threatening infections.

P. aeruginosa infections were noted in the literature in the 1800s when physicians began to report a “condition” causing a blue-green discoloration on bandages and associated with a “peculiar” odor. The cause of the discoloration was first characterized by Fordos in 1869, who extracted the blue crystalline pigment called pyocyanin.¹ In 1882, Gessard verified “the parasitic origin of this phenomenon” using Pasteur’s cultures and isolated the organism, which was originally called *Bacillus pyocyaneus*. Initially, this pathogen was regarded as “a curiosity without any influence upon human pathology,” and “old surgeons looked upon blue pus on their dressings as rather a favorable sign.”¹ In 1894, Williams provided one of the first reviews of case reports of *B. pyocyaneus* infections.¹ He described septic patients with “hemorrhagic spots of a port-wine color” and pustules, with recovery of the organism from these skin lesions. Subsequently, more case reports of infections caused by *B. pyocyaneus* appeared in the literature.^{1–3} In the 1940s, Haynes provided detailed microbiologic characteristics of *P. aeruginosa* that would distinguish it from *Pseudomonas fluorescens*. During the Vietnam War, *P. aeruginosa* was recorded as one of the three most common wound pathogens.^{4,5} By the mid-1990s, *P. aeruginosa* became of great concern as a pathogen associated with burn infections and war-related wounds. *P. aeruginosa* is now considered to be of most concern because it causes a variety of infections associated with considerable morbidity and mortality, usually occurring among immunocompromised hosts. Furthermore, single-drug and multidrug resistance rates are particularly

high for this pathogen, which severely limits the therapeutic options available to treat infected patients.

MICROBIOLOGY

The pseudomonads are aerobic gram-negative, motile rods. They are ubiquitous in soil, water, plants, and animals and have numerous important ecologic roles (Fig. 219.1). The German botanist Walter Migula first used the term *Pseudomonas*, which is derived from the Greek *pseudo*, meaning “false,” and *monas*, meaning “unit.” Although the etymology was never explained, it has been postulated that Migula created this name because the bacteria resembled the cells of nonflagellate *Monas* in size and motility.⁶ *P. aeruginosa* is an obligate aerobic rod-shaped bacterium measuring 0.5 to 1.0 μm in width and 1 to 3 μm in length. It grows as a single bacterium, although it can occur in short chains. It grows on many types of culture media, forming smooth round colonies with a characteristic grapelike or “corn-taco” odor and green-blue coloration. The coloration is due to the production of pyocyanin (blue) and pyoverdin (green). This distinct color explains the species name of “*aeruginosa*,” a Latin word meaning “verdigris” or “copper rust.” Some strains produce other pigments, including pyorubin (dark red) and pyomelanin (black). Isolates from patients with cystic fibrosis (CF) may have a distinct mucoid appearance. *P. aeruginosa* is oxidase positive and grows at 37°C to 42°C. Growth at 42°C allows differentiation from other *Pseudomonas* species, including *P. fluorescens* and *P. putida*. Identification of *P. aeruginosa* is based on colony morphology, coloration, oxidase positivity, and growth at 42°C.

Strains of *P. aeruginosa* can produce an extracellular polysaccharide, referred to as alginate. Overproduction of this substance leads to the formation of a mucoid colony phenotype, which is usually present among




FIG. 219.1 Functional and environmental range of *Pseudomonas* species. The *Pseudomonas*' common ancestor has encountered a wide range of abiotic and biotic environments, which has led to the evolution of a multitude of traits and lifestyles with significant overlap among species. UV, Ultraviolet. (Modified from Silby MW, Winstanley C, Godfrey SA, et al. *Pseudomonas* genomes: diverse and adaptable. FEMS Microbiol Rev. 2011;35:652–680.)

isolates recovered from patients with CF and other chronic infections. Isolates recovered from the environment or those causing nosocomial infections are usually nonmucoid.^{7–9}

The *Pseudomonas aeruginosa* Genome

The *P. aeruginosa* genome is large (over 6 million base pairs)¹⁰ and complex. In contrast to other large bacterial genomes, the *Pseudomonas* genome does not contain an abundance of gene duplication events but instead contains numerous distinct gene families. This finding explains the great genetic and functional diversity of this pathogen.^{11,12}

The genome is composed of a relatively invariable core genome, which contains 90% of the total genome and includes the conserved gene sequences that encode metabolic and pathogenic factors present in the majority of *P. aeruginosa* strains. The accessory genome is highly variable and includes genes found only in certain *P. aeruginosa* strains. The genetic elements in the accessory genome result in distinct *P. aeruginosa* phenotypes, with niche-specific adaption. These genetic elements include virulence factors, resistance genes, and genes encoding specific catabolic pathways that allow persistence in harsh environments (pollutants, pesticides).¹¹

Pseudomonas aeruginosa and the Human Microbiota

The Human Microbiome Project, established by the National Institutes of Health, has begun to characterize the human microbial communities

colonizing healthy individuals, using 454 FLX Titanium platform pyrosequencing of 16S ribosomal RNA genes. Analysis of the human microbiota among 242 healthy adults from five major body areas (oral cavity and oropharynx, stool, vagina, nares, and skin [inner elbows and behind the ears]) revealed that *P. aeruginosa* was completely absent from both the skin and nares and that *Pseudomonas* species (not identified at the species level) were present in miniscule abundance in the stool and oral cavity, although more abundant in the latter. These data suggest that *P. aeruginosa* is not a common bacterium that inhabits the human microbiota among healthy hosts.¹³

Alterations in the microbiome lead to a decrease in colonization resistance (increasing colonization with antimicrobial-resistant bacteria) and a decrease in resilience (the microbiome's ability to recover). The absence or near absence of *P. aeruginosa* in the healthy human microbiome strongly suggests that for detectable long-term colonization to occur, a perturbation of the microbiome is necessary. Antimicrobial agents, other medications (e.g., anticholinergics), gastrointestinal diseases, and diet may be some of the necessary factors required for *P. aeruginosa* colonization.

VIRULENCE FACTORS

An abundance of virulence factors have been identified among various *P. aeruginosa* strains. These factors are either present on the bacterial cell surfaces or are secreted (Table 219.1). The majority of virulence factors, however, have been identified in cell lines or animal models

TABLE 219.1 Virulence Factors of *Pseudomonas aeruginosa* and Their Role in Pathogenesis

VIRULENCE FACTOR	ROLE IN PATHOGENESIS
Pili and flagella ¹⁴	Attachment to host cells, motility, biofilm formation
Type I secretion system (alkaline protease) ¹⁵	Delivery of toxins to extracellular spaces
Type II secretion system (elastase, exotoxin A, phospholipase A, protease IV) ¹⁵	Cytotoxicity, inflammation, colonization
Type III secretion system (exotoxins S, T, U, and Y) ^{16,17}	Tissue injury
Endotoxin (lipopolysaccharide)	Resisting host innate defenses
Alginate ²¹	Antiphagocytic activity, resists opsonic killing
Pyocyanin ²²	Tissue damage, inhibition of lymphocyte proliferation
Pyoverdinin ²²	Binds iron
Quorum-sensing molecules ^{18,19}	Cell-to-cell communication regulating virulence and biofilm formation

and therefore their role in human disease has not been clearly established. Major virulence factors are discussed in the following sections. Virulence factors associated with CF are discussed in Chapter 71.

Pili

Pili allow the bacteria to adhere to cell surfaces, are involved in biofilm formation, and mediate motility. Five *pilA* alleles have been identified (groups I to V) among *P. aeruginosa* strains. Type IV pili (T4P) has been extensively investigated.¹⁴ This group of pili is unique in that its members can mediate motility independent of flagella. Many *P. aeruginosa* T4P-expressing strains exhibit “twitching motility,” a jerky movement with an estimated velocity of approximately 1 mm/h (equivalent to 500 cell lengths per hour, assuming the average length of a *P. aeruginosa* cell is 2 μ m).¹⁴ The functions of twitching motility include biofilm formation and exploration of surfaces. Given their role in adherence, T4P are deployed during the early stages of acute infection.¹⁴ Production of pili is frequently lost in chronic infections, such as CF, with selection of strains with other phenotypes better suited for that environmental niche.

P. aeruginosa possesses a single polar flagellum, which plays an important role in motility, colonization, and biofilm formation, similar to pili.

Type I and II Secretion Systems

The type I secretion system (T1SS) secretes toxins in a one-step process into extracellular spaces. The most important toxin studied in the T1SS is alkaline protease, which inhibits fibrin formation and promotes dissemination of *P. aeruginosa*. Secretion of toxins by type II secretion system (T2SS) is a two-step process whereby toxins are synthesized as precursor proteins and then cleaved. Toxins excreted by this system include exotoxin A, phospholipase C, protease IV, and elastase, which mediate cytotoxic effects and inflammatory processes and promote colonization.¹⁵

Type III Secretion Systems

The type 3 secretion system (T3SS) is a complex secretory system that directly injects exotoxins into the cell cytoplasm. Four proteins have been identified. Exotoxin U (ExoU) is a phospholipase, which induces apoptosis as well as causes necrosis of phagocytes and parenchymal cells. Exotoxin Y (ExoY) is an adenylate cyclase, which may disrupt the barrier function of pulmonary endothelial cells. Exotoxins T and S (ExoT and ExoS) are bifunctional proteins that affect target cell growth by inhibiting DNA synthesis and inducing changes in the cytoskeleton and cellular morphology, thus affecting adherence. Although all strains harbor T3SS genes, only a few are capable of secreting these effector proteins under the conditions tested. The expression of the T3SS in *P.*

aeruginosa isolates may confer a worse clinical outcome among humans than nonexpressors.^{16,17}

Quorum-Sensing Molecules

Quorum sensing (QS) allows cell-to-cell communication and involves signaling molecules called autoinducers. The usual steps of QS involve the production of autoinducers followed by their active or passive release into the environment. These autoinducers are then recognized by specific receptors, resulting in changes in gene regulation.¹⁸ This complex signaling network allows the “community” of *P. aeruginosa* bacteria to react to different signals and thereby adapt to different niches. There are three QS systems present in *P. aeruginosa* isolates: two are referred to as the LuxI/LuxR-type QS circuits, and the third is referred to as the *Pseudomonas* quinolone signal system.^{18,19} These QS systems control the expression of virulence factors, including elastase, exotoxin A, and proteases.

QS also controls biofilm formation (see Chapter 71). Biofilms are a type of growth mode that results in clusters of bacterial colonies, encased in a biopolymer matrix, that attach to surfaces. Biofilms are predominantly formed on implantable devices or during chronic infections, such as osteomyelitis and CF. One of the main functions of biofilms is to reduce the efficacy of antimicrobial agents by impeding the agents’ ability to reach the bacteria.^{19,20}

Other Virulence Factors

A variety of other virulence factors produced by *P. aeruginosa* have been described. Endotoxin, or lipopolysaccharide, is a virulence factor located on the outer portion of the outer membrane and provides resistance to host defenses. Pyoverdins are siderophores that compete with host proteins for iron chelation. Pyocyanin reacts with oxygen to form oxygen radicals, causing tissue damage, and inhibits both lymphocyte proliferation and cilia function. Lastly, alginate is an extracellular polysaccharide, which has antiphagocytic activity and resists killing by opsonization. It is a scavenger of free radicals that are released by macrophages and inhibits neutrophil chemotaxis and complement activation. Its secretion results in a mucoid morphology seen on culture plates.^{21,22}

EPIDEMIOLOGY

P. aeruginosa is implicated in both community- and hospital-acquired infections, although it is much more common in the latter. In the United States, *P. aeruginosa* is the 6th most common pathogen implicated in all hospital-acquired infections, as reported in 2016 by the National Healthcare Safety Network (NHSN).²³ It is the second most common pathogen associated with ventilator-associated pneumonia²⁴ and ranks third among causes of catheter-associated urinary tract infections. *P. aeruginosa* is the 5th most common pathogen causing surgical site infections and 10th most common pathogen causing catheter-related bloodstream infections (BSIs). Patients at high risk for *P. aeruginosa* hospital-acquired infections include those admitted to an intensive care unit (ICU) and those with burns, neutropenia, or CF. These patient populations are discussed in this section and also in Chapter 71.

Rates of *P. aeruginosa* implicated in infections in long-term acute care hospitals (LTACHs) are even higher than rates reported from hospital settings. LTACHs are defined as health care facilities that are accredited as acute care hospitals with an average annual length of stay of at least 25 days for Medicare patients, as per the Centers for Medicare and Medicaid Services. Patients admitted to these LTACHs usually require prolonged care after hospitalization, including hemodialysis, mechanical ventilation, intravenous medication, and wound care. In contrast to the hospital setting, *P. aeruginosa* is the most common cause of catheter-associated urinary tract infections and ventilator-associated pneumonia in LTACHs.²⁵ Even when rates of *P. aeruginosa* infections in LTACHs are compared with those from ICUs, *P. aeruginosa* is still a more frequent pathogen implicated in LTACH infections. For example, in 2010, 19% of catheter-associated urinary tract infections in patients in LTACHs were caused by *P. aeruginosa* compared with 9% to 12% of those occurring in ICUs. Ventilator-associated pneumonia caused by *P. aeruginosa* was also more frequent in LTACHs (35%) compared with that occurring in ICUs (17%–20%). The higher occurrence of *P. aeruginosa* infections

is likely due to an older, more debilitated patient population with excessive health care exposure.²⁵

Transmission Dynamics of *Pseudomonas aeruginosa* and Reservoirs

Acquisition of *P. aeruginosa* can occur both exogenously and endogenously. Exogenous acquisition occurs through contaminated hands of health care workers and environmental surfaces. An in-depth epidemiologic study characterized the exogenous transmission of multidrug-resistant (MDR) *P. aeruginosa* in six ICUs.²⁶ The investigators obtained cultures from hands, gloves, and gowns of health care workers during routine patient care activities, surveillance cultures from patients, and environmental samples from sinks, bedrails, vital sign monitors, supply carts, door handles, intravenous pumps, ventilators, and floors. Molecular typing, using pulsed-field gel electrophoresis, was performed to determine clonal relatedness among strains. In that study, MDR *Acinetobacter* species were the most common pathogens to contaminate health care workers' gloves and hands, occurring among 33% of interactions between health care workers and patients. *P. aeruginosa* was the second most common MDR pathogen to contaminate health care workers, which occurred during 17.4% of health care worker–patient interactions. Independent risk factors associated with health care worker contamination were presence of environmental contamination, duration in patients' room greater than 5 minutes, performing a physical examination, and contact with the mechanical ventilator. Environmental contamination was also very common: *P. aeruginosa* was recovered from 22% of rooms.²⁶ This study and many others emphasize that exogenous transmission plays a major role in the nosocomial acquisition of *P. aeruginosa* and that environmental contamination is central to its transmission to patients and health care workers.

Endogenous acquisition of a resistant strain of *P. aeruginosa* is defined as colonization with an antimicrobial-susceptible strain that subsequently becomes resistant primarily through antimicrobial selective pressure within the host. A study of imipenem-resistant *P. aeruginosa* transmission demonstrated that, among events that could be determined, endogenous acquisition accounted for 19% of identified acquisition events and that exogenous acquisition accounted for 31% of events.²⁷

As outlined earlier, environmental reservoirs contribute substantially to the spread of *P. aeruginosa*. The most common sites either have high moisture or humidity or are water related (Table 219.2). In the hospital setting, outbreaks of *P. aeruginosa* have been linked predominantly to water sources, including potable water, showerheads, and sinks (see Table 219.2).^{28,29} Other sources have included health care workers' artificial or long nails, intraocular lens solution, ultrasound transmission gel during transesophageal echocardiography, retained tissue in surgical instruments,³⁰ and soap dispensers. The ability of *P. aeruginosa* to form biofilms on surfaces increases its ability to survive on inanimate surfaces and makes it difficult to eradicate. Biofilms are microbial communities held together by structural polysaccharides (slime), which attach strongly

to surfaces. Biofilms produced by *P. aeruginosa* lead to antimicrobial tolerance and impede eradication by environmental cleaning agents.^{19,28}

True community-acquired infections among people without any prior exposure to a health care setting are rare. The rarity of community-acquired infections reflects the fact that *P. aeruginosa* is not part of the healthy human microbiota¹³ and that colonization occurs predominantly after hospitalization and antimicrobial exposure. Community-acquired infections, however, have been reported, including outbreaks from contaminated community reservoirs and among intravenous drug users. As with the hospital setting, water-related reservoirs are the main sources of *P. aeruginosa* in the community and include whirlpools, hot tubs, contact lenses, home humidifiers, water-damaged houses, swimming pools, loofah sponges, and even holy water.^{31,32} Contaminated recreational water, however, is among the most common sources of *P. aeruginosa* outbreaks.^{33,34} A review of outbreaks associated with recreational water, from 1971 to 2000, identified *P. aeruginosa* as the second most frequent causative pathogen, after *Cryptosporidium* species. *P. aeruginosa* was implicated in 36 outbreaks during the study period, with most of the outbreaks associated with whirlpool baths, hot tubs, and swimming pools. The main presentation is that of a superficial folliculitis that is pruritic and maculopapular and progresses to vesiculopustular within hours to days after exposure. It remits spontaneously. Conjunctivitis and otitis externa were also reported.³³ Common nonmedical terms associated with community-acquired *P. aeruginosa* infections include “hot tub rash,” “swimmer's ear,” and “hot hand-foot syndrome” (see “Skin and Soft Tissue Infections,” later).

Food products may also be sources of *P. aeruginosa* in both community and hospital settings, because this pathogen has been isolated in a variety of vegetables, including lettuce, mushrooms, and carrots, and is present in soil.^{35–37} A definitive link between the presence of *P. aeruginosa* in vegetables and subsequent colonization or infection, however, has not been clearly established. Nevertheless, it is plausible that ingestion of *P. aeruginosa*-contaminated raw vegetables by high-risk individuals, including neutropenic patients and other immunocompromised hosts and those with decreased gastric acidity or oropharyngeal invasive devices,^{38,39} may increase the risk for colonization.

ANTIMICROBIAL RESISTANCE

P. aeruginosa possesses a plethora of different resistance mechanisms. It is therefore not surprising that rates of antimicrobial resistance and multidrug resistance are among the highest in these organisms compared with other common human pathogens. Of even greater concern is the paucity of novel antimicrobial agents being developed to combat *P. aeruginosa* infections. The US Food and Drug Administration (FDA), however, approved ceftolozane-tazobactam in 2014. It has enhanced affinity for *P. aeruginosa* penicillin-binding proteins and appears to be unaffected by loss of porin channels or upregulation of efflux pumps. Over 80% of isolates resistant to ceftazidime or meropenem retain susceptibility to this novel agent. Currently, it is approved for use in urinary tract and intraabdominal infections,⁴⁰ and several reports suggest that it may also be useful for treating other severe infections caused by MDR *P. aeruginosa* with confirmed susceptibility to ceftolozane-tazobactam^{41,42} (see Chapter 21).

Rates of antimicrobial resistance among *P. aeruginosa* hospital isolates recovered from different types of health care–associated infections in the United States in 2014, as reported by the NHSN, were as follows: aminoglycosides, 7% to 21%; extended-spectrum cephalosporins, 10% to 27%; fluoroquinolones, 12% to 33%; carbapenems, 8% to 28%; and piperacillin ± tazobactam, 7% to 19%. Multidrug resistance, defined as resistance to three or more of these antimicrobial classes, was present among 4% to 20% of isolates.²³ Similarly to previous reports, isolates from ventilator-associated pneumonia had the highest rates, and those from surgical site infections had the lowest rates.

The 2016 Annual Report of the European Antimicrobial Resistance Surveillance Network (EARS-Net) reported rates of *P. aeruginosa* resistance to piperacillin ± tazobactam, ceftazidime, fluoroquinolones, aminoglycosides, and carbapenems. Overall, resistance rates to ceftazidime increased from 2013 to 2016. Conversely, the mean percentages of resistance for fluoroquinolones, aminoglycosides, and carbapenems significantly decreased over the same period of time, whereas resistance

TABLE 219.2 Environmental Reservoirs of *Pseudomonas aeruginosa*

Hospital Reservoirs of *P. aeruginosa*

Sinks, taps, showerheads
Potable water
Respiratory therapy equipment
Flower vases, ice makers
Hydrotherapy pools
Cleaning equipment (mops, buckets)
Bronchoscopes, endoscopes
Resuscitators
Water baths
Multidose vials

Community Reservoirs of *P. aeruginosa*

Home humidifiers
Whirlpools, hot tubs, spas
Swimming pools
Water-damaged homes

to piperacillin \pm tazobactam remained stable. In 2016, 33.9% of isolates were resistant to one or more of the five antimicrobial classes, 13.6% were resistant to three or more, and 4.4% were resistant to all five classes.⁴³ Resistance rates to three or more groups of antimicrobials varied considerably between European countries, with lowest percentages reported from Iceland, Denmark, Luxembourg, the United Kingdom, the Netherlands, and Norway (<3%) and highest percentages from Bulgaria and Romania (36%–49%). Percentages of single-drug resistance among *P. aeruginosa* isolates obtained from Latin America and the Asia-Pacific rim range from 23% to 41%.^{44,45}

In the United States in 2010, resistance rates in LTACH units were reported to be even higher than in ICUs. For example, MDR *P. aeruginosa* was more common among patients with catheter-associated urinary tract infections in LTACHs (25%) compared with those occurring in patients in ICUs (12%). Similarly, rates of multidrug resistance in patients with catheter-associated BSI were higher among those in LTACHs (16%) than those in ICUs (2%–9%).²⁵

Resistance Mechanisms

P. aeruginosa harbors numerous resistance mechanisms that either decrease penetration to the target site, alter the target site, or inactivate the antimicrobial agent using bacterial enzymes. These mechanisms can be broadly categorized as intrinsic,^{46–48} acquired, or adaptive, with overlap between categories.

Intrinsic Resistance

Decreased Permeability of the Outer Membrane

The semipermeable outer membrane of *P. aeruginosa* allows important nutrients to enter the cell through channels present on the cell membranes called porins. Numerous antimicrobial agents, including β -lactams, aminoglycosides, tetracyclines, fluoroquinolones, and carbapenems, enter the bacterial cell through these porins. Several families of porins have been characterized, including OprF, OprD, OprM, and TonB. Although OprF is a major porin in the outer cell membrane, allowing transport of large substrates, its role in antimicrobial resistance has not been definitively proven because loss of OprF porins in mutant strains does not alter antimicrobial penetration. Among the remaining three families, OprD has been the most studied. This channel allows entry of carbapenems but not of other β -lactams. Loss of OprD, however, may not have an equal impact on minimal inhibitory concentration increases for all carbapenems equally. Lastly, some porins in the family of OprM are presumed to be part of efflux systems and are discussed in the next section.

Not all antimicrobial agents use porins to enter the cell but instead decrease cell membrane stability by binding to lipopolysaccharides on the outer cell membrane. Examples of such agents include aminoglycosides and polymyxins.

Efflux Pumps

As the name implies, efflux pumps actively pump antimicrobial agents out of the bacteria. These pumps confer resistance to the great majority of antimicrobial agents (with the exception of polymyxins) and are the predominant systems for multidrug resistance among *P. aeruginosa*. Five superfamilies of efflux pumps have been identified, of which the resistance-nodulation-division (RND) family is among the most common. MexAB-OprM and MexXY-OprM efflux systems of the RND family confer intrinsic multidrug resistance to numerous antimicrobial agents, including fluoroquinolones, aminoglycosides, β -lactams, tetracyclines, tigecycline, and chloramphenicol. MexAB-OprM also confers resistance to meropenem but not imipenem, thus explaining differences in susceptibility patterns among different carbapenems. Similarly, the MexXY-OprM efflux system removes cefepime but not ceftazidime.⁴⁹

Antimicrobial-Modifying Enzymes

The majority of antimicrobial-modifying enzymes are acquired on plasmids and are discussed later in the chapter,⁴⁶ with the exception of AmpC, a chromosomally encoded inducible cephalosporinase. AmpC confers resistance to all β -lactams except fourth-generation cephalosporins and carbapenems. When AmpC is overproduced, through mutations, resistance may also be conferred to these antimicrobial classes.

Therapeutic failure due to the emergence of resistance during appropriate therapy can occur in over 50% of patients, especially those with serious *P. aeruginosa* infections and those with neutropenia or CF.

Acquired Resistance

Acquired resistance genes predominantly confer resistance to β -lactams and aminoglycosides.⁴⁷ Extended-spectrum β -lactamases (ESBLs) are plasmid mediated and confer resistance to penicillins, narrow- and extended-spectrum cephalosporins, aztreonam, and sometimes carbapenems. ESBL-producing *P. aeruginosa* strains have rapidly spread worldwide. ESBL families identified in *P. aeruginosa* include PER, VEB, GES, TEM, SHV, and CTX-M enzymes. GES-type enzymes also extend their activity to carbapenems. These enzymes have been recovered from isolates from China, South Africa, Brazil, and France.⁴⁷ Oxacillinase β -lactamases (OXAs) can either be narrow spectrum or broad spectrum and are weakly inhibited by clavulanic acid. Carbapenemase-hydrolyzing oxacillinases, which can be either acquired or naturally occurring, have also been identified in *P. aeruginosa* isolates, although less frequently than in *Acinetobacter* species. Resistance to carbapenems can also occur via metallo-carbapenemases and include the Verona integron-encoded metallo- β -lactamase (VIM), IMP, and New Delhi metallo- β -lactamase (NDM) families. These enzymes are of great concern because they are active against penicillins and cephalosporins as well as carbapenems. VIM-type isolates were first reported in *P. aeruginosa* isolates recovered in Italy in 1997 and subsequently spread into members of the Enterobacteriaceae, especially *Klebsiella pneumoniae*. Isolates with NDM enzymes may also carry aminoglycoside and fluoroquinolone resistance genes and remain susceptible only to colistin and polymyxin B.⁵⁰ Another enzyme active against carbapenems, present in *P. aeruginosa* isolates, is the *K. pneumoniae* carbapenemase (KPC) type. KPC-producing strains were first identified in Colombia and have since spread to Puerto Rico, China, and the United States, and most recently Brazil.⁵¹ Carbapenem-resistant isolates, regardless of the mechanism of resistance, result in serious infections. A longitudinal study from 1989 to 2006 demonstrated that rates of imipenem-resistant *P. aeruginosa* isolates increased from 13% to 20% and that infections caused by these resistant pathogens were associated with higher in-hospital mortality. Prior exposure to carbapenems increased the risk for imipenem-resistant *P. aeruginosa* infection by almost eightfold.⁵²

Aminoglycoside-Modifying Enzyme

Aminoglycoside-modifying enzymes are carried on multiple different genetic mobile elements, such as plasmids, transposons, and integrons. They confer resistance to all aminoglycosides, although, in general, amikacin may be less susceptible to these enzymes. The most common aminoglycoside-modifying enzymes are aminoglycoside nucleotidyl-transferase (2')-I, which confers resistance to gentamicin and tobramycin, and aminoglycoside acetyltransferase (6')-II, which also confers resistance to netilmicin.

TREATMENT OF PSEUDOMONAS AERUGINOSA INFECTIONS

Treatment of *P. aeruginosa* infections, especially BSIs, centers around the controversy of monotherapy versus combination therapy. A few earlier studies support the use of combination therapy since mortality rates were lower when two antimicrobial agents, instead of a single agent, were used to treat BSI caused by *P. aeruginosa*.⁵³ The main limitation of these studies is that in many the monotherapy study arm consisted of only an aminoglycoside, which is suboptimal for the treatment of *P. aeruginosa* BSI.⁵³ In a meta-analysis, which also concluded that combination therapy is superior to monotherapy, four of the five included studies used aminoglycosides in the monotherapy study arm.⁵⁴ There are numerous other limitations with older studies that support the use of combination therapy, including lack of double blinding and randomization, different sources of BSI, retrospective lack of adjustment for time to start of appropriate antimicrobial therapy, and duration of follow-up. Confounding by indication, whereby the severely ill patients receive combination therapy, is another limitation of studies addressing this issue. Another reason that is often cited for supporting combination therapy is the potential synergy between the two antimicrobial agents,

usually a β -lactam and an aminoglycoside. Although in vitro and animal studies show benefit of this combined regimen, clinical studies have provided conflicting data.^{55,56} Preventing the emergence of antimicrobial resistance is another reason often cited for supporting the use of combination therapy, but there are minimal data to support this statement. Administering the appropriate dose, at the correct frequency and for the optimal duration, is likely more important in preventing the emergence of resistance than combination therapy. Prompt initiation of the appropriate antimicrobial agents is also key to a successful outcome, as is removal of invasive devices if implicated.

Overall, the great majority of more recent studies do not show a survival benefit between combination therapy and monotherapy for definitive therapy.^{57–61} However, the main conclusion by most investigators is that large randomized clinical trials are needed to definitively answer the question of efficacy between combination therapy and monotherapy.

Use of combination therapy should be strongly considered in the severely ill patient for the empirical treatment of *P. aeruginosa* BSI, especially in those health care institutions with patients with a high rate of multidrug resistance. Using combination therapy will thus ensure that at least one antimicrobial agent is effective against the infecting *P. aeruginosa* strain. Deescalating to a single antimicrobial agent, once antimicrobial susceptibility profiles are available, should then be considered. Narrowing to a single agent is especially relevant when the combination therapy includes an aminoglycoside, because this regimen is associated with increased nephrotoxicity.^{60,61} Other adverse events that are more likely to occur with combination therapy, compared with monotherapy, include an increased risk for *Clostridioides difficile* (formerly *Clostridium difficile*) infection, further alterations in the protective effects of the human microbiota against colonization by other MDR organisms, and fungal infections.

Antimicrobial agents effective against *P. aeruginosa* and appropriate doses are listed in Table 219.3. Polymyxins (colistin, polymyxin B) should be reserved for MDR *P. aeruginosa*, when other alternatives are not

available, because this class of agents is inferior to other available antipseudomonal agents.⁶² Emergence of resistance during treatment has been documented for *K. pneumoniae* BSI and likely also occurs for *P. aeruginosa* infections.⁶³ High-dose continuous infusion of β -lactams has also been successfully used in the treatment of BSI caused by MDR *P. aeruginosa*. Moriyama and associates reported three patients infected with MDR *P. aeruginosa* strains who were successfully treated with a continuous infusion of ceftazidime (6.5–16.8 g/day) or aztreonam (8.4 g/day) and tobramycin.⁶⁴ The rationale behind this approach is that antibacterial activity of β -lactams depends on the time that the antimicrobial concentration is above the minimal inhibitory concentration of the bacteria. Using continuous infusion ensures that the concentration of the β -lactam will be above the minimal inhibitory concentration for the entire dosing interval, whereas intermittent dosing may cause the concentration to fall below the minimal inhibitory concentration. Future clinical studies, however, are required to further validate this treatment regimen.

Novel antipseudomonal agents are reviewed in detail by Wright and colleagues.⁶⁵ Several studies indicate that ceftolozane-tazobactam and ceftazidime-avibactam are effective in the treatment of MDR-*P. aeruginosa* BSI and other infections, although resistance can develop even with short courses of therapy.^{41,42,66,67} The combination of ceftazidime and the novel non- β -lactam β -lactamase inhibitor avibactam has been approved for the treatment of complicated urinary tract infections and intraabdominal infections.⁶⁵ The addition of avibactam to ceftazidime allows this combination to inhibit Ambler class A β -lactamases (including KPC), AmpC β -lactamases, and OXA-type Ambler class D β -lactamases, making this drug an excellent therapeutic option for infections caused by MDR gram-negative bacteria, including *P. aeruginosa*.⁶⁷ The recommended dose for patients with normal renal function is 2 g of ceftazidime and 500 mg of avibactam every 8 hours. Resistance to ceftazidime-avibactam among *P. aeruginosa* has been described, with higher rates among MDR isolates involving efflux mechanisms or metallo-carbapenemases.^{68,69}

INFECTIONS CAUSED BY *PSEUDOMONAS AERUGINOSA* Bloodstream Infections

BSIs are among the most serious infections caused by *P. aeruginosa*, with mortality rates reaching 60%. The nationwide Surveillance and Control of Pathogens of Epidemiological Importance (SCOPE), which included data from 49 US hospitals, reported that from 1995 to 2002 the incidence of *P. aeruginosa* nosocomial BSI was 2.1 per 10,000 hospital admissions. *P. aeruginosa* was the third most common gram-negative bacteria causing nosocomial BSI and accounted for 4.3% of all cases. In the ICUs, *P. aeruginosa* was the fifth most common isolate implicated in BSI, accounting for 4.7% of all cases, and was the seventh most common isolate in non-ICU wards, accounting for 3.8% of cases.⁷⁰ Outside the United States, *P. aeruginosa* is implicated in even more cases of nosocomial BSI.⁷¹ In a surveillance study that collected data from 16 Brazilian hospitals from 2007 to 2010 and used the same methodology as the SCOPE study, 8.9% of all nosocomial BSI were caused by *P. aeruginosa*.

Risk factors for *P. aeruginosa* BSI include immunodeficiency, prior hospitalization, previous antimicrobial exposure, advanced age, prior surgery, and invasive devices.^{72–74} Many of these risk factors represent an association with BSI, irrespective of the implicated pathogen.

Mortality rates for nosocomial BSI caused by *P. aeruginosa* are among the highest. The great majority of reported crude mortality percentages from large surveillance studies range from 39% to 60%. These percentages are similar to those caused by *Candida* species.^{70,71} Some studies, however, report lower mortality rates, ranging from 12% to 30%.^{75–77} The large range in mortality rates from different studies reflects the multitude of factors that affect outcomes associated with BSI. For *P. aeruginosa* BSI, advanced age, high Acute Physiology and Chronic Health Evaluation II (APACHE II) score, sepsis, poor functional status, polymicrobial bacteremia, and inappropriate initial antimicrobial therapy have all been associated with an increased risk for mortality.^{73,76–79}

Multidrug resistance is also a risk factor for increased mortality. Rates of mortality in infections due to MDR *P. aeruginosa* strains are

TABLE 219.3 Intravenous Antimicrobial Agents Effective Against *Pseudomonas aeruginosa*

ANTIMICROBIAL AGENT	INTRAVENOUS DOSE
Penicillins Plus β-Lactamase Inhibitor	
Ticarcillin-clavulanate	3.1 g q4h
Piperacillin-tazobactam	4.5 g q6h or 3.375 g q4h
Broad-Spectrum Cephalosporins \pm β-Lactamase Inhibitor	
Ceftazidime	2 g q8h
Cefepime	2 g q8–12h
Ceftolozane-tazobactam	1.5g q8h
Ceftazidime-avibactam	2.5 g q8h
Fluoroquinolones	
Ciprofloxacin	400 mg q8h
Levofloxacin	750 mg q24h
Carbapenems	
Imipenem	500 g q6h
Meropenem	1–2 g q8h
Doripenem	500 mg q8h
Monobactam	
Aztreonam	2 g q8h
Aminoglycosides	
Tobramycin	2 mg/kg loading dose, then 1.7 mg/kg q8h or 4–7 mg/kg q24h
Gentamicin	As for tobramycin
Amikacin	7.5 mg/kg q12h or 15 mg/kg q24h

For a review of novel antimicrobials with potential activity, refer to Wright et al.⁶⁵

twofold to threefold higher compared with non-MDR strains.^{75,80} Because inappropriate initial empirical therapy is a major contributor to these higher mortality rates, combination therapy for empirical treatment is warranted when multidrug resistance rates are high (see later).⁸¹

The predominant distinguishing feature of *P. aeruginosa* BSI is the occurrence of ecthyma gangrenosum. Although not pathognomonic for *P. aeruginosa*, the presence of these characteristic skin lesions should raise high suspicion for this pathogen (see “Skin and Soft Tissue Infections” later).

Infective Endocarditis

P. aeruginosa accounts for 3% of all cases of infective endocarditis (IE).⁸² Among non-HACEK pathogens (species other than *Haemophilus* species, *Aggregatibacter actinomycetemcomitans*, *Cardiobacterium hominis*, *Eikenella corrodens*, or *Kingella* species), *P. aeruginosa* is the second most common gram-negative pathogen causing IE after *Escherichia coli*.⁸³ There are no unique clinical characteristics of IE caused by *P. aeruginosa*, although the presence of ecthyma gangrenosum should raise suspicion. Complications are common, and mortality rates are high, ranging from 36% to 60%.⁸³

Intravenous drug users are the patient population at highest risk for *P. aeruginosa* IE. The great majority have used tripeleminamine and pentazocine with contaminated water or paraphernalia. Patients predominantly present with right-sided endocarditis. Complications are frequent and include sepsis, embolization, and congestive heart failure.⁸⁴ In this patient population, polymicrobial IE, with both *P. aeruginosa* and *Staphylococcus aureus*, can also occur.⁸⁴

Left-sided IE in patients without intravenous drug use also occurs, although infrequently. It is predominantly a nosocomial infection occurring after invasive procedures, including cardiac or urogenital procedures. Infected intravascular catheters have also been implicated in IE. Of note, valvular disease is not a necessary predisposing risk factor for *P. aeruginosa* left-sided IE.⁸⁵ Splenic abscesses, neurologic sequelae, and ring and annular abscesses are frequent complications.⁸⁶

The American Heart Association recommends, as per expert opinion, that medical treatment of *P. aeruginosa* IE should include an extended-spectrum penicillin (ticarcillin, piperacillin) and ceftazidime or cefepime in full doses in combination with high-dose tobramycin (8 mg/kg/day IV or IM in once-daily doses) with peak concentrations of 15 to 20 µg/mL and trough concentrations of less than or equal to 2 µg/mL. This combination should be given for at least 6 weeks. Use of ciprofloxacin in combination with an aminoglycoside should be used with caution because ciprofloxacin resistance can occur during therapy. Other regimens that have been successful in a small number of patients include the combination of imipenem and an aminoglycoside.⁸⁷

Case reports have also shown success with other combinations of antimicrobial agents. Among two renal transplant recipients with no valvular heart disease who developed *P. aeruginosa* nosocomial IE, 6 weeks of therapy with imipenem and ciprofloxacin was successful in the first patient and 2 weeks of imipenem and amikacin followed by 6 weeks of imipenem alone was successful for the second patient (amikacin was stopped early owing to nephrotoxicity). Both patients' conditions were stable at 6 months of follow-up.⁸⁸

The need for surgical intervention differs between right- and left-sided IE. Although medical treatment is usually sufficient for right-sided native valve IE, in refractory cases surgical intervention with partial tricuspid valvectomy or “vegetomy” without valve replacement may be necessary.⁸⁷ Left-sided IE and prosthetic valve IE usually require early surgical intervention, owing to the high risk for complications and high failure rates associated with medical therapy alone. Survival rates with medical therapy alone among patients with left-sided IE, for example, are only 14%.⁸⁵

Pneumonia Pneumonia Associated With Hospital Exposure

Three categories of *P. aeruginosa* pneumonia can occur as a result of hospital exposure: (1) hospital-acquired (nosocomial) pneumonia, defined as a pneumonia that occurs 48 hours or more after hospital admission and that was not incubating at the time of admission; (2) ventilator-associated

pneumonia, defined as a hospital-acquired pneumonia that develops more than 48 to 72 hours after endotracheal intubation; and (3) health care-associated pneumonia, defined as a pneumonia that occurs among nonhospitalized patients who (a) have had an acute care hospitalization for 2 or more days within 90 days of the infection, (b) live in a nursing home or long-term care facility, (c) received intravenous antimicrobial therapy, chemotherapy, or wound care in the previous 30 days of the current infection, or (d) attended a hospital or dialysis clinic in the previous 30 days of the current infection.⁸⁹ These definitions were developed for the 2005 American Thoracic Society (ATS) and Infectious Diseases Society of America (IDSA) guidelines.⁸⁹

Hospital-Acquired (Nosocomial) Pneumonia

P. aeruginosa is the most common gram-negative bacteria implicated in hospital-acquired pneumonia.²³ The great majority of *P. aeruginosa* hospital-acquired pneumonia cases are late onset, occurring after the fifth day of hospitalization, similar to hospital-acquired pneumonia caused by methicillin-resistant *S. aureus*.^{90,91} Factors that predispose hospitalized patients to hospital-acquired pneumonia include older age, duration of mechanical ventilation, prior antimicrobial exposure, transfer from a medical unit or ICU, and admission to a ward with a high prevalence of *P. aeruginosa*, reflecting colonization pressure.⁹¹ Mortality rates are extremely high, ranging from 42% to 87%. As with BSI, the wide range reflects the presence of comorbidities, inadequate initial therapy, severity of illness at presentation, and multidrug resistance. The presence of bacteremia is associated with particularly poor outcomes. Hospital-acquired pneumonia can be associated with mechanical ventilation or not, although the former is much more common.

Hospital-Acquired Pneumonia Among Ventilated Patients

P. aeruginosa is the second most common pathogen implicated in ventilator-associated pneumonia after *S. aureus*, as per the 2011 to 2014 NHSN data, which reported rates of hospital-acquired infections from 4515 hospitals in the United States.²³ Among all gram-negative bacteria, *P. aeruginosa* is the most common cause of ventilator-associated pneumonia.^{23,90}

Colonization with *P. aeruginosa* is a prerequisite for developing a *P. aeruginosa* ventilator-associated pneumonia. Once the patient becomes colonized, breaches in integrity of the airway mucosa caused by intubation result in tissue invasion by the bacteria, leading to pneumonia. Sources for *P. aeruginosa* colonization are either endogenous or exogenous. Endogenous sources include *P. aeruginosa* colonizing the oropharynx and stomach with subsequent aspiration.⁹² Exogenous sources include health care workers carrying the organism and inanimate surfaces. Contaminated water sources are a particularly common reservoir for exogenous acquisition of *P. aeruginosa* implicated in ventilator-associated pneumonia. Numerous *P. aeruginosa* outbreaks have been traced to contaminated faucets, sinks, and tap water. Contaminated bronchoscopes have also been implicated as a source of *P. aeruginosa* ventilator-associated pneumonia. These cases were linked to inadequate disinfection procedures, contaminated disinfecting machines, and manufacturing defects of bronchoscopes.⁹³

Antimicrobial resistance among *P. aeruginosa* strains implicated in ventilator-associated pneumonia is high. Reported percentages from US hospitals in 2014 are as follows: aminoglycosides, 18.2%; cefepime-ceftazidime, 25.7%; ciprofloxacin-levofloxacin, 31.9%; imipenem-meropenem, 28.4%; and piperacillin-tazobactam, 19.4%. Multidrug resistance, defined as resistance to three or more antimicrobial classes, was also frequent, occurring in 19.9% of strains.²³ These high rates support the use of combination therapy as empirical treatment for ventilator-associated pneumonia to ensure that at least one antimicrobial agent is active against *P. aeruginosa*. In a study of 100 consecutive patients with bacteremic *P. aeruginosa* pneumonia, adequate empirical combination therapy was associated with significantly lower 28-day mortality rates.⁹⁴

The ATS/IDSA guidelines recommend empirical antimicrobial therapy for ventilator-associated pneumonia with an antipseudomonal β-lactam plus either an antipseudomonal quinolone or an aminoglycoside. The choice of specific antimicrobial agents should take into account the local antimicrobial susceptibility patterns and avoiding agents to which

the patient was recently exposed. If aminoglycosides are chosen, consideration should be given toward a short (5-day) course, to minimize nephrotoxicity, when used in combination with a β -lactam to treat *P. aeruginosa* pneumonia.⁸⁹ Once susceptibility profiles are available, monotherapy also should be considered because data supporting combination therapy are lacking.^{89,94} Aerosolized antimicrobial agents (aminoglycosides, polymyxins) can be used as adjunctive therapy to systemic antimicrobial agents in patients with highly resistant *P. aeruginosa* ventilator-associated pneumonia.⁹⁵

Duration of antimicrobial therapy for *P. aeruginosa* ventilator-associated pneumonia should be longer than for other pathogens. A prospective randomized, double-blind study evaluated the efficacy of 8 days versus 15 days of antimicrobial therapy for ventilator-associated pneumonia and showed that, for ventilator-associated pneumonia caused by *P. aeruginosa*, an 8-day course was suboptimal and was associated with higher recurrence rates.⁹⁶ A 15-day course is therefore recommended.⁸⁹

Hospital-Acquired Pneumonia Among Nonventilated Patients

P. aeruginosa is also the most common gram-negative bacteria implicated in pneumonia among nonventilated hospitalized patients, accounting for 9% of all cases in this category.⁹⁰ These high percentages warrant antipseudomonal antimicrobial agents in the empirical treatment of these types of infections, similar to that used for patients with ventilator-associated pneumonia.⁸⁹

Health Care–Associated Pneumonia

Health care–associated pneumonia develops outside the hospital among patients who have had recent substantial exposure to the health care setting. *P. aeruginosa* is implicated in 2% to 25% of health care–associated pneumonia cases.^{97–99} Surveillance studies report that it is the most common or second most common gram-negative pathogen causing health care–associated pneumonia.^{98,99} Because *P. aeruginosa* strains implicated in this type of pneumonia are more likely to be resistant to antimicrobial agents because of prior health care exposure, the ATS/IDSA guidelines recommend that treatment regimens should follow those for health care–associated pneumonia, as outlined previously.⁸⁹ However, these recommendations may not apply in health care settings and countries with low rates of antimicrobial resistance; therefore treatment regimens recommended for community-acquired pneumonia may be more appropriate in these settings.⁹⁷

A retrospective, observational study compared characteristics of health care–associated and community-acquired pneumonia among patients admitted to a tertiary care hospital in South Korea from 2008 to 2010. Among 31% of patients with pneumonia in whom a pathogen was identified, *P. aeruginosa* was the second most common gram-negative bacteria, after *K. pneumoniae*, and accounted for 20% of health care–associated pneumonia and 11% of community-acquired pneumonia. These percentages were not statistically different. The percent of bacterial species, including *P. aeruginosa*, that were “potentially drug-resistant” was significantly higher among the health care–associated group (32%) than the community-acquired group (15%).⁹⁹ The investigators included the following bacteria as “potentially drug-resistant”: methicillin-resistant *S. aureus*, *Pseudomonas* species, *Acinetobacter* species, *Stenotrophomonas maltophilia*, and extended-spectrum β -lactamase–producing Enterobacteriaceae. Actual antimicrobial susceptibility profiles, however, were not determined.

Compared with community-acquired pneumonia, patients presenting with health care–associated pneumonia are more likely to be older and have more comorbidities, a higher pneumonia severity index, and a worse functional status at presentation.^{97,98} Mortality percentages are subsequently higher for health care–associated pneumonia compared with community-acquired pneumonia and range from 17% to 20%.^{98,99}

Community-Acquired Pneumonia

Reported percentages of *P. aeruginosa* implicated in community-acquired pneumonia vary from 0.3% to 17%.^{97–100} Although *P. aeruginosa* is not usually considered to be a common cause of community-acquired pneumonia, the high percentages reported by some studies suggest otherwise.^{98,99} A surveillance study on the microbiology of pneumonia

in 59 US hospitals from 2003 to 2004 found that *P. aeruginosa* was the causative pathogen for 17% of community-acquired pneumonia cases, a percentage similar to those for community-acquired pneumonia caused by *Streptococcus pneumoniae* (16.6%) and *Haemophilus* species (17.1%). In this study, the percentage for pneumonia caused by *P. aeruginosa* was similar for community-acquired pneumonia and health care–associated pneumonia.⁹⁸

Studies have identified numerous risk factors for *P. aeruginosa* community-acquired pneumonia, including chronic obstructive pulmonary disease, bronchiectasis, alcoholism, smoking, and frequent antibiotic therapy.^{101,102} An immunocompromised state, due to malignancy, transplantation, or immunosuppressive drugs, is also frequently associated with community-acquired pneumonia.^{102,103} Of note, many of these studies were performed before the inclusion of health care–associated pneumonia as a new category of pneumonia, and, therefore, many of these factors likely reflect an increased risk for health care–associated pneumonia and not true community-acquired pneumonia.

The ATS/IDSA guidelines for *P. aeruginosa* community-acquired pneumonia recommend an antipseudomonal β -lactam antimicrobial agent plus either ciprofloxacin, levofloxacin, or an aminoglycoside. An alternative recommended regimen is an aminoglycoside plus either ciprofloxacin or levofloxacin.¹⁰² Tapering to monotherapy once antimicrobial susceptibilities are available should then be considered. Studies defining the optimal duration of antimicrobial therapy are lacking. The ATS/IDSA guidelines state that longer durations of 3 to 7 days are rarely necessary but may be warranted if there are pulmonary cavities or other evidence of tissue necrosis.¹⁰²

Mortality rates associated with *P. aeruginosa* community-acquired pneumonia are higher than those for pneumonia caused by other pathogens. In one study specifically addressing *P. aeruginosa* community-acquired pneumonia, 28% of patients died, compared with 10% who died of community-acquired pneumonia caused by other pathogens.¹⁰¹

Cystic Fibrosis

CF and bronchiectasis are two pulmonary diseases that lead to chronic colonization with *P. aeruginosa* and recurrent pneumonia. CF is a genetic disease that affects multiple systems to varying degrees. Mutations in the CF transmembrane conductance regulator (*CFTR*) gene result in abnormal ion transport, leading to inefficient mucociliary clearance and an increased risk of bacterial infections.^{104–106} *P. aeruginosa* is one of the main bacterial pathogens associated with CF.¹⁰⁷ Initial events of *P. aeruginosa* infection in CF patients can occur early during childhood. Subsequently, the infection dynamics usually follow a model of recurrent episodes of infections that eventually lead to the establishment of a chronic infection in which *P. aeruginosa* can be continuously cultured from respiratory secretions.¹⁰⁶ The transition from an intermittent to a chronic infection state may occur in up to 80% of adult CF patients, and is frequently associated with the emergence of *P. aeruginosa* isolates with a mucoid phenotype, characterized by overproduction of the polysaccharide alginate.^{106,108} Environmental factors, such as oxidative stress and antimicrobial selective pressure, are major drivers of the changes in gene expression that lead to the mucoid-type *P. aeruginosa*. The chronic infection caused by mucoid *P. aeruginosa* isolates is a strong predictor of pulmonary function decline, and represents a significant clinical challenge because it is associated with biofilm formation and multidrug resistance.¹⁰⁹

Bone and Joint Infections

Bone and joint infections can occur via contiguous spread, direct inoculation, or secondary seeding from bacteremia, and occur frequently among intravenous drug users. Although *P. aeruginosa* is an infrequent pathogen implicated in bone and joint infections, certain specific presentations are more often seen with this pathogen. These entities include sternoclavicular septic arthritis, septic arthritis of the symphysis pubis with osteomyelitis, vertebral osteomyelitis, skull base osteomyelitis, and osteomyelitis associated with nail punctures. *P. aeruginosa* is also implicated in a majority of combat-related osteomyelitis cases.

Osteomyelitis caused by *P. aeruginosa* has a poor prognosis when compared with that caused by other pathogens. A study of 454 patients with osteomyelitis characterized risk factors for poor outcomes. *P.*