

Enhanced Meatal Care

Reducing meatal colonization would seem to be a reasonable measure to reduce the risk of CAUTI, given that ascension of uropathogens along the catheter-urethra interface is the predominant source of catheter-associated bacteriuria.⁶⁷ However, periurethral cleansing with chlorhexidine solution versus water before insertion of an indwelling urinary catheter did not reduce the rate of catheter-associated bacteriuria in two different trials.^{170,171} Moreover, results of large, randomized trials have shown no significant reduction in catheter-associated bacteriuria in men or women, compared with usual care (debris removal at daily baths), with one or more times daily meatal cleansing using nonantiseptic green soap or applications of povidone-iodine solution and ointment, polyantibiotic ointment or cream, or silver sulfadiazine 1% cream.^{14,18,20,65} Likewise, simultaneous interventions to block bacterial entry at the urethral insertion site, catheter-drainage tube junction, and outflow tube of the drainage bag were not effective in preventing catheter-associated bacteriuria.¹⁸ Possible reasons why enhanced meatal care has not been effective in reducing catheter-associated bacteriuria include the negative effect of increased catheter manipulation associated with the interventions, inadequate residual antiseptic activity of the topical agent, and lack of effect on the intraluminal route of infection.

Cranberry Products

Although there is conflicting evidence from randomized controlled trials about whether cranberry products are effective in reducing the risk of symptomatic UTIs in women with recurrent UTI, a Cochrane review concluded that cranberry did not significantly reduce the occurrence of symptomatic UTI in any subgroup, including people requiring catheterization.¹⁷² Randomized, placebo-controlled studies of cranberry in doses up to 2 g daily to prevent catheter-associated bacteriuria or CAUTI in patients with neurogenic bladders are mostly negative.^{168,172} The routine use of cranberry products for prevention of health care-associated UTI should be discouraged due to lack of clearly demonstrated efficacy in preventing catheter-associated ASB or CAUTI, problems of tolerance with long-term use, and cost. There are no published data on the use of cranberry products for prevention of ASB or CAUTI in catheterized adults without neurogenic bladder. More recently a double-blind, randomized, placebo-controlled trial of cranberry capsules in 185 noncatheterized women in nursing homes did not find any difference in the primary outcome of bacteriuria plus pyuria in the two arms of the study.¹⁷³

Bladder Irrigation With Antimicrobial Therapy or Saline

Bladder irrigation with antimicrobial agents such as povidone-iodine, chlorhexidine, neomycin, or polymyxin B sulfate has shown little overall benefit in the era of closed urinary drainage.^{14,20} In one demonstrative study, 187 adult patients who required short-term urinary catheterization were randomized to closed drainage with a triple-lumen, neomycin-polymyxin irrigated system, or with a double-lumen nonirrigated catheter system.¹⁴³ There was no significant difference in the rates of catheter-associated bacteriuria between the two groups, but uropathogens isolated from irrigated patients were significantly more resistant to the irrigating antimicrobial than those in the nonirrigated group. Bladder irrigation with saline or acidic solutions also does not have any clear benefit in reducing the risk of catheter blockage resulting from encrustation or in reducing symptomatic UTIs.¹⁷⁴ In summary, bladder irrigation does not appear to be effective in preventing or eradicating catheter-associated bacteriuria in the majority of patients with short-term or long-term indwelling catheterization, is time consuming, may damage the bladder mucosa, and may select for antimicrobial-resistant organisms. Catheter irrigation may have a role in reducing blood clots after urologic surgery, but this topic is outside the scope of this chapter.

Antimicrobial Drugs in the Drainage Bag

Once the drainage bag becomes contaminated, subsequent catheter-associated bacteriuria occurs in almost all patients who remain catheterized,¹⁶ and positioning of the drainage tube above the level of the bladder or below the level of the collection bag is a predictor for an increased risk of catheter-associated bacteriuria.²¹ Investigators have attempted

to prevent catheter-associated bacteriuria by sterilizing the drainage bag, but results of randomized trials evaluating the addition of chlorhexidine, hydrogen peroxide, or povidone-iodine to the bag suggest that this intervention is not effective.^{14,20,65,175} Such a strategy would not be expected to be effective if the integrity of the closed drainage system is carefully maintained because that should minimize bacterial entry into the drainage bag.

Routine Catheter Change

Urinary catheters readily develop biofilms on their inner and outer surfaces once they are inserted, and these biofilms protect uropathogens from antimicrobials and the host immune response.⁵⁶ In long-term catheterized patients, catheters are often changed routinely at periodic intervals (e.g., monthly) to reduce the risk of catheter-associated bacteriuria, as well as for aesthetic reasons. A meta-analysis found only one randomized, controlled trial of this practice; 17 men in a nursing home were randomized to catheter change only when obstructed/infected versus monthly scheduled changes as well as when necessary for obstruction/infection.¹⁷⁶ The trial was too small to provide reliable evidence about CAUTI or blockage. Moreover, it has been recommended that patients who experience repeated early catheter blockage should have their catheters changed every 7 to 10 days to avoid obstruction,¹⁷⁷ but this practice has not been evaluated in clinical trials. The practice of routine catheter change with the purpose of preventing infection or blockage, or both, is unlikely to change in the absence of data to address this issue.

Prophylactic Antimicrobial Therapy at Catheter Removal or Replacement

Fever and bacteremia can occur at the time of removal or replacement of a urethral catheter, and prophylactic antimicrobials are sometimes used to prevent such events. Among catheterized and bacteriuric women in LTCFs, transient fever is twice as common within 24 hours of catheter replacement, compared with other days.²⁹ Studies in chronically catheterized and bacteriuric men and women have shown that bacteremia occurs in 4% to 10% of patients after urethral catheter removal or replacement, but episodes are transient and asymptomatic.^{178–180} The effect of antimicrobial prophylaxis on prevention of bacteremia is unknown. A prospective, randomized, placebo-controlled trial of antimicrobial treatment of catheter-associated ASB persisting 48 hours after removal of short-term catheters in hospitalized women reported significantly improved microbiologic and clinical outcomes at 14 days in treated women.¹⁸¹ Seven (17%) of 42 women randomized to receive no therapy developed symptomatic UTI by 14 days, whereas no women in the treatment group became symptomatic. Whether these results are generalizable to current medical practice is unclear, because all had persistent bacteriuria documented at 48 hours after catheter removal; checking for bacteriuria postcatheterization is not a standard practice or even feasible in discharged patients. Screening for and eradicating postcatheterization catheter-associated ASB to prevent CAUTI is currently not recommended.

A meta-analysis of whether prophylactic antimicrobials at the time of removal of a short-term urinary catheter reduced the risk of symptomatic UTI found seven controlled studies on this topic, of which five were in surgical patients and six were randomized controlled trials.¹⁸² Overall, a benefit was seen with prophylactic antimicrobials, with a 5.8% absolute reduction in symptomatic UTI and a risk ratio of 0.45 (95% CI, 0.28–0.72). Nevertheless, prophylactic antimicrobial agents are not routinely recommended for catheter removal or replacement because it is unclear whether these findings apply to the more general medical inpatient population, and widespread implementation of prophylactic antimicrobials at the time of catheter removal has potential to cause harm in terms of cost, side effects, and resistance.

Preclinical Strategies for CAUTI Prevention

Many novel approaches have been studied *in vitro*, with the goal of prevention or eradication of the catheter-associated biofilm of uropathogens.¹⁸³ Of these, mannositides—small molecule inhibitors of the type 1 pilus adhesin, FimH—seem likely to move to clinical trials in

humans and could potentially impact prevention of UTI more generally. In a study in mice, mannoses provided significant protection against uropathogenic *E. coli* catheter-associated bacteriuria by preventing bacterial invasion and shifting the *E. coli* niche primarily to the extracellular milieu.¹⁸⁴ Mannosides can be targeted to the fimbrial adhesins on uropathogenic *E. coli* while having less affinity for the adhesins of commensal gut *E. coli*; thus mannoses were able to deplete the mouse gut of uropathogenic *E. coli* without perturbing the commensal *E. coli*.¹⁸⁵ A selective agent that could reduce colonization by urinary pathogens without disrupting the gut microbiota would have obvious advantages over prophylactic antimicrobials for UTI prevention.

HEALTH CARE–ASSOCIATED ASYMPTOMATIC BACTERIURIA

Routine Screening and Treatment

Screening and treatment of ASB have not been shown to be beneficial, will select for antimicrobial resistance, and are not recommended, except in pregnant women and patients who undergo traumatic genitourinary procedures associated with mucosal bleeding.^{5,186} Populations that have been extensively studied and for whom these recommendations apply include premenopausal, nonpregnant women; diabetic women; older people living in the community; elderly, institutionalized subjects; persons with SCI; and catheterized patients.⁵ For example, in 35 patients undergoing long-term catheterization, a prospective randomized trial of cephalexin or no antibiotic therapy for episodes of catheter-associated ASB caused by susceptible organisms reported no differences between the two groups in incidence and prevalence of catheter-associated bacteriuria, CAUTI, or catheter obstruction in patients followed up to 44 weeks.¹⁸⁷ In addition, 47% of reinfecting organisms in the cephalexin group compared with 26% in the control group were highly resistant to cephalexin. Even if treatment of catheter-associated ASB was found to be useful, one study in which daily catheter urine cultures were obtained found that 60% of 25 episodes of CAUTI occurred on the same day that catheter-associated bacteriuria was first detected,²⁶ complicating attempts at preemptive therapy. It is not known whether eradication of catheter-associated ASB might be beneficial in reducing cross-infection or inappropriate antimicrobial usage.

Prevention of Overtreatment of Asymptomatic Bacteriuria in Health Care Settings

The greatest impact that ASB has on clinical care is probably the excessive testing and treatment done for this condition in health care settings.¹⁸⁸ Prevention of inappropriate treatment of ASB is probably the second most beneficial strategy for patients related to health care–associated UTI, secondary only to avoidance of unnecessary urinary catheterization. Estimates vary a bit by site of care delivery, but studies in acute-care hospitals, LTCFs, SCI clinics, and emergency departments have documented that 20% to 83% of episodes of detected ASB are treated with antimicrobials.^{189–191} Thus prevention of inappropriate treatment of ASB is essential to antimicrobial stewardship in acute and long-term care settings, as endorsed by the American Board of Internal Medicine's Choosing Wisely campaign.^{192,193} A positive urine culture is a powerful stimulus for subsequent use of antimicrobials, as demonstrated by surveys of practitioners and also by an innovative study in which urine culture results from the microbiology laboratory were suppressed.^{194,195} A stewardship intervention that focused on the "culture of culturing,"¹⁹⁶ discouraging unnecessary cultures in patients without urinary-specific symptoms, successfully reduced both urine cultures and antimicrobial treatment of ASB in acute and long-term care wards of a large medical facility.¹⁹⁷ Unnecessary urine cultures in patients with ASB can lead to false elevation of the reported CAUTI rates, because a positive urine culture in a febrile patient must be reported to the NSHN as a CAUTI, regardless of whether the fever can be attributed to another cause.¹⁹⁸ The co-occurrence of fever and bacteriuria is particularly common in ICU and long-term care patients. A tertiary care academic medical center reduced CAUTI rates significantly in its ICUs through a multifaceted program that centered on the stewardship of urine culturing.¹⁹⁹ Likewise, emphasis on judicious use of urine cultures in a nationwide project in nursing homes was associated with a decrease in both urine cultures

and CAUTI.⁸⁵ Removing unnecessary urinary catheters through CAUTI prevention efforts may have the side benefit of also reducing testing and treatment for ASB, because cloudy urine or visible sediment in the urine is a stimulus for urine cultures among nursing personnel.^{194,200} A recent national study of over 4 million hospital admissions documented that 47% had a urinalysis and 27% had a urine culture during hospitalization²⁰¹; thus there is considerable room for improvement in the stewardship of urine testing as well as stewardship of antimicrobial use for ASB.

MANAGEMENT OF HEALTH CARE–ASSOCIATED URINARY TRACT INFECTION

The issues relevant to clinical management of health care–associated UTI, particularly CAUTI, include whether the catheter should be changed prior to urine collection for culture, whether the catheter should be changed during the course of treatment, choice of drugs, duration of therapy, and when to consider complicating factors that may impair response to antimicrobials. The wide variety of underlying conditions, diverse spectrum of possible etiologic agents, and paucity of controlled clinical trials with stratification according to specific complicating factors make generalizing about clinical management of CAUTI difficult. Antimicrobials alone may not be successful, if underlying anatomic, functional, or metabolic defects are not corrected.

Urine Culture and Catheter Replacement Before Treatment

CAUTIs, especially in patients with long-term catheterization, are often polymicrobial and caused by multidrug-resistant uropathogens, so urine cultures should be obtained prior to treatment to confirm that the empirical regimen provides appropriate coverage and to allow tailoring of the regimen based on antimicrobial susceptibility data.^{40,202} The culture should be obtained from a freshly placed catheter if the catheter has been in place for a few days because the catheter biofilm may result in spurious culture results.^{53,54} Moreover, clinical outcomes are improved if the catheter is replaced, as shown in a prospective, randomized, controlled trial in elderly nursing home residents with long-term indwelling catheters and CAUTI. This study demonstrated that patients whose catheters had been in place for longer than 2 weeks and who underwent catheter replacement before antimicrobial treatment had significantly shorter time to improved clinical status and significantly lower rates of polymicrobial catheter-associated bacteriuria and CAUTI after therapy, compared with those who did not undergo catheter replacement.²⁰³ These study findings support replacing the catheter and obtaining a urine culture from a freshly placed catheter before antimicrobial treatment for CAUTI if the catheter has been in place for at least 2 weeks and cannot be permanently removed.

Choice of Antimicrobial Agent

The choice of antimicrobial for empirical treatment should take into account the facility antibiogram, the patient's specific risk factors for having a multidrug-resistant urinary pathogen, the severity of illness, and the patient's underlying comorbidities.²⁰⁴ Risk factors for colonization with a multidrug-resistant organism, particularly an extended-spectrum β -lactamase-producing Enterobacteriaceae, include international travel to an area endemic for resistant Enterobacteriaceae, such as southern Asia, within the prior 12 months, and receipt of antibiotics to treat traveler's diarrhea during a prior international trip.^{205,206} Microbiologic data from urine cultures within the prior 2 years can be used to guide empirical antibiotic choice and improve the likelihood of choosing an antimicrobial agent effective against the current urinary pathogen.²⁰⁷ Health care exposure (such as hospitalization or stay in an LTCF) and receipt of antibiotics within the prior 6 months are also risk factors for colonization with resistant organisms.²⁰⁸ In nursing home residents, indwelling devices such as urinary catheters or feeding tubes are risk factors for colonization with, and thus infection by, multidrug-resistant organisms.²⁰⁹ Whether the patient is able to take oral agents is also a consideration in antimicrobial choice.

A general perspective on resistance trends in CAUTI-causing organisms can also be useful, particularly if a facility-specific antibiogram is not available. Data from the NSHN between 2011 and 2014 showed a

high prevalence of multidrug-resistant phenotypes among strains causing CAUTI.¹⁹ Thus in 2014, 34.8% of *E. coli* strains tested were resistant to fluoroquinolones and 15.5% to extended-spectrum cephalosporins (e.g., cefepime); 9.5% of *Klebsiella* spp. were resistant to carbapenems and 22.5% to extended-spectrum cephalosporins. Global studies from 2003 to 2010 of urology patients with health care–associated bacteriuria, which included ASB, found that slightly over 50% of urinary isolates were resistant to fluoroquinolones (excluding *Candida* from the possible pathogens).²¹⁰

Appropriate empirical treatment choices require that patients be categorized on a spectrum from mild illness and ability to take oral agents to severe illness and inability to take oral agents. In all patients a urine culture should be collected before starting antimicrobials so that the antimicrobial can be changed to a more appropriate agent if necessary. Outpatients who are not severely ill can be empirically treated with oral fluoroquinolones, cephalosporins, trimethoprim-sulfamethoxazole, or amoxicillin-clavulanate (Table 302.7). Consideration should be given to providing an initial dose(s) of a broader-spectrum parenteral agent, such as ceftriaxone, a carbapenem, or an aminoglycoside, if there is concern about antimicrobial resistance, while waiting for urine culture results. Patients at low risk for multidrug resistance and who are not critically ill but who require inpatient therapy can be treated with ceftriaxone or piperacillin-tazobactam. If a patient who is not critically ill has risk factors for an extended-spectrum β -lactamase–producing organism and no prior cultures with carbapenem-resistant Enterobacteriaceae, a carbapenem should be considered for empirical therapy. Patients who are critically ill should be empirically treated with an antipseudomonal carbapenem (meropenem, imipenem, or doripenem) and vancomycin or daptomycin. Patients with prior carbapenem-resistant Enterobacteriaceae cultures who are critically ill or unable to take an oral agent, may warrant, in consultation with an infectious diseases expert, empirical use of some of the newer, expensive, and very broad-spectrum agents such as ceftolozane-tazobactam, ceftazidime-avibactam, meropenem-vaborbactam, or plazomicin. The antimicrobial regimen should be tailored as appropriate when the infecting strain has been identified and antimicrobial susceptibilities are known.

Duration

Few studies have been performed that evaluate duration of treatment in populations with CAUTI or other complicated UTIs. Reviews of complicated UTI have recommended treatment durations from 7 to 10 days,¹⁴ 7 to 14 days,⁴⁰ and 7 to 21 days,²⁰² depending on the severity of the infection. However, it is desirable to limit the duration of treatment, especially for milder infections, to reduce the selection pressure for drug-resistant flora, especially in patients on long-term catheterization. In one small trial of women with lower urinary tract CAUTI, cure rates were comparable with a single-dose and a 10-day regimen of trimethoprim-sulfamethoxazole.¹⁸¹ However, single-dose therapy was ineffective in eradicating bacteriuria in institutionalized elderly men.²¹¹ In a randomized, double-blind, placebo-controlled trial comparing 3-day and 14-day regimens of ciprofloxacin for the treatment of mild CAUTI in 60 patients with SCI, there was no difference in clinical outcomes at long-term follow-up.²¹² Most recently, clinical and microbiologic success rates following treatment were almost identical in a noninferiority study of 619 patients with acute pyelonephritis or complicated UTI treated with a 5-day course of levofloxacin or a 10-day course of ciprofloxacin.²¹³ These data suggest that a 7-day regimen is reasonable for most patients with CAUTI, depending on their clinical response, and that shorter regimens, such as a 5-day regimen of a urinary fluoroquinolone, are likely to be sufficient in those patients who are less severely ill, are infected with uropathogens susceptible to the antimicrobial used, and have a rapid response to treatment.

Nephrostomy Tubes and Ureteral Stents

Patients with obstruction of the ureter or ureteropelvic junction, often due to stones, malignancy, or benign strictures, may have a ureteral catheter (stent) or a percutaneous nephrostomy tube inserted. A ureteral catheter may be inserted from the bladder during cystoscopy or through a percutaneous nephrostomy into the renal pelvis and further down into the ureter, with a drainage hole at the level of the renal pelvis.

TABLE 302.7 Empirical Management of Catheter-Associated Urinary Tract Infection

Antimicrobial Agent and Dosing

Mild to Moderate Illness, Can Tolerate Oral Therapy (Dosage Duration: 5–10 Days)^a

Ciprofloxacin 500 mg PO twice daily or 1 g (extended release) PO once daily^c
 Levofloxacin 750 mg PO once daily^c
 Cefpodoxime 200 mg PO q12h^d (or other third-generation cephalosporin)
 Trimethoprim-sulfamethoxazole (TMP-SMX) 160/800 mg PO q8h^e
 Amoxicillin-clavulanate immediate release 875 mg orally twice daily^d

Severe Illness or Unable to Tolerate Oral Therapy (Dosage Duration: 5–14 Days)^b

Ciprofloxacin 400 mg IV twice daily^c
 Levofloxacin 750 mg IV once daily^c
 Ceftriaxone 1–2 g IV once daily^d
 Cefepime 1 g IV twice daily^d
 Piperacillin-tazobactam 3.375 g IV (as piperacillin) q6h^d
 Meropenem 1 g IV q8h^d
 Imipenem-cilastatin 500 mg IV q6h^c
 Doripenem 500 mg IV q8h^d
 Ertapenem 1 g IV once daily^d
 Gentamicin 5–7 mg/kg IV once daily^e ± ampicillin 1–2 g IV q6h^d
 Ceftolozane-tazobactam 1.5 g IV (as both drugs) q8h^d
 Ceftazidime-avibactam 2.5 g IV (as both drugs) q8h^d
 Meropenem-vaborbactam 4 g IV (as both drugs) q8h (pregnancy category unknown)
 Plazomicin 15 mg/kg IV once daily^e

In Choosing an Empirical Agent, Consider the Following:

Collect a urine culture before beginning treatment to guide subsequent therapy
 Severity of illness and comorbidities—choose a broader agent for more severely ill patients or those who are immunocompromised
 Antimicrobial susceptibility of bacteriuria strains within past 2 years
 Antibiotic exposure within prior 6 months
 Local resistance data—in some locations, fluoroquinolones and/or TMP-SMX are no longer reasonable empirical choices given high resistance levels among *E. coli*
 Hospitalization or stay in a long-term care facility within past 6 months
 Travel to an area with high prevalence of drug resistance in prior 12 months
 Consider adding vancomycin if gram-positive organism suspected
 Use a carbapenem (meropenem, imipenem, doripenem, or ertapenem) if an extended-spectrum β -lactamase strain is known or suspected
 Tailor regimen based on susceptibility data, and transition to oral medications as soon as condition allows

^aConsider a concomitant single initial dose of ceftriaxone 1 mg IV/IM, ertapenem 1 g IV/IM, meropenem 1 g IV, gentamicin 5 mg/kg IV/IM, amikacin 15 mg/kg IV/IM, or tobramycin 5 mg/kg IV/IM, particularly in febrile patients or patients at risk for being unable to follow up. Duration of therapy can be 5 days in patients receiving fluoroquinolones. Other agents should be used for 7–10 days.

^bConsider an initial intravenous dose of a broader-spectrum agent, such as meropenem, particularly if using fluoroquinolones in a severely ill patient. Duration of therapy can be shortened to 5 days in patients receiving fluoroquinolones who have a rapid response. Other agents should be used for 7–14 days, with longer courses for patients with a slow response to therapy or with complications, such as an abscess. Doses in the table are for adults with normal renal function.

^cPregnancy category C—animal studies have shown an adverse effect on the fetus; use only if potential benefit justifies the potential risk to the fetus.

^dPregnancy category B—no clear risk to fetus based on animal or human studies, or both.

^ePregnancy category D—associated with human fetal risk; use only if the potential benefit justifies the potential risk to the fetus.

Modified from Hooton et al.² and Johnson and Russo.²⁰⁴

J-shaped curls at catheter ends are used to keep the catheter in place. Nephrolithiasis at the time of tube placement can trigger either systemic inflammatory response syndrome or sepsis; the risk of both is low but higher in patients who have a preexisting UTI or infected stone.²¹⁴ Intraoperative cultures of urine from the renal pelvis and stone material can be useful to guide postprocedure antibiotics.²¹⁵ Once a percutaneous nephrostomy tube is in place, colonization of the tube and urine with bacteria and/or *Candida* occurs commonly, often with more than one organism. Usually this colonization is asymptomatic, but it can lead to fever, pyelonephritis, renal abscess, or bacteremia if drainage from the renal pelvis becomes obstructed or, less often, if instrumentation of an infected urinary tract introduces organisms into the blood stream.²¹⁶ A clinical definition of nephrostomy-associated pyelonephritis, to help distinguish an actual infection from the more

common asymptomatic colonization, is the presence of fever, costovertebral angle tenderness, or flank pain associated with a positive urine culture, or a combination of these.²¹⁷ An obstructed, infected renal pelvis can rapidly evolve into a pyonephrosis (infection of the kidney with pus in the upper collecting system), which can be complicated by renal papillary necrosis or perinephric abscess. Infected urine in the pelvis can track along the nephrostomy tube into the perinephric space and even extend down the iliacus muscle. Treatment involves immediate drainage and appropriate antimicrobial drugs. Consideration should be given to exchange of the nephrostomy tube during treatment,²¹⁸ analogous to the recommendation to change an indwelling urethral catheter during treatment of a CAUTI. Penetration of the drug into the urine is not imperative in treating pyonephrosis or perinephric abscess but is relevant to treating bacteriuria or candiduria prior to a planned urologic procedure.²¹⁹ Fluconazole and flucytosine obtain high urine concentrations, while intravenous conventional amphotericin B achieves low concentrations. Other antifungals, such as the echinocandins, may have insufficient urine concentrations to affect candiduria but are active within the renal parenchyma. Amphotericin B irrigation of the renal pelvis as 50 mg/L in sterile water has been used for prophylaxis.²¹⁹ Toxicity is minimal but efficacy is unclear. When used as prophylaxis, an antifungal or antibacterial should be begun prior to the procedure and discontinued shortly after an uncomplicated procedure (duration of 24 hours or less).²²²

FUNGAL URINARY TRACT INFECTION

Yeasts, mostly *Candida* species, are commonly isolated from the urine of catheterized patients.²²³ NHSN data from 2011 to 2014 reported that *C. albicans* was the second most common organism identified in CAUTI, second only to *E. coli*; other *Candida* species (including all except *C. albicans*) ranked 10th, and *Candida glabrata* alone ranked 14th.¹⁹ In January 2015 the NHSN surveillance definitions for CAUTI changed to exclude *Candida* species and other fungal organisms as causative agents of CAUTI, so nationwide estimates on the prevalence of fungal CAUTI will be difficult to obtain after this date.

Among patients with nosocomial funguria enrolled in a multicenter prospective surveillance study, 85% had concomitant nonfungal infections (urinary and nonurinary); 90%, previous exposure to antimicrobial agents; 83%, urinary tract drainage devices; 39%, diabetes; 38%, urinary tract abnormalities; and 22%, a malignancy—only 11% had no obvious underlying illness.²²⁴ Although only 1.3% of the 530 patients with candiduria followed for 12 weeks had documented candidemia, the importance of comorbid conditions was reflected in the 20% mortality rate. Candiduria in itself may be a marker for severity of underlying illness, because multiple studies have shown that patients with candiduria have higher mortality rates, although not typically from *Candida* infection.²²⁵

Determination of the clinical significance of candiduria can be problematic because it can represent contamination of a voided specimen, colonization of catheters or stents, bladder infection, ascending kidney infection, or kidney infection associated with candidemia. Hematogenous dissemination is relatively much more likely to be the source of candiduria than is the case with bacteriuria.²²⁶ Neither colony count thresholds nor the presence of pseudohyphae in the urine are helpful to distinguish contamination from bladder infection. In noncatheterized women with candiduria, a catheter specimen may be indicated to rule out contamination with perineal flora.²²⁶ In the catheterized patient, pyuria is a nonspecific finding, but its absence suggests that candiduria is not causing tissue invasion.

Most health care–associated candiduria occurs in catheterized patients, and most episodes are asymptomatic. In the large prospective study of nosocomial funguria mentioned earlier, only 2% to 4% of patients had urinary symptoms,²²⁴ although comorbidities and urinary catheterization may complicate assessment of symptoms. However, catheterized patients with candiduria may have symptoms or signs such as suprapubic or flank pain, lower abdominal discomfort associated with prostatitis, or an enlarged and tender scrotum associated with epididymo-orchitis.

Ascending infection of the kidney and disseminated candidiasis are rarely associated with candiduria and usually occur in the setting of

obstruction of the urinary tract. In a retrospective study of 26 cases of candidemia associated with a well-defined urinary tract source, urinary tract abnormalities (mostly obstruction) were present in 88%, and 73% had undergone urinary tract procedures before the onset of candidemia.²²⁷ Episodes of candidemia were brief and low grade in intensity, although two of five in-hospital deaths were attributable to candidiasis. Paired urine and blood strains of *Candida*, however, may not be the same strain, as demonstrated in a case-control study in which 52% of paired strains were different by molecular typing.²²⁸ Other complications associated with fungal infections of the genitourinary tract include fever, fungus balls in the bladder or renal pelvis, renal and perirenal abscesses, emphysematous pyelitis or pyelonephritis, and papillary necrosis—many of these complications are more likely to occur in diabetics.

Few treatment studies have been performed in patients with candiduria,²²⁹ and there remain questions as to whom to treat, when to treat, and how long to treat. Asymptomatic candiduria rarely requires treatment because it often resolves spontaneously, morbidity is low, and treatment is often followed by rapid recurrence and may select out for resistant organisms.^{225,230} In the large prospective observational study mentioned previously, funguria cleared in 76% of 155 patients who had no specific therapy for funguria and in 35% of 116 patients who had their catheter removed as the only treatment.²²⁴ Other studies have shown spontaneous clearance rates of candiduria ranging from 29% to 62%.²²⁴ In a randomized, placebo-controlled trial of 316 hospitalized patients with asymptomatic or minimally symptomatic candiduria, a 2-week course of fluconazole resulted in significantly higher eradication rates than placebo (50% vs. 29%), but there was no significant difference in candiduria rates 2 weeks after completion of treatment.²³⁰ In placebo recipients who had their catheters removed, 41% had eradication of candiduria, compared with only 20% who had their catheters replaced. Pyelonephritis, candidemia, and fungus-related death were not observed in this study. The possibility of disseminated candidiasis should be considered in all hospitalized patients with candiduria, especially in the ICU, where candidemia is common and up to 80% of persons with candidemia will have accompanying candiduria.²³¹ However, candidemia occurs in less than 5% of ICU patients and thus most such patients with candiduria do not have candidemia.²³¹ If candiduria fails to resolve despite changing the catheter or stopping antibiotics (if feasible), a more deep-seated infection should be suspected, especially in critically ill patients, and imaging of the kidneys and collecting system is indicated to look for a renal abscess, fungus ball, or other urologic abnormality that may require treatment.²³¹ Antifungal treatment of candiduria should be reserved for those patients who have solid evidence of infection of the kidney or collecting system or disseminated candidiasis.²³¹ Treatment is also indicated in asymptomatic patients with neutropenia, infants with low birth weight, and patients who will undergo urologic manipulations because these conditions have a higher association with upper tract infection or dissemination.^{225,226} Recent data have raised questions as to whether asymptomatic candiduria in patients with renal transplants warrants treatment.²³² Of note, in 2000, a multicenter study in the United States reported that 43% of instances of candiduria were treated, likely reflecting overtreatment of funguria.²²⁴

Candiduria should be treated in symptomatic patients, and those with systemic signs or symptoms should be evaluated for disseminated infection with imaging and blood cultures. Oral fluconazole is the drug of choice for cystitis and pyelonephritis due to most species of *Candida* (but not resistant *C. glabrata*, *Candida krusei*, or other resistant yeasts); of note, doses adjusted for renal insufficiency may result in subtherapeutic concentrations.²²⁹ Amphotericin B and flucytosine can be used for *C. glabrata* resistant to fluconazole, and amphotericin B is the recommended agent for *C. krusei*.²²⁵ Other azoles that could serve as alternative choices for resistant isolates, including itraconazole, voriconazole, and posaconazole, are not recommended for *Candida* UTI because of minimal excretion into the urine, even though they achieve antifungal activity in renal parenchyma.²³¹ Systemic amphotericin B deoxycholate is no more effective than fluconazole for susceptible strains, and bladder irrigation with amphotericin B deoxycholate is rarely used. Lipid formulations of amphotericin B may not be the first choice for *Candida* UTI because of presumed low concentrations of the drug in renal tissue. Failure of the lipid complex formulation of amphotericin B therapy has

been described in the treatment of *Candida* pyelonephritis in experimental animals and patients.²²⁵ Echinocandins are minimally excreted and should not be considered first-line agents for candiduria. However, case reports suggest that caspofungin and micafungin may be effective in patients with *C. glabrata*-associated UTI.^{233,234} Even with apparently successful local or systemic antifungal therapy for candiduria, relapse is frequent, and this likelihood is increased by continued use of a urinary catheter. Persistent candiduria, especially in immunocompromised patients, warrants radiologic imaging of the kidneys to evaluate for hydronephrosis, fungal bezoars (common in critically ill neonates²²⁴), or perinephric abscesses associated with ascending infection.

SUMMARY

Health care-associated bacteriuria and candiduria are common, mostly associated with urinary catheterization, and usually asymptomatic. Whereas any symptomatic UTI should be treated, published data do not support routine screening to detect health care-associated bacteriuria or candiduria in asymptomatic patients because treatment does not appear to alter the natural course of infection and an increase in antimicrobial resistance often results. However, among asymptomatic patients with bacteriuria or funguria, we need a better understanding as to who has upper tract involvement or tissue invasion, how these affect the clinical course, and thus which patients, if any, might benefit

from screening and treatment. The quality of clinical trials needs to be improved because many are nonrandomized and underpowered and use nonspecific terminology and poorly described methodology. Studies of interventions to prevent catheter-associated bacteriuria need to address whether they prevent not only CAUTI (which many have not), but also whether prevention of catheter-associated ASB reduces inappropriate antimicrobial use, bacteremia, and cross-infection.

Strategies that are effective for prevention of catheter-associated bacteriuria are likely to also prevent catheter-associated candiduria. The most effective way to reduce health care-associated UTIs is to reduce urinary catheterization by restricting use to patients who have clear indications and by removing the catheter as soon as it is no longer needed. Implementation of strategies to reduce catheterization is likely to have more impact on catheter-associated bacteriuria than implementation of other strategies addressed in this chapter. However, use of multiple infection control techniques and strategies simultaneously (called *bundling*) likely offers the best opportunity to reduce the morbidity and mortality related to health care-associated infections.²³⁵ In those patients who do require urinary catheterization, an effective stewardship program for both urine testing and antimicrobial treatment in asymptomatic patients offers potential benefits to patients and the health care system. More research is needed in terms of how to best implement effective stewardship programs for bacteriuria.

Key References

The complete reference list is available online at Expert Consult.

- Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36:309–332.
- Hooton TM, Bradley SF, Cardenas DD, et al. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis*. 2010;50:625–663.
- Nicolle LE, Gupta K, Bradley SF, et al. Clinical Practice Guideline for the Management of Asymptomatic Bacteriuria: 2019 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2019 Mar 21. [Epub ahead of print].
- Warren JW. Catheter-associated urinary tract infections. *Infect Dis Clin North Am*. 1997;11:609–622.
- Kunin CM, McCormack RC. Prevention of catheter-induced urinary-tract infections by sterile closed drainage. *N Engl J Med*. 1966;274:1155–1161.
- Classen DC, Larsen RA, Burke JP, et al. Prevention of catheter-associated bacteriuria: clinical trial of methods to block three known pathways of infection. *Am J Infect Control*. 1991;19:136–142.
- Saint S, Lipsky BA. Preventing catheter-related bacteriuria: should we? Can we? How? *Arch Intern Med*. 1999;159:800–808.
- Maki DG, Tambyah PA. Engineering out the risk for infection with urinary catheters. *Emerg Infect Dis*. 2001;7:342–347.
- Tambyah PA, Maki DG. Catheter-associated urinary tract infection is rarely symptomatic: a prospective study of 1,497 catheterized patients. *Arch Intern Med*. 2000;160:678–682.
- Nicolle LE. Catheter-related urinary tract infection. *Drugs Aging*. 2005;22:627–639.
- Tenney JH, Warren JW. Bacteriuria in women with long-term catheters: paired comparison of indwelling and replacement catheters. *J Infect Dis*. 1988;157:199–202.
- Stamm WE. Catheter-associated urinary tract infections: epidemiology, pathogenesis, and prevention. *Am J Med*. 1991;91:655–715.
- Tambyah PA, Halvorson KT, Maki DG. A prospective study of pathogenesis of catheter-associated urinary tract infections. *Mayo Clin Proc*. 1999;74:131–136.
- Stark RP, Maki DG. Bacteriuria in the catheterized patient. What quantitative level of bacteriuria is relevant? *N Engl J Med*. 1984;311:560–564.
- Tambyah PA, Maki DG. The relationship between pyuria and infection in patients with indwelling urinary catheters: a prospective study of 761 patients. *Arch Intern Med*. 2000;160:673–677.
- Saint S, Meddings JA, Calfee D, et al. Catheter-associated urinary tract infection and the Medicare rule changes. *Ann Intern Med*. 2009;150:877–884.
- Smith PW, Bennett G, Bradley S, et al. SHEA/APIC guideline: infection prevention and control in the long-term care facility. *Infect Control Hosp Epidemiol*. 2008;29:785–814.
- Meddings J, Rogers MA, Macy M, et al. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis*. 2010;51:550–560.
- Saint S, Wiese J, Amory JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med*. 2000;109:476–480.
- Cornia PB, Amory JK, Fraser S, et al. Computer-based order entry decreases duration of indwelling urinary catheterization in hospitalized patients. *Am J Med*. 2003;114:404–407.
- Griffiths R, Fernandez R. Strategies for the removal of short-term indwelling urethral catheters in adults. *Cochrane Database Syst Rev*. 2007;(2):CD004011.
- Saint S, Kaufman SR, Rogers MA, et al. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc*. 2006;54:1055–1061.
- Niël-Weise BS, van den Broek PJ, da Silva EM, et al. Urinary catheter policies for long-term bladder drainage. *Cochrane Database Syst Rev*. 2012;(8):CD004201.
- Jamison J, Maguire S, McCann J. Catheter policies for management of long term voiding problems in adults with neurogenic bladder disorders. *Cochrane Database Syst Rev*. 2011;(12):CD004375.
- Moore KN, Fader M, Getliffe K. Long-term bladder management by intermittent catheterisation in adults and children. *Cochrane Database Syst Rev*. 2007;(4):CD006008.
- Lo E, Nicolle L, Coffin SE, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol*. 2014;35:464–479.
- Warren JW, Platt R, Thomas RJ, et al. Antibiotic irrigation and catheter-associated urinary-tract infections. *N Engl J Med*. 1978;299:570–573.
- Schumm K, Lam TB. Types of urethral catheters for management of short-term voiding problems in hospitalised adults. *Cochrane Database Syst Rev*. 2008;(2):CD004013.
- Drekonja DM, Kuskowski MA, Wilt TJ, et al. Antimicrobial urinary catheters: a systematic review. *Expert Rev Med Devices*. 2008;5:495–506.
- Jahn P, Beutner K, Langer G. Types of indwelling urinary catheters for long-term bladder drainage in adults. *Cochrane Database Syst Rev*. 2012;(10):CD004997.
- Lee BS, Bhuta T, Simpson JM, et al. Methenamine hippurate for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD003265.
- Jepson RG, Williams G, Craig JC. Cranberries for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD001321.
- Darouiche RO, Hull RA. Bacterial interference for prevention of urinary tract infection. *Clin Infect Dis*. 2012;55:1400–1407.
- U.S. Preventive Services Task Force. Screening for asymptomatic bacteriuria in adults: U.S. Preventive Services Task Force reaffirmation recommendation statement. *Ann Intern Med*. 2008;149:43–47.
- Stamm WE, Hooton TM. Management of urinary tract infections in adults. *N Engl J Med*. 1993;329:1328–1334.
- Raz R, Schiller D, Nicolle LE. Chronic indwelling catheter replacement before antimicrobial therapy for symptomatic urinary tract infection. *J Urol*. 2000;164:1254–1258.
- Kauffman CA, Vazquez JA, Sobel JD, et al. Prospective multi-center surveillance study of funguria in hospitalized patients. The National Institute for Allergy and Infectious Diseases (NIAID) Mycoses Study Group. *Clin Infect Dis*. 2000;30:14–18.
- Pappas PG, Kauffman CA, Andes DR, et al. Clinical Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;62:e1–e50.
- Sobel JD, Kauffman CA, McKinsey D, et al. Candiduria: a randomized, double-blind study of treatment with fluconazole and placebo. The National Institute of Allergy and Infectious Diseases (NIAID) Mycoses Study Group. *Clin Infect Dis*. 2000;30:19–24.
- Fisher JF, Sobel JD, Kauffman CA, et al. *Candida* urinary tract infections—treatment. *Clin Infect Dis*. 2011;52(suppl 6):S457–S466.

References

- Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36:309–332.
- Hooton TM, Bradley SF, Cardenas DD, et al. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis*. 2010;50:625–663.
- Platt R, Polk BF, Murdock B, et al. Mortality associated with nosocomial urinary-tract infection. *N Engl J Med*. 1982;307:637–642.
- Platt R, Polk BF, Murdock B, et al. Reduction of mortality associated with nosocomial urinary tract infection. *Lancet*. 1983;1:893–897.
- Nicolle LE, Gupta K, Bradley SF, et al. Clinical Practice Guideline for the Management of Asymptomatic Bacteriuria: 2019 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2019 Mar 21. [Epub ahead of print].
- Warren J, Anthony W, Hoopes J, et al. Cephalixin for susceptible bacteriuria in afebrile, long-term catheterized patients. *JAMA*. 1982;248:454–458.
- Breitenbucher RB. Bacterial changes in the urine samples of patients with long-term indwelling catheters. *Arch Intern Med*. 1984;144:1585–1588.
- Stevens V, Dumyati G, Fine LS, et al. Cumulative antibiotic exposures over time and the risk of *Clostridium difficile* infection. *Clin Infect Dis*. 2011;53:42–48.
- Tambyah PA, Maki DG. Catheter-associated urinary tract infection is rarely symptomatic: a prospective study of 1,497 catheterized patients. *Arch Intern Med*. 2000;160:678–682.
- Weber DJ, Sikkert-Bennett EE, et al. Incidence of catheter-associated and non-catheter-associated urinary tract infections in a healthcare system. *Infect Control Hosp Epidemiol*. 2011;32:822–823.
- Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases. 2014 National and State Healthcare-Associated Infections Progress Report. Atlanta: Centers for Disease Control and Prevention; 2016. Available at: www.cdc.gov/hai/progress-report/index.html.
- Fakih MG, Gould CV, Trautner BW, et al. Beyond infection: device utilization ratio as a performance measure for urinary catheter harm. *Infect Control Hosp Epidemiol*. 2016;37:327–333.
- Dudeck MA, Edwards JR, Allen-Bridson K, et al. National Healthcare Safety Network report, data summary for 2013, Device-associated Module. *Am J Infect Control*. 2015;43:206–221.
- Warren JW. Catheter-associated urinary tract infections. *Infect Dis Clin North Am*. 1997;11:609–622.
- Gould CV, Umscheid CA, Agarwal RK, et al. Healthcare Infection Control Practices Advisory Committee. *Guideline for Prevention of Catheter-Associated Urinary Tract Infections 2009*. Atlanta: Centers for Disease Control and Prevention; 2017. <https://www.cdc.gov/infectioncontrol/pdf/guidelines/cauti-guidelines.pdf>. update.
- Garibaldi RA, Burke JP, Dickman ML, et al. Factors predisposing to bacteriuria during indwelling urethral catheterization. *N Engl J Med*. 1974;291:215–219.
- Kunin CM, McCormack RC. Prevention of catheter-induced urinary-tract infections by sterile closed drainage. *N Engl J Med*. 1966;274:1155–1161.
- Classen DC, Larsen RA, Burke JP, et al. Prevention of catheter-associated bacteriuria: clinical trial of methods to block three known pathways of infection. *Am J Infect Control*. 1991;19:136–142.
- Weiner LM, Webb AK, Limbago B, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. *Infect Control Hosp Epidemiol*. 2016;37:1288–1301.
- Saint S, Lipsky BA. Preventing catheter-related bacteriuria: should we? Can we? How? *Arch Intern Med*. 1999;159:800–808.
- Maki DG, Tambyah PA. Engineering out the risk for infection with urinary catheters. *Emerg Infect Dis*. 2001;7:342–347.
- Saint S, Chenoweth CE. Biofilms and catheter-associated urinary tract infections. *Infect Dis Clin North Am*. 2003;17:411–432.
- Platt R, Polk BF, Murdock B, et al. Risk factors for nosocomial urinary tract infection. *Am J Epidemiol*. 1986;124:977–985.
- Finer G, Landau D. Pathogenesis of urinary tract infections with normal female anatomy. *Lancet Infect Dis*. 2004;4:631–635.
- Saint S. Clinical and economic consequences of nosocomial catheter-related bacteriuria. *Am J Infect Control*. 2000;28:68–75.
- Garibaldi RA, Mooney BR, Epstein BJ, et al. An evaluation of daily bacteriologic monitoring to identify preventable episodes of catheter-associated urinary tract infection. *Infect Control*. 1982;3:466–470.
- Hartstein AI, Garber SB, Ward TT, et al. Nosocomial urinary tract infection: a prospective evaluation of 108 catheterized patients. *Infect Control*. 1981;2:380–386.
- Tambyah PA, Maki DG. Catheter-associated urinary tract infection is rarely symptomatic: a prospective study of 1,497 catheterized patients. *Arch Intern Med*. 2000;160:678–682.
- Warren JW, Damron D, Tenney JH, et al. Fever, bacteremia, and death as complications of bacteriuria in women with long-term urethral catheters. *J Infect Dis*. 1987;155:1151–1158.
- Kreger BE, Craven DE, Carling PC, et al. Gram-negative bacteremia. III. Reassessment of etiology, epidemiology and ecology in 612 patients. *Am J Med*. 1980;68:332–343.
- Fortin E, Rocher I, Frenette C, et al. Healthcare-associated bloodstream infections secondary to a urinary focus: the Quebec provincial surveillance results. *Infect Control Hosp Epidemiol*. 2012;33:456–462.
- Kizilbash QF, Petersen NJ, Chen GJ, et al. Bacteremia and mortality with urinary catheter-associated bacteriuria. *Infect Control Hosp Epidemiol*. 2013;34:1153–1159.
- Conway LJ, Liu J, Harris AD, et al. Risk factors for bacteremia in patients with urinary catheter-associated bacteriuria. *Am J Crit Care*. 2016;26:43–52.
- Muder RR, Brennen C, Wagener MM, et al. Bacteremia in a long-term-care facility: a five-year prospective study of 163 consecutive episodes. *Clin Infect Dis*. 1992;14:647–654.
- Rudman D, Hontanosas A, Cohen Z, et al. Clinical correlates of bacteremia in a Veterans Administration extended care facility. *J Am Geriatr Soc*. 1988;36:726–732.
- Warren JW, Muncie HL Jr, Hebel JR, et al. Long-term urethral catheterization increases risk of chronic pyelonephritis and renal inflammation. *J Am Geriatr Soc*. 1994;42:1286–1290.
- Chant C, Smith OM, Marshall JC, et al. Relationship of catheter-associated urinary tract infection to mortality and length of stay in critically ill patients: a systematic review and meta-analysis of observational studies. *Crit Care Med*. 2011;39:1167–1173.
- Kunin CM, Douthitt S, Dancing J, et al. The association between the use of urinary catheters and morbidity and mortality among elderly patients in nursing homes. *Am J Epidemiol*. 1992;135:291–301.
- Kunin CM, Chin QF, Chambers S. Morbidity and mortality associated with indwelling urinary catheters in elderly patients in a nursing home—confounding due to the presence of associated diseases. *J Am Geriatr Soc*. 1987;35:1001–1006.
- Nicolle LE. Catheter-related urinary tract infection. *Drugs Aging*. 2005;22:627–639.
- Spivak ES, Burk M, Zhang R, et al. Management of bacteriuria in veterans affairs hospitals. *Clin Infect Dis*. 2017;65:910–917.
- Cope M, Cevallos ME, Cadle RM, et al. Inappropriate treatment of catheter-associated asymptomatic bacteriuria in a tertiary care hospital. *Clin Infect Dis*. 2009;48:1182–1188.
- Aslam S, Musher DM. An update on diagnosis, treatment, and prevention of *Clostridium difficile*-associated disease. *Gastroenterol Clin North Am*. 2006;35:315–335.
- Rotjanapan P, Dosa D, Thomas KS. Potentially inappropriate treatment of urinary tract infections in two Rhode Island nursing homes. *Arch Intern Med*. 2011;171:438–443.
- Schaberg DR, Haley RW, Highsmith AK, et al. Nosocomial bacteriuria: a prospective study of case clustering and antimicrobial resistance. *Ann Intern Med*. 1980;93:420–424.
- Jarlier V, Fosse T, Philippon A. Antibiotic susceptibility in aerobic gram-negative bacilli isolated in intensive care units in 39 French teaching hospitals (ICU study). *Intensive Care Med*. 1996;22:1057–1065.
- Bjork DT, Pelletier LL, Tigh RR. Urinary tract infections with antibiotic resistant organisms in catheterized nursing home patients. *Infect Control*. 1984;5:173–176.
- Wagenlehner FM, Krcmery S, Held C, et al. Epidemiological analysis of the spread of pathogens from a urological ward using genotypic, phenotypic and clinical parameters. *Int J Antimicrob Agents*. 2002;19:583–591.
- Hooton TM, Stamm WE. Diagnosis and treatment of uncomplicated urinary tract infection. *Infect Dis Clin North Am*. 1997;11:551–581.
- Kline KA, Lewis AL. Gram-positive uropathogens, polymicrobial urinary tract infection, and the emerging microbiota of the urinary tract. *Microbiol Spectr*. 2016;4.
- Warren JW, Tenney JH, Hoopes JM, et al. A prospective microbiologic study of bacteriuria in patients with chronic indwelling urethral catheters. *J Infect Dis*. 1982;146:719–723.
- Rahav G, Pinco E, Silbaq F, et al. Molecular epidemiology of catheter-associated bacteriuria in nursing home patients. *J Clin Microbiol*. 1994;32:1031–1034.
- Bergqvist D, Bronnestam R, Hedelin H, et al. The relevance of urinary sampling methods in patients with indwelling Foley catheters. *Br J Urol*. 1980;52:92–95.
- Tenney JH, Warren JW. Bacteriuria in women with long-term catheters: paired comparison of indwelling and replacement catheters. *J Infect Dis*. 1988;157:199–202.
- Mulvey MA, Schilling JD, Hultgren SJ. Establishment of a persistent *Escherichia coli* reservoir during the acute phase of a bladder infection. *Infect Immun*. 2001;69:4572–4579.
- Jacobsen SM, Stickler DJ, Mobley HL, et al. Complicated catheter-associated urinary tract infections due to *Escherichia coli* and *Proteus mirabilis*. *Clin Microbiol Rev*. 2008;21:26–59.
- Elliott TSJ, Reed L, Slack RCB, et al. Bacteriology and ultrastructure of the bladder in patients with urinary tract infections. *J Infect*. 1985;11:191–199.
- Elliott TSJ, Slack RCB, Bishop MC. Scanning electron microscopy of human bladder mucosa in acute and chronic urinary tract infection. *Br J Urol*. 1984;56:38–43.
- Rosen DA, Hooton TM, Stamm WE, et al. Detection of intracellular bacterial communities in human urinary tract infection. *PLoS Med*. 2007;4:e329.
- Robino L, Scavone P, Araujo L, et al. Intracellular bacteria in the pathogenesis of *Escherichia coli* urinary tract infection in children. *Clin Infect Dis*. 2014;59:e158–e164.
- Warren JW, Mobley HLT, Donnenberg MS. Host-parasite interactions and host defense mechanisms. In: Schrier RW, ed. *Diseases of the Kidney and Urinary Tract*. 7th ed. Philadelphia: Lippincott Williams & Wilkins; 2001: 903–921.
- Johnson JR. Microbial virulence determinants and the pathogenesis of urinary tract infection. *Infect Dis Clin North Am*. 2003;17:261–278.
- Trautner BW, Patterson JE, Petersen NJ, et al. Quality gaps in documenting urinary catheter use and infectious outcomes. *Infect Control Hosp Epidemiol*. 2013;34:793–799.
- Rutala WA, Kennedy VA, Loflin HB, et al. *Serratia marcescens* nosocomial infections of the urinary tract associated with urine measuring containers and urinometers. *Am J Med*. 1981;70:659–663.
- Stamm WE. Catheter-associated urinary tract infections: epidemiology, pathogenesis, and prevention. *Am J Med*. 1991;91:655–715.
- Ikäheimo R, Siitonen A, Kärkkäinen U, et al. Virulence characteristics of *Escherichia coli* in nosocomial urinary tract infection. *Clin Infect Dis*. 1993;16:785–791.
- Tambyah PA, Halvorson KT, Maki DG. A prospective study of pathogenesis of catheter-associated urinary tract infections. *Mayo Clin Proc*. 1999;74:131–136.
- Stark RP, Maki DG. Bacteriuria in the catheterized patient. What quantitative level of bacteriuria is relevant? *N Engl J Med*. 1984;311:560–564.
- Norden CW, Kass EH. Bacteriuria of pregnancy—a critical appraisal. *Annu Rev Med*. 1968;19:431–470.
- Gleckman R, Esposito A, Crowley M, et al. Reliability of a single urine culture in establishing diagnosis of asymptomatic bacteriuria in adult males. *J Clin Microbiol*. 1979;9:596–597.
- Nicolle LE, Harding GK, Kennedy J, et al. Urine specimen collection with external devices for diagnosis of bacteriuria in elderly incontinent men. *J Clin Microbiol*. 1988;26:1115–1119.
- Stamm WE, Counts GW, Running KR, et al. Diagnosis of coliform infection in acutely dysuric women. *N Engl J Med*. 1982;307:463–468.
- Hooton TM, Roberts PL, Cox ME, et al. Voided midstream urine culture and acute cystitis in premenopausal women. *N Engl J Med*. 2013;369:1883–1891.
- Lipsky BA, Ireton RC, Fihn SD, et al. Diagnosis of bacteriuria in men: specimen collection and culture interpretation. *J Infect Dis*. 1987;155:847–854.
- The prevention and management of urinary tract infections among people with spinal cord injuries. National Institute on Disability and Rehabilitation Research Consensus Statement. January 27–29, 1992. *J Am Paraplegia Soc*. 1992;15:194–204.
- Glob JF Jr, Claridge JA, Sando MJ, et al. Fever and leukocytosis in critically ill trauma patients: it's not the urine. *Surg Infect (Larchmt)*. 2008;9:49–56.
- Modi L, Juthani-Mehta M. Urinary tract infections in older women: a clinical review. *JAMA*. 2014;311:844–854.

78. Tambyah PA, Maki DG. The relationship between pyuria and infection in patients with indwelling urinary catheters: a prospective study of 761 patients. *Arch Intern Med.* 2000;160:673-677.
79. Steward DK, Wood GL, Cohen RL, et al. Failure of the urinalysis and quantitative urine culture in diagnosing symptomatic urinary tract infections in patients with long-term urinary catheters. *Am J Infect Control.* 1985;13:154-160.
80. Haley RW, Culver DH, White JW, et al. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol.* 1985;121:182-205.
81. Wald HL, Kramer AM. Nonpayment for harms resulting from medical care: catheter-associated urinary tract infections. *JAMA.* 2007;298:2782-2784.
82. Brown J, Doloresco Ili F, Mylotte JM. "Never events": not every hospital-acquired infection is preventable. *Clin Infect Dis.* 2009;49:743-746.
83. Krein SL, Kowalski CP, Harrod M, et al. Barriers to reducing urinary catheter use: a qualitative assessment of a statewide initiative. *JAMA Intern Med.* 2013;173:881-886.
84. Saint S, Greene MT, Krein SL, et al. A program to prevent catheter-associated urinary tract infection in acute care. *N Engl J Med.* 2016;374:2111-2119.
85. Mody L, Greene MT, Meddings J, et al. A national implementation project to prevent catheter-associated urinary tract infection in nursing home residents. *JAMA Intern Med.* 2017;177:1154-1162.
86. Gould CV, Umscheid CA, Agarwal RK, et al; Healthcare Infection Control Practices Advisory Committee (HICPAC). Guideline for prevention of catheter-associated urinary tract infections 2009. *Infect Control Hosp Epidemiol.* 2010;31:319-326.
87. Lo E, Nicolle LE, Coffin SE, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol.* 2014;35(suppl 2):S32-S47.
88. Jain M, Miller L, Belt D, et al. Decline in ICU adverse events, nosocomial infections and cost through a quality improvement initiative focusing on teamwork and culture change. *Qual Saf Health Care.* 2006;15:235-239.
89. Fakih MG, Watson SR, Greene MT, et al. Reducing inappropriate urinary catheter use: a statewide effort. *Arch Intern Med.* 2012;172:255-260.
90. Rosenthal VD, Todi SK, Álvarez-Moreno C, et al. Impact of a multidimensional infection control strategy on catheter-associated urinary tract infection rates in the adult intensive care units of 15 developing countries: findings of the International Nosocomial Infection Control Consortium. *Infection.* 2012;40:517-526.
91. Marra AR. Preventing catheter-associated urinary tract infection in the zero tolerance era. *Am J Infect Control.* 2011;39:817-821.
92. Saint S, Meddings JA, Calfee D, et al. Catheter-associated urinary tract infection and the Medicare rule changes. *Ann Intern Med.* 2009;150:877-884.
93. Saint S, Greene MT, Kowalski CP, et al. Preventing catheter-associated urinary tract infection in the United States: a national comparative study. *JAMA Intern Med.* 2013;173:874-879.
94. Fink R, Gilmartin H, Richard A, et al. Indwelling urinary catheter management and catheter-associated urinary tract infection prevention practices in Nurses Improving Care for Healthsystem Elders hospitals. *Am J Infect Control.* 2012;40:715-720.
95. Krein SL, Fowler KE, Ratz D, et al. Preventing device-associated infections in US hospitals: national surveys from 2005 to 2013. *BMJ Qual Saf.* 2015;24:385-392.
96. Lee GM, Kleinman K, Soumerai SB, et al. Effect of nonpayment for preventable infections in U.S. hospitals. *N Engl J Med.* 2012;367:1428-1437.
97. Smith PW, Bennett G, Bradley S, et al. SHEA/APIC guideline: infection prevention and control in the long-term care facility. *Infect Control Hosp Epidemiol.* 2008;29:785-814.
98. Meddings J, Saint S, Krein SL, et al. Systematic review of interventions to reduce urinary tract infection in nursing home residents. *J Hosp Med.* 2017;12:356-368.
99. Meddings J, Rogers MA, Macy M, et al. Systematic review and meta-analysis: reminder systems to reduce catheter-associated urinary tract infections and urinary catheter use in hospitalized patients. *Clin Infect Dis.* 2010;51:550-560.
100. Leuck AM, Wright D, Ellingson L, et al. Complications of Foley catheters—is infection the greatest risk? *J Urol.* 2012;187:1662-1666.
101. Jain P, Parada JP, David A, et al. Overuse of the indwelling urinary tract catheter in hospitalized medical patients. *Arch Intern Med.* 1995;155:1425-1429.
102. Munasinghe RL, Yazdani H, Siddique M, et al. Appropriateness of use of indwelling urinary catheters in patients admitted to the medical service. *Infect Control Hosp Epidemiol.* 2001;22:647-649.
103. Gardam AM, Amihod B, Orenstein P, et al. Overutilization of indwelling urinary catheters and the development of nosocomial urinary tract infections. *Clin Perform Qual Health Care.* 1998;6:99-102.
104. Gokula RR, Hickner JA, Smith MA. Inappropriate use of urinary catheters in elderly patients at a Midwestern community teaching hospital. *Am J Infect Control.* 2004;32:196-199.
105. Saint S, Wiese J, Amory JK, et al. Are physicians aware of which of their patients have indwelling urinary catheters? *Am J Med.* 2000;109:476-480.
106. Gokula RM, Smith MA, Hickner J. Emergency room staff education and use of a urinary catheter indication sheet improves appropriate use of Foley catheters. *Am J Infect Control.* 2007;35:589-593.
107. Stephan F, Sax H, Wachsmuth M, et al. Reduction of urinary tract infection and antibiotic use after surgery: a controlled, prospective, before-after intervention study. *Clin Infect Dis.* 2006;42:1544-1551.
108. Slappendel R, Weber EW. Non-invasive measurement of bladder volume as an indication for bladder catheterization after orthopaedic surgery and its effect on urinary tract infections. *Eur J Anaesthesiol.* 1999;16:503-506.
109. Lau H, Lam B. Management of postoperative urinary retention: a randomized trial of in-out versus overnight catheterization. *ANZ J Surg.* 2004;74:658-661.
110. Kunin C. Nosocomial urinary tract infections and the indwelling catheter. What is new and what is true? *Chest.* 2001;120:10-12.
111. Stevens E. Bladder ultrasound: avoiding unnecessary catheterizations. *Medsurg Nurs.* 2005;14:249-253.
112. Weber DJ, Kang J, Brown VM, et al. Preventing catheter-associated urinary tract infections: hospital location of catheter insertion. *Infect Control Hosp Epidemiol.* 2012;33:1057-1058.
113. Murphy C, Fader M, Prieto J. Interventions to minimise the initial use of indwelling urinary catheters in acute care: a systematic review. *Int J Nurs Stud.* 2014;51:4-13.
114. Huang WC, Wann SR, Lin SL, et al. Catheter-associated urinary tract infections in intensive care units can be reduced by prompting physicians to remove unnecessary catheters. *Infect Control Hosp Epidemiol.* 2004;25:974-978.
115. Apisarnthanarak A, Thongphubeth K, Sirinvaravong S, et al. Effectiveness of multifaceted hospital-wide quality improvement programs featuring an intervention to remove unnecessary urinary catheters at a tertiary care center in Thailand. *Infect Control Hosp Epidemiol.* 2007;28:791-798.
116. Saint S, Kaufman SR, Thompson M, et al. A reminder reduces urinary catheterization in hospitalized patients. *Jt Comm J Qual Patient Saf.* 2005;31:455-462.
117. Cornia PB, Amory JK, Fraser S, et al. Computer-based order entry decreases duration of indwelling urinary catheterization in hospitalized patients. *Am J Med.* 2003;114:404-407.
118. Topal J, Conklin S, Camp K, et al. Prevention of nosocomial catheter-associated urinary tract infections through computerized feedback to physicians and a nurse-directed protocol. *Am J Med Qual.* 2005;20:121-126.
119. Loeb M, Hunt D, O'Halloran K, et al. Stop orders to reduce inappropriate urinary catheterization in hospitalized patients: a randomized controlled trial. *J Gen Intern Med.* 2008;23:816-820.
120. Bruminhent J, Keegan M, Lakhani A, et al. Effectiveness of a simple intervention for prevention of catheter-associated urinary tract infections in a community teaching hospital. *Am J Infect Control.* 2010;38:689-693.
121. Elpern EH, Killen K, Ketchum A, et al. Reducing use of indwelling urinary catheters and associated urinary tract infections. *Am J Crit Care.* 2009;18:535-541.
122. Fakih MG, Dueweke C, Meisner S, et al. Effect of nurse-led multidisciplinary rounds on reducing the unnecessary use of urinary catheterization in hospitalized patients. *Infect Control Hosp Epidemiol.* 2008;29:815-819.
123. Griffiths R, Fernandez R. Strategies for the removal of short-term indwelling urethral catheters in adults. *Cochrane Database Syst Rev.* 2007;(2):CD004011.
124. Meddings J, Rogers MA, Krein SL, et al. Reducing unnecessary urinary catheter use and other strategies to prevent catheter-associated urinary tract infection: an integrative review. *BMJ Qual Saf.* 2014;23:277-289.
125. Kidd EA, Stewart F, Kassiss NC, et al. Urethral (indwelling or intermittent) or suprapubic routes for short-term catheterisation in hospitalised adults. *Cochrane Database Syst Rev.* 2015;(12):CD004203.
126. Saint S, Kaufman SR, Rogers MA, et al. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54:1055-1061.
127. Jamison J, Maguire S, McCann J. Catheter policies for management of long term voiding problems in adults with neurogenic bladder disorders. *Cochrane Database Syst Rev.* 2013;(11):CD004375.
128. Saint S, Kaufman SR, Rogers MA, et al. Condom versus indwelling urinary catheters: a randomized trial. *J Am Geriatr Soc.* 2006;54:1055-1061.
129. Consortium for Spinal Cord Medicine. Bladder management for adults with spinal cord injury: a clinical practice guideline for health-care providers. *J Spinal Cord Med.* 2006;29:527-573.
130. Guttman L, Frankel H. The value of intermittent catheterization in the early management of traumatic paraplegia and tetraplegia. *Paraplegia.* 1966;4:63-84.
131. Lapidis J, Diokno AC, Silber SJ, et al. Clean, intermittent self-catheterization in the treatment of urinary tract disease. *J Urol.* 1972;107:458-461.
132. Niel-Weise BS, van den Broek PJ, da Silva EM, et al. Urinary catheter policies for long-term bladder drainage. *Cochrane Database Syst Rev.* 2012;(8):CD004201.
133. Wyndaele JJ. Complications of intermittent catheterization: their prevention and treatment. *Spinal Cord.* 2002;40:536-541.
134. Duffy LM, Cleary J, Ahern S, et al. Clean intermittent catheterization: safe, cost-effective bladder management for male residents of VA nursing homes. *J Am Geriatr Soc.* 1995;43:865-870.
135. Jamison J, Maguire S, McCann J. Catheter policies for management of long term voiding problems in adults with neurogenic bladder disorders. *Cochrane Database Syst Rev.* 2011;(12):CD004375.
136. Moore KN, Fader M, Getliffe K. Long-term bladder management by intermittent catheterisation in adults and children. *Cochrane Database Syst Rev.* 2007;(4):CD006008.
137. Rognoni C, Tarricone R. Intermittent catheterisation with hydrophilic and non-hydrophilic urinary catheters: systematic literature review and meta-analyses. *BMC Urol.* 2017;17:4.
138. Cardenas DD, Moore KN, Dannels-McClure A, et al. Intermittent catheterization with a hydrophilic-coated catheter delays urinary tract infections in acute spinal cord injury: a prospective, randomized, multicenter trial. *PM R.* 2011;3:408-417.
139. Healy EF, Walsh CA, Cotter AM, et al. Suprapubic compared with transurethral bladder catheterization for gynecologic surgery: a systematic review and meta-analysis. *Obstet Gynecol.* 2012;120:678-687.
140. Lo E, Nicolle L, Coffin SE, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol.* 2014;35:464-479.
141. Carapeti EA, Andrews SM, Bentley PG. Randomised study of sterile versus non-sterile urethral catheterisation. *Ann R Coll Surg Engl.* 1996;78:59-60.
142. Kass EH. Asymptomatic infections of the urinary tract. *Trans Assoc Am Physicians.* 1956;69:56-64.
143. Warren JW, Platt R, Thomas RJ, et al. Antibiotic irrigation and catheter-associated urinary-tract infections. *N Engl J Med.* 1978;299:570-573.
144. Meddings J, Reichert H, Greene MT, et al. Evaluation of the association between Hospital Survey on Patient Safety Culture (HSOPS) measures and catheter-associated infections: results of two national collaboratives. *BMJ Qual Saf.* 2017;26:226-235.
145. Ahearn DG, Grace DT, Jennings MJ, et al. Effects of hydrogel/silver coatings on in vitro adhesion to catheters of bacteria associated with urinary tract infections. *Curr Microbiol.* 2000;41:120-125.
146. Darouiche RO, Safar H, Raad II. In vitro efficacy of antimicrobial-coated bladder catheters in inhibiting bacterial migration along catheter surface. *J Infect Dis.* 1997;176:1109-1112.
147. Johnson JR, Delavari P, Azar M. Activities of a nitrofurazone-containing urinary catheter and a silver hydrogel catheter against multidrug-resistant bacteria characteristic of catheter-associated urinary tract infection. *Antimicrob Agents Chemother.* 1999;43:2990-2995.
148. Desai DG, Liao KS, Cevallos ME, et al. Silver or nitrofurazone impregnation of urinary catheters has a minimal effect on uropathogen adherence. *J Urol.* 2010;184:2565-2571.
149. Johnson JR, Johnston B, Kuskowski MA. In vitro comparison of nitrofurazone- and silver alloy-coated Foley catheters for contact-dependent and diffusible inhibition of urinary tract infection-associated microorganisms. *Antimicrob Agents Chemother.* 2012;56:4969-4972.

150. Schumm K, Lam TB. Types of urethral catheters for management of short-term voiding problems in hospitalised adults. *Cochrane Database Syst Rev*. 2008;(2):CD004013.
151. Drekonja DM, Kuskowski MA, Wilt TJ, et al. Antimicrobial urinary catheters: a systematic review. *Expert Rev Med Devices*. 2008;5:495–506.
152. Johnson JR, Kuskowski MA, Wilt TJ. Systematic review: antimicrobial urinary catheters to prevent catheter-associated urinary tract infection in hospitalized patients. *Ann Intern Med*. 2006;144:116–126.
153. Jahn P, Beutner K, Langer G. Types of indwelling urinary catheters for long-term bladder drainage in adults. *Cochrane Database Syst Rev*. 2012;(10):CD004997.
154. Pickard R, Lam T, MacLennan G, et al. Types of urethral catheter for reducing symptomatic urinary tract infections in hospitalised adults requiring short-term catheterisation: multicentre randomised controlled trial and economic evaluation of antimicrobial- and antiseptic-impregnated urethral catheters (the CATHETER trial). *Health Technol Assess*. 2012;16:1–197.
155. Pickard R, Lam T, MacLennan G, et al. Antimicrobial catheters for reduction of symptomatic urinary tract infection in adults requiring short-term catheterisation in hospital: a multicentre randomised controlled trial. *Lancet*. 2012;380:1927–1935.
156. Lam TB, Omar MI, Fisher E, et al. Types of indwelling urethral catheters for short-term catheterisation in hospitalised adults. *Cochrane Database Syst Rev*. 2014;(9):CD004013.
157. Lusardi G, Lipp A, Shaw C. Antibiotic prophylaxis for short-term catheter bladder drainage in adults. *Cochrane Database Syst Rev*. 2013;(7):CD005428.
158. Rutschmann OT, Zwahlen A. Use of norfloxacin for prevention of symptomatic urinary tract infection in chronically catheterized patients. *Eur J Clin Microbiol Infect Dis*. 1995;14:441–444.
159. Trautner BW, Hull RA, Thornby JJ, et al. Coating urinary catheters with an avirulent strain of *Escherichia coli* as a means to establish asymptomatic colonization. *Infect Control Hosp Epidemiol*. 2007;28:92–94.
160. Prasad A, Cevallos ME, Riosa S, et al. A bacterial interference strategy for prevention of UTI in persons practicing intermittent catheterization. *Spinal Cord*. 2009;47:565–569.
161. Horwitz D, McCue T, Mapes AC, et al. Decreased microbiota diversity associated with urinary tract infection in a trial of bacterial interference. *J Infect*. 2015;71:358–367.
162. Hustinx WN, Mintjes-de Groot AJ, Verkooyen RP, et al. Impact of concurrent antimicrobial therapy on catheter-associated urinary tract infection. *J Hosp Infect*. 1991;18:45–56.
163. Gleckman R, Alvarez S, Joubert DW, et al. Drug therapy reviews: methenamine mandelate and methenamine hippurate. *Am J Hosp Pharm*. 1979;36:1509–1512.
164. Cronberg S, Welin CO, Henriksson L, et al. Prevention of recurrent acute cystitis by methenamine hippurate: double blind controlled crossover long term study. *Br Med J (Clin Res Ed)*. 1987;294:1507–1508.
165. Harding GK, Ronald AR. A controlled study of antimicrobial prophylaxis of recurrent urinary infection in women. *N Engl J Med*. 1974;291:597–601.
166. Lee BS, Bhuta T, Simpson JM, et al. Methenamine hippurate for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD003265.
167. Schiotz HA, Guttu K. Value of urinary prophylaxis with methenamine in gynecologic surgery. *Acta Obstet Gynecol Scand*. 2002;81:743–746.
168. Lee BB, Haran MJ, Hunt LM, et al. Spinal-injured neuropathic bladder antiseptic (SINBA) trial. *Spinal Cord*. 2007;45:542–550.
169. Kuhlmeier KV, Stover SL, Lloyd LK. Prophylactic antibiotic therapy for preventing urinary tract infections in spinal cord injury patients. *J Urol*. 1985;134:514–517.
170. Webster J, Hood RH, Burrage CA, et al. Water or antiseptic for periurethral cleansing before urinary catheterization: a randomized controlled trial. *Am J Infect Control*. 2001;29:389–394.
171. Cheung K, Leung P, Wong YC, et al. Water versus antiseptic periurethral cleansing before catheterization among home care patients: a randomized controlled trial. *Am J Infect Control*. 2008;36:375–380.
172. Jepson RG, Williams G, Craig JC. Cranberries for preventing urinary tract infections. *Cochrane Database Syst Rev*. 2012;(10):CD001321.
173. Juthani-Mehra M, Van Ness PH, Bianco L, et al. Effect of cranberry capsules on bacteriuria plus pyuria among older women in nursing homes: a randomized clinical trial. *JAMA*. 2016;316:1879–1887.
174. Shepherd AJ, Mackay WG, Hagen S. Washout policies in long-term indwelling urinary catheterisation in adults. *Cochrane Database Syst Rev*. 2017;(3):CD004012.
175. Thompson RL, Haley CE, Searcy MA, et al. Catheter-associated bacteriuria. Failure to reduce attack rates using periodic instillations of a disinfectant into urinary drainage systems. *JAMA*. 1984;251:747–751.
176. Priefer BA, Duthie EH Jr, Gambert SR. Frequency of urinary catheter changed and clinical urinary tract infection: study in hospital-based, skilled nursing home. *Urology*. 1982;20:141–142.
177. Kunin CM, Chin QF, Chambers S. Indwelling urinary catheters in the elderly. Relation of “catheter life” to formation of encrustations in patients with and without blocked catheters. *Am J Med*. 1987;82:405–411.
178. Jewes LA, Gillespie WA, Leadbetter A, et al. Bacteriuria and bacteraemia in patients with long-term indwelling catheters—a domiciliary study. *J Med Microbiol*. 1988;26:61–65.
179. Polastri F, Auckenthaler R, Loew F, et al. Absence of significant bacteremia during urinary catheter manipulation in patients with chronic indwelling catheters. *J Am Geriatr Soc*. 1990;38:1203–1208.
180. Bregenzer T, Frei R, Widmer AE, et al. Low risk of bacteremia during catheter replacement in patients with long-term urinary catheters. *Arch Intern Med*. 1997;157:521–525.
181. Harding GK, Nicolle LE, Ronald AR, et al. How long should catheter-acquired urinary tract infection in women be treated? A randomized controlled study. *Ann Intern Med*. 1991;114:713–719.
182. Marshall J, Carpenter CR, Fowler S, et al. Program CDCPE. Antibiotic prophylaxis for urinary tract infections after removal of urinary catheter: meta-analysis. *BMJ*. 2013;346:f3147.
183. Darouiche RO, Hull RA. Bacterial interference for prevention of urinary tract infection. *Clin Infect Dis*. 2012;55:1400–1407.
184. Guion PS, Cusumano CK, Kline KA, et al. Combinatorial small-molecule therapy prevents uropathogenic *Escherichia coli* catheter-associated urinary tract infections in mice. *Antimicrob Agents Chemother*. 2012;56:4738–4745.
185. Spaulding CN, Klein RD, Ruder S, et al. Selective depletion of uropathogenic *E. coli* from the gut by a FimH antagonist. *Nature*. 2017;546:528–532.
186. U.S. Preventive Services Task Force. Screening for asymptomatic bacteriuria in adults: U.S. Preventive Services Task Force reaffirmation recommendation statement. *Ann Intern Med*. 2008;149:43–47.
187. Warren JW, Anthony WC, Hoopes JM, et al. Cephalixin for susceptible bacteriuria in afebrile, long-term catheterized patients. *JAMA*. 1982;248:454–458.
188. Trautner BW. Asymptomatic bacteriuria: when the treatment is worse than the disease. *Nat Rev Urol*. 2011;9:85–93.
189. Trautner BW, Grigoryan L. Approach to a positive urine culture in a patient without urinary symptoms. *Infect Dis Clin North Am*. 2014;28:15–31.
190. Skelton F, Grigoryan L, Holmes SA, et al. Routine urine testing at the spinal cord injury annual evaluation leads to unnecessary antibiotic use: a pilot study and future directions. *Arch Phys Med Rehabil*. 2017.
191. Drekonja DM, Abbo LM, Kuskowski MA, et al. A survey of resident physicians’ knowledge regarding urine testing and subsequent antimicrobial treatment. *Am J Infect Control*. 2013;41:892–896.
192. American Geriatrics Society. *Ten Things Clinicians and Patients Should Question*. Philadelphia: Choosing Wisely/ABIM Foundation; 2015. <http://www.choosingwisely.org/societies/american-geriatrics-society>.
193. Infectious Diseases Society of America. *Five Things Physicians and Patients Should Question*. Philadelphia: Choosing Wisely/ABIM Foundation; 2015. <http://www.choosingwisely.org/societies/infectious-diseases-society-of-america>.
194. Trautner BW, Petersen NJ, Hysong SJ, et al. Overtreatment of asymptomatic bacteriuria: identifying provider barriers to evidence-based care. *Am J Infect Control*. 2014;42:653–658.
195. Leis JA, Rebick GW, Daneman N, et al. Reducing antimicrobial therapy for asymptomatic bacteriuria among noncatheterized inpatients: a proof-of-concept study. *Clin Infect Dis*. 2014;58:980–983.
196. Fakih MG, Khatib R. Improving the culture of culturing: critical asset to antimicrobial stewardship. *Infect Control Hosp Epidemiol*. 2017;38:377–379.
197. Trautner BW, Grigoryan L, Petersen NJ, et al. Effectiveness of an antimicrobial stewardship approach for urinary catheter-associated asymptomatic bacteriuria. *JAMA Intern Med*. 2015;175:1120–1127.
198. Al-Qas Hanna F, Sambirka O, Iyer S, et al. Clinician practice and the National Healthcare Safety Network definition for the diagnosis of catheter-associated urinary tract infection. *Am J Infect Control*. 2013;41:1173–1177.
199. Mullin KM, Kovacs CS, Fatica C, et al. A multifaceted approach to reduction of catheter-associated urinary tract infections in the intensive care unit with an emphasis on “stewardship of culturing”. *Infect Control Hosp Epidemiol*. 2017;38:186–188.
200. Trautner BW, Greene MT, Krein SL, et al. Infection prevention and antimicrobial stewardship knowledge for selected infections among nursing home personnel. *Infect Control Hosp Epidemiol*. 2017;38:83–88.
201. Horstman MJ, Spiegelman A, Naik AD, et al. National patterns of urine testing during inpatient admission. *Clin Infect Dis*. 2017;65:1199–1205.
202. Stamm WE, Hooton TM. Management of urinary tract infections in adults. *N Engl J Med*. 1993;329:1328–1334.
203. Raz R, Schiller D, Nicolle LE. Chronic indwelling catheter replacement before antimicrobial therapy for symptomatic urinary tract infection. *J Urol*. 2000;164:1254–1258.
204. Johnson JR, Russo TA. Acute pyelonephritis in adults. *N Engl J Med*. 2018;378:1162.
205. Arcilla MS, van Hattem JM, Haverkate MR, et al. Import and spread of extended-spectrum beta-lactamase-producing Enterobacteriaceae by international travellers (COMBAT study): a prospective, multicentre cohort study. *Lancet Infect Dis*. 2017;17:78–85.
206. Reuland EA, Sonder GJ, Stolte I, et al. Travel to Asia and traveller’s diarrhoea with antibiotic treatment are independent risk factors for acquiring ciprofloxacin-resistant and extended spectrum beta-lactamase-producing Enterobacteriaceae—a prospective cohort study. *Clin Microbiol Infect*. 2016;22:731 e731–731 e737.
207. Linsenmeyer K, Strymish J, Gupta K. Two simple rules for improving the accuracy of empiric treatment of multidrug-resistant urinary tract infections. *Antimicrob Agents Chemother*. 2015;59:7593–7596.
208. Hamprecht A, Rohde AM, Behnke M, et al. Colonization with third-generation cephalosporin-resistant Enterobacteriaceae on hospital admission: prevalence and risk factors. *J Antimicrob Chemother*. 2016;71:2957–2963.
209. Mody L, Krein SL, Saint S, et al. A targeted infection prevention intervention in nursing home residents with indwelling devices: a randomized clinical trial. *JAMA Intern Med*. 2015;175:714–723.
210. Cek M, Tandogdu Z, Wagenlehner F, et al. Healthcare-associated urinary tract infections in hospitalized urological patients—a global perspective: results from the GPIU studies 2003–2010. *World J Urol*. 2014;32:1587–1594.
211. Nicolle LE, Bjornson J, Harding GK, et al. Bacteriuria in elderly institutionalized men. *N Engl J Med*. 1983;309:1420–1425.
212. Dow G, Rao P, Harding G, et al. A prospective, randomized trial of 3 or 14 days of ciprofloxacin treatment for acute urinary tract infection in patients with spinal cord injury. *Clin Infect Dis*. 2004;39:658–664.
213. Peterson J, Kaul S, Khashab M, et al. A double-blind, randomized comparison of levofloxacin 750 mg once-daily for five days with ciprofloxacin 400/500 mg twice-daily for 10 days for the treatment of complicated urinary tract infections and acute pyelonephritis. *Urology*. 2008;71:17–22.
214. Rivera M, Viers B, Cockerill P, et al. Pre- and postoperative predictors of infection-related complications in patients undergoing percutaneous nephrolithotomy. *J Endourol*. 2016;30:982–986.
215. Singh P, Yadav S, Singh A, et al. Systemic inflammatory response syndrome following percutaneous nephrolithotomy: assessment of risk factors and their impact on patient outcomes. *Urol Int*. 2016;96:207–211.
216. Lara-Isa A, Medina-Polo J, Alonso-Isa M, et al. Urinary infections in patients with catheters in the upper urinary tract: microbiological study. *Urol Int*. 2017;98:442–448.
217. Bahu R, Chaffari AM, Hachem RY, et al. Nephrostomy tube related pyelonephritis in patients with cancer: epidemiology, infection rate and risk factors. *J Urol*. 2013;189:130–135.
218. Siddiq DM, Darouiche RO. Infectious complications associated with percutaneous nephrostomy catheters: do we know enough? *Int J Artif Organs*. 2012;35:898–907.
219. Tan WP, Turba UC, Deane LA. Renal fungus ball: a challenging clinical problem. *Urologia*. 2017;84:113–115.
220. Deleted in review.
221. Deleted in review.
222. Deshmukh S, Sternberg K, Hernandez N, et al. Compliance with American Urological Association

- guidelines for post-percutaneous nephrolithotomy antibiotics does not appear to increase rates of infection. *J Urol*. 2015;194:992–996.
223. Sobel JD, Fisher JF, Kauffman CA, et al. Candida urinary tract infections—epidemiology. *Clin Infect Dis*. 2011;52(suppl 6):S433–S436.
 224. Kauffman CA, Vazquez JA, Sobel JD, et al. Prospective multi-center surveillance study of funguria in hospitalized patients. The National Institute for Allergy and Infectious Diseases (NIAID) Mycoses Study Group. *Clin Infect Dis*. 2000;30:14–18.
 225. Pappas PG, Kauffman CA, Andes DR, et al. Clinical Practice Guideline for the Management of Candidiasis: 2016 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;62:e1–e50.
 226. Kauffman CA. Candiduria. *Clin Infect Dis*. 2005;41:S371–S376.
 227. Ang BS, Telenti A, King B, et al. Candidemia from a urinary tract source: microbiological aspects and clinical significance. *Clin Infect Dis*. 1993;17:662–666.
 228. Binelli CA, Moretti ML, Assis RS, et al. Investigation of the possible association between nosocomial candiduria and candidaemia. *Clin Microbiol Infect*. 2006;12:538–543.
 229. Lundstrom T, Sobel J. Nosocomial candiduria: a review. *Clin Infect Dis*. 2001;32:1602–1607.
 230. Sobel JD, Kauffman CA, McKinsey D, et al. Candiduria: a randomized, double-blind study of treatment with fluconazole and placebo. The National Institute of Allergy and Infectious Diseases (NIAID) Mycoses Study Group. *Clin Infect Dis*. 2000;30:19–24.
 231. Fisher JF, Sobel JD, Kauffman CA, et al. Candida urinary tract infections—treatment. *Clin Infect Dis*. 2011;52(suppl 6):S457–S466.
 232. Safdar N, Slattery WR, Knasinski V, et al. Predictors and outcomes of candiduria in renal transplant recipients. *Clin Infect Dis*. 2005;40:1413–1421.
 233. Sobel JD, Bradshaw SK, Lipka CJ, et al. Caspofungin in the treatment of symptomatic candiduria. *Clin Infect Dis*. 2007;44:e46–e49.
 234. Lagrotteria D, Rotstein C, Lee CH. Treatment of candiduria with micafungin: a case series. *Can J Infect Dis Med Microbiol*. 2007;18:149–150.
 235. Curtis LT. Prevention of hospital-acquired infections: review of non-pharmacological interventions. *J Hosp Infect*. 2008;69:204–219.

SHORT VIEW SUMMARY

Definition

- Viral hepatitis can be acquired by patients or health care workers (HCWs) during the process of health care delivery.

Epidemiology

- Transmission between patients and from patient to HCW or HCW to patient is rare, but the risk remains. Dialysis patients and workers are at particular risk, especially for hepatitis C virus (HCV) and, if unvaccinated, for hepatitis B virus (HBV).
- Increasingly, transmissions to patients are seen in long-term care settings and can be traced to inappropriate use of blood glucose monitoring devices.
- Transfusion-based risk of hepatitis E varies by geographic region. Implementation of universal screening of blood products varies accordingly.

Diagnosis

- Identification of transmission is difficult unless a cluster of cases occurs.
- Most investigations are conducted in conjunction with public health authorities.
- Use of a genotype to determine homology is useful in investigation.

Postexposure Treatment

- Hepatitis A: Vaccine is the preferred approach for postexposure prophylaxis (PEP) of otherwise healthy individuals. Vaccine and immunoglobulin are recommended for those with immunocompromising or chronic liver conditions.
- Hepatitis B: Give hepatitis B immune globulin (HBIG) to all exposed nonimmune persons (patients or HCWs) with or without initiation of vaccine series. The role of antivirals has not been determined.

- Hepatitis C: Monitor and treat only if the exposed person develops evidence of acute infection. Use of direct-acting antiviral (DAA) therapy has not been studied in this context.

Vaccination

- Hepatitis A vaccine is not routinely recommended for HCWs.
- Hepatitis B vaccine or formal declination is mandated for HCWs in the United States. Rapid decrease in measured antibody levels within months of immunization can occur despite persistence of immune protection. Booster doses are not recommended.
- No vaccine is available for HCV.

HISTORICAL BACKGROUND

The potential for bloodborne transmission of hepatitis B first was noted in 1885, when Lurman described jaundice in factory workers who had received smallpox vaccine prepared from “human lymph.”¹ More reports appeared in the subsequent decades as use of vaccines derived from human serum became more common.² In addition, more frequent use of phlebotomy equipment,³ insulin therapy,⁴ and intramuscular injection of antibiotics all led to small outbreaks of jaundice, which were ascribed to a transmissible “icterogenic” agent.

By the late 1940s, studies to clarify the modes of transmission were undertaken. Central to these was the use of human volunteers, who were given putatively infectious material intradermally, intranasally, or by ingestion of feces, and then observed for development of jaundice.^{2,5–7} From these landmark reports arose our current understanding of the basic principles of transmission of infectious hepatitis (hepatitis A) and serum hepatitis (hepatitis B).

The first report of occupational disease in health care workers (HCWs) was provided by Leibowitz and colleagues,⁸ who described jaundice in a blood bank nurse with numerous needle pricks on her hands and fingers. A spate of similar reports followed, describing occupationally acquired hepatitis among nurses, blood bank workers, phlebotomists, house staff, and others.^{9,10} Soon, the workers’ compensation boards of certain states ruled that viral hepatitis was a compensable occupational hazard. Improved understanding of routes of transmission, more comprehensive and rigorous infection control including needle disposal, and, for hepatitis B, vaccination of workers at risk have helped to decrease, but not eliminate, this occupational risk. A corollary risk, that of transmission of infection from infected HCWs, particularly surgeons, to nonimmune patients, has been described for hepatitis B^{11,12} and hepatitis C.¹³

CURRENT EPIDEMIOLOGY OF HEALTH CARE–ASSOCIATED HEPATITIS B AND C OUTBREAKS IN THE UNITED STATES

A health care–associated outbreak of hepatitis is defined as two or more cases that are epidemiologically and/or genetically linked.^{14,15} Since 2008, The Centers for Disease Control and Prevention (CDC) has maintained a Web-based inventory of health care–associated cases and outbreaks of hepatitis B and C.¹⁶ This source described 59 outbreaks affecting more than 450 people that were investigated between 2008 and 2016 (Fig. 303.1). In addition, 18 (15 hepatitis C and 3 hepatitis B) sporadic cases of patient-to-patient health care–associated transmission of hepatitis B and C were recognized during this time.

Health care–associated outbreaks of hepatitis mostly have been recognized in nonhospital settings, with a growing trend in long-term care facilities¹⁷ and ambulatory areas,¹⁸ especially outpatient hemodialysis units.¹⁶ Practices associated with transmission of hepatitis reveal a shift in etiology: syringe reuse and needlestick injuries have less commonly been the cause, and more cases have been traced to other health care exposures.¹⁹ These include improper use of blood glucose monitoring fingerstick devices on multiple patients,^{20,21} contamination of saline flush and multidose vials,¹⁸ and breaches in disinfection and sterilization of medical equipment.^{19,22}

Hemodialysis-related outbreaks of hepatitis C continue to occur, with most being attributed to environmental contamination or unsafe practices during vascular access or medication preparation or administration.^{16,23} Diversion of medication, wherein a practitioner personally administers an intravenous narcotic and then reuses the remainder of the vial of medication or the needle in a patient or patients, is also increasingly recognized as a source of transmission.²⁴ At least five




FIG. 303.1 (A and B) Hepatitis B and C outbreaks investigated between 2008 and 2016.

TABLE 303.1 Risk of Transmission, Prevention, and Postexposure Management of Health Care Workers With Occupational Exposure to Hepatitis Viruses

HEPATITIS	RATE ^a	PREVENTION	POSTEXPOSURE MANAGEMENT
A	10%–30%	Vaccine not recommended routinely for HCWs except in outbreak setting	Vaccine within 2 weeks of exposure. Immunoglobulin plus vaccine may be considered for HCWs with chronic liver or immunocompromising conditions.
B	HBeAg ⁻ source: 3% HBeAg ⁺ source: 22%–31%	HBV vaccination	HCWs with previous response to vaccine need no additional postexposure treatment regardless of level of exposure and time since vaccination. HCWs with no history of vaccination or nonresponders should receive HBIG and vaccine (for those with no history) as soon as possible after the exposure. HCWs with unknown response after primary series should have anti-HBs titers checked. <ul style="list-style-type: none"> • Levels <10 mIU/mL: Give HBIG × 1 and revaccinate • No further treatment for immune titers If anti-HBs titers cannot be readily obtained, first dose of HBIG should be administered. The role of antivirals is not defined.
C	3%	Immune globulin not recommended; see Chapter 154 for treatment	No role for direct-acting antivirals. Immediate baseline testing for hepatitis C antibody, then HCV RNA if antibody positive. If source patient confirmed with HCV infection, or with unknown status, repeat HCV RNA three weeks after exposure.
Delta	Unknown; outbreaks described only in dialysis units	HBV vaccination	Segregate HBsAg ⁺ dialysis patients by delta antibody status.
E	Unknown	Standard precautions	

^aRate of transmission from outbreak or needlestick exposure.

HBeAg, hepatitis B e antigen; HBIG, hepatitis B immune globulin; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HCV, hepatitis C virus; HCWs, health care workers.

diversion-related outbreaks have been recognized in the United States since 2009,¹⁶ including one from a traveling medical technician that spread across eight states, resulting in over 11,000 exposed individuals with 45 confirmed infections. This outbreak led to a criminal conviction of the source HCW.²⁵

Fecal-Oral Transmission Hepatitis A

HCWs are not considered to be at a higher occupational risk for hepatitis A, because nosocomial outbreaks and transmission from patient to care provider have been infrequent. The Advisory Committee on Immunization Practices (ACIP) does not recommend routine vaccination of HCWs to protect against hepatitis A.²⁶ The rare reports of nosocomial transmission have largely occurred in pediatric units and nurseries.^{27,28} The preponderance of hepatitis A outbreak reports in neonatal settings is due to the asymptomatic nature of infection in infants and young children

and prolonged shedding²⁹; the disease in index cases often goes unrecognized until secondary cases in adult contacts have occurred.³⁰ Rare outbreaks in adults have been related to high physical dependency or diarrhea or fecal incontinence.^{31–35}

Although dominantly an enterically transmitted infection, hepatitis A can rarely be acquired via blood and platelet transfusion from donors with incubating infection³⁶; in addition, a single transplant-associated case has also been described. Tertiary transmission of hepatitis A from a child who had initially contracted hepatitis A via multi-visceral organ transplant eventually led to infection in 2 of 42 home health nurses who provided direct care to the child.^{37,38}

Postexposure prophylaxis (PEP) by administration of intramuscular immune globulin to contacts has been used effectively for many years (Table 303.1).³⁹ Although not specifically assessed in health care-related exposures, vaccination within 2 weeks has been shown to be 80% to 90% effective in preventing clinical hepatitis A among immunocompetent

TABLE 303.2 Country-Specific Rates of Hepatitis E RNA Prevalence Among Screened Blood Products and Associated Risk of Transmission

COUNTRY (YEAR)	UNIT SCREENED	SAMPLES SCREENED	HEV RNA PREVALENCE	POSTTRANSFUSION HEV IN RECIPIENT	COMMENT
Netherlands (2013)	Mini-pools	59,474	0.076%	NA	Estimated duration of viremia in donors = 68 days
Germany (2012)	Mini-pools	16,125	0.08%	NA	
Spain (2015)	Individual	9998	0.03%	NA	
Denmark (2016)	Individual	25,637	0.04%	7/11 recipients without detectable viremia	Low copy numbers in donors (13 IU/mL)
Ireland (2016)	Individual	24,985	0.02%	NA	Universal screening implemented
France (2014)	Mini-pools	53,234	0.045%	NA	
United Kingdom (2014)	Mini-pools	225,000	0.035%	42% attack rate	Universal screening implemented
Australia (2017)	Pools	74,131	0.001%		
United States (2013)	Individual	1930	0%	No transmission	
United States (2016)	Individual	18,829	0.01%	NA	Low copy numbers in positive donors
United States (2017)	Mini-pools	128,028	0.002%	NA	Type 3a infections with low copy numbers

HEV, Hepatitis E virus; NA, not applicable.

adults younger than 40 years.⁴⁰ Vaccine was successfully used in two eligible nurses in the transplant recipient–related outbreak.³⁷ Since 2000, Canadian guidelines have routinely recommended vaccine as the preferred PEP for nonimmunocompromised patients older than 1 year. In United Kingdom, immune globulin is recommended to be given in conjunction with vaccine only for healthy adults older than 50 years, and those with chronic liver disease or an immunocompromising condition, regardless of age.⁴¹

Hepatitis E

First identified as a cause of acute hepatitis in Kashmir, India, hepatitis E virus (HEV) is primarily transmitted via contaminated water. The virus has caused several large-scale outbreaks among refugees and other displaced populations. Person-to-person spread among household contacts has been demonstrated only in resource-constrained settings, where transmission is attributed to poor sanitation and to communal hand-washing and eating practices.^{42–44}

In Europe, indigenous infection from zoonotic contact or acquisition via meat products (undercooked pork, pork liver sausage, game meat, and shellfish) is on the rise, with predominantly genotype 3a infections.⁴⁵ Currently in Europe, infections due to HEV are at least as common as hepatitis A among patients hospitalized with acute hepatitis.⁴⁶ Approximately half of all HEV cases are autochthonous in origin^{45–47} because risk of acquisition is strongly influenced by regional farming and dietary practices. Certain areas in eastern Japan, southwestern France, and the Netherlands have exceedingly high seropositivity rates as a result of these factors. A detailed description of the epidemiology of hepatitis E can be found in Chapter 178.

Seroprevalence of Hepatitis E in Developed Countries and Risk of Bloodborne Transmission

Seroprevalence of hepatitis E in the United States is estimated to be around 6.25%, with the highest rates in the Midwest.⁴⁸ Genotypes 3 and 4 prevail among the sporadic locally acquired infections. Based on population estimates from the National Health and Nutrition Examination Survey (NHANES), seroprevalence rates in US adults older than 30 years declined from 20% (1988–1994) to 8.4% (2009–2010); a decrease in seropositivity was seen among all age groups within this time period. Overall, anti-HEV immunoglobulin G (IgG) prevalence decreased from 9.6% to 5.2%.^{49,50}

Several assessments of blood donors in the United States have concluded that the transmission risk of HEV is minimal.^{48,51,52} Screening of blood products for HEV is not currently recommended in the United States. Blood donor screening data from several European countries have also demonstrated variable rates of hepatitis E viremia, ranging

from 0.02% to 0.08% with an associated transmission risk of 0.04% (Table 303.2).^{53–56} Some European countries have adopted universal screening of blood products for hepatitis E. A detailed discussion on this topic can be found in Chapter 304.

Other Nosocomial Transmission

Non-transfusion-related nosocomial HEV transmission is rare; an outbreak in Pakistan attributed to sharing of parenteral equipment affected 18 individuals, with 7 confirmed cases.⁵⁷ A single case of patient-to-patient spread of hepatitis E via contaminated hospital environment was suspected in a French hospital in which the donor patient with detectable virus in blood and stool had an overlapping stay with the case patient, who developed acute infection 3 weeks later. The strains were closely related by genotypic analysis.⁵⁸

Hepatitis B

Epidemiology

Hepatitis B was the first bloodborne disease recognized to pose an occupational hazard.^{8–10} In the prevaccine era, the prevalence of hepatitis B virus (HBV) among HCWs was 10-fold greater than in the general population. An early review found a preponderance of cases among pathologists, laboratory workers, and blood bank workers.¹⁰ Later studies added dentists, physicians, nurses, and laboratory and dialysis staff to the list of HCWs with increased risk.^{9,10,59–62}

Vaccine to prevent HBV infection became available in the United States in 1982. Introduction of the currently used recombinant HBV vaccine improved vaccine acceptance. Since 1991, it has been recommended as a universal childhood vaccination. In 1987, concern about occupational acquisition of human immunodeficiency virus (HIV) led the US Department of Labor, in conjunction with the Department of Health and Human Services, to recommend universal precautions to protect against exposure to body fluids.⁶³ Four years later, the Occupational Safety and Health Administration published the federal Bloodborne Pathogens Standard, which went into effect in early 1992.⁶⁴ This document mandated that HCWs with potential exposure to blood or other potentially infectious materials should be offered the hepatitis B vaccine series free of charge, should demonstrate immunity to hepatitis B, or should formally decline vaccination. In order to identify nonresponders and to intervene early, postvaccination serologic testing is recommended for HCWs prone to blood and body fluid exposures. Compliance with these recommendations has resulted in a 98% reduction in occupationally acquired hepatitis B,^{65,66} although rare cases continue to occur. Based on a survey conducted by the CDC between 2005 and 2010, 203 cases of hepatitis B were reported among HCWs; among these, only 40% were possibly related to occupational exposures. Complete vaccination

with three or more doses could be confirmed for 35 HCWs infected with hepatitis B, 1 had immune titers, and others were nonresponders or without additional information.⁶⁶

Incidence After Exposure

The risk of transmission from a single needlestick exposure varies according to the hepatitis B e antigen (HBeAg) status of the source case. The risk of developing clinical hepatitis ranges from 1% to 6% for HBeAg-negative blood to 22% to 31% for HBeAg-positive blood (see Table 303.1).^{66,67} In addition, transmission from a viremic hepatitis B surface antigen (HBsAg)-negative patient, even with low viral burden, may occur.⁶⁸

Reported Transmissions

Worker-to-patient transmission. Over the past several decades, dozens of episodes involving HCW-to-patient transmission of hepatitis B have been described, resulting in hundreds of secondary cases (range, 1–55 secondary cases per source case).^{11,12,66,68–73} In a series of 10 clusters reported from the United Kingdom, the transmission rate ranged from 0.3% to 9%.^{12,69} At least 42 of the 47 HCWs involved were dentists or surgeons. In a majority of the cases, no obvious infection-control breaches or overt percutaneous exposures were recognized. Lack of glove use during oral surgery and chronic skin conditions on the surgeon's exposed hands have been proposed as contributing factors in some of the older reports. No cases of dentist-to-patient transmission have been reported since 1987, which suggests the effectiveness of standard precautions and routine vaccination of HCWs,⁶⁹ although lack of case clustering may prevent recognition of sporadic transmission events.⁷³

In the United States, the last documented HBV transmission from a surgeon to patients occurred in 2009.⁷⁴ The source was an orthopedic surgeon with previously unrecognized infection and an HBV DNA level of almost 18 million IU/mL. Of note, he had twice received a three-dose HBV vaccination series but had not been evaluated for chronic HBV infection, despite failure to respond to the vaccine and recent immigration from a country with high rates of HBV. The last reported transmission during a surgical procedure occurred in Japan from an HBeAg-positive gynecologic surgeon with high level of virus in serum, sweat, and saliva.⁷⁵ In a classic outbreak from the 1990s,¹¹ 19 (13%) of 144 susceptible patients operated on by an HBeAg-positive thoracic surgery resident developed acute hepatitis B infection despite appropriate infection-control measures. Thirteen available isolates, including the surgeon's, were identical when compared by means of genetic analysis. Examination of the resident's surgical technique suggested that small cuts in his fingers, sustained while he tied sutures, resulted in entry of his blood into patients' wounds. Of note, the surgeon had declined hepatitis B vaccine 2 years earlier and then had become infected as the result of an occupational exposure.

In most but not all of the reported outbreaks, the source worker was HBeAg positive. However, in one series, four HBeAg-negative surgeons all transmitted disease.¹² The cases occurred in England, where restriction of HBeAg-positive surgeons (principally cardiothoracic, gynecologic, orthopedic, and abdominal) is strictly enforced. Confirmation of transmission was aided by HBV DNA sequencing of both the putative source and secondary cases.

In response to concerns around HCW-to-patient transmission, in 1991, the CDC promulgated recommendations for preventing transmission of HIV and HBV to patients during exposure-prone invasive procedures, which were modified by non-CDC experts in 2012 and are currently in revision.^{66,68,76} These and other guidelines addressing infected HCWs have attempted to balance patient safety against a worker's right to privacy. Most US and European guidelines now recommend that workers performing exposure-prone procedures maintain a low HBV DNA level, usually less than 1000 IU/mL.⁷⁶

Patient-to-worker transmission. Widespread transmission from a single patient to several HCWs is rare. In one instance, a patient in the preclinical window period for hepatitis B sustained severe trauma and underwent several operations.⁷⁷ At least four HCWs, including nurses and physicians, developed acute hepatitis temporally consistent with transmission from this patient. As with worker-to-patient transmission, clustered cases lead to the identification of transmission more

readily than individual transmissions. Other instances of transmission may go undetected and unreported.

Dialysis setting. For many years, dialysis patients and staff were at high risk for occupational acquisition of hepatitis B, given their high frequency of sharps injury or mucocutaneous exposure (5 instances per 10,000 dialysis procedures⁷⁸), the high titers of HBV in blood ($\geq 10^9$ /mL), and the ability of HBV to survive well in the environment.^{16,79} However, with segregation of patients by room, staff, and machine according to surface antigen (HBsAg) status; institution of active vaccination programs; monthly serologic testing of susceptible patients; and attention to disinfection, equipment, and cleaning procedures, this rate has decreased sharply.^{16,59,79–81} When rates were compared from a classic study conducted before the availability of vaccine,^{59,81} the incidence of new HBsAg among patients was found to have decreased from 3% to 0.05%, and the prevalence from 7.8% to 0.9%. During the same period, vaccination coverage of staff increased to 90% and HCW HBV incidence decreased from 2.6% to less than 0.5%. In US dialysis centers in 2002, the prevalence and incidence of HBV among dialysis patients had remained steady at 1% and 0.12%, respectively; 56% of patients and 90% of staff had been vaccinated.⁵⁹

Despite these gains, transmission continues in centers that fail to identify HBV-infected patients, share staff and equipment, or fail to vaccinate susceptible patients.^{16,79} Medication preparation close to treatment areas, infection-control lapses during vascular access, and inability to separate clean and contaminated environments continue to provide reservoirs of health care–based transmission of HBV.^{16,59,79} Nonimmune persons undergoing hemodialysis should be vaccinated with high-dose or double-dose vaccine formulations.⁸⁰ Serologic assessment of postvaccination titers 1 to 2 months after completion of the series is recommended. The determination for booster doses is made on an annual basis, after anti-HBs level assessment, and with the goal of maintaining titers over 10 mIU/mL. Clinically significant hepatitis has been reported in patients on dialysis who achieve a minimum protective antibody level but with subsequent decline.⁸² The approach for nonresponders is similar to others (refer to section “Postexposure Management”).

Other nosocomial transmissions. In some countries, nosocomial transmission due to unsafe practices continues to account for a substantial proportion of overall hepatitis B cases.^{83,84} In US and UK health care settings, transmission related to blood glucose monitoring devices,^{20,85} endoscopy equipment, reuse of single-dose vials in podiatry care, and jet injections has been increasingly reported,⁴ especially in long-term care.^{15,16} This group of exposures now account for up to 37% of all health care–associated acquisition of HBV among older adults in the United States.⁸⁶ In addition, patients (so-called medical tourists) who for financial or other reasons receive care in countries with a high prevalence of HBV may unknowingly invite the additional risk of virus acquisition.⁸⁸ Hepatitis B was transmitted to a patient operated on by the same oral surgeon and assistants who had performed tooth extraction in an HBeAg-positive patient 2 hours before. Thorough review of the case failed to disclose any overt infection-control breach.¹⁴

Vaccine Acceptance and Response Among Health Care Workers, and Duration of Immunity

The National Health Interview Survey (NHIS) estimated vaccination coverage of 63.8% for health care personnel in 2011.⁸⁹ Three out of every four HCWs with direct patient contact are vaccinated against hepatitis B. Approximately 88% of individuals achieve seroprotective titers after vaccination; suboptimal responses are related to older age, obesity, and smoking.⁹⁰ Among those who do not develop protective titers after the primary series (<10 mIU/mL), cumulative response on revaccination with additional three doses approaches 70%.⁶⁶ Preemployment serologic testing and HBsAg testing are recommended for HCWs from geographic regions with prevalence rates of $\geq 2\%$ regardless of vaccination status, and for those with known nonresponse to vaccine previously.⁶⁶

Knowledge of serostatus aids in decisions to revaccinate and postexposure management. Therefore, serologic testing is routinely recommended for those with an anticipated risk of occupational exposure.

*References 15, 16, 20, 21, 86, 87.

If protective levels are achieved 1 to 2 months after immunization, no further testing is necessary regardless of future exposures. For vaccinated HCWs with anti-HBs <10 mIU/mL immediately after completion, an additional dose or revaccination with the three-dose series is recommended. Any additional immunization beyond the sixth dose is not recommended for those without a response to two complete series. These individuals should be considered susceptible in the event of a significant exposure, and hepatitis B immune globulin (HBIG) administered promptly.^{91,92}

It is well recognized that antibody response wanes after completion of primary immunization series, regardless of initial response.^{93,94} The most precipitous drop in titers occurs in the years immediately after vaccination. In longitudinal reports spanning across at least 2 decades, persons with declining antibody level (to <10 mIU/mL) remain protected.^{95,96} Age at the time of vaccination and higher anti-HBs titers after vaccination correlate with duration of measurable immune titers.⁹³ In the longest cohort of vaccinees, including children and adults from 12 Alaskan communities, followed over a period of 30 years after primary hepatitis B vaccination series, approximately half of the individuals showed loss of measured protective antibody. Despite this, anamnestic response after booster challenge could be elicited in the vast majority (89%), confirming a protective response on exposure to HBV.⁹⁵ For the vaccinees without demonstrable anamnestic response, future susceptibility to acute or chronic HBV disease remains unclear.

Other studies conducted in areas of high endemicity have demonstrated breakthrough infection among vaccine responders marked by the development of antibodies against hepatitis B core protein.^{93,97} HCWs vaccinated as adults also demonstrate durable antibody response for up to 28 years after primary immunization.⁹⁸ Loss of protective titers was detected in 22% and did not correlate with interval since vaccination. Among those given a booster dose, 94% developed adequate levels. Immunized HCWs with occupational exposure to HBV have been shown to develop core and polymerase-specific CD8 (+) T-cell responses akin to acquired immunity after HBV infection.⁹⁹ These data collectively suggest that newly infected vaccinees typically develop subclinical disease and that vaccine-induced immunity confers protection from chronic HBsAg carriage and clinical hepatitis, but may not entirely prevent infection. Based on this, the CDC does not recommend routine revaccination of otherwise healthy HCWs.⁶⁶

In November 2017 a recombinant HPV vaccine with a novel CpG adjuvant was marketed for persons at least 18 years of age and is given as two doses 1 month apart. Immunogenicity is reported at least as good as the earlier two recombinant HBV vaccines. The vaccine, Heplisav-B, was recommended as an option in 2018 by the ACIP and may offer convenience for HCWs over the prior three-dose 16 weeks schedule.^{99a}

Postexposure Management

Management in exposed or susceptible workers has been well summarized elsewhere^{26,66} (see Table 303.1 and Chapter 145). Treatment should ideally be administered within 24 hours after exposure. Intramuscular HBIG was the original intervention for PEP.^{66,100} It is still used, in conjunction with initiation of a vaccine series, for management after exposure in unvaccinated persons and vaccine nonresponders. For vaccine responders, no postexposure intervention is required, regardless of the HBsAg or HBeAg status of the source. Vaccinated persons with an unknown response to vaccine should have their antibody titer checked immediately after exposure and should be treated according to the result (see Table 303.1). If HBIG and vaccine are indicated, titers should be checked at least 3 months after HBIG after antibody levels from passive immunization have waned, in order to accurately assess response to the vaccine. Nonresponders to hepatitis B vaccine should receive a second dose of HBIG 1 month later (HBIG dose, 0.06 mL/kg or 5.0 mL for adults).^{66,100} Vaccines and HBIG may be given at the same time but should be administered with separate needles and syringes and at different anatomic sites. Plain immune globulin does not contain sufficient titers of HBIG and should not be considered an acceptable alternative. The role of antiviral agents such as lamivudine, entecavir, and tenofovir in the management of nosocomial exposure has not been determined.

The best long-term management for the hepatitis B vaccine nonresponder is not known. All nonresponders to the first three-dose series should be tested for evidence of HBsAg if not already done at the time of hire.

Hepatitis C Seroprevalence

Most^{78,101} series suggest that many HCWs with increased risk for hepatitis B, including dialysis workers, laboratory workers, surgeons, nurses, and workers with the mentally impaired, have no increase in hepatitis C virus (HCV) seroprevalence. The overall prevalence of HCV among persons receiving dialysis is approximately 8% (nearly fivefold higher than the general population) and among dialysis center staff is around 1.7% (same as US general population).⁵⁹ Some studies, including a recent meta-analysis, have suggested that medical and laboratory staff and dental health workers may have an elevated risk.^{102,103}

Incidence After Occupational Exposure

Percutaneous exposure is the most common mode of health care–associated transmission. Risk is greatest with deep injuries from a hollow-bore needle that has been used to access the source patient's artery or vein.¹⁰⁴ Rare cases of acquisition from blood splash on the conjunctiva have been reported. HCV after exposure to body fluids other than blood or via contaminated environment has not been conclusively documented. Seroconversion occurs in 0% to 10% of nonimmune HCWs who sustain needlesticks from a source patient with hepatitis C.^{26,78,105–108} Maternal-fetal transmission rates are similar (5%–9%). Rates may vary because of differences in the diagnostic test used (antibody or HCV RNA) and nature of evaluated exposures. The highest transmission rate (10%) was from a study in which HCV RNA was used to detect infection in exposed workers.^{106,108} This high rate has not been duplicated, and most studies have placed the transmission rate at less than 3%. In a large case series from the United Kingdom, 626 percutaneous HCV exposures to blood or bloodstained fluids occurred from 1997 to 2007, resulting in a seroconversion rate of 2.2%.¹⁰⁴ Existing evidence supports a correlation between HCV RNA level in the source patient's blood and higher risk of transmission.¹⁰⁹ This association has not been established for other body fluids. A report of simultaneous transmission of HCV and HIV from a single needlestick was remarkable for the delayed time to seroconversion against each virus and the fulminant, fatal course of the HCV infection.¹¹⁰ This phenomenon appears to be extremely rare.

Reported Transmissions

Worker-to-patient transmission. HCWs have transmitted HCV to patients on several occasions, sometimes after acquiring their own infection from a different patient.^{111–113} A cardiac surgeon transmitted HCV to at least 5 of 222 patients; all infected patients underwent valve replacement.¹³ Molecular analysis showed significant homology between the surgeon's and the patients' viruses. The surgeon was treated until his HCV RNA level became undetectable. At that point, he was allowed to resume performing surgery.

Additional isolated cases of HCV transmission during thoracic, cardiac, and orthopedic surgery have been reported.¹¹⁴ In the UK experience, look-back investigations covering exposure-prone procedures from an HCV-infected source have yielded a transmission rate of 0.13% among susceptible patients. In Germany, an anesthesiologist at an outpatient gynecology clinic transmitted HCV to three patients during minor procedures on the same day; the precise cause of this cluster could not be determined. Additional testing did not reveal transmission events originating from the anesthesiologist.¹¹⁵

HCV infection in two Scottish tourists was traced back to an HCW; the two patients had received hemodialysis during a holiday in Spain.¹¹⁶ Most other transmissions involving nonsurgical personnel have occurred from drug diversion practices through cross-contamination of equipment and supplies by an HCV-infected HCW.¹¹⁷ In the last decade, three separate outbreaks in the United States, two from different radiology technicians and one from a nurse anesthetist, led to spread of HCV infection in 68 individuals.^{24,25,118} Ensuring secure access, tracking mechanisms, and safe administration of narcotics are

essential to eliminate HCV and other infection risks associated with drug diversion.

Transmission in dialysis setting. Despite improvements, transmission of HCV remains a substantial problem for dialysis patients throughout the world. In most series, HCV seroprevalence among dialysis patients is 2- to 10-fold higher than in the general population. The CDC reported regularly on dialysis-associated diseases in the United States through 2002. In their last survey, which included 96% of all US centers, representing more than 260,000 patients and 58,000 staff members, 63% of the institutions routinely tested patients for HCV, an increase from 39% in 1995.⁵⁹ HCV seroprevalence is even higher in South American, African, and Asian dialysis centers, exceeding 40% in some areas. Prevalence estimates have ranged from 5% to 10% in Europe and the United States.^{59,119} The prevalence of HCV among peritoneal dialysis patients, in contrast, is not elevated.¹²⁰

Because of their compromised immune response, dialysis patients may not mount a significant antibody response to HCV; therefore, concurrent testing with HCV RNA is recommended. Studies done among hemodialysis patients from Germany and France with an overall HCV prevalence of 7% found that 0.6% to 12.3% of HCV-infected patients were viremic without detectable antibody with third-generation assays.^{121,122} Cost remains a prohibiting factor for the routine use of HCV RNA in resource-limited settings. Select hepatitis C core antigen assays have demonstrated comparable diagnostic performance to nucleic acid amplification testing (NAAT) at HCV RNA levels >3000 IU/mL and may become an affordable option for broader screening in these areas.¹²³ The current recommendation by the CDC, endorsed by the National Kidney Foundation, is monthly measurement of alanine aminotransferase (ALT) levels combined with semiannual anti-HCV testing for susceptible dialysis patients as a more cost-effective approach.

Early reports demonstrated that nosocomial spread was caused by overt interruptions in infection control,¹²⁴ and one more recent report suggested that HCV was transmitted by contaminated hands of HCWs, demonstrating that the problem persists.^{125,126} In most newer reports, however, obvious breaches of infection control in dialysis centers and elsewhere have not been identified, suggesting that either subtle interruptions are responsible for spread or that HCV transmission is incompletely understood.¹⁶ Supporting the latter interpretation, some studies using genotypic analysis have not demonstrated homology among persons treated in the same or adjacent beds; rather, linked cases have been located throughout the dialysis center, suggesting widely dispersed transmission by means of an uncertain mechanism.¹²⁷

Recognized risk factors for acquisition of HCV include blood transfusion and the number of months for which a patient has required dialysis,¹²⁸ with the latter being more significant. The risk of blood transfusion has been reduced by the use of erythropoietin. Possible explanations for the association of acquisition of HCV with duration of dialysis include sharing of dialysis machines by HCV-infected and uninfected patients and reprocessing of dialyzers from HCV-infected patients.^{129,130} However, the current cleaning standards are extremely high, and the CDC therefore does not recommend against sharing machines or against reusing dialysis filters.⁵⁹

Some researchers have advocated keeping HCV-infected dialysis patients together, similar to the successful approach taken with HBV-infected patients. The CDC, however, does not advocate this approach because of the lack of sensitivity of the anti-HCV test, which means that not all infectious persons would be isolated, and because of the risk of superinfection or reinfection for those already infected.⁵⁹ Refer to Chapter 154 for further discussion.

Solid tumor transplantation. Transplantation of HCV-infected organs into HCV-infected or HCV-uninfected hosts is sometimes necessary.¹³¹ However, almost all susceptible recipients of HCV-positive organs eventually develop HCV infection,¹³² which may be severe.^{132,133} In most series, however, no adverse effect on overall survival of patients or grafts has been found, although recipients of HCV-positive organs had a higher rate of liver-related morbidity and mortality.^{132,133} Many organ banks now avoid using organs from HCV-positive donors except for lifesaving procedures such as heart, lung, and liver transplantation.¹³¹ Transmission of HCV to eight organ or tissue recipients from a single, seronegative, and NAAT-positive organ donor has been described.¹³⁴

The donor died early in the HCV “window period.” Introduction of tissue donor NAAT screening in conjunction with serologic testing has reduced the likelihood of undetected hepatitis C transmission; despite this, human error in interpreting results led to transmission in two kidney and a cardiopulmonary patch recipient from an antibody-negative, NAAT-positive donor.¹³⁵ In other transplant-related incidents, potential transmission via blood vessel conduit from a seropositive donor and through common use of perfusion machines has also been reported.¹³⁶ The use of active antivirals for organ recipients in these settings has not been studied.

Other nosocomial transmissions. As previously noted, in recent years more reports have described clustering of hepatitis C cases around various health care–related exposures, including transmissions related to endoscopy^{137,138}; myocardial perfusion¹³⁹; computed tomography scanning¹⁴⁰; reuse of saline flushes,¹⁴¹ needles and syringes,^{16,142} and multidose vials^{16,22,86,87,143}; and use of a spring-loaded fingerstick device.¹⁴⁴

In a highly publicized report from Las Vegas, several patients seen at one endoscopy clinic were found to have hepatitis C.¹³⁷ The infection was thought to be related to inappropriate reuse of syringes and the use in multiple persons of medication vials intended only for single use. In other reports from endoscopy clinics, the transmission may have occurred as a result of inadequate cleaning of the biopsy suction channel of the colonoscope or failure to autoclave some equipment, such as biopsy forceps.^{22,145} Hepatitis C transmission during a uterine dilation procedure was genotypically determined to have originated from a source patient who had undergone an operation earlier on the same day. The only common link was an anesthesiologist who prepared medications on a cart that was rolled between operating rooms throughout the day without cleaning and disinfection in between.¹⁴⁶

Transmission of hepatitis C between patients in a dental office has been documented.¹⁴⁷ The mechanism of transmission is unclear but might involve improperly sterilized instruments.

Several reports have suggested an association between development of HCV infection and hospitalization on an oncology or bone marrow transplant floor,^{18,148,149} or on a urology,¹⁵⁰ liver,¹⁵¹ or general ward.¹⁵² Although some studies are from relatively resource-poor countries, where reuse of needles may be the cause, the source of spread in other centers remains puzzling. Prolonged hospitalization is often a risk in these reports, and, in one large US outbreak, transmission was thought to be related to shared saline bags contaminated through syringe reuse.¹⁴⁸ Persistence of live HCV for several weeks on dried surfaces has been shown with use of a genetically engineered reporter strain.¹⁵³ Whether contact with contaminated surfaces can transmit hepatitis C is plausible but unproven.

Management

The CDC recommends determining the HCV serostatus of the source patient after any exposure.¹⁵⁴ For the exposed HCW, immediate anti-HCV testing within 48 hours is recommended. HCV RNA testing should be done if the initial testing result is positive. For those who are anti-HCV negative, an assessment for HCV RNA should be done at least 3 weeks after exposure to a source patient with HCV or unknown status. HCWs who test positive for HCV RNA at symptom onset or at postexposure testing at ≥3 weeks should be retested at 6 months for spontaneous clearance of infection.¹⁵⁴

Optimal management of a needlestick exposure is unknown, but each exposure need not be managed with antivirals. For the small subset of HCWs who seroconvert and remain with detectable RNA, newer direct-acting antivirals (DAAs) are highly efficacious at achieving a sustained virologic response. The role of DAAs has not been evaluated in the context of PEP, and they are not currently recommended for this purpose (see Chapter 154). Interferon- α (IFN- α) remains the only drug that has been studied in early infection; in a single report, HCV resolved in all 14 HCWs who were treated with IFN soon after acute infection.¹⁵⁵

Hepatitis D

Delta virus is a defective RNA virus that requires the presence of active HBV infection (acute HBV or HBsAg carrier state) to infect the liver. In a report from the 1980s,⁷² an HBV- and delta-infected source patient

regularly shared a dialysis machine with an asymptomatic HBsAg carrier. The latter patient subsequently developed acute delta hepatitis. A surgeon may also have become dually infected after a deep needlestick sustained while operating on the same source patient. Review identified several additional possible instances of delta hepatitis transmission in dialysis

centers. This review led to the current recommendation that patients and staff be vaccinated against HBV and that dialysis patients be separated according to delta virus status. An additional benefit of hepatitis B vaccination is that the incidence of hepatitis D has also substantially declined. Refer to Chapter 146.

Key References

The complete reference list is available online at Expert Consult.

12. Incident Investigation Teams, et al. Transmission of hepatitis B to patients from four infected surgeons without hepatitis B e antigen. *N Engl J Med*. 1997;336:178–184.
13. Esteban JI, Gomez J, Martell M, et al. Transmission of hepatitis C virus by a cardiac surgeon. *N Engl J Med*. 1996;334:555–560.
15. Thompson ND, Perz JF, Moorman AC, et al. Nonhospital health care-associated hepatitis B and C virus transmission: United States, 1998–2008. *Ann Intern Med*. 2009;150:33–39.
16. Healthcare-Associated Hepatitis B and C Outbreaks (≥ 2 cases) Reported to the Centers for Disease Control and Prevention (CDC) 2008–2016. <https://www.cdc.gov/hepatitis/outbreaks/pdfs/healthcareinvestigationtablepdf.pdf>. Accessed November 2017.
19. Perz JF, Thompson ND, Schaefer MK, et al. US outbreak investigations highlight the need for safe injection practices and basic infection control. *Clin Liver Dis*. 2010;14:137–151, x.
20. Centers for Disease Control, Prevention. Multiple outbreaks of hepatitis B virus infection related to assisted monitoring of blood glucose among residents of assisted living facilities—Virginia, 2009–2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:339–343.
22. Guh AY, Thompson ND, Schaefer MK, et al. Patient notification for bloodborne pathogen testing due to unsafe injection practices in the US health care settings, 2001–2011. *Med Care*. 2012;50:785–791.
24. Hellinger WC, Bacalis LP, Kay RS, et al. Health care-associated hepatitis C virus infections attributed to narcotic diversion. *Ann Intern Med*. 2012;156:477–482.
25. State of New Hampshire. Hepatitis C Outbreak Investigation, Exeter Hospital, Public Report, June 2013; 2013. <http://www.dhhs.nh.gov/dphs/cdcs/hepatitis/c/documents/hepc-outbreak-rpt.pdf>. Accessed January 15, 2018.
26. Immunizations of healthcare personnel. *MMWR Morb Mortal Wkly Rep*. 2011;60(RR07):1–45.
37. Wiseman R, Weil LM, Lozano C, et al. Notes from the field: health care-associated hepatitis A outbreak - Texas, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65:425–426.
42. Teshale EH, Grytdal SP, Howard C, et al. Evidence of person-to-person transmission of hepatitis E virus during a large outbreak in Northern Uganda. *Clin Infect Dis*. 2010;50:1006–1010.
48. Stramer SL, Moritz ED, Foster GA, et al. Hepatitis E virus: seroprevalence and frequency of viral RNA detection among US blood donors. *Transfusion*. 2016;56:481–488.
50. Teshale EH, Denniston MM, Drobeniuc J, et al. Decline in hepatitis E virus antibody prevalence in the United States from 1988–1994 to 2009–2010. *J Infect Dis*. 2015;211:366–373.
51. Xu C, Wang RY, Schechter CA, et al. An assessment of hepatitis E virus (HEV) in US blood donors and recipients: no detectable HEV RNA in 1939 donors tested and no evidence for HEV transmission to 362 prospectively followed recipients. *Transfusion*. 2013;53(10 Pt 2):2505–2511.
53. Hewitt PE, Ijaz S, Brailsford SR, et al. Hepatitis E virus in blood components: a prevalence and transmission study in southeast England. *Lancet*. 2014;384:1766–1773.
59. Finelli L, Miller JT, Tokars JI, et al. National surveillance of dialysis-associated diseases in the United States, 2002. *Semin Dial*. 2005;18:52–61.
62. Mast EE, Alter MJ. Prevention of hepatitis B virus infection among health-care workers. In: Ellis RE, ed. *Hepatitis B Vaccines in Clinical Practice*. New York: Marcel Dekker; 1993:295–307.
64. Occupational exposure to bloodborne pathogens—OSHA. Final rule. *Fed Regist*. 1991;56:64004–64182.
65. Agerton TB, Mahoney FJ, Polish LB, et al. Impact of the bloodborne pathogens standard on vaccination of healthcare workers with hepatitis B vaccine. *Infect Control Hosp Epidemiol*. 1995;16:287–291.
66. Schillie S, Murphy TV, Sawyer M, et al. CDC guidance for evaluating health-care personnel for hepatitis B virus protection and for administering postexposure management. *MMWR Recomm Rep*. 2013;62(RR-10):1–19.
67. Werner BG, Grady GE. Accidental hepatitis B-surface-antigen-positive inoculations. Use of e antigen to estimate infectivity. *Ann Intern Med*. 1982;97:367–369.
68. Centers for Disease Control, Prevention. Updated CDC recommendations for the management of hepatitis B virus-infected health-care providers and students. *MMWR Recomm Rep*. 2012;61(RR-3):1–12.
74. Enfield KB, Sharapov U, Hall KK, et al. Transmission of hepatitis B virus from an orthopedic surgeon with a high viral load. *Clin Infect Dis*. 2013;56:218–224.
76. Henderson DK, Dembry L, Fishman NO, et al. SHEA Guideline for management of healthcare workers who are infected with hepatitis B virus, hepatitis C virus and/or human immunodeficiency virus. *Infect Control Hosp Epidemiol*. 2010;31:203–232.
77. Shanson DC. Hepatitis B outbreak in operating-theatre and intensive care staff. *Lancet*. 1980;2:596.
78. Petrosillo N, Puro V, Jagger J, et al. The risks of occupational exposure and infection by human immunodeficiency virus, hepatitis B virus, and hepatitis C virus in the dialysis setting. Italian Multicenter Study on Nosocomial and Occupational Risk of Infections in Dialysis. *Am J Infect Control*. 1995;23:278–285.
80. CDC. Summarized from Recommendations of the Advisory Committee on Immunization Practices (ACIP); 2012. https://www.cdc.gov/dialysis/pdfs/vaccinating_dialysis_patients_and_patients_dec2012pdf.pdf. Accessed January 15, 2018.
81. Alter MJ, Faverio MS, Maynard JE. Impact of infection control strategies on the incidence of dialysis-associated hepatitis in the United States. *J Infect Dis*. 1986;153:1149–1151.
82. Stevens CE, Alter HJ, Taylor PE, et al. Hepatitis B vaccine in patients receiving hemodialysis. Immunogenicity and efficacy. *N Engl J Med*. 1984;311:496–501.
92. Simard EP, Miller JT, George PA, et al. Hepatitis B vaccination coverage levels among healthcare workers in the United States, 2002–2003. *Infect Control Hosp Epidemiol*. 2007;28:783–790.
95. Bruce MG, Bruden D, Hurlburt D, et al. Antibody levels and protection after hepatitis B vaccine: results of a 30-year follow-up study and response to a booster dose. *J Infect Dis*. 2016;214:16–22.
96. Simons BC, Spradling PR, Bruden DJ, et al. A longitudinal hepatitis B vaccine cohort demonstrates long-lasting hepatitis B virus (HBV) cellular immunity despite loss of antibody against HBV surface antigen. *J Infect Dis*. 2016;214:273–280.
98. Gara N, Abdalla A, Rivera E, et al. Durability of antibody response against hepatitis B virus in healthcare workers vaccinated as adults. *Clin Infect Dis*. 2015;60:505–513.
- 99a. Schillie S, Harris A, Link-Gelles R, et al. Recommendations of the Advisory Committee on Immunization Practices for use of a hepatitis B vaccine with a novel adjuvant. *MMWR Morb Mortal Wkly Rep*. 2018;67:455–458.
101. U.S. Public Health Service. Updated U.S. public health service guidelines for the management of occupational exposures to HBV, HCV, and HIV and recommendations for postexposure prophylaxis. *MMWR Recomm Rep*. 2001;50(RR-11):1–52.
104. Tomkins SE, Eloff J, Nichols T, et al. Occupational transmission of hepatitis C in healthcare workers and factors associated with seroconversion: UK surveillance data. *J Viral Hepat*. 2012;19:199–204.
109. Yazdanpanah Y, De Carli G, Miguere B, et al. Risk factors for hepatitis C virus transmission to health care workers after occupational exposure: a European case-control study. *Clin Infect Dis*. 2005;41:1423–1430.
112. Ross RS, Viazov S, Gross T, et al. Transmission of hepatitis C virus from a patient to an anesthesiology assistant to five patients. *N Engl J Med*. 2000;343:1851–1854.
121. Hinrichsen H, Leimenstoll G, Stegen G, et al. Prevalence and risk factors of hepatitis C virus infection in haemodialysis patients: a multicentre study in 2796 patients. *Gut*. 2002;51:429–433.
125. Girou E, Chevaliez S, Challine D, et al. Determinant roles of environmental contamination and noncompliance with standard precautions in the risk of hepatitis C virus transmission in a hemodialysis unit. *Clin Infect Dis*. 2008;47:627–633.
127. Fornis X, Fernandez-Llama P, Pons M, et al. Incidence and risk factors of hepatitis C virus infection in a haemodialysis unit. *Nephrol Dial Transplant*. 1997;12:736–740.
131. Pereira BJ, Levey AS. Hepatitis C virus infection in dialysis and renal transplantation. *Kidney Int*. 1997;51:981–999.
132. Pereira BJ, Milford EL, Kirkman RL, et al. Transmission of hepatitis C virus by organ transplantation. *N Engl J Med*. 1991;325:454–460.
133. Pereira BJ, Milford EL, Kirkman RL, et al. Prevalence of hepatitis C virus RNA in organ donors positive for hepatitis C antibody and in the recipients of their organs. *N Engl J Med*. 1992;327:910–915.
137. Fischer GE, Schaefer MK, Labus BJ, et al. Hepatitis C virus infections from unsafe injection practices at an endoscopy clinic in Las Vegas, Nevada, 2007–2008. *Clin Infect Dis*. 2010;51:267–273.
138. Muscarella LF, New York State health officials. Recommendations for preventing hepatitis C virus infection: analysis of a Brooklyn endoscopy clinic's outbreak. *Infect Control Hosp Epidemiol*. 2001;22:669.
145. Bronowicki JP, Venard V, Botte C, et al. Patient-to-patient transmission of hepatitis C virus during colonoscopy. *N Engl J Med*. 1997;337:237–240.
148. Macedo de Oliveira A, White KL, Leschinsky DP, et al. An outbreak of hepatitis C virus infections among outpatients at a hematology/oncology clinic. *Ann Intern Med*. 2005;142:898–902.
149. Sepkowitz KA. Risk to cancer patients from nosocomial hepatitis C virus. *Infect Control Hosp Epidemiol*. 2004;25:599–602.
154. Centers for Disease Control and Prevention. Testing for HCV infection: an update of guidance for clinicians and laboratorians. *MMWR Morb Mortal Wkly Rep*. 2013;62:362–365.

References

- MacCullum FO, Bauer DJ. Homologous serum jaundice: transmission experiments with human volunteers. *Lancet*. 1944;5:622–627.
- Seeff LB, Beebe GW, Hoofnagle JH, et al. A serologic follow-up of the 1942 epidemic of post-vaccination hepatitis in the United States Army. *N Engl J Med*. 1987;316:965–970.
- Mendelsohn K, Wits L. Transmission of infection during withdrawal of blood. *Br Med J*. 1945;5:625–626.
- Droller H. Outbreak of Hepatitis in a Diabetic Clinic. *Br Med J*. 1945;1:623–625.
- MacCullum FO, Bradley WH. Transmission of infective hepatitis to human volunteers. *Lancet*. 1944;2:228–231.
- Neefe JR, Gellis SS, Stokes J Jr. Homologous serum hepatitis and infectious (epidemic) hepatitis; studies in volunteers bearing on immunological and other characteristics of the etiological agents. *Am J Med*. 1946;1:3–22.
- Rosenbaum JR, Sepkowitz KA. Infectious disease experimentation involving human volunteers. *Clin Infect Dis*. 2002;34:963–971.
- Leibowitz S, Greenwald L, et al. Serum hepatitis in a blood bank worker. *J Am Med Assoc*. 1949;140:1331–1333.
- Byrne EB. Viral hepatitis: an occupational hazard of medical personnel. Experience of the Yale-New Haven Hospital, 1952 to 1965. *JAMA*. 1966;195:362–364.
- Trumbull ML, Greiner DJ. Homologous serum jaundice; an occupational hazard to medical personnel. *J Am Med Assoc*. 1951;145:965–967.
- Harpaz R, Von Seidlein L, Averhoff FM, et al. Transmission of hepatitis B virus to multiple patients from a surgeon without evidence of inadequate infection control. *N Engl J Med*. 1996;334:549–554.
- Incident Investigation Teams, et al. Transmission of hepatitis B to patients from four infected surgeons without hepatitis B e antigen. *N Engl J Med*. 1997;336:178–184.
- Esteban JI, Gomez J, Martell M, et al. Transmission of hepatitis C virus by a cardiac surgeon. *N Engl J Med*. 1996;334:555–560.
- Redd JT, Baumbach J, Kohn W, et al. Patient-to-patient transmission of hepatitis B virus associated with oral surgery. *J Infect Dis*. 2007;195:1311–1314.
- Thompson ND, Perz JF, Moorman AC, et al. Nonhospital health care-associated hepatitis B and C virus transmission: United States, 1998–2008. *Ann Intern Med*. 2009;150:33–39.
- Healthcare-Associated Hepatitis B and C Outbreaks (≥ 2 cases) Reported to the Centers for Disease Control and Prevention (CDC) 2008–2016. <https://www.cdc.gov/hepatitis/outbreaks/pdfs/healthcareinvestigationtablepdf.pdf>. Accessed November 2017.
- Calles DL, Collier MG, Khudyakov Y, et al. Hepatitis C virus transmission in a skilled nursing facility, North Dakota, 2013. *Am J Infect Control*. 2017;45:126–132.
- Greeley RD, Semple S, Thompson ND, et al. Hepatitis B outbreak associated with a hematology-oncology office practice in New Jersey, 2009. *Am J Infect Control*. 2011;39:663–670.
- Perz JF, Thompson ND, Schaefer MK, et al. US outbreak investigations highlight the need for safe injection practices and basic infection control. *Clin Liver Dis*. 2010;14:137–151, x.
- Centers for Disease Control, Prevention. Multiple outbreaks of hepatitis B virus infection related to assisted monitoring of blood glucose among residents of assisted living facilities—Virginia, 2009–2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:339–343.
- Centers for Disease Control, Prevention. Notes from the field: deaths from acute hepatitis B virus infection associated with assisted blood glucose monitoring in an assisted-living facility—North Carolina, August–October 2010. *MMWR Morb Mortal Wkly Rep*. 2011;60:182.
- Guh AY, Thompson ND, Schaefer MK, et al. Patient notification for bloodborne pathogen testing due to unsafe injection practices in the US health care settings, 2001–2011. *Med Care*. 2012;50:785–791.
- Muleta D, Kainer MA, Moore-Moravian L, et al. Hepatitis C outbreak in a dialysis clinic - Tennessee, 2014. *MMWR Morb Mortal Wkly Rep*. 2016;64:1386–1387.
- Hellinger WC, Bacalis LP, Kay RS, et al. Health care-associated hepatitis C virus infections attributed to narcotic diversion. *Ann Intern Med*. 2012;156:477–482.
- State of New Hampshire. Hepatitis C Outbreak Investigation, Exeter Hospital, Public Report, June 2013; 2013. <http://www.dhhs.nh.gov/dphs/cdcs/hepatitis/documents/hepc-outbreak-rpt.pdf>. Accessed January 15, 2018.
- Immunizations of health care personnel. *MMWR Morb Mortal Wkly Rep*. 2011;60(RR07):1–45.
- Drusin LM, Sohmer M, Groshen SL, et al. Nosocomial hepatitis A infection in a paediatric intensive care unit. *Arch Dis Child*. 1987;62:690–695.
- Doebbeling BN, Li N, Wenzel RP. An outbreak of hepatitis A among health care workers: risk factors for transmission. *Am J Public Health*. 1993;83:1679–1684.
- Rosenblum LS, Villarino ME, Nainan OV, et al. Hepatitis A outbreak in a neonatal intensive care unit: risk factors for transmission and evidence of prolonged viral excretion among preterm infants. *J Infect Dis*. 1991;164:476–482.
- Klein BS, Michaels JA, Rytel MW, et al. Nosocomial hepatitis A. A multirisk outbreak in Wisconsin. *JAMA*. 1984;252:2716–2721.
- Park JY, Lee JB, Jeong SY, et al. Molecular characterization of an acute hepatitis A outbreak among healthcare workers at a Korean hospital. *J Hosp Infect*. 2007;67:175–181.
- Guet L, Desbois D, Roque-Afonso AM, et al. Investigation of a severe nosocomial outbreak of hepatitis A among healthcare workers and adult patients. *J Hosp Infect*. 2011;78:70–71.
- Bohm SR, Berger KW, Hackert PB, et al. Hepatitis A outbreak among adults with developmental disabilities in group homes—Michigan, 2013. *MMWR Morb Mortal Wkly Rep*. 2015;64:148–152.
- Baptiste R, Koziol D, Henderson DK. Nosocomial transmission of hepatitis A in adult population. *Infect Control*. 1987;8:364–370.
- Jensenius M, Ringertz SH, Berild D, et al. Prolonged nosocomial outbreak of hepatitis A arising from an alcoholic with pneumonia. *Scand J Infect Dis*. 1998;30:119–123.
- Noble RCKM, Reeves SA, et al. Posttransfusion hepatitis A in neonatal intensive care unit. *JAMA*. 1984;252:2711–2715.
- Wiseman R, Weil LM, Lozano C, et al. Notes from the field: health care-associated hepatitis A outbreak - Texas, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65:425–426.
- Foster MA, Weil LM, Jin S, et al. Transmission of hepatitis A virus through combined liver-small intestine-pancreas transplantation. *Emerg Infect Dis*. 2017;23:590–596.
- Nelson NP. Updated dosing instructions for immune globulin (human) GamaSTAN S/D for hepatitis A virus prophylaxis. *MMWR Morb Mortal Wkly Rep*. 2017;66:959–960.
- Victor JC, Monto AS, Surdina TY, et al. Hepatitis A vaccine versus immune globulin for postexposure prophylaxis. *N Engl J Med*. 2007;357:1685–1694.
- Ontario Agency for Health Protection and Promotion (Public Health Ontario) PIDAC. Hepatitis A Post-exposure Prophylaxis. Toronto, ON: Queen's Printer for Ontario; October 2013. https://www.publichealthontario.ca/en/eRepository/Hepatitis_A_Post_exposure_Prophylaxis_2013.pdf. Accessed January 15, 2018.
- Teshale EH, Grytdal SP, Howard C, et al. Evidence of person-to-person transmission of hepatitis E virus during a large outbreak in Northern Uganda. *Clin Infect Dis*. 2010;50:1006–1010.
- Teshale EH, Howard CM, Grytdal SP, et al. Hepatitis E epidemic, Uganda. *Emerg Infect Dis*. 2010;16:126–129.
- Centers for Disease Control, Prevention. Investigation of hepatitis E outbreak among refugees - Upper Nile, South Sudan, 2012–2013. *MMWR Morb Mortal Wkly Rep*. 2013;62:581–586.
- Aspinall EJ, Couturier E, Faber M, et al. Hepatitis E virus infection in Europe: surveillance and descriptive epidemiology of confirmed cases, 2005 to 2015. *Euro Surveill*. 2017;22.
- Tholen AT, Schinkel J, Molenkamp R, et al. Hepatitis E virus as a Cause of Acute Hepatitis in The Netherlands. *PLoS ONE*. 2016;11:e0146906.
- Ijaz S, Said B, Boxall E, et al. Indigenous hepatitis E in England and Wales from 2003 to 2012: evidence of an emerging novel phylogroup of viruses. *J Infect Dis*. 2014;209:1212–1218.
- Stramer SL, Moritz ED, Foster GA, et al. Hepatitis E virus: seroprevalence and frequency of viral RNA detection among US blood donors. *Transfusion*. 2016;56:481–488.
- Nelson KE, Kmush B, Labrique AB. The epidemiology of hepatitis E virus infections in developed countries and among immunocompromised patients. *Expert Rev Anti Infect Ther*. 2011;9:1133–1148.
- Teshale EH, Denniston MM, Drobeniuc J, et al. Decline in hepatitis E virus antibody prevalence in the United States from 1988–1994 to 2009–2010. *J Infect Dis*. 2015;211:366–373.
- Xu C, Wang RY, Schecterly CA, et al. An assessment of hepatitis E virus (HEV) in US blood donors and recipients: no detectable HEV RNA in 1939 donors tested and no evidence for HEV transmission to 362 prospectively followed recipients. *Transfusion*. 2013;53(10 Pt 2):2505–2511.
- Roth NJ, Schafer W, Alexander R, et al. Low hepatitis E virus RNA prevalence in a large-scale survey of United States source plasma donors. *Transfusion*. 2017;57:2958–2964.
- Hewitt PE, Ijaz S, Brailsford SR, et al. Hepatitis E virus in blood components: a prevalence and transmission study in southeast England. *Lancet*. 2014;384:1766–1773.
- Hoad VC, Seed CR, Fryk JJ, et al. Hepatitis E virus RNA in Australian blood donors: prevalence and risk assessment. *Vox Sang*. 2017;112:614–621.
- Juhl D, Baylis SA, Blumel J, et al. Seroprevalence and incidence of hepatitis E virus infection in German blood donors. *Transfusion*. 2014;54:49–56.
- Domanovic D, Tedder R, Blumel J, et al. Hepatitis E and blood donation safety in selected European countries: a shift to screening? *Euro Surveill*. 2017;22.
- Siddiqui AR, Jooma RA, Smego RA Jr. Nosocomial outbreak of hepatitis E infection in Pakistan with possible parenteral transmission. *Clin Infect Dis*. 2005;40:908–909.
- Mansuy JM, Huynh A, Abravanel F, et al. Molecular evidence of patient-to-patient transmission of hepatitis E virus in a hematology ward. *Clin Infect Dis*. 2009;48:373–374.
- Finelli L, Miller JT, Tokars JJ, et al. National surveillance of dialysis-associated diseases in the United States, 2002. *Semin Dial*. 2005;18:52–61.
- Gibas A, Blewett DR, Schoenfeld DA, et al. Prevalence and incidence of viral hepatitis in health workers in the prehepatitis B vaccination era. *Am J Epidemiol*. 1992;136:603–610.
- Lewis TL, Alter HJ, Chalmers TC, et al. A comparison of the frequency of hepatitis-B antigen and antibody in hospital and nonhospital personnel. *N Engl J Med*. 1973;289:647–651.
- Mast EE, Alter MJ. Prevention of hepatitis B virus infection among health-care workers. In: Ellis RE, ed. *Hepatitis B Vaccines in Clinical Practice*. New York: Marcel Dekker; 1993:295–307.
- Mahoney FJ, Stewart K, Hu H, et al. Progress toward the elimination of hepatitis B virus transmission among health care workers in the United States. *Arch Intern Med*. 1997;157:2601–2605.
- Occupational exposure to bloodborne pathogens—OSHA. Final rule. *Fed Regist*. 1991;56:64004–64182.
- Agerton TB, Mahoney FJ, Polish LB, et al. Impact of the bloodborne pathogens standard on vaccination of health-care workers with hepatitis B vaccine. *Infect Control Hosp Epidemiol*. 1995;16:287–291.
- Schillie S, Murphy TV, Sawyer M, et al. CDC guidance for evaluating health-care personnel for hepatitis B virus protection and for administering postexposure management. *MMWR Recomm Rep*. 2013;62(RR-10):1–19.
- Werner BG, Grady GF. Accidental hepatitis-B-surface-antigen-positive inoculations. Use of e antigen to estimate infectivity. *Ann Intern Med*. 1982;97:367–369.
- Centers for Disease Control, Prevention. Updated CDC recommendations for the management of hepatitis B virus-infected health-care providers and students. *MMWR Recomm Rep*. 2012;61(RR-3):1–12.
- Bell DM, Shapiro CN, Ciesielski CA, et al. Preventing bloodborne pathogen transmission from health-care workers to patients. The CDC perspective. *Surg Clin North Am*. 1995;75:1189–1203.
- Garibaldi RA, Rasmussen CN, Holmes AW, et al. Hospital-acquired serum hepatitis. Report of an outbreak. *JAMA*. 1972;219:1577–1580.
- Gerety RJ. Hepatitis B transmission between dental or medical workers and patients. *Ann Intern Med*. 1981;95:229–231.
- Lettau LA, Smith JD, Williams D, et al. Transmission of hepatitis B with resultant restriction of surgical practice. *JAMA*. 1986;255:934–937.
- Spijkerman JJ, van Doorn LJ, Janssen MH, et al. Transmission of hepatitis B virus from a surgeon to his patients during high-risk and low-risk surgical procedures during 4 years. *Infect Control Hosp Epidemiol*. 2002;23:306–312.
- Enfield KB, Sharapov U, Hall KK, et al. Transmission of hepatitis B virus from an orthopedic surgeon with a high viral load. *Clin Infect Dis*. 2013;56:218–224.
- Sugimoto S, Nagakubo S, Ito T, et al. A case of acute hepatitis B related to previous gynecological surgery in Japan. *J Infect Chemother*. 2013;19:524–529.
- Henderson DK, Dembry L, Fishman NO, et al. SHEA Guideline for management of healthcare workers who are infected with hepatitis B virus, hepatitis C virus and/or human immunodeficiency virus. *Infect Control Hosp Epidemiol*. 2010;31:203–232.

77. Shanson DC. Hepatitis B outbreak in operating-theatre and intensive care staff. *Lancet*. 1980;2:596.
78. Petrosillo N, Puro V, Jagger J, et al. The risks of occupational exposure and infection by human immunodeficiency virus, hepatitis B virus, and hepatitis C virus in the dialysis setting. Italian Multicenter Study on Nosocomial and Occupational Risk of Infections in Dialysis. *Am J Infect Control*. 1995;23:278–285.
79. Centers for Disease Control, Prevention. Outbreaks of hepatitis B virus infection among hemodialysis patients—California, Nebraska, and Texas, 1994. *MMWR Morb Mortal Wkly Rep*. 1996;45:285–289.
80. CDC. Summarized from Recommendations of the Advisory Committee on Immunization Practices (ACIP); 2012. https://www.cdc.gov/dialysis/pdfs/vaccinating_dialysis_patients_and_patients_dec2012pdf. Accessed January 15, 2018.
81. Alter MJ, Favero MS, Maynard JE. Impact of infection control strategies on the incidence of dialysis-associated hepatitis in the United States. *J Infect Dis*. 1986;153:1149–1151.
82. Stevens CE, Alter HJ, Taylor PE, et al. Hepatitis B vaccine in patients receiving hemodialysis. Immunogenicity and efficacy. *N Engl J Med*. 1984;311:496–501.
83. Hutin YJ, Harpaz R, Drobeniuc J, et al. Injections given in healthcare settings as a major source of acute hepatitis B in Moldova. *Int J Epidemiol*. 1999;28:782–786.
84. Narendranathan M, Philip M. Reusable needles—a major risk factor for acute virus B hepatitis. *Trop Doct*. 1993;23:64–66.
85. Duffell EF, Milne LM, Seng C, et al. Five hepatitis B outbreaks in care homes in the UK associated with deficiencies in infection control practice in blood glucose monitoring. *Epidemiol Infect*. 2011;139:327–335.
86. Perz JF, Grytdal S, Beck S, et al. Case-control study of hepatitis B and hepatitis C in older adults: do healthcare exposures contribute to burden of new infections? *Hepatology*. 2013;57:917–924.
87. Centers for Disease Control, Prevention. Investigation of viral hepatitis infections possibly associated with health-care delivery—New York City, 2008–2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:333–338.
88. Harling R, Turbitt D, Millar M, et al. Passage from India: an outbreak of hepatitis B linked to a patient who acquired infection from health care overseas. *Public Health*. 2007;121:734–741.
89. Centers for Disease Control, Prevention. Noninfluenza vaccination coverage among adults—United States, 2011. *MMWR Morb Mortal Wkly Rep*. 2013;62:66–72.
90. Roome AJ, Walsh SJ, Cartter ML, et al. Hepatitis B vaccine responsiveness in Connecticut public safety personnel. *JAMA*. 1993;270:2931–2934.
91. Ghebrehewet S, Baxter D, Falconer M, et al. Intradermal recombinant hepatitis B vaccination (IDRV) for non-responsive healthcare workers (HCWs). *Hum Vaccin*. 2008;4:280–285.
92. Simard EP, Miller JT, George PA, et al. Hepatitis B vaccination coverage levels among healthcare workers in the United States, 2002–2003. *Infect Control Hosp Epidemiol*. 2007;28:783–790.
93. Wainwright RB, Bulkow LR, Parkinson AJ, et al. Protection provided by hepatitis B vaccine in a Yupik Eskimo population—results of a 10-year study. *J Infect Dis*. 1997;175:674–677.
94. McMahon BJ, Bruden DL, Petersen KM, et al. Antibody levels and protection after hepatitis B vaccination: results of a 15-year follow-up. *Ann Intern Med*. 2005;142:333–341.
95. Bruce MG, Bruden D, Hurlburt D, et al. Antibody levels and protection after hepatitis B vaccine: results of a 30-year follow-up study and response to a booster dose. *J Infect Dis*. 2016;214:16–22.
96. Simons BC, Spradling PR, Bruden DJ, et al. A longitudinal hepatitis B vaccine cohort demonstrates long-lasting hepatitis B virus (HBV) cellular immunity despite loss of antibody against HBV surface antigen. *J Infect Dis*. 2016;214:273–280.
97. Peto TJ, Mendy ME, Lowe Y, et al. Efficacy and effectiveness of infant vaccination against chronic hepatitis B in the Gambia Hepatitis Intervention Study (1986–90) and in the nationwide immunisation program. *BMC Infect Dis*. 2014;14:7.
98. Gara N, Abdalla A, Rivera E, et al. Durability of antibody response against hepatitis B virus in healthcare workers vaccinated as adults. *Clin Infect Dis*. 2015;60:505–513.
99. Werner JM, Abdalla A, Gara N, et al. The hepatitis B vaccine protects re-exposed health care workers, but does not provide sterilizing immunity. *Gastroenterology*. 2013;145:1026–1034.
- 99a. Schillie S, Harris A, Link-Gelles R, et al. Recommendations of the Advisory Committee on Immunization Practices for use of a hepatitis B vaccine with a novel adjuvant. *MMWR Morb Mortal Wkly Rep*. 2018;67:455–458.
100. Beasley RP, Hwang LY, Stevens CE, et al. Efficacy of hepatitis B immune globulin for prevention of perinatal transmission of the hepatitis B virus carrier state: final report of a randomized double-blind, placebo-controlled trial. *Hepatology*. 1983;3:135–141.
101. U.S. Public Health Service. Updated U.S. public health service guidelines for the management of occupational exposures to HBV, HCV, and HIV and recommendations for postexposure prophylaxis. *MMWR Recomm Rep*. 2001;50(RR-11):1–52.
102. Younai FS. Health care-associated transmission of hepatitis B & C viruses in dental care (dentistry). *Clin Liver Dis*. 2010;14:93–104, ix.
103. Westermann C, Peters C, Lisiak B, et al. The prevalence of hepatitis C among healthcare workers: a systematic review and meta-analysis. *Occup Environ Med*. 2015;72:880–888.
104. Tomkins SE, Elford J, Nichols T, et al. Occupational transmission of hepatitis C in healthcare workers and factors associated with seroconversion: UK surveillance data. *J Viral Hepat*. 2012;19:199–204.
105. Puro V, Petrosillo N, Ippolito G, et al. Occupational hepatitis C virus infection in Italian health care workers. Italian Study Group on Occupational Risk of Bloodborne Infections. *Am J Public Health*. 1995;85:1272–1275.
106. Mitsui T, Iwano K, Masuko K, et al. Hepatitis C virus infection in medical personnel after needlestick accident. *Hepatology*. 1992;16:1109–1114.
107. Lanphear BP, Linnemann CC Jr, Cannon CG, et al. Hepatitis C virus infection in healthcare workers: risk of exposure and infection. *Infect Control Hosp Epidemiol*. 1994;15:745–750.
108. Naggie S, Holland DP, Sulowski MS, et al. Hepatitis C virus postexposure prophylaxis in the healthcare worker: why direct-acting antivirals don't change a thing. *Clin Infect Dis*. 2017;64:92–99.
109. Yazdanpanah Y, De Carli G, Miguieres B, et al. Risk factors for hepatitis C virus transmission to health care workers after occupational exposure: a European case-control study. *Clin Infect Dis*. 2005;41:1423–1430.
110. Ridzon R, Gallagher K, Ciesielski C, et al. Simultaneous transmission of human immunodeficiency virus and hepatitis C virus from a needle-stick injury. *N Engl J Med*. 1997;336:919–922.
111. Olsen K, Dahl PE, Paulsen EJ, et al. Increased risk of transmission of hepatitis C in open heart surgery compared with vascular and pulmonary surgery. *Ann Thorac Surg*. 2010;90:1425–1431.
112. Ross RS, Viazov S, Gross T, et al. Transmission of hepatitis C virus from a patient to an anesthesiology assistant to five patients. *N Engl J Med*. 2000;343:1851–1854.
113. Cody SH, Nainan OV, Garfein RS, et al. Hepatitis C virus transmission from an anesthesiologist to a patient. *Arch Intern Med*. 2002;162:345–350.
114. Cardell K, Widell A, Fryden A, et al. Nosocomial hepatitis C in a thoracic surgery unit: retrospective findings generating a prospective study. *J Hosp Infect*. 2008;68:322–328.
115. Stark K, Hanel M, Berg T, et al. Nosocomial transmission of hepatitis C virus from an anesthesiologist to three patients—epidemiologic and molecular evidence. *Arch Virol*. 2006;151:1025–1030.
116. Roy KM, Galmes-Truyols A, Gimenez-Duran J, et al. Epidemiology and molecular investigation of hepatitis C infection following holiday haemodialysis. *J Hosp Infect*. 2012;82:158–163.
117. Bosch X. Hepatitis C outbreak astounds Spain. *Lancet*. 1998;351:1415.
118. Warner AE, Schaefer MK, Patel PR, et al. Outbreak of hepatitis C virus infection associated with narcotic diversion by an hepatitis C virus-infected surgical technician. *Am J Infect Control*. 2015;43:53–58.
119. Fabrizi E, Martin P, Dixit V, et al. Acquisition of hepatitis C virus in hemodialysis patients: a prospective study by branched DNA signal amplification assay. *Am J Kidney Dis*. 1998;31:647–654.
120. Puttering H, Vychtyl A. Hepatitis B and C in peritoneal dialysis patients. *Semin Nephrol*. 2002;22:351–360.
121. Hinrichsen H, Leimenstoll G, Stegen G, et al. Prevalence and risk factors of hepatitis C virus infection in haemodialysis patients: a multicentre study in 2796 patients. *Gut*. 2002;51:429–433.
122. Saune K, Kamar N, Miedouge M, et al. Decreased prevalence and incidence of HCV markers in haemodialysis units: a multicentric French survey. *Nephrol Dial Transplant*. 2011;26:2309–2316.
123. Freiman JM, Tran TM, Schumacher SG, et al. Hepatitis C core antigen testing for diagnosis of hepatitis C virus infection: a systematic review and meta-analysis. *Ann Intern Med*. 2016;165:345–355.
124. Niu MT, Alter MJ, Kristensen C, et al. Outbreak of hemodialysis-associated non-A, non-B hepatitis and correlation with antibody to hepatitis C virus. *Am J Kidney Dis*. 1992;19:345–352.
125. Girou E, Chevaliez S, Challine D, et al. Determinant roles of environmental contamination and noncompliance with standard precautions in the risk of hepatitis C virus transmission in a hemodialysis unit. *Clin Infect Dis*. 2008;47:627–633.
126. Savey A, Simon F, Izopet J, et al. A large nosocomial outbreak of hepatitis C virus infections at a hemodialysis center. *Infect Control Hosp Epidemiol*. 2005;26:752–760.
127. Fornis X, Fernandez-Llana P, Pons M, et al. Incidence and risk factors of hepatitis C virus infection in a haemodialysis unit. *Nephrol Dial Transplant*. 1997;12:736–740.
128. Hardy NM, Sandroni S, Danielson S, et al. Antibody to hepatitis C virus increases with time on hemodialysis. *Clin Nephrol*. 1992;38:44–48.
129. Rahnavardi M, Hosseini Moghaddam SM, Alavian SM. Hepatitis C in hemodialysis patients: current global magnitude, natural history, diagnostic difficulties, and preventive measures. *Am J Nephrol*. 2008;28:628–640.
130. Hayashi H, Okuda K, Yokosuka O, et al. Adsorption of hepatitis C virus particles onto the dialyzer membrane. *Artif Organs*. 1997;21:1056–1059.
131. Pereira BJ, Vevey AS. Hepatitis C virus infection in dialysis and renal transplantation. *Kidney Int*. 1997;51:981–999.
132. Pereira BJ, Milford EL, Kirkman RL, et al. Transmission of hepatitis C virus by organ transplantation. *N Engl J Med*. 1991;325:454–460.
133. Pereira BJ, Milford EL, Kirkman RL, et al. Prevalence of hepatitis C virus RNA in organ donors positive for hepatitis C antibody and in the recipients of their organs. *N Engl J Med*. 1992;327:910–915.
134. Tugwell BD, Patel PR, Williams IT, et al. Transmission of hepatitis C virus to several organ and tissue recipients from an antibody-negative donor. *Ann Intern Med*. 2005;143:648–654.
135. Centers for Disease Control, Prevention. Transmission of hepatitis C virus through transplanted organs and tissue—Kentucky and Massachusetts, 2011. *MMWR Morb Mortal Wkly Rep*. 2011;60:1697–1700.
136. Centers for Disease Control, Prevention. Potential transmission of viral hepatitis through use of stored blood vessels as conduits in organ transplantation—Pennsylvania, 2009. *MMWR Morb Mortal Wkly Rep*. 2011;60:172–174.
137. Fischer GE, Schaefer MK, Labus BJ, et al. Hepatitis C virus infections from unsafe injection practices at an endoscopy clinic in Las Vegas, Nevada, 2007–2008. *Clin Infect Dis*. 2010;51:267–273.
138. Muscarella LE, New York State health officials. Recommendations for preventing hepatitis C virus infection: analysis of a Brooklyn endoscopy clinic's outbreak. *Infect Control Hosp Epidemiol*. 2001;22:669.
139. Moore ZS, Schaefer MK, Hoffmann KK, et al. Transmission of hepatitis C virus during myocardial perfusion imaging in an outpatient clinic. *Am J Cardiol*. 2011;108:126–132.
140. Panella H, Rius C, Cayla JA, Barcelona Hepatitis C Nosocomial Research Working Group. Transmission of hepatitis C virus during computed tomography scanning with contrast. *Emerg Infect Dis*. 2008;14:333–336.
141. Arnold S, Melville SK, Morehead B, et al. Notes from the field: hepatitis C transmission from inappropriate reuse of saline flush syringes for multiple patients in an acute care general hospital—Texas, 2015. *MMWR Morb Mortal Wkly Rep*. 2017;66:258–260.
142. Comstock RD, Mallonee S, Fox JL, et al. A large nosocomial outbreak of hepatitis C and hepatitis B among patients receiving pain remediation treatments. *Infect Control Hosp Epidemiol*. 2004;25:576–583.
143. Germain JM, Carbone A, Thiers V, et al. Patient-to-patient transmission of hepatitis C virus through the use of multidose vials during general anesthesia. *Infect Control Hosp Epidemiol*. 2005;26:789–792.
144. Desenclos JC, Bourdiol-Razes M, Rolin B, et al. Hepatitis C in a ward for cystic fibrosis and diabetic patients: possible transmission by spring-loaded finger-stick devices for self-monitoring of capillary blood glucose. *Infect Control Hosp Epidemiol*. 2001;22:701–707.
145. Bronowicki JP, Venard V, Botte C, et al. Patient-to-patient transmission of hepatitis C virus during colonoscopy. *N Engl J Med*. 1997;337:237–240.

146. Apostolou A, Bartholomew ML, Greeley R, et al. Transmission of hepatitis C virus associated with surgical procedures - New Jersey 2010 and Wisconsin 2011. *MMWR Morb Mortal Wkly Rep.* 2015;64:165–170.
147. Weaver JM. Confirmed transmission of hepatitis C in an oral surgery office. *Anesth Prog.* 2014;61:93–94.
148. Macedo de Oliveira A, White KL, Leschinsky DP, et al. An outbreak of hepatitis C virus infections among outpatients at a hematology/oncology clinic. *Ann Intern Med.* 2005;142:898–902.
149. Sepkowitz KA. Risk to cancer patients from nosocomial hepatitis C virus. *Infect Control Hosp Epidemiol.* 2004;25:599–602.
150. Pekova LM, Teocharov P, Sakarev A. Clinical course and outcome of a nosocomial outbreak of hepatitis C in a urology ward. *J Hosp Infect.* 2007;67:86–91.
151. Forns X, Martinez-Bauer E, Feliu A, et al. Nosocomial transmission of HCV in the liver unit of a tertiary care center. *Hepatology.* 2005;41:115–122.
152. Martinez-Bauer E, Forns X, Armelles M, et al. Hospital admission is a relevant source of hepatitis C virus acquisition in Spain. *J Hepatol.* 2008;48:20–27.
153. Paintsil E, Binka M, Patel A, et al. Hepatitis C virus maintains infectivity for weeks after drying on inanimate surfaces at room temperature: implications for risks of transmission. *J Infect Dis.* 2014;209:1205–1211.
154. Centers for Disease Control and Prevention. Testing for HCV infection: an update of guidance for clinicians and laboratorians. *MMWR Morb Mortal Wkly Rep.* 2013;62:362–365.
155. Jaeckel E, Cornberg M, Wedemeyer H, et al. Treatment of acute hepatitis C with interferon alfa-2b. *N Engl J Med.* 2001;345:1452–1457.

Transfusion- and Transplantation-Transmitted Infections

Sridhar V. Basavaraju^a and Matthew J. Kuehnert

SHORT VIEW SUMMARY

Definition

- Transfusion- and transplant-transmitted infections occur when a pathogen is spread from the donor to recipients through blood, organs, or other tissue.

Epidemiology

- Transfusion- and transplant-transmitted infections are unusual but occur when screening does not prevent transmission, or when a donation is not screened for a pathogen. The risk of infection transmitted through organ transplantation is likely

greater than through blood or tissue owing to the lack of exclusion criteria for organ donors.

Microbiology

- Transfusion of blood product has transmitted infections caused by bacteria, viruses, parasites, and prions.
- Solid-organ transplantation has transmitted infections caused by viruses, bacteria, fungi, mycobacteria, and parasites.
- Encephalitis after solid-organ transplantation has resulted from transmission of West Nile

virus, lymphocytic choriomeningitis virus, rabies, and *Balamuthia mandrillaris*.

Diagnosis

- Diagnosis is dependent on clinical recognition that a recipient has an infection that could have been acquired from a transfusion or transplant.

Therapy and Prevention

- Better donor screening, or prompt recognition when transfusion- or transplant-transmitted infection occurs, can improve recipient outcomes.

TRANSFUSION-ASSOCIATED INFECTIONS

Beeson reported the first cases of transfusion-associated hepatitis in 1943, describing seven patients who developed illness 1 to 4 months after having received a red blood cell (RBC) or plasma transfusion.¹ Later identified as transfusion-transmitted hepatitis B virus (HBV), the advent of the acquired immunodeficiency syndrome (AIDS) epidemic in the 1980s followed by recognition of transfusion transmission of hepatitis C virus (HCV) in the 1990s and, most recently, reports of transfusion-transmitted West Nile virus (WNV) and Zika virus (ZIKV) infections in the 21st century have continued to draw attention to the safety of the blood supply, particularly the potential risk from emerging infectious agents.²⁻⁵

Although infections with the human immunodeficiency virus (HIV), HBV, HCV, WNV, and ZIKV have been the most publicized, a wide spectrum of other organisms, including viruses, bacteria, parasites, and prions, have been transmitted by transfusion (Table 304.1).⁶ Of these, transfusion transmission of bacterial infections is most common and earliest recognized in the history of transfusion, now is most often associated with platelet transfusions, and infrequently results in sepsis and death.⁷ Recognition of the threat of *Trypanosoma cruzi* transmission via transfusion and the availability of suitable testing have led to screening for this pathogen, which is the cause of Chagas disease. In addition to WNV, mosquito-borne pathogens, such as dengue virus, chikungunya virus, and most recently, ZIKV, have shown potential for transfusion transmission, although the burden of disease is unknown. Tick-borne agents also are recognized to pose an increasing risk to transfusion safety, including transmission of babesiosis and, most recently, anaplasmosis and ehrlichiosis. Transmission of variant Creutzfeldt-Jakob disease (vCJD) via transfusion has occurred in the United Kingdom.

An ongoing dilemma of the blood-banking community is a mandate to ensure a maximally safe blood supply while giving consideration to the cost of such measures. The current questions for screening donors that have been adopted by most blood banks are presented in Fig.

304.1, and the laboratory screening measures currently in place are listed in Table 304.2. As of 2014, pathogens or diseases screened solely through donor interview include malaria and vCJD. Real or perceived threats to the blood supply occur regularly, as emerging or reemerging infectious diseases raise concern over potential transmission. A previously published review summarizes potential blood supply risks from pathogens, the majority of which are not screened for with laboratory testing.⁸

Pathogen reduction technology (PRT) is now available to reduce the need to develop screening measures for new infectious threats. In December 2014, the US Food and Drug Administration (FDA) approved a PRT method for plasma and apheresis platelets that uses chemicals activated by ultraviolet light to inactivate nucleic acid material comprising DNA and RNA.^{9,10} However, there is no approved method for treatment of RBCs. Although this technology is likely to reduce the risk of transmitted infections of viruses, bacteria, and parasites, it will not likely eliminate it, because there are organisms occurring in high titers or that have structures (e.g., non-lipid-enveloped viruses, bacterial spores, prions) that resist or are not affected by the inactivation mechanism.^{9,10}

SCOPE OF BLOOD TRANSFUSION

According to estimates from the World Health Organization (WHO), more than 112 million units of blood were collected in 2013.¹¹ Less than half of donated blood is collected in developing and transitional countries, which are home to about 80% of the world's population. Of the 156 countries providing data to WHO for 2013, 13 were not able to screen all of their donated blood for one or more of the four infections (HIV, HBV, HCV, and syphilis) that are most widely recognized to be transmitted through blood and are recommended by WHO to be screened at donation.¹¹ A total of 126 countries have national guidelines on the appropriate clinical use of blood, and 70 countries have a national hemovigilance system to monitor adverse events associated with transfusion.¹¹

Surveys to determine blood product use in the United States, led by the National Heart and Lung Institute (now called the National Heart, Lung, and Blood Institute), began in 1971.¹² In the United States, surveys have reported the frequency of blood collection and utilization

^aThe findings and conclusions in this chapter are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

TABLE 304.1 List of Notable Infections Transmitted Through Blood Transfusion**Viruses**

Colorado tick fever virus
 Cytomegalovirus
 Dengue virus
 Epstein-Barr virus
 Hepatitis A virus
 Hepatitis B virus
 Hepatitis C virus
 Hepatitis E virus
 Human herpesvirus 8
 Human immunodeficiency virus 1 and 2
 Human T-lymphotropic virus 1 and 2
 Parvovirus B19
 Tick-borne encephalitis virus
 West Nile virus
 Zika virus

Bacteria

Anaplasma phagocytophilum
Brucella spp.
Coxiella burnetii
Ehrlichia
 Gram-positive organisms^a
 Gram-negative organisms^b
Rickettsia rickettsii
Treponema pallidum^c

Parasites

Babesia spp.
Leishmania spp.
Trypanosoma cruzi
Plasmodium spp.

Prions

Variant Creutzfeldt-Jakob disease

^aHigher risk in platelets (see Table 304.3); in addition, *Streptococcus gallolyticus* (bovis) is notable for being associated with colon cancer in the donor.

^bNoted in red blood cell units associated with cryophilic organisms, including *Pseudomonas fluorescens*, *Serratia liquefaciens*, and *Yersinia enterocolitica*.

^cNot thought a current risk; last reported transfusion transmission of syphilis in the United States was in 1966.

Modified from Perkins HA, Busch MP. Transfusion-associated infections: 50 years of relentless challenges and remarkable progress. *Transfusion*. 2010;50:2080–2099; and Stramer SL, Hollinger FB, Katz LM, et al. Emerging infectious disease agents and their potential threat to transfusion safety. *Transfusion*. 2009;49:15–295.

since the late 1980s, most recently through the National Blood Collection and Utilization Survey (NBCUS), which has been conducted biannually since 1998.^{12–14} In the 2017 NBCUS report, reflecting data collected in 2015, there were nearly 11 million whole-blood and 3.7 million apheresis collections, with approximately 19 million blood components transfused.¹⁴ In 2015, 11.33 million whole-blood and RBC units, 2.7 million units of plasma, 1.8 million units of apheresis platelets, 0.17 million units of whole-blood-derived platelets (measured in apheresis equivalent units), and 1.2 million units of cryoprecipitate were given.¹⁴ The number of RBC transfusions has continued to decline since 2008, perhaps reflecting the growing adoption of more aggressive hospital blood management practices. The number of transfused platelets has remained approximately the same, but transfusion of apheresis platelets has continued to increase, while the amount of whole-blood-derived or pooled platelets has continued to decline. In addition to contributions from the voluntary donor pool, plasma units also are collected annually from paid donors and are used to prepare immune globulin, albumin, and various other plasma-derived products. By 1987, the cost of collecting, processing, and transfusing patients exceeded \$3 billion; since then, costs have increased steadily with the addition of new screening tests and the implementation of leukoreduction.¹² In 2015, the mean price paid by a hospital was \$217 for a unit of leukocyte-reduced RBCs, \$60 for fresh-frozen plasma, and \$537 for apheresis platelets, so current costs are likely to exceed \$4 billion annually.^{14,15}

It has been estimated that the annual likelihood of an individual's receiving a transfusion increases dramatically with age; in 2015, the

TABLE 304.2 Laboratory Screening Performed for Pathogens or Diseases by US Blood Collection Centers**Required or Recommended by US Food and Drug Administration**

Hepatitis B surface antigen (HBsAg)
 Hepatitis B core antibody (anti-HBcAb)
 Hepatitis C virus antibody (anti-HCV)
 HIV-1 and HIV-2 antibody (anti-HIV-1 and anti-HIV-2)
 HIV p24 antigen
 HTLV-1 and HTLV-2 antibody (anti-HTLV-1 and anti-HTLV-2)
 Serologic test for syphilis
 NAAT for HIV-1, HBV, and HCV
 NAAT for West Nile virus
 Serologic test for *Trypanosoma cruzi* (one-time donor testing)

Additionally Required by Accrediting Organizations^a

Platelet screening for bacterial contamination^b

Performed Voluntarily

Serologic test for cytomegalovirus (on request)

Performed Under Investigational Protocols

PCR and serologic test for *Babesia microti*
 NAAT for Zika virus

^aIncluding AABB (formerly known as the American Association of Blood Banks) and the College of American Pathologists.

^bIn general, liquid culture media for apheresis units and pH/glucose indicators or bacterial antigen point-of-use tests for whole-blood-derived pooled units.

HBV, Hepatitis B virus; HCV, hepatitis C virus; HIV, human immunodeficiency virus; HTLV, human T-cell lymphotropic virus; NAAT, nucleic acid amplification testing; PCR, polymerase chain reaction.

transfusion rate in the US population was 35.3 units per 1000 people.¹⁴ Because of concern about contracting an infectious disease, there has been historic interest in autologous and donor-directed blood donation. However, donor-directed units, usually given by family members for a specific patient, have been shown to have higher rates of various infectious agents. There has been movement away from donor-directed donation and toward building a dedicated, voluntary repeat donor population.^{14,15} Viral infections are much less common among repeat donors compared with first-time whole-blood donors, and they may be even less common among donors associated with apheresis collection.^{16,17}

A majority of, but not all, blood components transfused in the United States are leukoreduced, a modification that is performed to filter out the majority of white blood cells. Leukoreduction is performed to reduce nonhemolytic transfusion reactions, but it may also reduce the risk for infectious disease transmission through removal of infected white blood cells, particularly for cell-associated agents.¹⁸

In 2015, only about 0.42% of the donated allogeneic blood supply was discarded on testing, usually because of detection of a potentially transmissible infection.¹⁴ This reflects a continued steady decline in units discarded owing to testing, perhaps indicating improved accuracy of screening tests and retention of repeat donors. The incidence of viral markers in donated units was increased among volunteers who donated after September 11, 2001, primarily owing to a large cohort of first-time and infrequent repeat donors.¹⁹

BLOODBORNE PATHOGENS

Transfusion-associated transmission risk has persisted for HIV, HBV, and HCV for two distinct reasons: (1) the incomplete sensitivity of the available screening tests, and (2) the “window” period, which is defined as the period between acute infection (and potential infectivity) and the point at which serologic tests can reliably detect infection.

The development and implementation of routine nucleic acid amplification testing (NAAT) has transformed blood bank screening for viral pathogens.^{3,20–22} NAAT can detect viral RNA after the first 10 to 14 days of infection, narrowing the window period by 7 to 10 days for HIV and by 50 to 60 days for HCV compared with serology. NAAT for blood donor screening, in addition to HIV, HBV, and HCV, is now

additionally required for WNV and ZIKV, and is voluntarily performed by plasma manufacturers in pools for parvovirus B19.²³⁻²⁶

With introduction of NAAT, the risk of acquiring HIV or HCV per unit of blood transfused has plummeted from approximately 1 per 500,000 for HIV and 1 per 100,000 for HCV before NAAT to a current rate of 1 per 2 million for both viruses. Detected cases are so unusual that the incidence can only be estimated statistically.²⁷ As remarkable as this advance is, a small window period remains, meaning that the risk for receiving one of these viruses in a unit of blood is not zero. Four cases of HIV transmission and many more cases of HCV transmission through transfusion have been recognized and reported to public health authorities since NAAT was implemented; the modeling

data estimate more cases than the number reported, so some cases likely go unrecognized.²⁸

The advent of NAAT also has had a substantial impact on the choice of serologic tests. In addition to sensitivity, blood bankers seek optimal specificity: False-positive or indeterminate test results exclude otherwise appropriate donors from subsequent donations and create anxiety due to incorrect diagnosis of infection. The introduction of an extremely sensitive test such as NAAT allows blood banks to focus on the specificity of other screening tests, thereby limiting the number of persons potentially excluded from the donor pool because of misleading test results. Although sensitivity varies little between the second- and third-generation antibody tests, the third-generation assay has superior

Full-Length Donor History Questionnaire

	Yes	No	
Are you			
1. Feeling healthy and well today?	<input type="checkbox"/>	<input type="checkbox"/>	
2. Currently taking an antibiotic?	<input type="checkbox"/>	<input type="checkbox"/>	
3. Currently taking any other medication for an infection?	<input type="checkbox"/>	<input type="checkbox"/>	
Please read the Medication Deferral List.			
4. Are you now taking or have you ever taken any medications on the Medication Deferral List?	<input type="checkbox"/>	<input type="checkbox"/>	
5. Have you read the educational materials?	<input type="checkbox"/>	<input type="checkbox"/>	
In the past 48 hours			
6. Have you taken aspirin or anything that has aspirin in it?	<input type="checkbox"/>	<input type="checkbox"/>	
In the past 6 weeks			
7. Female donors: Have you been pregnant or are you pregnant now? (Males: check "I am male.")	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> I am male
In the past 8 weeks have you			
8. Donated blood, platelets or plasma?	<input type="checkbox"/>	<input type="checkbox"/>	
9. Had any vaccinations or other shots?	<input type="checkbox"/>	<input type="checkbox"/>	
10. Had contact with someone who had a smallpox vaccination?	<input type="checkbox"/>	<input type="checkbox"/>	
In the past 16 weeks			
11. Have you donated a double unit of red cells using an apheresis machine?	<input type="checkbox"/>	<input type="checkbox"/>	
In the past 12 months have you			
12. Had a blood transfusion?	<input type="checkbox"/>	<input type="checkbox"/>	
13. Had a transplant such as organ, tissue, or bone marrow?	<input type="checkbox"/>	<input type="checkbox"/>	
14. Had a graft such as bone or skin?	<input type="checkbox"/>	<input type="checkbox"/>	
15. Come into contact with someone else's blood?	<input type="checkbox"/>	<input type="checkbox"/>	
16. Had an accidental needle-stick?	<input type="checkbox"/>	<input type="checkbox"/>	
17. Had sexual contact with anyone who has HIV/AIDS or has had a positive test for the HIV/AIDS virus?	<input type="checkbox"/>	<input type="checkbox"/>	
18. Had sexual contact with a prostitute or anyone else who takes money or drugs or other payment for sex?	<input type="checkbox"/>	<input type="checkbox"/>	
19. Had sexual contact with anyone who has ever used needles to take drugs or steroids, or anything <u>not</u> prescribed by their doctor?	<input type="checkbox"/>	<input type="checkbox"/>	
20. Had sexual contact with anyone who has hemophilia or has used clotting factor concentrates?	<input type="checkbox"/>	<input type="checkbox"/>	
21. Female donors: Had sexual contact with a male who has ever had sexual contact with another male? (Males: check "I am male.")	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> I am male
22. Had sexual contact with a person who has hepatitis?	<input type="checkbox"/>	<input type="checkbox"/>	
23. Lived with a person who has hepatitis?	<input type="checkbox"/>	<input type="checkbox"/>	
24. Had a tattoo?	<input type="checkbox"/>	<input type="checkbox"/>	
25. Had ear or body piercing?	<input type="checkbox"/>	<input type="checkbox"/>	

v.1.3

eff May 2008

FIG. 304.1 AABB (formerly American Association of Blood Banks) Blood Donor History Questionnaire, Version 1.3, May 2008. (From AABB. Full-Length Donor History Questionnaire. <http://www.aabb.org/tm/questionnaires/Documents/dhq/v2/DHQ%20v2.0.pdf>. Accessed May 29, 2019.)

Full-Length Donor History Questionnaire—cont'd

	Yes	No
26. Had or been treated for syphilis or gonorrhea?	<input type="checkbox"/>	<input type="checkbox"/>
27. Been in juvenile detention, lockup, jail, or prison for more than 72 hours?	<input type="checkbox"/>	<input type="checkbox"/>
In the past three years have you		
28. Been outside the United States or Canada?	<input type="checkbox"/>	<input type="checkbox"/>
From 1980 through 1996 ,		
29. Did you spend time that adds up to three (3) months or more in the United Kingdom? (Review list of countries in the UK.)	<input type="checkbox"/>	<input type="checkbox"/>
30. Were you a member of the U.S. military, a civilian military employee, or a dependent of a member of the U.S. military?	<input type="checkbox"/>	<input type="checkbox"/>
From 1980 to the present , did you		
31. Spend time that adds up to five (5) years or more in Europe? (Review list of countries in Europe.)	<input type="checkbox"/>	<input type="checkbox"/>
32. Receive a blood transfusion in the United Kingdom or France? (Review list of countries in the UK.)	<input type="checkbox"/>	<input type="checkbox"/>
From 1977 to the present , have you		
33. Received money, drugs, or other payment for sex?	<input type="checkbox"/>	<input type="checkbox"/>
34. Male donors: had sexual contact with another male, even once? (Females: check "I am female.")	<input type="checkbox"/>	<input type="checkbox"/>
Have you EVER		
35. Had a positive test for the HIV/AIDS virus?	<input type="checkbox"/>	<input type="checkbox"/>
36. Used needles to take drugs, steroids, or anything <u>not</u> prescribed by your doctor?	<input type="checkbox"/>	<input type="checkbox"/>
37. Used clotting factor concentrates?	<input type="checkbox"/>	<input type="checkbox"/>
38. Had hepatitis?	<input type="checkbox"/>	<input type="checkbox"/>
39. Had malaria?	<input type="checkbox"/>	<input type="checkbox"/>
40. Had Chagas disease?	<input type="checkbox"/>	<input type="checkbox"/>
41. Had babesiosis?	<input type="checkbox"/>	<input type="checkbox"/>
42. Received a dura mater (or brain covering) graft?	<input type="checkbox"/>	<input type="checkbox"/>
43. Had any type of cancer, including leukemia?	<input type="checkbox"/>	<input type="checkbox"/>
44. Had any problems with your heart or lungs?	<input type="checkbox"/>	<input type="checkbox"/>
45. Had a bleeding condition or a blood disease?	<input type="checkbox"/>	<input type="checkbox"/>
46. Had sexual contact with anyone who was born in or lived in Africa?	<input type="checkbox"/>	<input type="checkbox"/>
47. Been in Africa?	<input type="checkbox"/>	<input type="checkbox"/>
48. Have any of your relatives had Creutzfeldt-Jakob disease?	<input type="checkbox"/>	<input type="checkbox"/>

☐ I am female

v.1.3

eff May 2008

FIG. 304.1, cont'd

specificity, decreasing the proportion of persons with indeterminate antibody tests for HCV.²⁹ HBV NAAT has been adopted for blood donor screening.

Human Immunodeficiency Virus Type 1

More than 8000 persons in the United States have developed AIDS from receipt of blood or tissue.³⁰ In addition, at least 50% of all hemophiliacs in the United States and Europe became infected with HIV from 1978 to 1985, most from receipt of infected plasma factors, with the highest incidence in 1982, when there were 22 infections per 100 person-years.³¹ The first HIV screening test was introduced in 1985, and since 1987 few new infections among hemophiliacs have been reported.³¹ As noted, the introduction of NAAT screening has substantially lowered the risk for HIV transmission, with the last reported case occurring in 2008.²⁸

Redundant methods of testing also have been useful in confirming the status of infection for blood donor follow-up. NAAT has been useful in clarifying the HIV serostatus of persons with indeterminate results of Western blot tests.³² These reactions occur in about 1 of every 5000 donations and usually represent false-positive test results.³³

Human Immunodeficiency Virus Type 2

In June 1992, the FDA mandated screening for HIV-2.³⁴ Since then, few HIV-2–positive donors have been identified, and no cases of transmission have occurred in the United States, although transfusion-related HIV-2 cases have occurred elsewhere.^{35,36}

Human T-Cell Lymphotropic Virus Types 1 and 2

HTLV-1 and HTLV-2, unlike HIV-1 and HIV-2, are cell associated and therefore predominantly transmissible only with blood component transfusions.³⁷ Screening for both types is done with a single test. The rate of transmission of HTLV-1 decreased almost 10-fold (from 1 per 8,500 to 1 per 69,000) after introduction of the screening test.³⁸ Transplantation of solid organs from an HTLV-1–infected donor resulted in rapid progression to subacute myelopathy in three recipients; this phenomenon has not been described in transfusion recipients.³⁹

Other Retroviruses

A short-lived perceived threat to the blood supply was raised when xenotropic murine leukemia virus–related virus (XMRV) was linked to prostate cancer and then chronic fatigue syndrome. Furthermore,