

References

- Bodey GP, Buckley M, Sathe YS, et al. Quantitative relationships between circulating leukocytes and infection in patients with acute leukemia. *Ann Intern Med.* 1966;64:328–340.
- Sickles EA, Greene WH, Wiernik PH. Clinical presentation of infection in granulocytopenic patients. *Arch Intern Med.* 1975;135:715–719.
- Castagnola E, Fontana V, Cavaglia I, et al. A prospective study on the epidemiology of febrile episodes during chemotherapy-induced neutropenia in children with cancer or after hemopoietic stem cell transplantation. *Clin Infect Dis.* 2007;45:1296–1304.
- Trecarichi EM, Pagano L, Candoni A, et al. Current epidemiology and antimicrobial resistance data for bacterial bloodstream infections in patients with hematologic malignancies: an Italian multicentre prospective survey. *Clin Microbiol Infect.* 2015;21:337–343.
- Caselli D, Cesaro S, Fagioli F, et al. Incidence of colonization and bloodstream infection with carbapenem-resistant Enterobacteriaceae in children receiving antineoplastic chemotherapy in Italy. *Infect Dis (Lond).* 2016;48:152–155.
- Albiger B, Glasner C, Struelens MJ, et al. Carbapenemase-producing Enterobacteriaceae in Europe: assessment by national experts from 38 countries, May 2015. *Euro Surveill.* 2015;20.
- Tacconelli E, Cataldo MA, Dancer SJ, et al. ESCMID guidelines for the management of the infection control measures to reduce transmission of multidrug-resistant Gram-negative bacteria in hospitalized patients. *Clin Microbiol Infect.* 2014;20:1–55.
- Urrea M, Rives S, Cruz O, et al. Nosocomial infections among pediatric hematology/oncology patients: results of a prospective incidence study. *Am J Infect Control.* 2004;32:205–208.
- Simon A, Ammann RA, Bode U, et al. Healthcare-associated infections in pediatric cancer patients: results of a prospective surveillance study from university hospitals in Germany and Switzerland. *BMC Infect Dis.* 2008;8:70.
- Castagnola E, Cavaglia I, Pistorio A, et al. Bloodstream infections and invasive mycoses in children undergoing acute leukaemia treatment: a 13-year experience at a single Italian institution. *Eur J Cancer.* 2005;41:1439–1445.
- Castagnola E, Rossi MR, Cesaro S, et al. Incidence of bacteremias and invasive mycoses in children with acute non-lymphoblastic leukemia: results from a multi-center Italian study. *Pediatr Blood Cancer.* 2010;55:1103–1107.
- Orasch C, Weisser M, Mertz D, et al. Comparison of infectious complications during induction/consolidation chemotherapy versus allogeneic hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2010;45:521–526.
- Dettenkofer M, Wenzler-Röttele S, Babikir R, et al. Surveillance of nosocomial sepsis and pneumonia in patients with a bone marrow or peripheral blood stem cell transplant: a multicenter project. *Clin Infect Dis.* 2005;40:926–931.
- Velasco E, Byington R, Martins CA, et al. Comparative study of clinical characteristics of neutropenic and non-neutropenic adult cancer patients with bloodstream infections. *Eur J Clin Microbiol Infect Dis.* 2006;25:1–7.
- Vekemans M, Robinson J, Georgala A, et al. Low mannose-binding lectin concentration is associated with severe infection in patients with hematological cancer who are undergoing chemotherapy. *Clin Infect Dis.* 2007;44:1593–1601.
- Gafer-Gvili A, Fraser A, Paul M, et al. Meta-analysis: antibiotic prophylaxis reduces mortality in neutropenic patients. *Ann Intern Med.* 2005;142:979–995.
- Bucaneve G, Micozzi A, Menichetti F, et al. Levofloxacin to prevent bacterial infection in patients with cancer and neutropenia. *N Engl J Med.* 2005;353:977–987.
- Girmania C, Lo Coco F, Breccia M, et al. Infectious complications in patients with acute promyelocytic leukaemia treated with the AIDA regimen. *Leukemia.* 2003;17:925–930.
- Lehrnbecher T, Warwig D, Kaiser J, et al. Infectious complications in pediatric acute myeloid leukemia: analysis of the prospective multi-institutional clinical trial AML-BFM 93. *Leukemia.* 2004;18:72–77.
- Cullen M, Steven N, Billingham L, et al. Antibacterial prophylaxis after chemotherapy for solid tumors and lymphomas. *N Engl J Med.* 2005;353:988–998.
- Caira M, Girmania C, Fadda RM, et al. Invasive fungal infections in patients with acute myeloid leukemia and in those submitted to allogeneic hemopoietic stem cell transplant: who is at highest risk? *Eur J Haematol.* 2008;81:242–243.
- Pagano L, Caira M, Rossi G, et al. A prospective survey of febrile events in hematological malignancies. *Ann Hematol.* 2012;91:767–774.
- Haupt R, Romanengo M, Fears T, et al. Incidence of septicemias and invasive mycoses in children undergoing treatment for solid tumours: a 12-year experience at a single Italian institution. *Eur J Cancer.* 2001;37:2413–2419.
- Lehrnbecher T, Zimmermann M, Reinhardt D, et al. Prophylactic human granulocyte colony-stimulating factor after induction therapy in pediatric acute myeloid leukemia. *Blood.* 2007;109:936–943.
- Viscoli C, Herbrecht R, Akan H, et al. An EORTC Phase II study of caspofungin as first-line therapy of invasive aspergillosis in haematological patients. *J Antimicrob Chemother.* 2009;64:1274–1281.
- Nivoix Y, Velten M, Letscher-Bru V, et al. Factors associated with overall and attributable mortality in invasive aspergillosis. *Clin Infect Dis.* 2008;47:1176–1184.
- Mikulska M, Raiola AM, Bruno B, et al. Risk factors for invasive aspergillosis and related mortality in recipients of allogeneic SCT from alternative donors: an analysis of 306 patients. *Bone Marrow Transplant.* 2009;44:361–370.
- Fernández-Ruiz M, Meije Y, Manuel O, et al. ESCMID Study Group for Infections in Compromised Hosts (ESGICH) Consensus Document on the safety of targeted and biological therapies: an infectious diseases perspective (Introduction). *Clin Microbiol Infect.* 2018;pii:S1198-743X(18)30147-2.
- Drgona L, Gudiol C, Lanini S, et al. ESCMID Study Group for Infections in Compromised Hosts (ESGICH) Consensus Document on the safety of targeted and biological therapies: an Infectious Diseases perspective (Agents targeting lymphoid or myeloid cells surface antigens [II]: CD22, CD30, CD33, CD38, CD40, SLAMF-7 and CCR4). *Clin Microbiol Infect.* 2018;pii:S1198-743X(18)30232-5.
- Carlisle PS, Gucalp R, Wiernik PH. Nosocomial infections in neutropenic cancer patients. *Infect Control Hosp Epidemiol.* 1993;14:320–324.
- Sonis ST. The pathobiology of mucositis. *Nat Rev Cancer.* 2004;4:277–284.
- van Vliet MJ, Harmsen HJ, de Bont ES, et al. The role of intestinal microbiota in the development and severity of chemotherapy-induced mucositis. *PLoS Pathog.* 2010;6:e1000879.
- van Vliet MJ, Tissing WJ, et al. Chemotherapy treatment in pediatric patients with acute myeloid leukemia receiving antimicrobial prophylaxis leads to a relative increase of colonization with potentially pathogenic bacteria in the gut. *Clin Infect Dis.* 2009;49:262–270.
- Montassier E, Gastinne T, Vangay P, et al. Chemotherapy-driven dysbiosis in the intestinal microbiome. *Aliment Pharmacol Ther.* 2015;42:515–528.
- Castagnola E, Ruberto E, Guarino A. Gastrointestinal and liver infections in children undergoing antineoplastic chemotherapy in the years 2000. *World J Gastroenterol.* 2016;22:5853–5866.
- Huang Y, Yang W, Liu H, et al. Effect of high-dose methotrexate chemotherapy on intestinal Bifidobacteria, Lactobacillus and Escherichia coli in children with acute lymphoblastic leukemia. *Exp Biol Med (Maywood).* 2012;237:305–311.
- Taur Y, Xavier JB, Lipuma L, et al. Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis.* 2012;55:905–914.
- Montassier E, Al-Ghathier GA, Ward T, et al. Pretreatment gut microbiome predicts chemotherapy-related bloodstream infection. *Genome Med.* 2016;8:49.
- Agirbasli H, Ozcan SA, Gedikoğlu G. Fecal fungal flora of pediatric healthy volunteers and immunosuppressed patients. *Mycopathologia.* 2005;159:515–520.
- Gammelrud KW. Colonization by Candida in children with cancer, children with cystic fibrosis, and healthy controls. *Clin Microbiol Infect.* 2011;17:1875–1881.
- Zakhour R, Chaffari AM, Raad II. Catheter-related infections in patients with hematological malignancies: novel preventive and therapeutic strategies. *Lancet Infect Dis.* 2016;16:e241–e250.
- Kulkarni S, Wu O, Kasthuri R, et al. Centrally inserted external catheters and totally implantable ports for the delivery of chemotherapy: a systematic review and meta-analysis of device-related complications. *Cardiovasc Intervent Radiol.* 2014;37:990–1008.
- Samaras P, Dold S, Braun J, et al. Infectious port complications are more frequent in younger patients with hematologic malignancies than in solid tumor patients. *Oncology.* 2008;74:237–244.
- Chopra V, O'Horo JC, Rogers MA, et al. The risk of bloodstream infection associated with peripherally inserted central catheters compared with central venous catheters in adults: a systematic review and meta-analysis. *Infect Control Hosp Epidemiol.* 2013;34:908–918.
- Hord JD, Lawlor J, Werner E, et al. Central line associated blood stream infections in pediatric hematology/oncology patients with different types of central lines. *Pediatr Blood Cancer.* 2016;63:1603–1607.
- Lipitz-Snyderman A, Sepkowitz KA, Elkin EB, et al. Long-term central venous catheter use and risk of infection in older adults with cancer. *J Clin Oncol.* 2014;32:2351–2356.
- CDC Centers for Disease Control and Prevention. National Healthcare Safety Network (NHSN). Bloodstream Infection Event (Central Line-Associated Bloodstream Infection and Non-Central Line-Associated Bloodstream Infection). January 2018. http://www.cdc.gov/nhsn/PDFs/pscManual/4PSC_CLABSCurrent.pdf. Accessed March 16, 2018.
- Metzger KE, Rucker Y, Callaghan M, et al. The burden of mucosal barrier injury laboratory-confirmed bloodstream infection among hematology, oncology, and stem cell transplant patients. *Infect Control Hosp Epidemiol.* 2015;36:119–124.
- Epstein L, See I, Edwards JR, et al. Mucosal barrier injury laboratory-confirmed bloodstream infections (MBI-LCBI): descriptive analysis of data reported to National Healthcare Safety Network (NHSN), 2013. *Infect Control Hosp Epidemiol.* 2016;37:2–7.
- Zakhour R, Hachem R, Alawami HM, et al. Comparing catheter-related bloodstream infections in pediatric and adult cancer patients. *Pediatr Blood Cancer.* 2017;64.
- Dommett R, Chisholm J, Turner M, et al. Mannose-binding lectin genotype influences frequency and duration of infectious complications in children with malignancy. *J Pediatr Hematol Oncol.* 2013;35:69–75.
- Pfeil AM, Vulsteke C, Paridaens R, et al. Clinical and genetic risk factors for epirubicin-induced cardiac toxicity in early breast cancer patients. *BMC Cancer.* 2014;14:201.
- Ok M, Einsele H, Loeffler J. Genetic susceptibility to *Aspergillus fumigatus* infections. *Int J Med Microbiol.* 2011;301:445–452.
- Funakoshi T, Suzuki M, Muss HB. Infection risk in breast cancer patients treated with trastuzumab: a systematic review and meta-analysis. *Breast Cancer Res Treat.* 2015;149:321–330.
- Mansi L, Thierry-Vuillemin A, Nguyen T, et al. Safety profile of new anticancer drugs. *Expert Opin Drug Saf.* 2010;9:301–317.
- Wang Y, Wang M, Wang Q, et al. Incidence and risk of infections associated with EGFR-TKIs in advanced non-small-cell lung cancer: a systematic review and meta-analysis of randomized controlled trials. *Oncotarget.* 2017;8:29406–29415.
- Garcia CA, Wu S. Attributable Risk of Infection to mTOR Inhibitors Everolimus and Temsirolimus in the Treatment of Cancer. *Cancer Invest.* 2016;34:521–530.
- Bringham S, De Wit E, Dimopoulos MA. New agents in multiple myeloma: an examination of safety profiles. *Clin Lymphoma Myeloma Leuk.* 2017;17:391–407, e5.
- Zhang X, Ran Y, Shao Y, et al. Incidence and risk of severe infections associated with aflibercept in cancer patients: a systematic review and meta-analysis. *Br J Clin Pharmacol.* 2016;81:33–40.
- Carlson KR, Newsome SD, Kim EJ, et al. Progressive multifocal leukoencephalopathy associated with brentuximab vedotin therapy: a report of 5 cases from the Southern Network on Adverse Reactions (SONAR) project. *Cancer.* 2014;120:2464–2471.
- Teh BW, Harrison SJ, Worth LJ, et al. Infection risk with immunomodulatory and proteasome inhibitor-based therapies across treatment phases for multiple myeloma: a systematic review and meta-analysis. *Eur J Cancer.* 2016;67:21–37.
- Mikulska M, Lanini S, Gudiol C, et al. ESCMID Study Group for Infections in Compromised Hosts (ESGICH) Consensus Document on the safety of targeted and biological therapies: an infectious diseases perspective (agents targeting lymphoid cells surface antigens [I]: CD19, CD20 and CD52). *Clin Microbiol Infect.* 2018;pii:S1198-743X(18)30151-4.
- Gentile G, Andreoni M, Antonelli G, et al. Screening, monitoring, prevention, prophylaxis and therapy for hepatitis B virus reactivation in patients with hematologic malignancies and patients who underwent hematologic stem cell transplantation: a systematic review. *Clin Microbiol Infect.* 2017;23:916–923.
- Gea-Banacloche JC. Rituximab-associated infections. *Semin Hematol.* 2010;47:187–198.
- Dendle C, Gilbertson M, Korman TM, et al. Disseminated enteroviral infection associated with obintuzumab. *Emerg Infect Dis.* 2015;21:1661–1663.
- Pérez-Jacoste Asín MA, Fernández-Ruiz M, López-Medrano F, et al. Bacillus Calmette-Guérin (BCG) infection following intravesical BCG administration as

- adjuvant therapy for bladder cancer: incidence, risk factors, and outcome in a single-institution series and review of the literature. *Medicine (Baltimore)*. 2014;93:236–254.
67. Sha HH, Wang DD, Yan DL, et al. Chimeric antigen receptor T-cell therapy for tumour immunotherapy. *Biosci Rep*. 2017;37:pii: BSR20160332.
 68. Fitzgerald JC, Weiss SL, Maude SL, et al. Cytokine release syndrome after chimeric antigen receptor T cell therapy for acute lymphoblastic leukemia. *Crit Care Med*. 2017;45:e124–e131.
 69. Hill JA, Li D, Hay KA, et al. Infectious complications of CD19-targeted chimeric antigen receptor-modified T-cell immunotherapy. *Blood*. 2018;131:121–130.
 70. Capone A, Valle M, Proietti F, et al. Postoperative infections in cytoreductive surgery with hyperthermic intraperitoneal intraoperative chemotherapy for peritoneal carcinomatosis. *J Surg Oncol*. 2007;96:507–513.
 71. Sugarbaker PH, Alderman R, Edwards G, et al. Prospective morbidity and mortality assessment of cytoreductive surgery plus perioperative intraperitoneal chemotherapy to treat peritoneal dissemination of appendiceal mucinous malignancy. *Ann Surg Oncol*. 2006;13:635–644.
 72. Glehen O, Kwiatkowski F, Sugarbaker PH, et al. Cytoreductive surgery combined with perioperative intraperitoneal chemotherapy for the management of peritoneal carcinomatosis from colorectal cancer: a multi-institutional study. *J Clin Oncol*. 2004;22:3284–3292.
 73. Mortenson MM, Schneider PD, Khatri VP, et al. Immediate breast reconstruction after mastectomy increases wound complications: however, initiation of adjuvant chemotherapy is not delayed. *Arch Surg*. 2004;139:988–991.
 74. Penel N, Yazdanpanah Y, Chauvet M, et al. Prevention of surgical site infection after breast cancer surgery by targeted prophylaxis antibiotic in patients at high risk of surgical site infection. *J Surg Oncol*. 2007;96:124–129.
 75. Li Sol Y, Kim CW, Jeon UB, et al. Early infectious complications of percutaneous metallic stent insertion for malignant biliary obstruction. *AJR Am J Roentgenol*. 2010;194:261–265.
 76. Uchiyama K, Ueno M, Ozawa S, et al. Risk factors for postoperative infectious complications after hepatectomy. *J Hepatobiliary Pancreat Sci*. 2011;18:67–73.
 77. Kaibori M, Ishizak IM, Matsui K, et al. Postoperative infectious and non-infectious complications after hepatectomy for hepatocellular carcinoma. *Hepatogastroenterology*. 2011;58:1747–1756.
 78. Konishi T, Watanabe T, Kishimoto J, et al. Elective colon and rectal surgery differ in risk factors for wound infection: results of prospective surveillance. *Ann Surg*. 2006;244:758–763.
 79. Morris CD, Sepkowitz K, Fonshell C, et al. Prospective identification of risk factors for wound infection after lower extremity oncologic surgery. *Ann Surg Oncol*. 2003;10:778–782.
 80. Shiono S, Yoshida J, Nishimura M, et al. Risk factors of postoperative respiratory infections in lung cancer surgery. *J Thorac Oncol*. 2007;2:34–38.
 81. Horasan ES, Ersoz G, Horoz M, et al. Risk factors for infections caused by multidrug-resistant bacteria in patients with solid tumours. *Scand J Infect Dis*. 2011;43:107–111.
 82. Anatoliotaki M, Valatas V, Mantadakis E, et al. Bloodstream infections in patients with solid tumors: associated factors, microbial spectrum and outcome. *Infection*. 2004;32:65–71.
 83. Mikulska M, Del Bono V, Raiola AM, et al. Blood stream infections in allogeneic hematopoietic stem cell transplant recipients: reemergence of gram-negative rods and increasing antibiotic resistance. *Biol Blood Marrow Transplant*. 2009;15:47–53.
 84. Mikulska M, Viscoli C, Orasch C, et al. Fourth European Conference on Infections in Leukemia Group (ECIL-4), a joint venture of EBMT, EORTC, ICHS, ELN and ESGICH/ESCMID. Aetiology and resistance in bacteraemias among adult and paediatric haematology and cancer patients. *J Infect*. 2014;68:321–331.
 85. ECDC European Centre for Disease Prevention and Control. Surveillance of antimicrobial resistance in Europe 2016. Annual Report of the European Antimicrobial Resistance Surveillance Network (EARS-Net). Stockholm: ECDC; 2017. <https://ecdc.europa.eu/en/publications-data/antimicrobial-resistance-surveillance-europe-2016>. Accessed March 16, 2018.
 86. Averbuch D, Tridello G, Hoek J, et al. Antimicrobial resistance in gram-negative rods causing bacteremia in hematopoietic stem cell transplant recipients: intercontinental prospective study of the Infectious Diseases Working Party of the European Bone Marrow Transplantation Group. *Clin Infect Dis*. 2017;65:1819–1828.
 87. Vehreschild MJ, Hamprecht A, Peterson L, et al. A multicentre cohort study on colonization and infection with ESBL-producing Enterobacteriaceae in high-risk patients with haematological malignancies. *J Antimicrob Chemother*. 2014;69:3387–3392.
 88. Giacobbe DR, Del Bono V, Trecarichi EM, et al. Risk factors for bloodstream infections due to colistin-resistant KPC-producing *Klebsiella pneumoniae*: results from a multicenter case-control study. *Clin Microbiol Infect*. 2015;21:1106.e1–1106.e8.
 89. Weinstock DM, Conlon M, Iovino C, et al. Colonization, bloodstream infection, and mortality caused by vancomycin-resistant enterococcus early after allogeneic hematopoietic stem cell transplant. *Biol Blood Marrow Transplant*. 2007;13:615–621.
 90. Zahar JR, Farhat H, Chachaty E, et al. Incidence and clinical significance of anaerobic bacteraemia in cancer patients: a 6-year retrospective study. *Clin Microbiol Infect*. 2005;11:724–729.
 91. Frattino G, Molinari AC, Parodi S, et al. Central venous catheter-related complications in children with oncological/hematological diseases: an observational study of 418 devices. *Ann Oncol*. 2005;16:648–654.
 92. Jagarlamudi R, Kumar L, Kochupillai V, et al. Infections in acute leukemia: an analysis of 240 febrile episodes. *Med Oncol*. 2000;17:111–116.
 93. Hengartner H, Berger C, Nadal D, et al. Port-A-Cath infections in children with cancer. *Eur J Cancer*. 2004;40:2452–2458.
 94. Maki DG, Kluger DM, Crnich CJ. The risk of bloodstream infection in adults with different intravascular devices: a systematic review of 200 published prospective studies. *Mayo Clin Proc*. 2006;81:1159–1171.
 95. Cheong K, Perry D, Karapetis C, et al. High rate of complications associated with peripherally inserted central venous catheters in patients with solid tumours. *Intern Med J*. 2004;34:234–238.
 96. Harter C, Ostendorf T, Bach A, et al. Peripherally inserted central venous catheters for autologous blood progenitor cell transplantation in patients with haematological malignancies. *Support Care Cancer*. 2003;11:790–794.
 97. Abedin S, Kapoor G. Peripherally inserted central venous catheters are a good option for prolonged venous access in children with cancer. *Pediatr Blood Cancer*. 2008;51:251–255.
 98. Wolf HH, Leithäuser M, Maschmeyer G, et al. Central venous catheter-related infections in hematology and oncology: guidelines of the Infectious Diseases Working Party (AGIHO) of the German Society of Hematology and Oncology (DGHO). *Ann Hematol*. 2008;87:863–876.
 99. Gorschlüter M, Hahn C, Ziske C, et al. Low frequency of enteric infections by *Salmonella*, *Shigella*, *Yersinia* and *Campylobacter* in patients with acute leukemia. *Infection*. 2002;30:22–25.
 100. Kamboj M, Son C, Cantu S, et al. Hospital-onset *Clostridium difficile* infection rates in persons with cancer or hematopoietic stem cell transplant: a C3IC network report. *Infect Control Hosp Epidemiol*. 2012;33:1162–1165.
 101. Libshitz HI, Pannu HK, Elting LS, et al. Tuberculosis in cancer patients: an update. *J Thorac Imaging*. 1997;12:41–46.
 102. Viscoli C, Girmenia C, Marinus A, et al. Candidemia in cancer patients: a prospective, multicenter surveillance study by the Invasive Fungal Infection Group (IFIG) of the European Organization for Research and Treatment of Cancer (EORTC). *Clin Infect Dis*. 1999;28:1071–1079.
 103. Cornely OA, Gachot B, Akan H, et al. Epidemiology and mortality of fungemia in cancer patients—a clinical cohort of the Infectious Diseases Group (IDG) of the European Organization for Research and Treatment of Cancer (EORTC 65031) [oral presentation O109]. In: *Program and Abstracts of the 22nd European Congress of Clinical Microbiology and Infectious Diseases*. London, UK: European Society of Clinical Microbiology and Infectious Diseases; 2012.
 104. Alexander BD, Johnson MD, Pfeiffer CD, et al. Increasing echinocandin resistance in *Candida glabrata*: clinical failure correlates with presence of FKS mutations and elevated minimum inhibitory concentrations. *Clin Infect Dis*. 2013;56:1724–1732.
 105. Lamoth F, Kontoyiannis DP. The *Candida auris* alert: facts and perspectives. *J Infect Dis*. 2018;217:516–520.
 106. Pagano L, Cairn M, Candoni A, et al. The epidemiology of fungal infections in patients with hematologic malignancies: the SEIFEM-2004 study. *Haematologica*. 2006;9:1068–1075.
 107. Lortholary O, Gangneux JP, Sitbon K, et al. Epidemiological trends in invasive aspergillosis in France: the SAIF network (2005–2007). *Clin Microbiol Infect*. 2011;17:1882–1889.
 108. Cornely OA, Leguay T, Maertens J, et al. Randomized comparison of liposomal amphotericin B versus placebo to prevent invasive mycoses in acute lymphoblastic leukaemia. *J Antimicrob Chemother*. 2017;72:2359–2367.
 109. Kontoyiannis DP, Marr KA, Park BJ, et al. Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001–2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis*. 2010;50:1091–1100.
 110. Verweij PE, Chowdhary A, Melchers WJ, et al. Azole resistance in *Aspergillus fumigatus*: can we retain the clinical use of mold-active antifungal azoles? *Clin Infect Dis*. 2016;62:362–368.
 111. van der Linden JWM, Arendrup MC, Warris A, et al. Prospective multicenter international surveillance of azole resistance in *Aspergillus fumigatus*. *Emerg Infect Dis*. 2015;21:1041–1044.
 112. Alanio A, Denis B, Hamane S, et al. New therapeutic strategies for invasive aspergillosis in the era of azole resistance: how should the prevalence of azole resistance be defined? *J Antimicrob Chemother*. 2016;71:2075–2078.
 113. Camps SM, van der Linden JW, Li Y, et al. Rapid induction of multiple resistance mechanisms in *Aspergillus fumigatus* during azole therapy: a case study and review of the literature. *Antimicrob Agents Chemother*. 2012;56:10–16.
 114. Corzo-León DE, Satlin MJ, Soave R, et al. Epidemiology and outcomes of invasive fungal infections in allogeneic hematopoietic stem cell transplant recipients in the era of antifungal prophylaxis: a single-centre study with focus on emerging pathogens. *Mycoses*. 2015;58:325–336.
 115. Obeid KM, Aguilar J, Szpunar S, et al. Risk factors for *Pneumocystis jirovecii* pneumonia in patients with lymphoproliferative disorders. *Clin Lymphoma Myeloma Leuk*. 2012;12:66–69.
 116. Cordonnier C, Cesaro S, Maschmeyer G, et al. *Pneumocystis jirovecii* pneumonia: still a concern in patients with haematological malignancies and stem cell transplant recipients. *J Antimicrob Chemother*. 2016;71:2379–2385.
 117. Han XY. Epidemiologic analysis of reactivated cytomegalovirus antigenemia in patients with cancer. *J Clin Microbiol*. 2007;45:1126–1132.
 118. Medicines and Healthcare products Regulatory Agency. Drug Safety Update volume 10 issue 11, July 2017: 2. <https://www.gov.uk/drug-safety-update/bendamustine-levact-increased-mortality-observed-in-recent-clinical-studies-in-off-label-use-monitor-for-opportunistic-infections-hepatitis-b-reactivation#further-information>. Accessed March 16, 2018.
 119. Ljungman P, de la Camara R, Cordonnier C, et al. Management of CMV, HHV-6, HHV-7 and Kaposi-sarcoma herpesvirus (HHV-8) infections in patients with hematological malignancies and after SCT. *Bone Marrow Transplant*. 2008;4:227–240.
 120. Ng AP, Worth L, Chen L, et al. Cytomegalovirus DNAemia and disease: incidence, natural history and management in settings other than allogeneic stem cell transplantation. *Haematologica*. 2005;90:1672–1679.
 121. Hirsch HH, Martino R, Ward KN, et al. Fourth European Conference on Infections in Leukemia (ECIL-4): guidelines for diagnosis and treatment of human respiratory syncytial virus, parainfluenza virus, metapneumovirus, rhinovirus, and coronavirus. *Clin Infect Dis*. 2013;56:258–266.
 122. Chemaly RF, Ghosh S, Bodey GP, et al. Respiratory viral infections in adults with hematologic malignancies and human stem cell transplantation recipients: a retrospective study at a major cancer center. *Medicine (Baltimore)*. 2006;85:278–287.
 123. Kamar N, Izopet J, Tripon S, et al. Ribavirin for chronic hepatitis E virus infection in transplant recipients. *N Engl J Med*. 2014;370:1111–1120.
 124. Tavittan S, Peron JM, Huguet F, et al. Ribavirin for chronic hepatitis prevention among patients with hematologic malignancies. *Emerg Infect Dis*. 2015;21:1466–1469.
 125. Domanović D, Tedder R, Blümel J, et al. Hepatitis E and blood donation safety in selected European countries: a shift to screening? *Euro Surveill*. 2017;22:30514.
 126. Gafter-Gvili A, Fraser A, Paul M, et al. Antibiotic prophylaxis for bacterial infections in afebrile neutropenic patients following chemotherapy. *Cochrane Database Syst Rev*. 2012;(1):CD004386.
 127. Leibovici L, Paul M, Cullen M, et al. Antibiotic prophylaxis in neutropenic patients: new evidence, practical decisions. *Cancer*. 2006;107:1743–1751.
 128. Mikulska M, Averbuch D, Tissot F, et al. Fluoroquinolone prophylaxis in hematological cancer patients with neutropenia: ECIL critical appraisal of previous guidelines. *J Infect*. 2018;76:20–37.
 129. Mikulska M, Cordonnier C. Fluoroquinolone prophylaxis during neutropenia: what can we expect nowadays? *Clin Microbiol Infect*. 2018;pii:S1198-743X(18)30208-8.

130. Trecarichi EM, Tumbarello M, Caira M, et al. Multidrug resistant *Pseudomonas aeruginosa* bloodstream infection in adult patients with hematologic malignancies. *Haematologica*. 2011;96:e1–e3.
131. Girmenia C, Bertina A, Piciocchi A, et al. Incidence, risk factors and outcome of pre-engraftment gram-negative bacteremia after allogeneic and autologous hematopoietic stem cell transplantation: an Italian prospective multicenter survey. *Clin Infect Dis*. 2017;65:1884–1896.
132. Mikulska M, Raiola AM, Galaverna F, et al. Pre-engraftment bloodstream infections after allogeneic hematopoietic cell transplantation: impact of t cell-replete transplantation from a haploidentical donor. *Biol Blood Marrow Transplant*. 2018;24:109–118.
133. Slavin MA, Lingaratnam S, Mileskin L, et al. Use of antibacterial prophylaxis for patients with neutropenia. Australian Consensus Guidelines 2011 Steering Committee. *Intern Med J*. 2011;41:102–109.
134. Klastersky J, de Naurois J, Rolston K, et al. Management of febrile neutropenia: ESMO Clinical Practice Guidelines. *Ann Oncol*. 2016;27:v111–v118.
135. Afsharian P, Möllgård L, Hassan Z, et al. The effect of ciprofloxacin on cyclophosphamide pharmacokinetics in patients with non-Hodgkin lymphoma. *Eur J Haematol*. 2005;75:206–211.
136. Castagnola E, Boni L, Giacchino M, et al. A multicenter, randomized, double blind placebo-controlled trial of amoxicillin/clavulanate for the prophylaxis of fever and infection in neutropenic children with cancer. *Pediatr Infect Dis J*. 2003;22:359–365.
137. Laoprasopwattana K, Khwanna T, Suwankeeree P, et al. Ciprofloxacin reduces occurrence of fever in children with acute leukemia who develop neutropenia during chemotherapy. *Pediatr Infect Dis J*. 2013;32:e94–e98.
138. Wolf J, Tang L, Flynn PM, et al. Levofloxacin prophylaxis during induction therapy for pediatric acute lymphoblastic leukemia. *Clin Infect Dis*. 2017;65:1790–1798.
139. Girmenia C, Rossolini GM, Piciocchi A, et al. Infections by carbapenem-resistant *Klebsiella pneumoniae* in SCT recipients: a nationwide retrospective survey from Italy. *Bone Marrow Transplant*. 2015;50:282–288.
140. Bilinski J, Grzesiowski P, Sorensen N, et al. Fecal microbiota transplantation in patients with blood disorders inhibits gut colonization with antibiotic-resistant bacteria: results of a prospective, single-center study. *Clin Infect Dis*. 2017;65:364–370.
141. Hernández-Cruz B, Ponce-de-León-Rosales S, Sifuentes-Osorio J, et al. Tuberculosis prophylaxis in patients with steroid treatment and systemic rheumatic diseases. A case-control study. *Clin Exp Rheumatol*. 1999;17:81–87.
142. Sterling TR, Villarino ME, Borisov AS, et al. Three months of rifapentine and isoniazid for latent tuberculosis infection. *N Engl J Med*. 2012;365:2155–2166.
143. Lo Vecchio A, Schaffzin JK, Ruberto E, et al. Reduced central line infection rates in children with leukemia following caregiver training: a quality improvement study. *Medicine (Baltimore)*. 2016;95:e3946.
144. Liu Y, Zhang AQ, Cao L, et al. Taurolidine lock solutions for the prevention of catheter-related bloodstream infections: a systematic review and meta-analysis of randomized controlled trials. *PLoS ONE*. 2013;8:e79417.
145. Cornely OA, Maertens J, Winston DJ, et al. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med*. 2007;356:348–359.
146. Cornely OA, Ullmann AJ. Numbers needed to treat with posaconazole prophylaxis to prevent invasive fungal infection and death. *Clin Infect Dis*. 2008;46:1626–1627. author reply 1627–1628.
147. Maertens J, Marchetti O, Herbrecht R, et al. European guidelines for antifungal management in leukemia and hematopoietic stem cell transplant recipients: summary of the ECIL 3–2009 update. *Bone Marrow Transplant*. 2011;46:709–718.
148. Dekkers BGJ, Bakker M, van der Elst KCM, et al. Therapeutic drug monitoring of posaconazole: an update. *Curr Fungal Infect Rep*. 2016;10:51–61.
149. Ullmann AJ, Lipton JH, Vesole DH, et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med*. 2007;356:335–347.
150. Wingard JR, Carter SL, Walsh TJ, et al. Randomized, double-blind trial of fluconazole versus voriconazole for prevention of invasive fungal infection after allogeneic hematopoietic cell transplantation. *Blood*. 2010;116:5111–5118.
151. Marks DI, Pagliuca A, Kibbler CC, et al. Voriconazole versus itraconazole for antifungal prophylaxis following allogeneic hematopoietic stem-cell transplantation. *Br J Haematol*. 2011;15:318–327.
152. Brueggemann RJM, Alfenaar J-WC, Blijlevens NMA, et al. Clinical relevance of the pharmacokinetic interactions of azole antifungal drugs with other coadministered agents. *Clin Infect Dis*. 2009;48:1441–1458.
153. Tissot F, Agrawal S, Pagano L, et al. ECIL-6 guidelines for the treatment of invasive candidiasis, aspergillosis and mucormycosis in leukemia and hematopoietic stem cell transplant patients. *Haematologica*. 2017;102:433–444.
154. Rijnders BJ, Cornelissen JJ, Slobbe L, et al. Aerosolized liposomal amphotericin B for the prevention of invasive pulmonary aspergillosis during prolonged neutropenia: a randomized, placebo-controlled trial. *Clin Infect Dis*. 2008;46:1401–1408.
155. Bochennek KTL, Schedler N, Becker M, et al. Liposomal amphotericin B twice weekly as antifungal prophylaxis in paediatric haematological malignancy patients. *Clin Microbiol Infect*. 2011;17:1868–1874.
156. Cornely OA, Leguay T, Maertens J, et al. AmBiGuard Study Group. Randomized comparison of liposomal amphotericin B versus placebo to prevent invasive mycoses in acute lymphoblastic leukaemia. *J Antimicrob Chemother*. 2017;72:2359–2367.
157. Sipsas NV, Kontoyiannis DP. Clinical issues regarding relapsing aspergillosis and the efficacy of secondary antifungal prophylaxis in patients with hematological malignancies. *Clin Infect Dis*. 2006;42:1584–1591.
158. Reid AD, Chen SC-A, Worth LJ. *Pneumocystis jirovecii* pneumonia in non-HIV-infected patients: new risks and diagnostic tools. *Curr Opin Infect Dis*. 2011;24:534–544.
159. Maertens J, Cesaro S, Maschmeyer G, et al. ECIL guidelines for preventing *Pneumocystis jirovecii* pneumonia in patients with hematological malignancies and stem cell transplant recipients. *J Antimicrob Chemother*. 2016;71:2397–2404.
160. Marty FM, Ljungman P, Chemaly RF, et al. Letermovir prophylaxis for cytomegalovirus in hematopoietic-cell transplantation. *N Engl J Med*. 2017;377:2433–2444.
161. Huang H, Li X, Zhu J, et al. Entecavir vs lamivudine for prevention of hepatitis B virus reactivation among patients with untreated diffuse large B-cell lymphoma receiving R-CHOP chemotherapy: a randomized clinical trial. *JAMA*. 2014;312:2521–2530.
162. Timmer-Bonte JN, Tjan-Heijnen VC. Febrile neutropenia: highlighting the role of prophylactic antibiotics and granulocyte colony-stimulating factor during standard dose chemotherapy for solid tumors. *Anticancer Drugs*. 2006;17:881–889.
163. Wittman B, Horan J, Lyman GH. Prophylactic colony-stimulating factors in children receiving myelosuppressive chemotherapy: a meta-analysis of randomized controlled trials. *Cancer Treat Rev*. 2006;32:298–303.
164. Herbst C, Naumann F, Kruse EB, et al. Prophylactic antibiotics or G-CSF for the prevention of infections and improvement of survival in cancer patients undergoing chemotherapy. *Cochrane Database Syst Rev*. 2009;(1):CD007107.
165. Molica S, Musto P, Chiurazzi F, et al. Prophylaxis against infections with low-dose intravenous immunoglobulins (IVIg) in chronic lymphocytic leukemia. Results of a crossover study. *Haematologica*. 1996;81:121–126.
166. Raanani P, Gafter-Gvili A, Paul M, et al. Immunoglobulin prophylaxis in chronic lymphocytic leukemia and multiple myeloma: systematic review and meta-analysis. *Leuk Lymphoma*. 2009;50:764–772.
167. National Comprehensive Cancer Network (NCCN). NCCN Guidelines for Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. https://www.nccn.org/professionals/physician_gls/default.aspx#age. Accessed May 25, 2018.
168. Gyssens IC, Kern WV, Livermore DM, et al. The role of antibiotic stewardship in limiting antibacterial resistance among hematology patients. *Haematologica*. 2013;98:1821–1825.
169. Siegel JD, Rhinehart E, Jackson M, et al; Healthcare Infection Control Practices Advisory Committee. 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings. <https://www.cdc.gov/infectioncontrol/pdf/guidelines/isolation-guidelines.pdf>. Accessed March 16, 2018.
170. Bénet T, Nicolle MC, Thiebaut A, et al. Reduction of invasive aspergillosis incidence among immunocompromised patients after control of environmental exposure. *Clin Infect Dis*. 2007;45:682–686.
171. Berthelot P, Loulergue P, Raberin H, et al. Efficacy of environmental measures to decrease the risk of hospital-acquired aspergillosis in patients hospitalised in haematology wards. *Clin Microbiol Infect*. 2006;12:738–744.
172. Sonbol MB, Firwana B, Diab M, et al. The effect of a neutropenic diet on infection and mortality rates in cancer patients: a meta-analysis. *Nutr Cancer*. 2015;67:1230–1238.
173. van Dalen EC, Mank A, Leclercq E, et al. Low bacterial diet versus control diet to prevent infection in cancer patients treated with chemotherapy causing episodes of neutropenia. *Cochrane Database Syst Rev*. 2016;(4):CD006247.
174. U.S. Department of Agriculture's Food Safety and Inspection Service and the U.S. Department of Health and Human Services' Food and Drug Administration. September 2006 Slightly revised September 2011. <https://www.fda.gov/downloads/Food/FoodborneIllnessContaminants/UCM312793.pdf>. Accessed March 16, 2018.
175. Rubin LG, Levin MJ, Ljungman P, et al. 2013 IDSA clinical practice guideline for vaccination of the immunocompromised host. *Clin Infect Dis*. 2014;58:309–318.
176. Guidelines for vaccination of patients with hematological malignancies and HSCT recipients. September 2017. <http://www.ecil-leukaemia.com/telechargements/ECIL%207%20Vaccine%20Part%20I%20and%20II%20Final.pdf>. Accessed March 16, 2018.
177. Timitilli A, Bertoluzzo L, Micalizzi C, et al. Anti-varicella zoster vaccination in contacts of children receiving antineoplastic chemotherapy: a prospective pilot study. *Infect Med*. 2008;3:144–147.
178. Viscoli C. Management of infection in cancer patients. studies of the EORTC International Antimicrobial Therapy Group (IATG). *Eur J Cancer*. 2002;38:S82–S87.
179. Klastersky J, Ayme L, Maertens J, et al. Bacteraemia in febrile neutropenic cancer patients. *Int J Antimicrob Agents*. 2007;30:S51–S59.
180. Cattaneo C, Casari S, Bracchi F, et al. Recent increase in enterococci, viridans streptococci, *Pseudomonas* spp. and multiresistant strains from hematological patients, with a negative impact on outcome. Results of a 3-year surveillance study at a single institution. *Scand J Infect Dis*. 2010;42:324–332.
181. Averbuch D, Orasch C, Cordonnier C, et al. European guidelines for empirical antibacterial therapy for febrile neutropenic patients in the era of growing resistance: summary of the 2011 4th European Conference on Infections in Leukemia. *Haematologica*. 2013;98:1826–1835.
182. Lehnbecher T, Phillips R, Alexander S, et al. Guideline for the management of fever and neutropenia in children with cancer and/or undergoing hematopoietic stem-cell transplantation. *J Clin Oncol*. 2012;30:4427–4438.
183. Freifeld AG, Bow EJ, Sepkowitz KA, et al. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2011;52:e56–e93.
184. Kern WV, Marchetti O, Drgona L, et al. Oral antibiotics for fever in low-risk neutropenic patients with cancer: a double-blind, randomized, multicenter trial comparing single daily moxifloxacin with twice daily ciprofloxacin plus amoxicillin/clavulanic acid combination therapy—EORTC Infectious Diseases Group Trial XV. *J Clin Oncol*. 2013;31:1149–1156.
185. Boucher HW, Wilcox M, Talbot GH, et al. Once-weekly dalbavancin versus daily conventional therapy for skin infection. *N Engl J Med*. 2014;370:2169–2179.
186. Castagnola E, Mikulska M, Viscoli C. Prophylaxis and empirical therapy of infection in cancer patients. In: Mandell GL, Bennett JE, Dolin R, eds. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*. 7th ed. Philadelphia: Churchill Livingstone; 2010:3793–3807.
187. Averbuch D, Orasch C, Mikulska M, et al. Reply to "Comparison of antipseudomonal beta-lactams for febrile neutropenia empiric therapy: systematic review and network meta-analysis" by Horita et al. *Clin Microbiol Infect*. 2018;pii: S1198-743X(18)30066-1.
188. Drgona L, Paul M, Bucaneve G, et al. The need for aminoglycosides in combination with β -lactams for high-risk, febrile neutropenic patients with leukaemia. *EJC Supplements*. 2007;5:13–22.
189. Tumbarello M, Sali M, Trecarichi EM, et al. Bloodstream infections caused by extended-spectrum-beta-lactamase-producing *Escherichia coli*: risk factors for inadequate initial antimicrobial therapy. *Antimicrob Agents Chemother*. 2008;52:3244–3252.
190. Castagnola E, Caviglia I, Pescetto L, et al. Antibiotic susceptibility of gram-negatives isolated from bacteremia in children with cancer. Implications for empirical therapy of febrile neutropenia. *Future Microbiol*. 2015;10:357–364.
191. Mokart D, Slehoffer G, Lambert J, et al. De-escalation of antimicrobial treatment in neutropenic patients with severe sepsis: results from an observational study. *Intensive Care Med*. 2014;40:41–49.
192. Snyder M, Pasikhova Y, Baluch A. Early antimicrobial de-escalation and stewardship in adult hematopoietic stem cell transplantation recipients: retrospective review. *Open Forum Infect Dis*. 2017;4:ofx226.

193. Gustinetti G, Raiola A, Varaldo R, et al. De-escalation and discontinuation of empirical antibiotic treatment in a cohort of allogeneic hematopoietic stem cell transplantation recipients during the pre-engraftment period. *Biol Blood Marrow Transplant*. 2018;pii: S1083-8791(18)30131-9.
194. Pizzo PA, Robichaud KJ, Gill FA, et al. Duration of empiric antibiotic therapy in granulocytopenic patients with cancer. *Am J Med*. 1979;67:194–200.
195. Schimpff SC, Young V, Greene E, et al. Origin of Infection in acute non-lymphocytic leukemia: significance of hospital acquisition of potential pathogens. *Ann Intern Med*. 1972;77:707–715.
196. Harris VC, Haak BW, Boele van Hensbroek M, et al. The intestinal microbiome in infectious diseases: the clinical relevance of a rapidly emerging field. *Open Forum Infect Dis*. 2017;4:ofx144.
197. DiNubile MJ. Stopping antibiotic therapy in neutropenic patients. *Ann Intern Med*. 1988;108:289–292.
198. Santolaya ME, Villarreal M, Avendano LF, et al. Discontinuation of antimicrobial therapy for febrile, neutropenic children with cancer: a prospective study. *Clin Infect Dis*. 1997;25:92–97.
199. Klaassen RJ, Allen U, Doyle JJ. Randomized placebo-controlled trial of oral antibiotics in pediatric oncology patients at low-risk with fever and neutropenia. *J Pediatr Hematol Oncol*. 2000;22:405–411.
200. Aquino VM, Tkaczewski I, Buchanan GR. Early discharge of low-risk febrile neutropenic children and adolescents with cancer. *Clin Infect Dis*. 1997;25:74–78.
201. Slobbe L, Waal L, Jongman LR, et al. Three-day treatment with imipenem for unexplained fever during prolonged neutropenia in haematology patients receiving fluoroquinolone and fluconazole prophylaxis: a prospective observational safety study. *Eur J Cancer*. 2009;45:2810–2817.
202. Micol JB, Chahine C, Woerther PL, et al. Discontinuation of empirical antibiotic therapy in neutropenic acute myeloid leukaemia patients with fever of unknown origin: is it ethical? *Clin Microbiol Infect*. 2014;20:O453–O455.
203. Orasch C, Averbuch D, Mikulska M, et al. Discontinuation of empirical antibiotic therapy in neutropenic leukaemia patients with fever of unknown origin is ethical. *Clin Microbiol Infect*. 2015;21:e25–e27.
204. Aguilar-Guisado M, Espigado I, Martín-Peña A, et al. Optimisation of empirical antimicrobial therapy in patients with haematological malignancies and febrile neutropenia (How Long study): an open-label, randomised, controlled phase 4 trial. *Lancet Haematol*. 2017;4:e573–e583.
205. Goldberg E, Gafter-Gvili A, Robenshtok E, et al. Empirical antifungal therapy for patients with neutropenia and persistent fever: systematic review and meta-analysis. *Eur J Cancer*. 2008;44:2192–2203.
206. Bodey G, Buelmann B, Duguid W, et al. Fungal infections in cancer patients: an international autopsy survey. *Eur J Clin Microbiol Infect Dis*. 1992;11:99–109.
207. Beck-Sagué C, Jarvis WR. Secular trends in the epidemiology of nosocomial fungal infections in the United States, 1980–1990. National Nosocomial Infections Surveillance System. *J Infect Dis*. 1993;167:1247–1251.
208. Groll AH, Shah PM, Mentzel C, et al. Trends in the postmortem epidemiology of invasive fungal infections at a university hospital. *J Infect*. 1996;33:23–32.
209. Maertens J, Theunissen K, Verhoef G, et al. Galactomannan and computed tomography-based preemptive antifungal therapy in neutropenic patients at high risk for invasive fungal infection: a prospective feasibility study. *Clin Infect Dis*. 2005;41:1242–1250.
210. Cordonnier C, Pautas C, Maury S, et al. Empirical versus preemptive antifungal therapy for high-risk, febrile, neutropenic patients: a randomized, controlled trial. *Clin Infect Dis*. 2009;48:1042–1051.
211. Girmenia C, Micozzi A, Gentile G, et al. Clinically driven diagnostic antifungal approach in neutropenic patients: a prospective feasibility study. *J Clin Oncol*. 2010;28:667–674.
212. Duarte RF, Sánchez-Ortega I, Cuesta I, et al. Serum galactomannan-based early detection of invasive aspergillosis in hematology patients receiving effective antimold prophylaxis. *Clin Infect Dis*. 2014;59:1696–1702.
213. Maertens JA, Raad II, Marr KA, et al. Isavuconazole versus voriconazole for primary treatment of invasive mould disease caused by *Aspergillus* and other filamentous fungi (SECURE): a phase 3, randomised-controlled, non-inferiority trial. *Lancet*. 2016;387:760–769.
214. Marty FM, Ostrosky-Zeichner L, Cornely OA, et al. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect Dis*. 2016;16:828–837.
215. Verweij PE, Ananda-Rajah M, Andes D, et al. International expert opinion on the management of infection caused by azole-resistant *Aspergillus fumigatus*. *Drug Resist Updat*. 2015;21:22:30–40.
216. Norris LB, Kablaoui F, Brillhart MK, et al. Systematic review of antimicrobial lock therapy for prevention of central-line-associated bloodstream infections in adult and pediatric cancer patients. *Int J Antimicrob Agents*. 2017;50:308–317.
217. Castagnola E, Ginocchio F. Rescue therapy of difficult-to-treat indwelling central venous catheter-related bacteremias in cancer patients: a review for practical purposes. *Expert Rev Anti Infect Ther*. 2013;11:179–186.
218. Mermel LA, Allon M, Bouza E, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;49:1–45.
219. Caillot D, Couaillier J, Bernard A, et al. Increasing volume and changing characteristics of invasive pulmonary aspergillosis on sequential thoracic computed tomography scans in patients with neutropenia. *J Clin Oncol*. 2001;19:253–259.
220. van de Wetering MD, Weggelaar N, Offringa M, et al. Granulocyte transfusions in neutropenic children: a systematic review of the literature. *Eur J Cancer*. 2007;42:2082–2092.
221. West KA, Gea-Banacloche J, Stroneck D, et al. Granulocyte transfusions in the management of invasive fungal infections. *Br J Haematol*. 2017;177:357–374.
222. Price TH, Boeckh M, Harrison RW, et al. Efficacy of transfusion with granulocytes from G-CSF/dexamethasone-treated donors in neutropenic patients with infection. *Blood*. 2015;126:2153–2161.
223. Kadri SS, Remy KE, Strich JR, et al. Role of granulocyte transfusions in invasive fusariosis: systematic review and single-center experience. *Transfusion*. 2015;55:2076–2085.
224. Berghmans T, Paesmans M, Lafitte JJ, et al. Therapeutic use of granulocyte and granulocyte-macrophage colony-stimulating factors in febrile neutropenic cancer patients. A systematic review of the literature with meta-analysis. *Support Care Cancer*. 2002;10:181–188.
225. Wirth S, Emil SGS, Engeli A, et al. Moxifloxacin in pediatric patients with complicated intra-abdominal infections: results of the MOXIPEDIA randomized controlled study. *Pediatr Infect Dis J*. 2018;37:e207–e213.
226. Döring M, Cabanillas Stanchi KM, Queudeville M, et al. Efficacy, safety and feasibility of antifungal prophylaxis with posaconazole tablet in paediatric patients after haematopoietic stem cell transplantation. *J Cancer Res Clin Oncol*. 2017;143:1281–1292.
227. Mesini A, Faraci M, Giardino S, et al. Alternate day dosing of posaconazole tablets in children leads to efficient plasma levels. *Eur J Haematol*. 2018.
228. NAN ALERT. Risk of serious or fatal medication error. June 29, 2011. Available from: <https://www.nccmerp.org/sites/default/files/NANAlertJune2011.pdf>. Accessed March 16, 2018.
229. Delfino L, Puntillo F, Mosca A, et al. High dose, extended interval colistin administration in critically ill patients: is high dosing the right strategy? A preliminary study. *Clin Infect Dis*. 2012;54:1720–1726.
230. Plachouras D, Karvanen M, Friberg LE, et al. Population pharmacokinetic analysis of colistin methanesulfonate and colistin after intravenous administration in critically ill patients with infections caused by gram-negative bacteria. *Antimicrob Agents Chemother*. 2009;53:3430–3436.
231. Durante-Mangoni E, Signorile G, Andini R, et al. Colistin and rifampicin with colistin alone for the treatment of serious infections due to extensively drug-resistant *Acinetobacter baumannii*: a multicenter, randomized clinical trial. *Clin Infect Dis*. 2013;57:349–358.

Infections in Recipients of Hematopoietic Stem Cell Transplants

Jo-Anne H. Young and Celalettin Ustun

SHORT VIEW SUMMARY

Types of Donors of Stem Cells for Hematopoietic Stem Cell Transplantation (HSCT)

- Autologous: The recipient donates to self.
- Syngeneic: The donor is an identical twin sibling.
- Allogeneic: The donated material comes from a different individual than the recipient.

Specific Types of Allogeneic Donors

- Matched related or partially matched related donor, such as a sibling with the same or similar HLA type
- Unrelated donor or matched unrelated donor
- Haploidentical: parent, cousin, sibling, or child is the donor; one HLA haplotype matches. Although haploidentical or other HLA-mismatched transplants may lead to a high incidence of graft-versus-host disease (GVHD), administration of cyclophosphamide early posttransplant and T-cell depletion of the grafts may limit risks of GVHD.
- Cord: umbilical cord blood (UCB) usually partially HLA matched not matched for blood type; sometimes two cords used to provide blood with sufficient cells in adults
- Haplocord: haploidentical peripheral blood stem cells plus cord blood cells; haplocord engrafts rapidly but may not be sustained, yet provides neutrophil production until the cord engrafts

Types of Cells Used in HSCT

- Peripheral blood stem cells (PBSC), usually filgrastim (granulocyte colony-stimulating factor) with or without plerixafor mobilized; may be CD34 selected for T-cell depletion
- UCB: usually associated with delayed engraftment but less GVHD
- Bone marrow: collected by aspiration harvest from matched related or matched unrelated donor; associated with less chronic (c)GVHD compared with PBSC
- Donor lymphocyte infusion: donor cells sorted for lymphocytes; given after engraftment or after relapse for antitumor or antiviral effect but can stimulate GVHD

Preparation of Patient for Cell Infusion ("Conditioning Regimen")

- Myeloablative conditioning: total-body irradiation (TBI) plus cyclophosphamide or busulfan; can comprise chemotherapy alone or

TBI alone to induce immunosuppression in the host and have a good antitumor effect

- Nonmyeloablative or reduced-intensity conditioning: using lower-dose TBI or with antithymocyte globulin, most often with fludarabine to induce adequate immunosuppression to prevent graft rejection by host and has only minimal antitumor effect; success mostly relies on graft-versus-tumor effect

Outcomes

- Engraftment: absolute neutrophil count rises to more than 500 cells/ μ L by day 42
- Primary graft failure: no engraftment by day 42
- Mixed chimerism: coexistence of host and donor hematopoiesis
- Relapse: return of the underlying malignant condition

Complications

- Mucositis: mucosal inflammation that serves as a portal of entry for oral or intestinal infections; common with methotrexate-containing regimens. Palifermin may reduce mucositis
- Hemorrhagic cystitis: often occurs in conjunction with cyclophosphamide early after transplant, despite the use of mesna and forced diuresis; viral causes more common later after transplant, including adenovirus or BK virus
- Engraftment syndrome (immune reconstitution syndrome): fever with or without rash and/or pulmonary infiltrates at engraftment
- Venooclusive disease, also called sinusoidal obstruction syndrome: triad of jaundice, weight gain, and ascites leading to multiorgan failure (mostly in liver, kidney, and lung), usually occurring in the first 3 to 4 weeks after transplantation
- Diffuse alveolar hemorrhage: bleeding into alveoli, usually within the first month after transplantation; diagnosed by progressive bloody returns on lung lavage and treated with very-high-dose corticosteroids
- GVHD: inflammation and cell death (apoptosis) in skin, liver, gut, or lung. Acute form usually occurs until day 100. Chronic form generally occurs later and has features resembling, but distinct from, scleroderma. The risk is higher in the elderly and with grafts from partially matched donors. Calcineurin inhibitors (cyclosporine, tacrolimus), less often

sirolimus, and mycophenolate mofetil are started just before transplantation and continued for at least several months to decrease the incidence and severity of GVHD. High-dose corticosteroids (and other agents) constitute the core of severe GVHD management.

- Posterior reversible encephalopathy syndrome (PRES): neurotoxicity may present with headache, seizures, altered consciousness, and visual disturbance and is associated with typical bilateral white-matter abnormalities in vascular watershed areas in both occipital and parietal lobes. PRES can be resulted from hypertension, calcineurin inhibitors, or fludarabine use.
- Bronchiolitis obliterans organizing pneumonia and obliterative bronchiolitis: pneumonitis associated with small airway injury, sometimes associated with cGVHD

Antimicrobial Agents for Prophylaxis

- Acyclovir: low dose for herpes simplex virus (HSV), high dose for cytomegalovirus (CMV)
- Levofloxacin: used for prevention of bacterial infections, until fevers or infection develop, generally during severe neutropenia or high-dose steroid use
- Fluconazole: for prevention of candidiasis, in general starting at neutropenia and continuing throughout GVHD prophylaxis
- Trimethoprim-sulfamethoxazole: agent of choice for *Pneumocystis*; also covers *Toxoplasma* and some bacteria, such as *Nocardia*. Alternatives include aerosolized pentamidine, atovaquone, and dapsone. In general it is used starting after the first month and continuing throughout the first year or throughout GVHD prophylaxis.
- Penicillin VK: prevention against *Streptococcus pneumoniae* during active cGVHD. Regional penicillin resistance may lead to the use of levofloxacin for this indication.
- Voriconazole or posaconazole: given in place of fluconazole to prevent mold infections
- Echinocandins (micafungin or caspofungin): intravenously administered agents that may be substituted for azoles
- Lamivudine: prevention of hepatitis B virus (HBV) reactivation in anti-hepatitis B core antibody-positive, hepatitis B surface antigen-negative, and hepatitis B DNA-negative patient and/or with hepatitis

SHORT VIEW SUMMARY—cont'd

B–positive donors. Entecavir or tenofovir may be substituted for lamivudine.

- Entecavir or tenofovir: prevention of HBV reactivation in hepatitis B surface antigen–positive patient
- Ivermectin: two doses for patients from countries with high risk for *Strongyloides* infestation

Preemptive Therapy

- Ganciclovir, valganciclovir, or foscarnet: given to patients with CMV reactivation based on polymerase chain reaction (PCR) assay of blood or pp65 neutrophil antigen

Empirical Antibacterial Therapy During Fever and Neutropenia With Negative Workup

- Designated broad-spectrum agent, such as ceftazidime, piperacillin-tazobactam, or cefepime
- Addition of an aminoglycoside possibly required for institutional antibiotic resistance patterns
- Vancomycin if any cellulitis, dysfunction with an indwelling catheter, or hemodynamic

instability or colonized with methicillin-resistant *Staphylococcus aureus*.

Viral Diseases in the Transplant Recipient

- HSV: oral, esophageal, vaginal ulcers; autoinoculation of other skin sites
- Varicella-zoster virus: dermatomal zoster; may disseminate; rarely severe hepatitis or visceral zoster
- CMV: fever, viremia, cytopenias, pneumonitis, gastrointestinal disorders (e.g., hepatitis, esophageal ulcers, mucosal changes in the duodenum or colon), retinitis
- Human herpesvirus 6: fever, rash, viremia, pneumonitis, encephalitis; may be associated with delayed neutrophil recovery or prolonged thrombocytopenia
- Adenovirus: hemorrhagic cystitis, pneumonitis, hepatitis
- BK polyomavirus: hemorrhagic cystitis
- HBV and hepatitis C virus: hepatitis
- Major respiratory viruses: influenza, respiratory syncytial virus in the winter; parainfluenza virus (especially type 3) and rhinovirus year-round

- JC polyomavirus: progressive multifocal leukoencephalopathy, often fatal
- Epstein-Barr virus: viremia or posttransplant lymphoproliferative disorder

Other Infections

- *Clostridioides difficile* (formerly *Clostridium difficile*) diarrhea: stool toxin PCR assay considered the best test
- Intravascular central catheter–related bacteremia or tunnel infection
- Typhlitis: abdominal pain and cecal edema during neutropenia
- *Pneumocystis* infection: diffuse pneumonitis among those not taking prophylaxis
- Candidiasis: candidemia originating from the gut
- Infection with *Aspergillus* and other molds: lung field abnormalities, sinusitis in particular among patients with longer-duration neutropenia, cGVHD, and/or on high-dose steroid therapy
- Toxoplasmosis: brain lesions, can be fatal

The clinical approach to infections in patients undergoing hematopoietic stem cell transplantation (HSCT) involves an understanding of basic transplantation techniques, clinical syndromes, host defense defects at different times after transplantation, the natural history of individual infections, and the mechanisms underlying reconstitution of the immune system after transplantation. In general the dominant elements of infectious risks for bacterial, viral, fungal, and parasitic infections after HSCT depend on the pretransplantation exposure history (viral serostatus), whether the transplant is from an autologous or an allogeneic donor source, the intensity and content of preparative regimens, and the number of days after the transplantation under consideration. The distinguishing determinant of infectious risk between autologous and allogeneic grafts is the associated risk incurred by ongoing immunosuppression from prevention of graft-versus-host disease (GVHD), GVHD itself and its therapy; differing paces of humoral and cellular immune reconstitution that are highly associated with donor type also affect the risk. The time period after transplantation defines eras of differing transplantation complications and the evolution of the slowly resolving posttransplantation immunodeficiency: cutaneous and mucosal barrier breakdown, neutropenia, lymphopenia, hypogammaglobulinemia, or a combination of these. Many posttransplantation complications mimic infectious processes, and multiple infections may occur in the same patient at the same time. Therefore the patient undergoing HSCT should be examined in the context of pretransplantation infections, serologic profiles to document infection latency, intensity and content of “conditioning regimen,” available culture data from mucosal surfaces, contemporaneous transplantation complications in the patient’s institution, current antimicrobial prophylaxis, presence of active GVHD prophylaxis, and the current degree and duration of neutropenia and lymphopenia.

BASIC TRANSPLANTATION TECHNIQUES

HSCT involves the intravenous (IV) delivery of hematopoietic stem cells to a recipient whose hematopoietic and immune systems have been ablated or altered by a cytotoxic and immunosuppressive preparative regimen, commonly referred to as the conditioning regimen, given over the 4 to 10 days before HSCT. Hematopoietic stem cells are obtained from the patient (i.e., autologous) or from other individual(s) (i.e., allogeneic). Graft sources (stem cell collections sites) are bone marrow (BM; mostly

for pediatric transplantations), filgrastim-stimulated peripheral blood (PB), or umbilical cord blood (UCB).^{1,2}

Autologous HSCT is used with the intent of curative or prolongation of survival in the treatment of multiple myeloma, lymphomas, rarely certain leukemias, some high-risk solid tumors (e.g., neuroblastoma, germ cell tumors), and autoimmune disorders (scleroderma). Its efficacy depends purely on (1) conditioning regimen intensity and (2) disease sensitivity to the conditioning regimen. Hematopoietic cell support only makes sure that the recipient will start and continue to reproduce entire blood cell repertoire after myeloablative conditioning (i.e., hematopoiesis). However, its morbidity and mortality also arise from regimen-related toxic effects and early infections.

HSCT is an option with curative potential in treatment of malignancies (e.g., acute and chronic leukemias), preleukemic disorders (e.g., myelodysplastic syndrome), lymphomas, multiple myeloma, hemoglobinopathies or thalassemias, disorders of BM failure (e.g., aplastic anemia, Fanconi anemia), severe immunodeficiency syndromes, and inborn errors of metabolism (e.g., osteopetrosis, chronic granulomatous disease, Hurler syndrome, inherited leukodystrophies, and other lysosomal disorders).^{1,2} It is currently used investigationally in treatment of diseases such as epidermolysis bullosa.^{3,4} Allogeneic HSCT has been feasible for selected human immunodeficiency virus (HIV)-positive patients with malignant and nonmalignant diseases since highly active antiretroviral therapy became standard treatment for HIV infection.⁵

Allogeneic HSCT has the additional hazards of a higher risk of graft failure (prolonging neutropenia) or GVHD compared with autologous HSCT. Substantial improvements in the supportive care of severely immunosuppressed patients have evolved since the 1980s, and survival has markedly improved over time.⁶ As outcomes with transplantation improve, the use of transplantation as an option depends less on the availability of certain donor sources (i.e., whether there is an available matched donor) than in prior decades. Allogeneic donors can be related or unrelated and class I and II human leukocyte antigen (HLA) matched at loci of the major histocompatibility complex or mismatched. Matched unrelated adult volunteer donors (MUD) are available for more than half the general population (unfortunately still problematic for minorities), and unrelated UCB and HLA haploidentical related donors are used increasingly as a source of stem cells for transplantation reported by Center for International Blood and Marrow Transplant Research (CIBMTR) (<https://www.cibmtr.org/ReferenceCenter/SlidesReports/>

[SummarySlides/Pages/index.aspx](#) UCB). UCB has been shown to be an effective alternative for patients who lack a suitable adult donor,⁷ but its use is limited by finding a UCB unit with an adequate cell dose for the recipient's size and weight. Strategies such as transplanting two closely matched or ex vivo expanded UCB units have enabled transplantation for adults.⁸

Graft source may be a contributing factor when considering the likelihood of infectious complications. Reported infectious events before engraftment have been significantly higher with BM versus peripheral blood stem cells (PBSCs) sources, probably related to longer periods of neutropenia before engraftment.⁹ With UCB grafts, engraftment is even further delayed,¹⁰ incurring more infectious complications before engraftment.¹¹ Furthermore, immune recovery, in particular T cells, is also delayed after UCB HSCT.¹² Therefore infections are significantly higher with UCB grafts compared with MUD. In a CIBMTR study the incidences of bacterial infection at 1 year were 72% and 59% for UCB and MUD, respectively. Incidences of viral infection at 1 year were 68% and 45%, ($P < .0001$) for UCB and MUD, respectively.¹¹ Moreover, bacterial and viral, but not fungal, infections were more common after UCB than mismatched unrelated donor (MMUD) ($P = .0009$ and $< .0001$, respectively). In contrast, a French study showed the cumulative incidence of infection (72% vs. 57%) and mortality (8% vs. 3%) at 18 months was higher with MMUD grafts than UCB grafts¹³ due to higher rates of GVHD with PBSC HSCT, which contain a higher T-cell dose.^{9,13,14}

The conditioning regimen used to prepare the host is a major determinant of outcome because of variable host tissue injury and the potential for induction of prolonged immunodeficiency. Conditioning regimens may include immunosuppressive and cytotoxic chemotherapy alone or in combination with wide-field or total-body irradiation (TBI). Conditioning can damage mucosal surfaces, facilitating transmucosal entry of bloodstream infections (BSI).¹⁵ Indwelling IV catheters can also lead to disseminated BSI. Environmental exposure to airborne dusts and fungal spores can inoculate the sinuses or respiratory tree; therefore patients are required to stay in high-efficiency particulate air (HEPA)-filtered rooms. Prophylactic antibiotic therapies and nutritional changes can deplete commensal and potentially protective elements of the cutaneous and intestinal microbiome. The infectious risks for the patient undergoing HSCT are affected by transplantation complications, including the direct effects of this high-dose conditioning, such as mucositis, hemorrhagic cystitis, diarrhea, and hepatic venoocclusive disease (VOD); GVHD; and relapse of the underlying hematologic or oncologic disease.

Chemotherapy

Busulfan and melphalan are commonly used alkylating agents that are toxic to myeloid stem cells and mucosal and epithelial cells. Cyclophosphamide-containing regimens predispose patients to hemorrhagic cystitis. Fludarabine is less cytotoxic but intensely immunosuppressive and is often included in reduced-intensity conditioning (RIC) or nonmyeloablative (NMA) conditioning regimens.¹⁶ Antithymocyte globulin (ATG), which alters the function of or depletes T cells and other lymphocytes, can be used as conditioning regimen for aplastic anemia and, at times, for GVHD prevention or treatment.¹⁷ Serum sickness, a syndrome of fever, arthralgia, and rash, can occur with ATG or other xenoprotein therapy; it is treated with corticosteroid therapy. Antilymphoid antibodies, including alemtuzumab or rituximab affecting T and B cells, respectively, can induce prolonged and profound lymphopenia. Some GVHD prevention strategies include ex vivo graft manipulation (CD34⁺ or CD3⁺ selection) for T-lymphocyte depletion. A new approach is post-HSCT cyclophosphamide (on days 3 and 4 after grafting) to lyse proliferating alloreactive lymphoid cells, highly active in prevention of GVHD, made HLA haploidentical HSCT widely applicable. Cyclosporine, tacrolimus, mycophenolate mofetil, methotrexate, or sirolimus (and other agents) are often used for weeks to months after allografting to lessen the risks and severity of GVHD. Although all these may reduce the risks for GVHD, these maneuvers also substantially delay immune recovery, particularly in haploidentical (one HLA haplotype matched) or HLA-mismatched HSCT.

Irradiation

TBI may be administered as a single dose or more often "fractionated" in multiple doses given over several days. Diarrhea occurs in virtually all treated patients in the first week after irradiation. It may be treated symptomatically while stool studies are pending to rule out infectious causes. Severe oral mucositis occurs in many irradiated patients and is aggravated by prolonged neutropenia and the use of methotrexate for GVHD prophylaxis.¹⁵ As long as bleeding and oral inflammation do not compromise the patient's airway, mucositis is treated symptomatically. Keratinocyte growth factor (delivered clinically as palifermin) has proven activity in the prevention of oral and intestinal mucositis.^{18,19} Its use to limit mucositis may be of benefit in patients receiving highly toxic conditioning, including high-dose TBI, high-dose melphalan, or post-HSCT methotrexate.²⁰ Palifermin has not been shown to reduce GVHD, infections, or mortality.^{21,22} Diffuse alveolar hemorrhage (DAH) is another complication of TBI that is associated with decreased pulmonary function after HSCT.

Reduced-Intensity Conditioning or Nonmyeloablative Transplantation

RIC or NMA regimens have been developed with the goal of permitting donor-derived hematopoietic and immunologic reconstitution.^{23,24} These lesser-intensity regimens may provide a somewhat weaker anticancer effect and rely on the graft-versus-tumor effects to eradicate underlying malignancies. Doses of TBI are usually not more than 2 Gy, compared with 12 to 14 Gy in fully ablative transplants. Fludarabine, ATG, or lower doses of cytotoxic drugs may be used with induction of extended immune suppression.²⁵ This approach is often used for older patients or those with significant medical comorbid conditions. They are intensely immunosuppressive but less cytotoxic, resulting in less mucosal, enteric, and hepatic injury in the early weeks after transplantation, and perhaps because of that associated with fewer risks for GVHD.²⁶

Human Leukocyte Antigen Matching

In general, engraftment is most rapid, and thus neutropenia is briefest, when the patient and allogeneic donor are completely matched at all genetic HLA loci (most often considering only HLA-A, B, C, and DRB1). Similarly, identical twin (syngeneic) transplantations or those in which hematopoietic stem cells collected from the recipient (autologous) are associated with prompt neutrophil recovery. Allogeneic HSCT has the highest chance of prompt engraftment when fully HLA-matched sibling donor transplants are used, but less than 30% of intended recipients have a matched sibling donor available.^{27,28} Greater HLA mismatch augments risks for graft failure, acute and chronic (c)GVHD, and consequent prolonged immunodeficiency.²⁹

Killer immunoglobulin-like receptors (KIR) interact with HLA and other molecules (KIR ligands) that bind to and modulate the function of natural killer (NK) lymphoid cells. After HSCT, altered NK-cell function may modify risks for infection, GVHD, or malignant relapse. KIR ligands can be grouped based on their amino-acid sequence determining the KIR-binding epitopes, primarily in HLA-C and HLA-B molecules, but, owing to linkage disequilibrium, are not inherited coordinately with HLA. T-cell recovery and NK-cell proliferation and functional maturation are not altered by KIR ligand match or mismatch status.³⁰ However, recipients of stem cells from unrelated donors with an activating NK-cell immunoglobulin-like receptor (KIR) (B/x) genotype have decreased infectious complications believed to be due to enhanced NK-cell function.³¹ The role of KIR genotype in donor selection before HSCT, especially for less relapse, is of research interest at this time.

Prevention of Infection

Preventive strategies include protective isolation for reduced exposure to pathogens, enhancement of host immune reconstitution with hematopoietic growth factors, prophylaxis during high-risk periods with targeted antimicrobial chemotherapy, and suppression of subclinical infection with preemptive therapy, which is best facilitated by scheduled periodic surveillance.³²⁻³⁴ Prophylaxis or preemptive strategies are more effective than treatment after infection is established, and the mortality rate among patients with established infections continues to be high despite appropriate therapy. Prophylaxis can be tailored for patients

who have a higher risk of reactivation, therefore screening for lifelong infections, including syphilis, tuberculosis, toxoplasmosis, HIV, hepatitis B virus (HBV), hepatitis C virus (HCV), and herpes simplex virus (HSV) should be tested at screening period.³⁵ Of note, false-positive serum antibody results can occur after IVIG infusions, and false-negative serum antibody results can occur in the early months after transplantation.³⁶ For this reason, viral serostatus, as well as serostatus of a few nonviral infections, are checked before transplantation so that the risk of reactivation is clear.

After mucositis has cleared and oral alimentation has resumed, oral therapy is the preferred route for antiinfective prophylaxis.

CLINICAL SYNDROMES UNIQUE TO THE HEMATOPOIETIC STEM CELL RECIPIENT

Hemorrhagic Cystitis (HC)

HC is a common complication that can lead to gross hematuria, clots, urinary retention, and impairment of renal function. Severe forms (grade III–IV) of HC are associated with increased nonrelapse mortality.³⁷ Cystitis that occurs within the first weeks after marrow infusion usually is noninfectious in origin, caused instead by the administration of cyclophosphamide, especially at myeloablative doses in the conditioning regimen.^{37,38} Prophylactic measures include mesna (for binding and elimination of the alkylator metabolites of cyclophosphamide), forced diuresis, and continuous bladder irrigation. Supportive care for established cystitis may also necessitate large-bore catheter drainage or bladder irrigation and transfusions. Later in the posttransplantation period, severe GVHD, HLA-mismatched status (MMUD, UCB) and infection are contributing causes of cystitis.^{37,39,40} The majority of infectious agents inducing cystitis are viral, usually either the polyomavirus BK virus or adenovirus. Infection with HSV, CMV, the polyomavirus JC virus, human herpesvirus 6 (HHV-6), various bacteria, and *Strongyloides* occurs in lower frequencies.^{41–43} Although data are predominantly derived from studies in renal transplants, there is evidence that quinolone use (levofloxacin) does not have a preventive or therapeutic role for BK virus reactivation and disease.^{44–46} Neither BK viruria nor viremia confirm that BK virus is causal for hemorrhagic cystitis.⁴⁰ Polyomaviruses are shed in the urine in many HSCT patients without clinical symptoms (see Chapter 144).^{47,48} Higher urine or detectable blood viral loads of BK virus may indicate a higher risk for hemorrhagic cystitis.⁴⁹ No standard antiviral treatment is currently available for viruria caused by BK virus or adenovirus, although intravesicular cidofovir or low-dose IV cidofovir has been administered when ongoing hemorrhage or large mucosal clots persist despite continuous bladder irrigation.⁵⁰

Venoocclusive Disease (Sinusoidal Obstruction Syndrome)

VOD is a syndrome of liver toxicity that occurs at any time after the onset of the high-dose conditioning regimen, usually before day 30. It is characterized by painful hepatomegaly, 5% or greater weight gain, and hyperbilirubinemia (bilirubin levels >2 mg/dL).⁵¹ Severe VOD (sometimes called “sinusoidal obstruction syndrome” [SOS]) with marked jaundice or ascites leads to multiorgan failure involving the kidneys, heart, and lungs and high mortality.⁵² The only US Food and Drug Administration (FDA)-approved drug for severe VOD/SOS is defibrotide, which stabilizes endothelial cells by reducing endothelial cell activation and by protecting endothelial cells from further damage, resulting in the restoration of the thrombofibrinolytic balance.⁵³ Anticoagulant or antithrombotic therapies are mostly ineffective and associated with risk of serious bleeding.

Cyclophosphamide followed by TBI increases the risk for VOD because recurrent endothelial damage, along with reduced hepatocyte glutathione, increases vulnerability to radiation toxicity.^{54,55} Other risk factors for VOD include elevated aminotransferases before HSCT, mismatched or unrelated donor HSCT, and history of abdominal radiation therapy.⁵⁵ Genetic predispositions in urea cycle enzymes (e.g., glutathione-S-transferase) may, in some patients, increase risks for VOD.⁵⁴ Conditions that may mimic VOD include cholestasis in patients with septicemia; hepatic infiltration secondary to infection or tumor; pericardial tamponade; cytomegaloviral or other viral hepatitis; and

intraabdominal inflammatory diseases, such as pancreatitis, peritonitis, or cholecystitis. In addition, early GVHD and cyclosporine-induced cholestasis are noninfectious causes of liver toxicity that may coexist with or mimic VOD. Diagnosis of VOD may be difficult in early stages when treatment is more effective, and ultrasonographic assessment of hepatic portal venous flow and newer techniques, such as ultrasound point shear wave elastography, showed shear-wave velocity as significantly elevated in patients with VOD after an HSCT prospective study in 134 patients who received allogeneic HSCT between 2011 and 2016.⁵⁶ Transjugular liver biopsy and measuring corrected hepatic sinusoidal pressure gradient during the procedure with a relatively safe profile (three major complications in 141 HSCT patients) can be important to make correct diagnosis.⁵⁷ Hepatotoxic and nephrotoxic drugs should be avoided in patients with VOD.

Graft-Versus-Host Disease

Acute GVHD (aGVHD) is a major, life-threatening complication, developing in 40% to 80% of patients after allogeneic transplantation.⁵⁸ The risk for developing GVHD is higher in older patients and with partially matched or unrelated donor HSCT and is associated with the higher doses of donor T cells infused with filgrastim-mobilized peripheral blood grafts.⁵⁹ In aGVHD donor, T lymphocytes mount an immune attack against the recipient's tissues that is amplified by proinflammatory cytokines. The favorable antineoplastic impact accompanying GVHD is called the graft-versus-leukemia (GVL) effect.⁶⁰ T cells and possibly NK and other lymphoid cells mediate GVL effects after allogeneic HSCT by the production of inflammatory cytokines and by direct target lysis.

The primary target organs of aGVHD are skin, liver, and gastrointestinal tract (GIT), therefore its common clinical manifestations include rash, cholestatic hepatitis, nausea, vomiting, and diarrhea. aGVHD can be lethal, and one of the major clinical problems remains to predict who will have severe GVHD in real time. Recently, models to predict patients at risk of poor outcomes have been developed using both clinical and laboratory data.⁶¹

Effective immunosuppressive agents for the prevention of GVHD include calcineurin inhibitors (cyclosporine or tacrolimus) with or without methotrexate, mycophenolate mofetil, and sirolimus; they are usually started before transplantation in T-cell replete HSCT. T cells can be removed ex vivo (negative selection or CD34⁺ cell selection) or in vivo (e.g., ATG, alemtuzumab use in conditioning) from the graft; however, these methods may result in an increase of some complications (e.g., graft failure, infections such as CMV and *Aspergillus*, and relapse).^{62,63} In addition, posttransplant cyclophosphamide, used especially after haploidentical HSCT, has become an effective method for prevention of both acute and cGVHD.⁶⁴ GVHD itself can compound and prolong post-HSCT immunodeficiency. The corticosteroids or other immunosuppressive drugs used as treatment of GVHD may impair phagocytic function and directly worsen lymphopenia and cellular immune deficiency.^{65,66} Patients with acute and cGVHD have splenic dysfunction and thus an added risk for infection with encapsulated bacteria, such as *Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae*. T-cell depletion of the donor graft is less associated with GVHD but more with infection and sometimes with much-delayed immune recovery.

Hepatitis

Clinical hepatitis in HSCT recipients can range from fever accompanied by abdominal pain to fulminant illness. Infectious hepatitis must be distinguished from several common noninfectious causes, including liver dysfunction related to the “conditioning regimen” (VOD), aGVHD, cholestatic liver injury related to sepsis, and chemical hepatitis related to either drugs or hyperalimentation.⁶⁷

Clinically important viral hepatitis syndromes that occur after transplantation include acquisition or reactivation of infection with HBV, HCV, varicella-zoster virus (VZV), adenovirus, HSV, CMV, and HHV-6.^{68,69} Reactivation of HBV is more likely than HCV to result in fulminant hepatitis, although this occurs in only a minority of infected patients. Disseminated VZV and adenovirus infections may manifest as elevations in serum aminotransferase levels; these elevations can precede the appearance of other disease manifestations by several days. In rare cases liver biopsy with viral culture and polymerase chain reaction

(PCR) assay are needed to establish a diagnosis of severe hepatitis in the early posttransplantation period. Viruses such as hepatitis G virus and transfusion-transmitted viruses are not known to influence the outcome of HSCT. Hepatitis E virus, transmitted by the fecal-oral route, is now increasingly identified as a pathogen in HSCT, as a cause of both acute and chronic hepatitis.⁷⁰ Screening strategies for donors are not established, and diagnostics should include PCR. Therapy with ribavirin has been effective.⁷¹

HBV (surface antigen [HBsAg], surface antibody [HBsAb], and core antibody [HBcAb]) and HCV serologic profiles are tested in donor and recipient before HSCT. Patients who are HBsAg positive or HBcAb positive are at high risk for reactivation, whereas the risk is extremely rare in those with HBsAb positive. Pretransplantation imaging studies or liver biopsy may be needed to evaluate HCV-seropositive patients with abnormal liver enzyme levels or tender hepatomegaly. Donors and recipients with positivity for HBsAg and HBcAb should be tested for viral load with PCR studies for HBV DNA before transplantation because the risk for HBV hepatitis can be reduced by treatment to lower a detectable viral load. Lamivudine, entecavir, and tenofovir are commonly used to suppress HBV replication (see Chapter 145).⁷² Current stratifications of risk are discussed in Chapter 306. The more potent activity of entecavir has led to support for its use in preference to lamivudine, particularly in patients with HBsAg present in the serum at the time of immunosuppression.⁷³

A transplant from an HBV-infected donor can be used if no alternative donor is available or if the intended recipient is already seropositive.⁷⁴ HBV can be transmitted from an HBsAg-positive (or, less likely, an HBcAb-positive) donor to a recipient who is either HBV naïve or HBsAb positive but HBcAb negative. The risk for transmission is low when an HBV-positive donor has an undetectable viral load and/or has nonreplicative HBV.⁷⁵ If the recipient is HBV naïve before transplantation, the subsequent infection is more likely to have clinical consequences. If the transplant can be delayed, then HBV vaccination of the recipient or use of HBV immune globulin or both may reduce the likelihood of hepatitis after transplantation.⁷⁶ HBV immunity can be transferred from an HBsAb-positive donor to an HBV-naïve recipient.^{69,76} Through adoptive immune transfer, HBV infection can be cleared from an HBsAg-positive recipient by transplant from an HBsAb-positive (i.e., immune) donor, which generally follows a period of hepatitis (elevation of liver functions).⁶⁹

After transplantation the following recipients should be monitored periodically with PCR testing of HBV DNA viral load: (1) those with liver enzyme elevation suggestive of activation of HBV from latency, (2) those with transplants from HBV-infected donors, and (3) those with known infection before transplantation. High HBV viral load ($>10^5$ copies/mL) is the most important risk factor for clinically apparent reactivation in recipients positive for HBsAg.⁷⁷ Among recipients in whom PCR studies of HBV DNA yield positive results persistently after transplantation despite treatment, the risk for fatal liver disease may be up to 12%.⁷⁷

HCV-positive donors and recipients should undergo RNA viral load testing. Although best avoided, but if there are no other options, an HCV-infected individual can serve as a donor. In contrast to HBV, however, the rate of transmission of HCV from an HCV RNA-positive donor approaches 100%. Interferon (IFN) can be used to suppress HCV replication in donors,⁷⁸ but its limited efficacy, systemic and hematologic toxicity, and delayed response may not contain active HCV to render a donor suitable for donation. Direct-acting antiviral (DAA) therapies are highly active in HCV and have rendered IFN-containing regimens obsolete for almost all HCV genotypes, although the evidence comes from non-HSCT patients⁷⁹ (see Chapters 47 and 154). In a study, no patient receiving DAA therapy after transplantation discontinued HCV therapy, and efficacy was 85%.⁸⁰ However, dose adjustments per renal function and drug interactions may be needed.⁷⁶

In contrast to HBV, which induces immune-mediated hepatitis, HCV can induce direct hepatotoxicity.⁸¹ Hepatitis most commonly occurs 2 to 3 months after transplantation; severe hepatitis is rare. There is no strong evidence to support HCV as a cause of increasing incidence of VOD, especially when conditioning regimens are not risk factors for VOD, or GVHD. However, even without direct hepatic failure, HCV seems to be

associated with a higher nonrelapse mortality in the first year. Beyond 10 years, the long-term complication of HCV infection is cirrhosis.⁸² No data have demonstrated a correlation between hepatitis C genotype and type or severity of liver disease after transplantation. Because of the myelosuppressive effects of IFN and other antiviral agents, their use in the treatment of hepatitis C after HSCT is limited. However, in a study with 20-year follow-up, antiviral treatment (IFN with or without ribavirin) showed a strong trend to reduce the risk for severe liver complications (odds ratio, 0.33; 95% confidence interval, 0.11 to 1.03; $P = .058$).⁸³ In the study of 195 patients, 33 died, among which 6 died from liver complications. The cumulative incidence of severe liver complications (death from liver failure, cirrhosis, and liver transplantation) was 11.7% at 20 years after HSCT.⁸³ A new American Society for Blood and Marrow Transplantation task force recommends long-term liver function test (alanine aminotransferase) monitoring in all HCV-positive recipients/donors. HCV RNA monitoring should be measured in all patients at entry into care, and monitoring of viral load should be performed in patients receiving HCV treatment or liver dysfunction.⁷⁹

Pneumonia Syndromes

Infectious pneumonias must be distinguished from noninfectious pulmonary complications after HSCT, which can include pulmonary edema, pleural effusion, alveolar hemorrhage, radiation injury (pneumonitis or fibrosis), drug reactions, adult respiratory distress syndrome, idiopathic pneumonia syndrome, cytolytic thrombi (causing multiple peripheral lung nodules), obliterative bronchiolitis, bronchiolitis obliterans with organizing pneumonia, and cGVHD.⁸⁴ Management of noninfectious pneumonias requires that lower respiratory tract infection (LRTI) be ruled out or recognition of its coexistence. Their pathophysiologic processes may be distinct: Some syndromes may be more likely to respond therapeutically to high-dose corticosteroid therapy.

Diffuse alveolar hemorrhage (DAH) begins with dyspnea and alveolar infiltrates, with progressively bloody return during bronchoscopic examination and alveolar lavage.⁸⁵ Although DAH definition requires no infection, infection-associated alveolar hemorrhage and DAH are related clinical syndromes with similar clinical presentation, risks, and associated high mortality.

The syndrome usually occurs in the second to fourth weeks after HSCT, generally before platelet engraftment and perineutrophil engraftment.⁸⁶ Because DAH occurs around neutrophil engraftment, tissue damage due to leukocyte influx into lungs is suggested.⁸⁷ Delayed engraftment, thrombocytopenia, infection, toxic effects of drugs, TBI, intensely cytotoxic regimens, and solid malignancy have been implicated as risk factors.^{85,86} Very-high-dose corticosteroids are recommended, but DAH remains highly mortal.⁸⁶

Idiopathic pneumonia syndrome is a process of widespread alveolar epithelial injury that is characterized clinically by diffuse interstitial infiltrates and varying degrees of respiratory failure in the absence of active LRTI.⁸⁸ It is believed to be related to the chemotherapy or TBI, or both, used as part of the conditioning regimen, which induces proinflammatory cytokine release and increasing alveolar capillary permeability. Idiopathic pneumonia syndrome occurs in 8% to 17% of patients but may be more frequent after allogeneic than autologous transplantation and thus has been implicated, at least in animal models, as a component of the GVHD reaction. Mortality rates are 60% to 80%. Idiopathic pneumonia syndrome occurs classically during two peaks: one in the first few weeks and the other in the second and third month after transplantation. Corticosteroids and etanercept may yield a clinical response.⁸⁹ More frequent use of sensitive diagnostic techniques has revealed a high incidence of occult infections, including due to human rhinovirus, CMV, HHV-6, and *Aspergillus*.⁸⁹ The detection of these viruses may be associated with increased mortality. This diagnostic yield changes the “precision” of the diagnosis of idiopathic pneumonia, which should lack detectable pathogens.⁹⁰ The uncertain role of this detection on the applicability of immunosuppressive regimens is not yet clear.

Diarrhea

Diarrhea, a common symptom after transplantation, is primarily a result of noninfectious causes, such as regimen-related gut mucosal toxicity and GVHD. Diarrhea is associated with infection in less than 20% of cases.^{91,92}

The list of infectious agents responsible for diarrhea includes *Clostridioides difficile* (formerly *Clostridium difficile*), adenovirus, rotavirus, norovirus, enterovirus, coxsackievirus, HHV-6, *Escherichia coli*, *Salmonella*, *Giardia*, *Strongyloides*, *Cryptosporidium*, and *Campylobacter* spp. Infection with *C. difficile* occurs with increasing frequency.^{93,94} Outbreaks of diarrhea have been reported for *Cryptosporidium* and enterovirus. From other countries, reports of diarrhea have been associated with *Trichostrongylus* spp.

Typhlitis, or neutropenic enterocolitis, is an anaerobic infectious syndrome that is relatively common and may be associated with diarrhea during neutropenia.⁹⁵ Typhlitis is preceded by fever, abdominal pain, and right lower quadrant tenderness that may be accompanied by rebound tenderness. Appendicitis can mimic typhlitis. Computed tomography of the abdomen reveals right-sided colonic enlargement and inflammation with thickening of the mucosa. Therapy against anaerobic bacteria should be added to the medical regimen.

Rash

Skin eruptions are often noninfectious, occurring as a direct result of radiation effect from conditioning therapy or secondary to aGVHD or drug reaction.⁹⁶ Rashes from conditioning regimens can result in the sudden onset of marked erythema over large areas of the body and blistering on the hands and feet. A skin biopsy can assist in distinguishing infectious from noninfectious causes of rash. For all lesions suspected to be infectious, samples should be submitted for culture or biopsy. The most common infectious causes are VZV, catheter-related exit site or tunnel infections, and cutaneous manifestations of disseminated bacterial or fungal infections.⁹⁷ Focal areas of bacterial cellulitis may occur on the lower extremities in the setting of edema from heart failure, VOD, lymphedema, and impaired venous return.

Osteomyelitis

Osteomyelitis is uncommon after HSCT. The spectrum of organisms can include atypical mycobacteria, yeasts, and molds, in addition to bacteria.^{98,99} In rare instances osteomyelitis follows marrow aspiration from the sternum or marrow harvest from the iliac crest.^{99,100} When prolonged pain and fever occur after BM harvest, osteomyelitis caused by *Staphylococcus aureus* should be considered.

PATTERNS OF IMMUNOSUPPRESSION AT DIFFERENT TIMES AFTER MYELOABLATIVE HSCT

Historically, three risk periods of immunologic deficiency occur predictably in graft recipients after HSCT (Fig. 307.1). They are the preengraftment period, the early postengraftment period (until day 100), and the late postengraftment period (after day 100). An understanding of the immune deficiencies in each risk period and the period of peak risk for individual infections that are observed with standard infection prophylaxis helps the clinician recognize uncommon manifestations of these infectious pathogens (Table 307.1).

Preengraftment Risk Period

Infection present in the transplant candidate before conditioning may have important adverse effects after transplantation, so screening and diagnosis of symptomatic respiratory and GI disease is an important intervention. For infections that might resolve, delay of conditioning should be considered to ameliorate the risk associated with conditioning during active infection.^{101,102} The preengraftment risk period begins with the onset of conditioning therapy and continues until approximately days 20 to 40 after transplantation. By definition, graft failure is declared if there is no neutrophil recovery by day 42. Pretransplantation neutropenia is associated with increased infection-related mortality.¹⁰³ Bacterial infections are common during this time of profound neutropenia and lymphopenia, necessitating prophylactic and promptly administered empirical systemic antibiotic therapy (see Chapter 306).¹⁰⁴ Prophylactic systemic antibiotics (often a fluoroquinolone, such as levofloxacin) can be administered when the neutrophil count drops to less than 500/mm³ and continued until the neutrophil count recovers to prevent bacterial infection. GI decontamination with nonabsorbable antibiotics was used in the past but is now rarely performed.

Prophylactic antibiotic use has shifted the spectrum of GI microbiota to potentially pathogenic organisms, such as *C. difficile* and enterococci.^{93,105}

The etiologic agents of bacteremia have shifted to more gram-positive organisms; in particular, coagulase-negative staphylococci (CoNS), enterococci, and viridans group streptococci are often isolated from bloodstream cultures of febrile neutropenic HSCT recipients.¹⁰⁶ Mechanical barrier defects caused by mucositis and central catheters predispose patients to BSI by allowing access for skin-colonizing organisms and GI mucosal flora to otherwise sterile body sites.¹⁰⁷ Colonization with vancomycin-resistant enterococci or other multidrug-resistant pathogens may predispose patients to bacteremia.^{105,108} Recipients of autologous and allogeneic grafts may develop a similar spectrum of infections during the preengraftment period; the major transplantation-related complications occurring in this risk period (mucositis, severe neutropenia, and VOD/SOS) are similarly frequent with all types of transplantations. However, the less frequent use of TBI and methotrexate and the more rapid neutrophil recovery after autologous transplantation of peripheral blood stem cells have markedly decreased the risks for mucositis and serious bacteremia in this subpopulation of patients. Similarly, the use of RIC regimens before NMA allografting has lessened the risks for early bacteremia.

During this risk period, HSV is predictably reactivated in 80% of patients who are HSV seropositive. Most such infections occur before week 4 after transplantation. Prophylactic acyclovir, 400 mg twice daily (5 mg/kg twice daily for children), has minimized this clinical infection.^{109,110}

Candidemia and early-onset aspergillosis occur in less than 5% of patients with neutropenia. The risk is greater in patients with slow engraftment or extended neutropenia before transplantation. With fluconazole (200–400 mg/day),^{111–113} voriconazole,¹¹⁴ or micafungin¹¹⁵ (50 mg IV once daily) prophylaxis, *Candida albicans* infections have been mostly eliminated during this risk period; however, *Candida krusei* and *Candida glabrata* have emerged as fluconazole-resistant pathogens.¹¹⁶

Some allogeneic transplantation candidates are at higher risk for mold infections during the preengraftment period than others; their transplantation course begins with mold prophylaxis, possibly beginning even 2 to 4 weeks before HSCT.¹¹⁷ These patients can include those with extended pre-HSCT neutropenia, such as acute leukemia, who have undergone serial chemotherapy before transplantation; those with myelodysplastic syndrome; those with aplastic or Fanconi anemia; and others. Fungal infection that occurred within 6 to 9 months before transplantation may not be cured and could reactivate. Patients with more remote fungal infections can receive a standard regimen of fungal prophylaxis. If patients have a history of aspergillosis within 4 months of transplantation or have suspect pulmonary nodules without a specific diagnosis,¹¹⁸ they should receive secondary fungal chemoprophylaxis (i.e., ongoing maintenance antifungal therapy) and undergo rescanning before and after transplantation.

Use of hematopoietic growth factors has reduced the incidence of bacteremia by shortening the duration of neutropenia, but these agents have not been shown to improve outcome in established infections.

Adjunctive therapy with granulocyte transfusions has been used in some centers for treatment of serious infections that develop in patients with neutropenia.¹¹⁹ Use of colony-stimulating factors with dexamethasone to prime granulocyte donors and increase the collection yield is being evaluated. Evidence of efficacy and therefore the indications for use of this expensive and labor-intensive supportive measure are uncertain and still under study.

Routine culture of hematopoietic progenitor cell products yields low rates of recovery of bacterial organisms, most often *Corynebacterium* spp. or staphylococci. Appropriate testing of collections of graft products includes routine culture of hematopoietic progenitor cells before HSCT, but patients receiving culture-positive harvests usually do so without clinically adverse outcomes.^{120,121}

Postengraftment Risk Period

The postengraftment period begins with neutrophil recovery and continues until day 100, when early B- and T-lymphocyte functional recovery may be apparent. Reconstituted T lymphocytes have abnormal function for approximately 18 months, as evidenced by CD4 deficiency




FIG. 307.1 Phases of predictable opportunistic infections among patients undergoing hematopoietic stem cell transplantation. Immune defects predisposing to infection are bordered by color (pink, neutropenia; blue, lymphopenia; green, hypogammaglobulinemia). Barrier defects predisposing to infection are shaded in color (gold, mucosal breakdown; gray, skin breakdown). Contribution of defects to infections occurring with high incidence are designated by border color (for immune defects) or shading (for barrier defects) or both. CMV, Cytomegalovirus; GVHD, graft-versus-host disease; HSV, herpes simplex virus; VOD, venoocclusive disease; VZV, varicella-zoster virus. (Modified from Van Burik JA, Freifeld AG. Infection in the severely immunocompromised host. In: Abeloff MD, Armitage JO, Niederhuber JE, et al, eds. Clinical Oncology. 3rd ed. Philadelphia: Churchill Livingstone; 2004:942.)

and by in vitro antigen and mitogen proliferative responses. However, T-lymphocyte reconstitution may be blunted by the effects of GVHD or CMV and their attendant treatments (corticosteroids, calcineurin inhibitors, anti-T-lymphocyte therapy, and ganciclovir products). As a result, the rate of infection during this risk period is higher in recipients of allogeneic grafts, who are more likely to develop GVHD or CMV, than in recipients of autologous-syngeneic grafts. Another consequence of GVHD during this risk period is disruption of the GI mucosa, which can permit transmural entry of pathogens and lead to bacteremia or fungemia.

Late-onset aspergillosis may also occur during this risk period in up to 10% of patients, especially those with continuing GVHD, those receiving high-dose corticosteroids, and those with poor graft function.¹²² Advanced-generation azoles that have activity against filamentous molds are effective prophylaxis against deep mycoses in patients at high risk.^{114,123} Careful surveillance is required for these high-risk groups; serum galactomannan screening is insufficiently sensitive for reliable early detection of *Aspergillus* infections.³² Although infrequent, infection with agents of mucormycosis may occur as they are resistant to the advanced azoles (e.g., voriconazole).

TABLE 307.1 Infections After HSCT in Order of Occurrence

ORGANISM	PEAK TIME PERIOD OF RISK (WEEKS AFTER HSCT)	USUAL PROPHYLAXIS	INCIDENCE
Preengraftment Risk Period (1–4 Weeks)			
Herpes simplex virus (seropositive) ^{109,222–224}	1–2	Acyclovir or valacyclovir	5%–9%
Gram-positive bacteremia (most commonly coagulase-negative staphylococci, viridans group streptococci, and enterococci) ^{105,153,162–164,179–181,184–186}	1–4	Prophylactic broad-spectrum antibiotics	20%–30%
Gram-negative bacteremia ^{104–106,147,164,165,167,169,183,184}	1–4	Prophylactic broad-spectrum antibiotics	5%–10%
<i>Clostridioides difficile</i> (formerly <i>Clostridium difficile</i>) diarrhea ^{93,190,191,462}	1–5	—	5%–10%
<i>Candida</i> ^{111–115,344,357,360,361}	1–4	Fluconazole, micafungin, voriconazole	Systemic infection: <5% colonization: 30%
<i>Aspergillus</i> and other molds ^{114,115,122,345,346,367}	1–4	HEPA air filtration, itraconazole, voriconazole, posaconazole, micafungin, or low-dose amphotericin	<5%
Respiratory viruses ^{285,287,289,291–295,299,300,302,304}	2–5	Isolation, hand washing	15%
Idiopathic pneumonia syndrome ⁸⁴	2–4	—	8%–17%
Early Postengraftment (4–26 Weeks) and Late Postengraftment (26–52 Weeks) Risk Periods			
Cytomegalovirus (seropositive) ^{62,132,134,141,229–232}	7–26	Ganciclovir or foscarnet	<5% (end-organ disease), up to 40% (antigenemia/viremia)
Varicella-zoster virus (seropositive) ^{255–257,264}	4–52	— ^a	≤50%
<i>Aspergillus</i> and other molds ^{122,123,346,363,367}	4–26	Itraconazole, posaconazole, or voriconazole	10%–15%
BK virus ^{37,46–48}	—	—	≤50% (shedding)
Adenovirus ^{268,269,271,273,274}	—	—	1%–5%
<i>Pneumocystis jirovecii</i> ^{124,126,352}	4–104	Trimethoprim-sulfamethoxazole	<1%
<i>Toxoplasma gondii</i> (seropositive) ^{425,427,428,431}	2–8	— ^{b,c}	2%–7%
Infrequent Infections (May Span Multiple Risk Periods)			
Herpes simplex virus (seronegative)	—	—	<2%
Cytomegalovirus (seronegative) ^{220,226,227}	—	Blood product screening or filtration, letermovir	1%–4%
Varicella-zoster virus (seronegative)	—	—	<3%
<i>Streptococcus pneumoniae</i> ^{d,175}	—	Vaccination, penicillin ^{e,f}	<1%
<i>Haemophilus influenzae</i> ^{d,441,442}	—	Vaccination, penicillin ^{e,f}	—
<i>Neisseria meningitidis</i> ^{d,441,442}	—	Penicillin ^f	—
Human herpesvirus 6 ^{323,325–327,329–332,334,335,463}	—	Ganciclovir or foscarnet	5%–60%, few treated
Epstein-Barr virus ^{312,317,318}	—	—	<1% (disease)
<i>Nocardia</i> spp. ^{213,214}	—	— ^c	<1%
<i>Legionella</i> spp. ^{209–212,216}	—	—	<1%
<i>Mycobacterium</i> spp. ^{196–199,206,207}	—	Screening, then prophylaxis	<1%
<i>Listeria monocytogenes</i> ^{217–219}	—	—	<1%

^aAntiviral medications used to prevent other viral infections may be acting as prophylaxis against reactivation of varicella-zoster virus.

^bProphylaxis with pyrimethamine-sulfadoxine may be used for seropositive patients in countries with a high rate of seroprevalence.²⁹⁹

^cThe sulfa component of *Pneumocystis* prophylaxis may be acting as prophylaxis against *Toxoplasma* or *Nocardia* infection.

^dRisk is increased in patients with chronic graft-versus-host disease.

^eEfficacy in transplant recipients is undetermined.

^fIncreasing penicillin resistance may indicate a need for macrolides or extended-spectrum quinolones.

HEPA, High-efficiency particulate air (filter); HSCT, hematopoietic stem cell transplantation.

Prophylaxis of *Pneumocystis jirovecii* infection with trimethoprim-sulfamethoxazole (TMP-SMX), dapsone, atovaquone, or aerosolized pentamidine is required for 6 to 12 months or, if cGVHD is continuing, longer.^{124–128} GI dysfunction should preclude the use of atovaquone, the absorption of which is poor in that context.

Reactivation of CMV predictably occurs in 20% to 40% of patients who are CMV seropositive. Transmission to seronegative recipients from seropositive donors is uncommon.⁶² Surveillance for reactivation of CMV has been improved by the use of scheduled CMV testing with either pp65 antigenemia assay or PCR testing for serum DNA.¹²⁹

Ganciclovir or other antiviral therapy initiated preemptively at subclinical indications of reactivation has reduced the incidence of end-organ disease caused by CMV to only 5% to 10% of seropositive recipients.^{130–132} Continuing GVHD or delayed immune recovery after partially matched- or unrelated-donor HSCT can lead to later-onset CMV infection and may indicate a need for prolonged CMV surveillance.

Late Risk Period

The late posttransplantation risk period begins at approximately day 100 and ends when the patient regains normal immunity, 18 to 36

months after HSCT.^{133–139} In general, clinical immune recovery is demonstrable by the end of the first year after transplantation as long as the patient is no longer taking immunosuppressive medication and remains free from GVHD. For patients with continuing cGVHD, this period persists as long as therapy for cGVHD is required and includes dysfunction of lymphocyte, macrophage, and humoral immunity. VZV reactivation, infections with encapsulated bacteria (*S. pneumoniae*, *N. meningitidis*, and *H. influenzae*), and invasive aspergillosis or other invasive tissue mold infections may develop in this late risk period. Malignant disease relapse, regardless of its tempo or the choice of therapy, impairs the restoration of immunocompetence. Survival after relapse may be lengthy for some more indolent malignancies (e.g., chronic myelogenous leukemia, chronic lymphocytic leukemia, follicular lymphomas), but their post-HSCT immunodeficiency will persist. Therefore, even after relapse, ongoing surveillance and therapy, as in post-HSCT patients in remission, is still required.

The most common clinical infection syndromes include sinusitis, bronchitis, pneumonia, and otitis media caused by respiratory viruses or bacteria. CMV disease may develop; therefore CMV surveillance must be continued in seropositive recipients with cGVHD.^{134,135} In CMV-infected patients with a risk of late disease identified by prior CMV disease, prophylaxis, or acute or cGVHD treated with the equivalent of more than 0.5 mg/kg of prednisone, no advantage has been identified between a PCR-based preemptive strategy or valganciclovir prophylaxis.¹³⁶ Late infections may be more common among patients with unrelated donors than among patients whose donors were family members, even in the absence of GVHD. Approximately 50% of late pneumonias in patients with ongoing cGVHD are caused by noninfectious interstitial pneumonitis. Lung histopathologic studies reveal organizing obliterative bronchitis that may respond to corticosteroid therapy.

IMMUNODEFICIENCY AFTER NONMYELOABLATIVE HSCT

NMA HSCT is associated with less disruption of mucosal barriers, shorter periods of severe neutropenia, fewer episodes of bacteremia in the first 30 days, and a trend toward fewer episodes of bacteremia during the first 100 days after HSCT.¹⁴⁰ However, this type of transplantation can still be associated with severe GVHD, often necessitating high-dose corticosteroid treatment. It is often used for older patients and those with compromised organ function and poor performance status. GVHD, CMV disease, and invasive fungal infection may be delayed 1 to 2 months, but the overall incidences of these conditions are similar to those with conventional myeloablative HSCT during the first year after HSCT.^{140–142} Patients undergoing NMA HSCT may need surveillance for CMV and fungal infections well beyond day 100, as well as preemptive or prophylactic treatment similar to that for myeloablative HSCT recipients between day 100 and 1 year after HSCT.

MEASURES TO REDUCE RISKS FOR INFECTION

Protective measures that should be discussed before HSCT and reinforced during the recovery period involve travel, crowds, and pets.¹⁴³ With regard to travel, there are no particular restrictions, but strategies to minimize transmission of infectious diseases have been summarized.³² Some social situations, such as sitting in a crowded movie theater or classroom, may increase the risk for acquiring a viral respiratory illness. Turning away from individuals who are coughing or sneezing, or even quickly donning a mask, may be helpful in preventing transmission of airborne, droplet-based infection. Patients need instruction to remember to augment infection prevention by washing their hands as soon as possible after being close to someone with a cold. Because outbreaks of noroviruses have involved cruise ships, and because other types of outbreaks (e.g., *Staphylococcus*) are commonly associated with the close living quarters of this type of vacation, cruise ships may be unwise vacation choices.

Healthy dogs and cats are considered acceptable pets. However, immunosuppressed patients should not get a new pet or be responsible for scooping cat litter because of potential *Toxoplasma* cyst exposure. Similarly, such patients should not play in sandboxes because these areas are concentrated sites that outdoor cats may use as litter boxes. Because

reptiles of many sorts have been reported to be infected with *Salmonella*, such patients should not touch these animals or their aquarium homes. The water of tropical fish tanks may carry *Mycobacterium marinum*. Psittacine birds can transmit *Chlamydia* *psittaci*.

Hand washing or the use of alcohol-based hand rub disinfectant is the mainstay of infection prevention in the hospital or clinic.¹⁴⁴ Persons entering the patient's room to perform examination or touch the patient (including visitors and health care workers) should wash or disinfect their hands outside the room. During respiratory virus season, infection control personnel will often add extra signage to doorways and other places on the wards to remind visitors of the importance of hand washing. Patients should not have direct contact with staff and/or visitors with respiratory viral infections. Routine use of gown, gloves, or masks, or a combination of these, is not required in the presence of a neutropenic transplant recipient, but ongoing caution to prevent interperson or nosocomial transmission is essential.

NATURAL HISTORY OF INDIVIDUAL INFECTIONS AFTER HSCT

With advances in infection prevention strategies, the risk periods for some infections are changing. It is important to understand the natural history of individual infections as they occur in the HSCT recipient and how it may be distinct from that in other immunocompromised patient populations. Infections that occur with a high incidence among HSCT recipients justify prophylaxis during the applicable risk period or empirical treatment during the course of infection (see Table 307.1). However, interactions between antibacterial prophylaxis, gut GVHD, and the microbiome remain to be addressed.^{145–147} Moreover, antimicrobial resistance in gram-negative rods can be a concern. A European, Australian, and Asian study found 655 gram-negative rods—caused infection episodes in 591 patients (Enterobacteriaceae, 73%; nonfermentative rods, 24%; and 3% others). Half of the rods were fluoroquinolone and noncarbapenem resistant; 18.5% carbapenem resistant; and 35.2% multidrug resistant.¹⁴⁸

Bacterial Infections

It is important to acknowledge that the incidence of BSI in HSCT recipients is high, mainly comprises CoNS,^{149,150} and that bacteremia may occur without fever because of the use of corticosteroids.¹⁵¹ The use of surveillance blood cultures without clinical indication in HSCT recipients is common, but the yield of significant results is low and leads to unnecessary interventions and increased cost.¹⁵²

Gram-positive organisms, higher in the past, currently account for half of bacteremias occurring after HSCT.^{105,150,153–155} Although the primary reservoir for these organisms is believed to be the skin, colonization of the GIT may be an additional source. Therefore a BSI can be a central line-associated bloodstream infection (CLABSI) or a mucosal-barrier injury laboratory-confirmed bloodstream infection (MBI-LCBI), defined by the Centers for Disease Control and Prevention.¹⁵⁶ A substantial evolution has occurred in the understanding of the sources of bacteremia in patients with mucosal-barrier injury. One important effect of this change is seen in determining the threshold and expectations related to changes of catheters, which would previously have been considered the default source of bacteremia.¹⁵⁷ *Staphylococcus epidermidis* (i.e., CoNS)¹⁵⁸ is the species most commonly recovered in culture from the skin and nose. Oropharyngeal organisms include *Streptococcus pyogenes*, *Streptococcus mitis*, and *Enterococcus* spp. (both vancomycin-sensitive and vancomycin-resistant species).¹⁵⁵ The troublesome incidence of vancomycin-resistant enterococcal infection continues to rise without a concomitant improvement in the ability to determine its significance or optional therapeutic approaches.^{159,160} although integration of colonization status is helpful in guiding rational use of empirical therapy.¹⁶¹ Unlike catheter-associated infections with *S. aureus*, *Candida* spp., or some gram-negative bacilli, most gram-positive bacteremias can be managed successfully without removal of the intravascular device.^{162,163} Methicillin-resistant *S. aureus* (MRSA), although rare in neutropenic infections, but may be associated with high mortality (≈7%–50%), in particular with MRSA outbreaks.¹⁴⁹ If the patient does not respond to initial antibiotic management or if there is tenderness or erythema along a catheter tunnel tract, the catheter should be removed. In rare cases, adjunctive surgical

débridement of the skin tunnel is needed. Catheter removal and surgical débridement are often required when a tunnel infection is caused by rapidly growing mycobacteria.

Gram-negative organisms are the second most frequent cause of BSI. The incidence of infection with *Pseudomonas* spp. is low, in part because of the use of antipseudomonal antibiotics for prophylaxis against, and initial empirical therapy for, neutropenic fever.¹⁰⁵ However, the virulence and potential for sepsis syndrome remain high for gram-negative organisms, including *E. coli*, *Klebsiella* spp., *Enterobacter* spp., and *Pseudomonas* spp. As a result, gram-negative infections preengraftment can be associated with a higher mortality rate.^{164,165} Remodeling of sanitary and water supply systems decreased multidrug-resistant *Pseudomonas aeruginosa* infections (from 18.9%–3%).¹⁶⁶ Carbapenem-resistant Enterobacteriaceae (CRE) has been an emerging issue worldwide (*Klebsiella pneumoniae* carbapenemase-producing–*K. pneumoniae* [KPC-Kp] in the United States) and reported as 0.4% and 2% in a retrospective multicenter Italian study after autologous and allogeneic HSCT, respectively.¹⁶⁷ Although the incidence of CRE was low, the mortality rates, in particular after allogeneic HSCT, were very high (16% and 64%, respectively). Gut colonization by multidrug-resistant gram-negative bacteria is an independent risk factor for the development of intestinal aGVHD.¹⁶⁸ CRE occurs in very sick patients as shown by a recent study that found risk factors for CRE included breakthrough infection while taking a carbapenem medication, longer hospitalization, requiring intensive care support, and previous other antibiotic therapy.¹⁴⁸ Another study from Europe showed that CRE before engraftment was the only risk factor for increased mortality after haploidentical HSCT in multivariate analysis.¹⁶⁴ In the United States, CRE infections are almost absent.

Although bacteremias have historically occurred during the neutropenic period, bacteremias continue to develop in patients with long-term central IV catheters, in patients with ongoing immunosuppression resulting from GVHD or its therapy, and in those with neutropenia secondary to graft failure or drug-related marrow suppression (e.g., ganciclovir).^{169,170}

Bacterial meningitis was more common after allogeneic HSCT compared with autologous HSCT, caused by *Streptococcus pneumoniae* in 11 patients, *Neisseria meningitidis* in 2, and *Streptococcus mitis* in 1 patient.¹⁷¹

Encapsulated Bacteria

Functional hyposplenism is not uncommon and is limited to patients with cGVHD.¹⁷² cGVHD is also associated with decreased immunoglobulin M (IgM) memory B cells,¹⁷³ that are present in the marginal zone of the spleen and may produce immunoglobulins with specificity for polysaccharide antigens.¹⁷⁴ Therefore, after allogeneic HSCT, the risk for life-threatening bacterial sepsis with encapsulated organisms is increased. Invasive pneumococcal infection may occur months to years after HSCT. The annual incidence is 8 per 1000 allogeneic HSCT recipients and is higher (21/1000) among those with cGVHD.¹⁷⁵ Penicillin or macrolide prophylaxis may be indicated until immunosuppression is discontinued.^{138,176} Reports of frequent penicillin-resistant pneumococcal infections in the general population have prompted a change in prophylaxis from penicillin to a quinolone in some centers.^{177,178}

Viridans-Group Streptococci

Viridans-group streptococcal bacteremias, mostly caused by *S. mitis*, may carry a high mortality rate early after HSCT.^{179–182} Although consistently reported that it occurred in the very early severely neutropenic phase of HSCT (median, ≈4 days), the incidence of breakthrough *S. mitis* BSI varied between 3% and 16% during fluoroquinolone prophylaxis.^{183–185} Poor dental hygiene and myeloablative conditioning are risk factors for *S. mitis* bacteremia in HSCT patients.^{184,186} Normally antibiotic sensitive, these organisms may be resistant to ciprofloxacin, levofloxacin, and penicillin in patients depending on prophylactic antibiotics. Vancomycin is the drug of choice for HSCT patients. Oral ulcerations caused by HSV reactivation during conditioning are believed to be an entry point, corroborated by a decreased incidence of viridans-group streptococcal septicemia after active prophylaxis of HSV infections with acyclovir.¹⁸⁰

Clostridioides difficile

Increased rates of *C. difficile*-associated diarrhea or infection (CDI) have become a major issue for many hospitalized populations. The incidence of *C. difficile* after allogeneic HSCT is reported to be between 10% and 34%.^{187–189} The incidence is lower after autologous HSCT (6%).¹⁸⁷ Although most CDIs occur before engraftment, CDI continues to occur after engraftment as well.¹⁹⁰ Prior hospitalization, relapsed underlying disease and chemotherapy, receipt of any high-risk antimicrobials (e.g., cephalosporin and GVHD) were found as risk factors.^{190,191} Of interest, CDI may also increase gut GVHD.¹⁸⁷ At Memorial Sloan Kettering Cancer Center, the incidence of early disease during the period 2005–10 was 11.9%, and no support was found for an increased pathogenicity of the NAPI/027 strain.¹⁸⁸ The risk for recurrent disease in allogeneic HSCT, which is as high as 31%, is related to antimicrobial use for indications other than the treatment of CDI.^{188,192} The relative roles of colonization or acquisition during hospitalization are unclear, but rates of disease during hospitalization are significantly elevated in those colonized at admission.^{193,194} Although little data support any choice between standard therapies in HSCT, the role of fecal microbiota transplantation should be explored. A case report of nasojejunal instillation of donor stool had a successful result.¹⁹⁵

Mycobacteria

Mycobacteria are an infrequent cause of infection after HSCT, but it is important to identify them because treatment requires medication that would not be used empirically. The rapidly growing nontuberculous mycobacteria are responsible for skin/soft tissue infection, catheter exit site infections, tunnel infections, bacteremia (with waterborne *Mycobacterium mucogenicum*), or pneumonia.^{196,197} Patients with cGVHD are at high risk for both nontuberculous mycobacteria¹⁹⁸ and tuberculosis.¹⁹⁹ Infection with *Mycobacterium tuberculosis* occurs predominantly in countries with high endemic rates; the number of cases may be increasing worldwide as transplantation becomes available globally.^{200–203} Transplantation candidates and donors who are at risk for reactivation of latent tuberculosis can be readily identified during the pretransplantation evaluation on the basis of residence in endemic areas or close contact with another person with known or suspected tuberculosis. For these transplantation candidates, screening is recommended with tuberculin skin testing or an ex vivo IFN- γ release assay, such as QuantiFERON-TB Gold (Qiagen, Germantown, MD).^{204–207} Clinically significant infection can be prevented by extended-duration antituberculous prophylaxis when QuantiFERON-TB Gold assays or reactive tuberculin skin testing yields positive results. Potential transplantation patients and donors should also receive screening if they have a history of abnormal chest radiographic findings before HSCT, have recently traveled to a foreign country for longer than 3 months, have been employed in an institution with tuberculous clients, have a history of alcoholism or IV drug use, or are seropositive for HIV. QuantiFERON-TB Gold has 96% to 99% specificity that is unaffected by the vaccine for bacillus Calmette-Guérin.²⁰⁸ For patients with a positive result of a screening test but no signs of active tuberculosis and no previous antituberculous therapy, a chest radiograph and liver function studies should be obtained, in addition to peritransplantation and 9-month posttransplantation therapy with isoniazid and pyridoxine. A recent study from South Korea showed that tuberculosis was higher (2.9%) than the general population and that isoniazid prophylaxis, starting at the time of the positive IFN- γ release assay result, did not decrease the incidence of tuberculosis after allogeneic HSCT.¹⁹⁹

Intracellular Bacteria

Legionellosis^{209–212} and nocardiosis^{213,214} are uncommon, but both can manifest as pneumonia or lung nodules in the HSCT patient. Skin nodules and serositis (peritonitis, pericarditis) can also occur. The mean time to legionellosis after HSCT was 390 days and perhaps is associated with cGVHD. Most of the patients who had nocardiosis (median time was 10 months) were on prednisone, had lymphopenia, and received a myeloablative conditioning regimen.²¹⁴ Detection of *Legionella* by direct fluorescent antibody (DFA) assays has proved unreliable in the HSCT setting because of false-positive results and because it does not detect a high proportion of disease caused by *Legionella* species. These

species can include *Legionella feeleyi*, *Legionella micdadei*, and *Legionella bozemanii*.²¹² Urinary antigen testing is the most common test used for diagnosis, whereas nucleic acid–based tests are increasingly available for use in *Legionella*.²¹⁵ Culture remains the gold standard for diagnosis, but this takes time and its sensitivity varies significantly.²¹⁶ Infection can persist or relapse after 3 weeks of appropriate antimicrobial therapy (e.g., fluoroquinolone), which suggests that prolonged antibiotic treatment is indicated for HSCT recipients with legionellosis.²¹¹ Medical therapies for nocardiosis often consist of administration of sulfonamide in combination with a synergistic agent, such as ceftazidime or imipenem-cilastatin; adjunctive surgical débridement may be useful for catheter-related infections with this organism. Nocardiosis is associated with high mortality (up to 50%).²¹⁴ The role of other intracellular bacterial agents as pathogens has not been well defined, but *Listeria monocytogenes* infection may manifest as bacteremia or meningitis.^{217–219}

Viral Infections

Certain viral infections are preventable. Administration of acyclovir for HSV-seropositive patients during the preengraftment period is widely accepted. CMV-safe (serologically screened or filtered) blood product transfusions for CMV-seronegative patients have proved very effective in preventing transfusion-acquired CMV infections.^{220,221} Periodic (e.g., weekly) CMV diagnostic surveillance for CMV-seropositive patients during the postengraftment period and prompt institution of antiviral therapy are essential. CMV-seronegative patients (with seronegative allogeneic donors) are at comparatively low risk; therefore diagnostic monitoring continues for only 6 to 10 weeks. Scrupulous hand washing and avoidance of crowds to prevent transmission of respiratory viral and other infections (in the hospital or ambulatory clinic) remain the mainstay of effective infection control practice for the HSCT recipient.

Herpes Simplex Virus

Among HSV-seropositive recipients during the first month after transplantation, the incidence of HSV reactivation can be reduced from 80% to less than 5% through the use of acyclovir or valacyclovir, initiated at the time of conditioning and continued until mucositis has healed.^{109,222,223} The majority of postengraftment HSV infections are confined to the oropharynx, although the infection occasionally extends directly to squamous epithelial surfaces in the upper esophagus, larynx, or skin in the perioral or perianal areas. Patients who do not respond to acyclovir after engraftment, and particularly those who have received prolonged or repeated courses of acyclovir, may have acyclovir-resistant HSV. Foscarnet or cidofovir may be beneficial in that setting. In uncommon cases, HSV infection causes Bell palsy, hepatitis, or encephalitis. Valacyclovir achieves predictably higher drug levels than does acyclovir after oral administration.^{223,224}

Cytomegalovirus

CMV reactivation before day 100 continues to be associated with increased nonrelapse mortality, with relative risks by disease category ranging from 1.61 to 1.95 in a review of CIBMTR data from 2003–10.²²⁵ The historical incidence rate of primary CMV infection among the CMV-seronegative HSCT recipient can be reduced from 40% down to 4% or less by use of CMV-safe blood products during transfusions.^{226,227} A cellular graft is associated with CMV transmission when a CMV-seronegative

recipient receives a CMV-seropositive donor graft.²²⁸ T-cell–depleted graft, HLA-mismatched transplantation, steroid treatment, and acute or cGVHD are risk factors for CMV infections. The incidence of CMV reactivation and viremia, traditionally 70% in CMV-seropositive patients with allogeneic transplants and 45% among patients with autologous transplants, can be reduced down to between 20% and 40% by the use of preemptive or prophylactic antiviral therapy.^{130,229–236} With the most common approach, prophylactic acyclovir combined with preemptive early ganciclovir product therapy, the median time of onset of CMV end-organ disease has been delayed from 1 to 2 months toward 4 to 6 months after HSCT; this indicates that CMV surveillance must be longer in groups at high risk (see Chapter 137).^{134,135}

Weekly screening enables identification of patients who might benefit most from preemptive therapy with ganciclovir products (Table 307.2). CMV pp65 leukocyte antigen and quantitative DNA PCR testing are excellent methods for early CMV detection, although antigen testing is unreliable during neutropenia. Laboratories are converting CMV viral load to log₁₀ international units to decrease interinstitutional variability. Once CMV is identified by an early detection method, most patients are treated with 7 to 14 days of induction ganciclovir therapy (5 mg/kg IV twice daily), followed by maintenance therapy (ganciclovir, 5 mg/kg IV once daily, or valganciclovir, 900 mg orally once daily), for several weeks beyond negative CMV test results (Table 307.3). Maintenance therapy may need to be continued for patients with persistent detection of virus or those with profound immunosuppression caused by active GVHD.

Oral valganciclovir is a safe and effective ganciclovir prodrug, with the valine ester cleaved during the first pass through the liver, and it can be considered for the patient who needs induction and/or long-term maintenance and is otherwise taking oral medications without difficulty.¹³⁶ A valganciclovir dosage of 900 mg once per day produces blood-level drug exposure similar to those produced by an IV 5 mg/kg dose of ganciclovir.

Foscarnet can be used empirically for patients who have marrow suppression from ganciclovir and fail to respond to this drug or who have concurrent HHV-6 viremia. Foscarnet is administered in doses of 90 mg/kg IV every 12 hours for induction and 90 mg/kg IV every 24 hours for maintenance, or renal dose equivalents. Good urine flow can minimize irritation of the urethra and labia by foscarnet. The serum biochemical abnormalities (chelation of calcium and phosphate) accompanying foscarnet therapy necessitates hospital observation and very careful electrolyte monitoring, at least through the initial days of its use.

Letermovir, an antiviral drug inhibiting CMV replication by binding to components of the terminase complex,²³⁷ when used for prophylaxis at a dose of 480 mg per day (or 240 mg per day in patients taking cyclosporine), significantly reduced CMV infection (CMV disease or CMV viremia leading to preemptive treatment) compared with placebo (37.5% vs. 60.6%, $p < .001$) in adults after allogeneic HSCT.²³⁸ Nausea, edema, and atrial fibrillation were more common in the letermovir group.

End-organ manifestations of CMV disease include pneumonitis (63%), enteritis (26%), and, in rare cases, retinitis (5%).^{131,239} CMV pneumonitis occurs in less than 5% of CMV-seropositive allogeneic patients who receive ganciclovir preemptive therapy during the first 100 days.¹³¹ CMV DNA load in bronchoalveolar lavage (BAL) fluid can

TABLE 307.2 Weekly Screening Schedule for Initiation of Preemptive Cytomegalovirus (CMV) Therapy After Hematopoietic Stem Cell Transplantation

CMV SEROSTATUS OF RECIPIENT	CMV SEROSTATUS OF DONOR	BLOOD PRODUCTS	DURATION OF WEEKLY SURVEILLANCE
Seronegative	Seronegative	CMV safe ^a	Weeks 2–12
Seronegative	Seropositive	CMV safe ^a	Weeks 2–12
Seronegative	None (autologous)	CMV safe ^a	Weeks 2–5
Seropositive	Seronegative or seropositive	CMV untested	Weeks 2–12, then every 2–4 weeks until week 26 ^b
Seropositive	None (autologous)	CMV untested	Weeks 2–5

^aBlood filtered to remove neutrophils or from seronegative donor.

^bPossible indications for testing beyond week 26 include high risk for late CMV disease (e.g., in patients treated with corticosteroids for graft-versus-host disease).

TABLE 307.3 Suggestions for Management of Possible CMV Infection After HSCT

INDICATION	STRATEGY ^a	COMMENT
Prevention		
Allogeneic Transplant		
Seropositive recipient	Preemptive ganciclovir ^b induction for subclinical viremia, 5 mg/kg bid for 7–14 days, followed by 5 mg/kg daily until the end of maintenance or ganciclovir prophylaxis ^c at engraftment	Some cases of CMV disease may occur shortly after ganciclovir discontinuation. CMV reactivation might be delayed, occurring later after HSCT.
Seronegative recipient with seropositive donor	Preemptive ganciclovir induction for subclinical viremia, 5 mg/kg bid for 7–14 days, followed by 5 mg/kg daily until the end of maintenance and seronegative or filtered blood products	Prophylaxis at engraftment is not recommended because of the low incidence of posttransplantation infection.
Seronegative recipient with seronegative donor	Seronegative or filtered blood products	
Autologous Transplant		
Seropositive recipient	Early ganciclovir induction of subclinical viremia, 5 mg/kg ganciclovir bid for 7 days, followed by 5 mg/kg daily for 14 days of maintenance	Because of the very low risk in some settings, monitoring is not uniformly advocated.
Seronegative recipient	Seronegative or filtered blood products	
Treatment of Disease		
CMV pneumonitis	Ganciclovir induction, 5 mg/kg bid for 14–21 days, followed by 5 mg/kg daily for at least 3–4 wk of maintenance plus IVIG every other day for the duration of induction	Extended maintenance throughout periods of severe immunosuppression (i.e., GVHD treatment) may be considered.
Gastrointestinal disease	Ganciclovir induction, 5 mg/kg bid for 14–21 days, followed by 5 mg/kg daily for at least 3–4 wk of maintenance	If deep ulcerations are present, maintenance may be required for a longer time.
Marrow failure	Foscarnet, 90 mg/kg bid for 14 days, followed by 90 mg/kg daily for 2 wk plus G-CSF	Ganciclovir plus IVIG has also been used.
Retinitis	Ganciclovir, 5 mg/kg bid for 14–21 days, followed by 5 mg/kg daily for at least 3–4 wk	Extended maintenance may be required.

^aRegimens should be accompanied by weekly monitoring with antigenemia or PCR-based nucleic acid testing.

^bOral 900-mg doses of valganciclovir produce blood levels that are similar to those for the standard intravenous dose (5 mg/kg) of ganciclovir. Foscarnet and cidofovir are acceptable alternatives. Renal dose adjustment is required for all antiviral agents.

^cFoscarnet, high-dose acyclovir, or valacyclovir are acceptable alternatives. Renal dose adjustment required for all antiviral agents.

bid, Twice daily; *CMV*, cytomegalovirus; *G-CSF*, granulocyte colony-stimulating factor; *GVHD*, graft-versus-host disease; *HSCT*, hematopoietic stem cell transplantation; *IVIG*, intravenous immune globulin; *PCR*, polymerase chain reaction.

be used to differentiate CMV pneumonitis from pulmonary shedding.²⁴⁰ Among patients with CMV pneumonitis, the mortality rate is as high as 50%, even when prompt antiviral treatment is combined with intravenous immune globulin (IVIG) or CMV-specific immune globulin.²⁴¹ CMV viremia and pneumonitis are rare before engraftment.^{242,243} Anorexia, nausea, vomiting, and sometimes diarrhea characterize CMV gastroenteritis; the diagnosis is made by endoscopy and biopsy with immunoperoxidase staining of CMV-infected cells.²⁴⁴ CMV disease of the GIT is often associated with GVHD of the specific organ.²⁴⁴ Response to therapy is not ensured, even with ganciclovir, but CMV GI disease is not strongly associated with increased nonrelapse mortality.²⁴⁵ Although CMV retinitis is common among patients infected with HIV, it is quite uncommon among HSCT recipients.²⁴⁶

Once an end organ has established disease from CMV, the infection is difficult to treat. CMV pneumonitis is treated with a combination of ganciclovir at induction doses for 14 to 21 days and IVIG (500 mg/kg every other day for the duration of induction, 14–21 days) and then with maintenance ganciclovir. Standard IVIG is generally used because CMV-specific immune globulin is scarce and costly and has not been shown to improve outcome. CMV enteritis is treated with ganciclovir at induction doses for 3 weeks or longer.²⁴⁷ Treatment of protracted CMV enteritis might include ganciclovir and a longer duration of ganciclovir maintenance therapy to facilitate GI healing.²⁴⁴

Development of a CMV-specific cytotoxic T-lymphocyte response is critical for the reconstitution of normal immunity and protection from late CMV disease.²⁴⁸ Long-term IVIG delays recovery of CMV immunity. For patients who remain at risk for late disease, CMV monitoring should be continued beyond day 100. Patients who are treated with acyclovir followed by ganciclovir or those treated with serial ganciclovir courses may be at increased risk for developing genotypic resistance. Clinical resistance episodes should be treated with foscarnet (or cidofovir) until genotype assays for mutations in

the *UL97* or *UL54* gene are able to confirm virologic resistance.²⁴⁹ An illustrative case in the development of drug resistance identifies the donor-negative/recipient-positive setting, the prolonged period to development of a CMV-specific response, and GVHD with extended immunosuppression as risks.²⁵⁰ Extended ganciclovir therapy appears to delay recovery of cytotoxic T-lymphocyte (CTL) activity, either by a direct effect on lymphocytes or by limitation of the amount of antigen exposure to lymphocytes. CMV-specific cytotoxic T cells were used in 50 patients and reduced the need for pharmacotherapy (17% vs. 36%, $P = .01$) and in the total number of treatment days in the cohort receiving CTL (3.4 days vs. 8.9 days, $P = .03$), but this treatment did not reduce CMV reactivation rates. There was also no increase in GVHD.²⁵¹ Currently, CMV-specific T cells from third parties are being evaluated for CMV and EBV prevention or treatment in the HSCT setting.^{252,253} Brincidofovir, CMX001, an oral bioavailable lipid acyclic nucleoside phosphonate, significantly reduced the incidence of CMV events at a dose of 100 mg twice weekly compared with placebo (10% vs. 37%, $P = .002$) in a randomized HSCT study of CMV seropositive patients.²⁵⁴

Varicella-Zoster Virus

VZV infection is a primary occurrence (5% of occurrences) or a reactivation (95%) in 40% of patients at any time in the first year after transplantation.^{255,256} Conditioning regimen intensity may not be important regarding VZV reactivation.²⁵⁷ VZV can be effectively prevented with acyclovir prophylaxis (800 mg twice a day orally),²⁵⁸ but VZV prophylaxis is not used at all transplantation centers because only 30% to 50% of adult patients and 25% of pediatric patients develop this infection during the first year after transplantation. The median time of onset is 5 months after transplantation. Prolonged antiviral prophylaxis may delay the onset of VZV but is not associated with rebound VZV.²⁵⁸ Localized zoster may manifest atypically with a few vesicles, or skin lesions may

appear as atypical vesicles; therefore laboratory confirmation of VZV reactivation is recommended.

Manifestations of VZV disease are most often dermatomal shingles but may include hemorrhagic pneumonia, hepatitis, abdominal pain, central nervous system (CNS) disease, thrombocytopenia, and retinal necrosis.^{259–263} Disseminated or visceral varicelliform zoster may manifest as low back pain or acute abdominal pain before the appearance of skin lesions. GVHD is a strong predisposing factor for VZV dissemination. Visceral varicella-zoster infections are rare but can be fatal.²⁶⁴ Most fatal cases of disseminated or abdominal zoster occur in patients who were treated with suboptimal doses of acyclovir or for whom therapy was initiated relatively late in the course of infection. High-dose acyclovir (10 mg/kg IV every 8 hours) has been the treatment of choice for disseminated VZV infection. Valacyclovir and famciclovir can be used as step-down treatment after IV acyclovir or as initial treatment of localized infection. Although uncommon, patients who are already seropositive may acquire a second primary VZV infection. VZV vaccination is recommended at the 2-year anniversary visit for patients who have been free of immunosuppressive medications for several months, unless the underlying hematologic or oncologic disease is in relapse.^{265,266} The vaccine used should be the lower-plaque-forming unit version (Varivax) used for prevention of chickenpox among children, not the higher-titer (Zostavax) vaccine used for immunocompetent older adults.

VZV is a fastidious virus and may not withstand the time required to transport the specimen to the diagnostic laboratory. By scraping the base of a vesicle and examining the cells by DFA with VZV-specific monoclonal antibodies, clinicians can best diagnose lesions of herpes zoster and chickenpox. The Tzanck smear is less sensitive and is no longer recommended. Tissue diagnosis can be made through histologic examination, immunohistochemical techniques, or culture. PCR assay, if available, is the laboratory technique of choice.

When a VZV-seronegative patient receives a significant exposure to a person with active or incubating chickenpox, a course of acyclovir with or without varicella-zoster immune globulin (VariZIG) is recommended to prevent chickenpox.²⁶⁷ Acyclovir or valacyclovir for 3 to 22 days after exposure seems appropriate, based on institutional experience and extrapolation from the treatment of VZV disease. For a VZV-seropositive patient living in the same dwelling as someone with an index case of active chickenpox or shingles, acyclovir is reported to be useful in preventing new infection. VariZIG is usually not given to exposed seropositive patients, but antiviral prophylaxis is often recommended, depending on the exposure, the length of time since the transplantation, and the level of immunosuppression. For lower-risk seropositive patients, acyclovir or valacyclovir may be appropriate, although the efficacy in this situation is unknown. Patients already receiving empirical ganciclovir for CMV reactivation do not need further antiviral agents.

Adenovirus

Adenovirus infection reactivates in approximately 12% of allogeneic and approximately 6% of autologous adult HSCT patients.^{268,269} Chronic shedding can occur in the absence of clinical disease, but adenovirus can also be acquired from respiratory droplet transmission.²⁷⁰ In its most common clinical manifestation in this setting, adenovirus is a cause of hemorrhagic cystitis.⁴¹ Systemic infection in the lungs, liver, GIT, and kidneys occurs in 18% to 20% of infected patients. GVHD is a risk factor for the occurrence of clinically apparent adenovirus infection after HSCT.²⁷¹ In addition, allogeneic patients who do not receive ganciclovir (seronegative for CMV or seropositive without need for ganciclovir) are at higher risk for developing adenovirus infection than are patients who did receive ganciclovir, even though ganciclovir has no activity against adenovirus.²⁶⁸ Alternative grafts, mismatched HSCT, T-cell depletion, lymphopenia, and GVHD were found to be risk factors.²⁷²

Immunofluorescence, shell vial, or conventional tube culture of blood, urine, stool, or tissue can be used to diagnose adenovirus. PCR testing is helpful in diagnosis of infection but in general is not cost-effective for surveillance because many patients are asymptomatic.^{273,274} Adenovirus viral load can be used as a valid surrogate end point for disease severity and prognosis.²⁷⁵ Given that no effective therapy is available for adenoviral infections, although cidofovir or the lipid-conjugated derivative

brincidofovir (CMX001) have been used in patients able to tolerate the potential toxicities.^{276,277} Some centers use quantitative viral load monitoring, followed by preemptive treatment²⁷⁸ that might be correlated with faster virologic clearance and improved outcomes when started at a lower adenovirus viral load.²⁷² Brincidofovir, with less nephrotoxicity, may have shown effectiveness even when cidofovir failed patients in case reports.^{275,279} Adenovirus-specific cytotoxic T cells are becoming available, but the generation of virus-specific T cells can be costly and time consuming. Newer techniques allowed generation of donor-derived adenovirus specific T cells within 2 weeks, and in first in-human studies these cells cleared cidofovir-resistant adenovirus (i.e., increasing titers) in a few patients.²⁸⁰ Third-party prepared “off-the-shelves” products showed promising safety and efficacy profile.^{253,281}

Norovirus

New diagnostic testing of stool has found norovirus to be a frequent cause of diarrhea in stem cell transplant recipients.²⁸² Crampy abdominal pain, nausea, or vomiting may accompany the diarrhea. Duration of diarrhea can be quite prolonged, even months, and create diagnostic confusion with GVHD or *C. difficile* diarrhea. Virus shedding may persist after resolution of symptoms.²⁸³ Because of the low inoculum needed to cause disease and the known persistence of norovirus on surfaces, some facilities with a known norovirus problem have required gloves for care providers and hand washing, not just hand cleanser, of persons exiting the norovirus patient's room.²⁸⁴

Respiratory Viruses

Patients who have undergone HSCT and develop a respiratory viral infection typically present with rhinorrhea and nasal congestion and may also have fever, cough, throat pain, headache, or myalgias.²⁸⁵ The common pathogens in such patients include respiratory syncytial virus (RSV), parainfluenza virus, and, to a lesser extent, influenza virus, rhinovirus, human bocavirus, human metapneumovirus, and coronavirus.^{285–296} Current rapid test methods allow detection of RSV, parainfluenza, influenza, and other viruses in respiratory specimens within 48 hours (see Chapter 16). Respiratory virus infections commonly occur during the winter season and cause pneumonia in up to 50% of patients, although progression to lower tract infection is less common with NMA conditioning regimens.²⁸⁹ In contrast, parainfluenza virus type 3 infections and rhinovirus may occur throughout the year²⁹³ and nosocomial outbreaks of RSV have occurred at other times than the established winter season. Influenza, most often type A, infrequently progresses to pneumonia. Current diagnostic strategies that use commercially available multiplex pathogen panels will detect, but not distinguish, human rhinovirus from an emerging pathogen, such as enterovirus D68, an increasingly important pathogen in HSCT recipients.²⁹⁷ Prophylactic or early initiation of oseltamivir therapy for upper respiratory tract disease may prevent progression to lower respiratory tract disease.²⁹⁴ In a recent study of high-risk patients (e.g., most of them underwent alternative donor or HLA-mismatched donor HSCT), coronavirus was not rare (23%) and not benign (e.g., about one-third of patients progressed to LRTI, 19% were hospitalized, and 5% died).²⁹⁸

RSV and parainfluenza are associated with a high incidence of progression from upper to lower tract disease among infected patients. Upper respiratory tract illness with parainfluenza usually resolves without serious sequelae. LRTI has a mortality rate of 80% for RSV and 30% to 35% for parainfluenza virus.^{287,299,300} Lymphopenia is a risk factor for progression to LRTI.³⁰¹ Therapy with oral or aerosolized ribavirin or a combination of ribavirin and IVIG has been used for RSV LRTI and seems to decrease lung-related mortality,³⁰² although rigorous trials to determine its efficacy are not available.^{285,303} Oral ribavirin is used in patients with upper respiratory tract infections who are at risk for progression to LRTI.^{304,305} The survival rate appears to be higher when treatment is initiated before significant hypoxia is present, and ribavirin may help to decrease the viral burden.^{303,306} There are only anecdotal case reports regarding the effectiveness of ribavirin for treatment of other respiratory viruses, including parainfluenza, adenovirus, and influenza. A meta-analysis found no decrease in mortality of LRTI with ribavirin for parainfluenza infections.³⁰⁷ Preemptive therapy with ribavirin in patients with positive nasopharyngeal cultures for RSV appears

promising. Patients who develop respiratory viral pneumonia before engraftment have poorer outcomes. Aerosolized ribavirin (Virazole) is administered via facemask to adults through a small-particle aerosol generator. Contamination of the patient's room with this possibly teratogenic agent is of concern to pregnant hospital staff (see Chapter 45). The inconvenience, lack of efficacy data, difficult-to-interpret indications, and enormous cost of aerosolized ribavirin have led many institutions to adopt oral ribavirin as an alternative.^{308,309}

Prevention of exposure is critical because treatment is not very effective. Protection involves the use of frequent hand washing by hospital staff and droplet isolation of patients with colds or respiratory tract symptoms. In addition, family members and health care workers with upper respiratory tract symptoms should be separated from patients. Vaccination of family members, health care workers, and other close contacts against influenza may help control exposures. Amantadine or rimantadine prophylaxis has limited usefulness because of the widespread development of resistance. Oseltamivir or inhaled zanamivir provides useful prophylaxis against both influenza A and B especially within 48 hours of exposure and appeared useful when used in a housing facility for HSCT recipients.³¹⁰ Immune globulin prophylaxis with RSV-specific polyclonal or monoclonal antibody, which is useful in infants at high risk, has not been sufficiently evaluated in the HSCT setting.³¹¹ During the respiratory virus season, all patients with respiratory symptoms should have a sample taken from the nasopharynx to be evaluated for respiratory viruses.

Epstein-Barr Virus

The majority of Epstein-Barr virus (EBV) reactivation is subclinical and requires no therapy. UCB transplants, aGVHD, HLA-mismatched HSCT, and T-cell-depleted HSCT perhaps lead to prolonged T-cell lymphopenia and augment the risk for EBV reactivation.^{312–315} The findings that the detection of EBV DNA in peripheral blood lymphocytes by PCR assay is highly correlated with EBV-related transplantation lymphoproliferative disorder (PTLD)³¹⁶ and that rituximab is effective in EBV-related PTLD³¹⁴ led to investigate whether preemptive treatment is effective. Quantitative diagnostic monitoring of EBV viral load accompanied by preemptive therapy may reduce the risk for progression to life-threatening PTLD.³¹⁷ In most cases, high viral loads are associated with progression to PTLD.³¹⁸ Infusions of rituximab or nonirradiated donor leukocytes may be effective treatment for allograft recipients with high-titer EBV viremia or PTLD.^{312,319,320} However, methods, sampling, and the threshold to start preemptive rituximab treatment remain to be standardized.^{321,322}

Human Herpesvirus 6

HHV-6 can reactivate after HSCT from the latent state in organs/cells (latent in organs/cells such as brain, kidneys, salivary glands, T-lymphocytes, BM progenitor cells), as can other herpesviruses, and probably cause fever, rash, BM suppression, fatal meningoencephalitis, and interstitial pneumonitis in less than 2% of HSCT patients.³²³ A distinctive limbic encephalitis from HHV-6 can cause confusion, memory loss, hallucinations, seizures, and even status epilepticus.^{324–326} Magnetic resonance imaging shows bilateral signal change in the medial temporal lobe, which may be more prominent in the week(s) after clinical presentation. HHV-6 is found in the cerebrospinal fluid (CSF) and blood.³²⁴ In patients clinically suspected for CNS infection, HHV-6 positivity in CNS should be considered as active infection regardless of blood test results. Severe hyponatremia, caused by a syndrome of inappropriate secretion of antidiuretic hormone, developed in patients with CNS infection.^{325–327} IV foscarnet is recommended. Permanent neurologic sequelae can occur.³²⁸ Recipients of UCB may have more viremia than do other populations.^{329,330} HHV-6 appears to reactivate commonly, occurring in 46% of HSCT patients according to culture diagnosis and as many as 100% of patients according to PCR assay of blood.³³¹ Many episodes of reactivation detected by DNA PCR may be asymptomatic, and the value of therapy for subclinical viremia and hence surveillance is unknown.³³² It may also represent latent infection.³³³ High-level HHV-6 viremia is associated with aGVHD and nonrelapse mortality.³³⁴ Most strains of HHV-6 identified after HSCT appear to be caused by the B variant in blood or urine, although the A variant has been correlated

with pneumonitis.³³⁵ HHV-6 has greater than 60% DNA homology with CMV, and treatment of documented infection is usually initiated with induction doses of foscarnet. Responses to antiviral therapy are not universal, and benefits of antiviral therapy have not been rigorously determined.³³⁶ Third-party virus-specific T cells seem promising in HHV-6 and BK virus treatment.²⁵³

Parvovirus

Parvovirus B19 (human parvovirus [HPV] B19) is a rare cause of refractory anemia with erythroid hypoplasia or BM after HSCT.^{337–340} Skin rash confusion with GVHD rash has been reported.³⁴¹ Asymptomatic patients should not be tested.³⁴² Antibody or PCR tests detect parvovirus, although PCR assay may yield positive findings for months after the acute infection. Use of single-patient rooms on HSCT wards may be preventing transmission of this contagious virus to other patients undergoing HSCT, and the administration of IVIG for other reasons may be treating subclinical infections (see Chapter 147). In a study, 21 (1%) of 2123 blood products tested positive for the presence of HPV B19 DNA-transfused immunocompromised hematology patients, including HSCT patients. Fourteen patients received these positive blood products but did not have symptomatic infections.³⁴³

Fungal Infections

Invasive fungal infections are important causes of morbidity and mortality. The major causes of invasive fungal disease include *Candida* spp., *Aspergillus* spp., and, less frequently, the non-*Aspergillus* filamentous molds. Patients undergoing allogeneic transplantation are at 10-fold increased risk for invasive fungal infection compared with patients receiving an autologous graft. Systemic fluconazole prophylaxis (at least 200 mg/day) or low-dose amphotericin B (0.1–0.3 mg/kg daily) can decrease the incidence of systemic candidiasis.^{111–113,344,345} Advanced-generation azoles and micafungin prophylaxis extend the spectrum of organisms covered to include molds, but their general value or cost-effectiveness for all allograft recipients has not been shown.³⁴⁶ Empirical therapy for febrile neutropenic patients is discussed in Chapter 306. As prophylaxis with triazoles has become more prevalent, awareness of changes in the epidemiology of fungal infections is necessary. In a report of transplants from 2002–11, *Aspergillus* (31 episodes) and candidemia (15 episodes) were most common, but 16 episodes were caused by *Mucorales* and other uncommon pathogens, including yeasts such as *Trichosporon* and unidentified molds.³⁴⁷

Pneumocystis

P. jirovecii infection³⁴⁸ usually manifests as pneumonia with dyspnea, cough, fever, and bilateral infiltrates in the majority of infected patients.¹²⁴ It can occur after both autologous and allogeneic transplantation, although the frequency is lower for the former. (1→3)- β -D-glucan in serum or BAL can be useful in diagnosis,^{349,350} although microscopic evaluation of BAL is the gold standard. Before the use of routine prophylaxis, *Pneumocystis* infection occurred in approximately 7% of patients who underwent allogeneic HSCT, at a median of 1 to 3 months after transplantation, and was associated with a 5% risk for death.^{351,352} A large survey of CIBMTR data from 1995–2005 found rates of 0.63% and 0.28% in allogeneic and autologous transplant recipients, respectively. This low incidence was accompanied by a significantly higher risk of death.³⁵³ Prophylaxis with TMP-SMX has resulted in negligible rates of infection. For patients who do not tolerate medications containing sulfa, prophylaxis options include desensitization with TMP-SMX and use of dapsone,^{128,354} atovaquone,³⁵⁵ and inhaled pentamidine.^{125,356} The treatment of choice for *P. jirovecii* infection is TMP-SMX.^{124,126}

Candida

Candidiasis is an infection acquired from endogenous organisms colonizing the GIT; it usually manifests as fungemia or visceral candidiasis (see Chapter 256). Before fluconazole prophylaxis, the onset of candidiasis occurred at a median of 2 to 3 weeks after transplantation, and *Candida* spp. were second in frequency to *Aspergillus* spp. as the cause of brain abscess after HSCT.^{357,358} The current cumulative incidence rate of invasive candidiasis during the first year after HSCT is probably less

than 5%.³⁵⁹ Risk factors for invasive candidiasis include neutropenia, breakdown of the normal mucosal barriers, central venous catheter, and the use of broad-spectrum antibiotics or corticosteroids. *C. albicans* infections are successfully prevented when fluconazole is given as prophylaxis from the time of conditioning until either engraftment or day 75 after HSCT. The strategy of prolonged fluconazole therapy has been associated with improved survival rates, although the mechanism of the observed benefits is uncertain.^{113,360,361} The benefit of fluconazole prophylaxis is less clear for autologous transplants, for which the degree of mucositis is less.

With the use of fluconazole since the 1990s, the number of *Candida* infections has decreased.³⁴⁴ The spectrum of colonizing and infecting *Candida* organisms has shifted from *C. albicans* and *Candida tropicalis* to include *C. krusei*, *C. glabrata*, and *Candida parapsilosis*.¹¹⁶ *C. krusei* is innately resistant to fluconazole. *Candida auris* is an emerging candidal species that is often multidrug resistant and is difficult to identify with standard laboratory methods.³⁶²

Aspergillus

Aspergillus and other mold infections are acquired exogenously, by inhalation of spores into the respiratory tract from the environment and in some localities may occur with higher frequency during the summer.¹²² Common sites of initial infection include the lung and sinuses, although contiguous or hematogenous extension to the CNS or other internal organs may occur. With the use of fluconazole prophylaxis during the preengraftment period, invasive aspergillosis emerged as the leading fungal infection found at autopsy among patients who underwent HSCT.³⁴⁴ Preengraftment prophylaxis with mold-active therapy is now being used for patients who may have *Aspergillus* incubating at the time of transplantation owing to prior infection or prolonged neutropenia.^{114,115,363} Postengraftment prophylaxis with posaconazole or voriconazole has led to a decrease in mortality among patients with a high risk for invasive aspergillosis.^{114,123,364} A CIBMTR analysis showed that prior invasive fungal infections (mostly *Aspergillus* and *Candida* infections) were not absolute contraindications for allogeneic HSCT, but patients with invasive fungal infections before transplantation did have a higher rate of invasive fungal infection at 1 year after transplant (24% vs. 17% control subjects, $P < .001$) and a higher rate of nonrelapse mortality (relative risk, 1.27 [1.09–1.49], $P = .002$).³⁶⁵

The incidence of invasive aspergillosis among patients undergoing HSCT ranges from 4% to 15%.^{122,366,367} The onset of *Aspergillus* infection after HSCT occurs in a bimodal distribution, with the first peak at 2 to 3 weeks (during neutropenia) and the second at 3 to 4 months after HSCT, usually in conjunction with persisting GVHD.¹²² Postengraftment aspergillosis can occur after 6 months, again alongside cGVHD but also with CMV. Older age is associated with the acquisition of aspergillosis during either the preengraftment or postengraftment risk periods. Donor type, male gender, and summer season are recognized risk factors for preengraftment aspergillosis, whereas construction in the vicinity of the hospital, GVHD and attendant corticosteroid therapy, lymphopenia, CMV, respiratory virus infection, and multiple myeloma are significant risk factors for the development of postengraftment aspergillosis.^{122,368} Early aspergillosis is temporally associated with neutropenia; therefore infection among autologous HSCT patients is rare after engraftment. The estimated 1-year survival rate among patients with proven invasive aspergillosis is 7% to 30%, although more aggressive, prolonged, or combination antifungal therapies may be improving these outcomes.

Preventive strategies focus on reducing both environmental and host risk factors. The use of HEPA-filtered air systems or laminar airflow rooms during the preengraftment risk period aids in the prevention of infection, particularly for allograft recipients. HEPA filters are capable of removing particles greater than 0.2 μm in diameter, such as mold spores. The patient's room is continuously maintained at positive pressure in relation to the corridor, which enhances the barrier effect. For transport out of HEPA-filtered rooms or after discharge, tight-fitting facemasks reproduce this barrier and are sometimes used, at least for the early post-HSCT period.

Patients might ask whether portable HEPA filters should be purchased for use after the hospitalization. This extra measure can be implemented on an individual basis. Units can be obtained for each of the rooms

that the patient will occupy during the day and night, and each unit is sized for the room it will be placed in. There is no evidence of the clinical efficacy of these filters out of the hospital setting in preventing acquisition of airborne mold infections. For outpatients they are probably of little value and of considerable expense.

Other prevention strategies, including nasal and aerosolized amphotericin B, have not been studied in controlled trials.

The availability of accurate early diagnostic tests for invasive fungal infections lags behind those for other types of infections. The *Aspergillus* galactomannan test is most useful for patients not already taking antifungal therapy that includes coverage for molds, which is a minority of high-risk allogeneic recipients.³⁶⁹ Other antigen- and nucleic acid-based diagnostic tests have been studied for early diagnosis of invasive tissue mold infection, but they have not demonstrated usefulness in clinical practice. Most have not been tested in large numbers of clinical samples from HSCT recipients. Blood cultures for molds rarely yield positive findings of mold organisms, except in the case of *Fusarium*.

A high index of suspicion in persistently febrile neutropenic patients and timely computed tomography of the chest to detect new infiltrates are important for early detection of invasive pulmonary aspergillosis. A small ground-glass halo around the lung lesion or pleura-based or nodular infiltrates on computed tomographic scans are highly suggestive of aspergillosis or other mold infection in a neutropenic host.¹¹⁸ Bronchoscopy with cytologic examination and culture of lavage fluid for fungi, as well as other organisms common to immunocompromised hosts, is important. Galactomannan assay of the BAL fluid may have augmented diagnostic value. A lack of clinical or radiographic response during empirical antifungal therapy may necessitate therapeutic drug level monitoring and/or diagnostic tissue sampling.³⁷⁰ Minimally invasive surgery (video-assisted thoracoscopic surgery) is associated with less morbidity than is open-lung biopsy.

Patients who have undergone HSCT and have suspected invasive mold infections should promptly begin taking a mold-active antifungal agent while diagnostic procedures are being arranged. A lack of clinical or radiographic response during proven infection may necessitate a switch to an agent from a different class or to combination therapy. Combination treatment of fungal infections with echinocandins, azoles, and polyene agents is common, whereas results of a large randomized clinical trial failed to show a benefit of anidulafungin added to voriconazole.³⁷¹ Echinocandin agents might be fungistatic rather than fungicidal in the case of mold infections because their interruption of cell wall synthesis is limited to actively growing hyphae. Echinocandins are not preferred for primary therapy for invasive aspergillosis but are used in salvage or combination therapy.

For documented invasive tissue mold infection, therapy is usually continued until some weeks (4–6) after lesions are resolved or stable, immunocompetence has improved, and the patient is afebrile. Although amphotericin B had been the gold standard antifungal agent since the 1960s, voriconazole produced superior outcomes in treatment for aspergillosis in 53% of patients, in contrast to 32% of patients treated with amphotericin B, followed by other antifungal therapy.³⁷² Treatment of CNS mold infections should include voriconazole, which, on the basis of a few patients studied, attains CSF levels approximately 50% of plasma levels and CNS tissue levels approximately 200% of plasma levels (see Chapter 40A). In a randomized study, isavuconazole was noninferior to voriconazole (with fewer adverse effects) for the primary treatment of suspected invasive mold infections in 527 patients with hematologic disorders, including those undergoing allogeneic HSCT.³⁷³ All-cause mortality at day 42 in these patients was similar (19% with isavuconazole vs. 20% with voriconazole).

After initial control of an *Aspergillus* infection, subsequent maintenance therapy for the duration of immunosuppression has been advocated to reduce the risk for reactivation. Multiple drug-drug interactions occur with the azoles, and adjustments may be required for immunosuppressive agents. Transient visual disturbances or hallucinations can occur with voriconazole. Difficulty in achieving therapeutic plasma drug levels complicates the administration of itraconazole and posaconazole. Itraconazole solution has improved oral bioavailability over the capsule and can be used, although blood level monitoring may be needed to ensure adequate absorption. Posaconazole (delayed-release version) and

isavuconazole have good antimold activity but are new to the market and may be expensive.

Aspergillosis of the CNS most frequently occurs among patients who have had refractory pulmonary disease despite antifungal treatment and has an associated poor prognosis; survival is approximately 24%.³⁷⁴

Concerns have arisen, predominantly in Europe, about azole resistance in *Aspergillus fumigatus*. As HSCT patients are at high risk, if these developments become generalized, an organized approach to mitigating risk might include intensified clinical awareness, antifungal susceptibility testing, or molecular diagnostics and genotyping.^{375,376}

Other Yeasts

Malassezia furfur causes tinea versicolor, catheter-related fungemia, and sometimes pneumonia.^{377–379} Response to either topical or systemic therapy is slow; recovery of granulocyte counts is usually associated with resolution.³⁷⁸ Catheter removal and discontinuation of IV lipids are important for a successful outcome in cases of fungemia. Trichosporonosis has manifested as fungemia, skin lesions, pneumonitis, and arthritis.^{380,381} Fungemia, usually acquired via an IV catheter, has been reported with *Trichosporon* and *Rhodotorula* spp., *Cryptococcus laurentii*, and *Hansenula anomala*. Meningitis with *Cryptococcus neoformans*, perhaps due to delayed recovery of CD4 cells and inverted CD4:CD8 ratio,³⁸² is unusual in contrast to its frequent occurrence among patients infected with HIV. Widespread anti-*Candida* prophylaxis with fluconazole may contribute to the low frequency of these infections.

Other Molds

Non-*Aspergillus* molds are rare but lead to high mortality.³⁸³ Non-*Aspergillus* molds, such as *Alternaria*, *Pseudallescheria/Scedosporium*, *Paecilomyces*, *Fusarium*, and *Phialophora* spp., are infrequent causes of invasive tissue infections whose clinical appearance is similar to that of *Aspergillus* infection. They are indistinguishable from *Aspergillus* hyphae in tissue sections; thus culture is required for identification. Disseminated fusariosis, most common within 100 days after allogeneic HSCT and associated with neutropenia, is a highly fatal infection for patients who have undergone HSCT, manifesting as positive blood cultures, skin lesions, or endophthalmitis.^{383–385} Whereas UCB HSCT might be associated with a higher incidence, more recent transplants (those transplanted after 2002) had a lower incidence.³⁸³ Successful resolution is usually associated with neutrophil recovery in addition to antifungal therapy.³⁸⁵ In the case of fusarial endophthalmitis, enucleation of the affected eye may be required.³⁸⁶

Mucormycosis is also uncommon after HSCT (allogeneic, 0.58%, and autologous, <0.1%, at 1 year), but it mimics aspergillosis clinically and may occur long after HSCT.^{387–389} The cause of one hepatic infection included over-the-counter herbal medication.³⁹⁰ Acute or cGVHD, steroid use, diabetes, advanced disease at the time of HSCT, and older age are risk factors for mucormycosis.³⁸³ Patients receiving voriconazole prophylaxis are at risk for breakthrough infection with invasive mucormycosis, but the risk for such breakthrough infections appears low.^{391–394} Posaconazole may be an effective maintenance treatment for infections caused by certain species of Mucorales after response to amphotericin B.³⁹⁵ Isavuconazole is FDA approved for the treatment of invasive mucormycosis in addition to treatment of invasive aspergillosis, but the approval is controversial because of the small number of patients: 21 with initial therapy and 16 with salvage therapy.³⁹⁶

Clinically significant infections caused by the dimorphic fungi, including coccidioidomycosis, histoplasmosis, and blastomycosis, are unusual even in hyperendemic areas of the United States. A review of the incidence of coccidioidomycosis in HSCT recipients in an endemic area from 2003–13 revealed an overall incidence of 2.3% (11/426), with 8 (73%) being de novo infection, and an overall mortality of 55%.³⁹⁷

Parasitic Infections

Parasitic infection after HSCT usually manifests as reactivation of toxoplasmosis (mostly reported in United States and Europe),³⁹⁸ although Chagas disease (Latin America),^{399–403} malaria (Africa),^{404–409} strongyloidiasis,⁴¹⁰ schistosomiasis,⁴¹¹ *Clonorchis* infection,⁴¹² giardiasis,⁴¹³ cryptosporidiosis,^{414–417} pulmonary microsporidiosis (now classified as

fungus),^{418–420} leishmaniasis (Mediterranean countries, Latin America), and *Acanthamoeba* and *Trichomonas* meningoencephalitis^{421–424} have also been reported. Routine blood smears before HSCT cannot be used to rule out malarial transmission. In Hong Kong, *Clonorchis sinensis* infection was identified in only 1% of screening stool examinations performed 7 days before HSCT.⁴¹² None of the patients had symptoms related to clonorchiasis; patients received praziquantel (25 mg/kg orally three times for 1 day) before HSCT, and subsequent stool examinations did not reveal the presence of ova.

Toxoplasmosis is infrequent after HSCT, occurring in 2% to 7% of patients who are seropositive before transplantation.^{425–428} Although the parasite can be transmitted as a primary infection through marrow, blood products, or donor solid organs, toxoplasmosis in patients who have undergone HSCT is mostly the result of reactivation of prior infection.⁴²⁹ GVHD is a risk factor for the suppression of cell-mediated immunity that is critical for host defense against *Toxoplasma gondii*.⁴³⁰ The clinical presentation includes fever, headaches, encephalitis with focal cerebral lesions, meningitis, pneumonitis, myocarditis, chorioretinitis, pancytopenia, hepatosplenomegaly, or lymphadenopathy.⁴³¹ Most cases of toxoplasmosis occur within 100 days after HSCT but also reactivate when *Pneumocystis* prophylaxis is complete or interrupted. Immunosuppression can affect the enhancement of CNS lesions. Parasitemia is a feature of reactivation that may be identified in tissue culture, although many diagnoses are now made with PCR assay.^{432,433} In countries where the prevalence of this infection is high, pretransplant detection of DNA using PCR assay may indicate those at highest risk for infection after transplant.⁴³⁴ Stereotactic brain biopsy is also useful in diagnosis of the infection.⁴³⁵ Mild increase in protein or mononuclear cells in CSF is common, and detecting DNA of *Toxoplasma* is highly specific but less sensitive for infection. Morphologically seeing tachyzoites is very rare. The identifiable risk period is 2 to 8 weeks after HSCT, and toxoplasmosis is more common among those receiving myeloablative conditioning.⁴³⁶ Seropositive patients not receiving TMP-SMX (160 mg of TMP/800 mg of SMZ, three times a week) are at risk for breakthrough toxoplasmosis.⁴³⁷ Pyrimethamine or atovaquone can be considered for prophylaxis. For these patients *Toxoplasma* reactivation can be monitored through PCR assay during the first 1 to 3 months after HSCT.⁴³⁸ Serologic monitoring for seroconversion (emergence of specific IgA or IgM) or reactivation (>threefold increase in IgG) is not reliable and useful. However, in countries where the prevalence is low, routine prophylaxis is not justified. Treatment of infection includes long-term pyrimethamine and sulfadiazine (or clindamycin in intolerance) with folinic acid or high-dose TMP-SMZ. Avoiding contact with possible infected sources (e.g., animal-cat feces, undercooked meat, soil, and water) is important to prevent primary infection.⁴³⁹

METHODS OF IMMUNE SYSTEM RECONSTITUTION AFTER HSCT

Vaccination

Patients undergoing autologous or allogeneic HSCT eventually lose immunity to the common childhood diseases and should be serially reimmunized 6 and 24 months after transplantation (Table 307.4; see Fig. 307.1).^{266,440} The efficacy of vaccination is influenced by the time elapsed since transplantation, the nature of the hematopoietic graft, the presence of GVHD, and the use of serial immunization.⁴⁴¹ There have been no reports of exacerbation of GVHD after immunization of patients who underwent HSCT. A national survey of HSCT immunization practices revealed that vaccines were underutilized and schedules for revaccination varied.⁴⁴² To ensure compliance, it is recommended that all transplant recipients be immunized on the same schedule, regardless of cell source.⁴⁴³

Patients who have undergone HSCT should receive all indicated nonlive vaccines, regardless of transplant type or presence of GVHD. Such patients should be revaccinated every 10 years with the combined tetanus-diphtheria-pertussis toxoid. Vaccination with acellular pertussis does not generate enough immunity, so a more effective vaccine is needed.⁴⁴⁴ At 6 months they should also be immunized against polio by the inactivated intramuscular vaccine, *H. influenzae* type B, hepatitis B, and *S. pneumoniae*. If the patient was previously immunized, only

TABLE 307.4 Suggested Schedule for Vaccination After HSCT

VACCINE	TIME PERIOD FOR IMMUNIZATION AFTER HSCT		
	6–12 MONTHS	14 MONTHS	24 MONTHS
Inactivated Vaccines			
Diphtheria, tetanus, pertussis ⁴⁴⁴	X	X	X
<i>Haemophilus influenzae</i> serotype B conjugate	X	X	X
Hepatitis B ^{76,445}	X	X	X
Pneumococcal	X (13-valent conjugated)		X ^a (13-valent conjugated)
Inactivated poliovirus	X	X	X
Influenza ⁴⁴⁶	Lifelong, seasonal administration of inactivated vaccine, beginning before HSCT and resuming ≥6 mo after HSCT, is recommended		
Hepatitis A	Routine administration is not indicated. If hepatitis A vaccination is given, two doses, given 6–12 mo apart, are required		
Varicella recombinant (Shingrix) ^{450–452}	US Food and Drug Administration approved in 2017		
Meningococcal conjugate	Routine administration is not indicated		
Human papillomavirus ⁴⁴⁷	Administration is recommended for men and women younger than 27 yr		
Rabies	Routine administration is not indicated; any decision to use should be individualized		
Live Replication-Competent Vaccines			
Measles-mumps-rubella	Administration is not indicated		X ^{a,b}
Varicella-zoster (Varivax) ⁴⁴⁹	Administration is not indicated		X ^{a,b}
Yellow fever virus	Routine administration is not indicated; any decision to use should be individualized		

^aOptional.^bIn patients with no active graft-versus-host disease or immunosuppressive therapy. HSCT, Hematopoietic stem cell transplantation.

one dose of hepatitis B vaccine should be given. Postvaccine titers of hepatitis B virus should be documented to ensure response and adequate protection, even when the vaccine is given at the specified time interval after HSCT.⁴⁴⁵ At 2 years a second dose of pneumococcal vaccine is optional; it provides a second opportunity to vaccinate persons who failed to respond to the first dose, especially patients with cGVHD. Lifelong, seasonal administration of influenza vaccine should begin before HSCT and resume by 2 to 6 months after HSCT. Children younger than 9 years who are receiving influenza vaccination for the first time require two doses yearly. Adults do not benefit from receiving two influenza vaccine doses.⁴⁴⁶ Influenza vaccine for HSCT ward employees, clinical health care workers, and household contacts may be especially necessary within the first year or for patients with ongoing GVHD, in whom protective responses may be impaired. For health care workers directly caring for HSCT recipients, the live-attenuated intranasal vaccine should be avoided to limit inadvertent exposure to patients. Vaccination with the quadrivalent human papillomavirus vaccine has not been studied but may be appropriate for male and female long-term transplant recipients ages 9 to 26 years and perhaps for women older than 26 years who are at risk for squamous intraepithelial lesions.^{447,448}

Live-virus vaccines, such as measles-mumps-rubella (MMR) and varicella, should not be given to transplant recipients with active GVHD or ongoing immunosuppressive therapy; the first doses are given to transplant recipients more than 24 months after HSCT who are taking no immunosuppressive medications and are presumed immunocompetent. A second MMR dose should be given 6 to 12 months later; however, the benefit of a second dose in this population has not been evaluated.

Vaccination against varicella is desired. Often, vaccination with live-attenuated VZV vaccine (Varivax, not Zostavax) is used for VZV-seronegative patients who no longer require immunosuppressive therapy and are free of GVHD.⁴⁴⁹ When varicella vaccination is given to persons older than 13 years, two doses, given 4 to 8 weeks apart, are required. Vaccination of HSCT recipients who have developed reactivation shingles after transplantation is not needed. Susceptible family members should receive VZV vaccine to minimize

chickenpox exposure for VZV-seronegative transplant recipients. In 2017 a new nonlive vaccine was licensed for use (Shingrix).^{450,451} Adoption of this vaccine will probably be forthcoming by many transplant centers.^{452–454}

Routine administration of hepatitis A, meningococcal, and rabies vaccines is not indicated. Hepatitis A vaccine is recommended for transplant recipients with chronic liver disease, including hepatitis C infection or cGVHD, or who live in hepatitis A-endemic areas or in areas experiencing outbreaks. If given, hepatitis A vaccination requires two doses, given 6 to 12 months apart. For transplant recipients with potential occupational exposure to rabies, preexposure rabies vaccination should be delayed until at least 12 months, if not 24 months, after HSCT.

Immunoglobulin Replacement

The major defect in humoral immunity is the absence of specific antibody production. Antibody levels in the first year after HSCT are affected primarily by pretransplantation levels in the recipient and, to a lesser degree, in the donor.⁴⁵⁵ Among patients with cGVHD, reduced production of opsonizing antibody and of all classes of IgG and IgA antibodies is seen.⁴⁵⁶ This immunodeficiency is further complicated by poor splenic function and is associated with risks for recurrent pneumococcal infections and episodes of bronchitis or pneumonia. IVIG does not prevent infections when given weekly during the preengraftment or late risk periods, but it does reduce rates of septicemia and localized infection when given in the postengraftment risk period after transplantation.^{457–459} It may modulate the severity of GVHD.⁴⁶⁰ Replacement IVIG (200–500 mg/kg every 1–2 weeks) may be beneficial for patients with IgG levels less than 400 mg/dL; however, in one prospective trial routine use of IVIG delayed the recovery of antigen (viral)-specific immunity.⁴⁵⁸

The role of hyperimmune globulin for prevention of specific infections is less clear. High-titer CMV globulin for prevention of CMV infection and treatment of end-organ CMV disease has proved to be of clear benefit in comparison with IVIG. However, antiviral drugs are effective in providing protection against CMV disease. Therefore, because of its limited availability and cost considerations, the use of CMV-specific

globulin is minimal at most transplantation centers. Hyperimmune RSV globulin provided only a very modest increase in neutralizing antibody when given in the first 6 weeks after HSCT.³¹¹ Hepatitis B, human rabies, and tetanus immune globulin should be used as needed in the event of exposures. VariZIG is a human polyclonal IgG

commercially available for intramuscular or IV administration for patients at high risk within the first 10 days of exposure.⁴⁶¹ VariZIG administration may extend the varicella incubation period from 10 days to as much as 28 days, is expensive, and is not uniformly effective in preventing chickenpox in patients who have undergone HSCT.

Key References

The complete reference list is available online at Expert Consult.

9. Young JA, Logan BR, Wu J, et al. Infections after transplantation of bone marrow or peripheral blood stem cells from unrelated donors. *Biol Blood Marrow Transplant.* 2016;22:359–370.
62. van Burik JA, Carter SL, Freifeld AG, et al. Higher-risk of cytomegalovirus and *Aspergillus* infections in recipients of T cell depleted unrelated bone marrow: analysis of infectious complications in patients treated with T cell depletion versus immune suppressive therapy to prevent graft-versus-host disease. *Biol Blood Marrow Transplant.* 2007;13:1487–1498.
70. Versluis J, Pas SD, Agteresch HJ, et al. Hepatitis E virus: an underestimated opportunistic pathogen in recipients of allogeneic hematopoietic stem cell transplantation. *Blood.* 2013;122:1079–1086.
72. Lau GK, He ML, Fong DY, et al. Preemptive use of lamivudine reduces hepatitis B exacerbation after allogeneic hematopoietic cell transplantation. *Hepatology.* 2002;36:702–709.
80. Kyvernitakis A, Mahale P, Popat UR, et al. Hepatitis C virus infection in patients undergoing hematopoietic cell transplantation in the era of direct-acting antiviral agents. *Biol Blood Marrow Transplant.* 2016;22:717–722.
93. Alonso CD, Treadway SB, Hanna DB, et al. Epidemiology and outcomes of *Clostridium difficile* infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis.* 2012;54:1053–1063.
101. Campbell AP, Guthrie KA, Englund JA, et al. Clinical outcomes associated with respiratory virus detection before allogeneic hematopoietic stem cell transplant. *Clin Infect Dis.* 2015;61:192–202.
105. Taur Y, Xavier JB, Lipuma L, et al. Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis.* 2012;55:905–914.
114. Wingard JR, Carter SL, Walsh TJ, et al. Randomized, double-blind trial of fluconazole versus voriconazole for prevention of invasive fungal infection after allogeneic hematopoietic cell transplantation. *Blood.* 2010;116:5111–5118.
115. van Burik JA, Ratanatharathorn V, Stepan DE, et al. Micafungin versus fluconazole for prophylaxis against invasive fungal infections during neutropenia in patients undergoing hematopoietic stem cell transplantation. *Clin Infect Dis.* 2004;39:1407–1416.
123. Ullmann AJ, Lipton JH, Vesole DH, et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med.* 2007;356:335–347.
136. Boeckh M, Nichols WG, Chemaly RF, et al. Valganciclovir for the prevention of complications of late cytomegalovirus infection after allogeneic hematopoietic cell transplantation: a randomized trial. *Ann Intern Med.* 2015;162:1–10.
141. Junghans C, Boeckh M, Carter RA, et al. Incidence and outcome of cytomegalovirus infections following nonmyeloablative compared with myeloablative allogeneic stem cell transplantation, a matched control study. *Blood.* 2002;99:1978–1985.
155. Vydra J, Shanley RM, George I, et al. Enterococcal bacteremia is associated with increased risk of mortality in recipients of allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis.* 2012;55:764–770.
170. Bock AM, Cao Q, Ferrieri P, et al. Bacteremia in blood or marrow transplantation patients: clinical risk factors for infection and emerging antibiotic resistance. *Biol Blood Marrow Transplant.* 2013;19:102–108.
181. Razonable RR, Litzow MR, Khaliq Y, et al. Bacteremia due to viridans group streptococci with diminished susceptibility to levofloxacin among neutropenic patients receiving levofloxacin prophylaxis. *Clin Infect Dis.* 2002;34:1469–1474.
187. Alonso CD, Treadway SB, Hanna DB, et al. Epidemiology and outcomes of *Clostridium difficile* infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis.* 2012;54:1053–1063.
211. Harrington RD, Woolfrey AE, Bowden R, et al. Legionellosis in a bone marrow transplant center. *Bone Marrow Transplant.* 1996;18:361–368.
213. van Burik JA, Hackman R, Nadeem S, et al. Nocardiosis after bone marrow transplantation: a retrospective study. *Clin Infect Dis.* 1997;24:1154–1160.
225. Teira P, Battiwalla M, Ramanathan M, et al. Early cytomegalovirus reactivation remains associated with increased transplant related mortality in the current era: a CIBMTR analysis. *Blood.* 2016;127:2427–2438.
266. Ljungman P, Cordonnier C, Einsele H, et al. Vaccination of hematopoietic cell transplant recipients. *Bone Marrow Transplant.* 2009;44:521–526.
287. Ustun C, Slaby J, Shanley RM, et al. Human parainfluenza virus infection after hematopoietic stem cell transplantation: risk factors, management, mortality, and changes over time. *Biol Blood Marrow Transplant.* 2012;18:1580–1588.
306. Boeckh M, Englund J, Li Y, et al. Randomized controlled multicenter trial of aerosolized ribavirin for respiratory syncytial virus upper respiratory tract infection in hematopoietic cell transplant recipients. *Clin Infect Dis.* 2007;44:245–249.
310. Vu D, Peck AJ, Nichols WG, et al. Safety and tolerability of oseltamivir prophylaxis in hematopoietic stem cell transplant recipients: a retrospective case-control study. *Clin Infect Dis.* 2007;45:187–193.
323. Zerr DM, Corey L, Kim HW, et al. Clinical outcomes of human herpesvirus 6 reactivation after hematopoietic stem cell transplantation. *Clin Infect Dis.* 2005;40:932–940.
330. Zerr DM, Fann JR, Breiger D, et al. HHV-6 reactivation and its effect on delirium and cognitive functioning in hematopoietic cell transplantation recipients. *Blood.* 2011;117:5243–5249.
337. Eid AJ, Brown RA, Patel R, et al. Parvovirus B19 infection after transplantation: a review of 98 cases. *Clin Infect Dis.* 2006;43:40–48.
344. van Burik JA, Leisenring W, Myerson D, et al. The effect of prophylactic fluconazole on the clinical spectrum of fungal diseases in bone marrow transplant recipients with special attention to hepatic candidiasis: an autopsy study of 355 patients. *Medicine (Baltimore).* 1998;77:246–254.
353. Williams KM, Ahn KW, Chen M, et al. The incidence, mortality and timing of *Pneumocystis jirovecii* pneumonia after hematopoietic cell transplantation: a CIBMTR analysis. *Bone Marrow Transplant.* 2016;51:573–580.
371. Marr KA, Schlamm HT, Herbrecht R, et al. Combination antifungal therapy for invasive aspergillosis: a randomized trial. *Ann Intern Med.* 2015;162:81–89.
392. Marty FM, Cosimi LA, Baden LR. Breakthrough zygomycosis after voriconazole treatment in recipients of hematopoietic stem-cell transplants. *N Engl J Med.* 2004;350:950–952.
397. Mendoza N, Noel P, Blair JE. Diagnosis, treatment, and outcomes of coccidioidomycosis in allogeneic stem cell transplantation. *Transpl Infect Dis.* 2015;17:380–388.
440. Small TN, Cowan MJ. Immunization of hematopoietic stem cell transplant recipients against vaccine-preventable diseases. *Expert Rev Clin Immunol.* 2011;7:193–203.
446. Karras NA, Weeres M, Sessions W, et al. A randomized trial of one versus two doses of influenza vaccine after allogeneic transplantation. *Biol Blood Marrow Transplant.* 2013;19:109–116.

References

- Anasetti C, Logan BR, Lee SJ, et al. Peripheral-blood stem cells versus bone marrow from unrelated donors. *N Engl J Med*. 2012;367:1487–1496.
- Ballen KK, Koreth J, Chen YB, et al. Selection of optimal alternative graft source: mismatched unrelated donor, umbilical cord blood, or haploidentical transplant. *Blood*. 2012;119:1972–1980.
- Dvorak CC, Cowan MJ. Hematopoietic stem cell transplantation for primary immunodeficiency disease. *Bone Marrow Transplant*. 2008;41:119–126.
- Burt RK, Loh Y, Pearce W, et al. Clinical applications of blood-derived and marrow-derived stem cells for nonmalignant diseases. *JAMA*. 2008;299:925–936.
- Gupta V, Tomblyn M, Pedersen TL, et al. Allogeneic hematopoietic cell transplantation in human immunodeficiency virus-positive patients with hematologic disorders: a report from the Center for International Blood and Marrow Transplant Research. *Biol Blood Marrow Transplant*. 2009;15:864–871.
- Gooley TA, Chien JW, Pergam SA, et al. Reduced mortality after allogeneic hematopoietic cell transplantation. *N Engl J Med*. 2010;363:2091–2101.
- Brunstein CG, Wagner JE. Cord blood transplantation for adults. *Vox Sang*. 2006;91:195–205.
- de Lima M, McNiece I, Robinson SN, et al. Cord-blood engraftment with ex vivo mesenchymal-cell coculture. *N Engl J Med*. 2012;367:2305–2315.
- Young JA, Logan BR, Wu J, et al. Infections after transplantation of bone marrow or peripheral blood stem cells from unrelated donors. *Biol Blood Marrow Transplant*. 2016;22:359–370.
- Brunstein CG, Eapen M, Ahn KW, et al. Reduced-intensity conditioning transplantation in acute leukemia: the effect of source of unrelated donor stem cells on outcomes. *Blood*. 2012;119:5591–5598.
- Ballen K, Woo Ahn K, Chen M, et al. Infection rates among acute leukemia patients receiving alternative donor hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2016;22:1636–1645.
- Bejanyan N, Brunstein C, Cao Q, et al. Immune reconstitution (IR) after allogeneic hematopoietic cell transplantation (alloHCT): comparing results in recipients of unrelated umbilical cord blood (UCB) to those with an HLA-matched sibling donor peripheral blood (MSD PB). *Blood*. 2016;4590a.
- Servais S, Lengline E, Porcher R, et al. Long-term immune reconstitution and infection burden after mismatched hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2014;20:507–517.
- Lucchini G, Perales MA, Vey P. Immune reconstitution after cord blood transplantation: peculiarities, clinical implications and management strategies. *Cytotherapy*. 2015;17:711–722.
- Filicco J, Lazarus HM, Flomenberg N. Mucosal injury in patients undergoing hematopoietic progenitor cell transplantation: new approaches to prophylaxis and treatment. *Bone Marrow Transplant*. 2003;31:1–10.
- Chaudhury S, Auerbach AD, Kernan NA, et al. Fludarabine-based cytoreductive regimen and T-cell-depleted grafts from alternative donors for the treatment of high-risk patients with Fanconi anaemia. *Br J Haematol*. 2008;140:644–655.
- Macmillan ML, Couriel D, Weisdorf DJ, et al. A phase 2/3 multicenter randomized clinical trial of ABX-CBL versus ATG as secondary therapy for steroid-resistant acute graft-versus-host disease. *Blood*. 2007;109:2657–2662.
- Tsirigotis P, Triantafyllou K, Girkas K, et al. Keratinocyte growth factor is effective in the prevention of intestinal mucositis in patients with hematological malignancies treated with high-dose chemotherapy and autologous hematopoietic SCT: a video-capsule endoscopy study. *Bone Marrow Transplant*. 2008;42:337–343.
- Stiff P. Mucositis associated with stem cell transplantation: current status and innovative approaches to management. *Bone Marrow Transplant*. 2001;27(suppl 2):S3–S11.
- Spielberger R, Stiff P, Bensinger W, et al. Palifermin for oral mucositis after intensive therapy for hematologic cancers. *N Engl J Med*. 2004;351:2590–2598.
- Levine JE, Blazar BR, DeFor T, et al. Long-term follow-up of a phase I/II randomized, placebo-controlled trial of palifermin to prevent graft-versus-host disease after related donor allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2008;14:1017–1021.
- Blazar BR, Weisdorf DJ, DeFor T, et al. Phase 1/2 randomized, placebo-control trial of palifermin to prevent graft-versus-host disease after allogeneic hematopoietic stem cell transplantation. *Blood*. 2006;108:3216–3222.
- Majhail NS, Brunstein CG, Tomblyn M, et al. Reduced-intensity allogeneic transplant in patients older than 55 years: unrelated umbilical cord blood is safe and effective for patients without a matched related donor. *Biol Blood Marrow Transplant*. 2008;14:282–289.
- Warlick E, Ahn KW, Pedersen TL, et al. Reduced intensity conditioning is superior to nonmyeloablative conditioning for older chronic myelogenous leukemia patients undergoing hematopoietic cell transplant during the tyrosine kinase inhibitor era. *Blood*. 2012;119:4083–4090.
- Niederwieser D, Maris M, Shizuru JA, et al. Low-dose total body irradiation and fludarabine followed by hematopoietic cell transplantation from HLA-matched or mismatched unrelated donors and postgrafting immunosuppression with cyclosporine and mycophenolate mofetil can induce durable complete chimerism and sustained remissions in patients with hematologic diseases. *Blood*. 2003;101:1620–1629.
- Scott BL, Pasquini MC, Logan BR, et al. Myeloablative versus reduced-intensity hematopoietic cell transplantation for acute myeloid leukemia and myelodysplastic syndromes. *J Clin Oncol*. 2017;35:1154–1161.
- Petersdorf EW, Hansen JA, Martin PJ, et al. Major-histocompatibility-complex class I alleles and antigens in hematopoietic cell transplantation. *N Engl J Med*. 2001;345:1794–1800.
- Rubinstein P. HLA matching for bone marrow transplantation—how much is enough? *N Engl J Med*. 2001;345:1842–1844.
- Anasetti C, Amos D, Beatty PG, et al. Effect of HLA compatibility on engraftment of bone marrow transplants in patients with leukemia or lymphoma. *N Engl J Med*. 1989;320:197–204.
- Weisdorf D, Cooley S, Devine S, et al. T cell-depleted partial matched unrelated donor transplant for advanced myeloid malignancy: KIR ligand mismatch and outcome. *Biol Blood Marrow Transplant*. 2012;18:937–943.
- Tomblyn M, Young JA, Haagenson MD, et al. Decreased infections in recipients of unrelated donor hematopoietic cell transplantation from donors with an activating KIR genotype. *Biol Blood Marrow Transplant*. 2010;16:1155–1161.
- Freifeld AG, Bow EJ, Sepkowitz KA, et al. Infectious Diseases Society of America. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2011;52:427–431, and e56–e93.
- Tomblyn M, Chiller T, Einsele H, et al. Guidelines for preventing infectious complications among hematopoietic cell transplantation recipients: a global perspective. *Biol Blood Marrow Transplant*. 2009;15:1143–1238.
- Yokoe D, Casper C, Dubberke E, et al. Infection prevention and control in health-care facilities in which hematopoietic cell transplant recipients are treated. *Bone Marrow Transplant*. 2009;44:495–507.
- Ullmann AJ, Schmidt-Hieber M, Bertz H, et al. Infectious diseases in allogeneic haematopoietic stem cell transplantation: prevention and prophylaxis strategy guidelines 2016. *Ann Hematol*. 2016;95:1435–1455.
- Arnold DM, Crowther MA, Meyer RM, et al. Misleading hepatitis B test results due to intravenous immunoglobulin administration: implications for a clinical trial of rituximab in immune thrombocytopenia. *Transfusion*. 2010;50:2577–2581.
- Lunde LE, Dasaraju S, Cao Q, et al. Hemorrhagic cystitis after allogeneic hematopoietic cell transplantation: risk factors, graft source and survival. *Bone Marrow Transplant*. 2015;50:1432–1437.
- Korkmaz A, Topal T, Oter S. Pathophysiological aspects of cyclophosphamide and ifosfamide induced hemorrhagic cystitis; implication of reactive oxygen and nitrogen species as well as PARP activation. *Cell Biol Toxicol*. 2007;23:303–312.
- Yaghobi R, Ramzi M, Delghani S. The role of different risk factors in clinical presentation of hemorrhagic cystitis in hematopoietic stem cell transplant recipients. *Transplant Proc*. 2009;41:2900–2902.
- Dalianis T, Ljungman P. Full myeloablative conditioning and an unrelated HLA mismatched donor increase the risk for BK virus-positive hemorrhagic cystitis in allogeneic hematopoietic stem cell transplanted patients. *Anticancer Res*. 2011;31:939–944.
- Akiyama H, Kurosu T, Sakashita C, et al. Adenovirus is a key pathogen in hemorrhagic cystitis associated with bone marrow transplantation. *Clin Infect Dis*. 2001;32:1325–1330.
- Silva Lde P, Patash PA, Saliba RM, et al. Hemorrhagic cystitis after allogeneic hematopoietic stem cell transplants is the complex result of BK virus infection, preparative regimen intensity and donor type. *Haematologica*. 2010;95:1183–1190.
- Childs R, Sanchez C, Engler H, et al. High incidence of adenovirus- and polyomavirus-induced hemorrhagic cystitis in bone marrow allotransplantation for hematologic malignancy following T cell depletion and cyclosporine. *Bone Marrow Transplant*. 1998;22:889–893.
- Knoll GA, Humar A, Fergusson D, et al. Levofloxacin for BK virus prophylaxis following kidney transplantation: a randomized clinical trial. *JAMA*. 2014;312:2106–2114.
- Lee BT, Gabardi S, Grafals M, et al. Efficacy of levofloxacin in the treatment of BK viremia: a multicenter, double-blinded, randomized, placebo-controlled trial. *Clin J Am Soc Nephrol*. 2014;9:583–589.
- Cesaro S, Dalianis T, Hanssen Rinaldo C, et al. ECIL guidelines for the prevention, diagnosis and treatment of BK polyomavirus-associated haemorrhagic cystitis in haematopoietic stem cell transplant recipients. *J Antimicrob Chemother*. 2018;73:12–21.
- Priftakis P, Bogdanovic G, Kokhaei P, et al. BK virus quantification in urine samples of bone marrow transplanted patients is helpful for diagnosis of hemorrhagic cystitis, although wide individual variations exist. *J Clin Virol*. 2003;26:71–77.
- Ambalathingal GR, Francis RS, Smyth MJ, et al. BK polyomavirus: clinical aspects, immune regulation, and emerging therapies. *Clin Microbiol Rev*. 2017;30:503–528.
- Erard V, Kim HW, Corey L, et al. BK DNA viral load in plasma: evidence for an association with hemorrhagic cystitis in allogeneic hematopoietic cell transplant recipients. *Blood*. 2005;106:1130–1132.
- Bridges B, Donegan S, Badros A. Cidofovir bladder instillation for the treatment of BK hemorrhagic cystitis after allogeneic stem cell transplantation. *Am J Hematol*. 2006;81:535–537.
- Litzow MR, Repoussis PD, Schroeder G, et al. Veno-occlusive disease of the liver after blood and marrow transplantation: analysis of pre- and post-transplant risk factors associated with severity and results of therapy with tissue plasminogen activator. *Leuk Lymphoma*. 2002;43:2099–2107.
- Carreras E, Diaz-Beya M, Rosinol L, et al. The incidence of veno-occlusive disease following allogeneic hematopoietic stem cell transplantation has diminished and the outcome improved over the last decade. *Biol Blood Marrow Transplant*. 2011;17:1713–1720.
- Richardson PG, Riches ML, Kernan NA, et al. Phase 3 trial of defibrotide for the treatment of severe veno-occlusive disease and multi-organ failure. *Blood*. 2016;127:1656–1665.
- Carreras E. How I manage sinusoidal obstruction syndrome after haematopoietic cell transplantation. *Br J Haematol*. 2015;168:481–491.
- McDonald GB, Hinds MS, Fisher LD, et al. Veno-occlusive disease of the liver and multiorgan failure after bone marrow transplantation: a cohort study of 355 patients. *Ann Intern Med*. 1993;118:255–267.
- Echeveste TF, Pimpoll LG, Martinez JP, et al. Point shear wave elastography for the diagnosis of sinusoidal obstruction syndrome/venoocclusive disease (SOD/VOD): a prospective study. *Europena Radiol*. 2017;C-1271.
- Kis B, Pamarthi V, Fan CM, et al. Safety and utility of transjugular liver biopsy in hematopoietic stem cell transplant recipients. *J Vasc Interv Radiol*. 2013;24:85–89.
- Weisdorf D. Graft vs. host disease: pathology, prophylaxis and therapy GVHD overview. *Best Pract Res Clin Haematol*. 2008;21:99–100.
- Weisdorf D. Should all unrelated donors for transplantation be matched? *Best Pract Res Clin Haematol*. 2008;21:79–83.
- Weisdorf D, Zhang MJ, Arora M, et al. Graft-versus-host disease induced graft-versus-leukemia effect: greater impact on relapse and disease-free survival after reduced intensity conditioning. *Biol Blood Marrow Transplant*. 2012;18:1727–1733.
- Holtan SG, MacMillan ML. A risk-adapted approach to acute GVHD treatment: are we there yet? *Bone Marrow Transplant*. 2016;51:172–175.
- van Burik JA, Carter SL, Freifeld AG, et al. Higher-risk of cytomegalovirus and Aspergillus infections in recipients of T cell depleted unrelated bone marrow: analysis of infectious complications in patients treated with T cell depletion versus immune suppressive therapy to prevent graft-versus-host disease. *Biol Blood Marrow Transplant*. 2007;13:1487–1498.
- Saad A, Lamb LS. Ex vivo T-cell depletion in allogeneic hematopoietic stem cell transplant: past, present and future. *Bone Marrow Transplant*. 2017;52:1241–1248.
- Luznik L, O'Donnell PV, Symons HJ, et al. HLA-haploidentical bone marrow transplantation for hematologic malignancies using nonmyeloablative

- conditioning and high-dose, posttransplantation cyclophosphamide. *Biol Blood Marrow Transplant.* 2008;14:641–650.
65. Kim SS. Treatment options in steroid-refractory acute graft versus-host disease following hematopoietic stem cell transplantation. *Ann Pharmacother.* 2007;41:1436–1444.
66. Fraser CJ, Scott Baker K. The management and outcome of chronic graft-versus-host disease. *Br J Haematol.* 2007;138:131–145.
67. Levitsky J, Sorrell ME. Hepatic complications of hematopoietic cell transplantation. *Curr Gastroenterol Rep.* 2007;9:60–65.
68. Arai S, Lee LA, Vogelsang GB. A systematic approach to hepatic complications in hematopoietic stem cell transplantation. *J Hematother Stem Cell Res.* 2002;11:215–229.
69. Ustun C, Koc H, Karayalcin S, et al. Hepatitis B virus infection in allogeneic bone marrow transplantation. *Bone Marrow Transplant.* 1997;20:289–296.
70. Versluis J, Pas SD, Agteresch HJ, et al. Hepatitis E virus: an underestimated opportunistic pathogen in recipients of allogeneic hematopoietic stem cell transplantation. *Blood.* 2013;122:1079–1086.
71. Kamar N, Izopet J, Tripot S, et al. Ribavirin for chronic hepatitis E virus infection in transplant recipients. *N Engl J Med.* 2014;370:1111–1120.
72. Lau GK, He ML, Fong DY, et al. Preemptive use of lamivudine reduces hepatitis B exacerbation after allogeneic hematopoietic cell transplantation. *Hepatology.* 2002;36:702–709.
73. Di Bisceglie AM, Lok AS, Martin P, et al. Recent US Food and Drug Administration warnings on hepatitis B reactivation with immune-suppressing and anticancer drugs: just the tip of the iceberg? *Hepatology.* 2015;61:703–711.
74. Tomblyn M, Chen M, Kukreja M, et al. No increased mortality from donor or recipient hepatitis B- and/or hepatitis C-positive serostatus after related-donor allogeneic hematopoietic cell transplantation. *Transpl Infect Dis.* 2012;14:468–478.
75. Ustun C, Idilman R, Gurman G, et al. Hematopoietic stem cell transplantation from non-replicative hepatitis B virus carriers is safe. *J Hepatol.* 1999;31:202–209.
76. Idilman R, Ustun C, Karayalcin S, et al. Hepatitis B virus vaccination of recipients and donors of allogeneic peripheral blood stem cell transplantation. *Clin Transplant.* 2003;17:438–443.
77. Lau GK, Leung YH, Fong DY, et al. High hepatitis B virus DNA viral load as the most important risk factor for HBV reactivation in patients positive for HBV surface antigen undergoing autologous hematopoietic cell transplantation. *Blood.* 2002;99:2324–2330.
78. Vance EA, Soiffer RJ, McDonald GB, et al. Prevention of transmission of hepatitis C virus in bone marrow transplantation by treating the donor with alpha-interferon. *Transplantation.* 1996;62:1358–1360.
79. Torres HA, Chong PP, De Lima M, et al. Hepatitis C virus infection among hematopoietic cell transplant donors and recipients: American Society for Blood and Marrow Transplantation Task Force recommendations. *Biol Blood Marrow Transplant.* 2015;21:1870–1882.
80. Kyvernitis A, Mahale P, Popat UR, et al. Hepatitis C virus infection in patients undergoing hematopoietic cell transplantation in the era of direct-acting antiviral agents. *Biol Blood Marrow Transplant.* 2016;22:717–722.
81. Rizza SA, Challagundla KB, Natesampillai S, et al. TRAIL dependent fratricidal killing of gp120 primed hepatocytes by HCV core expressing hepatocytes. *PLoS ONE.* 2011;6:e27171.
82. Peffault de Latur R, Levy V, Asselah T, et al. Long-term outcome of hepatitis C infection after bone marrow transplantation. *Blood.* 2004;103:1618–1624.
83. Ljungman P, Locasciulli A, de Soria VG, et al. Long-term follow-up of HCV-infected hematopoietic SCT patients and effects of antiviral therapy. *Bone Marrow Transplant.* 2012;47:1217–1221.
84. Solh M, Arat M, Cao Q, et al. Late-onset noninfectious pulmonary complications in adult allogeneic hematopoietic cell transplant recipients. *Transplantation.* 2011;91:798–803.
85. Majhail NS, Parks K, DeFor TE, et al. Diffuse alveolar hemorrhage and infection-associated alveolar hemorrhage following hematopoietic stem cell transplantation: related and high-risk clinical syndromes. *Biol Blood Marrow Transplant.* 2006;12:1038–1046.
86. Alrawi EB, Keklik F, Cao Q, et al. Umbilical cord blood transplantation and delay engraftment are associated with increased risks of diffuse alveolar hemorrhage. *Blood.* 2016;222a.
87. Weisdorf DJ. Diffuse alveolar hemorrhage: an evolving problem? *Leukemia.* 2003;17:1049–1050.
88. Panoskaltis-Mortari A, Griese M, Madtes DK, et al. An official American Thoracic Society research statement: noninfectious lung injury after hematopoietic stem cell transplantation: idiopathic pneumonia syndrome. *Am J Respir Crit Care Med.* 2011;183:1262–1279.
89. Yanik GA, Ho VT, Levine JE, et al. The impact of soluble tumor necrosis factor receptor etanercept on the treatment of idiopathic pneumonia syndrome after allogeneic hematopoietic stem cell transplantation. *Blood.* 2008;112:3073–3081.
90. Seo S, Renaud C, Kuypers JM, et al. Idiopathic pneumonia syndrome after hematopoietic cell transplantation: evidence of occult infectious etiologies. *Blood.* 2015;125:3789–3797.
91. Kamboj M, Mihiu CN, Sepkowitz K, et al. Work-up for infectious diarrhea after allogeneic hematopoietic stem cell transplantation: single specimen testing results in cost savings without compromising diagnostic yield. *Transpl Infect Dis.* 2007;9:265–269.
92. van Kraaij MG, Dekker AW, Verdonck LF, et al. Infectious gastroenteritis: an uncommon cause of diarrhoea in adult allogeneic and autologous stem cell transplant recipients. *Bone Marrow Transplant.* 2000;26:299–303.
93. Alonso CD, Treadway SB, Hanna DB, et al. Epidemiology and outcomes of *Clostridium difficile* infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis.* 2012;54:1053–1063.
94. Zacharioudakis IM, Ziakas PD, Mylonakis E. *Clostridium difficile* infection in the hematopoietic unit: a meta-analysis of published studies. *Biol Blood Marrow Transplant.* 2014;S1083–S1879.
95. Davila ML. Neutropenic enterocolitis. *Curr Opin Gastroenterol.* 2006;22:44–47.
96. Canninga-Van Dijk MR, Sanders CJ, Verdonck LF, et al. Differential diagnosis of skin lesions after allogeneic haematopoietic stem cell transplantation. *Histopathology.* 2003;42:313–330.
97. van Burik JA, Colven R, Spach D. Cutaneous aspergillosis. *J Clin Microbiol.* 1998;36:3115–3121.
98. Ferra C, Doebbeling BN, Hollis RJ, et al. *Candida tropicalis* vertebral osteomyelitis: a late sequela of fungemia. *Clin Infect Dis.* 1994;19:697–703.
99. Miyakis S, Velegaki A, Delikou S, et al. Invasive *Acremonium strictum* infection in a bone marrow transplant recipient. *Pediatr Infect Dis J.* 2006;25:273–275.
100. Barasch A, Mosier KM, D'Ambrosio JA, et al. Posttransplantation osteomyelitis in a bone marrow transplant recipient. *Oral Surg Oral Med Oral Pathol.* 1993;75:391–396.
101. Campbell AP, Guthrie KA, Englund JA, et al. Clinical outcomes associated with respiratory virus detection before allogeneic hematopoietic stem cell transplant. *Clin Infect Dis.* 2015;61:192–202.
102. van Montfrans J, Schulz L, Versluis B, et al. Viral PCR positivity in stool before allogeneic hematopoietic cell transplantation is strongly associated with acute intestinal graft-versus-host disease. *Biol Blood Marrow Transplant.* 2015;21:772–774.
103. Scott BL, Park JY, Deeg HJ, et al. Pretransplant neutropenia is associated with poor-risk cytogenetic features and increased infection-related mortality in patients with myelodysplastic syndromes. *Biol Blood Marrow Transplant.* 2008;14:799–806.
104. Engelhard D, Akova M, Boeckh MJ, et al. Bacterial infection prevention after hematopoietic cell transplantation. *Bone Marrow Transplant.* 2009;44:467–470.
105. Taur Y, Xavier JB, Lipuma L, et al. Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis.* 2012;55:905–914.
106. Collin BA, Leather HL, Wingard JR, et al. Evolution, incidence, and susceptibility of bacterial bloodstream isolates from 519 bone marrow transplant patients. *Clin Infect Dis.* 2001;33:947–953.
107. Marena C, Zecca M, Carenni ML, et al. Incidence of, and risk factors for, nosocomial infections among hematopoietic stem cell transplantation recipients, with impact on procedure-related mortality. *Infect Control Hosp Epidemiol.* 2001;22:510–517.
108. Oliveira AL, de Souza M, Carvalho-Dias VM, et al. Epidemiology of bacteremia and factors associated with multi-drug-resistant gram-negative bacteremia in hematopoietic stem cell transplant recipients. *Bone Marrow Transplant.* 2007;39:775–781.
109. Wade JC, Newton B, Flournoy N, et al. Oral acyclovir for prevention of herpes simplex virus reactivation after marrow transplantation. *Ann Intern Med.* 1984;100:823–828.
110. Zaia J, Baden L, Boeckh MJ, et al. Viral disease prevention after hematopoietic cell transplantation. *Bone Marrow Transplant.* 2009;44:471–482.
111. Goodman JL, Winston DJ, Greenfield RA, et al. A controlled trial of fluconazole to prevent fungal infections in patients undergoing bone marrow transplantation. *N Engl J Med.* 1992;326:845–851.
112. MacMillan ML, Goodman JL, DeFor TE, et al. Fluconazole to prevent yeast infections in bone marrow transplantation patients: a randomized trial of high versus reduced dose, and determination of the value of maintenance therapy. *Am J Med.* 2002;112:369–379.
113. Slavin MA, Osborne B, Adams R, et al. Efficacy and safety of fluconazole prophylaxis for fungal infections after marrow transplantation—a prospective, randomized, double-blind study. *J Infect Dis.* 1995;171:1545–1552.
114. Wingard JR, Carter SL, Walsh TJ, et al. Randomized, double-blind trial of fluconazole versus voriconazole for prevention of invasive fungal infection after allogeneic hematopoietic cell transplantation. *Blood.* 2010;116:5111–5118.
115. van Burik JA, Ratanatharathorn V, Stepan DE, et al. Micafungin versus fluconazole for prophylaxis against invasive fungal infections during neutropenia in patients undergoing hematopoietic stem cell transplantation. *Clin Infect Dis.* 2004;39:1407–1416.
116. Hachem R, Hanna H, Kontoyiannis D, et al. The changing epidemiology of invasive candidiasis: *Candida glabrata* and *Candida krusei* as the leading causes of candidemia in hematologic malignancy. *Cancer.* 2008;112:2493–2499.
117. Marr KA, Bow E, Chiller T, et al. Fungal infection prevention after hematopoietic cell transplantation. *Bone Marrow Transplant.* 2009;44:483–487.
118. Wingard JR, Hiemenz JW, Jantz MA. How I manage pulmonary nodular lesions and nodular infiltrates in patients with hematologic malignancies or undergoing hematopoietic cell transplantation. *Blood.* 2012;120:1791–1800.
119. Hübel K, Carter R, Liles W, et al. Granulocyte transfusion therapy for infections in candidates and recipients of hematopoietic cell transplantation: a comparative analysis of feasibility and outcome of community donors versus related donors. *Transfusion.* 2002;42:1414–1421.
120. Lazarus HM, Magalhaes-Silverman M, Fox RM, et al. Contamination during in vitro processing of bone marrow for transplantation: clinical significance. *Bone Marrow Transplant.* 1991;7:241–246.
121. Nasser RM, Hajjar I, Sandhaus LM, et al. Routine cultures of bone marrow and peripheral stem cell harvests: clinical impact, cost analysis, and review. *Clin Infect Dis.* 1998;27:886–888.
122. Wald A, Leisenring W, van Burik JA, et al. Epidemiology of *Aspergillus* infections in a large cohort of patients undergoing bone marrow transplantation. *J Infect Dis.* 1997;175:1459–1466.
123. Ullmann AJ, Lipton JH, Vesole DH, et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med.* 2007;356:335–347.
124. Tuan IZ, Dennison D, Weisdorf DJ. *Pneumocystis carinii* pneumonia following bone marrow transplantation. *Bone Marrow Transplant.* 1992;10:267–272.
125. Link H, Vohringer HF, Wingen F, et al. Pentamidine aerosol for prophylaxis of *Pneumocystis carinii* pneumonia after BMT. *Bone Marrow Transplant.* 1993;11:403–406.
126. Souza JB, Boeckh M, Gooley TA, et al. High rates of *Pneumocystis carinii* pneumonia in allogeneic blood and marrow transplant recipients receiving dapsone prophylaxis. *Clin Infect Dis.* 1999;29:1467–1471.
127. Fishman JA. Prevention of infection caused by *Pneumocystis carinii* in transplant recipients. *Clin Infect Dis.* 2001;33:1397–1405.
128. Sangiolo D, Storer B, Nash R, et al. Toxicity and efficacy of daily dapsone as *Pneumocystis jirovecii* prophylaxis after hematopoietic stem cell transplantation: a case-control study. *Biol Blood Marrow Transplant.* 2005;11:521–529.
129. Yakushiji K, Gondo H, Kamezaki K, et al. Monitoring of cytomegalovirus reactivation after allogeneic stem cell transplantation: comparison of an antigenemia assay and quantitative real-time polymerase chain reaction. *Bone Marrow Transplant.* 2002;29:599–606.
130. Boeckh M, Gooley TA, Myerson D, et al. Cytomegalovirus pp65 antigenemia-guided early treatment with ganciclovir versus ganciclovir at engraftment after allogeneic marrow transplantation: a randomized double-blind study. *Blood.* 1996;88:4063–4071.
131. Walker CM, van Burik JA, DeFor TE, et al. Cytomegalovirus infection after allogeneic transplantation: comparison of cord blood with peripheral blood and marrow graft sources. *Biol Blood Marrow Transplant.* 2007;13:1106–1115.
132. Munoz-Cobo B, Solano C, Costa E, et al. Dynamics of cytomegalovirus (CMV) plasma DNAemia in initial and

- recurrent episodes of active CMV infection in the allogeneic stem cell transplantation setting: implications for designing preemptive antiviral therapy strategies. *Biol Blood Marrow Transplant*. 2011;17:1602–1611.
133. Sullivan KM, Mori M, Sanders J, et al. Late complications of allogeneic and autologous marrow transplantation. *Bone Marrow Transplant*. 1992;10(suppl 1):127–134.
 134. Boeckh M, Leisenring W, Riddell SR, et al. Late cytomegalovirus disease and mortality in recipients of allogeneic hematopoietic stem cell transplants: importance of viral load and T-cell immunity. *Blood*. 2003;101:407–414.
 135. Beck JC, Wagner JE, DeFor TE, et al. Impact of cytomegalovirus reactivation after umbilical cord blood transplantation. *Biol Blood Marrow Transplant*. 2010;16:215–222.
 136. Boeckh M, Nichols WG, Chemaly RF, et al. Valganciclovir for the prevention of complications of late cytomegalovirus infection after allogeneic hematopoietic cell transplantation: a randomized trial. *Ann Intern Med*. 2015;162:1–10.
 137. Ochs L, Shu XO, Miller J, et al. Late infections after allogeneic bone marrow transplantations: comparison of incidence in related and unrelated donor transplant recipients. *Blood*. 1995;86:3979–3986.
 138. Mackall C, Fry T, Gress R, et al. Background to hematopoietic cell transplantation, including post transplant immune recovery. *Bone Marrow Transplant*. 2009;44:457–462.
 139. Robin M, Porcher R, De Castro Araujo R, et al. Risk factors for late infections after allogeneic hematopoietic stem cell transplantation from a matched related donor. *Biol Blood Marrow Transplant*. 2007;13:1304–1312.
 140. Junghanss C, Marr KA, Carter RA, et al. Incidence and outcome of bacterial and fungal infections following nonmyeloablative compared with myeloablative allogeneic hematopoietic stem cell transplantation: a matched control study. *Biol Blood Marrow Transplant*. 2002;8:512–520.
 141. Junghanss C, Boeckh M, Carter RA, et al. Incidence and outcome of cytomegalovirus infections following nonmyeloablative compared with myeloablative allogeneic stem cell transplantation, a matched control study. *Blood*. 2002;99:1978–1985.
 142. Hagen EA, Stern H, Porter D, et al. High rate of invasive fungal infections following nonmyeloablative allogeneic transplantation. *Clin Infect Dis*. 2003;36:9–15.
 143. Yokoe D, Casper C, Dubberke E, et al. Safe living after hematopoietic cell transplantation. *Bone Marrow Transplant*. 2009;44:509–519.
 144. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings: recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *Infect Control Hosp Epidemiol*. 2002;23(suppl 12):S3–S40.
 145. Farowski F, Buckner V, Vehreschild JJ, et al. Impact of choice, timing, sequence and combination of broad-spectrum antibiotics on the outcome of allogeneic haematopoietic stem cell transplantation. *Bone Marrow Transplant*. 2018;53:52–57.
 146. Routy B, Letendre C, Enot D, et al. The influence of gut decontamination prophylactic antibiotics on acute graft-versus-host disease and survival following allogeneic hematopoietic stem cell transplantation. *Oncoimmunology*. 2017;6:e125806.
 147. Bilinski J, Robak K, Peric Z, et al. Impact of gut colonization by antibiotic-resistant bacteria on the outcomes of allogeneic hematopoietic stem cell transplantation: a retrospective, single-center study. *Biol Blood Marrow Transplant*. 2016;22:1087–1093.
 148. Averbuch D, Tridello G, Hoek J, et al. Antimicrobial resistance in gram-negative rods causing bacteremia in hematopoietic stem cell transplant recipients: intercontinental prospective study of the Infectious Diseases Working Party of the European Bone Marrow Transplantation Group. *Clin Infect Dis*. 2017;65:1819–1828.
 149. Shaw BE, Boswell T, Byrne JL, et al. Clinical impact of MRSA in a stem cell transplant unit: analysis before, during and after an MRSA outbreak. *Bone Marrow Transplant*. 2007;39:623–629.
 150. Mikulska M, Viscoli C, Orasch C, et al. Aetiology and resistance in bacteraemias among adult and paediatric haematology and cancer patients. *J Infect*. 2014;68:321–331.
 151. Kameda K, Kimura S, Akahoshi Y, et al. High incidence of afebrile bloodstream infection detected by surveillance blood culture in patients on corticosteroid therapy after allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2016;22:371–377.
 152. Neshler L, Chemaly RF, Shah DP, et al. Utility of routine surveillance blood cultures in asymptomatic allogeneic hematopoietic stem cell transplant recipients with indwelling central venous catheters at a comprehensive cancer center. *Am J Infect Control*. 2014;42:1084–1088.
 153. Elishoov H, Or R, Strauss N, et al. Nosocomial colonization, septicemia, and Hickman/Broviac catheter-related infections in bone marrow transplant recipients: a 5-year prospective study. *Medicine (Baltimore)*. 1998;77:83–101.
 154. Engelhard D, Elishoov H, Strauss N, et al. Nosocomial coagulase-negative staphylococcal infections in bone marrow transplantation recipients with central vein catheter: a 5-year prospective study. *Transplantation*. 1996;61:430–434.
 155. Vydra J, Shanley RM, George I, et al. Enterococcal bacteremia is associated with increased risk of mortality in recipients of allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis*. 2012;55:764–770.
 156. Dandoy CE, Ardura MI, Papanicolaou GA, et al. Bacterial bloodstream infections in the allogeneic hematopoietic cell transplant patient: new considerations for a persistent nemesis. *Bone Marrow Transplant*. 2017;52:1091–1106.
 157. Metzger KE, Rucker Y, Callaghan M, et al. The burden of mucosal barrier injury laboratory-confirmed bloodstream infection among hematology, oncology, and stem cell transplant patients. *Infect Control Hosp Epidemiol*. 2015;36:119–124.
 158. Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. *Clin Microbiol Rev*. 2014;27:870–926.
 159. Tavazee M, Rybicki L, Mossad S, et al. Risk factors for vancomycin-resistant *Enterococcus* bacteremia and its influence on survival after allogeneic hematopoietic cell transplantation. *Bone Marrow Transplant*. 2014;49:1310–1316.
 160. Webb BJ, Healy R, Majers J, et al. Prediction of bloodstream infection due to vancomycin-resistant *Enterococcus* in patients undergoing leukemia induction or hematopoietic stem-cell transplantation. *Clin Infect Dis*. 2017;64:1753–1759.
 161. Becker K, Pagnier I, Schuhen B, et al. Does nasal colonization by methicillin-resistant coagulase-negative staphylococci and methicillin-susceptible *Staphylococcus aureus* strains occur frequently enough to represent a risk of false-positive methicillin-resistant *S. aureus* determinations by molecular methods? *J Clin Microbiol*. 2006;44:229–231.
 162. Schots R, Trulleman F, Van Riet I, et al. The clinical impact of early gram-positive bacteremia and the use of vancomycin after allogeneic bone marrow transplantation. *Transplantation*. 2000;69:1511–1514.
 163. Arns da Cunha C, Weisdorf D, Shu XO, et al. Early gram-positive bacteremia in BMT recipients: impact of three different approaches to antimicrobial prophylaxis. *Bone Marrow Transplant*. 1998;21:173–180.
 164. Mikulska M, Raiola AM, Galaverna F, et al. Pre-engraftment bloodstream infections after allogeneic hematopoietic stem cell transplantation: impact of T cell-replete transplantation from a haploidentical donor. *Biol Blood Marrow Transplant*. 2018;24:109–118.
 165. Girmenia C, Bertina A, Picocchi A, et al. Incidence, risk factors and outcome of pre-engraftment gram-negative bacteremia after allogeneic and autologous hematopoietic stem cell transplantation: an Italian Prospective Multicenter Survey. *Clin Infect Dis*. 2017;65:1884–1896.
 166. Kossow A, Kampmeier S, Willems S, et al. Control of multidrug-resistant *Pseudomonas aeruginosa* in allogeneic hematopoietic stem cell transplant recipients by a novel bundle including remodeling of sanitary and water supply systems. *Clin Infect Dis*. 2017;65:935–942.
 167. Girmenia C, Rossolini GM, Picocchi A, et al. Infections by carbapenem-resistant *Klebsiella pneumoniae* in SCT recipients: a nationwide retrospective survey from Italy. *Bone Marrow Transplant*. 2015;50:282–288.
 168. Peric Z, Vranjes VR, Durakovic N, et al. Gut colonization by multidrug-resistant gram-negative bacteria is an independent risk factor for development of intestinal acute graft-versus-host disease. *Biol Blood Marrow Transplant*. 2017;23:1221–1222.
 169. Hakki M, Limaye AP, Kim HW, et al. Invasive *Pseudomonas aeruginosa* infections: high rate of recurrence and mortality after hematopoietic cell transplantation. *Bone Marrow Transplant*. 2007;39:687–693.
 170. Bock AM, Cao Q, Ferrieri P, et al. Bacteremia in blood or marrow transplantation patients: clinical risk factors for infection and emerging antibiotic resistance. *Biol Blood Marrow Transplant*. 2013;19:102–108.
 171. van Veen KE, Brouwer MC, van der Ende A, et al. Bacterial meningitis in hematopoietic stem cell transplant recipients: a population-based prospective study. *Bone Marrow Transplant*. 2016;51:1490–1495.
 172. Rozmus J, Mallhi K, Ke J, et al. Functional hyposplenism after hematopoietic stem cell transplantation. *Bone Marrow Transplant*. 2015;50:1343–1347.
 173. D'Orsogna LJ, Wright MP, Krueger RG, et al. Allogeneic hematopoietic stem cell transplantation recipients have defects of both switched and IgM memory B cells. *Biol Blood Marrow Transplant*. 2009;15:795–803.
 174. Weller S, Braun MC, Tan BK, et al. Human blood IgM “memory” B cells are circulating splenic marginal zone B cells harboring a prediversified immunoglobulin repertoire. *Blood*. 2004;104:3647–3654.
 175. Engelhard D, Cordonnier C, Shaw PJ, et al. Early and late invasive pneumococcal infection following stem cell transplantation: a European Bone Marrow Transplantation survey. *Br J Haematol*. 2002;117:444–450.
 176. Schutze GE, Mason EO Jr, Wald ER, et al. Pneumococcal infections in children after transplantation. *Clin Infect Dis*. 2001;33:16–21.
 177. Haddad PA, Repka TL, Weisdorf DJ. Penicillin-resistant *Streptococcus pneumoniae* septic shock and meningitis complicating chronic graft versus host disease: a case report and review of the literature. *Am J Med*. 2002;113:152–155.
 178. Tauro S, Dobie D, Richardson G, et al. Recurrent penicillin-resistant pneumococcal sepsis after matched unrelated donor transplantation for refractory T cell lymphoma. *Bone Marrow Transplant*. 2000;26:1017–1019.
 179. Steiner M, Villablanca J, Kersey J, et al. Viridans streptococcal shock in bone marrow transplantation patients. *Am J Hematol*. 1993;42:354–358.
 180. Ringden O, Heimdahl A, Lonnqvist B, et al. Decreased incidence of viridans streptococcal septicemia in allogeneic bone marrow transplant recipients after the introduction of acyclovir. *Lancet*. 1984;314:744.
 181. Razonable RR, Litzow MR, Khaliq Y, et al. Bacteremia due to viridans group streptococci with diminished susceptibility to levofloxacin among neutropenic patients receiving levofloxacin prophylaxis. *Clin Infect Dis*. 2002;34:1469–1474.
 182. Almyroudis NG, Fuller A, Jakubowski A, et al. Pre- and post-engraftment bloodstream infection rates and associated mortality in allogeneic hematopoietic stem cell transplant recipients. *Transpl Infect Dis*. 2005;7:11–17.
 183. Busca A, Cavechia I, Locatelli F, et al. Blood stream infections after allogeneic stem cell transplantation: a single-center experience with the use of levofloxacin prophylaxis. *Transpl Infect Dis*. 2012;14:40–48.
 184. Blennow O, Ljungman P, Sparrelid E, et al. Incidence, risk factors, and outcome of bloodstream infections during the pre-engraftment phase in 521 allogeneic hematopoietic stem cell transplantations. *Transpl Infect Dis*. 2014;16:106–114.
 185. Kimura M, Araoka H, Yoshida A, et al. Breakthrough viridans streptococcal bacteremia in allogeneic hematopoietic stem cell transplant recipients receiving levofloxacin prophylaxis in a Japanese hospital. *BMC Infect Dis*. 2016;16:372.
 186. Graber CJ, de Almeida KN, Atkinson JC, et al. Dental health and viridans streptococcal bacteremia in allogeneic hematopoietic stem cell transplant recipients. *Bone Marrow Transplant*. 2001;27:537–542.
 187. Alonso CD, Treadway SB, Hanna DB, et al. Epidemiology and outcomes of *Clostridium difficile* infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis*. 2012;54:1053–1063.
 188. Kamboj M, Xiao K, Katsas A, et al. *Clostridium difficile* infection after allogeneic hematopoietic stem cell transplant: strain diversity and outcomes associated with NAP1/027. *Biol Blood Marrow Transplant*. 2014;20:1626–1633.
 189. Schuster MG, Cleveland AA, Dubberke ER, et al. Infections in hematopoietic cell transplant recipients: results from the Organ Transplant Infection Project, a multicenter, prospective, cohort study. *Open Forum Infect Dis*. 2017;4:ofx050.
 190. Dubberke ER, Reske KA, Olsen MA, et al. Risk for *Clostridium difficile* infection after allogeneic hematopoietic stem cell transplant remains elevated in the postengraftment period. *Transplant Direct*. 2017;3:e145.
 191. Aldrete SD, Kraft CS, Magee MJ, et al. Risk factors and epidemiology of *Clostridium difficile* infection in hematopoietic stem cell transplant recipients during the peritransplant period. *Transpl Infect Dis*. 2017;19.
 192. Mani S, Rybicki L, Jagadeesh D, et al. Risk factors for recurrent *Clostridium difficile* infection in allogeneic hematopoietic stem cell transplant recipients. *Bone Marrow Transplant*. 2016;51:713–717.
 193. Jain T, Crosswell C, Urdy-Cornejo V, et al. *Clostridium difficile* colonization in hematopoietic stem cell transplant recipients: a prospective study of the epidemiology and outcomes involving toxigenic and nontoxigenic strains. *Biol Blood Marrow Transplant*. 2016;22:157–163.
 194. Brumhant J, Wang ZX, Hu C, et al. *Clostridium difficile* colonization and disease in patients undergoing hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2014;20:1329–1334.

195. Neemann K, Eichele DD, Smith PW, et al. Fecal microbiota transplantation for fulminant *Clostridium difficile* infection in an allogeneic stem cell transplant patient. *Transpl Infect Dis*. 2012;14:E161–E165.
196. Gaviria JM, Garcia PJ, Garrido SM, et al. Nontuberculous mycobacterial infections in hematopoietic stem cell transplant recipients: characteristics of respiratory and catheter-related infections. *Biol Blood Marrow Transplant*. 2000;6:361–369.
197. Roy V, Weisdorf D. Mycobacterial infections following bone marrow transplantation: a 20 year retrospective review. *Bone Marrow Transplant*. 1997;19:467–470.
198. Beswick J, Shin E, Michels F, et al. Incidence and risk factors for nontuberculous mycobacterial infection after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2018;24:366–372.
199. Lee HJ, Lee DG, Choi SM, et al. The demanding attention of tuberculosis in allogeneic hematopoietic stem cell transplantation recipients: high incidence compared with general population. *PLoS ONE*. 2017;12:e0173250.
200. Maeda T, Kusumi E, Kami M, et al. Disseminated tuberculosis following reduced-intensity cord blood transplantation for adult patients with hematological diseases. *Bone Marrow Transplant*. 2005;35:91–97.
201. Lee J, Lee MH, Kim WS, et al. Tuberculosis in hematopoietic stem cell transplant recipients in Korea. *Int J Hematol*. 2004;79:185–188.
202. George B, Mathews V, Srivastava V, et al. Tuberculosis among allogeneic bone marrow transplant recipients in India. *Bone Marrow Transplant*. 2001;27:973–975.
203. Budak-Alpdogan T, Tangun Y, Kalayoglu-Besik S, et al. The frequency of tuberculosis in adult allogeneic stem cell transplant recipients in Turkey. *Biol Blood Marrow Transplant*. 2000;6:370–374.
204. Pai M, Zwerling A, Menzies D. Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection: an update. *Ann Intern Med*. 2008;149:177–184.
205. Menzies D, Pai M, Comstock G. Meta-analysis: new tests for the diagnosis of latent tuberculosis infection: areas of uncertainty and recommendations for research. *Ann Intern Med*. 2007;146:340–354.
206. Mazurek GH, Jereb J, Lobue P, et al. Guidelines for using the QuantiFERON-TB Gold test for detecting *Mycobacterium tuberculosis* infection, United States. *MMWR Recomm Rep*. 2005;54(RR-15):49–55.
207. Kobashi Y, Obase Y, Fukuda M, et al. Clinical reevaluation of the QuantiFERON TB-2G test as a diagnostic method for differentiating active tuberculosis from nontuberculous mycobacteriosis. *Clin Infect Dis*. 2006;43:1540–1546.
208. Lee SS, Liu YC, Huang TS, et al. Comparison of the interferon gamma release assay and the tuberculin skin test for contact investigation of tuberculosis in BCG-vaccinated health care workers. *Scand J Infect Dis*. 2008;40:373–380.
209. Kugler JW, Armitage JO, Helms CM, et al. Nosocomial legionnaires' disease: occurrence in recipients of bone marrow transplants. *Am J Med*. 1983;74:281–288.
210. Oren I, Zuckerman T, Avivi I, et al. Nosocomial outbreak of *Legionella pneumophila* serogroup 3 pneumonia in a new bone marrow transplant unit: evaluation, treatment and control. *Bone Marrow Transplant*. 2002;30:175–179.
211. Harrington RD, Woolfrey AE, Bowden R, et al. Legionellosis in a bone marrow transplant center. *Bone Marrow Transplant*. 1996;18:361–368.
212. Schwelke JR, Hackman R, Bowden R. Pneumonia due to *Legionella micdadei* in bone marrow transplant recipients. *Rev Infect Dis*. 1990;12:824–828.
213. van Burik JA, Hackman R, Nadeem S, et al. Nocardiosis after bone marrow transplantation: a retrospective study. *Clin Infect Dis*. 1997;24:1154–1160.
214. Shannon K, Pasikhova Y, Ibekweh Q, et al. Nocardiosis following hematopoietic stem cell transplantation. *Transpl Infect Dis*. 2016;18:169–175.
215. Mercante JW, Winchell JM. Current and emerging *Legionella* diagnostics for laboratory and outbreak investigations. *Clin Microbiol Rev*. 2015;28:95–133.
216. Sivagnanam S, Pergam SA. Legionellosis in Transplantation. *Curr Infect Dis Rep*. 2016;18:9.
217. Martino R, Lopez R, Pericas R, et al. Listeriosis in bone marrow transplant recipient. *Clin Infect Dis*. 1996;23:419–420.
218. Chang J, Powles R, Mehta J, et al. Listeriosis in the bone marrow transplant recipient. *Clin Infect Dis*. 1995;21:1289–1290.
219. Long SG, Leyland MJ, Milligan DW. Listeria meningitis after bone marrow transplantation. *Bone Marrow Transplant*. 1993;12:537–539.
220. Ljungman P, Larsson K, Kumlien G, et al. Leukocyte depleted, unscreened blood products give a low risk for CMV infection and disease in CMV seronegative allogeneic stem cell transplant recipients with seronegative stem cell donors. *Scand J Infect Dis*. 2002;34:347–350.
221. Bowden RA, Slichter SJ, Sayers MH, et al. Use of leukocyte-depleted platelets and cytomegalovirus-seronegative red blood cells for prevention of primary cytomegalovirus infection after marrow transplant. *Blood*. 1991;78:246–250.
222. Saral R, Burns WH, Laskin OL, et al. Acyclovir prophylaxis of herpes simplex virus infections: a randomized, double-blind, controlled trial in bone marrow transplant recipients. *N Engl J Med*. 1981;305:63–67.
223. Dignani MC, Mykietiak A, Michelet M, et al. Valacyclovir prophylaxis for the prevention of herpes simplex virus reactivation in recipients of progenitor cells transplantation. *Bone Marrow Transplant*. 2002;29:263–267.
224. Liesveld JL, Abboud CN, Ifthikharuddin JJ, et al. Oral valacyclovir versus intravenous acyclovir in preventing herpes simplex virus infections in autologous stem cell transplant recipients. *Biol Blood Marrow Transplant*. 2002;8:662–665.
225. Teira P, Battiwalla M, Ramanathan M, et al. Early cytomegalovirus reactivation remains associated with increased transplant related mortality in the current era: a CIBMTR analysis. *Blood*. 2016;127:2427–2438.
226. Nichols WG, Price TH, Gooley T, et al. Transfusion-transmitted cytomegalovirus infection after receipt of leukoreduced blood products. *Blood*. 2003;101:4195–4200.
227. Bowden RA, Slichter SJ, Sayers M, et al. A comparison of filtered leukocyte-reduced and cytomegalovirus seronegative blood products for the prevention of transfusion-associated CMV infection after marrow transplant. *Blood*. 1995;86:3598–3603.
228. Pergam SA, Xie H, Sandhu R, et al. Efficiency and risk factors for CMV transmission in seronegative hematopoietic stem cell recipients. *Biol Blood Marrow Transplant*. 2012;18:1391–1400.
229. Goodrich JM, Mori M, Gleaves CA, et al. Early treatment with ganciclovir to prevent cytomegalovirus disease after allogeneic bone marrow transplantation. *N Engl J Med*. 1991;325:1601–1607.
230. Reusser P, Einsele H, Lee J, et al. Randomized multicenter trial of foscarnet versus ganciclovir for preemptive therapy of cytomegalovirus infection after allogeneic stem cell transplantation. *Blood*. 2002;99:1159–1164.
231. Goodrich JM, Bowden RA, Fisher L, et al. Ganciclovir prophylaxis to prevent cytomegalovirus disease after allogeneic marrow transplant. *Ann Intern Med*. 1993;118:173–178.
232. Winston DJ, Ho WG, Bartoni K, et al. Ganciclovir prophylaxis of cytomegalovirus infection and disease in allogeneic bone marrow transplant recipients: results of a placebo-controlled, double-blind trial. *Ann Intern Med*. 1993;118:179–184.
233. Prentice HG, Gluckman E, Powles RL, et al. Impact of long-term acyclovir on cytomegalovirus infection and survival after allogeneic bone marrow transplantation. European Acyclovir for CMV Prophylaxis Study Group. *Lancet*. 1994;343:749–753.
234. Burns LJ, Miller W, Kandaswamy C, et al. Randomized clinical trial of ganciclovir vs acyclovir for prevention of cytomegalovirus antigenemia after allogeneic transplantation. *Bone Marrow Transplant*. 2002;30:945–951.
235. Milano F, Pergam SA, Xie H, et al. Intensive strategy to prevent CMV disease in seropositive umbilical cord blood transplant recipients. *Blood*. 2011;118:5689–5696.
236. Winston DJ, Yeager AM, Chandrasekar PH, et al. Randomized comparison of oral valacyclovir and intravenous ganciclovir for prevention of cytomegalovirus disease after allogeneic bone marrow transplantation. *Clin Infect Dis*. 2003;36:749–758.
237. Melendez DP, Razonable RR. Letermovir and inhibitors of the terminase complex: a promising new class of investigational antiviral drugs against human cytomegalovirus. *Infect Drug Resist*. 2015;8:269–277.
238. Marty FM, Ljungman P, Chemaly RF, et al. Letermovir prophylaxis for cytomegalovirus in hematopoietic-cell transplantation. *N Engl J Med*. 2017;377:2433–2444.
239. Ljungman P, Boeckh M, Hirsch HH, et al. Definitions of cytomegalovirus infection and disease in transplant patients for use in clinical trials. *Clin Infect Dis*. 2017;64:87–91.
240. Boeckh M, Stevens-Ayers T, Travi G, et al. Cytomegalovirus (CMV) DNA quantitation in bronchoalveolar lavage fluid from hematopoietic stem cell transplant recipients with CMV pneumonia. *J Infect Dis*. 2017;215:1514–1522.
241. Reed EC, Bowden RA, Dandiker PS, et al. Treatment of cytomegalovirus pneumonia with ganciclovir and intravenous cytomegalovirus immunoglobulin in patients with bone marrow transplants. *Ann Intern Med*. 1988;109:783–788.
242. Limaye AP, Bowden RA, Myerson D, et al. Cytomegalovirus disease occurring before engraftment in marrow transplant recipients. *Clin Infect Dis*. 1997;24:830–835.
243. Limaye AP, Huang ML, Leisenring W, et al. Cytomegalovirus DNA load in plasma for the diagnosis of CMV disease before engraftment in hematopoietic stem-cell transplant recipients. *J Infect Dis*. 2001;183:377–382.
244. van Burik JA, Lawatsch EJ, DeFor TE, et al. Cytomegalovirus enteritis among hematopoietic stem cell transplant recipients. *Biol Blood Marrow Transplant*. 2001;7:674–679.
245. Green ML, Leisenring W, Stachel D, et al. Efficacy of a viral load-based, risk-adapted, preemptive treatment strategy for prevention of cytomegalovirus disease after hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2012;18:1687–1699.
246. Crippa F, Corey L, Chuang EL, et al. Virological, clinical, and ophthalmologic features of cytomegalovirus retinitis after hematopoietic stem cell transplantation. *Clin Infect Dis*. 2001;32:214–219.
247. Reed EC, Shepp DH, Dandiker PS, et al. Ganciclovir treatment of cytomegalovirus infection of the gastrointestinal tract after marrow transplantation. *Bone Marrow Transplant*. 1988;3:199–206.
248. Walter EA, Greenberg PD, Gilbert MJ, et al. Reconstitution of cellular immunity against cytomegalovirus in recipients of allogeneic bone marrow by transfer of T-cell clones from the donor. *N Engl J Med*. 1995;333:1038–1044.
249. Erice A, Borrell N, Li W, et al. Ganciclovir susceptibilities and analysis of UL97 region in cytomegalovirus isolates from bone marrow recipients with CMV disease after antiviral prophylaxis. *J Infect Dis*. 1998;178:531–534.
250. Herling M, Schröder L, Awerkwies S, et al. Persistent CMV infection after allogeneic hematopoietic stem cell transplantation in a CMV-seronegative donor-to-positive recipient constellation: development of multidrug resistance in the absence of anti-viral cellular immunity. *J Clin Virol*. 2016;74:57–60.
251. Blyth E, Clancy L, Simms R, et al. Donor-derived CMV-specific T cells reduce the requirement for CMV-directed pharmacotherapy after allogeneic stem cell transplantation. *Blood*. 2013;121:3745–3758.
252. Boeckh M. New insights into cytomegalovirus infection after allogeneic hematopoietic stem cell transplant. *Clin Adv Hematol Oncol*. 2017;15:515–517.
253. Tzannou I, Papadopoulos A, Naik S, et al. Off-the-shelf virus-specific T cells to treat BK virus, human herpesvirus 6, cytomegalovirus, Epstein-Barr virus, and adenovirus infections after allogeneic hematopoietic stem-cell transplantation. *J Clin Oncol*. 2017;35:3547–3557.
254. Marty FM, Winston DJ, Rowley SD, et al. CMX001 to prevent cytomegalovirus disease in hematopoietic-cell transplantation. *N Engl J Med*. 2013;369:1227–1236.
255. Arvin AM. Varicella-zoster virus: pathogenesis, immunity, and clinical management in hematopoietic cell transplant recipients. *Biol Blood Marrow Transplant*. 2000;6:219–230.
256. Steer CB, Szer J, Sasadeusz J, et al. Varicella-zoster infection after allogeneic bone marrow transplantation: incidence, risk factors and prevention with low-dose acyclovir and ganciclovir. *Bone Marrow Transplant*. 2000;25:657–664.
257. Blennow O, Fjaertoft G, Winiarski J, et al. Varicella-zoster reactivation after allogeneic stem cell transplantation without routine prophylaxis—the incidence remains high. *Biol Blood Marrow Transplant*. 2014;20:1646–1649.
258. Erard V, Guthrie KA, Varley C, et al. One-year acyclovir prophylaxis for preventing varicella-zoster virus disease after hematopoietic cell transplantation: no evidence of rebound varicella-zoster virus disease after drug discontinuation. *Blood*. 2007;110:3071–3077.
259. David DS, Tegtmeyer BR, O'Donnell MR, et al. Visceral varicella-zoster after bone marrow transplantation: report of a case series and review of the literature. *Am J Gastroenterol*. 1998;93:810–813.
260. Schiller GJ, Nimer SD, Gajewski JL, et al. Abdominal presentation of varicella zoster infection in recipients of allogeneic bone marrow transplantation. *Bone Marrow Transplant*. 1991;7:489–491.
261. Tenenbaum T, Kramm CM, Laws HJ, et al. Pre-eruptive varicella zoster virus encephalitis in two children after haematopoietic stem cell transplantation. *Med Pediatr Oncol*. 2002;38:288–289.
262. Verdonck LF, Cornelissen JJ, Dekker AW, et al. Acute abdominal pain as a presenting symptom of varicella zoster virus infection in recipients of bone marrow transplants. *Clin Infect Dis*. 1993;16:190–191.