

- doses in young women: a randomized clinical trial. *JAMA*. 2013;309:1793–1802.
395. Laprise JF, Markowitz LE, Chesson HW, et al. Comparison of 2-dose and 3-dose 9-valent human papillomavirus vaccine schedules in the United States: a cost-effectiveness analysis. *J Infect Dis*. 2016;214:685–688.
  396. Iversen OE, Miranda MJ, Ulied A, et al. Immunogenicity of the 9-valent HPV vaccine using 2-dose regimens in girls and boys vs a 3-dose regimen in women. *JAMA*. 2016;316:2411–2421.
  397. Ogilvie G, Sauvageau C, Dionne M, et al. Immunogenicity of 2 vs 3 doses of the quadrivalent human papillomavirus vaccine in girls aged 9 to 13 years after 60 months. *JAMA*. 2017;317:1687–1688.
  398. Markowitz LE, Liu G, Hariri S, et al. Prevalence of HPV after introduction of the vaccination program in the United States. *Pediatrics*. 2016;137:e20151968.
  399. Tabrizi SN, Brotherton JM, Kaldor JM, et al. Assessment of herd immunity and cross-protection after a human papillomavirus vaccination programme in Australia: a repeat cross-sectional study. *Lancet Infect Dis*. 2014;14:958–966.
  400. Herrero R, Quint W, Hildesheim A, et al. Reduced prevalence of oral human papillomavirus (HPV) 4 years after bivalent HPV vaccination in a randomized clinical trial in Costa Rica. *PLoS ONE*. 2013;8:e68329.
  401. Drolet M, Benard E, Boily MC, et al. Population-level impact and herd effects following human papillomavirus vaccination programmes: a systematic review and meta-analysis. *Lancet Infect Dis*. 2015;15:565–580.
  402. Brotherton JM, Fridman M, May CL, et al. Early effect of the HPV vaccination programme on cervical abnormalities in Victoria, Australia: an ecological study. *Lancet*. 2011;377:2085–2092.
  403. Powell SE, Hariri S, Steinau M, et al. Impact of human papillomavirus (HPV) vaccination on HPV 16/18-related prevalence in precancerous cervical lesions. *Vaccine*. 2012;31:109–113.
  404. Benard VB, Castle PE, Jenison SA, et al. Population-based incidence rates of cervical intraepithelial neoplasia in the human papillomavirus vaccine era. *JAMA Oncol*. 2017;3:833–837.
  405. Lewis RM, Markowitz LE, Gargano JW, et al. Prevalence of genital human papillomavirus among sexually experienced males and females aged 14–59 years, United States, 2013–2014. *J Infect Dis*. 2018;217:869–877.
  406. Joura EA, Giuliano AR, Iversen OE, et al. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. *N Engl J Med*. 2015;372:711–722.
  407. Macartney KK, Chiu C, Georgousakis M, et al. Safety of human papillomavirus vaccines: a review. *Drug Saf*. 2013;36:393–412.
  408. Phillips A, Patel C, Pillsbury A, et al. Safety of human papillomavirus vaccines: an updated review. *Drug Saf*. 2018;41:329–346.
  409. Moreira ED Jr, Block SL, Ferris D, et al. Safety profile of the 9-valent HPV vaccine: a combined analysis of 7 phase III clinical trials. *Pediatrics*. 2016;138.
  410. Grimaldi-Bensouda L, Rossignol M, Kone-Paut I, et al. Risk of autoimmune diseases and human papilloma virus (HPV) vaccines: six years of case-referent surveillance. *J Autoimmun*. 2017;79:84–90.
  411. Miranda S, Chaignot C, Collin C, et al. Human papillomavirus vaccination and risk of autoimmune diseases: a large cohort study of over 2 million young girls in France. *Vaccine*. 2017;35:4761–4768.
  412. Gee J, Sukumaran L, Weintraub E, et al. Risk of Guillain-Barre syndrome following quadrivalent human papillomavirus vaccine in the vaccine safety datalink. *Vaccine*. 2017;35:5756–5758.
  413. Arana JE, Harrington T, Cano M, et al. Post-licensure safety monitoring of quadrivalent human papillomavirus vaccine in the vaccine adverse event reporting system (VAERS), 2009–2015. *Vaccine*. 2018;36:1781–1788.
  414. Petrosky E, Bocchini JA Jr, Hariri S, et al. Use of 9-valent human papillomavirus (HPV) vaccine: updated HPV vaccination recommendations of the advisory committee on immunization practices. *MMWR Morb Mortal Wkly Rep*. 2015;64:300–304.
  415. Meites E, Kempe A, Markowitz LE. Use of a 2-dose schedule for human papillomavirus vaccination - updated recommendations of the advisory committee on immunization practices. *MMWR Morb Mortal Wkly Rep*. 2016;65:1405–1408.
  416. Markowitz LE, Dunne EF, Saraiya M, et al. Human papillomavirus vaccination: recommendations of the advisory committee on immunization practices (ACIP). *MMWR Recomm Rep*. 2014;63(RR-05):1–30.
  417. Bonde U, Joergensen JS, Lamont RF, et al. Is HPV vaccination in pregnancy safe? *Hum Vaccin Immunother*. 2016;12:1960–1964.
  418. Swedish KA, Factor SH, Goldstone SE. Prevention of recurrent high-grade anal neoplasia with quadrivalent human papillomavirus vaccination of men who have sex with men: a nonconcurrent cohort study. *Clin Infect Dis*. 2012;54:891–898.
  419. Joura EA, Garland SM, Paavonen J, et al. Effect of the human papillomavirus (HPV) quadrivalent vaccine in a subgroup of women with cervical and vulvar disease: retrospective pooled analysis of trial data. *BMJ*. 2012;344:e1401.

# JC, BK, and Other Polyomaviruses: Progressive Multifocal Leukoencephalopathy (PML)

C. Sabrina Tan and Igor J. Koralnik

## SHORT VIEW SUMMARY

### JC VIRUS

#### Definition

- JC virus (JCV) is a ubiquitous human polyomavirus that causes several central nervous system diseases in immunocompromised patients, including progressive multifocal leukoencephalopathy (PML), JCV granule cell neuronopathy, JCV encephalopathy, and JCV meningitis.

#### Epidemiology

- JCV infects 40% to 86% of the general population worldwide.
- JCV can be detected in the urine of one-third of healthy and immunosuppressed individuals.
- PML can occur in up to 5% of untreated patients with acquired immunodeficiency syndrome.
- Up to 82% of PML patients are infected with human immunodeficiency virus.

#### Microbiology

- JCV is a member of the Polyomaviridae family, *Orthopolyomavirus* genus.
- It is a double-stranded DNA virus without an envelope.
- After primary infection, JCV remains latent in the kidney tubular epithelial cells.
- Reactivation of JCV causes a lytic infection of oligodendrocytes in the brain, leading to PML.

#### Diagnosis

- Definitive Diagnosis of PML
  - JCV DNA is detected in cerebrospinal fluid by polymerase chain reaction (PCR) assay,

or JCV proteins are detected in brain tissues through histopathologic assessment.

- Possible Diagnosis of PML
  - Magnetic resonance imaging findings and clinical presentation are consistent with PML in the absence of detection of JCV in cerebrospinal fluid.

#### Therapy

- There are no effective antiviral medications.
- Immune reconstitution can boost host cellular immune response to better control JC viral replication.

#### Prevention

- Measures should be taken to prevent immunosuppression.

### BK VIRUS

#### Definition

- BK virus (BKV) is a ubiquitous human polyomavirus that causes hemorrhagic cystitis in hematopoietic stem cell transplantation patients and nephropathy in kidney transplantation recipients.

#### Epidemiology

- BKV infects 82% to 90% of the general population worldwide.
- BKV can be detected in the urine of up to 20% of asymptomatic healthy individuals.
- BKV nephropathy occurs in up to 10% of kidney transplant patients.

- BKV-associated disease can occur in up to 15.9% of allogeneic stem cell transplant recipients.

#### Microbiology

- BKV is a member of the Polyomaviridae family, *Orthopolyomavirus* genus.
- It is a double-stranded DNA virus without an envelope.
- After primary infection, BKV remains latent in the kidney tubular epithelial cells.
- Reactivation of BKV causes hemorrhagic cystitis and nephropathy.

#### Diagnosis

- BKV DNA is detected through PCR assay in blood and sustained viremia in urine.
- Cytopathologic changes may be detected in a kidney biopsy specimen.

#### Therapy

- There is no effective antiviral medication.
- Reduction in immunosuppression can boost host cellular immune response to better control BKV replication.

#### Prevention

- Early detection of viral reactivation in either urine or blood with PCR assay is an indicator for a preemptive reduction in immunosuppression to help reduce occurrence of renal disease.

The human polyomaviruses, JC virus (JCV) and BK virus (BKV), are ubiquitous in most human populations throughout the world and do not cause disease in immunocompetent individuals. As much as 90% and 86% of the general adult population are seropositive for BKV and JCV, respectively.<sup>1,2</sup> In individuals with immunosuppression, JCV is the etiologic agent of a demyelinating disease of the central nervous system (CNS)—progressive multifocal leukoencephalopathy (PML)<sup>3,4</sup>—whereas BKV causes nephropathy, hemorrhagic cystitis, and ureteral stenosis. PML is an acquired immunodeficiency syndrome (AIDS)-defining opportunistic infection.<sup>5</sup> BKV nephropathy is a cause of allograft loss in kidney transplant recipients. The polyomavirus family has expanded with the discoveries of new viruses and their associated diseases.

## VIROLOGY

### History

Polyomaviruses are small (45-nm) nonenveloped viruses that are composed of 72 capsomeres with icosahedral symmetry containing

a circular double-stranded DNA. The Polyomaviridae family is composed of three genera: two mammalian genera, *Orthopolyomavirus* and *Wukupolyomavirus*; and one avian genus, *Avipolyomavirus*.<sup>6</sup> The polyomaviruses are ubiquitous in nature and are species specific, including humans (JCV, BKV), monkeys (simian virus 40 [SV40]), and mice (murine polyomavirus). In humans, JCV was first isolated from the brain of a patient with PML whose initials were J.C.<sup>7</sup> Similarly, B.K. were the initials of a kidney transplant patient in whom BKV-associated ureteral stenosis was first described.<sup>8</sup> Since 2007, the Polyomaviridae family has expanded to include newly discovered viruses. These were named after the site of discovery—WUPyV (Washington University) and KIPyV (Karolinska Institute)<sup>9,10</sup>; their geographic origins—MWPyV (Malawi)<sup>11</sup> and STLPyV (St. Louis)<sup>12</sup>; the diseases that they cause—MCV (Merkel cell carcinoma)<sup>13</sup> and TSPyV (trichodysplasia spinulosa)<sup>14</sup>; or their order of discovery—HPyV6, HPyV7, HPyV9, HPyV12, and HPyV 13 (human polyomaviruses 6, 7, 9, 12, 13).<sup>15–17</sup>


## Epidemiology

Both JCV and BKV have worldwide distribution, including geographically isolated populations with little exposure to other infections.<sup>18,19</sup>

Although seroprevalence for both BKV and JCV increases rapidly with age,<sup>20</sup> primary infection of each virus occurs independently.<sup>21</sup> Acquisition of both viruses through vertical transmission has been documented.<sup>22</sup> After primary infection, BKV and JCV can be detected in renal tubular epithelial cells, where they may remain. Although JC viruria occurs independent of the host's immune status, JC viruria is usually detected only in individuals with immunosuppression. JCV is detected in the urine of 20% to 30% of healthy and immunosuppressed individuals alike, with or without PML.<sup>23,24</sup> However, JCV viremia has been detected in 20% to 40% of individuals who are human immunodeficiency virus (HIV) positive without PML and in 60% to 80% of patients with PML.<sup>25,26</sup> Conversely, BK viruria occurs in 0% to 20% of asymptomatic immunocompetent individuals,<sup>1</sup> but viral shedding in the urine is higher in individuals with immunosuppression (10%–60%),<sup>27–29</sup> correlating to the degrees of immunosuppression.<sup>30</sup> Specifically, in HIV-positive patients, BK viral load in the urine increases with decreased CD4<sup>+</sup> T-cell counts.<sup>31,32</sup> Although BKV is usually not detected in the peripheral blood of patients with either immunocompetence or immunosuppression, the detection of BKV DNA in the plasma of renal transplant patients is indicative of an increased risk of development of BKV-induced nephropathy.<sup>27,33</sup> In a cohort analysis of 491 patients undergoing hematopoietic stem cell transplant (HSCT), investigators reported that BKV-associated disease occurred in 15.9% and was associated with significant and prolonged morbidity.<sup>34</sup> Patients who underwent reduced-intensity, double umbilical cord blood cell transplantation had BK viremia frequently detected (15/27) within 100 days of transplantation.<sup>35</sup>

## Genome

Polymaviruses have a circular double-stranded genomic DNA of approximately 5 kb. The genomes of BKV and JCV have been used extensively in human population migration studies because of their extremely conserved coding regions. JCV and BKV share approximately 70% genome homology,<sup>36</sup> and all polymaviruses maintain the feature of coding DNA from both strands. The early proteins, large T and small t, are transcribed counterclockwise from one strand, whereas the late proteins, VP1, VP2, VP3, and agnoprotein, are transcribed clockwise from the opposite strand (Fig. 144.1). The genome can be divided into three distinct areas: (1) the early genes region, which includes the large T and small t antigens, regulatory proteins responsible for viral transformation, replication, and regulation of gene expression; (2) the late genes region, which encodes the capsid proteins VP1, VP2, and VP3



**FIG. 144.1 Polymavirus genomic map (~5 kb).** RR, Regulatory region. (Courtesy Dr. Yiping Chen.)

and the agnoprotein; and (3) the noncoding regulatory region. The agnoprotein is important both in regulation of JCV transcription and translation and in dysregulation of host cell cycle and DNA repairs.<sup>37,38</sup> Similar to the origin of DNA replication, the noncoding regulatory region also contains several binding sites for nuclear factors, which play important roles in JCV transcription. Deletions, insertions, and rearrangements in the regulatory region are associated with tissue tropism and virulence. The archetype regulatory region is mainly found in JCV isolated from kidney cells and consequently from urine. It is thought to be the type from which all other JCV regulatory region sequences have evolved. The rearranged type of regulatory region of JCV is principally detected in the brain and cerebrospinal fluid (CSF) of patients with PML.<sup>39</sup> Whereas the rearranged type of regulatory region is found in the peripheral blood of patients with PML, a mixture of both types is present in the bone marrow.<sup>25,40</sup> Which type of regulatory region is associated with primary infection is not known.

## Receptors and Cell Entry

### JC Virus

Data suggest that N-linked glycoprotein containing an  $\alpha(2,6)$ -linked sialic acid may act as a JCV receptor.<sup>41</sup> In addition, evidence from cultured cell lines shows that JCV uses the serotonergic 5-HT2A receptors for cell entry.<sup>42</sup> Pharmacologic blockade of the serotonergic receptors restricts JCV infection *in vitro*. The 5-HT2A receptor is present on glial cells and astrocytes and on B lymphocytes, platelets, and kidney epithelial cells. JCV also uses clathrin-coated pits on cell surfaces after binding and adsorption, mediated by the capsid protein VP1. Cell entry occurs via endocytosis and results in fusion of virus-carrying vesicle with the nuclear membrane.<sup>43</sup>

### BK Virus

BKV uses an N-linked glycoprotein containing an  $\alpha(2,3)$ -linked sialic acid as a receptor. Entry into the cell occurs via caveolae-mediated endocytosis.<sup>44</sup>

## New Human Polymaviruses

Seroprevalences for the newer human polymaviruses range from 25% to 90%.<sup>15,45–48</sup> The KIPyV and WUPyV polymaviruses were first isolated from respiratory samples with high-throughput screens.<sup>10,49</sup> Subsequently, molecular evidence indicated a worldwide distribution for both viruses in the respiratory secretions of children, with incidence rates as high as 2.5% in Australia<sup>50</sup> and 7% in South Korea.<sup>51</sup> For WUPyV, A study analyzing nasopharyngeal aspirates in the United Kingdom showed a bimodal age distribution of KIPyV and detection of the WUPyV in samples from patients younger than 15 years old.<sup>52</sup> Although both WUPyV and KIPyV could be colonizers of the respiratory tracts, KIPyV was associated with respiratory symptoms in transplant recipients.<sup>53</sup> HPyV6, HPyV7, MWPyV, and STLPyV were also all recovered from nonsterile sites such as skin and stool. Therefore the proof of their existence as bona fide infectious agents remains to be established. Like KIPyV, these viruses could be part of the human microbiota that causes problems only in immunosuppressed patients. For example, HPyV7 was found to be the cause of a pruritic rash in a lung transplant recipient.<sup>54</sup> Lastly, the detection of HPyV9 in normally sterile blood distinguished this virus as a possible human pathogen.

The Merkel cell polyomavirus (MCV), discovered in 2008, and trichodysplasia spinulosa polyomavirus (TSPyV), discovered in 2010, are both associated with human skin diseases. MCV can be detected in up to 80% of tissues with Merkel cell carcinoma, an aggressive and deadly neuroectodermal tumor that principally affects patients with immunosuppression.<sup>13</sup> Molecular studies indicate that MCV shares a similar genome with the other human polymaviruses and that integration of MCV into the tumor genome most likely takes place before clonal expansion of tumor cells. MCV is most similar to the African green monkey lymphotropic polyomavirus. TSPyV viral particles were detected with electron microscopy on a hair follicle from a heart transplant patient with trichodysplasia spinulosa, a rare skin disease affecting immunocompromised patients.<sup>14</sup>

Interesting to note, unlike JCV and BKV, these newer human polymaviruses do not contain a gene that encodes the agnoprotein.

The agnoprotein was initially named because its function was unknown. However, this protein appears to have important regulatory functions of both polyomaviruses and host cells. The significance of the agnoprotein deletion in these polyomaviruses remains to be determined.

## PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY AND OTHER JC VIRUS-ASSOCIATED SYNDROMES

### Epidemiology

PML was initially described in 1958.<sup>55</sup> The disease was estimated to occur in 0.07% of individuals with hematologic malignant diseases,<sup>56</sup> such as leukemia and lymphoma, or rarely in patients with solid-organ cancer or organ transplants. Since the 1980s, in the era of HIV infection, PML has been recognized as a major opportunistic infection that affects up to 5% of patients with AIDS.<sup>5</sup> Despite the advent of combined antiretroviral therapy (ART), PML still occurs in 0.6 to 1.3 per 1000 HIV-positive person-years.<sup>4</sup> Currently, 82% of patients with PML have AIDS, 8.4% have hematologic malignant diseases, 2.8% are solid-organ transplant recipients, 0.95% have chronic inflammatory diseases, and 6% have no defined risk factor for PML.<sup>4</sup> Indeed, 38 cases of PML have been reported in HIV-negative individuals with minimal or occult immunosuppression.<sup>57</sup> In HIV-negative individuals, the incidence rate of PML per 100,000 person-years is 0.3 in the general population, 1.0 in patients with rheumatoid arthritis, 2.4 in those with systemic lupus erythematosus, 10.8 in those with autoimmune vasculitis, 8.3 in those with non-Hodgkin lymphoma and chronic lymphocytic leukemia, and 35.4 in bone marrow transplant recipients.<sup>58</sup> Lastly, the newest group of patients with PML has emerged among patients treated with immunomodulatory medications for malignant diseases or autoimmune diseases, such as patients treated with natalizumab and dimethyl fumarate for multiple sclerosis (MS) or Crohn disease,<sup>59–63</sup> rituximab for lymphoma<sup>64,65</sup> or lupus,<sup>66</sup> and efalizumab or fumaric acid for psoriasis.<sup>67,68</sup>

Natalizumab-treated MS patients are at increased risk of developing PML. As of November 30, 2017, 756 cases of natalizumab-related PML had been documented worldwide. Risk of PML increases in patients with longer natalizumab treatment duration, prior use of other immunosuppressants, and seropositivity for JCV. The incidence of natalizumab-associated PML is highest after 49 to 72 months of treatment in JCV-seropositive patients with a prior history of immunosuppressive treatment, where it reached 1.3%.<sup>69</sup>

### Pathogenesis

Primary infection with JCV is asymptomatic, after which the virus becomes clinically latent. JCV DNA is detected with polymerase chain reaction (PCR) assay in the tonsillar tissues of 39% of healthy individuals, which suggests a possible oral or respiratory route of infection.<sup>70,71</sup> Detection of JCV in urine of healthy adults indicates that the kidney

is a site of latency. Immunohistochemical staining shows viral proteins in the kidney tubular epithelial cells. Detection of JCV DNA in the bone marrow<sup>72</sup> and brain<sup>73</sup> of patients without PML suggests that these are also possible additional sites of latency.

In patients with PML, JCV infects oligodendrocytes and astrocytes in the brain and rarely the spinal cord, which causes a lytic infection that results in the destruction of the myelin sheath. Pathologic examinations of PML lesions show extensive demyelination of the affected areas. Because of the presence of oligodendrocytes and astrocytes in the hemispheric cortex, demyelination can also occur within the cortical gray matter.

### Clinical Manifestations Classic Progressive Multifocal Leukoencephalopathy

Because PML causes multifocal demyelination of the white matter of the CNS, the resulting neurologic deficits correspond to the location of the lesions. Although initial symptoms of PML can vary greatly, depending on the affected anatomic brain region, the predominant symptoms include coordination difficulties, gait imbalance, cognitive dysfunction, visual problems, and limb paresis. The optic nerves and the spinal cord of the CNS are usually spared, but incidental postmortem findings of PML lesions have been discovered in the spinal cord of a patient who was HIV positive with hemispheric PML.<sup>74</sup> Furthermore, although PML lesions are generally located in the white matter, seizures, which are usually considered to be of cortical origin, can occur in up to 44% of PML survivors and are associated with juxtacortical demyelinated lesions and with hyperintense cortical signal (HCS) demonstrated on T1-weighted magnetic resonance images. The histologic correlate of HCS is leukocortical encephalitis, which recapitulates the triad of PML (demyelination, astrogliosis, and infiltration with phagocytic macrophages, albeit in the cortical gray matter) (Table 144.1).<sup>75–77</sup>

### Progressive Multifocal Leukoencephalopathy–Immune Reconstitution Inflammatory Syndrome

Inflammatory PML has been frequently detected when the immune response is rapidly restored with ART in patients with HIV. Patients usually have an increasing CD4<sup>+</sup> cell count and a decreasing plasma HIV viral load. Unlike classic PML, up to 57% of the PML–immune reconstitution inflammatory syndrome (IRIS) lesions may display contrast enhancement with magnetic resonance imaging (MRI),<sup>4,78</sup> which indicates a breakdown of the blood-brain barrier, along with worsening of the initial presenting symptoms. Conditions may stabilize after the initial worsening of symptoms, but fatal outcome has been reported.<sup>79,80</sup> Magnetic resonance spectroscopy (MRS) helped define a unique metabolic pattern in PML lesions that showed an increased ratio of lipids to creatine consistent with inflammation. Combined with the presence of contrast enhancement in PML lesions, this finding yielded a 79%

**TABLE 144.1 JC Virus–Associated Diseases**

| CLINICAL PRESENTATION      | CLASSIC PML  | PML-IRIS  | JCV GCN   | JCV E  | JCV MENINGITIS   |
|----------------------------|--|---|---|--|--|
| Onset                      | Subacute   | Immune recovery   | Chronic   | Subacute   | Chronic  |
| Magnetic resonance imaging | Asymmetrical, well-demarcated, nonenhancing subcortical white matter lesions; hyperintense on T2-weighted and FLAIR images; hypointense on T1-weighted images                          | Contrast-enhancing lesions and possible mass effect                           | Cerebellar atrophy  | Cortical lesions   | Enlarged ventricles  |
| Neurologic symptoms        | Based on location  | Based on location and inflammation  | Cerebellar syndrome   | Encephalopathy   | Headache   |
| Histology                  | Demyelinating lesions often at gray-white matter junction; JCV detected in enlarged oligodendrocytes; bizarre astrocytes; presence of CD8 <sup>+</sup> T cells near JCV-infected cells | Demyelination similar to classic PML but with marked inflammatory infiltrates | Focal areas of cell loss in granule cell layer of cerebellum; JCV detected in enlarged granule cell neurons | Focal areas of cell loss in hemispheric cortex; JCV detected in enlarged pyramidal neurons | Productive infection of meningeal and choroid plexus cells |

E, Encephalopathy; FLAIR, fluid-attenuated inversion recovery; GCN, granule cell neuropathy; IRIS, immune reconstitution inflammatory syndrome; JCV, JC virus; PML, progressive multifocal leukoencephalopathy.



probability of IRIS compared with 13% in the absence of these criteria.<sup>81</sup>

In autopsy samples of HIV-positive patients with PML-IRIS, mainly CD8<sup>+</sup> lymphocytes were detected in the lesions, along with a paucity of CD4<sup>+</sup> lymphocytes.<sup>80,82,83</sup> The reported time at onset of PML-IRIS ranged from 4 to 108 weeks after start of ART.<sup>79,80</sup> In some patients, ART was started as treatment for PML<sup>79,80,82–86</sup>; in others, PML was diagnosed at the time of IRIS presentation.<sup>79,87</sup>

PML-IRIS has also been reported in HIV-negative patients, such as in natalizumab-treated MS patients after plasma exchange to remove natalizumab.<sup>88</sup>

### JC Virus Granule Cell Neuronopathy

Demyelination of white matter in the cerebellum is well described in patients with PML. In addition to oligodendrocytes and astrocytes, JCV can also infect and destroy cerebellar granule cell neurons. This neuronal infection can result in a novel syndrome characterized by cerebellar atrophy, gait ataxia, and incoordination, without associated demyelination.<sup>89</sup> This novel syndrome, distinct from PML, is called JCV granule cell neuronopathy (GCN).<sup>90,91</sup> JCV GCN appears to be caused by a JCV variant harboring a 10-bp deletion in the carboxyl terminus of the VP1 gene.<sup>92</sup> A histologic survey indicated that infection of granule cell neurons may be found in up to half of patients with PML.<sup>93</sup> This finding was confirmed in other JCV GCN isolates from various geographic origins, and other mutations occurring in this area of the VP1 gene were characterized, strengthening its association with infection of GCN.<sup>4,94</sup> More recently, JCV GCN has also been diagnosed in natalizumab-treated MS patients.<sup>95,96</sup>

### JC Virus Encephalopathy

JCV can also infect hemispheric cortical pyramidal neurons. The clinical presentation differs from both classic PML and JCV GCN. Brain lesions are initially restricted to the gray matter at MRI, and the patient presents with a global cognitive decline and aphasia, rather than with focal neurologic deficits such as sensory or motor dysfunction.<sup>97</sup> The JCV strain isolated from cortical samples had a unique deletion in the agnoprotein gene.<sup>4,98</sup>

This deletion impairs VP1 production and virion expression, which may explain a restrictive infection of JCV in cortical pyramidal neurons.<sup>99,100</sup>

### JC Virus Meningitis

Although JCV is not routinely tested for in the CSF of patients with meningeal symptoms, several studies have documented JCV as the only pathogen present in the CSF of patients with typical meningitis symptoms, including neck stiffness and diplopia.<sup>101–103</sup> Whether these cases result from JCV primary infection or reactivation is unclear. A case report of JCV meningitis in an HIV-negative patient had reported productive JCV infection of meningeal and choroid plexus cells with limited infiltration of the cerebral parenchyma.<sup>104,105</sup> Because JCV PCR is not routinely performed in the CSF of patients with meningitis who do not have CNS lesions, the exact incidence of JCV meningitis is unknown.

### Diagnosis

The diagnosis of PML can be established through either histopathologic or clinical pathways. Whereas the first requires demonstration of JCV infection in the brain, the latter is based on clinical and radiologic findings consistent with PML and not better explained by other disorders, coupled with demonstration of JCV DNA by PCR in CSF. Diagnostic classification includes certain, probable, possible, and not PML.<sup>106</sup>

Because PML remains a relatively rare disease, it is often misdiagnosed, with a median time from initial symptoms to diagnosis of 74 days.<sup>107</sup>

### Imaging

Radiographically, PML brain lesions typically appear as multiple white matter lesions, usually sparing the cortex, that do not conform to particular vascular territories. No associated mass effect or contrast enhancement is generally seen. Both computed tomography (CT) and MRI can be used in the diagnosis of PML. Lesions appear in the white

matter as hypodense or patchy on CT scans; at MRI, they are areas of hyperintense (bright) signal on T2-weighted and fluid-attenuated inversion recovery (FLAIR) images and show as hypointense (dark) signal on T1-weighted images. MRI is more sensitive than CT and is the imaging method of choice for diagnosis of PML. As the name *progressive multifocal leukoencephalopathy* suggests, multiple lesions are usually present; they are often located in the subcortical white matter or in the cerebellar peduncles.<sup>108,109</sup> Lesions may also be found in gray matter structures, such as the basal ganglia or thalamus, which also contain myelinated fibers. Furthermore, HCS can also be seen overlying white matter lesions of PML.<sup>77</sup> In addition, PML may have atypical presentations, including unifocal and miliary patterns.<sup>110,111</sup> Although features of MRI appearance are not linked to survival time, the rare detection of mass effect predicts poor prognosis.<sup>112</sup> Lastly, atypical PML lesions, including ones with contrast enhancement, are seen in natalizumab-associated PML<sup>113</sup> and in inflammatory PML that results from IRIS (Fig. 144.2).<sup>79</sup>

### Brain Biopsy


Brain biopsy is the gold standard for the diagnosis of PML. This test has a sensitivity that ranges from 64% to 96% and a specificity of 100%. Histologic examination shows demyelinated areas, with reactive gliosis and enlarged and bizarre astrocytes, and macrophages that contain phagocytosed myelin and cellular debris. These lesions are located in both cortical and subcortical regions of the brain.<sup>114</sup> JCV can be detected in infected oligodendrocytes with enlarged amphophilic nuclei located at the periphery of the lesions. Intraparenchymal and perivascular infiltrates by CD8<sup>+</sup> T cells are usually present in PML-IRIS (Fig. 144.3).<sup>80</sup>

### Cerebrospinal Fluid Polymerase Chain Reaction Assay

Brain biopsy may not be feasible in some patients, depending on the location of the lesions and the disability of the patient. In addition, biopsy carries an inherent morbidity and mortality risk. Therefore CSF examination is more commonly used as an alternative method of diagnosis. Patients with PML generally have a nonspecific CSF profile, including mild pleocytosis, with slightly elevated protein and normal glucose levels. The detection of JCV DNA with PCR assay of CSF had a sensitivity of 72% to 92% and a specificity of 92% to 100%<sup>115</sup> before extensive availability of ART. However, in the post-ART era, the sensitivity of PCR detection of CSF has decreased significantly to 58% with no changes in the specificity in patients with HIV presenting with PML while on ART.<sup>116</sup>

### Prognosis

PML is a fatal disease in most cases. Before the ART era, only 10% of patients who were HIV positive with PML lived longer than a year after diagnosis.<sup>117,118</sup> After the extensive use of ART, the 1-year survival rate has increased to 50%.<sup>118</sup> A number of studies have delineated several prognostic markers, which include patients who present with a lower JCV CSF burden,<sup>119</sup> detectable JCV-specific immune response in blood and CSF,<sup>120–123</sup> and development of inflammatory immune response in the CNS.<sup>124,125</sup> Because positive JCV serology is detected in most individuals, including those with PML, the humoral immune response alone is not able to prevent viral reactivation. However, cellular immune response to JCV has been detected in patients with PML, and a strong response is associated with better prognosis. Studies have documented the presence of cytotoxic T lymphocytes (CTLs) that recognize two human leukocyte antigen (HLA) A\*0201-restricted epitopes of the major capsid protein VP1 in patients with PML who survived more than 1 year.<sup>121,126–128</sup> Presence of JCV-specific CTLs was associated with an increase in 1-year survival of PML from 46% to 73%.<sup>129</sup> In a study of 66 patients, PML survivors were 22.2 times more likely to have both CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses against JCV early after disease onset than PML progressors.<sup>130</sup> Furthermore, elevation of myo-inositol, a marker of inflammation detected with MRS in PML lesions, was associated with a better prognosis.<sup>125,131</sup> Finally, hyperperfusion within PML lesions, measured with arterial spin labeling MRI, was associated with a worse prognosis.<sup>132</sup>



**FIG. 144.2** Magnetic resonance images of patient with human immunodeficiency virus infection with classic progressive multifocal leukoencephalopathy at presentation. (A) Fluid-attenuated inversion recovery (FLAIR) image shows lesions in right and left frontal lobes (arrows). (B) No enhancement is evident after gadolinium injection on a T1-weighted image (arrows). (C) FLAIR image after antiretroviral therapy shows progressive multifocal leukoencephalopathy-immune reconstitution inflammatory syndrome displaying progression of lesions in both hemispheres (arrows). (D) Peripheral enhancement can be seen in T1-weighted image after gadolinium enhancement (arrows).

## Therapy

No specific treatment exists for PML. However, in patients infected with HIV, initiation or optimization of ART has been associated with an increase in PML survival rate at 1 year from 10% to 50%.<sup>118</sup> For patients who are HIV negative, improvement of the immune status through reduction of immunosuppression may allow for the adaptive immune system to take control of the infection. Because a strong cellular immunity against JCV has been associated with a better clinical outcome, immunotherapies aimed at boosting this immune response may become a treatment option for PML.<sup>133</sup> Immune reconstitution after allogeneic HSCT has been associated with selective control of JCV reactivation.<sup>134</sup> Cytarabine has shown an *in vitro* effect in controlling JCV replication and multiplication.<sup>135</sup> One randomized controlled clinical trial with patients who were HIV positive with PML did not show efficacy,<sup>136</sup> but one small retrospective study did show stabilization after intravenous treatment with cytarabine<sup>137</sup> in 7 of 19 HIV-negative patients (37%) with PML who had leukemia or lymphoma. Cidofovir is another potent antiviral agent that has shown *in vitro* activities against murine polyomavirus and SV40, but it has not been tested against JCV or BKV. However, multiple studies in different centers with cidofovir in patients both with and without HIV have not shown a significant effect in changing the disease course.<sup>118,138–140</sup> Lastly, because *in vitro* studies revealed that JCV infects cells via the serotonin receptor 5-HT<sub>2A</sub>,


mirtazapine, a serotonin receptor blocker, has been considered as a potential candidate for containment of JCV infection.<sup>42</sup> However, clinical improvement of PML with mirtazapine remains anecdotal so far.<sup>141</sup> A large pharmacologic screening study suggested that mefloquine, an antimalarial medication, has the ability to inhibit JCV replication in cell culture.<sup>142</sup> A multicenter study examining the role of mefloquine in the treatment of PML was terminated ahead of schedule because of lack of efficacy.<sup>143</sup>

Immunotherapy against JCV is a potential future option. This includes administration of interleukin 7 (IL-7) which boosted both CD4<sup>+</sup> and CD8<sup>+</sup> T-cell counts in a patient with idiopathic lymphocytopenia who survived PML.<sup>144</sup>

Another option is adoptive transfer therapy, in which virus-specific T cells are stimulated either from patient's own lymphocytes or from a matching donor's lymphocytes. Broadly neutralizing antibodies against multiple JCV VP1 capsid mutations could potentially be infused as therapy.<sup>144a</sup> JCV-specific vaccine against a specific strain was given to a PML patient and induced cognate neutralizing antibodies.<sup>145</sup>

In treatment of patients with inflammatory PML, in whom brain swelling and mass effect can be present, a short course of corticosteroids including prednisone, dexamethasone, and methylprednisolone may be used sparingly for mitigation of the inflammation surrounding the





**FIG. 144.3** Histologic features of JC virus (JCV) infection in the central nervous system. (A) Classic progressive multifocal leukoencephalopathy (PML), with demyelinating lesion in white matter (arrow) surrounded by multiple JCV-infected glial cells (dark blue, arrowheads). (B) PML-immune response inflammatory syndrome, with marked lymphocytic perivascular infiltrates (arrows). (C) JCV granule cell neuronopathy in a patient with human immunodeficiency virus infection with focal JCV infection of granule cell neurons (dark blue, arrows). (D) JCV encephalopathy with hemispheric cortical neurons (brown) infected with JCV (blue, arrows). (A, C, and D courtesy Dr. Christian Wüthrich; B courtesy Dr. Françoise Gray.)

lesion, especially with a risk for brain herniation. However, the use of these medications in PML-IRIS cases remains a matter of debate.<sup>146,147</sup> In patients with HIV, no evidence exists that interruption of ART during corticosteroid treatment is beneficial. In published case studies, most clinicians chose to continue ART after the diagnosis of PML-IRIS. Short-term interruption of ART can be used temporarily to reduce the uncontrolled immune reaction. However, clinicians need to be aware that disruptions in ART have been associated with higher HIV viral load rebound and lower CD4 counts.<sup>148,149</sup> Furthermore, whether a recurrence of IRIS may develop when ART is restarted is also unclear. Corticosteroids, including dexamethasone, prednisone, and hydrocortisone, have been used anecdotally for their antiinflammatory and immunosuppressive effects.<sup>150</sup> However, no clear guidelines exist for corticosteroid treatments of patients with PML-IRIS. Therefore these medications should be used only in cases with clear clinical or radiologic worsening attributable to IRIS, including life-threatening situations associated with cerebral edema and impending brain herniation.

### NEPHROPATHY AND OTHER BK VIRUS-ASSOCIATED DISEASES

#### Epidemiology

Although BKV-induced nephropathy rarely occurs in a native kidney,<sup>151,152</sup> the prevalence rate of this condition in kidney transplant recipients ranges from 1% to 10%.<sup>153,154</sup> Diagnosis is made an average of 44 weeks after transplantation, with a peak around 24 weeks.<sup>27</sup> With the use of potent immunosuppressive medications, such as tacrolimus and mycophenolate, a trend has been seen toward increasing prevalence of BKV-induced nephropathy in renal transplant patients.<sup>155,156</sup> Incidences of BK viremia and viruria decrease significantly 1 year after transplantation, diminishing the risk for BKV reactivation.<sup>157</sup> In addition, the incidence rate of BKV-induced ureteral stenosis in renal transplant patients is estimated to be 3%.<sup>158</sup> Lastly, hemorrhagic cystitis is the most

prevalent BKV-associated complication and occurs in 10% to 25% of bone marrow transplant recipients.<sup>159,160</sup>

#### Pathogenesis

BKV primary infection occurs in childhood and is asymptomatic. Thereafter, the virus remains latent in the kidney. Most cases of asymptomatic BK viruria are not associated with nephropathy or hemorrhagic cystitis. Therefore BKV pathogenesis is the result of several factors, including host predisposition, target organ damage, and immunosuppression. During the initial posttransplant period, severe therapeutic immunosuppressive conditioning regimens are a trigger for reactivation of BKV in renal cells. Such reactivation may lead to productive infection and lysis of kidney tubular epithelial cells. Although the ability of the host to mount BKV-specific immunity can be limited by immunosuppressive medications, the resulting cell lysis and necrosis mediated by the immune cells may trigger further renal dysfunction. In patients with renal transplantation, BK nephropathy is associated with risk factors including age, sex, race, BK serostatus of the recipients, ABO incompatibility, and types of immunosuppression used.<sup>161–164</sup> In HSCT patients, BKV seropositivity, older age, high anti-BKV immunoglobulin G (IgG) levels before transplant, and lower total lymphocyte count have been associated with increased risk for BKV reactivation.<sup>165,166</sup> Furthermore, in the late engraftment period, the recovery of immune function may produce additional injury from inflammation caused by immune reconstitution, resulting in hemorrhagic cystitis.<sup>167</sup> Lastly, alloimmune dysregulation may play an important role because BKV-induced hemorrhagic cystitis is rarely detected in autologous transplant recipients who received similar conditioning regimens.<sup>168</sup>

#### Clinical Manifestations

##### Nephropathy

BKV has a tropism for cells of the genitourinary tract. Infection with BKV in patients with immunocompromise can result in asymptomatic

hematuria, both hemorrhagic and nonhemorrhagic cystitis, and ureteral stenosis. Furthermore, BKV can induce interstitial nephritis in patients with HIV or renal transplantation. However, in patients with a renal allograft, BKV-induced renal tubular epithelial infection results in nephropathy, which is not associated with any specific immunosuppressive medication. BK viruria and viremia usually precede nephropathy. The factors associated with detection of BKV in urine and blood include high donor BK antibody titer and, possibly, absence of the HLA-C7 class I allele in both donor and recipient.<sup>169</sup> In addition, a higher risk for development of BKV nephropathy is associated with older age, male gender, comorbidity of diabetes mellitus, white race, and placement of ureteral stent.<sup>154,170</sup> Onset of BKV nephropathy after transplant ranges from 6 days to 5 years, with a mean time of 10 to 13 months.<sup>171,172</sup> Clinical manifestations of BKV nephropathy are similar to that of graft rejection, with slowly increasing serum creatinine levels without symptoms. Hematuria and fever can be detected in some patients.<sup>173,174</sup> Laboratory findings suggest renal insufficiency and urinary abnormalities. A rise in quantitative BKV serology can be seen with reactivation of the virus; however, this increase does not prevent progression of disease.<sup>169</sup>

### Ureteral Stenosis

Renal transplant patients with ureteral stenosis do not usually present with pain or discomfort because the transplanted kidney is not innervated. However, patients can present with urinary obstruction and laboratory findings of elevated serum creatinine levels.

### Hemorrhagic Cystitis

BK viruria in HSCT patients is associated with increased mortality.<sup>175</sup> Because of BKV tropism for renal urinary tract cells, up to 50% of HSCT patients develop BK viruria within 2 months of transplant,<sup>36,159</sup> usually after engraftment.<sup>176</sup> The rate of BK viruria is no different between allogeneic versus autologous grafts,<sup>177</sup> but only a portion of those with viruria develop hemorrhagic cystitis (10%–25%), ureteral stenosis, and interstitial nephritis. Furthermore, most cases of hemorrhagic cystitis occur in allogeneic HSCT recipients who also have graft-versus-host disease,<sup>168</sup> which indicates that immune reconstitution is part of the pathogenesis. Myeloablative conditioning regimen and graft-versus-host diseases have been identified as risk factors that predispose HSCT patients to BK viruria and hemorrhagic cystitis.<sup>178–180</sup> Diagnosis of BKV-induced hemorrhagic cystitis is considered when postengraftment HSCT patients develop hematuria, dysuria, urgency, frequency, or suprapubic pain. Severe bleeding and clot formation can lead to complications, including urinary tract obstruction and renal failure.

### Infections Outside the Renal System

BKV has been reported to disseminate to extrarenal organs. Multiple case reports have described encephalitis in immunosuppressed patients with detection of BKV in their CSF and brain tissues with PCR assay.<sup>181–185</sup> Furthermore, BKV-associated pneumonia and pneumonitis are also reported in HSCT patients after hemorrhagic cystitis.<sup>186–188</sup> Detection of BKV in the urine has also been associated with genitourinary tumors.<sup>189</sup>

### Diagnosis

#### Urine Polymerase Chain Reaction Assay

In the appropriate clinical setting, detection of BK viruria and renal insufficiency is diagnostic of BKV-induced nephropathy. Detection of BKV in urine often precedes viral detection in the blood, and sustained viruria, defined as detection in two or more consecutive urine samples, can be 100% sensitive and 94% specific for BK viremia.<sup>190</sup> Urine cytologic evaluation that reveals decoy cells (enlarged nucleus with a single large basophilic intranuclear inclusion) is an indication of viral infection, but is not specific to BKV infection because these cells are also associated with CMV and adenovirus infections.<sup>191,192</sup> Although detection of BKV DNA with PCR assay in the urine of patients with hemorrhagic cystitis is sensitive, it is nonspecific because asymptomatic BK viruria is common. However, the presence of an increased BK viral load in urine along with hematuria supports the diagnosis.

### Renal Biopsy

Renal biopsy is essential in the diagnosis of nephropathy. Histologic examination of renal biopsy shows viral replication in the tubular epithelium cells with large intranuclear inclusions and cell detachment (Fig. 144.4).<sup>193</sup> These cytopathic changes are initially localized to the medulla and distal tubules, with progression to the proximal tubules. However, renal biopsy is associated with a false-negative rate of up to 30% because of the focal nature of the disease.

### Plasma Polymerase Chain Reaction Assay

BKV DNA can be detected in the plasma of kidney transplant recipients.<sup>194</sup> However, this test is more useful in ruling out BKV nephropathy than in diagnosing it. BKV PCR in blood has a negative predictive value of 100% but a positive predictive value of only 50% for BKV nephropathy.<sup>36</sup> One study, however, indicated that plasma BK viral load above 10<sup>4</sup> copies/mL has a sensitivity and specificity of 93% in prediction of histologic manifestations of nephropathy.<sup>195</sup> Unlike in nephropathy, BKV is usually not detected in the blood of patients with hemorrhagic cystitis or ureteral stenosis. However, one study found that BK viremia in HSCT patients is an independent predictor for development of post-HSCT renal impairment.<sup>196</sup>

### Prognosis


BKV-induced nephropathy is associated with irreversible graft failure in 1% to 10% of cases.<sup>70</sup> Histologic findings of mild viral cytopathic changes with minimal inflammatory infiltrates or fibrosis correlated with better prognosis for allograft outcome.<sup>195</sup> Studies have shown<sup>197</sup> decreased incidence of nephropathy with regular and early screening for BK viruria, viremia, and urine cytologic detection of virus-infected decoy cells and subsequent reduction of immunosuppressive agents.<sup>33,198–207</sup> Poor outcome from BKV-induced nephropathy is influenced by higher BKV serum viral load, immunosuppression with tacrolimus, and delayed diagnosis.<sup>208</sup> In post-renal transplant patients with BKV reactivation, those patients with BKV-specific cellular responses curtail viral replication within a median duration of 1 month without other interventions.<sup>209</sup> Furthermore, in patients who developed BKV nephropathy, a robust BKV-specific cellular immune response correlates with decreased viruria and viremia, whereas the humoral immune response correlated with decreased viremia only.<sup>210</sup>

### Prevention and Therapy Nephropathy

Systematic screening for BKV reactivation in post-renal transplant recipients is effective in preventing nephropathy. Detection of BKV in blood is usually followed by reduction of immunosuppressants to revive adaptive immune responses against BKV.<sup>211</sup> Screening for BKV in urine can detect reactivation earlier because viruria almost always occurs before viremia. However, not all patient with viruria will progress to develop viremia.

Treatment of BKV-induced nephropathy is reduction of immunosuppression. Multiple medications, including cidofovir, leflunomide, quinolones, and intravenous immune globulin (IVIG), have shown varying success in small clinical studies.<sup>212–216</sup> However, patients in these studies also had concomitant decrease of immunosuppressive medications. Because no specific antiviral therapy exists for BKV, reduction of immunosuppression is the therapy of choice<sup>200</sup> and has to be balanced with the increase in risk for graft rejection. The clearance of BKV from plasma is a surrogate marker for resolution of renal tissue pathologic conditions. Fluoroquinolones, such as ciprofloxacin, can inhibit BKV replication *in vitro* by inhibiting BKV-encoded DNA gyrase.<sup>217</sup> Two small retrospective studies demonstrated decreased BKV infection after renal transplantation in the group with exposure to fluoroquinolones.<sup>218,219</sup> However, two recently conducted randomized, placebo-controlled studies showed that 3 months of treatment with levofloxacin did not reduce the rate of BK viruria, but did result in increased resistance to levofloxacin in bacterial isolates.<sup>220,221</sup> Antibody-based treatment of BKV was based on the finding that neutralizing antibodies against BKV are specific to viral capsid sequences.<sup>222,223</sup> Several case series have demonstrated a potential role for IVIG in





**FIG. 144.4 Histology of BK virus nephropathy.** (A) Hematoxylin and eosin stain demonstrating tubular epithelial cells with viral inclusions and interstitial inflammation. (B) Simian virus 40 (SV40) antibody stain showing BKV-infected cells. (C) Electron microscopy image showing BK virus. (Courtesy I. Stillman, Department of Pathology, Beth Israel Deaconess Medical Center, Boston MA.)

reducing BK viral load in the plasma of renal transplant recipients.<sup>224–226</sup> A randomized double-blinded study is currently underway to determine the efficacy of IVIG in treatment of BK viremia in renal transplant recipients. Lastly, potential future treatments include switching immunosuppression to mTOR inhibitors and changes in T-cell transfer of immunity.

### Ureteral Stenosis

Post-renal transplant patients with the occurrence of BKV-associated ureteral stenosis can benefit from reduced immune suppression. Further treatments primarily involve surgical interventions that relieve the obstruction.

### Hemorrhagic Cystitis

Treatment of hemorrhagic cystitis is symptomatic and includes continuous bladder irrigation, analgesia, hyperhydration, forced diuresis, and transfusion to maintain platelet levels above 50,000 cells/mm<sup>3</sup> and hematocrit values greater than 25%. Intravesicular cidofovir has been used in treatment of hemorrhagic cystitis in several case reports and retrospective studies.<sup>227–232</sup> Although this agent may provide some symptomatic relief, there is no definitive decrease of urine BKV load with treatment.<sup>233</sup> Randomized controlled trials are needed to clarify benefits of intravesicular cidofovir use in hemorrhagic cystitis. Prophylaxis with ciprofloxacin resulted in reduced incidence of BKV-associated hemorrhagic cystitis after HSCT in a small retrospective study.<sup>234</sup>

### Key References

The complete reference list is available online at Expert Consult.

- Tan CS, Koralnik IJ. Progressive multifocal leukoencephalopathy and other disorders caused by JC virus: clinical features and pathogenesis. *Lancet Neurol*. 2010;9:425–437.
- Gheuens S, Wuthrich C, Koralnik IJ. Progressive multifocal leukoencephalopathy: why gray and white matter. *Annu Rev Pathol*. 2013;8:189–215.
- Padgett BL, Walker DL, Zur Rhein GM, et al. Cultivation of papova-like virus from human brain with progressive multifocal leukoencephalopathy. *Lancet*. 1971;1:1257–1260.
- Gardner SD, Field AM, Coleman DV, et al. New human papovavirus (B.K.) isolated from urine after renal transplantation. *Lancet*. 1971;1:1253–1257.
- Gaynor AM, Nissen MD, Whitley DM, et al. Identification of a novel polyomavirus from patients with acute respiratory tract infections. *PLoS Pathog*. 2007;3:e64.
- Astrom KE, Mancall EL, Richardson EP. Progressive multifocal leukoencephalopathy. *Brain*. 1958;81:93–127.
- van der Meijden E, Janssens RW, Lauber C, et al. Discovery of a new human polyomavirus associated with trichodysplasia spinulosa in an immunocompromised patient. *PLoS Pathog*. 2010;6:e1001024.
- DeCaprio JA, Garcea RL. A cornucopia of human polyomaviruses. *Nat Rev Microbiol*. 2013;11:264–276.

20. Knowles WA, Pipkin P, Andrews N, et al. Population-based study of antibody to the human polyomaviruses BKV and JCV and the simian polyomavirus SV40. *J Med Virol*. 2003;71:115–123.
24. Kitamura T, Aso Y, Kuniyoshi N, et al. High incidence of urinary JC virus excretion in nonimmunosuppressed older patients. *J Infect Dis*. 1990;161:1128–1133.
25. Koralnik IJ, Schmitz JE, Lifton MA, et al. Detection of JC virus DNA in peripheral blood cell subpopulations of HIV-1-infected individuals. *J Neurovirol*. 1999;5:430–435.
27. Nickeleit V, Klimkait T, Binet IF, et al. Testing for polyomavirus type BK DNA in plasma to identify renal-allograft recipients with viral nephropathy. *N Engl J Med*. 2000;342:1309–1315.
29. Bressollette-Bodin C, Coste-Burel M, Hourmant M, et al. A prospective longitudinal study of BK virus infection in 104 renal transplant recipients. *Am J Transplant*. 2005;5:1926–1933.
33. Hirsch HH, Knowles W, Dickenmann M, et al. Prospective study of polyomavirus type BK replication and nephropathy in renal-transplant recipients. *N Engl J Med*. 2002;347:488–496.
42. Elphick GF, Querbes W, Jordan JA, et al. The human polyomavirus, JCV, uses serotonin receptors to infect cells. *Science*. 2004;306:1380–1383.
49. Feng H, Shuda M, Chang Y, et al. Clonal integration of a polyomavirus in human merkel cell carcinoma. *Science*. 2008;319:1096–1100.
58. Amend KL, Turnbull B, Foskett N, et al. Incidence of progressive multifocal leukoencephalopathy in patients without HIV. *Neurology*. 2010;75:1326–1332.
69. Bloomgren G, Richman S, Hotermans C, et al. Risk of natalizumab-associated progressive multifocal leukoencephalopathy. *N Engl J Med*. 2012;366:1870–1880.
72. Tan CS, Dezube BJ, Bhargava P, et al. Detection of JC virus DNA and proteins in the bone marrow of HIV-positive and HIV-negative patients: implications for viral latency and neurotropic transformation. *J Infect Dis*. 2009;199:881–888.
73. Tan CS, Ellis LC, Wuthrich C, et al. JC virus latency in the brain and extraneural organs of patients with and without progressive multifocal leukoencephalopathy. *J Virol*. 2010;84:9200–9209.
75. Lima MA, Drislane FW, Koralnik IJ. Seizures and their outcome in progressive multifocal leukoencephalopathy. *Neurology*. 2006;66:262–264.
78. Post MJ, Thurnher MM, Clifford DB, et al. CNS-immune reconstitution inflammatory syndrome in the setting of HIV infection, 1. Overview and discussion of progressive multifocal leukoencephalopathy-immune reconstitution inflammatory syndrome and cryptococcal-immune reconstitution inflammatory syndrome. *AJNR Am J Neuroradiol*. 2013;34:1297–1307.
80. Vendrely A, Bienvenu B, Gasnault J, et al. Fulminant inflammatory leukoencephalopathy associated with HAART-induced immune restoration in AIDS-related progressive multifocal leukoencephalopathy. *Acta Neuropathol (Berl)*. 2005;109:449–455.
81. Gheuens S, Ngo L, Wang X, et al. Metabolic profile of PML lesions in patients with and without IRIS: an observational study. *Neurology*. 2012;79:1041–1048.
91. Koralnik IJ, Wuthrich C, Dang X, et al. JC virus granule cell neuronopathy: a novel clinical syndrome distinct from progressive multifocal leukoencephalopathy. *Ann Neurol*. 2005;57:576–580.
93. Wuthrich C, Cheng YM, Joseph JT, et al. Frequent infection of cerebellar granule cell neurons by polyomavirus JC in progressive multifocal leukoencephalopathy. *J Neuropathol Exp Neurol*. 2009;68:15–25.
94. Dang X, Vidal JE, Oliveira AC, et al. JC virus granule cell neuronopathy is associated with VP1 C terminus mutants. *J Gen Virol*. 2012;93(Pt 1):175–183.
97. Wuthrich C, Dang X, Westmoreland S, et al. Fulminant JC virus encephalopathy with productive infection of cortical pyramidal neurons. *Ann Neurol*. 2009;65:742–748.
106. Berger JR, Aksamit AJ, Clifford DB, et al. PML diagnostic criteria: consensus statement from the AAN neuroinfectious disease section. *Neurology*. 2013;80:1430–1438.
109. Whiteman ML, Post MJ, Berger JR, et al. Progressive multifocal leukoencephalopathy in 47 HIV-seropositive patients: neuroimaging with clinical and pathologic correlation. *Radiology*. 1993;187:233–240.
121. Du Pasquier RA, Kuroda MJ, Schmitz JE, et al. Low frequency of cytotoxic T lymphocytes against the novel HLA-A\*0201-restricted JC virus epitope VP1(p36) in patients with proven or possible progressive multifocal leukoencephalopathy. *J Virol*. 2003;77:11918–11926.
122. Lima MA, Marzocchetti A, Autissier P, et al. Frequency and phenotype of JC virus-specific CD8<sup>+</sup> T lymphocytes in the peripheral blood of patients with progressive multifocal leukoencephalopathy. *J Virol*. 2007;81:3361–3368.
126. Koralnik IJ, Du Pasquier RA, Letvin NL. JC virus-specific cytotoxic T lymphocytes in individuals with progressive multifocal leukoencephalopathy. *J Virol*. 2001;75:3483–3487.
127. Koralnik IJ, Du Pasquier RA, Kuroda MJ, et al. Association of prolonged survival in HLA-A2<sup>+</sup> progressive multifocal leukoencephalopathy patients with a CTL response specific for a commonly recognized JC virus epitope. *J Immunol*. 2002;168:499–504.
128. Du Pasquier RA, Kuroda MJ, Zheng Y, et al. A prospective study demonstrates an association between JC virus-specific cytotoxic T lymphocytes and the early control of progressive multifocal leukoencephalopathy. *Brain*. 2004;127:1970–1978.
136. Hall CD, Dafni U, Simpson D, et al. Failure of cytarabine in progressive multifocal leukoencephalopathy associated with human immunodeficiency virus infection. *AIDS* clinical trials group 243 team [see comments]. *N Engl J Med*. 1998;338:1345–1351.
146. Tan K, Roda R, Ostrow L, et al. PML-IRIS in patients with HIV infection: clinical manifestations and treatment with steroids. *Neurology*. 2009;72:1458–1464.
154. Hirsch HH, Brennan DC, Drachenberg CB, et al. Polyomavirus-associated nephropathy in renal transplantation: interdisciplinary analyses and recommendations. *Transplantation*. 2005;79:1277–1286.
159. Arthur RR, Shah KV, Baust SJ, et al. Association of BK viremia with hemorrhagic cystitis in recipients of bone marrow transplants. *N Engl J Med*. 1986;315:230–234.
179. Giraud G, Priftakis P, Bogdanovic G, et al. BK-viremia and haemorrhagic cystitis are more frequent in allogeneic haematopoietic stem cell transplant patients receiving full conditioning and unrelated-HLA-mismatched grafts. *Bone Marrow Transplant*. 2008;41:737–742.
194. Boudreaux AA, Courtemanche C, Latulippe E, et al. Screening for polyomavirus associated nephropathy in renal transplantation with blood viral load measurement. *J Clin Virol*. 2009;45:318–321.
195. Drachenberg CB, Papadimitriou JC, Hirsch HH, et al. Histological patterns of polyomavirus nephropathy: correlation with graft outcome and viral load. *Am J Transplant*. 2004;4:2082–2092.
197. Schaub S, Hirsch HH, Dickenmann M, et al. Reducing immunosuppression preserves allograft function in presumptive and definitive polyomavirus-associated nephropathy. *Am J Transplant*. 2010;10:2615–2623.
203. Hammann AL, Oqvist B, Wahlgren J, et al. Systematic screening of BK virus by real-time PCR prevents BK virus associated nephropathy in renal transplant recipients. *J Med Virol*. 2011;83:1959–1965.
204. Almeras C, Vetromile F, Garrigue V, et al. Monthly screening for BK viremia is an effective strategy to prevent BK virus nephropathy in renal transplant recipients. *Transpl Infect Dis*. 2011;13:101–108.
210. Chen Y, Trofe J, Gordon J, et al. Interplay of cellular and humoral immune responses against BK virus in kidney transplant recipients with polyomavirus nephropathy. *J Virol*. 2006;80:3495–3505.
213. Williams JW, Javadi B, Kadambi PV, et al. Leflunomide for polyomavirus type BK nephropathy. *N Engl J Med*. 2005;352:1157–1158.
228. Eisen DP, Fraser IR, Sung LM, et al. Decreased viral load and symptoms of polyomavirus-associated chronic interstitial cystitis after intravesical cidofovir treatment. *Clin Infect Dis*. 2009;48:e86–e88.
231. Cesaro S, Hirsch HH, Faraci M, et al. Cidofovir for BK virus-associated hemorrhagic cystitis: a retrospective study. *Clin Infect Dis*. 2009;49:233–240.
234. Miller AN, Glode A, Hogan KR, et al. Efficacy and safety of ciprofloxacin for prophylaxis of polyomavirus BK virus-associated hemorrhagic cystitis in allogeneic hematopoietic stem cell transplantation recipients. *Biol Blood Marrow Transplant*. 2011;17:1176–1181.

## References

- Knowles WA. Discovery and epidemiology of the human polyomaviruses BK virus (BKV) and JC virus (JCV). *Adv Exp Med Biol*. 2006;577:19–45.
- Weber T, Trebst C, Frye S, et al. Analysis of the systemic and intrathecal humoral immune response in progressive multifocal leukoencephalopathy. *J Infect Dis*. 1997;176:250–254.
- Tan CS, Koralnik IJ. Progressive multifocal leukoencephalopathy and other disorders caused by JC virus: clinical features and pathogenesis. *Lancet Neurol*. 2010;9:425–437.
- Gheuens S, Wuthrich C, Koralnik IJ. Progressive multifocal leukoencephalopathy: why gray and white matter. *Annu Rev Pathol*. 2013;8:189–215.
- Berger JR, Kaszovitz B, Post MJ, et al. Progressive multifocal leukoencephalopathy associated with human immunodeficiency virus infection: a review of the literature with a report of sixteen cases. *Ann Intern Med*. 1987;107:78–87.
- Johne R, Buck CB, Allander T, et al. Taxonomical developments in the family polyomaviridae. *Arch Virol*. 2011;156:1627–1634.
- Padgett BL, Walker DL, ZuRhein GM, et al. Cultivation of papova-like virus from human brain with progressive multifocal leukoencephalopathy. *Lancet*. 1971;1:1257–1260.
- Gardner SD, Field AM, Coleman DV, et al. New human papovavirus (B.K.) isolated from urine after renal transplantation. *Lancet*. 1971;1:1253–1257.
- Allander T, Andreasson K, Gupta S, et al. Identification of a third human polyomavirus. *J Virol*. 2007;81:4130–4136.
- Gaynor AM, Nissen MD, Whitley DM, et al. Identification of a novel polyomavirus from patients with acute respiratory tract infections. *PLoS Pathog*. 2007;3:e64.
- Siebrasse EA, Reyes A, Lim ES, et al. Identification of MW polyomavirus, a novel polyomavirus in human stool. *J Virol*. 2012;86:10321–10326.
- Lim ES, Reyes A, Antonio M, et al. Discovery of STL polyomavirus, a polyomavirus of ancestral recombinant origin that encodes a unique T antigen by alternative splicing. *Virology*. 2013;436:295–303.
- Feng H, Shuda M, Chang Y, et al. Clonal integration of a polyomavirus in human Merkel cell carcinoma. *Science*. 2008;319:1096–1100.
- van der Meijden E, Janssens RW, Lauber C, et al. Discovery of a new human polyomavirus associated with trichodysplasia spinulosa in an immunocompromised patient. *PLoS Pathog*. 2010;6:e1001024.
- Schwalter RM, Pastrana DV, Pumphrey KA, et al. Merkel cell polyomavirus and two previously unknown polyomaviruses are chronically shed from human skin. *Cell Host Microbe*. 2010;7:509–515.
- Scuda N, Hofmann J, Calvignac-Spencer S, et al. A novel human polyomavirus closely related to the African green monkey-derived lymphotropic polyomavirus. *J Virol*. 2011;85:4586–4590.
- DeCaprio JA, Garcea RL. A cornucopia of human polyomaviruses. *Nat Rev Microbiol*. 2013;11:264–276.
- Brown P, Tsai T, Gajdusek DC. Seroepidemiology of human papovaviruses: discovery of virgin populations and some unusual patterns of antibody prevalence among remote peoples of the world. *Am J Epidemiol*. 1975;102:331–340.
- Gossai A, Waterboer T, Nelson HH, et al. Seroepidemiology of human polyomaviruses in a US population. *Am J Epidemiol*. 2016;183:61–69.
- Knowles WA, Pipkin P, Andrews N, et al. Population-based study of antibody to the human polyomaviruses BKV and JCV and the simian polyomavirus SV40. *J Med Virol*. 2003;71:115–123.
- Shah KV, Daniel RW, Warszawski RM. High prevalence of antibodies to BK virus, an SV40-related papovavirus, in residents of Maryland. *J Infect Dis*. 1973;128:784–787.
- Boldorini R, Allegrini S, Miglio U, et al. Serological evidence of vertical transmission of JC and BK polyomaviruses in humans. *J Gen Virol*. 2011;92:1044–1050.
- Doerries K. Human polyomavirus JC and BK persistent infection. *Adv Exp Med Biol*. 2006;577:102–116.
- Kitamura T, Aso Y, Kuniyoshi N, et al. High incidence of urinary JC virus excretion in nonimmunosuppressed older patients. *J Infect Dis*. 1990;161:1128–1133.
- Koralnik IJ, Schmitz JE, Lifton MA, et al. Detection of JC virus DNA in peripheral blood cell subpopulations of HIV-1-infected individuals. *J Neurovirol*. 1999;5:430–435.
- Tornatore C, Berger JR, Houff SA, et al. Detection of JC virus DNA in peripheral lymphocytes from patients with and without progressive multifocal leukoencephalopathy. *Ann Neurol*. 1992;31:454–462.
- Nickeleit V, Klimkait T, Binet IF, et al. Testing for polyomavirus type BK DNA in plasma to identify renal-allograft recipients with viral nephropathy. *N Engl J Med*. 2000;342:1309–1315.
- Howell DN, Smith SR, Butterly DW, et al. Diagnosis and management of BK polyomavirus interstitial nephritis in renal transplant recipients. *Transplantation*. 1999;68:1279–1288.
- Bressollette-Bodin C, Coste-Burel M, Hourmant M, et al. A prospective longitudinal study of BK virus infection in 104 renal transplant recipients. *Am J Transplant*. 2005;5:1926–1933.
- Ahsan N, Shah KV. Polyomaviruses and human diseases. *Adv Exp Med Biol*. 2006;577:1–18.
- Knowles WA, Pillay D, Johnson MA, et al. Prevalence of long-term BK and JC excretion in HIV-infected adults and lack of correlation with serological markers. *J Med Virol*. 1999;59:474–479.
- Gluck TA, Knowles WA, Johnson MA, et al. BK virus-associated haemorrhagic cystitis in an HIV-infected man. *AIDS*. 1994;8:391–392.
- Hirsch HH, Knowles W, Dickenmann M, et al. Prospective study of polyomavirus type BK replication and nephropathy in renal-transplant recipients. *N Engl J Med*. 2002;347:488–496.
- Rorij NM, Shea MM, Satyanarayana G, et al. BK virus disease after allogeneic stem cell transplantation: a cohort analysis. *Biol Blood Marrow Transplant*. 2014;20:564–570.
- Satyanarayana G, Hammond SP, Broge TA, et al. BK polyomavirus reactivation after reduced-intensity double umbilical cord blood cell transplantation. *Transpl Immunol*. 2015;32:116–120.
- Hirsch HH, Steiger J. Polyomavirus BK. *Lancet Infect Dis*. 2003;3:611–623.
- Safak M, Barrucco R, Darbinyan A, et al. Interaction of JC virus agno protein with T antigen modulates transcription and replication of the viral genome in glial cells. *J Virol*. 2001;75:1476–1486.
- Khalili K, White MK, Sawa H, et al. The agnoprotein of polyomaviruses: a multifunctional auxiliary protein. *J Cell Physiol*. 2005;204:1–7.
- Takahashi K, Sekizuka T, Fukumoto H, et al. Deep-sequence identification and role in virus replication of a JC virus quasispecies in patients with progressive multifocal leukoencephalopathy. *J Virol*. 2017;91.
- Marzocchetti A, Wuthrich C, Tan CS, et al. Rearrangement of the JC virus regulatory region sequence in the bone marrow of a patient with rheumatoid arthritis and progressive multifocal leukoencephalopathy. *J Neurovirol*. 2008;14:455–458.
- Mayberry CL, Nelson CDS, Maginnis MS. JC polyomavirus attachment and entry: potential sites for PML therapeutics. *Curr Clin Microbiol Rep*. 2017;4:132–141.
- Elphick GF, Querbes W, Jordan JA, et al. The human polyomavirus, JCV, uses serotonin receptors to infect cells. *Science*. 2004;306:1380–1383.
- Eash S, Manley K, Gasparovic M, et al. The human polyomaviruses. *Cell Mol Life Sci*. 2006;63:865–876.
- Dugan AS, Eash S, Atwood WJ. Update on BK virus entry and intracellular trafficking. *Transpl Infect Dis*. 2006;8:62–67.
- Nguyen NL, Le BM, Wang D. Serologic evidence of frequent human infection with WU and KI polyomaviruses. *Emerg Infect Dis*. 2009;15:1199–1205.
- Carter JJ, Paulson KG, Wipf GC, et al. Association of merkel cell polyomavirus-specific antibodies with merkel cell carcinoma. *J Natl Cancer Inst*. 2009;101:1510–1522.
- van der Meijden E, Kazem S, Burgers MM, et al. Seroepidemiology of trichodysplasia spinulosa-associated polyomaviruses. *Emerg Infect Dis*. 2011;17:1355–1363.
- Trusch F, Klein M, Finsterbusch T, et al. Seroepidemiology of human polyomavirus 9 and cross-reactivity to African green monkey-derived lymphotropic polyomavirus. *J Gen Virol*. 2012;93:698–705.
- Allander T, Andreasson K, Gupta S, et al. Identification of a third human polyomavirus. *J Virol*. 2007;81:4130–4136.
- Bialasiewicz S, Whitley DM, Lambert SB, et al. A newly reported human polyomavirus, KI virus, is present in the respiratory tract of Australian children. *J Clin Virol*. 2007;40:15–18.
- Han TH, Chung JY, Koo JW, et al. WU polyomavirus in children with acute lower respiratory tract infections, South Korea. *Emerg Infect Dis*. 2007;13:1766–1768.
- Abadi Kiasari B, Valley PJ, Corless CE, et al. Age-related pattern of KI and WU polyomavirus infection. *J Clin Virol*. 2008;43:123–125.
- Bialasiewicz S, Rockett RJ, Barraclough KA, et al. Detection of recently discovered human polyomaviruses in a longitudinal kidney transplant cohort. *Am J Transplant*. 2016;16:2734–2740.
- Ho J, Jedrych JJ, Feng H, et al. Human polyomavirus 7-associated pruritic rash and viremia in transplant recipients. *J Infect Dis*. 2015;211:1560–1565.
- Astrom KE, Mancall EL, Richardson EP. Progressive multifocal leukoencephalopathy. *Brain*. 1958;81:93–127.
- Power C, Gladden JG, Halliday W, et al. AIDS- and non-AIDS-related PML association with distinct p53 polymorphism. *Neurology*. 2000;54:743–746.
- Gheuens S, Pierone G, Peeters P, et al. Progressive multifocal leukoencephalopathy in individuals with minimal or occult immunosuppression. *J Neurol Neurosurg Psychiatry*. 2010;81:247–254.
- Amend KL, Turnbull B, Foskett N, et al. Incidence of progressive multifocal leukoencephalopathy in patients without HIV. *Neurology*. 2010;75:1326–1332.
- Langer-Gould A, Atlas SW, Green AJ, et al. Progressive multifocal leukoencephalopathy in a patient treated with natalizumab. *N Engl J Med*. 2005;353:375–381.
- Kleinschmidt-DeMasters BK, Tyler KL. Progressive multifocal leukoencephalopathy complicating treatment with natalizumab and interferon beta-1a for multiple sclerosis. *N Engl J Med*. 2005;353:369–374.
- Van Assche G, Van Ranst M, Sciort R, et al. Progressive multifocal leukoencephalopathy after natalizumab therapy for Crohn's disease. *N Engl J Med*. 2005;353:362–368.
- Nieuwkamp DJ, Murk JL, van Oosten BW, et al. PML in a patient without severe lymphocytopenia receiving dimethyl fumarate. *N Engl J Med*. 2015;372:1474–1476.
- Rosenkranz T, Novas M, Terborg C. PML in a patient with lymphocytopenia treated with dimethyl fumarate. *N Engl J Med*. 2015;372:1476–1478.
- Rey J, Belmecheri N, Bouayed N, et al. JC papovavirus leukoencephalopathy after first line treatment with CHOP and rituximab. *Haematologica*. 2007;92:e101.
- Bonavita S, Conforti R, Russo A, et al. Infratentorial progressive multifocal leukoencephalopathy in a patient treated with fludarabine and rituximab. *Neurol Sci*. 2008;29:37–39.
- Harris HE. Progressive multifocal leukoencephalopathy in a patient with systemic lupus erythematosus treated with rituximab. *Rheumatology (Oxford)*. 2008;47:224–225.
- Ermis U, Weis J, Schulz JB. PML in a patient treated with fumaric acid. *N Engl J Med*. 2013;368:1657–1658.
- Genetech: Important Safety Information. [http://www.genetech.com/gene/products/information/pdf/raptiva\\_dhpc.pdf](http://www.genetech.com/gene/products/information/pdf/raptiva_dhpc.pdf). Accessed 2008.
- Bloomgren G, Richman S, Hotermans C, et al. Risk of natalizumab-associated progressive multifocal leukoencephalopathy. *N Engl J Med*. 2012;366:1870–1880.
- Monaco MC, Jensen PN, Hou J, et al. Detection of JC virus DNA in human tonsil tissue: evidence for site of initial viral infection. *J Virol*. 1998;72:9918–9923.
- Goudsmit J, Wertheim-van Dillen P, van Strien A, et al. The role of BK virus in acute respiratory tract disease and the presence of BKV DNA in tonsils. *J Med Virol*. 1982;10:91–99.
- Tan CS, Dezuze BJ, Bhargava P, et al. Detection of JC virus DNA and proteins in the bone marrow of HIV-positive and HIV-negative patients: implications for viral latency and neurotropic transformation. *J Infect Dis*. 2009;199:881–888.
- Tan CS, Ellis LC, Wuthrich C, et al. JC virus latency in the brain and extraneural origins of patients with and without progressive multifocal leukoencephalopathy. *J Virol*. 2010;84:9200–9209.
- Bernal-Cano E, Joseph JT, Koralnik IJ. Spinal cord lesions of progressive multifocal leukoencephalopathy in an acquired immunodeficiency syndrome patient. *J Neurovirol*. 2007;13:474–476.
- Lima MA, Drislane FW, Koralnik IJ. Seizures and their outcome in progressive multifocal leukoencephalopathy. *Neurology*. 2006;66:262–264.
- Miskin DP, Herman ST, Ngo LH, et al. Predictors and characteristics of seizures in survivors of progressive multifocal leukoencephalopathy. *J Neurovirol*. 2016;22:464–471.
- Khoury MN, Alsop DC, Agnihotri SP, et al. Hyperintense cortical signal on magnetic resonance imaging reflects focal leukocortical encephalitis and seizure risk in progressive multifocal leukoencephalopathy. *Ann Neurol*. 2014;75:659–669.
- Post MJ, Thurnher MM, Clifford DB, et al. CNS-immune reconstitution inflammatory syndrome in the setting of HIV infection. I. Overview and discussion of progressive multifocal leukoencephalopathy-immune reconstitution inflammatory syndrome and cryptococcal-immune reconstitution inflammatory syndrome. *AJNR Am J Neuroradiol*. 2013;34:1297–1307.
- Du Pasquier RA, Koralnik IJ. Inflammatory reaction in progressive multifocal leukoencephalopathy: harmful or beneficial? *J Neurovirol*. 2003;9:25–31.
- Vendrey A, Bienvenu B, Gasnault J, et al. Fulminant inflammatory leukoencephalopathy associated with HAART-induced immune restoration in AIDS-related



- progressive multifocal leukoencephalopathy. *Acta Neuropathol (Berl)*. 2005;109:449–455.
81. Gheuens S, Ngo L, Wang X, et al. Metabolic profile of PML lesions in patients with and without IRIS: an observational study. *Neurology*. 2012;79:1041–1048.
  82. Venkataramana A, Pardo CA, McArthur JC, et al. Immune reconstitution inflammatory syndrome in the CNS of HIV-infected patients. *Neurology*. 2006;67:383–388.
  83. D'Amico R, Sarkar S, Yusuff J, et al. Immune reconstitution after potent antiretroviral therapy in AIDS patients with progressive multifocal leukoencephalopathy. *Scand J Infect Dis*. 2007;39:347–350.
  84. Martinez JV, Mazzioti JV, Efron ED, et al. Immune reconstitution inflammatory syndrome associated with PML in AIDS: a treatable disorder. *Neurology*. 2006;67:1692–1694.
  85. Rushing EJ, Liappis A, Smirniotopoulos JD, et al. Immune reconstitution inflammatory syndrome of the brain: case illustrations of a challenging entity. *J Neuropathol Exp Neurol*. 2008;67:819–827.
  86. Kastrop O, Wanke I, Esser S, et al. Evolution of purely infratentorial PML under HAART: negative outcome under rapid immune reconstitution. *Clin Neurol Neurosurg*. 2005;107:509–513.
  87. Hoffmann C, Horst HA, Albrecht H, et al. Progressive multifocal leukoencephalopathy with unusual inflammatory response during antiretroviral treatment. *J Neurol Neurosurg Psychiatry*. 2003;74:1142–1144.
  88. Scarpazza C, Prosperini L, De Rossi N, et al. To do or not to do? plasma exchange and timing of steroid administration in progressive multifocal leukoencephalopathy. *Ann Neurol*. 2017;82:697–705.
  89. Hecht JH, Glenn OA, Wara DW, et al. JC virus granule cell neuronopathy in a child with CD40 ligand deficiency. *Pediatr Neurol*. 2007;36:186–189.
  90. Du Pasquier RA, Corey S, Margolin DH, et al. Productive infection of cerebellar granule cell neurons by JC virus in an HIV+ individual. *Neurology*. 2003;61:775–782.
  91. Koralnik IJ, Wuthrich C, Dang X, et al. JC virus granule cell neuronopathy: a novel clinical syndrome distinct from progressive multifocal leukoencephalopathy. *Ann Neurol*. 2005;57:576–580.
  92. Dang X, Koralnik IJ. A granule cell neuron-associated JC virus variant has a unique deletion in the VP1 gene. *J Gen Virol*. 2006;87:2533–2537.
  93. Wuthrich C, Cheng YM, Joseph JT, et al. Frequent infection of cerebellar granule cell neurons by polyomavirus JC in progressive multifocal leukoencephalopathy. *J Neuropathol Exp Neurol*. 2009;68:15–25.
  94. Dang X, Vidal JE, Oliveira AC, et al. JC virus granule cell neuronopathy is associated with VP1 C terminus mutants. *J Gen Virol*. 2012;93(Pt 1):175–183.
  95. Agnihotri SP, Dang X, Carter JL, et al. JCV GCN in a natalizumab-treated MS patient is associated with mutations of the VP1 capsid gene. *Neurology*. 2014;83:727–732.
  96. Schippling S, Kempf C, Buchele F, et al. JC virus granule cell neuronopathy and GCN-IRIS under natalizumab treatment. *Ann Neurol*. 2013;74:622–626.
  97. Wuthrich C, Dang X, Westmoreland S, et al. Fulminant JC virus encephalopathy with productive infection of cortical pyramidal neurons. *Ann Neurol*. 2009;65:742–748.
  98. Dang X, Wuthrich C, Gordon J, et al. JC virus encephalopathy is associated with a novel agnoprotein-deletion JCV variant. *PLoS ONE*. 2012;7:e35793.
  99. Ellis LC, Norton E, Dang X, et al. Agnogene deletion in a novel pathogenic JC virus isolate impairs VP1 expression and virion production. *PLoS ONE*. 2013;8:e80840.
  100. Wuthrich C, Koralnik IJ. Frequent infection of cortical neurons by JC virus in patients with progressive multifocal leukoencephalopathy. *J Neuropathol Exp Neurol*. 2012;71:54–65.
  101. Viallard JF, Ellie E, Lazaro E, et al. JC virus meningitis in a patient with systemic lupus erythematosus. *Lupus*. 2005;14:964–966.
  102. Blake K, Pillay D, Knowles W, et al. JC virus associated meningoencephalitis in an immunocompetent girl. *Arch Dis Child*. 1992;67:956–957.
  103. Behzad-Behbahani A, Klapper PE, Vallely PJ, et al. BKV-DNA and JCV-DNA in CSF of patients with suspected meningitis or encephalitis. *Infection*. 2003;31:374–378.
  104. Agnihotri SP, Wuthrich C, Dang X, et al. A fatal case of JC virus meningitis presenting with hydrocephalus in a human immunodeficiency virus-seronegative patient. *Ann Neurol*. 2014;76:140–147.
  105. Miskin DP, Koralnik IJ. Novel syndromes associated with JC virus infection of neurons and meningeal cells: no longer a gray area. *Curr Opin Neurol*. 2015;28:288–294.
  106. Berger JR, Aksamit AJ, Clifford DB, et al. PML diagnostic criteria: consensus statement from the AAN neuroinfectious disease section. *Neurology*. 2013;80:1430–1438.
  107. Miskin DP, Ngo LH, Koralnik IJ. Diagnostic delay in progressive multifocal leukoencephalopathy. *Ann Clin Transl Neurol*. 2016;3:386–391.
  108. Skiest DJ. Focal neurological disease in patients with acquired immunodeficiency syndrome. *Clin Infect Dis*. 2002;34:103–115.
  109. Whiteman ML, Post MJ, Berger JR, et al. Progressive multifocal leukoencephalopathy in 47 HIV-seropositive patients: neuroimaging with clinical and pathologic correlation. *Radiology*. 1993;187:233–240.
  110. Horger M, Beschoner R, Beck R, et al. Common and uncommon imaging findings in progressive multifocal leukoencephalopathy (PML) with differential diagnostic considerations. *Clin Neurol Neurosurg*. 2012;114:1123–1130.
  111. Gheuens S, Buggy BP, Tlomak W, et al. A game of viral hide and seek: miliary PML masquerading as EBV encephalitis in an HIV(+) patient. *Clin Neurol Neurosurg*. 2013;115:1861–1863.
  112. Post MJ, Yiannoutsos C, Simpson D, et al. Progressive multifocal leukoencephalopathy in AIDS: are there any MR findings useful to patient management and predictive of patient survival? AIDS clinical trials group, 243 team. *AJNR Am J Neuroradiol*. 1999;20:1896–1906.
  113. Youssry TA, Pelletier D, Cadavid D, et al. Magnetic resonance imaging pattern in natalizumab-associated progressive multifocal leukoencephalopathy. *Ann Neurol*. 2012;72:779–787.
  114. Moll NM, Rietsch AM, Ransohoff AJ, et al. Cortical demyelination in PML and MS: similarities and differences. *Neurology*. 2008;70:336–343.
  115. Cinque P, Scarpellini P, Vago L, et al. Diagnosis of central nervous system complications in HIV-infected patients: cerebrospinal fluid analysis by the polymerase chain reaction. *AIDS*. 1997;11:1–17.
  116. Marzocchetti A, Di Giambenedetto S, Cingolani A, et al. Reduced rate of diagnostic positive detection of JC virus DNA in cerebrospinal fluid in cases of suspected progressive multifocal leukoencephalopathy in the era of potent antiretroviral therapy. *J Clin Microbiol*. 2005;43:4175–4177.
  117. Berger JR, Pall L, Lanska D, et al. Progressive multifocal leukoencephalopathy in patients with HIV infection. *J Neurovirol*. 1998;4:59–68.
  118. Antinori A, Cingolani A, Lorenzini P, et al. Clinical epidemiology and survival of progressive multifocal leukoencephalopathy in the era of highly active antiretroviral therapy: data from the Italian registry investigative neuro AIDS (IRINA). *J Neurovirol*. 2003;9:47–53.
  119. Yiannoutsos CT, Major EO, Curfman B, et al. Relation of JC virus DNA in the cerebrospinal fluid to survival in acquired immunodeficiency syndrome patients with biopsy-proven progressive multifocal leukoencephalopathy. *Ann Neurol*. 1999;45:816–821.
  120. Koralnik IJ. Overview of the cellular immunity against JC virus in progressive multifocal leukoencephalopathy. *J Neurovirol*. 2002;8:59–65.
  121. Du Pasquier RA, Kuroda MJ, Schmitz JE, et al. Low frequency of cytotoxic T lymphocytes against the novel HLA-A\*0201-restricted JC virus epitope VP1(p36) in patients with proven or possible progressive multifocal leukoencephalopathy. *J Virol*. 2003;77:11918–11926.
  122. Lima MA, Marzocchetti A, Autissier P, et al. Frequency and phenotype of JC virus-specific CD8<sup>+</sup> T lymphocytes in the peripheral blood of patients with progressive multifocal leukoencephalopathy. *J Virol*. 2007;81:3361–3368.
  123. Du Pasquier RA, Autissier P, Zheng Y, et al. Presence of JC virus-specific CTL in the cerebrospinal fluid of PML patients: rationale for immune-based therapeutic strategies. *AIDS*. 2005;19:2069–2076.
  124. Berger JR, Levy RM, Flomenhoft D, et al. Predictive factors for prolonged survival in acquired immunodeficiency syndrome-associated progressive multifocal leukoencephalopathy. *Ann Neurol*. 1998;44:341–349.
  125. Katz-Brull R, Lenkinski RE, Du Pasquier RA, et al. Elevation of myoinositol is associated with disease containment in progressive multifocal leukoencephalopathy. *Neurology*. 2004;63:897–900.
  126. Koralnik IJ, Du Pasquier RA, Letvin NL. JC virus-specific cytotoxic T lymphocytes in individuals with progressive multifocal leukoencephalopathy. *J Virol*. 2001;75:3483–3487.
  127. Koralnik IJ, Du Pasquier RA, Kuroda MJ, et al. Association of prolonged survival in HLA-A2<sup>+</sup> progressive multifocal leukoencephalopathy patients with a CTL response specific for a commonly recognized JC virus epitope. *J Immunol*. 2002;168:499–504.
  128. Du Pasquier RA, Kuroda MJ, Zheng Y, et al. A prospective study demonstrates an association between JC virus-specific cytotoxic T lymphocytes and the early control of progressive multifocal leukoencephalopathy. *Brain*. 2004;127:1970–1978.
  129. Marzocchetti A, Tompkins T, Clifford DB, et al. Determinants of survival in progressive multifocal leukoencephalopathy. *Neurology*. 2009;73:1551–1558.
  130. Gheuens S, Bord E, Kesari S, et al. Role of CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses against JC virus in the outcome of patients with progressive multifocal leukoencephalopathy (PML) and PML with immune reconstitution inflammatory syndrome. *J Virol*. 2011;85:7256–7263.
  131. Chang L, Ernst T, Tornatore C, et al. Metabolite abnormalities in progressive multifocal leukoencephalopathy by proton magnetic resonance spectroscopy. *Neurology*. 1997;48:836–845.
  132. Khoury MN, Gheuens S, Ngo L, et al. Hyperperfusion in progressive multifocal leukoencephalopathy is associated with disease progression and absence of immune reconstitution inflammatory syndrome. *Brain*. 2013;136(Pt 11):3441–3450.
  133. Marzocchetti A, Lima M, Tompkins T, et al. Efficient in vitro expansion of JC virus-specific CD8(+) T-cell responses by JCV peptide-stimulated dendritic cells from patients with progressive multifocal leukoencephalopathy. *Virology*. 2009;383:173–177.
  134. Tan SC, Broge TA, Ngo L, et al. Immune reconstitution after allogeneic hematopoietic stem cell transplantation is associated with selective control of JC virus reactivation. *Biol Blood Marrow Transplant*. 2014;20:992–999.
  135. Hou J, Major EO. The efficacy of nucleoside analogs against JC virus multiplication in a persistently infected human fetal brain cell line. *J Neurovirol*. 1998;4:451–456.
  136. Hall CD, Dafni U, Simpson D, et al. Failure of cytarabine in progressive multifocal leukoencephalopathy associated with human immunodeficiency virus infection. AIDS clinical trials group 243 team [see comments]. *N Engl J Med*. 1998;338:1345–1351.
  137. Aksamit AJ. Treatment of non-AIDS progressive multifocal leukoencephalopathy with cytosine arabinoside. *J Neurovirol*. 2001;7:386–390.
  138. Marra CM, Rajcic N, Barker DE, et al. A pilot study of cidofovir for progressive multifocal leukoencephalopathy in AIDS. *AIDS*. 2002;16:1791–1797.
  139. Osorio S, de la Camara R, Golbano N, et al. Progressive multifocal leukoencephalopathy after stem cell transplantation, unsuccessfully treated with cidofovir. *Bone Marrow Transplant*. 2002;30:963–966.
  140. Herrlinger U, Schwarzler F, Beck R, et al. Progressive multifocal leukoencephalopathy: cidofovir therapy in three patients with underlying hematological disease. *J Neurol*. 2003;250:612–614.
  141. Verma S, Cikurel K, Koralnik IJ, et al. Mirtazapine in progressive multifocal leukoencephalopathy associated with polycythemia vera. *J Infect Dis*. 2007;196:709–711.
  142. Brickelmaier M, Lugovskoy A, Kartikayan R, et al. Identification and characterization of mefloquine efficacy against JC virus in vitro. *Antimicrob Agents Chemother*. 2009;53:1840–1849.
  143. Clifford DB, Nath A, Cinque P, et al. A study of mefloquine treatment for progressive multifocal leukoencephalopathy: results and exploration of predictors of PML outcomes. *J Neurovirol*. 2013;19:351–358.
  144. Miskin DP, Chalkias SG, Dang X, et al. Interleukin-7 treatment of PML in a patient with idiopathic lymphocytopenia. *Neurol Neuroimmunol Neuroinflamm*. 2016;3:e213.
  - 144a. Jelcic I, Combautier B, Jelcic I, et al. Broadly neutralizing human monoclonal JC polyomavirus VP1-specific antibodies as candidate therapeutics for progressive multifocal leukoencephalopathy. *Sci Transl Med*. 2015;7:306ra150.
  145. Ray U, Cinque P, Gerevini S, et al. JC polyomavirus mutants escape antibody-mediated neutralization. *Sci Transl Med*. 2015;7:306ra151.
  146. Tan K, Roda R, Ostrow L, et al. PML-IRIS in patients with HIV infection: clinical manifestations and treatment with steroids. *Neurology*. 2009;72:1458–1464.
  147. Berger JR. Steroids for PML-IRIS: a double-edged sword. *Neurology*. 2009;72:1454–1455.
  148. Li X, Margolick JB, Conover CS, et al. Interruption and discontinuation of highly active antiretroviral therapy in the multi-center AIDS cohort study. *J Acquir Immune Defic Syndr*. 2005;38:320–328.
  149. Steingrover R, Pogany K, Fernandez Garcia E, et al. HIV-1 viral rebound dynamics after a single treatment interruption depends on time of initiation of highly active antiretroviral therapy. *AIDS*. 2008;22:1583–1588.

150. Czock D, Keller F, Rasche FM, et al. Pharmacokinetics and pharmacodynamics of systemically administered glucocorticoids. *Clin Pharmacokinet*. 2005;44:61–98.
151. Limaye AP, Smith KD, Cook L, et al. Polyomavirus nephropathy in native kidneys of non-renal transplant recipients. *Am J Transplant*. 2005;5:614–620.
152. Barber CE, Hewlett TJ, Geldenhuys L, et al. BK virus nephropathy in a heart transplant recipient: case report and review of the literature. *Transpl Infect Dis*. 2006;8:113–121.
153. Nicleleit V, Hirsch HH, Zeiler M, et al. BK-virus nephropathy in renal transplants-tubular necrosis, MHC-class II expression and rejection in a puzzling game. *Nephrol Dial Transplant*. 2000;15:324–332.
154. Hirsch HH, Brennan DC, Drachenberg CB, et al. Polyomavirus-associated nephropathy in renal transplantation: interdisciplinary analyses and recommendations. *Transplantation*. 2005;79:1277–1286.
155. Ramos E, Drachenberg CB, Portocarrero M, et al. BK virus nephropathy diagnosis and treatment: experience at the University of Maryland renal transplant program. *Clin Transpl*. 2002;143–153.
156. Dharnidharka VR, Cherikh WS, Abbott KC. An OPTN analysis of national registry data on treatment of BK virus allograft nephropathy in the United States. *Transplantation*. 2009;87:1019–1026.
157. Koukoulaki M, Grispou E, Pistolas D, et al. Prospective monitoring of BK virus replication in renal transplant recipients. *Transpl Infect Dis*. 2009;11:1–10.
158. Coleman DV, Mackenzie EF, Gardner SD, et al. Human polyomavirus (BK) infection and ureteric stenosis in renal allograft recipients. *J Clin Pathol*. 1978;31:338–347.
159. Arthur RR, Shah KV, Baust JS, et al. Association of BK viremia with hemorrhagic cystitis in recipients of bone marrow transplants. *N Engl J Med*. 1986;315:230–234.
160. Rorije NM, Shea MM, Satyanarayana G, et al. BK virus disease after allogeneic stem cell transplantation: a cohort analysis. *Biol Blood Marrow Transplant*. 2014;20:564–570.
161. Sharif A, Alachkar N, Bagnasco S, et al. Incidence and outcomes of BK virus allograft nephropathy among ABO- and HLA-incompatible kidney transplant recipients. *Clin J Am Soc Nephrol*. 2012;7:1320–1327.
162. Manitpitkul W, Drachenberg C, Ramos E, et al. Maintenance immunosuppressive agents as risk factors for BK virus nephropathy: a case-control study. *Transplantation*. 2009;88:83–88.
163. Steubl D, Baumann M, Schuster T, et al. Risk factors and interventional strategies for BK polyomavirus infection after renal transplantation. *Scand J Urol Nephrol*. 2012;46:466–474.
164. Sood P, Senanayake S, Sujiet K, et al. Donor and recipient BKV-specific IgG antibody and posttransplantation BKV infection: a prospective single-center study. *Transplantation*. 2013;95:896–902.
165. Wong AS, Chan KH, Cheng VC, et al. Relationship of pretransplantation polyoma BK virus serologic findings and BK viral reactivation after hematopoietic stem cell transplantation. *Clin Infect Dis*. 2007;44:830–837.
166. Drew RJ, Walsh A, Ni Laoi B, et al. BK virus (BKV) plasma dynamics in patients with BKV-associated hemorrhagic cystitis following allogeneic stem cell transplantation. *Transpl Infect Dis*. 2013;15:276–282.
167. Leung AY, Yuen KY, Kwong YL. Polyoma BK virus and haemorrhagic cystitis in haematopoietic stem cell transplantation: a changing paradigm. *Bone Marrow Transplant*. 2005;36:929–937.
168. Leung AY, Yuen KY, Cheng VC, et al. Clinical characteristics of and risk factors for herpes zoster after hematopoietic stem cell transplantation. *Haematologica*. 2002;87:444–446.
169. Bohl DL, Storch GA, Ryschewitsch C, et al. Donor origin of BK virus in renal transplantation and role of HLA C7 in susceptibility to sustained BK viremia. *Am J Transplant*. 2005;5:2213–2221.
170. Siparsky NF, Kushnir LF, Gallichio MH, et al. Ureteral stents: a risk factor for polyomavirus BK viremia in kidney transplant recipients undergoing protocol screening. *Transplant Proc*. 2011;43:2641–2644.
171. Randhawa PS, Finkelstein S, Scantlebury V, et al. Human polyoma virus-associated interstitial nephritis in the allograft kidney. *Transplantation*. 1999;67:103–109.
172. Ramos E, Drachenberg CB, Papadimitriou JC, et al. Clinical course of polyoma virus nephropathy in 67 renal transplant patients. *J Am Soc Nephrol*. 2002;13:2145–2151.
173. Vasudev B, Hariharan S, Hussain SA, et al. BK virus nephritis: risk factors, timing, and outcome in renal transplant recipients. *Kidney Int*. 2005;68:1834–1839.
174. Dall A, Hariharan S. BK virus nephritis after renal transplantation. *Clin J Am Soc Nephrol*. 2008;3:S68–S75.
175. Abudayeh A, Hamdi A, Lin H, et al. Symptomatic BK virus infection is associated with kidney function decline and poor overall survival in allogeneic hematopoietic stem cell recipients. *Am J Transplant*. 2016;16:1492–1502.
176. Bedi A, Miller CB, Hanson JL, et al. Association of BK virus with failure of prophylaxis against hemorrhagic cystitis following bone marrow transplantation. *J Clin Oncol*. 1995;13:1103–1109.
177. Azzi A, Cesaro S, Laszlo D, et al. Human polyomavirus BK (BKV) load and hemorrhagic cystitis in bone marrow transplantation patients. *J Clin Virol*. 1999;14:79–86.
178. Giraud G, Bogdanovic G, Priftakis P, et al. The incidence of hemorrhagic cystitis and BK-viremia in allogeneic hematopoietic stem cell recipients according to intensity of the conditioning regimen. *Haematologica*. 2006;91:401–404.
179. Giraud G, Priftakis P, Bogdanovic G, et al. BK-viremia and hemorrhagic cystitis are more frequent in allogeneic haematopoietic stem cell transplant patients receiving full conditioning and unrelated-HLA-mismatched grafts. *Bone Marrow Transplant*. 2008;41:737–742.
180. Dalianis T, Ljungman P. Full myeloablative conditioning and an unrelated HLA mismatched donor increase the risk for BK virus-positive hemorrhagic cystitis in allogeneic hematopoietic stem cell transplanted patients. *Anticancer Res*. 2011;31:939–944.
181. Behre G, Becker M, Christopeit M. BK virus encephalitis in an allogeneic hematopoietic stem cell recipient. *Bone Marrow Transplant*. 2008;42:499.
182. Lopes da Silva R, Ferreira I, Teixeira G, et al. BK virus encephalitis with thrombotic microangiopathy in an allogeneic hematopoietic stem cell transplant recipient. *Transpl Infect Dis*. 2011;13:161–167.
183. Green J, Saigal G, Rojas CP, et al. Rare presentation of BK encephalitis in a child: imaging and pathological findings. *Pediatr Radiol*. 2012;42:1145–1148.
184. Jorgensen GE, Hammann AL, Bratt G, et al. Identification of a unique BK virus variant in the CNS of a patient with AIDS. *J Med Virol*. 2003;70:14–19.
185. Bakri FG, Bahou YG, Al-Sammarrat FA, et al. Fatal encephalitis due to BK virus in a patient with common variable immunodeficiency: a case report. *J Clin Virol*. 2013;57:363–369.
186. Akazawa Y, Terada Y, Yamane T, et al. Fatal BK virus pneumonia following stem cell transplantation. *Transpl Infect Dis*. 2012;14:E142–E146.
187. Yapa HM, McLornan DP, Raj K, et al. Pneumonitis post-haematopoietic stem cell transplant—cytopathology clinches diagnosis. *J Clin Virol*. 2012;55:278–281.
188. Sandler ES, Aquino VM, Goss-Shoet E, et al. BK papova virus pneumonia following hematopoietic stem cell transplantation. *Bone Marrow Transplant*. 1997;20:163–165.
189. Papadimitriou JC, Randhawa P, Rinaldo CH, et al. BK polyomavirus infection and renourinary tumorigenesis. *Am J Transplant*. 2016;16:398–406.
190. Babel N, Fendt J, Karaivanov S, et al. Sustained BK viremia as an early marker for the development of BKV-associated nephropathy: analysis of 4128 urine and serum samples. *Transplantation*. 2009;88:89–95.
191. Kahan AV, Coleman DV, Koss LG. Activation of human polyomavirus infection—detection by cytologic technics. *Am J Clin Pathol*. 1980;74:326–332.
192. Traystman MD, Gupta PK, Shah KV, et al. Identification of viruses in the urine of renal transplant recipients by cytomorphology. *Acta Cytol*. 1980;24:501–510.
193. Drachenberg RC, Drachenberg CB, Papadimitriou JC, et al. Morphological spectrum of polyoma virus disease in renal allografts: diagnostic accuracy of urine cytology. *Am J Transplant*. 2001;1:373–381.
194. Boudreaux AA, Courtemanche C, Latulippe E, et al. Screening for polyomavirus associated nephropathy in renal transplantation with blood viral load measurement. *J Clin Virol*. 2009;45:318–321.
195. Drachenberg CB, Papadimitriou JC, Hirsch HH, et al. Histological patterns of polyomavirus nephropathy: correlation with graft outcome and viral load. *Am J Transplant*. 2004;4:2082–2092.
196. O'Donnell PH, Swanson K, Josephson MA, et al. BK virus infection is associated with hematuria and renal impairment in recipients of allogeneic hematopoietic stem cell transplants. *Biol Blood Marrow Transplant*. 2009;15:1038–1048, e1.
197. Schaub S, Hirsch HH, Dickenmann M, et al. Reducing immunosuppression preserves allograft function in presumptive and definitive polyomavirus-associated nephropathy. *Am J Transplant*. 2010;10:2615–2623.
198. Burgos D, Lopez V, Cabello M, et al. Polyomavirus BK nephropathy: the effect of an early diagnosis on renal function or graft loss. *Transplant Proc*. 2006;38:2409–2411.
199. Almeras C, Foulongne V, Garrigue V, et al. Does reduction in immunosuppression in viremic patients prevent BK virus nephropathy in de novo renal transplant recipients? A prospective study. *Transplantation*. 2008;85:1099–1104.
200. Brennan DC, Agha I, Bohl DL, et al. Incidence of BK with tacrolimus versus cyclosporine and impact of preemptive immunosuppression reduction. *Am J Transplant*. 2005;5:582–594.
201. Chung BH, Hong YA, Kim HG, et al. Clinical usefulness of BK virus plasma quantitative PCR to prevent BK virus associated nephropathy. *Transpl Int*. 2012;25:687–695.
202. Chakera A, Dyar OJ, Hughes E, et al. Detection of polyomavirus BK reactivation after renal transplantation using an intensive decoy cell surveillance program is cost-effective. *Transplantation*. 2011;92:1018–1023.
203. Hammann AL, Oqvist B, Wahlgren J, et al. Systematic screening of BK virus by real-time PCR prevents BK virus associated nephropathy in renal transplant recipients. *J Med Virol*. 2011;83:1959–1965.
204. Almeras C, Vetromile F, Garrigue V, et al. Monthly screening for BK viremia is an effective strategy to prevent BK virus nephropathy in renal transplant recipients. *Transpl Infect Dis*. 2011;13:101–108.
205. Gautam A, Patel V, Pelletier L, et al. Routine BK virus surveillance in renal transplantation—a single center's experience. *Transplant Proc*. 2010;42:4088–4090.
206. Renoult E, Coutlee F, Paquet M, et al. Evaluation of a preemptive strategy for BK polyomavirus-associated nephropathy based on prospective monitoring of BK viremia: a kidney transplantation center experience. *Transplant Proc*. 2010;42:4083–4087.
207. Knight RJ, Gaber LW, Patel SJ, et al. Screening for BK viremia reduces but does not eliminate the risk of BK nephropathy: a single-center retrospective analysis. *Transplantation*. 2013;95:949–954.
208. Schwarz A, Linnenweber-Held S, Heim A, et al. Factors influencing viral clearing and renal function during polyomavirus BK-associated nephropathy after renal transplantation. *Transplantation*. 2012;94:396–402.
209. Schachner T, Muller K, Stein M, et al. BK virus-specific immunity kinetics: a predictor of recovery from polyomavirus BK-associated nephropathy. *Am J Transplant*. 2011;11:2443–2452.
210. Chen Y, Trofe J, Gordon J, et al. Interplay of cellular and humoral immune responses against BK virus in kidney transplant recipients with polyomavirus nephropathy. *J Virol*. 2006;80:3495–3505.
211. Kidney Disease: Improving Global Outcomes Transplant Work G. KDIGO clinical practice guideline for the care of kidney transplant recipients. *Am J Transplant*. 2009;9:S1–S155.
212. Leung AY, Chan MT, Yuen KY, et al. Ciprofloxacin decreased polyoma BK virus load in patients who underwent allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis*. 2005;40:528–537.
213. Williams JW, Javadi B, Kadambi PV, et al. Lefunomide for polyomavirus type BK nephropathy. *N Engl J Med*. 2005;352:1157–1158.
214. Sener A, House AA, Jevnikar AM, et al. Intravenous immunoglobulin as a treatment for BK virus associated nephropathy: one-year follow-up of renal allograft recipients. *Transplantation*. 2006;81:117–120.
215. Randhawa PS. Anti-BK virus activity of ciprofloxacin and related antibiotics. *Clin Infect Dis*. 2005;41:1366–1367, author reply 1367.
216. Krisl JC, Taber DJ, Pilch N, et al. Lefunomide efficacy and pharmacodynamics for the treatment of BK viral infection. *Clin J Am Soc Nephrol*. 2012;7:1003–1009.
217. Ferrazzi E, Peracchi M, Biasolo MA, et al. Antiviral activity of gyrase inhibitors norfloxacin, coumermycin A1 and nalidixic acid. *Biochem Pharmacol*. 1988;37:1885–1886.
218. Gabardi S, Waikar SS, Martin S, et al. Evaluation of fluoroquinolones for the prevention of BK viremia after renal transplantation. *Clin J Am Soc Nephrol*. 2010;5:1298–1304.
219. Wojciechowski D, Chandra R, Chandran S, et al. Ciprofloxacin prophylaxis in kidney transplant recipients reduces BK virus infection at 3 months but not at 1 year. *Transplantation*. 2012;94:1117–1123.
220. Knoll GA, Humar A, Fergusson D, et al. Levofloxacin for BK virus prophylaxis following kidney transplantation. *JAMA*. 2014;312:2106–2114.
221. Fyfe I. Transplantation: levofloxacin treatment does not prevent BK virus infection. *Nat Rev Nephrol*. 2015;11:65.
222. Pastrana DV, Brennan DC, Cuburu N, et al. Neutralization serotyping of BK polyomavirus infection in kidney transplant recipients. *PLoS Pathog*. 2012;8:e1002650.
223. Pastrana DV, Ray U, Magaldi TG, et al. BK polyomavirus genotypes represent distinct serotypes with distinct entry tropism. *J Virol*. 2013;87:10105–10113.
224. Vu D, Shah T, Ansari J, et al. Efficacy of intravenous immunoglobulin in the treatment of persistent BK viremia and BK virus nephropathy in renal transplant recipients. *Transplant Proc*. 2015;47:394–398.

225. Kable K, Davies CD, O'Connell PJ, et al. Clearance of BK virus nephropathy by combination antiviral therapy with intravenous immunoglobulin. *Transplant Direct*. 2017;3:e142.
226. Shah T, Vu D, Naraghi R, et al. Efficacy of intravenous immunoglobulin in the treatment of persistent BK viremia and BK virus nephropathy in renal transplant recipients. *Clin Transpl*. 2014;109–116.
227. Bridges B, Donegan S, Badros A. Cidofovir bladder instillation for the treatment of BK hemorrhagic cystitis after allogeneic stem cell transplantation. *Am J Hematol*. 2006;81:535–537.
228. Eisen DP, Fraser IR, Sung LM, et al. Decreased viral load and symptoms of polyomavirus-associated chronic interstitial cystitis after intravesical cidofovir treatment. *Clin Infect Dis*. 2009;48:e86–e88.
229. Ganguly N, Clough LA, Dubois LK, et al. Low-dose cidofovir in the treatment of symptomatic BK virus infection in patients undergoing allogeneic hematopoietic stem cell transplantation: a retrospective analysis of an algorithmic approach. *Transpl Infect Dis*. 2010;12:406–411.
230. Rao KV, Buie LW, Shea T, et al. Intravesicular cidofovir for the management of BK virus-associated cystitis. *Biol Blood Marrow Transplant*. 2009;15:391–392.
231. Cesaro S, Hirsch HH, Faraci M, et al. Cidofovir for BK virus-associated hemorrhagic cystitis: a retrospective study. *Clin Infect Dis*. 2009;49:233–240.
232. Philippe M, Ranchon F, Gilis L, et al. Cidofovir in the treatment of BK Virus-associated hemorrhagic cystitis after allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2016;22:723–730.
233. Mackey MC. Intravesicular cidofovir for the treatment of polyomavirus-associated hemorrhagic cystitis. *Ann Pharmacother*. 2012;46:442–446.
234. Miller AN, Glode A, Hogan KR, et al. Efficacy and safety of ciprofloxacin for prophylaxis of polyomavirus BK virus-associated hemorrhagic cystitis in allogeneic hematopoietic stem cell transplantation recipients. *Biol Blood Marrow Transplant*. 2011;17:1176–1181.



## Hepatitis B Virus

Chloe Lynne Thio and Claudia Hawkins

## SHORT VIEW SUMMARY

**Definition**

- Hepatitis B virus (HBV) causes chronic hepatitis B, which can lead to progressive liver disease, including cirrhosis and hepatocellular carcinoma.

**Epidemiology**

- Nearly 5% percent of the world's population has chronic hepatitis B, but prevalence varies widely (see Table 145.1).
- The highest HBV prevalence is in Asia, Africa, and parts of the Middle East.
- Chronic hepatitis B is the leading cause of end-stage liver disease worldwide.
- Transmission of HBV is through percutaneous, sexual routes, or perinatal. Perinatal transmission is the leading cause of HBV worldwide.

**Microbiology**

- HBV is a partially double-stranded DNA virus in the Hepadnaviridae family that primarily infects hepatocytes.
- HBV replication is through an RNA intermediate, so it has a reverse transcriptase.

**Clinical Presentation and Diagnosis**

- Clinical presentation of HBV varies from asymptomatic to fulminant hepatitis with liver failure.

- The risk of developing chronic hepatitis B is inversely proportional to age of acquisition of infection. Chronic hepatitis B is also more likely in high-risk patient groups, such as injection drug users, men who have sex with men, and human immunodeficiency virus–infected individuals.
- Laboratory diagnosis of HBV is by enzyme immunoassays, which detect various HBV antigens and antibodies, and by real-time polymerase chain reaction to detect HBV DNA (see Table 145.2).

**Therapy**

- Anti-HBV agents significantly reduce complications of chronic hepatitis B, including liver cirrhosis and hepatocellular carcinoma. These therapies are also effective in reducing the recurrence of HBV infection in transplant recipients and other immunocompromised hosts.
- HBV is treated with 180 µg of pegylated interferon-α (PEG IFN-α) weekly for 48 weeks or with nucleos(t)ide analogues, often indefinitely (see Table 145.4).
- The first-line nucleos(t)ide agents are entecavir 0.5 mg daily or tenofovir disoproxil fumarate

(TDF) 300 mg daily or tenofovir alafenamide 25 mg daily.

- New drugs are in early phase trials

**Prevention**

- Vaccination with the hepatitis B vaccine is recommended in all infants and children and for nonimmune adults at high risk for infection (see Table 145.6).
- Infants born to mothers with chronic hepatitis B should receive the hepatitis B vaccine and hepatitis B immune globulin (HBIG) at birth. Mothers with HBV DNA >1962 IU/mL should receive TDF in the third trimester of pregnancy to reduce the risk of transmission to the infant.
- Nonimmune individuals who have percutaneous, sexual, ocular, or mucous membrane exposure to HBV-infected fluids should receive postexposure prophylaxis with HBIG and hepatitis B vaccine.


**OVERVIEW**

Hepatitis B virus (HBV) infects more than 200 million people worldwide. It is the leading cause of chronic hepatitis, cirrhosis, and hepatocellular carcinoma (HCC), and these sequelae of chronic infection account for more than 800,000 deaths annually. The outcome of infection and spectrum of illness varies widely. During the acute phase, infections range from asymptomatic hepatitis to icteric hepatitis, including fulminant hepatitis. Once chronic infection is established, the spectrum of illness ranges from the asymptomatic, inactive hepatitis to progressive liver disease that results in the sequelae of end-stage liver disease, including cirrhosis and HCC. Even though HBV cannot be cultured easily, enough is known about the viral life cycle that specific antivirals are available to control viral replication in patients; however, no cure is available yet for hepatitis B. Identification of the correlates of protective immunity has facilitated the widespread use of a highly effective vaccine, which by preventing chronic carriage of the virus and the subsequent development of HCC, represents the first true cancer vaccine.

**Historical Background and Classification**

Hippocrates recognized the spread of jaundice by infectious agents as early as 4000 BC. Recently, the oldest known HBV was isolated from the mummified remains of a child discovered in the 16th century from

Naples, Italy.<sup>1</sup> The early modern cases of HBV infection were linked to the use of conventional viral vaccines, which were prepared from or contained human serum. In 1885 Lurman<sup>2</sup> described the appearance of jaundice in 15% of 1289 shipyard workers who received smallpox vaccine prepared from human lymph. Epidemics of hepatitis were also recorded after the administration of yellow fever vaccine, which was stabilized with human serum.<sup>3</sup> In the early part of the 20th century the increasing use of contaminated syringes and needles by diabetics on insulin and by patients treated for syphilis at venereal disease clinics elevated the importance of serum hepatitis.<sup>4,5</sup> This led to the association of hepatitis B with blood and blood products, as well as to its distinction from infectious hepatitis, caused by hepatitis A virus, a member of the Picornaviridae family.<sup>6</sup> The first hint of viral etiology came from the studies of Blumberg,<sup>7</sup> who reported the discovery of a human antigen in Australian Aborigines termed Australian (Au) antigen. Subsequently, the Au antigen came to be known as hepatitis B surface antigen (HBsAg), and its association with acute hepatitis was established. For this discovery, Dr. Baruch Blumberg received the Nobel Prize in Physiology and Medicine in 1976. In 1971 Dane, an electron microscopist, visualized the presence of 22-nm HBsAg subviral particles along with the complete 42-nm virus particles in the blood of hepatitis B patients (Fig. 145.1).<sup>8</sup>



**FIG. 145.1** Structure of hepatitis B virus (HBV) and hepatitis B surface antigen (HBsAg) particles. (A) Electron micrograph of negatively stained HBV. (B) Diagram of 42-nm HBV showing partially duplex DNA genome with a covalently linked protein at the 5' end of the complete positive strand. (C) Diagram of 22-nm HBsAg filament. (Modified from Flint SJ, Enquist LW, Racaniello VR, et al. Principles of Virology. 2nd ed. Washington, DC: American Society for Microbiology Press; 2004:808.)

Due to their unique biologic and molecular characteristics, as well as their liver tropism, HBV and HBV-like animal viruses were given the status of a new family designated Hepadnaviridae (*hepatotropic DNA viruses*). HBV, the human pathogen, and other mammalian hepatitis viruses with sequence homology and similar genome organization are grouped in the genus *Orthohepadnavirus*. The genus *Avihepadnavirus* includes viruses that infect ducks, geese, and herons. The hepadnavirus animal models, which include duck hepatitis B virus,<sup>9</sup> woodchuck hepatitis virus, and ground squirrel hepatitis virus, have been extensively studied and have contributed to the current knowledge of the molecular biology of HBV infection and replication. However, these viral genomes are considerably divergent from human HBV, and there are limited tools to study the immune reaction to the virus in these animals. The primary host for HBV is human, but HBsAg has been detected in other primates, such as chimpanzees, gibbons, orangutans, African green monkeys, and squirrel monkeys. However, the use of these animals for the study of pathogenesis is both difficult and expensive. Thus a convenient animal model for HBV remains unavailable; however, the discovery of the HBV receptor on hepatocytes has led to development of a more efficient tissue culture system that should facilitate new drug discovery. HBV was molecularly cloned from patients' sera in 1979 and its complete DNA sequence determined.<sup>10–12</sup>

## Virology

HBV is a small DNA virus, whose 3200-kilobase (kb) partially double-stranded DNA genome is maintained in a circular conformation.<sup>13</sup> Partially duplex DNA molecules represent incomplete synthesis of the viral DNA during morphogenesis (Fig. 145.2).<sup>14</sup> One of the unique features of HBV infection is the production of large quantities of subviral spherical and filamentous HBsAg particles, which are devoid of HBV DNA and so are noninfectious, in addition to complete virus particles. Under electron microscopy, these are distinguished by a diameter of 42 nm for complete virus (Dane) particles and 22-nm spherical and filamentous structures for subviral particles (see Fig. 145.1).<sup>8</sup> HBsAg is expressed on the exterior of all these particles. HBsAg subviral particles reach a titer of  $10^{13}$ /mL, whereas viral particles range in titer from  $10^4$ /mL to  $10^9$ /mL. All of these particles circulate in blood and permit convenient diagnosis of viral antigen by enzyme-linked immunosorbent assay or radioimmunoassay.<sup>15</sup> HBsAg forms the viral component of the lipoprotein envelope, which encloses a core shell containing the viral DNA genome and the virus-encoded polymerase protein. The nucleocapsid or core is composed of a 21-kilodalton (kDa) basic phosphoprotein commonly known as hepatitis B core antigen (HBcAg).<sup>16</sup> A cell-derived kinase activity has been shown to be associated with the virion particles,<sup>17</sup> but the functional significance of this enzyme is not understood.

The subviral particles are composed of surface antigen polypeptides and lipids derived from hepatocyte membranes. The lipid content is


approximately 30% by weight and includes phospholipids, cholesterol, cholesterol esters, and triglycerides.<sup>18,19</sup> Because of their high immunogenicity, purified HBsAg particles can be used as an HBV vaccine. HBsAg elicits neutralizing antibodies, which offer protection from reinfection.<sup>20</sup> The high titers of HBsAg in patients during natural infection can potentially serve to adsorb neutralizing antibody or promote T-cell exhaustion, thus protecting the virus from host defenses.<sup>21</sup>

## Attachment, Entry, and Hepatotropism

HBV primarily infects hepatocytes. The pre-S1 domain of the large HBsAg protein promotes attachment and entry of HBV into the hepatocyte via a liver cell-specific receptor recently identified as Na (sodium) taurocholate cotransporting polypeptide (NTCP), which is an integral membrane protein used in bile acid transport.<sup>22,23</sup> There is evidence that hepatocyte entry may be a multistep process, including binding to heparan sulfate proteoglycans,<sup>24</sup> which are found on a variety of cells, and clathrin-mediated endocytosis.<sup>25</sup> The viral envelope fuses with the endosomal membrane and delivers core particles into the cytoplasm, which are transported to the nuclear pore complex, allowing viral DNA to enter the nucleus.<sup>26</sup> In the nucleus the partial strand is repaired, forming a double-stranded covalently closed circular (ccc) DNA form (Fig. 145.3), which is the template for replication (see “Replication”).<sup>9</sup> Liver specificity is also displayed at the level of viral gene expression, which is controlled largely by the promoters and enhancers (see Fig. 145.2).

## Viral Genome

HBV uses unique transcriptional and translational strategies to maximize the limited coding capacity of its genome. The HBV DNA consists of the complete (negative) strand, which contains a protein covalently linked to its 5' terminus, and the incomplete (positive) strand, displaying variable length and bearing a 5' capped oligoribonucleotide (see Figs. 145.1 and 145.2).<sup>27,28</sup> The asymmetry of the DNA strands reflects the incomplete synthesis of DNA during maturation of viral particles. HBV DNA codes for four overlapping open reading frames (ORFs): S, for the surface antigen or envelope gene (HBsAg); C, for the nucleocapsid (core), and “e” antigen gene (HBcAg, HBeAg); P, for the polymerase gene; and X, for the HBx gene (see Fig. 145.2). The surface antigen ORF contains three in-frame initiator codons from which small or major (S), middle (M), and large (L) kDa HBsAg polypeptides are synthesized, respectively. These polypeptides are variably glycosylated to yield several species: S, p24/gp26; M, p30/gp33/gp37; and L, p39/gp42 kDa HBsAg polypeptides, respectively. L and M HBsAg contain characteristic pre-S1 and pre-S2 domains, and hence these proteins are also referred to as pre-S1 and pre-S2 proteins, respectively. The subviral particles are composed of mostly S and lesser amounts of M polypeptides and few or no L polypeptides. The Dane particles, which represent the complete virus, contain all three HBsAg forms.



**FIG. 145.2** Hepatitis B virus (HBV) genome organization, map of viral transcripts, and proteins. The partially double-stranded 3.2-kilobase viral DNA is shown in the inner circle. The single-stranded (ss) region is indicated in yellow-orange. The extent of the ss region varies from molecule to molecule. HBV-encoded overlapping genes are indicated in the outer circles in various colors. Four promoter regions preceding a corresponding gene are indicated as S1p, S2p, Cp, and Xp. Two enhancer elements (I and II) are also shown. Viral transcripts are indicated in the outermost circles (*thin lines*). The three forms of HBsAg, HBcAg, and HBeAg (surface, core, and e antigen, respectively) polypeptides are also shown. (Modified from Murray PR, Rosenthal KS, Kobayashi GS, Pfaller MA. *Hepatitis viruses*. In: Murray PA, Rosenthal KS, Kobayashi GS, et al, eds. *Medical Microbiology*. 4th ed. St. Louis: Mosby; 2002:591–605.)

The C region, or the core ORF, contains two in-frame initiator codons, with the first adenine-thymidine-guanine (ATG) start codon responsible for e antigen polypeptide synthesis (HBeAg) (see Fig. 145.2). The second ATG serves as an initiation codon for the HBcAg polypeptide. HBeAg is secreted from cells and accumulates in serum as an immunologically distinct soluble antigen and serves as a marker of ongoing viral replication. Both gene products (core and e polypeptides) are made from the same reading frame. The P and X ORFs encode the polymerase and the HBx proteins, respectively.


Before the advent of current molecular methods, HBV was divided into serotypes on the basis of the antibody response to HBsAg. However, HBV is now divided into 10 genotypes (A–J), based on genetic diversity of at least 8% in the HBV genome.<sup>29,30</sup> Within these genotypes are subgenotypes, which differ by a minimum of 4%. These genotypes and subgenotypes are distributed geographically, with genotypes A in North America, Europe, and parts of Africa; genotypes B and C in Asia; genotype D in India, the Middle East, the Mediterranean region, and parts of Africa; genotype E in Africa; genotype F in Central and South America; genotype G in France, Germany, and North America. Genotype H is found in Central America. Genotype I, a novel recombination between genotypes A, C, and G, is found in Vietnam and Laos.<sup>31</sup> Genotype J

was identified in Japan.<sup>32</sup> There are accumulating data that these genotypic differences are associated with disease and treatment outcomes. Genotypes G and C are associated with more severe liver disease and higher HCC risk.<sup>33</sup> Genotype C also has slower HBeAg seroconversion compared with genotype B.<sup>34,35</sup> Genotypes A and B are more responsive to pegylated interferon- $\alpha$  (PEG IFN- $\alpha$ ) therapy than are genotypes C and D.<sup>36</sup>

### Transcription

The HBV genome maximizes the use of its limited genomic size by using multiple overlapping reading frames and multiple initiation codons to generate antigenically different proteins. The cccDNA is a stable episomal form of HBV in the nucleus that is associated with histones and other proteins, serves as the substrate for viral transcription, and uses host polymerase II (pol II) in the synthesis of viral transcripts (see Fig. 145.3). The expression of the four different genes, designated C, S, P, and X, is regulated by four promoter elements (S1p, S2p, Cp, and Xp) and two enhancer elements (enhancer I and II; see Fig. 145.2).<sup>15</sup> These transcriptional regulatory elements direct the synthesis of multiple viral transcripts that are approximately 3.5, 2.4/2.1, and 0.8 kb in length, respectively. All viral transcripts





**FIG. 145.3** Hepatitis B virus life cycle. The virion attaches to the hepatocyte through the Na (sodium) taurocholate cotransporting polypeptide (NTCP) receptor. It is brought into the cytoplasm via clathrin-mediated endocytosis. The surface protein binds to the endosomal membrane, allowing release of the capsid, which is transported to the nuclear pore complex, where the capsid dissociates and the DNA is translocated to the nucleus. In the nucleus the partial double strand is completed, forming a covalently closed circular form called cccDNA. The negative strand of such cccDNA is the template for transcription by cellular RNA polymerase II of a longer-than-genome-length RNA, called the *pregenome*, and shorter subgenomic transcripts. Viral messenger RNAs (mRNAs) are transported from the nucleus. The hepatitis B surface antigens (HBsAg) encoding viral mRNAs are translated by ribosomes bound to the endoplasmic reticulum (ER), and the proteins enter the secretory pathway. The pregenome RNA is translated at low efficiency to produce a 90-kilodalton polymerase protein, *P*, which possesses reverse transcriptase activity. This protein then binds to a specific site at the 5' end of its own transcript, where viral DNA synthesis is eventually initiated. The pregenomic RNA also serves as mRNA for the capsid protein. Concurrently with capsid formation, the RNA-P protein complex is packaged and reverse transcription begins with synthesis of negative-strand DNA, followed by positive-strand DNA synthesis. Before the completion of +/- strand synthesis, core particles mature, and these structures acquire envelopes by budding into the ER, where viral morphogenesis is completed. Progeny-enveloped virions are released from the cell by exocytosis or are recycled to the nucleus, where the process is repeated. (Modified from Flint SJ, Enquist LW, Racaniello VR, et al. *Principles of Virology*. 2nd ed. Washington, DC: American Society for Microbiology Press; 2004:809.)

are unspliced, capped, and polyadenylated. All the transcripts are encoded on one strand of the DNA and coterminate at an identical polyadenylation site.<sup>37</sup>

The pre-S1 (S1p) promoter directs the synthesis of the 2.4-kb transcripts, which code for the large envelope (L HBsAg) polypeptide. This promoter contains binding sites for two liver-enriched transcription factors, hepatocyte nuclear factor 3 (HNF-3) and HNF-1, which are primarily responsible for the liver-specific activity of this promoter.<sup>38</sup> The pre-S2 or S (S2p) promoter directs the transcription of multiple species of messenger RNAs (mRNAs) coding for pre-S2/M and S HBsAg polypeptides. Collectively, these RNA species are approximately 2.1 kb in length with 5' heterogeneous ends. The mRNAs, which initiate downstream of the pre-S2 initiator ATG, code for the S, or the major polypeptide. The S2p promoter also displays liver specificity and is controlled by the enhancer II element located about 2000 base pairs away (see Fig. 145.2).<sup>39</sup> The S2p promoter is stronger than the S1p promoter, which results in the synthesis of an excess of the major S over the L (pre-S1) and M (pre-S2) forms of the HBsAg. This regulation is especially critical for synthesizing the appropriate levels of the three forms of surface proteins within the cell. The basis for the differential

regulation of these promoters is governed by mechanisms that involve both positive and negative *cis*-acting elements and the *trans*-acting transcriptional factors.<sup>15,40</sup>

The core/pregenome promoter (Cp) governs the expression of two longer-than-genome-length transcripts (3.5 kb) designated precore (pre-C) and core (C) RNAs. The slightly longer precore RNA directs the translation of HBeAg polypeptide. The shorter C RNA is used for the translation of core and polymerase proteins. The ATG for the polymerase (reverse transcriptase) is located several hundred nucleotides from the 5' end (see Fig. 145.2). The C RNA, after its translation into core polypeptides, packages its own RNA, which then functions as a pregenome RNA (pregenomic RNA). The Cp promoter contains binding sites for several liver-enriched and ubiquitous transcription factors.<sup>40</sup> Immediately juxtaposed to this lies enhancer II, which, in concert with enhancer I, plays a regulatory role in the overall HBV gene expression. A complex scheme of transcriptional regulation appears to operate within these control elements.<sup>41</sup>

The X promoter (Xp) is located immediately downstream of enhancer I and regulates the synthesis of the low abundance 0.8-kb X transcripts that correlate with similar levels of HBx protein synthesis in infected

cells.<sup>42,43</sup> Both ubiquitous and liver-enriched factors binding to enhancer I regulate the biosynthesis of X transcripts.

## Translation

HBV, which encodes three major viral transcripts, also uses unique translational strategies to maximize the limited coding capacity of its genome. The S region contains three in-frame translational start codons from which the synthesis of three distinct surface antigen polypeptides is regulated (L/pre-S1, M/pre-S2, and S). These proteins exhibit distinct amino termini and differ with respect to the extent that they are post-translationally modified by glycosylation.<sup>15</sup> The S protein is referred to as the major surface antigen because it represents approximately 85% of the HBsAg that is produced by the virus. The pre-S1 (L) and pre-S2 (M) proteins are present at an abundance of approximately 15% and 1% to 2%, respectively.

The pre-C and C mRNAs are translated into e and core polypeptides (HBe/cAg), respectively. The pre-C protein contains a signal peptide sequence encoding 19 amino-acid residues, which targets this protein to the secretory pathway in the endoplasmic reticulum (ER).<sup>44</sup> Further processing of the protein includes cleavage of the signal peptide, and several amino-acid residues at the carboxyl terminus, leading to the production of a 17-kDa HBeAg (see Fig. 145.2). HBeAg is secreted and found in the serum of patients and serves as a marker of active replication in chronic hepatitis. Although the function of HBeAg is not clearly understood, one study demonstrated that it downregulated Toll-like receptor 2 expression on hepatocytes and monocytes, leading to a decrease in cytokine expression.<sup>45</sup> These data suggest a role for HBeAg in modifying the innate immune response to HBV. HBeAg is dispensable for replication, as HBV carriers with pre-C mutant viruses arise spontaneously and are both infectious and pathogenic.<sup>46,47</sup> The C RNA initiates downstream of the pre-C ATG and translates into HBCAg and less frequently into polymerase. Because polymerase ATG is located several hundred nucleotides downstream of the 5' end (see Fig. 145.2), the ribosomes must access this initiator codon by a mechanism different from ribosome scanning. Because this process is inefficient, one polymerase protein is translated for every 200 to 300 molecules of core polypeptides.

HBV polymerase protein consists of 832 amino acids and is composed of four domains: terminal protein (TP), spacer, reverse transcriptase (RT), and ribonuclease H (RNase H).<sup>15</sup> The TP domain represents the portion of the polymerase protein that is covalently bound to negative-strand DNA. The spacer domain or tether region connects the TP domain to the RT, which contains the catalytic domain for the polymerase. The RT domain is followed by an RNase H domain, which functions to hydrolyze the pregenomic RNA template after reverse transcription within the nucleocapsid.<sup>48</sup>

## Replication

Although HBV is a DNA-containing virus, the replication occurs via an RNA intermediate, a characteristic that places hepadnaviruses close to retroviruses. Much of our understanding of HBV replication strategy comes from the classic experiments of Summers and Mason,<sup>49</sup> which demonstrated that HBV amplifies via reverse transcription of an RNA intermediate and that these events occur within the subviral core particles in the cytoplasm. The complex mechanism by which pregenomic RNA is converted to partially double-stranded virion DNA begins with the encapsidation of pregenomic RNA by the core polypeptide along with the polymerase (Fig. 145.4).<sup>50,51</sup> The pregenomic RNA is greater than genome length because it contains terminally redundant 200-nucleotide sequences, which include an epsilon ( $\epsilon$ ) stem-loop structure, and a short sequence of 11 to 12 nucleotides termed DR1 (see Fig. 145.4).<sup>15</sup> However, only the 5' end  $\epsilon$  stem-loop structure contains a binding site for the HBV polymerase to bind to its own molecule. The interaction between the  $\epsilon$  signal of the pregenomic RNA and the polymerase leads to the formation of a three-base DNA priming sequence. This primer sequence then transfers to the DR1 near the 3' end of the pregenomic RNA (pgRNA) to initiate elongation of the negative-strand DNA and concurrent digestion of pgRNA by RNase H of the polymerase. After this, positive-strand synthesis occurs. The positive strand is extended to variable lengths to yield mature HBV DNA. The cessation of the positive-strand synthesis at various stages of its synthesis coincides with

the maturation of the core particles and their entry into the ER, leading to the packaging of partially duplex genomic DNA found in hepatitis B (HB) virions (see Fig. 145.3). This entry prevents access to cytoplasmic deoxynucleotide pools needed for completion of DNA synthesis. Matured core particles face two options at this stage: They either enter the ER, undergo virion packaging with HBsAg polypeptides, and bud from the membrane as 42-nm Dane particles; or they enter the nucleus and deliver the partially duplex DNA and repeat the cycle of replication (see Fig. 145.3). cccDNA serves as a substrate of all viral mRNA synthesis; it is the stable form of HBV DNA that is most resistant to antiviral treatment and the host immune response.<sup>15</sup> In summary, the salient features of HBV replication are cccDNA as the replication template for all viral RNAs, the use of pregenomic RNA as a template for reverse transcription occurring within core particles; incomplete positive-strand synthesis that results in a partially duplex DNA genome found within the hepatitis B virion particles; and mature core particles either enter nucleus or are secreted out of the hepatocyte.<sup>52</sup>

## HBx

HBx is a regulatory protein whose primary role is to enhance transcription of the HBV genome by promoting degradation of a host restriction factor, SMC5/6.<sup>53</sup> It is also involved in several cellular pathways. The cellular targets are numerous, and the list of the cellular functions HBx has been shown to modulate continues to grow.<sup>15</sup> Overall, the functions of HBx are characterized as transcriptional *trans*-activators. Nuclear factor kappa B (NF- $\kappa$ B) was one of the first HBx-responsive elements identified.<sup>54</sup> Because the NF- $\kappa$ B site does not exist within the HBV genome, it is believed that the *trans*-activating potential of HBx extends to cellular target genes involved in inflammation. Other HBx-responsive transcription factors include activating transcription factor 2/cAMP response element binding protein (ATF-2/CREB), activation protein 1 (AP-1), signal transducer and activator of transcription (STAT)-3, and nuclear factor of activated T cells (NF-AT).<sup>55,56</sup> Within the cytoplasm, it also localizes to the mitochondria and induces oxidative stress via calcium signaling.<sup>57,58</sup> These activities may mediate mitochondria-mediated liver injury. HBx has also been shown to modulate cellular signal transduction pathways.<sup>59,60</sup> It was shown to activate sarcoma (Src) kinase and stimulate the Ras-Raf–mitogen-activated protein kinase (MAPK) signal transduction pathway.<sup>61</sup> The physiologic significance of any of these observations remains to be established in the context of human infections. HBx is not known to be directly oncogenic, but on the basis of its role in activation of transcription factors, elevation of reactive oxygen species, and inhibition of apoptosis,<sup>62</sup> it is considered a potential contributor to the processes of liver oncogenesis.<sup>63</sup>

## Morphogenesis and Assembly


Envelopment of the nucleocapsid occurs in the ER or ER-Golgi intermediate compartments.<sup>15</sup> All three forms of the surface proteins are synthesized in the ER as integral transmembrane proteins. M protein is not important for morphogenesis but is required for infectivity. The N-terminal domain of L is critical for virion morphogenesis, serving to translocate the newly-synthesized viral cores across the ER membranes.<sup>64</sup> After budding from the ER the virions are secreted via vesicular transport through the remaining compartments of the secretory pathways and are eventually released into the bloodstream. During chronic hepatitis, accumulation of the L surface protein-containing particles has been associated with the pathologic phenotype of ground-glass hepatocytes.<sup>15</sup>

## NATURAL HISTORY AND PATHOGENESIS OF DISEASE

### Natural History

If HBV is acquired in adulthood, ~95% mount a successful immune response that results in recovery and production of protective antibodies (anti-HBs). In contrast, this only occurs in 5% to 10% of perinatally acquired infection. However, recovery is not synonymous with viral clearance because cccDNA is still present in some hepatocytes, so reactivation can occur (see "Reactivation").

The phases of chronic hepatitis B are in flux, with the newest guidelines delineating phases based on serologic markers. These phases result from the interplay between the virus and the host; however, it is not necessary



**FIG. 145.4** Hepatitis B virus replication pathway. The terminally redundant pregenomic RNA (pgRNA) (top line) is capped and polyadenylated. Boxes 1 and 2 represent DR1 and DR2, respectively. The horizontal bracket labeled *R* represents the terminally redundant region of the RNA. The epsilon (ε) stem-loops are indicated at the termini. pgRNA packaging into cores is initiated by the interaction of polymerase (P) protein with the 5' copy of ε, which leads to formation of 3-base pair priming sequence. This primer sequence translocates to the DR1 of the 3' end of the pgRNA, initiating positive synthesis (step 2). As DNA synthesis continues the pgRNA template is degraded by the associated ribonuclease H activity of the polymerase (step 3). Elongation of negative-strand DNA is finished on complete copying of the pgRNA template (step 4). The product is a terminally redundant complete negative-strand DNA species. The redundancies (8–10 nucleotides) are labeled *r*. At this time the primer for positive-strand synthesis is generated from the 5'-terminal 15 to 18 nucleotides of the pregenomic RNA. The primer is capped (CAP) and includes the sequence to the 3' end of DR1. Elongation of the positive strand results in a duplex linear genome. In the majority of cases the primer is translocated and base pairs with the DR2 sequence near the 5' end of negative-strand DNA. The positive-strand synthesis is initiated, and then elongation begins. On reaching the 5' end of negative-strand DNA, an intramolecular template exchange occurs, resulting in a circular DNA genome (step 5). This exchange is facilitated by the short terminal redundancy in negative-strand DNA. The positive-strand DNA synthesis then continues for a variable distance, resulting in the relaxed circular partially duplex form of the genome found in mature virions. (Modified from Tavis JE. The replication strategy of the hepadnaviruses. *Viral Hepatitis Rev.* 1996;2:205–218.)

to go through all of the phases.<sup>65</sup> The first phase, the HBeAg-positive HBV infection phase (also known as the immunotolerant phase), is characterized by limited immune response against the virus, so there is minimal elevation of serum aminotransferases and minimal liver inflammation despite circulating HBsAg, HBeAg, and high levels of HBV DNA. In cases of perinatal infection, this phase may last for decades, during which there are low rates of spontaneous seroconversion to anti-HBe. The cumulative rate of spontaneous HBeAg clearance during this phase is estimated to be approximately 2% during the first 3 years of life and only 15% 20 years after perinatal infection.<sup>66,67</sup> In contrast,

among immunocompetent adults, this phase is typically present only during the incubation period.

During the second phase of the infection the HBeAg-positive chronic hepatitis B phase (also known as the immune active phase), there is an increase in immune response, accompanied by increased aminotransferases and liver inflammation and high HBV DNA levels. This is thought to be the period in which there is augmentation of both innate and acquired HBV immunity, leading to cytolytic destruction of hepatocytes, as the peak of the immune response coincides with the aminotransferase elevations.<sup>68</sup> The result of this increased immune response is usually



conversion from HBe antigenemia to anti-HBe. In perinatally acquired infection, this period usually occurs in the second to third decade of life, and HBeAg seroconversion rates may approach 10% to 20% per year.<sup>66,69</sup> Individuals with active chronic hepatitis B remain in this phase because the immune reaction is not sufficiently vigorous to result in production of protective antibodies and viral control. Some individuals with chronic hepatitis B can eventually experience HBeAg seroconversion. In a study of 1536 Alaskan natives who acquired HBV as adults and did not receive treatment, two-thirds of subjects cleared HBeAg and developed anti-HBe during 12 years of follow-up without reversion to the HBeAg-positive state.<sup>70</sup> The likelihood of seroconversion is inversely related to the serum aminotransferase. Spontaneous seroconversion occurs in 50% of those with serum aminotransferases greater than five times the upper limit of normal (ULN), compared with <10% in those with lower aminotransferases.<sup>71</sup> Spontaneous seroconversion is marked by the appearance of HBcAg-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells,<sup>72</sup> but the precise factors preceding this event are unknown, and seroconversion cannot be linked to specific viral mutants, as was previously believed.<sup>73</sup> Males and older individuals are also more likely to have spontaneous seroconversion, and viral genotype may play a role.<sup>35</sup> Because immunoglobulin M (IgM) anti-HBc may increase with alanine aminotransferase (ALT) flares during this phase, acute hepatitis B may be erroneously diagnosed.

The third phase, known as the HBeAg-negative HBV infection phase (also known as the inactive phase), is characterized by absence of HBeAg, presence of anti-HBe, normal aminotransferases, and low or undetectable HBV DNA. Termination of virus replication is associated in most patients with biochemical and histologic regression of inflammatory activity.<sup>74</sup> Some individuals will also clear HBsAg, although this is unusual, occurring in <1% of patients per year in adults and 0.05% to 0.8% in infection acquired in infancy or childhood.<sup>70</sup> Individuals who remain in this phase have excellent prognosis and minimal liver injury. During this phase, seroreversion to HBeAg positivity and return to the second phase occurs 10% to 40% of the time.

The fourth phase is the HBeAg-negative chronic hepatitis B phase, where HBeAg remains negative but viral replication increases, resulting in high levels of HBV DNA and elevations of ALT. During this phase the HBV DNA levels are moderately high but generally lower than in those with HBeAg-positive chronic hepatitis B.

## Pathogenesis

It has been difficult to understand the mechanism for viral persistence and immune-mediated liver injury because research in this area is hampered by the lack of a robust culture system or a small animal model. Most studies suggest that HBV is not directly cytopathic to the hepatocyte,<sup>76,77</sup> which is also supported by the existence of asymptomatic hepatitis B carriers with normal liver histology and function. Both human and animal studies have now shown that the liver injury mediated by HBV is initiated by a virus-specific cellular and humoral immune response. In >95% of immunocompetent adults the immune response is vigorous, polyclonal, and multispecific and results in acute, self-limited hepatitis with reduction of viral load and the development of long-lasting humoral and cellular immunity. Persistent infection is associated with necroinflammatory activity, which eventually leads to cirrhosis.

## Acute Hepatitis B

Natural recovery from acute hepatitis B probably depends on multiple components of cellular immune responses, including natural killer (NK) cells, natural killer T (NK-T) cells, and virus-specific CD4<sup>+</sup> T cells and CD8<sup>+</sup> cytotoxic T lymphocytes (CTL). However, a hallmark of acute hepatitis B is failure to induce an innate cytokine response mediated by pattern recognition receptors.<sup>78,79</sup> In self-limited infections the noncytopathic response leads to lowering of the HBV DNA by >90% within a few weeks of peak HBV replication and before liver damage occurs.<sup>80</sup> Both NK and NK-T cells also contribute to clearance through production of IFN- $\alpha/\beta$ .<sup>81</sup> Acute HBV infection is also accompanied by a strong and transient expansion of CD4<sup>+</sup> T cells directed against multiple epitopes within the HBV. HBc is the dominant antigen recognized by CD4<sup>+</sup> T cells in most cases of acute, resolving HBV infection.<sup>82</sup> These HBc-specific CD4<sup>+</sup> T cells provide help for the production of antibody

to HBsAg and are also associated with the development of a vigorous and polyclonal cytotoxic CTL capable of recognizing different epitopes within the HBV genome.<sup>83</sup> Individuals who successfully resolve an HBV infection, either spontaneously or after IFN therapy, maintain these broad and strong peripheral CTL responses.<sup>84</sup> CD4<sup>+</sup> and CTL memory in the presence of low levels of persisting HBV DNA has been shown to persist up to 23 years after infection despite markers of serologic recovery. Both virus-specific CD4<sup>+</sup> and CTL contribute to control of HBV replication through both direct cytolysis of infected cells but, more important, through production of cytokines that control viral replication.<sup>85</sup> Analysis of the NK, CTL, and CD4<sup>+</sup> T-cell responses in the incubation phase has demonstrated that CTL response increases in parallel with ALT, consistent with the previously observed notions that CTL activity is responsible for liver injury, and peaks about the time that the HBV DNA titers begin to fall.<sup>86</sup> Recently, regulatory T cells were found to influence the immune response during acute HBV infection by decreasing cytokine production of effector T cells, thereby down-regulating the antiviral activity. This limits the liver injury during acute infection but, as a consequence, prolongs viral clearance.<sup>87</sup>

The development of surface antibody (anti-HBs) follows the disappearance of HBsAg and marks recovery from acute hepatitis B, which occurs in 95% of adults after acute infection. Anti-HBs is sufficient for protection against HBV infection, as demonstrated by the success of the current HBV vaccines, even if it is not the sole operative mechanism clearing acute infection. The “a” determinant is the predominant B-cell epitope common to all HBV subtypes. Antibodies against this epitope confer immunity to all HBV subtypes. Coexistence of HBsAg and anti-HBs is reported in up to 24% of chronically infected individuals, in which case the anti-HBs is directed against one of the subtypic determinants, and a mutation preventing neutralization of the virus has occurred. Other surface antigens that stimulate antibody responses include the pre-S1 and pre-S2 antigens. Antibody to these develops during recovery and can be detected before anti-HBs; however, routine serologic assays are not readily available. Mutation in the S gene usually occurs in the “a” determinant, which allows the virus to escape antibody neutralization. This may occur after the use of hepatitis B immune globulin (HBIG) for passive immunization of newborns and transplant recipients.

## Chronic Hepatitis B

Persistent HBV infection may result because of the failure of initial innate and adaptive immune responses. In animal models HBeAg may be tolerogenic,<sup>88</sup> and because HBeAg and HBcAg are cross-reactive at the T-cell level, deletion of the CD4<sup>+</sup> HBc-specific T-cell responses results in ineffective CTL responses to HBcAg.

Adult immunocompetent individuals who fail to resolve acute HBV infection also have less vigorous CD4<sup>+</sup> T-cell and CTL responses. In contrast to acute resolving infection, peripheral CD4<sup>+</sup> T-cell and CTL responses in those individuals who develop chronic infection are weak and more narrowly focused.<sup>89</sup> Despite these weak CTL responses, HBV-specific CD4<sup>+</sup> and CD8<sup>+</sup> T cells can be isolated from the livers of chronic hepatitis B patients that are capable of *ex vivo*, class-I-restricted cytolytic activity in response to envelope and core peptides. The cytotoxicity mediated by these cells appears to be sufficiently strong to cause liver injury but not strong enough to eradicate virus and cccDNA from all hepatocytes.<sup>90,91</sup> In addition to the virus-specific cells, the inflamed liver contains other cells that may participate in hepatocyte damage.<sup>92</sup> CD8<sup>+</sup> T cells from patients with chronic hepatitis B exhibit multiple coinhibitory molecules, including programmed cell death protein 1 (PD-1) and cytotoxic T-lymphocyte antigen 4 (CTLA-4). Upregulation of these exhaustion markers may occur in the liver microenvironment, where high HBV antigen concentrations exist, because one study found that HBV-specific CD8<sup>+</sup> T cells in the liver express higher levels of PD-1 compared with circulating T cells.<sup>93</sup>

The mechanisms by which HBV evades the immune response and results in chronic infection in these adults remain obscure. In part this is determined by host genetic factors because persons with certain genetic polymorphisms and human leukocyte antigen (HLA) alleles appear to be more susceptible to chronic infection,<sup>94-97</sup> and certain HLA-DP alleles have been associated with recovery from acute HBV infection.<sup>98</sup> It has

also been suggested that the size of initial viral inoculum and viral kinetics may be such that the immune system is overwhelmed by the virus and becomes “exhausted.”<sup>99</sup>

## Hepatocellular Carcinoma

Although epidemiologic evidence supports the role of HBV as a causal agent of liver cancer (see “Epidemiology of Hepatitis B” later),<sup>100</sup> the molecular mechanisms addressing the link between the viral infection and the development of HCC have been highly debated.<sup>101</sup> Despite a large body of work on this subject, a clear view of how HBV infection triggers events that lead to liver oncogenesis remains elusive. HCC development, like other cancers, proceeds in multiple steps that correlate with specific lesions associated with livers of patients with HCC. These include altered hepatic foci, dysplastic (neoplastic) nodules, and low- and high-grade HCCs. These lesions are characterized as exhibiting different levels of cell differentiation. Similar to other well-characterized human cancers, HCC progresses through these individual stages. The molecular switches associated with each step of HCC development still need to be identified.

HBV-related HCCs are derived from the clonal expansion of a single transformed cell or cancerous cells.<sup>100</sup> Whereas 80% of HBV-associated HCC tumors contain integrated viral DNA, most of the HBV genes are either truncated or transcriptionally inactivated.<sup>15</sup> After integration into the host chromosomes, the HBV DNA control elements, such as enhancers or promoters, can act in *cis* to activate a cellular oncogene. HBV integration was initially thought to be a random event, but recent next-generation sequencing has identified recurrent sites for integration alteration of the *TERT*, *MLL4*, *CCNE1*, *NTRK2*, *IRAK2*, and *p42MAPK1* genes, all of which are important for cell growth and differentiation.<sup>102</sup> It is thought that the random but multiple integration events might frequently lead to genomic instability by causing chromosomal aberrations, such as amplifications, translocations, and deletions, all of which have the potential for activating pathways that lead to liver neoplasia. Indeed, in HBV-associated liver tumors, the HBV DNA integrants are highly rearranged with deletions, inversions, and sequence reiterations.<sup>103</sup> HBV DNA insertion into cellular genes encoding for proteins that are necessary for control of cell signaling and proliferation has been demonstrated.<sup>104</sup>

The pleiotropic functions of HBx in HCC have been implicated in the processes of liver oncogenesis.<sup>105</sup> HBx modulates the expression and activities of numerous genes, as well as epigenetic molecules (e.g., microRNAs) and events (e.g., methylation and acetylation), leading to deregulation of various pathways and functions. Other functions include its effects on DNA repair processes, antiapoptotic properties, and potential to cause direct oxidative damage to host DNA by elevating the intracellular levels of reactive oxidant species in cells.<sup>15,106</sup> HBx activates a number of nuclear transcription pathways and can modulate cytoplasmic signal transduction pathways, including Ras-Raf-MAPK, Janus kinase/STAT (Jun kinase [JNK]/STAT), resulting in hepatic cell proliferation. In addition, HBx can directly inactivate or indirectly downregulate various tumor suppressors, such as p53, or senescence-related factors.<sup>107</sup> Because most of the HBV genes are extinguished in advanced tumors, the viral role in the development of HCC is most likely to be at the stage(s) of initiation or promotion of hepatocarcinogenesis. This would preclude the presence of HBx in maintaining the transformed phenotype.

Finally, HBV-associated HCC may be due to the repeated cellular division associated with the inflammatory response.<sup>15,108</sup> HCC associated with HBV is most common in the context of cirrhosis, although it can occur in all stages of liver disease. Cirrhosis is the result of years of inflammation and associated repair processes, during which there is considerable cell killing and repeated hepatocyte regeneration. In all types of liver damage, there is evidence of enhanced production of free radicals or significant decrease of antioxidant defense, or both.<sup>109</sup> Oxidative stress induced by inflammation and accumulation of viral surface proteins over a long period may give rise to high mutation rates in infected hepatocytes.

## EPIDEMIOLOGY OF HEPATITIS B


Approximately 3.6% of the world's population (~257 million people) is chronically infected with HBV, and more than 800,000 deaths due

to HBV occur annually.<sup>110,111</sup> The prevalence of chronic HBV varies widely, ranging from 0.01% to 2% in the low-prevalence areas (United Kingdom, United States, Canada, Western Europe, and Japan), 2% to 5% in the lower intermediate areas (parts of South America, North Africa, Russia), 5% to 8% in the higher intermediate areas (South Africa, China, and Thailand), and >8% in areas of high prevalence (most of sub-Saharan Africa and a number of countries in the Western Pacific region; Fig. 145.5 and Table 145.1). The largest number of people living with chronic HBV live in the Western Pacific region (>95 million) and the African region (>75 million).<sup>110</sup> In some parts of Europe, typically considered a low-prevalence area, high HBV prevalence rates have been observed in persons migrating from countries with intermediate/high HBV prevalence.<sup>112</sup> Specific patient groups, such as injection drug users, men who have sex with men (MSM), and human immunodeficiency virus (HIV)-infected patients have been shown to have significantly higher rates of chronic HBV infection than the general population.<sup>113</sup> The risk of chronic infection is inversely proportional to age of acquisition of infection. In areas of high seroprevalence, HBV is more likely to be acquired perinatally, with an attendant high risk of chronic infection of 90%. For infections acquired between the ages of 6 months and 5 years, the risk of chronic infection is between 20% and 60%, and for infections acquired by immunocompetent adults the risk is approximately 5%.<sup>114</sup>

The epidemiology of HBV is changing with the advent of universal vaccination programs adopted by many countries. Over the past 40 to 50 years decreases in HBV prevalence in countries of South-East Asia, the Western Pacific, and the Eastern Mediterranean have been observed, whereas increases have been observed in parts of Eastern Europe and Africa (Nigeria) (8.55%–9.8%). In the United States the estimated number of new symptomatic infections per year was more than 10 per 100,000 in the mid-1980s, whereas after the adoption of universal vaccination for infants and “catch-up” vaccination for older children in 1991, this number was estimated to be about 1.1 per 100,000 in 2015.<sup>115</sup> The overall incidence of acute hepatitis B has declined in all age groups over this period, although it remains highest in persons age 30 to 39 years (2.6 cases/100,000 population) (Fig. 145.6).<sup>115</sup> Despite these declines, there are still between an estimated 850,000 and 2.2 million HBV carriers in the United States, of whom about 25% will develop serious sequelae during their lifetime, and 1800 will die annually due to complications of chronic liver disease, including HCC.<sup>115,116</sup>

## ROUTES OF TRANSMISSION

Hepatitis B replicates to high titer in the blood ( $10^8$ – $10^{10}$  virions/mL), especially during the acute phase of illness. Any parenteral or mucosal exposure to infected blood thus represents a potential risk for acquisition of hepatitis B and accounts for the 100 times more efficient transmission of HBV compared with HIV after needlestick exposure.<sup>117</sup> HBV is also found in other body fluids to a variable degree, including semen, saliva, cervical secretions, and tears, and can survive up to 7 days on environmental surfaces.<sup>117a</sup> Thus exposure to even minute amounts of blood or contaminated secretions may transmit virus, and infection can occur in settings of prolonged, close personal contact, such as occurs between children, or among residents of institutions for the developmentally disabled, probably due to inapparent contact of infected secretions with nonintact skin. HBV is not found in urine, sweat, or stools. The typical mode of transmission of HBV varies in part with the prevalence of infection (see Table 145.1). In high- and intermediate-endemicity areas, such as sub-Saharan Africa and South East Asia, the predominant route of transmission is perinatal or horizontal during childhood (i.e., close contact excluding parenteral, sexual, or perinatal exposure), whereas in more low-prevalence areas, transmission occurs via injection drug use and high-risk sexual behaviors. Persons at increased risk of acquiring HBV infection include members of the following groups: illicit drug users (injecting, inhaling, snorting, pill popping), heterosexual men and women and MSM with multiple partners, household contacts and sexual partners of HBV carriers, infants born to HBV-infected mothers, patients and staff in custodial institutions for the developmentally disabled, recipients of certain plasma-derived products (including patients with congenital coagulation defects) before 1992, hemodialysis patients, health and public safety workers who have contact with blood, travelers



**FIG. 145.5** Global prevalence of hepatitis B surface antigenemia. (From Schweitzer A, Horn J, Mikolajczyk RT, et al. Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *Lancet*. 2015;386:1546–1555.)

**TABLE 145.1** Global Seroprevalence Rates and Modes of Transmission of Hepatitis B

| CHARACTERISTIC       | HIGH   | INTERMEDIATE  | LOW   |
|----------------------|--|---|---|
| Carrier rate (%)     | $\geq 8$   | 2–7.99  | $< 2$   |
| Distribution         | Alaskan Eskimos, sub-Saharan Africa, Somalia, Yemen, Djibouti, Haiti, Kyrgyzstan, Papua New Guinea, Solomon Islands, Nauru, Vietnam, | Eastern and southern Europe except Kyrgyzstan, Eastern Mediterranean, Western Pacific region except Australia, Japan, Papua New Guinea, Vietnam, Southeast and Central Asia, except Nepal and India | United States, Canada, Western Europe, Australia, Mexico, South and Central America except Peru, Suriname, Columbia and Belize; Nepal, India; Japan |
| Age at infection     | Perinatal and early childhood  | Childhood   | Adult   |
| Mode of transmission | Maternal and perinatal   | Percutaneous, horizontal  | Sexual, percutaneous  |

to regions with intermediate or high rates of HBV ( $\geq 2\%$ ), and persons born in areas of high HBV endemicity and their children. Persons at risk of exposure to HBV should be screened for markers of HBV infection and receive hepatitis B vaccine if seronegative (see “[Prevention of Hepatitis B Virus Infection](#)” later).


### CLINICAL MANIFESTATIONS AND PROGNOSIS

The spectrum of clinical manifestations of HBV infection varies in both acute and chronic infection. During the acute phase, manifestations range from subclinical or anicteric hepatitis to icteric hepatitis and, in some cases, fulminant hepatitis. During the chronic phase, manifestations range from an asymptomatic carrier state to the signs and symptoms of cirrhosis and HCC. Extrahepatic manifestations can also occur with both acute and chronic infection.


### Acute Hepatitis B

After exposure to HBV, there is an incubation period of 1 to 4 months.<sup>118</sup> Acute hepatitis B is a clinical syndrome indistinguishable from other acute hepatitis and often consists of a flulike syndrome with malaise, fatigue, anorexia, nausea, vomiting, and right upper quadrant discomfort. Serum sickness–like manifestations may be present before the onset of jaundice. Physical signs include jaundice (present in almost all patients) and tender hepatomegaly. The likelihood of developing icteric illness is inversely proportional to age. Symptomatic hepatitis rarely develops in children younger than 1 year, in 10% of children younger than 5 years, and between 30% and 80% of adults.<sup>114</sup> Most reported cases of HBV are the result of icteric illness, but it is believed that approximately two-thirds of cases either result in subclinical illness or are only diagnosed in retrospect. The acute illness may be more severe in the setting of other coinfections, such as simultaneous acquisition of hepatitis D (see





**FIG. 145.6** Incidence of symptomatic, acute hepatitis B per 100,000 population by age group: United States, 2000–2015. (Modified from Centers for Disease Control and Prevention. Surveillance summaries: surveillance for viral hepatitis—United States, 2015. <https://www.cdc.gov/hepatitis/statistics/2015surveillance/pdfs/2015HepSurveillanceRpt.pdf>. Accessed December 21, 2017.)



**FIG. 145.7** Typical course of hepatitis B. *Left*, Typical course of acute hepatitis B. *Right*, Chronic hepatitis B. *HBe*, Hepatitis B envelope; *HBsAg*, hepatitis B surface antigen; *IgM*, immunoglobulin M.

Chapter 146), or with underlying liver disease, such as alcoholic liver disease. The symptoms and jaundice generally disappear in 1 month, but some patients have prolonged fatigue even after resolution of the elevated serum aminotransferases. See Fig. 145.7 for an illustration of a typical clinical and laboratory course of acute hepatitis B compared with that of chronic hepatitis B. In acute HBV, elevations in the concentration of ALT and aspartate aminotransferase (AST) levels with values up to 1000 to 2000 U/L can be seen, with ALT being higher than AST. The serum bilirubin concentration may be normal in patients with anicteric hepatitis. The prothrombin time (PT) is the best indicator of prognosis, with a high PT indicative of fulminant liver failure.<sup>119</sup> Among patients who recover, normalization of serum aminotransferases usually occurs within 1 to 4 months. Persistent elevation of serum ALT for more than 6 months indicates progression to chronic hepatitis. Rates of progression to chronic disease are similar whether or not there is symptomatic acute disease.

Acute HBV is characterized by the development of IgM antibody against the core antigen, followed by the development of immunoglobulin G (IgG) anti-HBc (see “Screening and Diagnosis of HBV Infection” later). Markers of viral replication, such as HBsAg and HBV DNA in serum, are present at the same time as anti-HBc. Loss of HBV DNA, HBeAg, HBsAg, and IgM anti-HBc and the development of anti-HBe and anti-HBs all characterize immunity. In a recent longitudinal study

of Chinese patients with acute HBV, the median HBsAg clearance time from the onset of symptoms was 6 weeks, and 90% of the patients achieved HBsAg seroconversion in 19 weeks.<sup>120</sup> Patients who recover from acute hepatitis B are usually not cured of infection because a significant number of patients will have HBV DNA detectable by polymerase chain reaction (PCR) many years after clinical recovery.<sup>121</sup> They generally have lifelong protection from disease unless there is significant immunosuppression with loss of protective immune responses, such as in the setting of HIV, chemotherapy, or bone marrow transplantation.<sup>122,123</sup>

### Fulminant Hepatitis

Fulminant hepatitis B is rare, occurring in about 1% of patients, and causes <10% of fulminant liver failure in the United States.<sup>124,125</sup> Patients typically present with rapidly progressive acute hepatitis <28 days from the time of symptom onset, accompanied by signs of liver failure, such as coagulopathy, encephalopathy, and cerebral edema. Mortality rates from fulminant HBV as high as 70% have been reported, although the rates have decreased substantially with advances in liver transplantation and critical care management.<sup>126–128</sup> Poor prognostic factors for transplant-free survival include a lower mean arterial pressure on admission and low platelet count.<sup>124</sup> Laboratory testing may not reveal HBsAg due to early clearance but will have IgM anti-HBc and a positive HBV DNA.

Recently, there have been increasing reports of fulminant HBV in the setting of HBV reactivation with chemotherapy or immunosuppressants, with an associated high mortality (5%–22%).<sup>129–131</sup> Fulminant hepatitis is also more common in the setting of coinfection with hepatitis D virus (HDV).<sup>132</sup>

The pathogenesis of fulminant hepatic failure is not clear but may be related to massive immune responses against the virus. Staining of the liver in fulminant HBV often reveals few copies of HBV, suggesting that it is the exuberant immune response that is primarily responsible for the pathogenesis.<sup>133</sup> In addition to host factors relating to immune status, several variants of HBV have also been implicated in several outbreaks of fulminant HBV, including pre-S and S gene variations; BCP; and precore and core gene variations, which result in the decrease or absence of HBeAg; *pol* gene; and X gene variations.<sup>134</sup> Particular HBV genotypes, such as B1 and F (in the context of HDV infection), have also been shown to be associated with fulminant HBV.<sup>135</sup>

### Chronic Hepatitis B

Chronic hepatitis B is defined by at least 6 months of persistent HBsAg. Many patients with chronic hepatitis B are not diagnosed as a result of follow-up after a case of icteric illness or due to specific symptoms, but rather as a result of incidental elevations in serum aminotransferases or due to membership in a specific risk category. Symptoms, if present, can be as nonspecific as fatigue, unless cirrhosis or HCC is present. Other less common symptoms include nausea, right upper quadrant tenderness, anorexia, myalgias, and arthralgias. Symptoms often do not correlate with severity of disease, levels of serum aminotransferases, or hepatic injury on liver biopsy. The physical examination may be normal, or there may be an enlarged liver. The presence of jaundice, splenomegaly, ascites, encephalopathy, or pedal edema suggests cirrhosis.

In patients with chronic hepatitis B, serum aminotransferases may be normal, although most patients with HBeAg-positive or HBeAg-negative chronic hepatitis B have at least mild-to-moderate elevations. Patients will have markers of viral replication, including HBsAg and often HBeAg or HBV DNA as well (see “[Screening and Diagnosis of HBV Infection](#)” later). During flares of disease activity or just before seroconversion to anti-HBe status, there may be marked elevations in the serum aminotransferases to more than 20 times normal. Flares of disease activity may be due to changes in the level of baseline HBV replication, followed by an immune response against the virus. In some patients this results in repopulation of the viral species with a new variant.<sup>136</sup> Patients with flares of disease activity should be followed for the development of anti-HBe, which signals control of viral replication. Cirrhosis should be suspected if there is evidence of hypersplenism, manifested as decreased platelet count, or impaired hepatic synthetic function, indicated by hypoalbuminemia, hyperbilirubinemia, or decreased albumin. Similar to the range of findings observed in other laboratory tests, findings on liver biopsy range from minimal inflammation to cirrhosis. About 2% to 5% of patients with HBV-related compensated cirrhosis develop decompensated disease each year. Decompensation can occur over time or can result from hepatitis flares.<sup>137</sup> Clinical signs and symptoms of decompensated disease include ascites, jaundice, hepatic encephalopathy, or variceal bleeding.

There are no characteristic pathologic findings that can distinguish hepatitis B from other forms of viral hepatitis, although liver biopsies can be stained for the presence of HBsAg and HBeAg. The most characteristic feature of chronic hepatitis B is the ground-glass hepatocyte, which is thought to be due to accumulation of HBsAg within the ER.<sup>138</sup> Differentiation from alcoholic liver disease can be made by characteristic patterns of steatosis, Mallory bodies, and micronodular cirrhosis, whereas chronic hepatitis C is marked by steatosis and characteristic lymphoid follicles within portal tracts. Immunosuppressed hosts may have a variant of HBV known as fibrosing cholestatic hepatitis, which reveals periportal fibrosis, hepatocyte ballooning, bile stasis, and mild or absent inflammation.<sup>139</sup>

### Precore or HBe-Negative Mutants

Early reports described patients in whom HBeAg was absent, although HBV DNA continued to be present at a high level, typically in association

with severe liver disease.<sup>140–143</sup> This is due to the development of HBV pre-C or core promoter mutants that cannot produce HBeAg (see “[Translation](#)” earlier).<sup>144</sup> The most frequent pre-C mutation is a guanine (G) to adenine (A) change at nucleotide 1986, which produces a translation stop codon at amino-acid position 28 in the pre-C region and abolishes production of HBeAg.<sup>145</sup> A1762T and G1764A are frequently occurring mutations that take place in the basal core promoter (BCP) region. All these mutations abrogate HBeAg synthesis without affecting the replication of the virus, which persists and results in chronic inflammation. Several large cohort and case-control studies have demonstrated an association between BCP A1762T/G1764A mutations and an increased risk of liver disease progression and HCC.<sup>146</sup> At the time of initial presentation, 29% to 38% of HBeAg-negative chronic hepatitis B patients have cirrhosis.<sup>147,148</sup> These patients have very low rates of spontaneous seroconversion of <0.5% per year<sup>149</sup> and are more difficult to treat.<sup>141</sup> Originally, HBeAg-negative hepatitis was thought to be predominantly a geographic phenomenon because it is more prevalent in certain parts of the world, especially Asia and the Mediterranean. However, studies suggest that the likelihood of HBeAg-negative chronic hepatitis B is related to duration of infection, as suggested by the older age at presentation,<sup>147</sup> which would make it appear to be of higher prevalence in areas where perinatal transmission predominates. HBeAg-negative disease is also more frequent with genotypes C, D, and F than with genotype A and B infection and presumably contribute to genotype specific immunopathogenesis.<sup>135,150</sup>

### Prognosis of Untreated Chronic Hepatitis B

The likelihood of morbidity and mortality in chronic hepatitis B is directly related to the development of cirrhosis. Among chronic HBsAg-positive patients, the risk of developing cirrhosis ranges from 1 to 5.4 per 100 person-years, with a 5-year cumulative probability of progression ranging from 8% to 20%.<sup>147,151,152</sup> For persons who clear HBsAg, the prognosis is good, although, perhaps surprisingly, not entirely benign. In one study of 189 patients who were noncirrhotic at the time of HBsAg clearance, 3 (1.6%) developed cirrhosis, 2 (1.1%) developed HCC, and 1 died of HCC. These complications all developed in patients with concurrent hepatitis C virus (HCV) or HDV infection, however.<sup>153</sup> In the absence of cirrhosis the long-term prognosis even of HBsAg-positive patients is good. In a 16-year follow-up study of 317 HBsAg-positive blood donors from Montreal, for example, only 3 died from HBV-related cirrhosis and none developed HCC.<sup>154</sup> In a US-based study of 6689 patients with a chronic hepatitis B who did not receive therapy and who were followed for 10 years, 3.1% died from HBV-related causes, with the highest percentage (15.5%) in those older than 65 years.<sup>155</sup> However, in another study of HBsAg-positive patients in England and Wales, during a mean of 22 years of follow-up, 17.4% of deaths were due to HCC or liver disease.<sup>156</sup>

Multiple variables account for the wide estimates of risk. A study from Taiwan (REVEAL [Risk Evaluation of Viral Load Elevation and Associated Liver Disease/Cancer]–HBV) of 3582 people with chronic hepatitis B followed for 11 years found that the strongest predictor of progression to cirrhosis was the HBV DNA level, which was independent of the HBeAg status, and inflammation as represented by ALT values.<sup>157</sup> Individuals with >10<sup>6</sup> copies/mL of HBV DNA had a 36.2% cumulative incidence of cirrhosis compared with 4.5% in those with <300 copies/mL. Most of the people in this study acquired HBV in infancy or early childhood; thus this study may not be applicable to adult-acquired HBV. The rate of progression to cirrhosis may be higher in patients who are HBeAg negative compared with those who are HBeAg positive,<sup>141</sup> although these findings may be confounded by a longer duration of disease in HBeAg-negative patients and presence of precore and BCP mutations.<sup>158,159</sup> Patients with more severe inflammation and fibrosis at the time of presentation are also more likely to progress to cirrhosis. Fattovich and colleagues<sup>71</sup> found that 30% of patients with moderate, chronic active hepatitis developed cirrhosis on histology after 6 years, whereas 50% of patients with severe, chronic active hepatitis (bridging necrosis) developed cirrhosis on histology after 4 years. Consumption of alcohol also increases both the risk of cirrhosis and HCC.<sup>160</sup> Infection with HCV, when accompanied by active replication of

both viruses, also increases the rate of progression to cirrhosis,<sup>161,162</sup> as does HDV.<sup>163</sup>

For patients with compensated cirrhosis, survival is 84% at 5 years and 68% at 10 years without treatment.<sup>164</sup> Once cirrhosis develops the risk of decompensation is 20% to 25% per year without treatment.<sup>71,165</sup> High HBV DNA levels and HBeAg seropositivity and persistence have been found to be independently associated with decompensated disease and death.<sup>166–168</sup> The prognosis is poor after the development of decompensation, with estimated survival rates of only 55% to 70% at 1 year and 14% to 35% at 5 years.<sup>151,169</sup>

## Hepatocellular Carcinoma

Another major cause of mortality in chronic hepatitis B is HCC, which has a poor prognosis unless caught in a small, surgically resectable stage. HCC is the sixth most common cancer and the fourth leading cause of cancer-related deaths worldwide. An estimated 800,000 new cases occur each year, with chronic hepatitis B accounting for 18% (Europe) to 65% (China) of total cases.<sup>170</sup> The lifetime risk for untreated HBV chronic carriers is currently estimated to be about 20% to 40% and about 10- to 25-fold higher compared with non-HBV-infected controls.<sup>171,172</sup> In a recent systematic review of 66 studies of HCC in HBV-infected persons from Europe and North America, summary estimates of incidence rates according to liver disease status (95% confidence intervals [CIs]) were 0.05 (0.03 to 0.08), 0.31 (0.22 to 0.41), 0.42 (0.27 to 0.56), and 2.97 (2.35 to 3.59) in inactive carriers, asymptomatic carriers, subjects with chronic hepatitis, and with compensated cirrhosis, respectively.<sup>173</sup> Studies suggest that the interval between acquisition of HBV and development of HCC spans several decades (average, 30 years).<sup>174</sup> However, that interval is much shorter in parts of sub-Saharan Africa, where 43% of HBV-related HCC has been shown to develop in adults younger than 40 years.<sup>174a</sup> Coinfection with HDV, HCV, or HIV augments the risk of HCC development in HBV infection. More recently, patients with genotype C (Asians) and A1 (Africans),<sup>175</sup> as well as those with the A1762T and G1764A basal core promoter and pre-S gene mutations, have been shown to have a higher risk of HCC.<sup>135,176</sup> Other risk factors include older age, male gender, heavy alcohol consumption, metabolic syndrome, exposure to aflatoxin B, and family history of HCC.<sup>177–180</sup> Levels of HBV DNA are also a risk factor, with levels  $>10^5$  copies/mL ( $\approx 20,000$  IU/mL) being strongly linked to the development of HCC.<sup>177</sup> In a recent study the importance of serum HBV DNA level as a predictor of HCC was even greater in HBeAg-negative patients with normal ALTs and no cirrhosis at baseline.<sup>181</sup> HBsAg levels have also been shown to positively correlate with the development of HCC and better predict HCC in HBeAg-negative patients with low HBV DNA. In the Taiwanese Elucidation of Risk Factors for Disease Control or Advancement in Taiwanese Hepatitis B Carriers (ERADICATE-B) study, among a subset of 1068 HBeAg-negative individuals with HBV DNA levels  $\geq 2000$  IU/mL, the risk of HCC was significantly increased in these patients with an HBsAg level  $\leq 1000$  IU/mL compared with those with a level  $<1000$  IU/mL (hazard ratio, 5.4; 95% CI, 2.1 to 14.2).<sup>182</sup> Clearance of HBsAg by the age of 50 years appears to decrease the risk of HCC.<sup>183</sup> Finally, treatment with nucleoside analogues decreases the risk of HCC about 30% to 80%.<sup>180</sup> To date, no studies have shown convincing data for an association between occult HBV, defined as the presence of HBV DNA without serologic markers of HBV infection, and the presence of HCC.

## Extrahepatic Manifestations of Hepatitis B

Extrahepatic manifestations can occur in approximately 20% of patients with acute and chronic hepatitis B and are thought to be mediated by circulating immune complexes. Acute hepatitis may be manifested in 10% to 20% of patients as a serum sickness–like illness with fever, skin rash, arthralgias, and polyarthritides, typically occurring just before the onset and subsiding with the development of jaundice. The skin rash can be of virtually any type, including erythematous, macular, maculopapular, urticarial, or petechial. The most common extrahepatic clinical manifestations of chronic hepatitis B include sensorimotor neuropathies, myalgias, arthralgias, Sjögren syndrome, glomerulonephritis, uveitis, and Raynaud syndrome.<sup>184</sup> A number of autoantibodies can also be

seen, including cryoglobulinemia and rheumatoid factor. Polyarteritis nodosa (PAN) is a rare complication of chronic hepatitis B, which usually manifests within the first 6 months of infection. It is a vasculitis of small- to medium-size arteries and typically presents with fever, rash, hypertension, eosinophilia, abdominal pain, renal disease, and polyarthritides. In one review the frequency of HBV infection in PAN cases was 33.7%; however, the overall incidence is declining.<sup>185</sup> Treatment of PAN usually includes the combination of plasma exchanges, anti-HBV therapies, and a short course of steroids in cases of life-threatening organ involvement.<sup>186–188</sup> Recovery from vasculitis can occur in up to 81% when HBeAg seroconversion and decreased viral replication are achieved.<sup>189</sup> HBV-related cryoglobulinemic vasculitis has been successfully treated with nucleoside therapies.<sup>190,191</sup> Glomerular disease also occurs as a manifestation of HBV infection. Nephrotic syndrome secondary to membranous or membranoproliferative glomerulonephritis is a rare complication of HBV infection occurring predominantly in children with active viral replication. The typical presentation is with nephrotic-range proteinuria. Approximately 30% to 60% of children with HBV-related membranous nephropathy undergo spontaneous remission, usually in association with HBeAg to anti-HBe seroconversion.<sup>192</sup> Progression to renal failure can occur, particularly in adults. The prognosis of HBV-related membranous nephropathy is variable. In adults it may lead to progressive renal insufficiency<sup>193</sup>; however, it can also be successfully treated. In a recent meta-analysis of 82 patients, treatment with IFN- $\alpha$  (five studies) or lamivudine (one study) resulted in proteinuria remission in 65.2% of the patients.<sup>194</sup>

## CLINICAL MANIFESTATIONS AND NATURAL HISTORY IN SPECIAL HOSTS

### Individuals With Human Immunodeficiency Virus Infection

Because of common parenteral routes of transmission, HBV and HIV are frequently seen in concert. Since the introduction of antiretroviral therapies (ARTs), the incidence of acute HBV infection in persons living with HIV has been declining. Earlier studies reported the incidence of HBV as high as 12.2 cases per 1000 person-years, although lower in those receiving either ART with lamivudine, ART without lamivudine, or one or more doses of HBV vaccine.<sup>195</sup> In a recent study of 2375 HBV-uninfected MSM enrolled in the Multicenter AIDS Cohort Study, the overall unadjusted HBV incidence rate was 9.6 per 1000 person-years. Incidence rates were significantly lower in the highly active antiretroviral therapy (HAART) era than in the pre-HAART era among both those infected with HIV (incidence rate ratio [IRR], 0.2 [95% CI, 0.1 to 0.4]) and not infected with HIV (IRR, 0.3 [95% CI, 0.2 to 0.4]).<sup>196</sup> Of interest, this study also found that receiving effective ART (HIV RNA  $< 400$  copies/mL) was protective against acute hepatitis B, whereas men receiving ART who had HIV RNA  $\geq 400$  copies/mL were not protected. In fact, the incidence rate of acute hepatitis B in HIV-infected men with effective ART was equivalent to HIV-uninfected men.<sup>196</sup> Current estimates are that between 40% and 60% of individuals who are HIV seropositive in the United States are positive for some marker of past HBV infection, and 4% to 10% are HBsAg positive.<sup>197</sup> In the regions with high HBV endemicity, up to 25% of HIV-infected persons are coinfecting with chronic HBV.<sup>198</sup> Patients with HIV and acquired immunodeficiency syndrome are more likely to have chronic HBV infection, increased replication manifested as higher HBV DNA, and increased likelihood of HBeAg positivity.<sup>199</sup> They are also more likely to have occult HBV; therefore HIV patients should be screened for both HBsAg and anti-HBc. The availability of ARTs with activity against both HIV and HBV has resulted in improved outcomes for patients with HIV/HBV coinfection. However, the risk of liver disease and overall mortality remains higher compared with those with either infection alone, even with effective virologic suppression.<sup>200–202</sup> In fact, as ART has reduced HIV-related deaths, liver-related deaths have now emerged as one of the leading causes of deaths in persons living with HIV. Many of these liver-related deaths are due to HCC, which among HIV/HBV coinfecting persons is approximately fivefold to sixfold higher than in the general population, although less in the era of ART.<sup>203</sup> Other factors likely contribute to this increased risk of HCC, such as obesity, age,



HCV coinfection, and metabolic factors. Lower CD4 counts in HIV-HBV coinfecting persons have also been associated with high HBV DNA levels and an increased risk of HCC and mortality.<sup>204–206</sup> Finally, isolated anti-HBc is found in approximately 16% to 20% of HIV-infected patients.<sup>207</sup> One study demonstrated that the isolated anti-HBc serologic pattern is usually stable, but when changes from isolated anti-HBc occur, the transition is usually to anti-HBc and anti-HBs positivity, suggesting that isolated anti-HBc is due to intermittently low levels of anti-HBs.<sup>207</sup> Rates of occult hepatitis B in those with isolated anti-HBc are variable, ranging from <5% up to 90%.<sup>207–210</sup>

### Hepatitis B After Liver Transplantation

Reinfection of the allograft accounts for the majority of cases of HBV after liver transplantation in patients with chronic HBV. HBV can also be acquired de novo in patients who are HBsAg negative from anti-HBc-positive donors, with rates of infection as high as 77% in the absence of HBIG or antiviral prophylaxis.<sup>211–213</sup> This includes previously vaccinated recipients in whom HBV acquisition can also occur.<sup>214,215</sup> Recipients who are anti-HBs positive and anti-HBc positive are at the least risk of de novo infection occurring in 0% to 4%. Reactivation of HBV has also been reported in recipients who are HBsAg negative and anti-HBc positive pretransplant. Before the use of prophylactic HBIG and nucleos(t)ide therapy at the time of transplantation, graft reinfection rates approached 80% to 100% and was a major cause of allograft dysfunction, cirrhosis, and graft failure.<sup>216</sup> Reinfection was strongly correlated with high levels of HBV DNA, antiviral resistance, and HBeAg seropositivity.<sup>217</sup> The use of HBIG and nucleos(t)ide therapy, considered the standard of care in transplantation programs, has resulted in significant reductions in recurrence of HBV infection in the allograft, and post-transplant survival has improved markedly.<sup>218</sup> In a large study conducted between 1988 and 2002, the 1-, 5-, and 10-year patient survival rates were 91%, 81%, and 73%,<sup>219</sup> respectively, which are comparable with other conditions leading to transplantation. Specific HBV therapies used in liver transplantation are discussed in more detail under “Management in Special Populations” and “Liver Transplantation.”

### Hepatitis B After Other Types of Transplantation

HBV is associated with high morbidity and mortality in renal transplant recipients, which is thought to be due to persistent viral replication and reduced HBsAg seroconversion as a result of the effects of immunosuppression. HBV reactivation has occurred in renal transplant patients with previously resolved HBV infection characterized by negative HBsAg and positive anti-HBc, with or without anti-HBs, but the overall risk is low.<sup>220</sup> Reactivation is higher among those without anti-HBs compared with those with anti-HBs.<sup>221</sup> All renal transplant recipients who are HBsAg positive should receive antiviral therapy with nucleos(t)ide therapy (entecavir or tenofovir alafenamide [TAF] preferred) before or at the time of transplantation and continue indefinitely. Renal transplant recipients with anti-HBc with or without anti-HBs should have HBV DNA measured every 3 to 6 months after renal transplantation and receive nucleos(t)ide therapy if viral reactivation occurs but before onset of clinical hepatitis.<sup>222</sup> The risk of de novo infection in HBV naïve kidney recipients depends on use of prophylaxis and HBV immunity in the recipient and ranges from 0% to 27%. HBV reactivation is low among patients receiving transplants from anti-HBc-positive donors (1%–4%) and may present subclinically.<sup>223</sup> HBV prophylaxis before transplantation protects against HBV acquisition in nonimmune renal transplant recipients whose donors are anti-HBc positive. Recipients of bone marrow transplantations are also at risk of hepatitis B reactivation, which may present as a severe flare at the time of withdrawal of immunosuppression<sup>224</sup> or as progressive chronic liver disease.<sup>225</sup> HBV reactivation has been reported in 10% to 21% of allogeneic hematopoietic stem cell transplantation (HSCT) recipients with resolved HBV infection and without antiviral prophylaxis and 45% in HBsAg-positive auto-HSCT patients.<sup>226,227</sup> Lamivudine in these patients has been shown to reduce the incidence and severity of HBV reactivation.<sup>228,229</sup> Of interest, there have been case reports of hepatitis B cure after bone marrow transplantation due to the transfer of HBV-specific immune cells in the graft.<sup>230,231</sup>

### Coinfection With Hepatitis C

Rates of HCV coinfection among patients with HBV range from 3% to 18%, depending on the geographic area and the epidemiologic risk group.<sup>232</sup> HCV infection is especially frequent in HBV endemic areas, in intravenous (IV) drug users, and among persons infected with HIV.<sup>233</sup> Superinfection and acute infection with HCV can result in suppression of HBV replication and clearance of HBsAg and HBeAg.<sup>75,234–236</sup> In turn, inhibition of HCV replication has been observed in patients with chronic HCV superinfected with HBV and higher rates of HCV RNA clearance have been reported in HBV/HCV coinfecting compared with HCV mono-infected individuals.<sup>237,238</sup> However, when both viruses are replicating, the liver disease is usually more severe than in patients infected by HBV or HCV alone.<sup>239–241</sup> Patients with dual HBV and HCV infection may also have a higher rate of HCC compared with patients infected by either virus alone.<sup>242</sup> Treatment of HCV in HCV-HBV coinfecting individuals has resulted in significant declines in HCC, all-cause mortality, and liver-related mortality.<sup>243</sup> In the past, treatment of HCV with PEG IFN was associated with both relatively high rates of HBsAg seroclearance (≈11%) and HBV reactivation (≈62%).<sup>244,245</sup> HBV reactivation has also been demonstrated in HBV-HCV coinfecting patients during and after HCV treatment, with directly acting antivirals (DAA) in patients who are HBsAg positive with undetectable or low HBV DNA and in patients who are isolated anti-HBc positive.<sup>246,247,248</sup> The risk of reactivation is highest among HBsAg-positive individuals; reactivation among patients with isolated anti-HBc is limited to case reports.<sup>249</sup> Current Infectious Diseases Society of America/American Association for the Study of Liver Disease (AASLD) guidelines recommend all patients initiating HCV DAA therapy should be assessed for HBV coinfection with HBsAg testing and for evidence of prior infection with anti-HBs and anti-HBc. In HBsAg-positive patients HBV therapy should be initiated for patients who meet criteria for treatment (see “Treatment”); for patients whose HBV DNA level is below treatment initiation criteria, either prophylactic antiviral therapy or monitoring of HBV DNA levels during and immediately after DAA therapy with preemptive therapy is recommended.<sup>250</sup>

### SCREENING AND DIAGNOSIS OF HBV INFECTION

Infection with HBV is associated with characteristic patterns of hepatitis B antigens and antibodies. In addition to confirming the stage of HBV infection, proper interpretation of the available tests will aid in the monitoring of patients and selection for antiviral therapy.

#### Acute Hepatitis

The diagnosis of acute hepatitis B is based on the detection of HBsAg and IgM anti-HBc (Table 145.2; see Fig. 145.7). During the replicative phase of infection, HBeAg and HBV DNA are also present. Recovery is accompanied by the disappearance of markers of HBV replication and the appearance of antibodies to these proteins.

HBsAg is the serologic hallmark of HBV infection and is the first marker to appear after acute infection. HBsAg appears in serum 1 to 10 weeks after an acute exposure to HBV, before the onset of symptoms or elevation of serum ALT. In patients who subsequently recover, HBsAg usually becomes undetectable after 4 to 6 months, followed by development of anti-HBs, as early as 8 months after infection. In most immunocompetent patients who recover, anti-HBs persists for life, thereby conferring long-term immunity.

HBeAg is an intracellular antigen that is not detectable in serum. Anti-HBc can be detected throughout the course of HBV infection, and its presence signifies natural infection. During acute infection, anti-HBc is detectable at the time symptoms appear (9–21 weeks after exposure) and is predominantly IgM class. It can be the sole marker of HBV infection during what is called the *window period* between the disappearance of HBsAg and the appearance of anti-HBs. Although IgM anti-HBc is usually an indicator of acute HBV infection, it may remain detectable up to 2 years after the acute infection and can be detectable during HBV flares. IgG anti-HBc persists even in individuals with HBs antigenemia, indicating that the presence of this antibody does not confer protection against viral replication.

**TABLE 145.2 Interpretation of Serologic Tests in Hepatitis B**

| TEST         | ACUTE HEPATITIS B | IMMUNITY THROUGH INFECTION <sup>a</sup> | IMMUNITY THROUGH VACCINATION | CHRONIC HEPATITIS B |
|--------------|-------------------|---|------------------------------|---------------------|
| HBsAg        | +                 | –                                       | –                            | +                   |
| Anti-HBs     | –                 | +                                       | +                            | –                   |
| HBeAg        | +                 | –                                       | –                            | ± <sup>b</sup>      |
| Anti-HBe     | –                 | ±                                       | –                            | ±                   |
| Anti-HBc     | +                 | +                                       | –                            | +                   |
| IgM anti-HBc | +                 | –                                       | –                            | –                   |
| HBV DNA      | +                 | –                                       | –                            | +                   |
| ALT          | Elevated          | Normal                                  | Normal                       | Normal or elevated  |

<sup>a</sup>On occasion, individuals with past infection have isolated anti-HBc only. The presence of an isolated immunoglobulin G anti-HBc may indicate a window period during acute infection or remote prior infection with loss of HBsAg or anti-HBs. In such cases an HBV DNA test may prove useful.

<sup>b</sup>Chronic hepatitis B with a precore mutant is HBeAg<sup>–</sup> and anti-HBe<sup>+</sup>.

ALT, Alanine aminotransferase; anti-HBc, antibody to hepatitis B core antigen; anti-HBe, antibody to hepatitis B e antigen; anti-HBs, antibody to hepatitis B surface antigen; DNA, deoxyribonucleic acid; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; IgM, immunoglobulin M.

## Past Hepatitis B Infection and Vaccination

Anti-HBs is the protective antibody and is present both after recovery from acute infection and successful vaccination. These two can be distinguished by testing for total anti-HBc (Table 145.3). Recovery from an HBV infection is characterized by the presence of both anti-HBs and total anti-HBc. Immunity to HBV after vaccination is indicated by the presence of anti-HBs only.

## Chronic Hepatitis B Infection

The diagnosis of chronic hepatitis B is based on the persistence of HBsAg for more than 6 months. Additional tests for HBV replication—HBeAg and serum HBV DNA—should be performed in those with chronic hepatitis B to further classify the disease and to determine the need for treatment. In adults there are two groups of patients with chronic hepatitis B who do not need treatment but who should be monitored. The first are those who are HBeAg negative with an undetectable HBV DNA and normal serum aminotransferases. These patients are in the inactive phase of chronic hepatitis B (previously known as “inactive carriers”) and should have aminotransferases tested every 6 to 12 months to detect reactivation. The annual rate of reactivation is about 4%, with pre-C mutations, male sex, and age older than 30 years being predictors of reactivation.<sup>251</sup> The second group is patients who are in the immune-tolerant phase of infection (see “Natural History” earlier) with HBeAg and high HBV DNA levels but normal or minimally elevated serum aminotransferases. They should also be monitored every 6 to 12 months with aminotransferase testing.

Those with progressive liver disease who should be considered for treatment can be either HBeAg negative or positive, have detectable HBV DNA, and usually have elevated serum aminotransferase concentrations. The HBV DNA level in HBeAg-positive chronic hepatitis B patients tends to be higher than in those with HBeAg-negative chronic hepatitis B.

In HBeAg-positive patients, development of anti-HBe can occur at any point during chronic hepatitis B infection; it occurs spontaneously in a small percentage of patients per year. It is often accompanied by an increase in serum aminotransferases, followed by a disappearance of HBeAg and HBV DNA from the serum and improvement in liver inflammation. However, some patients continue to have active liver disease and detectable HBV DNA in serum after HBeAg seroconversion.<sup>252</sup>

## Measures of Hepatitis B Replication

PCR-based assays are the standard way to quantify HBV replication. Several real-time PCR assays are available now and have lower limits of sensitivity of approximately 10 IU/mL and upper limits of 4 × 10<sup>9</sup>

**TABLE 145.3 Selection of Patients for Treatment in Chronic Hepatitis B**

| HBeAg | HBV DNA <sup>a</sup> | ALT <sup>b</sup> | TREATMENT STRATEGY   |
|-------|----------------------|------------------|--|
| +     | >20,000 IU/mL        | <2 × ULN         | Observe patient<br>Consider liver biopsy or noninvasive measurement of liver disease if age > 40, ALT > 1 but <2 × ULN, family history HCC, HIV positive; treat if moderate-to-severe inflammation or fibrosis |
| +     | >20,000 IU/mL        | >2 × ULN         | Observe 3–6 mo for spontaneous HBeAg seroconversion before treatment<br>Treatment with PEG IFN-α (48 wk) or NUC (indefinite in most; minimum 6–12 mo after HBeAg seroconversion)                               |
| –     | >20,000 IU/mL        | >2 × ULN         | Treatment with NUC (end point not defined; likely indefinite)<br>PEG IFN-α (48 wk) second line   |
| –     | >2000 IU/mL          | 1 to <2 × ULN    | Consider liver biopsy or noninvasive measurement of liver disease and treatment if moderate-to-severe inflammation or fibrosis   |
| –     | >2000 IU/mL          | <ULN             | Observe; treat if HBV DNA or ALT increase  |
| +/-   | +                    | Cirrhosis        | Compensated: treat with NUC if HBV DNA detectable<br>Decompensated: NUC; consider liver transplantation  |
| +/-   | –                    | Cirrhosis        | Compensated: observe<br>Decompensated: consider liver transplantation  |

<sup>a</sup>Conversion factor to copies/mL = 5.6 (20,000 IU/mL is approximately 10<sup>5</sup> copies/mL).

<sup>b</sup>Also use moderate-to-severe necroinflammation on liver biopsy as guide.

ALT, Alanine aminotransferase; anti-HBe, antibody to hepatitis e antigen; DNA, deoxyribonucleic acid; HBeAg, hepatitis B e antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HIV, human immunodeficiency virus; NUC, nucleoside or nucleotide; PEG IFN-α, pegylated interferon-α; ULN, upper limit of normal.

Modified from Terrault N, Bzowej NH, Chang KM, et al. AASLD Guidelines for the treatment of chronic hepatitis B. Hepatology. 2016;63:261–283.

IU/mL. The major role of HBV DNA assays is to establish the phase of the infection and assess candidacy for treatment. In those who are treated, monitoring HBV DNA is essential for assessing the response. HBV DNA is also useful in several other circumstances. The first is distinguishing the window period of acute infection from chronic infection in those patients who are IgG anti-HBc only. HBV DNA is also useful in cases of fulminant hepatitis, where there may be undetectable levels of HBsAg on presentation. HBV DNA testing is also indicated in patients in whom there may be a false-negative HBsAg test, which occurs due to mutants in the surface protein that are not detected by all HBsAg assays.<sup>253</sup>

The enhanced sensitivity of the newer tests, especially PCR-based methods, has also raised questions about the significance of low-level viremia. Recovery from acute infection was formerly thought to be accompanied by complete clearance of HBV from serum. However, use of PCR-based methods has demonstrated that some individuals may have very low-level replication in the absence of any biochemical or histologic markers of liver injury.<sup>254</sup> The significance of this is not known, although some studies have suggested that such “occult” hepatitis B is associated with progressive liver disease. The threshold level that is associated with a risk of progression of chronic liver disease is not known. An arbitrary value of  $\geq 20,000$  IU/mL or  $10^5$  copies/mL has been suggested as a diagnostic criterion for chronic hepatitis,<sup>254</sup> but lower levels may be seen in patients with ongoing liver injury, especially in HBeAg-negative patients. Because levels of HBV DNA may fluctuate over time in a given individual, the use of HBV DNA testing as a sole determination of future prognosis is not recommended.<sup>255</sup> Recently there has been increasing interest in alternative biomarkers of HBV infection, such as hepatitis B core-related antigen (HBcrAg) or HBV RNA, that may better define the activity of chronic HBV infection, monitor infection, and predict long-term outcomes.<sup>256–258</sup> However, more data are needed on these markers before they can be incorporated in routine clinical care.

### Isolated anti-HBc

The isolated presence of anti-HBc without other HBV serologic markers is not uncommon, being present in 0.4% to 2.3% of blood donors in low-prevalence areas<sup>259,260</sup> and in 10% to 20% of the population in endemic countries.<sup>261</sup> Isolated anti-HBc can be found in patients during the window period of acute hepatitis, many years after resolution of acute hepatitis due to a decline of anti-HBs to undetectable titers, or rarely, after years of chronic infection with decline of HBsAg to titers below the limit of detection. Individuals with isolated anti-HBc should have repeated testing done to exclude false-positive results and IgM anti-HBc to exclude acute hepatitis if that is suspected. If there are unexplained aminotransferase elevations in a patient with isolated anti-HBc, HBV DNA testing is recommended. Isolated anti-HBc is more common in people coinfecting with HIV and in those with both chronic and cleared HCV.<sup>207</sup> The clinical significance of isolated anti-HBc with or without a low-level positive HBV DNA test is not clear; however, such patients should be considered potentially infectious because transmission of HBV from blood and organ donors with anti-HBc alone has been reported.<sup>262,263</sup>

## MANAGEMENT OF HEPATITIS B

### Chronic Hepatitis B

The goals of antiviral therapy in hepatitis B are reduction of the morbidity and mortality due to liver disease. The beneficial effects of antiviral therapy on reducing the incidence of cirrhosis and HCC have been shown for all of the approved anti-HBV medications. In a meta-analysis the risk reduction for cirrhosis with antiviral treatment was 0.39 (95% CI; 0.20 to 0.75) in observational studies and was 0.55 (95% CI; 0.38 to 0.78) in randomized clinical trials.<sup>250</sup> Reductions in death, HCC, and hepatic decompensation were also demonstrated.

Because reduction in morbidity and mortality is challenging to study, a variety of end points have been used as surrogates to define response to treatment in clinical trials, including normalization of serum aminotransferases (biochemical response), improvement in liver histology (histologic responses), achieving an undetectable HBV DNA in the serum (virologic response), and loss of HBeAg with or without anti-HBe

(serologic response).<sup>250</sup> Elimination of HBsAg is rare but important because the risk of HCC continues to be elevated above baseline among patients with persistent HBs antigenemia.<sup>183</sup> The rate of HBsAg loss with IFN or tenofovir disoproxil fumarate (TDF) alone after 48 weeks of treatment is in the range of 3% to 4%.<sup>36,264,265</sup> Because long-term studies have demonstrated that either seroconversion to anti-HBe or durable suppression of HBV DNA is associated with reduction in cirrhosis,<sup>250,266–268</sup> an improvement in survival,<sup>269–272</sup> and reduction of the rates of HCC,<sup>273</sup> these are the most common surrogate markers used in both clinical practice and clinical trials to monitor the efficacy of therapy. Quantitative HBeAg and HBsAg have also been studied, but they are not clearly superior to the existing markers in terms of predicting long-term outcome. Markers such as circulating HBcrAg and pregenomic RNA are being assessed as additional potential markers (see “Measures of Hepatitis B Replication”).

### Acute Hepatitis B

Treatment in acute hepatitis B is generally supportive. Medication lists should be reviewed, and patients should be reminded to avoid medications metabolized by the liver if possible or limit the doses. This is particularly true for agents such as acetaminophen, which patients may be taking to minimize discomfort and fever. Treatment of fulminant hepatitis is also supportive, including liver transplantation for those patients who do not appear to have spontaneous recovery. Treatment with antiviral agents is not indicated for acute hepatitis B because approximately 95% will spontaneously seroconvert to anti-HBs; however, there is some evidence of benefit for cases of acute fulminant HBV. One randomized controlled trial of lamivudine versus placebo in acute severe hepatitis B showed greater decline in HBV DNA at 4 weeks with lamivudine but did not show long-term clinical benefit.<sup>274</sup> Several case series support that lamivudine, entecavir, and TDF may be beneficial when given before the onset of liver failure.<sup>275,276</sup> TAF should also be effective, but it has not been studied in fulminant, acute hepatitis B. If given, treatment for fulminant hepatitis B should be continued for 3 months after HBsAg seroconversion or for at least 12 months after HBeAg seroconversion if HBsAg loss does not occur. Treatment is also recommended for protracted severe acute hepatitis when increase in international normalized ratio and jaundice last for more than 4 weeks.<sup>250</sup>

### Selection of Patients for Treatment

(Also see Chapter 117.)

The currently available treatments all have limited short- and long-term efficacy, as well as substantial costs, both in terms of side effects (e.g., as in the case of IFN) and financial considerations. Thus not all patients warrant treatment. Current recommendations are to treat all cirrhotic patients with detectable HBV DNA. For noncirrhotic patients, treatment should be initiated with detectable HBV DNA—either HBeAg positive with a HBV DNA  $>20,000$  IU/mL or, in the case of HBeAg-negative disease, with HBV DNA  $>2000$  IU/mL—and evidence of liver disease (serum aminotransferases greater than two times the ULN [30 U/L for men and 19 U/L for women] or moderate-to-severe necroinflammation on biopsy) (see Table 145.3).<sup>250</sup> These HBV DNA cutoffs were initially arbitrary, but there is evidence that at these levels the risk of progressive liver disease and HCC increases.<sup>157</sup> Because HBV DNA levels can fluctuate, serial monitoring of levels is important if treatment is not initiated.

### Interferon

IFN- $\alpha$  is the recombinant version of one or more proteins that are naturally produced by the body in response to viral infection and has been used in the treatment of hepatitis B for more than 25 years. The precise mechanism of action as an anti-HBV therapeutic has not been established. One study demonstrated that IFN- $\alpha$  decreases transcription of pregenomic RNA and subgenomic RNA from cccDNA.<sup>277</sup> PEG IFN- $\alpha$ , given as a once-a-week injection of 180  $\mu$ g, has supplanted standard IFN- $\alpha$  for treatment of hepatitis B. Factors predictive of a high response to PEG IFN- $\alpha$  are low HBV DNA, aminotransferase elevation greater than twice the ULN, age, female sex, and genotype A or B (compared with genotype C or D).<sup>278</sup> There is also some evidence that IL28B genotype may be associated with response to PEG IFN- $\alpha$ .<sup>279</sup> Treatment of patients



with normal ALT results in response rates of <10%,<sup>280</sup> suggesting an important contribution of the immune response in IFN-based therapy.

PEG IFN- $\alpha$  is administered as subcutaneous injections. Patients should be treated for at least 48 weeks, and there is some evidence that up to 72 weeks will increase the rate of sustained response in those with HBeAg-negative disease.<sup>281</sup> In HBeAg-positive patients, 1-year rates of HBeAg seroconversion are 27%<sup>36,278</sup>, the durability of the response is improved if PEG IFN- $\alpha$  is continued for at least 24 weeks after seroconversion. Quantitative HBeAg is a predictor of response to PEG IFN- $\alpha$  in HBeAg-positive disease, with levels >20,000 IU/mL at 24 weeks being associated with decreased likelihood of HBeAg seroconversion.<sup>282</sup> If HBeAg or HBsAg loss occur, the durability of the response in one study of 172 HBeAg-positive patients followed for 3 years was 81% and 30%, for HBeAg and HBsAg, respectively.<sup>283</sup> In HBeAg-negative disease, the rate of sustained undetectable HBV DNA, measured 24 weeks after 48 weeks of treatment, is approximately 19%. Given the toxicity of IFN- $\alpha$ , most providers do not use it in HBeAg-negative disease as the tolerability of long-term nucleos(t)ides is better.

Acute side effects of PEG IFN- $\alpha$  therapy include flulike illness, fever, myalgias, headache, and fatigue. Other side effects seen after prolonged dosing include leukopenia and thrombocytopenia, hair loss, and changes in mood, including depression, which can be severe. IFN therapy can also lead to the development of autoantibodies, such as antithyroid antibodies, and worsening of other autoimmune disorders.<sup>284</sup> PEG IFN- $\alpha$  is contraindicated in decompensated cirrhosis because it may precipitate flares with subsequent decompensation.<sup>285</sup>

## Nucleoside and Nucleotide Analogues

Recognition that HBV, like HIV, has an RT step in its life cycle led to the testing of many nucleoside agents (Table 145.4). Because these agents are better tolerated than IFN- $\alpha$ , they are used long term for both viral suppression and improvement in histologic disease. Unlike IFN- $\alpha$ , nucleos(t)ide agents can be used in the setting of hepatic decompensation<sup>286</sup> and may even prevent or delay the need for transplantation.<sup>287,288</sup> However, resistance to some of these agents develops over time (see “Viral Resistance” later). The currently approved agents in this category include lamivudine, adefovir dipivoxil, telbivudine, entecavir, TDF, and TAF (also see Chapter 47). Emtricitabine, which is coformulated with TDF and TAF as an anti-HIV therapeutic, is also active against HBV but has not been approved for HBV treatment. These agents all suppress HBV replication but cannot directly eliminate cccDNA; thus, once therapy is discontinued, relapse of viremia occurs in the majority of HBeAg-positive patients who do not achieve HBeAg seroconversion and in nearly all HBeAg-negative patients.

## Lamivudine

Lamivudine is the negative enantiomer of 2',3'-dideoxy-3'-thiacytidine (see Chapter 47). It is phosphorylated by host enzymes, and the incorporation of the triphosphate form into DNA results in premature chain termination, which inhibits HBV DNA synthesis. Lamivudine is administered at 100 mg/day in those without HIV infection and is well tolerated, with side effects no different than placebo. In patients coinfecting with HIV, 300 mg/day is given to inhibit both HIV and HBV replication. Dose reduction is necessary for patients with renal insufficiency (creatinine clearance [CrCl] < 50 mL/min). Several large, randomized clinical trials in patients with HBeAg-positive and HBeAg-negative chronic hepatitis B demonstrate the efficacy of lamivudine (see Table 145.4).<sup>289,290</sup> A multinational study of Asian patients showed that HBeAg seroconversion rates increased over time from 17% to 27% at 2 years to 50% by 5 years.<sup>291,292</sup> An important predictor of HBeAg seroconversion at 1 year is the pretreatment aminotransferase level, ranging from 5% in patients with ALT less than two times the ULN to 64% in those with ALT greater than five times the ULN.<sup>293</sup> Patients with HBeAg-negative hepatitis B appear to have similar response rates,<sup>294–296</sup> as do patients who have previously failed IFN- $\alpha$  therapy.<sup>297</sup> Genotype, as with all the oral agents, is not a known predictor of response. Resistance to lamivudine develops when used as monotherapy limiting the durability of response; thus it is not a recommended first-line agent (see “Viral Resistance” later).

## Adefovir

Adefovir dipivoxil is the oral prodrug of adefovir, a phosphonate nucleotide analogue of adenosine monophosphate (see Chapter 47). It is effective against both wild-type and lamivudine-resistant HBV<sup>298–300</sup>; however, it is the least potent of the oral agents (see Table 145.4) and is not a recommended first-line agent. Adefovir is administered at 10 mg daily with adjustment for renal insufficiency, and it should not be given with TDF or TAF.

In clinical trials, headache and abdominal pain are the most common side effects. Nephrotoxicity occurs in 3% of patients with compensated liver disease after 4 to 5 years of continued adefovir and in 47% of patients who underwent a liver transplant.<sup>301,302</sup>

## Entecavir

Entecavir is a guanosine analogue that inhibits all three functions of the HBV polymerase: priming, negative-strand reverse transcription, and synthesis of the positive-strand HBV DNA (also see Chapter 47). It received US Food and Drug Administration (FDA) approval for the treatment of HBeAg-negative and HBeAg-positive chronic

**TABLE 145.4 Approved Agents for Treatment of Chronic Hepatitis B With 48-Week Response**

|                                   | PEG IFN- $\alpha$           | LAMIVUDINE            | ADEFOVIR                    | ENTECAVIR  | TELBIVUDINE           | TENOFOVIR DF/AF   |
|-----------------------------------|-----------------------------|-----------------------|-----------------------------|--|-----------------------|---|
| Route                             | Subcutaneous                | Oral                  | Oral                        | Oral   | Oral                  | Oral  |
| Dose                              | 180 $\mu$ g/wk              | 100 mg/d <sup>b</sup> | 10 mg/d <sup>b</sup>        | 0.5 mg/d <sup>b</sup> (1 mg/d if lamivudine resistant) | 600 mg/d <sup>b</sup> | 300 mg/d <sup>b</sup> for TDF and 25 mg/d for TAF           |
| Duration (wk)                     | 48                          | $\geq$ 48             | $\geq$ 48                   | $\geq$ 48  | $\geq$ 48             | $\geq$ 48   |
| Tolerability                      | Fair-poor: flulike symptoms | Good                  | Good: follow renal function | Good   | Good                  | Good: follow renal function and bone mineral density on TDF |
| HBeAg seroconversion              | 27%                         | 16%–21%               | 12%                         | 21%  | 22%                   | 21%   |
| Undetectable HBV DNA <sup>c</sup> | 25%–63%                     | 40%–73%               | 21%–51%                     | 67%–90%  | 60%–88%               | 76%–93%   |
| ALT normalization                 | 38%                         | 41%–75%               | 53%                         | 72%  | 65%                   | 74%   |
| HBsAg loss                        | 3%                          | <1%                   | 0%                          | 2%   | <1%                   | 3%  |
| Viral resistance                  | None                        | 15%–30%               | Minimal                     | None <sup>d</sup>                                      | 6%                    | 0%  |

<sup>a</sup>All data are for 1 year unless otherwise noted.

<sup>b</sup>Dose adjustment for creatinine clearance.

<sup>c</sup>Higher end of range for HBeAg-negative disease.

<sup>d</sup>None; otherwise, 7% if preexisting lamivudine resistance after 48 weeks and 50% after 5 years.

ALT, Alanine aminotransferase; anti-HBe, antibody to hepatitis B e antigen; HBV, hepatitis B virus; HBsAg, hepatitis B surface antigen; PEG IFN- $\alpha$ , pegylated interferon- $\alpha$ ; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate.

Modified from Lok AS, McMahon BJ. Chronic hepatitis B. *Hepatology*. 2009;50:661–662.

hepatitis B in 2005. As one of the most potent anti-HBV agents, it is one of the first-line therapies for chronic hepatitis B in patients without lamivudine-resistant HBV. The recommended dose of entecavir is 0.5 mg/day for those who are treatment naïve and 1 mg/day in those with prior lamivudine experience because the activity is lower against lamivudine-resistant HBV. The dose should be adjusted for those with CrCl <50 mL/min. In a double-blind controlled trial of 709 HBeAg-positive patients, the 354 who received 48 weeks of entecavir had a decline in HBV DNA of 6.9 log<sub>10</sub> copies/mL, which was superior to the 5.4 log<sub>10</sub> copies/mL in the 355 who received lamivudine.<sup>303</sup> Furthermore, 67% in the entecavir group compared with 36% in the lamivudine group achieved HBV DNA that was <300 copies/mL, and histologic improvement occurred in 72% and 62% in the entecavir and lamivudine groups, respectively. However, HBeAg seroconversion was similar between the groups. Results after 96 weeks of treatment showed similar efficacy rates.<sup>304</sup> Similarly, in the HBeAg-negative patients, 70% of the 331 in the entecavir group had histologic improvement compared with 61% of the 317 in the lamivudine group. An undetectable HBV DNA was achieved in 90% in the entecavir group compared with 72% in the lamivudine group. In a study from China, after 7 years of entecavir undetectable HBV DNA was achieved by 96.8% of the HBeAg-positive patients and 100% of the HBeAg-negative patients.<sup>305</sup> Side effects to entecavir include headache, abdominal pain, and diarrhea.

Entecavir inhibits HIV-1 replication and can select for the lamivudine-resistant HIV mutation M184V when given as monotherapy<sup>306</sup>; thus HIV status should be checked before initiating therapy with entecavir. Furthermore, it should not be given to an HIV-coinfected patient without a suppressive HAART regimen.

### Telbivudine

Telbivudine is an L-nucleoside analogue that is structurally related to lamivudine (see Chapter 47) and thus has overlapping resistance patterns at the rtM204 position. Telbivudine is administered at 600 mg daily and has potent activity. However, due to relatively high rates of resistance and cross-resistance with lamivudine, it is not a first-line agent. It is controversial as to whether it inhibits HIV-1 replication. The GLOBE trial, the registration trial for telbivudine, was an international study randomizing patients to telbivudine (*n* = 680) or lamivudine (*n* = 687) daily.<sup>307</sup> At 52 weeks the proportion with undetectable HBV DNA (<300 copies/mL) was greater in the telbivudine than the lamivudine group (60% vs. 40%, respectively). At week 104 telbivudine continued to be superior, but the proportion with undetectable HBV DNA decreased to 56%. These changes resulted from the development of resistance at rtM204I, which reached 25% at week 104. Results of the HBeAg-negative group also showed superiority of telbivudine over lamivudine and demonstrated poor durability of response due to development of resistance, which occurred in 8.6% of the telbivudine group.

### Tenofovir Disoproxil Fumarate (TDF) and Tenofovir Alafenamide (TAF)

TDF was approved for treatment of hepatitis B in 2008 and is also approved for treatment of HIV-1 (see Chapters 47 and 128). TDF is potent, and resistance has not clearly been identified; thus it is a recommended first-line agent in patients with and without lamivudine-resistant HBV. TDF is dosed at 300 mg daily and needs to be adjusted for renal insufficiency. In two phase III trials comparing TDF with adefovir, viral suppression occurred more often in those receiving TDF than adefovir dipivoxil in both HBeAg-negative (93% vs. 63%, *P* < .001) and HBeAg-positive patients (76% vs. 13%, *P* < .001) after 48 weeks.<sup>264</sup> Of significance, more HBeAg-positive patients treated with TDF than those treated with adefovir dipivoxil had normalized ALT levels (68% vs. 54%, *P* = .03) and loss of HBsAg (3% vs. 0%, *P* = .02). TDF produced a similar HBV DNA response in patients regardless of prior lamivudine use. Continued use of TDF up to 7 years demonstrates that 99% of the HBeAg-positive and HBeAg-negative patients on treatment have HBV DNA <29 IU/mL.<sup>308</sup> No signature TDF-resistance mutations have been identified. The major side effects of TDF are renal insufficiency with increases in serum creatinine and decrease in bone mineral density.

TAF has equivalent anti-HBV activity to TDF but has less kidney and bone toxicity.<sup>309,310,311</sup> It was approved for treatment against hepatitis B in November 2016 and is given at a dose of 25 mg once daily. Two randomized, double-blinded studies compared TAF with TDF for the treatment of HBeAg-positive and HBeAg-negative chronic hepatitis B. After 48 weeks of treatment, HBV DNA of <29 IU/mL was achieved in 94% of the TAF group and in 93% of the TDF group in the HBeAg-negative study.<sup>310</sup> The proportions were 64% and 67% for TAF and TDF, respectively, in the HBeAg-positive study.<sup>311</sup> In both studies the patients receiving TAF had significantly smaller declines in bone mineral density and lower changes in glomerular filtration rate.

### Other Agents and Combination Therapy

The other agent that is approved for HIV treatment but is also active against HBV is emtricitabine. Emtricitabine is similar to lamivudine, but it has a longer half-life. Mutations that confer resistance to lamivudine also confer resistance to emtricitabine (see Chapters 47 and 128).

Combination therapy for hepatitis B is attractive because it may increase potency and decrease the rates for developing resistance; however, these concepts have not been clearly demonstrated to date, so combination therapy is not generally recommended. PEG IFN- $\alpha$  and lamivudine for 48 weeks had greater HBV DNA decline compared with monotherapy with either one alone; however, the relapse rate in the combination group was higher, so 24 weeks after the end of treatment, there was no difference between the PEG IFN- $\alpha$  group alone or with lamivudine.<sup>36,265</sup> Another study showed that adding PEG IFN- $\alpha$  for 24 weeks after 24 weeks of entecavir improved HBV DNA, HBeAg, and HBsAg responses compared with continuing entecavir for 48 weeks; however, this study did not have a PEG IFN- $\alpha$  monotherapy arm.<sup>312</sup> Other studies have not shown a benefit of PEG IFN- $\alpha$  added after 12 weeks of entecavir compared with PEG IFN- $\alpha$  alone.<sup>313,314</sup> In a randomized, open-label study, the combination of TDF and PEG IFN- $\alpha$  demonstrated greater HBsAg loss (9%) compared with either TDF or PEG IFN- $\alpha$  alone (0% and 2.8%, respectively); however the rates were low in all groups.<sup>315</sup> Another randomized open-label study of HBeAg-positive patients on TDF or entecavir for more than 12 months with HBV DNA <2000 IU/mL found that adding PEG IFN- $\alpha$  compared with continuing the nucleoside alone for 48 weeks had greater HBeAg seroconversion, but it was not statistically significantly different (18% vs. 8%, *P* = .31).<sup>316</sup>

Given the expansion in the number of oral agents available, it is desirable to consider combining these as is done in HIV therapy. The oral anti-HBV drugs all target the same part of the HBV life cycle; thus the situation is not entirely analogous to HIV, in which drugs target different aspects of the viral life cycle. In a randomized, open-label study combining the two most potent agents, entecavir and TDF compared with entecavir alone, there was no difference after 96 weeks of therapy between the 197 subjects receiving combination therapy and the 182 subjects receiving entecavir alone for the primary end point (<50 IU/mL) of HBV DNA (83.2% and 76.4%, respectively, *P* = .088).<sup>317</sup> However, when stratified by HBeAg status, the HBeAg-positive patients who received combination therapy were more likely to have a virologic response compared with the monotherapy group (*P* = .046), which was entirely attributable to those with baseline HBV DNA >10<sup>8</sup> IU/mL. There were no differences in HBeAg or HBsAg seroconversion in any groups. Thus this study suggests that combination therapy may be most beneficial in those with high HBV DNA levels, but more studies of longer duration are needed to confirm this.

### Viral Resistance

The major limitation in the use of oral agents for hepatitis B is the development of resistance, which has been observed with the use of all approved oral agents except tenofovir (DF and AF), albeit at different times and frequencies. Resistance mutations differ based on the structural group of the drug, which include: L-nucleosides, D-cyclopentanes, and acyclic nucleotides. Cross-resistance occurs between agents in the same group and may decrease sensitivity between groups. The L-nucleosides group consists of lamivudine, emtricitabine, and telbivudine, with the primary resistance mutation occurring at position rt204 of the HBV polymerase, which is essential for polymerase activity, and leads to

substitution of isoleucine (I) or valine (V) or serine (S) for methionine.<sup>291,318,319</sup> Of the L-nucleosides, genotypic resistance develops most rapidly to lamivudine, with 14% to 32% of patients having evidence of mutations after 1 year of treatment.<sup>289,290</sup> Cumulative resistance increases over time, with one study demonstrating rates of 15%, 38%, 55%, 67%, and 69%, at 1, 2, 3, 4, and 5 years, respectively.<sup>291</sup> HBV mutants with the rtM204V/I appear to have reduced replication capacity in vitro and in vivo compared with wild-type HBV.<sup>320</sup> However, the development of a compensatory mutation at rtL180M restores replication capacity, at least in vitro.<sup>320</sup> The rates of lamivudine resistance in patients treated for HBeAg-negative chronic hepatitis B appear to be more variable (0%–27% at 1 year and 10%–56% at 2 years).<sup>296,321</sup>

Although the rates of resistance to telbivudine are lower than to those of lamivudine (11% vs. 26% at 2 years), resistance is high enough to preclude its widespread use.<sup>322</sup> Telbivudine is not active against HBV strains bearing lamivudine mutations rtL180M/rtM204V/I but remained active against the rtM204V single mutant in vitro, potentially explaining the difference in resistance profiles between telbivudine and lamivudine. Against HBV genomes with known telbivudine-resistance mutations, rtM204I and rtL80I/M204I, telbivudine, lamivudine, and entecavir lost 353- to >1000-fold activity, whereas adefovir and TDF exhibited no more than threefold to fivefold change.<sup>323</sup>

The second group, D-cyclopentanes, consists of entecavir, which has a high genetic barrier to resistance as it requires the presence of multiple mutations, including the rtM204V and rtL180M mutations, along with either the rtT184S/A/I/L or rtC202G/C or rtM250 mutations. In patients who are treatment naïve and do not have a preexisting rtM204V or rtL180M mutation, the rates of resistance are <1% per year with a 5-year cumulative rate of 1.2%.<sup>324</sup> However, in patients with the characteristic lamivudine mutations rtM204V and rtL180M, the cumulative incidence is 51% after 5 years; therefore treatment with tenofovir (DF or AF) is preferred over entecavir in patients with lamivudine-resistant HBV. Due to overlapping resistance mutations with lamivudine, lamivudine should be discontinued when entecavir is started, to theoretically decrease the risk for developing entecavir resistance.

The third group, acyclic nucleotides, consists of tenofovir (DF and AF) and adefovir dipivoxil. Resistance to adefovir dipivoxil occurs at residue rtN236T and rtA181V/T of the HBV polymerase. After 240 weeks of treatment the cumulative probability of resistance was 29% for HBeAg-negative patients. In patients with prior lamivudine-resistant mutations, cumulative incidence of adefovir dipivoxil resistance was higher—43% at 4 years.<sup>325</sup> In HBV cell lines expressing adefovir dipivoxil resistance mutations rtN236T and rtA181V, telbivudine remained active as shown by respective changes of 0.5-fold (rtN236T) and 1.0-fold (rtA181V and rtA194T).<sup>323</sup>

TDF resistance rates appear low, but precise mutations with confirmed resistance to TDF have not been characterized. Through 5 years of therapy, TDF maintains HBV DNA suppression, and there is no clear evidence of resistance.<sup>326</sup> The adefovir resistance mutations rtA181T/V and rtN236T confer decreased sensitivity to TDF. In one retrospective study, patients on TDF with these mutations were less likely to achieve HBV DNA <400 copies/mL than those without the mutations.<sup>327</sup> Because TAF is a different prodrug than TDF, the resistance patterns will be the same.

Resistance is suspected with the reappearance of HBV DNA in a patient who was previously undetectable or with a >1 log<sub>10</sub> increase in HBV DNA in a patient who is compliant with therapy. Serum aminotransferases may or may not become elevated, and there are rare instances of acute exacerbations and hepatic decompensation.<sup>328</sup> Commercial assays are available to detect resistance mutations. If resistance occurs, current recommendations are to change therapy to the most effective agent that does not share cross-resistance. In general, the L-nucleosides (lamivudine, telbivudine, and entecavir) are not cross-resistant with the acyclic nucleotides (adefovir, tenofovir DF/AF), with tenofovir DF or AF preferred. In cases of multidrug resistance a combination of entecavir and tenofovir DF or AF is recommended.<sup>329,330</sup>

Given the overlapping reading frames, mutations in the HBV polymerase can lead to changes in the surface antigen, including in the immunogenic “a” determinant. One study demonstrated that a triple mutant that occurs on lamivudine monotherapy (rtV173L/rtL180M/

rtM204V) leads to the surface changes sE164D/I195M. These changes in the surface antigen lead to reduced binding of anti-HBs in vitro, and thus the virus behaves as a vaccine escape mutant.<sup>331</sup>

## Monitoring and Treatment Duration of Nucleos(t)ide Therapy

Virologic response to nucleos(t)ide analogue therapy is based on HBV DNA levels and should be evaluated every 3 months after initiating or changing therapy until the HBV DNA is undetectable and then monitored every 3 to 6 months. In addition to HBV DNA, HBeAg (in HBeAg-positive patients), and ALT should also be monitored (Table 145.5). If there is less than a 1-log IU/mL decline in the first 3 months, this is defined as primary nonresponse, which is most commonly due to noncompliance, especially if a patient is on entecavir, TDF, or TAF. If a patient with a primary nonresponse is compliant, then resistance testing is appropriate to determine a change in the treatment strategy. After 12 months of therapy a virologic response is defined as having an undetectable HBV DNA by a sensitive PCR assay. However, if after 12 months of therapy in a patient who is compliant, the HBV DNA is detectable but has declined more than 1 log<sub>10</sub>, then this is defined as a partial virologic response. In this case a treatment-naïve patient receiving entecavir, TDF, or TAF can continue the same therapy because there is a low rate of resistance. In a patient receiving a less potent drug, changing to entecavir, TDF, or TAF is recommended. In patients with low-level persistent viremia on entecavir, TDF, or TAF, there is little evidence that adding a second drug is beneficial. Virologic breakthrough can occur at any time and is defined as >1 log IU/mL increase in HBV DNA from the nadir value on therapy. The main causes of virologic breakthrough are noncompliance with therapy or viral resistance (see “Viral Resistance” earlier).

The duration of therapy for most individuals is lifelong, but there are some patients for whom discontinuation of therapy can be considered. For patients with HBeAg-positive chronic hepatitis B who have HBeAg seroconversion, current recommendations are to treat for 1 year after seroconversion and then consider discontinuing therapy if HBV DNA is undetectable and ALT is normal.<sup>250</sup> The exception

**TABLE 145.5 Virologic Response to Nucleos(t)ide Treatment**

| RESPONSE                   | EVALUATION TIME (MONTHS) | CRITERIA   | THERAPEUTIC STRATEGY   |
|----------------------------|--------------------------|--|--|
| Primary nonresponse        | 3                        | <1 log <sub>10</sub> decline   | Assess compliance. If compliant, obtain resistance test and change treatment.  |
| Partial virologic response | 12                       | HBV DNA detectable but >1 log <sub>10</sub> decline from baseline          | If treatment naïve and taking entecavir or tenofovir DF/AF, then continue.<br>If treatment naïve and taking less potent drug, change to entecavir or tenofovir DF/AF.<br>If treatment experienced on entecavir, either change to or add tenofovir DF/AF. |
| Virologic response         | 12                       | HBV DNA undetectable by sensitive PCR assay                                | Continue treatment and monitor every 3–6 mo.   |
| Virologic breakthrough     | Any time                 | >1 log <sub>10</sub> increase in serum HBV DNA level from on therapy nadir | Obtain resistance test and change therapy (see “Viral Resistance” in text).  |

AF, Adefovir; DF, disoproxil fumarate; DNA, deoxyribonucleic acid; HBV, hepatitis B virus; PCR, polymerase chain reaction.



to this is that patients with cirrhosis should not be considered for discontinuation of therapy. Once HBeAg seroconversion has occurred, the rate of durable response after discontinuation ranges from 30% to 80%.<sup>332</sup> However, due to the possibility of relapse, some recommend continuing therapy until HBeAg seroconversion, which is infrequent. In one study recurrent viremia occurred in 90% of the 39 patients who stopped therapy a median of 12 months after seroconversion, compared with 0% in those who continued therapy, so monitoring every 3 months with serum aminotransferases and HBV DNA is important after discontinuation of therapy.<sup>250,333</sup> There is some evidence that quantitative HBeAg may be useful to predict who will relapse after therapy discontinuation.<sup>334</sup> If a patient does not have HBeAg seroconversion but has suppressed HBV DNA replication, then therapy should be continued indefinitely because there is evidence of progressive enhancement of the HBeAg seroconversion rate with longer duration of treatment, and patients may continue to derive histologic benefit.<sup>267</sup> In HBeAg-negative patients therapy should be continued indefinitely because relapse occurs in the majority of cases except in rare instances of HBeAg seroconversion. However, the European guidelines suggest that, in selected noncirrhotic HBeAg negative patients with undetectable HBV DNA for at least 3 years, discontinuation of therapy could be considered with close monitoring.<sup>65</sup> In support of this, one study of HBeAg-negative patients with undetectable HBV DNA for  $\geq 3.5$  years found that 13 of 21 people who discontinued TDF remained off therapy through 144 weeks.<sup>335</sup> In both HBeAg-negative and HBeAg-positive disease, therapy discontinuation can be considered 6 to 12 months after HBeAg seroconversion, which only occurred in 8% after 6 years of TDF.

## MANAGEMENT IN SPECIAL POPULATIONS

### Liver Transplantation

Recurrent hepatitis B after liver transplantation can occur in patients with a history of acute or chronic hepatitis B before transplantation or in HBsAg-negative recipients of anti-HBc-positive donors. Historically, rates of recurrence of HBV infection were high (90%), and the consequences of reinfection were devastating, such that HBV was considered a relative contraindication to liver transplantation. HBIG was introduced in the early 1980s and nucleosides in the early 1990s, with progressive reduction in likelihood of HBV recurrence with each intervention. Using HBIG alone, recurrence rates ranged from 20% to 25%, depending on the schedule and trough anti-HBs titer. The addition of oral nucleos(t)ide therapy (lamivudine) to HBIG was shown to further decrease recurrence rates to as low as 10% and improve mortality.<sup>336,337</sup> Similar to nontransplant patients, however, treatment with lamivudine alone resulted in the emergence of lamivudine resistance and severe and even fatal posttransplantation disease.<sup>338</sup> Tenofovir (DF or AF) or entecavir are now the preferred nucleos(t)ide analogues and are  $>95\%$  effective in protecting the grafts from recurrent disease. Both have been shown to suppress lamivudine-resistant variants<sup>298</sup> and even resolution of graft failure in those patients who have them.<sup>339</sup> In a recent study evaluating long-term outcomes in 265 chronic HBV liver transplant recipients treated with entecavir without HBIG, followed for a median of 59 months, 92% were negative for HBsAg at 8 years, and 100% had undetectable HBV DNA. Eighty-five percent of the patients were alive at 9 years.<sup>340</sup> In a randomized controlled trial comparing TDF and emtricitabine with and without HBIG in orthotopic liver transplant recipients, followed for a median of 3.4 years, no recurrences of HBV occurred in either arm, suggesting therapy with TDF and emtricitabine without HBIG is effective at reducing HBV recurrence.<sup>341</sup> Although an increasing number of transplant centers are using nucleos(t)ide analogues alone for post-transplant prophylaxis, HBIG continues to be recommended in certain high-risk patients, including individuals with high HBV DNA levels, known antiviral resistance, higher risk for recurrence of HCC, HIV coinfection, and HDV coinfection.<sup>342</sup> Finally, although TDF is most studied in posttransplant patients, TAF is a better alternative due to its safer renal profile. In summary, with currently available combination therapy, survival is excellent in patients undergoing liver transplantation for chronic hepatitis B, even in those with active viral replication pretransplantation.

Vaccination after transplantation has variable success; in one study, active immunization after liver transplantation induced anti-HBs titers ( $>10$  IU/mL) in 82% of liver transplant candidates with inactive HBV disease<sup>343</sup>; however, in a later study of liver transplant recipients with cirrhosis who received conventional HBV vaccine after transplantation, only 18% achieved protective anti-HBs titers.<sup>344</sup> Other reports of conventional vaccine failure after liver transplantation have been reported.<sup>345,346</sup> The use of repeated immunization with recombinant HBV vaccine emulsified in novel adjuvants in patients who were HBV DNA negative after orthotopic liver transplantation resulted in anti-HBs levels ranging from 721 to 83,121 IU/L, depending on the concentration, in 80%, which remained stable after discontinuation of HBIG.<sup>346</sup> However, in a more recent study of eight liver transplant recipients with similar characteristics but in whom HBIG was stopped, the use of an adjuvant containing HBV vaccine resulted in development of anti-HBs in only one patient.<sup>347</sup>

## Human Immunodeficiency Virus

The efficacy of the nucleos(t)ide agents does not seem to be compromised by coinfection with HIV.<sup>348</sup> However, with lamivudine monotherapy, drug resistance develops more rapidly than in HBV monoinfection.<sup>349</sup> Lamivudine-resistant HBV is common in HIV coinfection because it has been a backbone agent for HIV therapy for years. Tenofovir (DF and AF) has potent activity against both HIV and HBV<sup>350</sup> and is also effective against lamivudine-resistant HBV.<sup>351–353</sup> The combination of TDF or TAF and emtricitabine (or TDF and lamivudine) is therefore now recommended in the treatment of chronic HBV in HIV-infected individuals.<sup>354</sup> TAF is less likely to adversely impact renal function and bone mineral density without compromising efficacy and thus is preferred over TDF when CrCl  $> 30$  mL/min. The three approved fixed-dose TAF-containing regimens include elvitegravir/cobicistat/emtricitabine/TAF; emtricitabine/rilpivirine/TAF; and emtricitabine/TAF. Current Department of Health and Human Services (DHHS) HIV treatment guidelines recommend initiation of ART in all HIV-infected patients, including those with HIV/HBV coinfection.<sup>354</sup> If TDF or TAF cannot be safely used, entecavir should be used with a fully suppressive ART regimen because entecavir can also inhibit HIV replication and select for the HIV drug-resistance mutation M184V.<sup>306</sup> A higher dose of entecavir (1 mg) should be used in patients who have already received lamivudine or have known lamivudine resistance. However, due to the relatively high rate of resistance of entecavir in those with lamivudine resistance (see “Viral Resistance” earlier), HBV DNA should also be monitored frequently.<sup>354</sup> If HBV needs treatment and a fully active anti-HIV regimen cannot be given, then drugs that are active only against HBV and do not lead to HIV drug resistance, such as adefovir dipivoxil or PEG IFN, are recommended. Telbivudine is not recommended, given its intermediate rate of resistance in HBV monoinfection and unknown rates in HIV coinfection. Treatment is indefinite unless HBsAg seroconversion occurs.

## Recipients of Immunosuppressive Therapies

A number of cytotoxic chemotherapeutic agents, immunomodulators, and immunosuppressants have been associated with reactivation of HBV. Agents that have been most commonly associated are prednisone, anthracyclines, and anti-CD20 blockers, specifically rituximab or ofatumumab as part of chemotherapy.<sup>355</sup> The frequency of reactivation in patients undergoing cytotoxic chemotherapy ranges from 14% to 72% in different studies and case series, depending on serologic status of the patient and potency of immunosuppressants. Patients at risk for reactivation include those with resolved infection (HBsAg negative, anti-HBs, and anti-HBc positive), those with inactive chronic hepatitis B (HBsAg positive, HBV DNA undetectable), and those with untreated chronic active hepatitis B (HBsAg positive, HBV DNA detectable). Tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) inhibitors, including infliximab, etanercept, and adalimumab and low-dose steroids, have a low-to-moderate risk of reactivation. Very few cases of reactivation have been reported with the use of immunosuppressants, including azathioprine and methotrexate.<sup>355</sup> Anthracyclines carry a moderate risk for HBV reactivation in HBsAg-negative, anti-HBc-positive patients and a high

risk in HBsAg-positive patients. The risk of reactivation of HBV associated with rituximab, a monoclonal antibody directed against B-cell marker CD20 and increasingly used in standard chemotherapy regimens for many rheumatologic disorders, carries the highest risk of reactivation in patients who are anti-HBc positive, with or without anti-HBs. In a recent meta-analysis of case series and reviews of HBV reactivation in lymphoproliferative disorders associated with rituximab, there was a more-than-fivefold higher rate of HBV reactivation in patients with a history of prior HBV exposure (anti-HBc positive, with or without anti-HBs) who received rituximab-based therapy, compared with those who did not.<sup>356</sup> In a study of 46 HBsAg-negative/anti-HBc-positive patients (the majority of whom were anti-HBs positive), 25% of patients receiving rituximab and cyclophosphamide-hydroxydaunorubicin-Oncovin (vincristine)-prednisone (CHOP) for diffuse large B-cell lymphoma reactivated HBV (became HBsAg positive) compared with 0% who received CHOP alone.<sup>357</sup> In 2013 the FDA issued a black box warning concerning the risk of HBV reactivation in patients receiving anti-CD20 monoclonal antibodies after more than 100 cases of fatal HBV-related liver failure associated with rituximab or ofatumumab were reported in postmarketing data from the FDA adverse-event report system between 1997–2012.<sup>357a</sup>

Prophylaxis is recommended in HBsAg-positive patients at highest risk of reactivation, if untreated, especially if they are taking B-cell-depleting drugs, anthracycline derivatives, or high-dose steroids or receiving bone marrow transplants.<sup>358</sup> Anti-HBc-positive patients who are HBsAg positive or negative and at moderate risk of reactivation, that is, patients on TNF- $\alpha$  inhibitors, low-dose steroids, and anthracycline inhibitors (if HBsAg negative) can be closely monitored or receive prophylaxis.<sup>355</sup> Although lamivudine is the most widely studied antiviral agent (associated with reductions in risk of HBV reactivation of 79%–100%), tenofovir (DF or AF) and entecavir are now preferred, especially when the duration of prophylaxis is expected to be longer than 1 year. Entecavir and TDF have higher antiviral potency and lower rate of resistance with long-term use. In a randomized trial in patients with diffuse large B-cell lymphoma receiving chemotherapy with rituximab-CHOP (R-CHOP), entecavir was found to be superior to lamivudine for the prevention of HBV reactivation and HBV-related hepatitis in those with prechemotherapy HBV DNA  $\geq 2000$  IU/mL. TDF has also shown to be effective in reducing HBV reactivation in anti-HBc-positive patients with hematologic malignancies treated with rituximab.<sup>359</sup> The optimal duration of prophylactic therapy is unknown; current recommendations are to start preferred antiviral drugs with immunosuppressive medication and continue 6 to 12 months after the last dose of treatment. For patients receiving B-cell-depletion agents, therapy should be continued 12 to 18 months after the last dose, as reactivations as late as 17 months after completion of therapy have been reported.<sup>250,360,361</sup> Among patients with higher levels of baseline HBV DNA ( $>2000$  IU/mL), the American Association for the Study of Liver Disease (AASLD) recommends continuing antiviral therapy until they reach treatment end points for HBV infection.<sup>250</sup> Among bone marrow transplantation recipients, the risk of reactivation of HBV in HBsAg-positive patients is up to 54%.<sup>358</sup> The use of preemptive antivirals therefore, in these situations, is also recommended. Of interest, a donor bone marrow with natural immunity (anti-HBs and anti-HBc positive) may result in clearance of HBsAg.<sup>362</sup>

### Treatment of Hepatitis B Virus During Pregnancy

All women presenting for prenatal care should be routinely tested for HBsAg early in their pregnancy. There are at least three indications for treatment of HBV during pregnancy: (1) to treat active liver disease in situations when therapy cannot be delayed, (2) the continuation of HBV treatment in a mother who becomes pregnant while on HBV therapy, and (3) to prevent transmission of HBV from mother to child in mothers with high circulating viremia. All treatment decisions need to take into consideration the risks and benefits for mother and fetus (potential exposure to teratogenic drugs). TDF (category B) and lamivudine (category C) have been studied most extensively in pregnancy. In the Antiretroviral Pregnancy Registry and the Development of Antiretroviral Therapy study (DART), birth defects among women exposed to HBV therapy (mostly TDF and lamivudine) were similar to that in the general

population in the first and second trimesters and to other classes of antiretrovirals.<sup>363</sup> Similarly, to date no adverse outcomes have been reported with telbivudine (category B).<sup>364</sup> A recent systematic review and meta-analysis of antiviral therapy in chronic HBV found no evidence of fetal adverse effects or congenital defects associated with TDF.<sup>364</sup> There are minimal data on the safety of adefovir or entecavir or TAF (category C) in pregnancy; thus they should be avoided in pregnancy until more clinical data are available. Interferon is contraindicated in pregnancy. Based on data demonstrating a decrease in perinatal transmission, current guidelines recommend treatment of pregnant women in the third trimester when HBV DNA  $> 200,000$  IU/mL (see also “Prevention of Perinatal Transmission” later). The guidelines also recommend that pregnant women in the immunoactive phase are managed the same as nonpregnant women, with the caveat that only drugs tested in pregnant women should be used.<sup>250</sup>

### OTHER MANAGEMENT ISSUES IN CHRONIC HEPATITIS B

Patients with chronic hepatitis B should be counseled about disease-modifying factors, as well as means to prevent spread of HBV to other persons. For example, patients should be counseled about the means of spread of delta hepatitis and hepatitis C to avoid superinfection with these viruses. Patients should also be counseled to consume minimal, if any alcohol, in the absence of data regarding safe levels of consumption and because consumption of large amounts of alcohol is clearly a risk factor for more rapid progression to cirrhosis. Other major issues in the management of patients with chronic hepatitis B include prevention of hepatitis A, prevention of spread of HBV, and surveillance for HCC.

#### Hepatitis A Vaccination

The official recommendation of the Advisory Committee for Immunization Practices (ACIP) in the United States is that all persons with chronic liver disease be vaccinated against hepatitis A virus (HAV).<sup>365</sup> The data supporting this recommendation are not strong<sup>366</sup> because at least one study revealed that the risk of fulminant HAV is significantly increased only in patients with underlying hepatitis C and not hepatitis B.<sup>367</sup> However, the guidelines of the AASLD call for immunization of all patients with chronic HBV against HAV.

#### Screening and Vaccination of Contacts

Sexual and household contacts of persons with HBV are at increased risk of infection. All sexual partners and household contacts should be tested for HBV and vaccinated if seronegative. Until the immunization series is complete, sexual partners should use barrier methods. Both patients and contacts should be counseled regarding the modes of transmission and advised on methods to prevent household transmission, including avoiding sharing of items that might be contaminated with small amounts of blood, such as toothbrushes, and the need to cover open wounds. Pregnant women or women who want to become pregnant and are infected with HBV should also be counseled on the risk of transmission to the newborn and the method to prevent such transmission. Postexposure prophylaxis with HBIG and hepatitis B vaccine should be given to infants born to HBsAg-positive mothers, unvaccinated infants whose mothers or primary caregivers have acute hepatitis B, sexual contacts of persons with hepatitis B, and health care workers after occupational exposure to HBsAg-positive blood depending on their vaccination and vaccine response status. Household and sexual contacts of persons with chronic HBV infection should be vaccinated.

Recommendations for the health care worker infected with HBV vary from country to country. Although there is general agreement that individuals with HBeAg or HBV DNA  $> 10,000$  IU/mL pose the greatest risk of transmission,<sup>368</sup> there have been documented cases of transmission from health care workers in the absence of HBeAg and during “low-risk” procedures.<sup>369,370</sup> Updated Centers for Disease Control and Prevention (CDC) guidelines published in 2012 recommend that most chronically HBV-infected providers and students who adopt standard infection control precautions do not require curtailing their practices. Previous recommendations were updated to also include the following: (1) prenotification of a health care provider's or student's HBV status is no longer required; (2) the use of HBV DNA serum levels rather than

HBeAg status is recommended to monitor infectivity; and (3) for those health care professionals requiring oversight (i.e., health care worker performing highly exposure-prone procedures), there are specific suggestions for composition of expert review panels and threshold value of serum HBV DNA considered “safe” for practice (<1000 IU/mL).<sup>371</sup>

### Surveillance for Hepatocellular Carcinoma

In multiple longitudinal studies, carriers of HBsAg have been shown to be at increased risk of developing HCC.<sup>372</sup> The risk is higher among males and those who are older, have a family history of HCC, consume alcohol regularly, have elevated serum ALT levels, have positive HBeAg status, have hepatitis B DNA >10<sup>4</sup> copies/mL (≈2000 IU/mL), have genotype C virus, have cirrhosis, and have a longer duration of infection.<sup>177,373,374</sup> The goal of HCC screening is to detect small, surgically resectable tumors because the prognosis for more advanced lesions is poor. Current recommendations are that HBsAg-positive persons at high risk for developing HCC should be periodically screened; these include Asian males older than 40 years, Asian females older than 50 years, African/North American blacks older than 20 years, those with a family history of HCC, those with cirrhosis, and persons with ALT elevations or HBV DNA >2000 IU/mL, or both.<sup>250</sup> In a single, randomized, controlled trial conducted in China comparing surveillance using ultrasonography and  $\alpha$ -fetoprotein levels with no surveillance, the incidence of HCC-related mortality was reduced by 37%.<sup>375</sup> However, there are no randomized trials of HCC surveillance in US populations, and several studies show that  $\alpha$ -fetoprotein determination lacks adequate sensitivity and specificity. Therefore the current recommendation for HCC screening is with ultrasonography alone every 6 months.<sup>375</sup> The role of other imaging technologies, such as computed tomography, in screening have not been adequately assessed. Although the risk of HBV is decreased in patients achieving viral suppression, it is not eliminated, especially in the setting of cirrhosis. A study following Chinese patients with chronic hepatitis B on adefovir for 10 years found that the incidence of HCC in cirrhotic patients was 43.1% compared with 7.1% in noncirrhotics.<sup>376</sup> Furthermore, cirrhotic patients who achieved virologic suppression had a lower incidence of HCC than those who did not (27.8% vs. 62.2% 10-year incidence), and the same pattern was seen in noncirrhotic patients (4.0% vs. 13.4%). Thus, even with virologic suppression, HCC occurs and surveillance every 6 months is recommended.

### PREVENTION OF HEPATITIS B VIRUS INFECTION

The HBV vaccine is effective in preventing HBV and its complications. It has been shown in multiple studies to be safe, efficacious, and immunogenic, with seroconversion rates of >90% in healthy adults and seroprotection rates in infants born to HBV-infected mothers of 88.5% to 95.8%.<sup>377–379</sup> Successful vaccination prevents HBV infection and therefore reduces HCC-related mortality.<sup>380</sup> In Taiwan, which was an early adopter of universal HBV vaccination in children, the average annual incidence of HCC in 6- to 14-year-olds declined from 0.70 per 100,000 children between 1981 and 1986, before widespread vaccination, to 0.36 between 1990 and 1994, after initiation of widespread vaccination ( $P < .01$ ).<sup>381</sup> The corresponding rates of mortality from HCC also decreased.

### Active Immunization

Vaccines against hepatitis B have been available since 1982. The first vaccines were plasma derived and have now been completely replaced by recombinant vaccines. The vaccines contain small HBsAg as the major component produced in transfected yeast cells. Hepatitis B vaccines are available in monovalent forms for use in adults and for birth doses (Engerix-B [GSK; Philadelphia, PA] and Recombivax HB [Merck; Kenilworth, NJ]) or in combination with other vaccines for infant vaccination, including diphtheria-tetanus-pertussis (DTP), *Haemophilus influenzae* type b (Hib), and inactivated polio vaccine (IPV) (Pediarix; GSK). Pediarix can only be administered before age 6 weeks through 6 years of age. A combination vaccine (TWINRIX R; GSK), which expresses both HBsAg and hepatitis A, is also available and is approved for use in persons  $\geq 18$  years of age in the United States and Europe

(see Chapter 316). Despite high efficacy of recombinant vaccines, certain populations remain at risk of suboptimal or nonresponse to these vaccines (see “Efficacy”). Recently, the first and only two-dose hepatitis B vaccine, Heplisav-B, was approved for use in adults 18 years of age and older. This novel vaccine uses HBsAg adjuvanted with immunostimulatory phosphorothioate oligodeoxyribonucleotide sequences that stimulate the innate immune system through Toll-like receptor 9 (TLR-9). The approval of Heplisav-B was based on data from three phase III noninferiority trials of nearly 10,000 adult participants who received the vaccine.<sup>382,383</sup> Results from the three trials have demonstrated high levels of protection early after administration with the largest trial, which included 6665 participants, demonstrating a statistically significantly higher rate of protection of 95% compared with 81% for Engerix-B. In a subgroup analysis of 961 participants with type 2 diabetes, Heplisav-B demonstrated a statistically significantly higher rate of protection of 90% compared with 65% for Engerix-B.<sup>384</sup> Additional subgroup analyses revealed significantly higher rates of protection among smokers, persons with body mass index  $\geq 30$  kg/m<sup>2</sup>, and older age groups who are typically at risk of suboptimal responses with recombinant vaccines. Adults younger than 40 years had a similar response.<sup>385</sup> Heplisav-B has not been studied in persons infected with HIV.

### Indications for Vaccination

All persons at high risk of acquiring HBV (see “Routes of Transmission” earlier) should be offered vaccination if nonimmune. Recent best-practice advice from the American College of Physicians and CDC recommends the following at-risk patient groups should be vaccinated: (1) all unvaccinated adults (including pregnant women) at risk for infection due to sexual, percutaneous, or mucosal exposure; (2) health care and public safety workers at risk for blood exposure; (3) adults with chronic liver disease and end-stage renal disease (including hemodialysis patients); (4) adults with HIV infection; (5) travelers to HBV-endemic regions; and (6) any adults seeking protection from HBV infection.<sup>386,387</sup> In 2011 the ACIP updated their guidelines to recommend vaccination in all adults age 19 to 59 years with diabetes. Universal vaccination of all infants has been recommended in the United States since 1991, and has since been adopted by many other countries and incorporated into routine childhood immunization programs. As of 2015, 185 (95%) of World Health Organization (WHO) member states vaccinated infants against hepatitis B as part of their vaccination schedules; global coverage with three doses of HBV vaccine during infancy reached 84%. Since the introduction of the HBV vaccine (ranging from the 1980s–2000s, varying by country) and 2015, the prevalence of HBV infection among children younger than 5 years has fallen from 4.7% to 1.3%.<sup>388</sup> In the United States in 2014, 92% of infants had been fully vaccinated with three doses of HBV vaccine.<sup>389</sup> In October 2016 ACIP recommended that all US-born infants who weigh at least 2000 g receive a dose of HepB vaccine before 24 hours of age.<sup>390</sup>

### Dose Regimen

Two single-antigen hepatitis B vaccines are currently licensed in the United States: Engerix-B and Recombivax HB. Engerix-B is formulated to contain 20  $\mu$ g HBsAg/mL, and Recombivax HB contains 10  $\mu$ g HBsAg/mL (Table 145.6). The recommendation for adults is to administer 1 mL of either Engerix-B or Recombivax HB in three doses at months 0, 1 to 2, and then at 6 to 12 months. In infants, three doses of 0.5 mL of vaccine are required to complete the course. For adolescents (11–19 years of age), three doses of 0.5 mL of Recombivax HB or 1 mL of Engerix-B is recommended.<sup>391</sup> Either vaccine can be interchanged during the series of injections. An optional two-dose regimen of 1 mL Recombivax HB has also been approved for adolescents age 11 to 15 years, with a second dose given 4 to 6 months after the first dose.<sup>392</sup> The recently approved Heplisav-B is administered in two doses over 1 month at 0 and 4 weeks, although there are no formal guidelines currently regarding who should receive this vaccine over the other two. However, since safety in pregnancy has not been assessed, pregnant women should receive either Engerix-B or Recombivax HB. Vaccines should be administered intramuscularly because deposition of the vaccine into adipose tissue results in a lower seroconversion rate.<sup>393</sup> Adverse reactions are uncommon, and most consist of soreness at the



**TABLE 145.6 Doses and Schedules of Licensed Hepatitis B Vaccines**

| HEPATITIS B VACCINES    | AGE                                  | DOSE                                | VOLUME             | SCHEDULE   |
|-------------------------|--------------------------------------|-------------------------------------|--------------------|--|
| Engerix-B               | <20 yr                               | 10 µg                               | 10 µg/0.5 mL       | Infants <sup>b</sup> : 1st dose within 24 h of birth, 1–4, 6–18 mo<br>Older children: 0, 1–2, 4 mo       |
|                         | >20 yr                               | 20 µg                               | 20 µg/1 mL         | 0, 1, 6 mo   |
|                         | Diabetes 19–59 yr <sup>c</sup>       | 20 µg                               | 20 µg/1 mL         | 0, 1, 6 mo   |
|                         | Dialysis and other immunocompromised | 40 µg                               | 2–20 µg/1 mL doses | 0, 1, 2, 6 mo  |
| Recombivax HB           | <20 yr                               | 5 µg                                | 5 µg/0.5 mL        | Infants <sup>b</sup> : 1st dose within 24 h of birth, 1–4, 6–18 mo<br>Older children: 0, 1–2, 4 mo       |
|                         | 11–15 yr                             | 10 µg                               | 10 µg/1 mL         | 0, 4–6 mo  |
|                         | >20 yr                               | 10 µg                               | 10 µg/1 mL         | 0, 1, 6 mo   |
|                         | Diabetes 19–59 yr <sup>c</sup>       | 10 µg                               | 10 µg/1 mL         | 0, 1, 6 mo   |
|                         | Dialysis and other immunocompromised | 40 µg <sup>d</sup>                  | 40 µg/1 mL         | 0, 1, 6 mo   |
| Heplisav-B <sup>e</sup> | >18 yr                               | 20 µg                               | 20 µg/0.5 mL doses | 0, 1 mo  |
| COMBINATION VACCINES    | AGE <sup>f</sup>                     | ANTIGEN                             | VOLUME             | SCHEDULE   |
| Pediarix                | 6 wk–6 yr                            | Engerix-B, Infanrix (DTaP), and IPV | 0.5 mL             | For newborns, give monovalent hepatitis B vaccine within 24 h of birth, then give Pediarix at 2, 4, 6 mo |
| Twinrix                 | >18 yr                               | Havrix (HAV) and Engerix-B (20 µg)  | 1 mL               | 0, 1, 6 mo   |

<sup>a</sup>All vaccines should be administered intramuscularly in the deltoid. <https://www.cdc.gov/hepatitis/hbv/hbvfaq.htm#vaccFAQ>.

<sup>b</sup>Infants born to hepatitis B surface antigen (HBsAg)-positive mothers should have hepatitis B immune globulin (HBIG) within 12 h of delivery, along with vaccine at a separate site. If mother's HBsAg status is unknown, administer vaccine within 12 h and test mother. If mother is HBsAg positive, administer HBIG within 1 wk.

<sup>c</sup>Hepatitis B vaccination may be administered at the discretion of the treating clinician to unvaccinated adults with diabetes mellitus who are age ≥60 yr (see <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6050a4.htm>).

<sup>d</sup>Special formulation.

<sup>e</sup>Safety and effectiveness of Heplisav-B have not been established in adults on hemodialysis.

<sup>f</sup>Birth dose should be monovalent vaccine only; subsequent doses can be combination.

DTaP, Diphtheria, tetanus, acellular pertussis (vaccine); IPV, inactivated polio vaccine.

injection site. Low-grade fever, malaise, headache, and myalgias are seen in <1% of vaccinees. The vaccine can be administered during pregnancy.

## Efficacy

A protective level of immune response after vaccination is defined as a titer of anti-HBs of >10 IU/L. Although this was somewhat arbitrary, clinical studies suggest that a decrease in titer below this level is associated with a risk of infection. Both Recombivax HB and Engerix-B administered over three doses, as described previously, result in a protective antibody response in >95% of adults, infants, and children. In persons receiving the newly approved Heplisav-B, 95% of patients exhibited a seroprotective response 8 weeks after the second dose of vaccine.<sup>382</sup> Because the response rate in healthy individuals is so high, routine postvaccination testing in persons receiving Recombivax HB and Engerix-B is not recommended, except among immunocompetent individuals who are at high risk of repeated exposure to HBV, such as health care workers, IV drug users, chronic hemodialysis patients, and individuals who are at risk for recurrent exposure to HBV (e.g., sexual partners of carriers, MSM). Testing should be performed 1 to 2 months after the vaccine series in these individuals. For these individuals who have no or inadequate anti-HBs titers after a primary series, the recommendation is to administer one or more doses of vaccine. After one to two doses, up to 25% of previous nonresponders or hyporesponders may have adequate titers. Almost all healthy adults who do not respond to a primary three-dose series with anti-HBs concentrations of >10 IU/L will respond to a three-dose revaccination series.<sup>394</sup> Postvaccination testing is also recommended in infants born to HBsAg-positive mothers, which should be performed at age 9 to 12 months to avoid detection of anti-HBs from HBIG administered during infancy.<sup>395</sup> Vaccination response is decreased in certain populations, including those older than 40 years; those who are obese, have particular genetic polymorphisms, and have a lack of sleep; and those who smoke, have HIV or any form of immunosuppression, diabetes mellitus, renal failure, chronic liver disease, improper administrations (e.g., administration into the buttock or subcutaneous injection), and freezing of the vaccine.<sup>396,397</sup>

## Efficacy in Patients With Human Immunodeficiency Virus Infection

In patients with HIV response rates vary between 18% and 71%.<sup>398–400</sup> Although a strict correlation between response and immune status is not always found, response is better with CD4 T-cell counts >500 cells/mL.<sup>401,402</sup> US guidelines do not recommend delaying HBV vaccination in patients with HIV infection. HIV-infected persons who are nonresponders to HBV vaccine are not protected from infection and have similar rates of HBV infection as unvaccinated individuals with HIV, unlike immunocompetent individuals who may have some level of protection from the vaccine, even if anti-HBs titers fall below protective levels.<sup>403</sup> This is thought to be due to decreased numbers of antigen-specific memory B cells in HIV-infected individuals. A number of vaccination strategies to boost immunity in HIV-infected individuals have been examined, including accelerated schedules (four vaccine doses at 0, 1, 2, and 6 months),<sup>404</sup> intradermal administration, and rescue vaccination after primary vaccination, with variable success. Another strategy is use of adjuvants as stimulators of immunogenicity of the vaccine, although studies show mixed results.<sup>405–407</sup> Currently DHHS recommends the standard three-dose HBV vaccination, although other vaccines strategies are recommended by some experts, including the use of double-dose vaccine at 0, 1, and 6 months, with or without additional doses.<sup>408</sup>

## Efficacy in End-Stage Renal Disease

In patients on chronic hemodialysis the response rate to recombinant vaccines is 50% to 60%.<sup>409</sup> In addition, anti-HBs levels decline more rapidly compared with healthy individuals.<sup>410</sup> Intradermal injection has been shown to be associated with higher seroconversion rates than intramuscular administration.<sup>411,412</sup> The CDC recommends all end-stage renal disease patients receive double doses of conventional recombinant HBV vaccine, preferably before the onset of dialysis; administration of a fourth vaccine dose may be necessary to increase the seroprotection rate (see Table 145.6). Testing 1 to 2 months after vaccination is recommended for hemodialysis patients to determine their response to the vaccine.<sup>413</sup> They should also undergo yearly anti-HBs

testing and receive a single booster dose of vaccine if the titer falls below 10 IU/L.

### Durability of Response

There is excellent durability of response after a successful primary HBV vaccine series with Recombivax HB and Engerix-B. No long-term durability data are available from Heplisav-B. In a study from Alaska, one of the longest post-HBV vaccine cohorts, 94% of Native Alaskan children and adults who had received plasma-derived HBV vaccine at a minimum of 6 months of age in the early 1980s had evidence of immunity at 22 years of follow-up, even though only 60% of the vaccinated cohort retained detectable levels of anti-HBs. Complete protection was conferred for up to 22 years in those immunized as children and adults, and for up to 15 years in those immunized as infants.<sup>414</sup> A recently published study extended follow-up to 30 years and indicated that >90% of vaccines had evidence of protective immunity, based on anti-HBs levels  $\geq 10$  IU/L or responses to booster doses of vaccine.<sup>415</sup> A companion study indicated that evidence of T-cell immunity to HBV was present even if anti-HBs levels were not detected.<sup>416,417</sup> Long-term data from Taiwan, where 20% of the population is infected with chronic hepatitis B, indicate that the vaccine provides long-term protection up to 20 years in adults.<sup>418</sup> Among HIV-infected individuals the durability of a response to HBV vaccination is significantly associated with higher anti-HBs titers after primary vaccination, with anti-HBs responses >100 IU/L associated with the most prolonged seroprotection.<sup>419</sup> Breakthrough infections have been reported in successfully vaccinated individuals who were subsequently exposed to HBV after a full vaccine series; however, the number of such cases is few, and clinically significant infections are rare.

### Postexposure Immunoprophylaxis

Postexposure prophylaxis with HBIG and vaccine is recommended for all nonimmune individuals who have percutaneous, sexual contact, ocular, or mucous membrane exposure to blood, including human bites that penetrate the skin (see Chapter 316), where the source is known to be at high risk of being HBsAg positive. The first dose of 0.06 mL/kg should be administered as soon as possible, preferably within 12 hours, although there is a window period of up to 24 hours.<sup>420</sup> The first vaccine dose should be given at the same time, although in a different site, followed by the remainder of the series. For individuals who are vaccinated but who do not have documentation of adequate titers of anti-HBs, recommendations are to administer both HBIG and a vaccine booster dose pending documentation of adequate anti-HBs. Individuals who have failed to respond to a vaccine series require two doses of HBIG 1 month apart.

### Prevention of Perinatal Transmission

In women with chronic HBV infection, the risk of transmission of HBV infection to their child without prophylaxis is high and varies according to HBeAg status of mothers from 12% for HBeAg-negative/anti-HBe-positive mothers, to 25% for HBeAg-negative/anti-HBe-negative mothers, to 70% to 90% for HBeAg-positive mothers.<sup>421–423</sup> Maternal serum HBV DNA levels also correlate with the risk of transmission.<sup>424</sup> It is recommended that all neonates born to mothers with chronic hepatitis B receive HBIG 0.5 mL IM at birth and hepatitis B vaccine within 12 hours of birth, followed by vaccine dose at 1 and 6 months, which together are 85% to 95% effective in preventing perinatal HBV infection.<sup>422,423</sup> In a large systematic review and meta-analysis the addition of HBIG decreased risk of transmission from mother to infant compared

with HBV vaccine alone<sup>425</sup>; however, it may not be necessary in HBsAg-positive/HBeAg-negative mothers.<sup>426,427</sup> Administration of a “birth-dose” HBV vaccine (within 12 hours of birth), compared with administration after 12 hours of birth, is associated with further reductions in HBV transmission, although birth-dose coverage remains low in many developing countries (39% in 2016).<sup>428</sup> As of 2016, 97 WHO member states have introduced the hepatitis B birth-dose vaccine.<sup>429</sup> Despite the relative success of immunoprophylaxis, a significant proportion of infants remain at risk for HBV transmission. Ten percent to 30% of infants develop HBV infection when mothers have high levels of viremia (defined as HBV DNA  $>2 \times 5 \log_{10}$  IU/mL).<sup>430,431,432</sup>

Updated 2016 AASLD HBV treatment guidelines now recommend antiviral therapy beginning in the third trimester, to reduce the risk of perinatal transmission of hepatitis B in HBsAg-positive pregnant women with an HBV DNA level  $>200,000$  IU/mL.<sup>250</sup> TDF remains the first-line recommended therapy based on its safety and efficacy, although telbivudine and lamivudine are also recommended. In a recent randomized, placebo-controlled trial, Pan and colleagues<sup>433</sup> evaluated 200 HBeAg-positive mothers with HBV DNA levels  $>2 \times 5 \log_{10}$  IU/mL and compared TDF therapy (300 mg daily) from 30 to 32 weeks of gestation to 4 weeks postpartum versus no drugs (control subjects). All infants received standard immunoprophylaxis at birth. A reduction in mother-to-child transmission for TDF compared with control subjects was found (0% vs. 7%;  $P=.01$ ).<sup>433</sup> Other studies with lamivudine and telbivudine have shown similar effects.<sup>434–436</sup> In a recent systematic review of safety studies with TDF, many found no difference in fetal or maternal adverse effects.<sup>436</sup> Resistant variants have been shown in up to 19% of women receiving lamivudine in prior studies,<sup>437</sup> and these resistant variants rarely occur in women receiving telbivudine.<sup>438</sup> More data are needed on the efficacy and safety of entecavir and TAF in pregnancy before they can be recommended. Additional information is also needed on optimal timing of antiviral therapy during pregnancy, the optimal HBV DNA thresholds to treat patients with antivirals, and resistance patterns in infants with nucleos(t)ide therapy. There is no evidence that cesarean section prevents maternal-infant transmission, and thus routine cesarean section is not recommended.<sup>439</sup> Neither breastfeeding nor amniocentesis appears to increase the risk of transmission.<sup>440</sup>

### Hepatitis B Surface Antigen Escape Mutants

HBs mutants have been described in infants infected with HBV after passive-active vaccination,<sup>441</sup> in vaccinated adults, in liver transplant recipients who have received prolonged courses of HBIG to prevent recurrence of HBV in the allograft,<sup>442,443</sup> and in immunocompromised patients with resolved HBV infection receiving dasatinib.<sup>444,445</sup> The most common mutations are the glycine-to-arginine substitution at codon 145 (G145R) and the sD144A in the “a” determinant sequence of HBsAg, which decrease binding of HBsAg to anti-HBs. The prevalence of these escape mutants is increasing over time, but the clinical and epidemiologic importance and the impact on current vaccination strategies are unclear. Selection of these HBs antigen mutants appears to be a rare event, and their prevalence even in HBV endemic areas is low.<sup>446,447</sup> In a chimpanzee model of infection, the current vaccines appear to protect against the spread of HBs mutants.<sup>448</sup>

### ACKNOWLEDGMENTS

We want to thank Aleem Siddiqui for his contributions to this chapter in the sixth edition of this book and Margaret Koziel for her contributions to the seventh edition.

### Key References

The complete reference list is available online at Expert Consult.

8. Dane DS, Cameron C, Briggs M. Virus-like particles in serum of patients with Australia antigen associated hepatitis. *Lancet*. 1970;2:695–698.
15. Seeger C, Zoulim F, Mason WS. Hepadnaviruses. In: Knipe DM, Howley PM, eds. *Fields Virology*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2013.
22. Yan H, Zhong G, Xu G, et al. Sodium taurocholate cotransporting polypeptide is a functional receptor for human hepatitis B and D virus. *Elife*. 2012;1:e00049.
36. Lau GK, Piratvisuth T, Luo KX, et al. Peginterferon Alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2005;352:2682–2695.
49. Summers J, Mason WS. Replication of the genome of a hepatitis B-like virus by reverse transcription of an RNA intermediate. *Cell*. 1982;29:403–415.
52. Tu T, Budzinska MA, Shackel NA, et al. HBV DNA integration: molecular mechanisms and clinical implications. *Viruses*. 2017;9:pil:E75.
53. Decorsière A, Mueller H, van Breugel PC, et al. Hepatitis B virus X protein identifies the Smc5/6 complex as a host restriction factor. *Nature*. 2016;531:386–389.
70. McMahon BJ, Holck P, Bulkow L, et al. Serologic and clinical outcomes of 1536 Alaska Natives chronically infected with hepatitis B virus. *Ann Intern Med*. 2001;135:759–768.

80. Guidotti LG, Rochford R, Chung J, et al. Viral clearance without destruction of infected cells during acute HBV infection. *Science*. 1999;284:825–829.
98. Kamatani Y, Wattanapokayakit S, Ochi H, et al. A genome-wide association study identifies variants in the HLA-DP locus associated with chronic hepatitis B in Asians. *Nat Genet*. 2009;41:591–595.
102. Sung WK, Zheng H, Li S, et al. Genome-wide survey of recurrent HBV integration in hepatocellular carcinoma. *Nat Genet*. 2012;44:765–769.
110. Schweitzer A, Horn J, Mikolajczyk RT, et al. Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *Lancet*. 2015;386:1546–1555.
115. Centers for Disease Control and Prevention. Surveillance for Viral Hepatitis—United States, 2015. <https://www.cdc.gov/hepatitis/statistics/2015surveillance/commentary.htm>. Accessed December 7, 2017.
- 117a. Centers for Disease Control and Prevention. Hepatitis B Information for Health Professionals: Hepatitis B FAQs for Health Professionals. <https://www.cdc.gov/hepatitis/hbv/hbvfaq.htm>. Accessed December 5, 2017.
121. Rehermann B, Ferrari C, Pasquinelli C, et al. The hepatitis B virus persists for decades after patients' recovery from acute viral hepatitis despite active maintenance of a cytotoxic T lymphocyte response. *Nat Med*. 1996;2:1104–1108.
135. Lin CL, Kao JH. Natural history of acute and chronic hepatitis B: the role of HBV genotypes and mutants. *Best Pract Res Clin Gastroenterol*. 2017;31:249–255.
157. Iloeje UH, Yang HI, Su J, et al. Predicting cirrhosis risk based on the level of circulating hepatitis B viral load. *Gastroenterology*. 2006;130:678–686.
168. Iloeje UH, Yang HI, Jen CL, Risk Evaluation of Viral Load Elevation and Associated Liver Disease/Cancer-Hepatitis B Virus Study Group, et al. Risk and predictors of mortality associated with chronic hepatitis B infection. *Clin Gastroenterol Hepatol*. 2007;5:921–931.
170. Global Burden of Cancer Collaboration; Fitzmaurice C, Allen C, Barber RM, et al. Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 32 cancer groups, 1990 to 2015: a systematic analysis for the Global Burden of Disease Study. *JAMA Oncol*. 2017;3:524–548.
173. Raffetti F, Fattovich G, Donato F. Incidence of hepatocellular carcinoma in untreated subjects with chronic hepatitis B: a systematic review and meta-analysis. *Liver Int*. 2016;36:1239–1251.
- 174a. Yang JD, Gyedu A, Afihene MY, et al. Hepatocellular carcinoma occurs at an earlier age in Africans, particularly in association with chronic hepatitis B. *Am J Gastroenterol*. 2015;110:1629–1631.
177. Chen CJ, Yang HI, Su J, et al. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA*. 2006;295:65–73.
180. Papatheodoridis GV, Chan HLY, Hansen BE, et al. Risk of hepatocellular carcinoma in chronic hepatitis B: assessment and modification with current antiviral therapy. *J Hepatol*. 2015;62:956–967.
182. Tseng TC, Liu CJ, Yang HC, et al. High levels of hepatitis B surface antigen increase risk of hepatocellular carcinoma in patients with low HBV load. *Gastroenterology*. 2012;142:1140–1149.e3.
196. Falade-Nwulia O, Seaberg EC, Snider AE, et al. Incident hepatitis B in HIV-infected and HIV-uninfected men who have sex with men from pre-HAART to HAART periods: a cohort study. *Ann Intern Med*. 2015;163:673–680.
200. Thio CL, Seaberg EC, Skolasky R Jr, et al. HIV-1, hepatitis B virus, and risk of liver-related mortality in the Multicenter Cohort Study (MACS). *Lancet*. 2002;360:1921–1926.
202. Singh K, et al. Hepatitis B virus co-infection: epidemiology, pathogenesis, and treatment. *AIDS*. 2017;31:2035–2052.
223. Choi J, Lim YS. Characteristics, prevention and management of hepatitis B virus reactivation in HBV-infected patients who require immunosuppressive therapy. *J Infect Dis*. 2017;216(S8):S778–S784.
247. Calvaruso V, Ferraro D, Licata A, et al. HBV reactivation in patients with HCV/HBV cirrhosis on treatment with direct-acting antivirals. *J Viral Hepat*. 2018;25:72–79.
250. Terrault NA, Bzowej NH, Chang KM, et al. AASLD guidelines for the treatment of chronic hepatitis B. *Hepatology*. 2016;61:261–283.
264. Marcellin P, Heathcote EJ, Buti M, et al. Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med*. 2008;359:2442–2455.
303. Chang TT, Gish RG, de Man R, et al. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2006;354:1001–1010.
306. McMahon MA, Jilek BL, Brennan TP, et al. The HBV drug entecavir—effects on HIV-1 replication and resistance. *N Engl J Med*. 2007;356:2614–2621.
310. Buti M, Gane E, Seto WK, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of patients with HBeAg-negative chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol*. 2016;1:196–206.
311. Chan HL, Fund S, Seto WK, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of HBeAg-positive chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol*. 2016;1:185–195.
317. Lok AS, Trinh H, Carosi G, et al. Efficacy of entecavir with or without tenofovir disoproxil fumarate for nucleos(t)ide-naïve patients with chronic hepatitis B. *Gastroenterology*. 2012;143:619–628.
340. Fung J, Wong T, Chok K, et al. Long-term outcomes of entecavir therapy for chronic Hepatitis B after liver transplantation: results up to 8 years. *Hepatology*. 2017;66:1036–1044.
354. Department of Health and Human Services; Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents. <http://aidsinfo.nih.gov/contentfiles/lvguidelines/adultandadolescentgl.pdf>. Accessed December 12, 2017.
355. Perrillo RP, Gish R, Falck-Ytter YT. American Gastroenterological Association. Institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology*. 2015;148:221–244.e3.
360. Zhou K, Terrault N. Management of hepatitis B in special populations. *Best Pract Res Clin Gastroenterol*. 2017;31:311–320.
361. Huang H, Li X, Zhu J, et al. Entecavir vs lamivudine for prevention of hepatitis B virus reactivation among patients with untreated diffuse large B-cell lymphoma receiving R-CHOP chemotherapy: a randomized clinical trial. *JAMA*. 2014;312:2521–2530.
374. Fattovich G, Stroffolini T, Zagni I, et al. Hepatocellular carcinoma in cirrhosis: incidence and risk factors. *Gastroenterology*. 2004;127(5 suppl 1):S35–S50.
381. Chang MH, Chen CJ, Lai MS, et al. Universal hepatitis B vaccination in Taiwan and the incidence of hepatocellular carcinoma in children. Taiwan Childhood Hepatoma Study Group. *N Engl J Med*. 1997;336:1855–1859.
384. Jackson S, Lentino J, Kopp J, et al. Immunogenicity of a two-dose investigational hepatitis B vaccine, HBsAg-1018, using a toll-like receptor 9 agonist adjuvant compared with a licensed hepatitis B vaccine in adults. *Vaccine*. 2018;36:668–674. <https://doi.org/10.1016/j.vaccine.2017.12.038>.
386. Abara WE, Qaseem A, Schillie S, et al. Hepatitis B vaccination, screening and linkage to care: best practice advice from the American College of Physicians and Centers for Disease Control and Prevention. *Ann Intern Med*. 2017;167:794–804.
387. Schillie S, Vellozzi C, Reingold A, et al. Prevention of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices. *MMWR Recomm Rep*. 2018;67:1–31.
392. From the Centers for Disease Control and Prevention. Alternate two dose hepatitis B vaccination schedule for adolescents aged 11 to 15 years. *JAMA*. 2000;283:2100.
404. Launay O, van der Vliet D, Rosenberg AR, et al. Safety and immunogenicity of 4 intramuscular double doses and 4 intradermal low doses vs. standard hepatitis B vaccine regimen in adults with HIV-1. *JAMA*. 2011;305:1432–1440.
408. Rey D, Piroth L, Wendling MJ, et al. Safety and immunogenicity of double-dose versus standard-dose hepatitis B revaccination in non-responding adults with HIV-1 (ANRS HB04 B-BOOST): a multicentre, open-label, randomised controlled trial. *Lancet Infect Dis*. 2015;15:1283–1291.
416. Simons BC, Spradling PR, Bruden DJ, et al. A Longitudinal hepatitis B vaccine cohort demonstrates long-lasting hepatitis B virus (HBV) cellular immunity despite loss of antibody against HBV surface antigen. *J Infect Dis*. 2016;214:273–280.
427. Lu Y, Liang XF, Wang FZ, et al. Hepatitis B vaccine alone may be enough for preventing hepatitis B virus transmission in neonates of HBsAg (+)/HBeAg (–) mothers. *Vaccine*. 2017;35:40–45.
431. Patton H, Tran TT. Management of hepatitis B during pregnancy. *Nat Rev Gastroenterol Hepatol*. 2014;11:402–409.
433. Pan CQ, Duan Z, Dai E, et al. China Study Group for the Mother-to-Child Transmission of Hepatitis B. Tenofovir to prevent hepatitis B transmission in mothers with high viral load. *N Engl J Med*. 2016;374:2324–2334.



## References

- Patterson RZ, Klunk J, Fornaciari G, et al. The paradox of HBV evolution as revealed from a 16th century mummy. *PLoS Pathog*. 2018;14:e1006750.
- Lurman A. Eine icterus epidemic. *Berlin Klinische Wochenschrift*. 1885;22:20–23.
- Findlay G, MacCallum F. Note on acute hepatitis and yellow fever immunization. *Trans Soc Trop Med Hyg*. 1937;31:297.
- Bigger J, Dubi S. Jaundice in syphilitics under treatment. *Lancet*. 1943;1:457.
- Flaum A, Malmros H, Persson E. Eine nosocomiale icterus epidemic. *Acta Med Scand Suppl*. 1926;16:544.
- Feinstone S, Kapikian A, Purcell R, et al. Transfusion associated hepatitis not due to viral hepatitis type A or B. *N Engl J Med*. 1975;282:767.
- Blumberg B, Alter H, Visnitch S. A “new” antigen in leukemia sera. *JAMA*. 1965;191:541–546.
- Dane DS, Cameron C, Briggs M. Virus-like particles in serum of patients with Australia antigen associated hepatitis. *Lancet*. 1970;2:695–698.
- Mason WS, Seal G, Summers J. Virus of Pekin ducks with structural and biological relatedness to human hepatitis B virus. *J Virol*. 1980;36:829–836.
- Sninsky JJ, Siddiqui A, Robinson WS, et al. Cloning and endo-nuclease mapping of the hepatitis B viral genome. *Nature*. 1979;279:346–348.
- Galibert F, Mandart E, Fitoussi F, et al. Nucleotide sequence of the hepatitis B virus genome (subtype ayw) cloned in *E. coli*. *Nature*. 1979;281:646–650.
- Valenzuela P, Gray P, Quiroga M, et al. Nucleotide sequence of the gene coding for the major protein of hepatitis B virus surface antigen. *Nature*. 1979;280:815–819.
- Landers TA, Greenberg HB, Robinson WS. Structure of hepatitis B Dane particle DNA and nature of the endogenous DNA polymerase reaction. *J Virol*. 1977;23:368–376.
- Robinson WS, Clayton DA, Greenman RL. DNA of a human hepatitis B virus candidate. *J Virol*. 1974;14:384–391.
- Seeger C, Zoulim F, Mason WS. Hepadnaviruses. In: Knipe DM, Howley PM, eds. *Fields Virology*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2013.
- Roosinck MJ, Siddiqui A. In vivo phosphorylation and protein analysis of hepatitis B virus core antigen. *J Virol*. 1987;61:955–961.
- Albin C, Robinson WS. Protein kinase activity in hepatitis B virus. *J Virol*. 1980;34:297–302.
- Gavilanes F, Gonzalez-Ros JM, Peterson DL. Structure of hepatitis B surface antigen: characterization of the lipid components and their association with the viral proteins. *J Biol Chem*. 1982;257:7770–7777.
- Peterson DL. Isolation and characterization of the major protein and glycoprotein of hepatitis B surface antigen. *J Biol Chem*. 1981;256:6975–6983.
- Gerin JL, Alexander H, Shih JW, et al. Chemically synthesized peptides of hepatitis B surface antigen duplicate the d/y specificities and induce subtype-specific antibodies in chimpanzees. *Proc Natl Acad Sci USA*. 1983;80:2365–2369.
- Rydell GE, Prakash K, Norder H, et al. Hepatitis B surface antigen on subviral particles reduces the neutralizing effect of anti-HBs antibodies on hepatitis B viral particles in vitro. *Virology*. 2017;509:67–70.
- Yan H, Zhong G, Xu G, et al. Sodium taurocholate cotransporting polypeptide is a functional receptor for human hepatitis B and D virus. *Elife*. 2012;1:e00049.
- Glebe D, Urban S, Knoop EV, et al. Mapping of the hepatitis B virus attachment site by use of infection-inhibiting preS1 lipopeptides and tupaia hepatocytes. *Gastroenterology*. 2005;129:234–245.
- Schulze A, Gripon P, Urban S. Hepatitis B virus infection initiates with a large surface protein-dependent binding to heparan sulfate proteoglycans. *Hepatology*. 2007;46:1759–1768.
- Huang HC, Chen CC, Chang WC, et al. Entry of hepatitis B virus into immortalized human primary hepatocytes by clathrin-dependent endocytosis. *J Virol*. 2012;86:9443–9453.
- Rabe B, Vlachou A, Pante N, et al. Nuclear import of hepatitis B virus capsids and release of the viral genome. *Proc Natl Acad Sci USA*. 2003;100:9849–9854.
- Gerlich WH, Robinson WS. Hepatitis B virus contains protein attached to the 5' terminus of its complete DNA strand. *Cell*. 1980;21:801–809.
- Lien JM, Aldrich CE, Mason WS. Evidence that a capped oligoribonucleotide is the primer for duck hepatitis B virus plus-strand DNA synthesis. *J Virol*. 1986;57:229–236.
- Okamoto H, Tsuda F, Sakugawa H, et al. Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J Gen Virol*. 1988;69:2575–2583.
- Lin CL, Kao JH. The clinical implications of hepatitis B virus genotype: recent advances. *J Gastroenterol Hepatol*. 2011;26(suppl 1):123–130.
- Tran TT, Trinh TN, Abe K. New complex recombinant genotype of hepatitis B virus identified in Vietnam. *J Virol*. 2008;82:5657–5663.
- Tatematsu K, Tanaka Y, Kurbanov F, et al. A genetic variant of hepatitis B virus divergent from known human and ape genotypes isolated from a Japanese patient and provisionally assigned to new genotype J. *J Virol*. 2009;83:10538–10547.
- Yang HI, Yeh SH, Chen PJ, et al. Associations between hepatitis B virus genotype and mutants and the risk of hepatocellular carcinoma. *J Natl Cancer Inst*. 2008;100:1134–1143.
- Kao JH, Chen PJ, Lai MY, et al. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology*. 2000;118:554–559.
- Chu CJ, Hussain M, Lok AS. Hepatitis B virus genotype B is associated with earlier HBeAg seroconversion compared with hepatitis B virus genotype C. *Gastroenterology*. 2002;122:1756–1762.
- Lau GK, Piratvisuth T, Luo KX, et al. Peginterferon Alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2005;352:2682–2695.
- Russnak R, Ganem D. Sequences 5' to the polyadenylation signal mediate differential poly(A) site use in hepatitis B viruses. *Genes Dev*. 1990;4:764–776.
- Raney AK, Milich DR, McLachlan A. Characterization of hepatitis B virus major surface antigen gene transcriptional regulatory elements in differentiated hepatoma cell lines. *J Virol*. 1989;63:3919–3925.
- Zhou DX, Yen TS. Differential regulation of the hepatitis B virus surface gene promoters by a second viral enhancer. *J Biol Chem*. 1990;265:20731–20734.
- Lopez-Cabrera M, Letovsky J, Hu KQ, et al. Multiple liver-specific factors bind to the hepatitis B virus core/pregenomic promoter: trans-activation and repression by CCAAT/enhancer binding protein. *Proc Natl Acad Sci USA*. 1990;87:5069–5073.
- Guo W, Chen M, Yen TS, et al. Hepatocyte-specific expression of the hepatitis B virus core promoter depends on both positive and negative regulation. *Mol Cell Biol*. 1993;13:443–448.
- Siddiqui A, Jameel S, Mapoles J. Expression of hepatitis B virus X gene in mammalian cells. *Proc Natl Acad Sci USA*. 1987;84:2513–2517.
- Treinin M, Laub O. Identification of a promoter element located upstream from the hepatitis B virus X gene. *Mol Cell Biol*. 1987;7:545–548.
- Ou JH, Laub O, Rutter WJ. Hepatitis B virus gene function: the precore region targets the core antigen to cellular membranes and causes the secretion of the e antigen. *Proc Natl Acad Sci USA*. 1986;83:1578–1582.
- Visvanathan K, Skinner NA, Thompson AJ, et al. Regulation of Toll-like receptor-2 expression in chronic hepatitis B by the precore protein. *Hepatology*. 2007;45:102–110.
- Bonino F, Brunetto MR, Rizzetto M, et al. Hepatitis B virus unable to secrete e antigen. *Gastroenterology*. 1991;100:1138–1141.
- Gunther S, Meisel H, Reip A, et al. Frequent and rapid emergence of mutated pre-C sequences in HBV from e-antigen positive carriers who seroconvert to anti-HBe during interferon treatment. *Virology*. 1992;187:271–279.
- Radziwill G, Tucker W, Schaller H. Mutational analysis of the hepatitis B virus P gene product: domain structure and RNase H activity. *J Virol*. 1990;64:613–620.
- Summers J, Mason WS. Replication of the genome of a hepatitis B-like virus by reverse transcription of an RNA intermediate. *Cell*. 1982;29:403–415.
- Pollack JR, Ganem D. An RNA stem-loop structure directs hepatitis B virus genomic RNA encapsidation. *J Virol*. 1993;67:3254–3263.
- Wang GH, Zoulim F, Leber EH, et al. Role of RNA in enzymatic activity of the reverse transcriptase of hepatitis B viruses. *J Virol*. 1994;68:8437–8442.
- Tu T, Budzinska MA, Shackel NA, et al. HBV DNA integration: molecular mechanisms and clinical implications. *Viruses*. 2017;9:pii:E75.
- Decorsiere A, Mueller H, van Bruegel PC, et al. Hepatitis B virus X protein identifies the Smc5/6 complex as a host restriction factor. *Nature*. 2016;531:386–389.
- Siddiqui A, Gaynor R, Srinivasan A, et al. Trans-activation of viral enhancers including long terminal repeat of the human immunodeficiency virus by the hepatitis B virus X protein. *Virology*. 1989;169:479–484.
- Benn J, Su F, Doria M, et al. Hepatitis B virus HBx protein induces transcription factor AP-1 by activation of extracellular signal-regulated and c-Jun N-terminal mitogen-activated protein kinases. *J Virol*. 1996;70:4978–4985.
- Waris G, Siddiqui A. Interaction between STAT-3 and HNF-3 leads to the activation of liver-specific hepatitis B virus enhancer 1 function. *J Virol*. 2002;76:2721–2729.
- Rahmani Z, Huh KW, Lasher R, et al. Hepatitis B virus X protein colocalizes to mitochondria with a human voltage-dependent anion channel, HVDAC3, and alters its transmembrane potential. *J Virol*. 2000;74:2840–2846.
- Shirakata Y, Koike K. Hepatitis B virus X protein induces cell death by causing loss of mitochondrial membrane potential. *J Biol Chem*. 2003;278:22071–22078.
- Cross JC, Wen P, Rutter WJ. Transactivation by hepatitis B virus X protein is promiscuous and dependent on mitogen-activated cellular serine/threonine kinases. *Proc Natl Acad Sci USA*. 1993;90:8078–8082.
- Kekule AS, Lauer U, Weiss L, et al. Hepatitis B virus transactivator HBx uses a tumour promoter signalling pathway. *Nature*. 1993;361:742–745.
- Benn J, Schneider RJ. Hepatitis B virus HBx protein activates Ras-GTP complex formation and establishes a Ras, Raf, MAP kinase signaling cascade. *Proc Natl Acad Sci USA*. 1994;91:10350–10354.
- Elmore LW, Hancock AR, Chang SF, et al. Hepatitis B virus X protein and p53 tumor suppressor interactions in the modulation of apoptosis. *Proc Natl Acad Sci USA*. 1997;94:14707–14712.
- Ng SA, Lee C. Hepatitis B virus X gene and hepatocarcinogenesis. *J Gastroenterol*. 2011;46:974–990.
- Huovila AP, Eder AM, Fuller SD. Hepatitis B surface antigen assembles in a post-ER, pre-Golgi compartment. *J Cell Biol*. 1992;118:1305–1320.
- European Association for the Study of the Liver. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol*. 2017;67:370–398.
- Lok AS, Lai CL, Wu PC, et al. Spontaneous hepatitis B e antigen to antibody seroconversion and reversion in Chinese patients with chronic hepatitis B virus infection. *Gastroenterology*. 1987;92:1839–1843.
- Chang MH, Hsu HY, Hsu HC, et al. The significance of spontaneous hepatitis B e antigen seroconversion in childhood: with special emphasis on the clearance of hepatitis B e antigen before 3 years of age. *Hepatology*. 1995;22:1387–1392.
- Yang PL, Althage A, Chung J, et al. Hydrodynamic injection of viral DNA: a mouse model of acute hepatitis B virus infection. *Proc Natl Acad Sci USA*. 2002;99:13825–13830.
- Liaw YF, Chu CM, Lin DY, et al. Age-specific prevalence and significance of hepatitis B e antigen and antibody in chronic hepatitis B virus infection in Taiwan: a comparison among asymptomatic carriers, chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma. *J Med Virol*. 1984;13:385–391.
- McMahon BJ, Holck P, Bulkow L, et al. Serologic and clinical outcomes of 1536 Alaska Natives chronically infected with hepatitis B virus. *Ann Intern Med*. 2001;135:759–768.
- Fattovich G, Brollo L, Giustina G, et al. Natural history and prognostic factors for chronic hepatitis type B. *Gut*. 1991;32:294–298.
- Tsai SL, Chen PJ, Lai MY, et al. Acute exacerbations of chronic type B hepatitis are accompanied by increased T cell responses to hepatitis B core and e antigens: implications for hepatitis B e antigen seroconversion. *J Clin Invest*. 1992;89:87–96.
- Schulte-Frohlinde E, Foster GR. Spontaneous seroconversion in chronic hepatitis B: role of mutations in the precore/core gene. *Dig Dis Sci*. 1998;43:1714–1718.
- Fattovich G, Ruggie M, Brollo L, et al. Clinical, virologic and histologic outcome following seroconversion from HBeAg to anti-HBe in chronic hepatitis type B. *Hepatology*. 1986;6:167–172.
- Liaw YF, Lin SM, Sheen IS, et al. Acute hepatitis C virus super-infection followed by spontaneous HBeAg seroconversion and HBsAg elimination. *Infection*. 1991;19:250–251.
- Ferrari C, Missale G, Boni C, et al. Immunopathogenesis of hepatitis B. *J Hepatol*. 2003;39(suppl 1):S36–S42.
- Chang JJ, Lewin SR. Immunopathogenesis of hepatitis B virus infection. *Immunol Cell Biol*. 2007;85:16–23.
- Wieland S, Thimme R, Purcell RH, et al. Genomic analysis of the host response to hepatitis B virus infection. *Proc Natl Acad Sci USA*. 2004;101:6669–6674.
- Dunn C, Peppas D, Khanna P, et al. Temporal analysis of early immune responses in patients with acute hepatitis B virus infection. *Gastroenterology*. 2009;137:1289–1300.
- Guidotti LG, Rochford R, Chung J, et al. Viral clearance without destruction of infected cells during acute HBV infection. *Science*. 1999;284:825–829.
- Baron JL, Gardiner L, Nishimura S, et al. Activation of a non-classical NKT cell subset in a transgenic mouse

- model of hepatitis B virus infection. *Immunity*. 2002;16:583–594.
82. Ferrari C, Bertoletti A, Penna A, et al. Identification of immunodominant T cell epitopes of the hepatitis B virus nucleocapsid antigen. *J Clin Invest*. 1991;88:214–222.
  83. Penna A, Chisari FV, Bertoletti A, et al. Cytotoxic T lymphocytes recognize an HLA-A2-restricted epitope within the hepatitis B virus nucleocapsid antigen. *J Exper Med*. 1991;174:1565–1570.
  84. Penna A, Artini M, Cavalli A, et al. Long-lasting memory T cell responses following self-limited acute hepatitis B. *J Clin Invest*. 1996;98:1185–1194.
  85. Guidotti LG, Chisari FV. Noncytolytic control of viral infections by the innate and adaptive immune response. *Annu Rev Immunol*. 2001;19:65–91.
  86. Guidotti LG, Chisari FV. Cytokine-induced viral purging—role in viral pathogenesis. *Curr Opin Microbiol*. 1999;2:388–391.
  87. Stross L, Gunther J, Gasteiger G, et al. Foxp3+ regulatory T cells protect the liver from immune damage and compromise virus control during acute, experimental hepatitis B virus infection. *Hepatology*. 2012;56:873–883.
  88. Milich DR, Schodel F, Peterson DL, et al. Characterization of self-reactive T cells that evade tolerance in hepatitis B e antigen transgenic mice. *Eur J Immunol*. 1995;25:1663–1672.
  89. Bertoletti A, Sette A, Chisari FV, et al. Natural variants of cytotoxic epitopes are T-cell receptor antagonists for antiviral cytotoxic T cells. *Nature*. 1994;369:407–410.
  90. Barnaba V, Franco A, Alberti A, et al. Recognition of hepatitis B virus envelope proteins by liver-infiltrating lymphocytes in chronic HBV infection. *J Immunol*. 1989;143:2650–2655.
  91. Barnaba V, Franco A, Alberti A, et al. Selective killing of hepatitis B envelope antigen-specific B cells by class I-restricted, exogenous antigen-specific T lymphocytes. *Nature*. 1990;345:258–260.
  92. Maini MK, Boni C, Lee CK, et al. The role of virus-specific CD8(+) cells in liver damage and viral control during persistent hepatitis B virus infection. *J Exp Med*. 2000;191:1269–1280.
  93. Fiscaro P, Valdatta C, Massari M, et al. Antiviral intrahepatic T-cell responses can be restored by blocking programmed death-1 pathway in chronic hepatitis B. *Gastroenterology*. 2010;138:682–693.
  94. Thurs MR, Kwiatkowski D, Allsopp CE, et al. Association between an MHC class II allele and clearance of hepatitis B virus in the Gambia. *N Engl J Med*. 1995;332:1065–1069.
  95. Frodsham AJ. Host genetics and the outcome of hepatitis B viral infection. *Transpl Immunol*. 2005;14:183–186.
  96. Li Y, Si L, Zhai Y, et al. Genome-wide association study identified 8p21.3 associated with persistent hepatitis B virus infection among Chinese. *Nat Commun*. 2016;7:11664.
  97. Thio CL, Astemborski J, Thomas R, et al. Interaction between RANTES promoter variant and CCR5Delta32 favors recovery from hepatitis B. *J Immunol*. 2008;181:7944–7947.
  98. Kamatani Y, Watanapokayakit S, Ochi H, et al. A genome-wide association study identifies variants in the HLA-DP locus associated with chronic hepatitis B in Asians. *Nat Genet*. 2009;41:591–595.
  99. Chisari FV. Rous-Whipple Award Lecture. Viruses, immunity, and cancer: lessons from hepatitis B. *Am J Pathol*. 2000;156:1117–1132.
  100. Beasley RP, Hwang LY, Lin CC, et al. Hepatocellular carcinoma and hepatitis B virus: a prospective study of 22,707 men in Taiwan. *Lancet*. 1981;2:1129–1133.
  101. Cougot D, Neuveut C, Buendia MA. HBV induced carcinogenesis. *J Clin Virol*. 2005;34(suppl 1):S75–S78.
  102. Sung WK, Zheng H, Li S, et al. Genome-wide survey of recurrent HBV integration in hepatocellular carcinoma. *Nat Genet*. 2012;44:765–769.
  103. Buendia MA. Hepatitis B viruses and cancerogenesis. *Biomed Pharmacother*. 1998;52:34–43.
  104. Murakami Y, Saigo K, Takashima H, et al. Large scaled analysis of hepatitis B virus (HBV) DNA integration in HBV related hepatocellular carcinomas. *Gut*. 2005;54:1162–1168.
  105. Caselmann W, Koshy R. *Transactivators of HBV, Signal Transduction and Tumorigenesis*. London: Imperial College Press; 1998.
  106. Zoulim F, Saputelli J, Seeger C. Woodchuck hepatitis virus X protein is required for viral infection in vivo. *J Virol*. 1994;68:2026–2030.
  107. Leverro M, Zucman-Rossi J. Mechanisms of HBV-induced hepatocellular carcinoma. *J Hepatol*. 2016;64(suppl 1):S84–S101.
  108. Brunetto MR, Giarin MM, Oliveri F, et al. Wild-type and e antigen-minus hepatitis B viruses and course of chronic hepatitis. *Proc Natl Acad Sci USA*. 1991;88:4186–4190.
  109. Hussain S, Hofseth L, Harris C. Radical causes of cancer. *Nat Rev*. 2003;3:276–285.
  110. Schweitzer A, Horn J, Mikolajczyk RT, et al. Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *Lancet*. 2015;386:1546–1555.
  111. World Health Organization. WHO hepatitis B factsheet. <https://www.who.int/en/news-room/fact-sheets/detail/hepatitis-b>.
  112. Radun D. Epidemiology of hepatitis B in Germany. [http://www.vhpb.org/files/html/Meetings\\_and\\_publications/Presentations/S34Radunberlin03.pdf](http://www.vhpb.org/files/html/Meetings_and_publications/Presentations/S34Radunberlin03.pdf).
  113. van Houdt R, Bruisten SM, Speksnijder AG, et al. Unexpectedly high proportion of drug users and men having sex with men who develop chronic hepatitis B infection. *J Hepatol*. 2012;57:529–533.
  114. McMahon BJ, Alward WL, Hall DB, et al. Acute hepatitis B virus infection: relation of age to the clinical expression of disease and subsequent development of the carrier state. *J Infect Dis*. 1985;151:599–603.
  115. Centers for Disease Control and Prevention. Surveillance for Viral Hepatitis—United States, 2015. <https://www.cdc.gov/hepatitis/statistics/2015surveillance/commentary.htm>. Accessed December 7, 2017.
  116. Kowdley KV, Wang CC, Welch S, et al. Prevalence of chronic hepatitis B among foreign-born persons living in the United States by country of origin. *Hepatology*. 2012;56:422–433.
  117. Gerberding JL. Incidence and prevalence of human immunodeficiency virus, hepatitis B virus, hepatitis C virus, and cytomegalovirus among health care personnel at risk for blood exposure: final report from a longitudinal study. *J Infect Dis*. 1994;170:1410–1417.
  - 117a. Centers for Disease Control and Prevention. Hepatitis B Information for Health Professionals: Hepatitis B FAQs for Health Professionals. <https://www.cdc.gov/hepatitis/hbv/hbvfaq.htm>. Accessed December 5, 2017.
  118. Hoofnagle JH. Type B hepatitis: virology, serology and clinical course. *Semin Liver Dis*. 1981;1:7–14.
  119. Lee WM. Medical progress: hepatitis B virus infection. *N Engl J Med*. 1997;337:1733–1745.
  120. Du X, Liu Y, Ma L, et al. Virological and serological features of acute hepatitis B in adults. *Medicine (Baltimore)*. 2017;96:e6088.
  121. Reherrmann B, Ferrari C, Pasquini C, et al. The hepatitis B virus persists for decades after patients' recovery from acute viral hepatitis despite active maintenance of a cytotoxic T lymphocyte response. *Nat Med*. 1996;2:1104–1108.
  122. Al-Taie OH, Mork H, Gassel AM, et al. Prevention of hepatitis B flare-up during chemotherapy using lamivudine: case report and review of the literature. *Ann Hematol*. 1999;78:247–249.
  123. Bessesen M, Ives D, Condrey L, et al. Chronic active hepatitis B exacerbations in human immunodeficiency virus-infected patients following development of resistance to or withdrawal of lamivudine. *Clin Infect Dis*. 1999;28:1032–1035.
  124. Schiodt FV, Atillasoy E, Shakil AO, et al. Etiology and outcome for 295 patients with acute liver failure in the United States. *Liver Transpl Surg*. 1999;5:29–34.
  125. Wai CT, Fontana RJ, Polson J, et al. Clinical outcome and virological characteristics of hepatitis B-related acute liver failure in the United States. *J Viral Hepat*. 2005;12:192–198.
  126. Sako A, Yasunaga H, Horiguchi H, et al. Acute hepatitis B in Japan: incidence, clinical practices and health policy. *Hepatol Res*. 2011;41:39–45.
  127. Arteaga-Rodriguez A, Carrasco-Garrido P, Lopez de Andres A, et al. Trends of acute hepatitis B hospitalizations, comorbidities, fatality rate, and costs associated with the hospitalization in Spain (2001–2006). *Eur J Gastroenterol Hepatol*. 2010;22:961–966.
  128. Liaw YF, Chu CM. Hepatitis B virus infection. *Lancet*. 2009;373:582–592.
  129. Umemura T, Tanaka E, Kiyosawa K, et al. Japan de novo Hepatitis B Research Group. Mortality secondary to fulminant hepatic failure in patients with prior resolution of hepatitis B virus infection in Japan. *Clin Infect Dis*. 2008;47:e52–e56.
  130. Mindikoglu AL, Regav A, Schiff ER. Hepatitis B virus reactivation after cytotoxic chemotherapy: the disease and its prevention. *Clin Gastroenterol Hepatol*. 2006;4:1076–1081.
  131. Hayashi K, Ishigami M, Ishizu Y, et al. Clinical characteristics and molecular analysis of hepatitis B virus reactivation in hepatitis B surface antigen-negative patients during or after immunosuppressive or cytotoxic chemotherapy. *J Gastroenterol*. 2016;51:1081–1089.
  132. Wu JC, Chen CL, Hou MC, et al. Multiple viral infection as the most common cause of fulminant and subfulminant viral hepatitis in an area endemic for hepatitis B: application and limitations of the polymerase chain reaction. *Hepatology*. 1994;19:836–840.
  133. Wright TL, Mamish D, Combs C, et al. Hepatitis B virus and apparent fulminant non-A, non-B hepatitis. *Lancet*. 1992;339:952–955.
  134. Liang TJ, Hasegawa K, Munoz SJ, et al. Hepatitis B virus precore mutation and fulminant hepatitis in the United States: a polymerase chain reaction-based assay for the detection of specific mutation. *J Clin Invest*. 1994;93:550–555.
  135. Lin CL, Kao JH. Natural history of acute and chronic hepatitis B: the role of HBV genotypes and mutants. *Best Pract Res Clin Gastroenterol*. 2017;31:249–255.
  136. Liu CJ, Chen PJ, Lai MY, et al. A prospective study characterizing full-length hepatitis B virus genomes during acute exacerbation. *Gastroenterology*. 2003;124:80–90.
  137. Chu CM, Liaw YF. Hepatitis B virus-related cirrhosis: natural history and treatment. *Semin Liver Dis*. 2006;26:142–152.
  138. Gerber MA, Hadziyannis S, Vissoulis C, et al. Electron microscopy and immunoelectronmicroscopy of cytoplasmic hepatitis B antigen in hepatocytes. *Am J Pathol*. 1974;75:489–502.
  139. Harrison RF, Davies MH, Goldin RD, et al. Recurrent hepatitis B in liver allografts: a distinctive form of rapidly developing cirrhosis. *Histopathology*. 1993;23:21–28.
  140. Carman WF, Jacyna MR, Hadziyannis S, et al. Mutation preventing formation of hepatitis B e antigen in patients with chronic hepatitis B infection. *Lancet*. 1989;2:588–591.
  141. Fattovich G, Brolo L, Alberti A, et al. Long-term follow-up of anti-HBe-positive chronic active hepatitis B. *Hepatology*. 1988;8:1651–1654.
  142. Gunther S, Sommer G, Von Breunig F, et al. Amplification of full-length hepatitis B virus genomes from samples from patients with low levels of viremia: frequency and functional consequences of PCR-introduced mutations. *J Clin Microbiol*. 1998;36:531–538.
  143. Hawkins AE, Gilson RJ, Bickerton EA, et al. Conservation of precore and core sequences of hepatitis B virus in chronic viral carriers. *J Med Virol*. 1994;43:5–12.
  144. Tong SP, Li JS, Vitvitski L, et al. Evidence for a base-paired region of hepatitis B virus pregenome encapsidation signal which influences the patterns of precore mutations abolishing HBe protein expression. *J Virol*. 1993;67:5651–5655.
  145. Hadziyannis SJ, Vassilopoulos D. Immunopathogenesis of hepatitis B e antigen negative chronic hepatitis B infection. *Antiviral Res*. 2001;52:91–98.
  146. Alexopoulou A, Karayiannis P. HBeAg negative variants and their role in the natural history of chronic hepatitis B virus infection. *World J Gastroenterol*. 2014;20:7644–7652.
  147. Zarski JP, Marcellin P, Cohard M, et al. Comparison of anti-HBe-positive and HBe-antigen-positive chronic hepatitis B in France. French Multicentre Group. *J Hepatol*. 1994;20:636–640.
  148. Di Marco V, Lo Iacono O, Camma C, et al. The long-term course of chronic hepatitis B. *Hepatology*. 1999;30:257–264.
  149. Hsu YS, Chien RN, Yeh CT, et al. Long-term outcome after spontaneous HBeAg seroconversion in patients with chronic hepatitis B. *Hepatology*. 2002;35:1522–1527.
  150. Chan HL, Hussain M, Lok AS. Different hepatitis B virus genotypes are associated with different mutations in the core promoter and precore regions during hepatitis B e antigen seroconversion. *Hepatology*. 1999;29:976–984.
  151. Fattovich G, Giustina G, Schalm SW, et al. Occurrence of hepatocellular carcinoma and decompensation in western European patients with cirrhosis type B. The EUROHEP Study Group on Hepatitis B Virus and Cirrhosis. *Hepatology*. 1995;21:77–82.
  152. Moreno-Otero R, Garcia-Monzon C, Garcia-Sanchez A, et al. Development of cirrhosis after chronic type B hepatitis: a clinicopathologic and follow-up study of 46 HBeAg-positive asymptomatic patients. *Am J Gastroenterol*. 1991;86:560–564.
  153. Chen YC, Sheen IS, Chu CM, et al. Prognosis following spontaneous HBsAg seroclearance in chronic hepatitis B patients with or without concurrent infection. *Gastroenterology*. 2002;123:1084–1089.
  154. Villeneuve JP, Desrochers M, Infante-Rivard C, et al. A long-term follow-up study of asymptomatic hepatitis B surface antigen-positive carriers in Montreal. *Gastroenterology*. 1994;106:1000–1005.
  155. Szpakowski JL, Tucker LY. Causes of death in patients with hepatitis B: a natural history cohort study in the United States. *Hepatology*. 2013;58:21–30.
  156. Crook PD, Jones ME, Hall AJ. Mortality of hepatitis B surface antigen-positive blood donors in England and Wales. *Int J Epidemiol*. 2003;32:118–124.



157. Iloeje UH, Yang HI, Su J, et al. Predicting cirrhosis risk based on the level of circulating hepatitis B viral load. *Gastroenterology*. 2006;130:678–686.
158. Tseng TC, Liu CJ, Yang HC, et al. Higher proportion of viral basal core promoter mutant increases the risk of liver cirrhosis in hepatitis B carriers. *Gut*. 2015;64:292–302.
159. Liu CJ, Chen BF, Chen PJ, et al. Role of hepatitis B virus precore/core promoter mutations and serum viral load on noncirrhotic hepatocellular carcinoma: a case-control study. *J Infect Dis*. 2006;194:594–599.
160. Donato F, Tagger A, Gelatti U, et al. Alcohol and hepatocellular carcinoma: the effect of lifetime intake and hepatitis virus infections in men and women. *Am J Epidemiol*. 2002;155:323–331.
161. Liaw YF. Hepatitis C virus superinfection in patients with chronic hepatitis B virus infection. *J Gastroenterol*. 2002;37(suppl 13):S65–S68.
162. Crespo J, Lozano JL, de la Cruz F, et al. Prevalence and significance of hepatitis C viremia in chronic active hepatitis B. *Am J Gastroenterol*. 1994;89:1147–1151.
163. Fattovich G, Boscaro S, Noventa F, et al. Influence of hepatitis delta virus infection on progression to cirrhosis in chronic hepatitis type B. *J Infect Dis*. 1987;155:931–935.
164. Readli G, Fattovich G, Hadziyannis S, et al. Survival and prognostic factors in 366 patients with compensated cirrhosis type B: a multicenter study. The Investigators of the European Concerted Action on Viral Hepatitis (EUROHEP). *J Hepatol*. 1994;21:656–666.
165. Liaw YF, Lin DY, Chen TJ, et al. Natural course after the development of cirrhosis in patients with chronic type B hepatitis: a prospective study. *Liver*. 1989;9:235–241.
166. Chen YC, Chu CM, Yeh CT, et al. Natural course following the onset of cirrhosis in patients with chronic hepatitis B: a long-term follow-up study. *Hepatol Int*. 2007;1:267–273.
167. Fattovich G, Pantalena M, Zagni I, et al. Effect of hepatitis B and C virus infections on the natural history of compensated cirrhosis: a cohort study of 297 patients. *Am J Gastroenterol*. 2002;97:2886–2895.
168. Iloeje UH, Yang HI, Jen CL, et al. Risk Evaluation of Viral Load Elevation and Associated Liver Disease/Cancer-Hepatitis B Virus Study Group. Risk and predictors of mortality associated with chronic hepatitis B infection. *Clin Gastroenterol Hepatol*. 2007;5:921–931.
169. de Jongh FE, Janssen HL, de Man RA, et al. Survival and prognostic indicators in hepatitis B surface antigen-positive cirrhosis of the liver. *Gastroenterology*. 1992;103:1630–1635.
170. Global Burden of Cancer Collaboration; Fitzmaurice C, Allen C, Barber RM, et al. Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 32 cancer groups, 1990 to 2015: a systematic analysis for the Global Burden of Disease Study. *JAMA Oncol*. 2017;3:524–548.
171. Chen CJ, Chen DS. Interaction of hepatitis B virus, chemical carcinogen, and genetic susceptibility: multistage hepatocarcinogenesis with multifactorial etiology. *Hepatology*. 2002;36:1046–1049.
172. Beasley RP. Hepatitis B virus. The major etiology of hepatocellular carcinoma. *Cancer*. 1988;61:1942–1956.
173. Raffetti E, Fattovich G, Donato F. Incidence of hepatocellular carcinoma in untreated subjects with chronic hepatitis B: a systematic review and meta-analysis. *Liver Int*. 2016;36:1239–1251.
174. Lai MY, Chen DS, Chen PJ, et al. Status of hepatitis B virus DNA in hepatocellular carcinoma: a study based on paired tumor and nontumor liver tissues. *J Med Virol*. 1988;25:249–258.
- 174a. Yang JD, Gyedu A, Afihene MY, et al. Hepatocellular carcinoma occurs at an earlier age in Africans, particularly in association with chronic hepatitis B. *Am J Gastroenterol*. 2015;110:1629–1631.
175. Kramvis A. Molecular characterisation of the genotypes and mutants of hepatitis B virus from South Africa. *S Afr J Epidemiol Infect*. 2008;23:29–32.
176. Yang HI, Yeh SH, Chen PJ, et al. Associations between hepatitis B virus genotype and mutants and the risk of hepatocellular carcinoma. *J Natl Cancer Inst*. 2008;100:1134–1143.
177. Chen CJ, Yang HI, Su J, et al. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA*. 2006;295:65–73.
178. Huang Y, Wang Z, An S, et al. Role of hepatitis B virus genotypes and quantitative HBV DNA in metastasis and recurrence of hepatocellular carcinoma. *J Med Virol*. 2008;80:591–597.
179. Yu MW, Shih WL, Lin CL, et al. Body-mass index and progression of hepatitis B: a population-based cohort study in men. *J Clin Oncol*. 2008;26:5576–5582.
180. Papatheodoridis GV, Chan HLY, Hansen BE, et al. Risk of hepatocellular carcinoma in chronic hepatitis B: assessment and modification with current antiviral therapy. *J Hepatol*. 2015;62:956–967.
181. Iloeje UH, Yang HI, Chen CJ. Natural history of chronic hepatitis B: what exactly has REVEAL revealed? *Liver Int*. 2012;32:1333–1341.
182. Tseng TC, Liu CJ, Yang HC, et al. High levels of hepatitis B surface antigen increase risk of hepatocellular carcinoma in patients with low HBV load. *Gastroenterology*. 2012;142:1140–1149.e3.
183. Yuen MF, Wong DK, Fung J, et al. HBsAg seroclearance in chronic hepatitis B in Asian patients: replicative level and risk of hepatocellular carcinoma. *Gastroenterology*. 2008;135:1192–1199.
184. Cacoub P, Saadoun D, Bourlière M, et al. Hepatitis B virus genotypes and extrahepatic manifestations. *J Hepatol*. 2005;43:764–770.
185. Guillemin L, Mahr A, Callard P, et al. Hepatitis B virus-associated polyarteritis nodosa: clinical characteristics, outcome, and impact of treatment in 115 patients. *Medicine (Baltimore)*. 2005;84:313–322.
186. Avaris E, Savas B, Tozun N, et al. Successful treatment of polyarteritis nodosa related to hepatitis B virus with interferon alpha as first-line therapy. *J Hepatol*. 1998;28:525–526.
187. Simsek H, Telatar H. Successful treatment of hepatitis B virus-associated polyarteritis nodosa by interferon alpha alone. *J Clin Gastroenterol*. 1995;20:263–265.
188. Cacoub P, Terrier B. Hepatitis B-related autoimmune manifestations. *Rheum Dis Clin North Am*. 2009;35:125–137.
189. Guillemin L, Lhote F, Cohen P, et al. Polyarteritis nodosa related to hepatitis B virus. A prospective study with long-term observation of 41 patients. *Medicine (Baltimore)*. 1995;74:238–253.
190. Mazzaio C, Dal Maso L, Urraro T, et al. Hepatitis B virus related cryoglobulinemic vasculitis: a multicentre open label study from the Gruppo Italiano di Studio delle Crioglobulinemie—GISC. *Dig Liver Dis*. 2016;48:780–784.
191. Viganò M, Martin P, Cappelletti M, et al. HBV-associated cryoglobulinemic vasculitis: remission after antiviral therapy with entecavir. *Kidney Blood Press Res*. 2014;39:65–73.
192. Gilbert RD, Wiggelinkhuizen J. The clinical course of hepatitis B virus-associated nephropathy. *Pediatr Nephrol*. 1994;8:11–14.
193. Lhotta K. Beyond hepatorenal syndrome: glomerulonephritis in patients with liver disease. *Semin Nephrol*. 2002;22:302–308.
194. Fabrizi F, Dixit V, Martin P. Meta-analysis: anti-viral therapy of hepatitis B virus-associated glomerulonephritis. *Aliment Pharmacol Ther*. 2006;24:781–788.
195. Kellerman SE, Hanson DL, McNaghten AD, et al. Prevalence of chronic hepatitis B and incidence of acute hepatitis B infection in human immunodeficiency virus-infected subjects. *J Infect Dis*. 2003;188:571–577.
196. Falade-Nwulia O, Seaberg EC, Snider AE, et al. Incident hepatitis B in HIV-infected and HI-uninfected men who have sex with men from pre-HAART to HAART periods: a cohort study. *Ann Intern Med*. 2015;163:673–680.
197. Chun HM, Fieberg AM, Hulsiek KH, et al. Infectious Disease Clinical Research Program HIV Working Group. Epidemiology of hepatitis B virus infection in a US cohort of HIV-infected individuals during the past 20 years. *Clin Infect Dis*. 2010;50:426–436.
198. Thio CL. Hepatitis B and human immunodeficiency virus coinfection. *Hepatology*. 2009;49(suppl):S138–S145.
199. Gilson RJ, Hawkins AE, Beecham MR, et al. Interactions between HIV and hepatitis B virus in homosexual men: effects on the natural history of infection. *AIDS*. 1997;11:597–606.
200. Thio CL, Seaberg EC, Skolasky R Jr, et al. HIV-1, hepatitis B virus, and risk of liver-related mortality in the Multicenter Cohort Study (MACS). *Lancet*. 2002;360:1921–1926.
201. Coffin CS, Osiowy C, Myers RP, et al. Virology and clinical sequelae of long-term antiviral therapy in a North American cohort of hepatitis B virus (HBV)/human immunodeficiency virus type 1 (HIV-1) co-infected patients. *J Clin Virol*. 2013;57:103–108.
202. Singh K, et al. Hepatitis B virus co-infection: epidemiology, pathogenesis, and treatment. *AIDS*. 2017;31:2035–2052.
203. Sun H-Y, Sheng W-H, Tsai M-S, et al. Hepatitis B virus coinfection in human immunodeficiency virus-infected patients: a review. *World J Gastroenterol*. 2014;20:14598–14614.
204. Thio CL, Smeaton L, Saulynas M, et al. Characterization of HIV-HBV coinfection in a multinational HIV-infected cohort. *AIDS*. 2013;27:191–201.
205. Clifford GM, Rickenbach M, Polesel J, et al. Swiss HIV Cohort. Influence of HIV-related immunodeficiency on the risk of hepatocellular carcinoma. *AIDS*. 2008;22:2135–2141.
206. Altfeld M, Rockstroh JK, Addo M, et al. Reactivation of hepatitis B in a long-term anti-HBs-positive patient with AIDS following lamivudine withdrawal. *J Hepatol*. 1998;29:306–309.
207. Witt MD, Lewis RJ, Rieg G, et al. Predictors of the Isolated Hepatitis B core antibody pattern in HIV-infected and -uninfected men in the multicenter AIDS cohort study. *Clin Infect Dis*. 2013;56:606–612.
208. Santos EA, Yoshida CF, Rolla VC, et al. Frequent occult hepatitis B virus infection in patients infected with human immunodeficiency virus type 1. *Eur J Clin Microbiol Infect Dis*. 2003;22:92–98.
209. Shire NJ, Rouster SD, Stanford SD, et al. The prevalence and significance of occult hepatitis B virus in a prospective cohort of HIV-infected patients. *J Acquir Immune Defic Syndr*. 2007;44:309–314.
210. Tsui JJ, French AL, Seaberg EC, et al. Prevalence and long-term effects of occult hepatitis B virus infection in HIV-infected women. *Clin Infect Dis*. 2007;45:736–740.
211. Chazouillères O, Mamish D, Kim M, et al. “Occult” hepatitis B virus as source of infection in liver transplant recipients. *Lancet*. 1994;343:142–146.
212. Bohorquez HE, Cohen AJ, Girgrah N, et al. Liver transplantation in hepatitis B core-negative recipients using livers from hepatitis B core-positive donors: a 13-year experience. *Liver Transpl*. 2013;19:611–618.
213. Yen RD, Bonatti H, Mendez J, et al. Case report of lamivudine-resistant hepatitis B virus infection post liver transplantation from a AT B core antibody donor. *Am J Transplant*. 2006;6:1077–1083.
214. Skagen CL, Jou JH, Said A. Risk of de novo hepatitis in liver recipients from hepatitis B core antibody-positive grafts—a systematic analysis. *Clin Transplant*. 2011;25:E243–E249.
215. Yen RD, Bonatti H, Mendez J, et al. Case report of lamivudine-resistant hepatitis B virus infection post liver transplantation from a hepatitis B core antibody donor. *Am J Transplant*. 2006;6:1077–1083.
216. Todo S, Demetris AJ, Van Thiel D, et al. Orthotopic liver transplantation for patients with hepatitis B virus-related liver disease. *Hepatology*. 1991;13:619–626.
217. Marzano A, Gaia S, Ghisetti V, et al. Viral load at the time of liver transplantation and risk of hepatitis B virus recurrence. *Liver Transpl*. 2005;11:402–409.
218. Saab S, Waterman B, Chi AC, et al. Comparison of different immunoprophylaxis regimens after liver transplantation with hepatitis B core antibody-positive donors: a systematic review. *Liver Transpl*. 2010;16:300–307.
219. Steinmuller T, Seehofer D, Rayes N, et al. Increasing applicability of liver transplantation for patients with hepatitis B-related liver disease. *Hepatology*. 2002;35:1528–1535.
220. Mahboobi N, Tabatabaei SV, Blum HE, et al. Renal grafts from anti-hepatitis B core-positive donors: a quantitative review of the literature. *Transplant Infect Dis*. 2012;14:445–451.
221. Kanaan N, Kabamba B, Maréchal C, et al. Significant rate of hepatitis B reactivation following kidney transplantation in patients with resolved infection. *J Clin Virol*. 2012;55:233–238.
222. Villamil F, Cairo F. Solid organ transplantation and hepatitis B virus. *Clin Liver Dis*. 2013;23:38–40.
223. Choi J, Lim YS. Characteristics, prevention and management of hepatitis B virus reactivation in HBV-infected patients who require immunosuppressive therapy. *J Infect Dis*. 2017;216(S8):S778–S784.
224. Bird GL, Smith H, Portmann B, et al. Acute liver decompensation on withdrawal of cytotoxic chemotherapy and immuno-suppressive therapy in hepatitis B carriers. *Q J Med*. 1989;73:895–902.
225. Locasciulli A, Alberti A, Bandini G, et al. Allogeneic bone marrow transplantation from HBsAg+ donors: a multicenter study from the Gruppo Italiano Trapianto di Midollo Osseo (GITMO). *Blood*. 1995;86:3236–3240.
226. Seto WK, Chan TS, Hwang YY, et al. Hepatitis B reactivation in occult liver carriers undergoing hematopoietic stem cell transplantation: a prospective study. *Hepatology*. 2017;65:1451–1461.
227. Ramos CA, Saliba RM, de Pádua Silva L, et al. Resolved hepatitis B virus infection is not associated with worse outcome after allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2010;16:686–694.
228. Lau GK, He ML, Fong DY, et al. Preemptive use of lamivudine reduces hepatitis B exacerbation after allogeneic hematopoietic cell transplantation. *Hepatology*. 2002;36:702–709.



229. Huang H, Cai Q, Lin T, et al. Lamivudine for the prevention of hepatitis B virus reactivation after high-dose chemotherapy and autologous hematopoietic stem cell transplantation for patients with advanced or relapsed non-Hodgkin's lymphoma: single institution experience. *Expert Opin Pharmacother*. 2009;10:2399–2406.
230. Ilan Y, Nagler A, Adler R, et al. Ablation of persistent hepatitis B by bone marrow transplantation from a hepatitis B-immune donor. *Gastroenterology*. 1993;104:1818–1821.
231. Lau GK, Suri D, Liang R, et al. Resolution of chronic hepatitis B and anti-HBs seroconversion in humans by adoptive transfer of immunity to hepatitis B core antigen. *Gastroenterology*. 2002;122:614–624.
232. Jamma S, Hussain G, Lau DT. Current concepts of HBV/HCV coinfection: coexistence, but not necessarily in harmony. *Curr Hepat Rep*. 2010;9:260–269.
233. Tyson GL, Kramer JR, Duan Z, et al. Prevalence and predictors of hepatitis B virus coinfection in a United States cohort of hepatitis C virus infected patients. *Hepatology*. 2013;58:538–545.
234. Shih CM, Lo SJ, Miyamura T, et al. Suppression of hepatitis B virus expression and replication by hepatitis C virus core protein in HuH-7 cells. *J Virol*. 1993;67:5823–5832.
235. Sagnelli E, Coppola N, Scolastico C, et al. Virologic and clinical expressions of reciprocal inhibitory effect of hepatitis B, C, and delta viruses in patients with chronic hepatitis. *Hepatology*. 2000;32:1106–1110.
236. Liaw YF, Tsai SL, Chang JJ, et al. Displacement of hepatitis B virus by hepatitis C virus as the cause of continuing chronic hepatitis. *Gastroenterology*. 1994;106:1048–1053.
237. Sagnelli E, Coppola N, Messina V, et al. HBV superinfection in hepatitis C virus chronic carriers, viral interaction, and clinical course. *Hepatology*. 2002;36:1285–1291.
238. French AL, Hotton A, Young M, et al. Isolated hepatitis B core antibody status is not associated with accelerated liver disease progression in HIV/hepatitis C coinfection. *J Acquir Immune Defic Syndr*. 2016;72:274–280.
239. Weltman MD, Brododihardjo A, Crewe EB, et al. Coinfection with hepatitis B and C or B, C and delta viruses results in severe chronic liver disease and responds poorly to interferon-alpha treatment. *J Viral Hepat*. 1995;2:39–45.
240. Zarski JP, Bohn B, Bastie A, et al. Characteristics of patients with dual infection by hepatitis B and C viruses. *J Hepatol*. 1998;28:27–33.
241. Sagnelli E, Coppola N, Pisaturo M, et al. HBV superinfection in HCV chronic carriers: a disease that is frequently severe but associated with the eradication of HCV. *Hepatology*. 2009;49:1090–1097.
242. Sun CA, Farzadegan H, You SL, et al. Mutual confounding and interactive effects between hepatitis C and hepatitis B viral infections in hepatocellular carcinogenesis: a population-based case-control study in Taiwan. *Cancer Epidemiol Biomarkers Prev*. 1996;5:173–178.
243. Hung CH, Lu SN, Wang JH, et al. Sustained HCV clearance by interferon-based therapy reduces hepatocellular carcinoma in hepatitis B and C dually-infected patients. *Antivir Ther*. 2011;16:959–968.
244. Liu CJ, Chuang WL, Lee CM, et al. An open label, comparative, multicenter study of peginterferon alfa-2a plus ribavirin in the treatment of patients with chronic hepatitis C/hepatitis B co-infection versus those with chronic hepatitis C monoinfection. *Gastroenterology*. 2009;136:496–504.
245. Yu ML, Lee CM, Chen CL, et al. Sustained HCV clearance and increased HBsAg seroclearance in patients with dual chronic hepatitis C and B during posttreatment follow-up. *Hepatology*. 2013;57:2135–2142.
246. Aggeletopoulou I, Konstantakis C, Manolakopoulos S, et al. Risk of hepatitis B reactivation in patients treated with direct-acting antivirals for hepatitis C. *World J Gastroenterol*. 2017;23:4317–4323.
247. Calvaruso V, Ferraro D, Licata A, et al. HBV reactivation in patients with HCV/HBV cirrhosis on treatment with direct-acting antivirals. *J Viral Hepat*. 2018;25:72–79.
248. Mücke VT, Mücke MM, Peiffer KH, et al. No evidence of hepatitis B virus reactivation in patients with resolved infection treated with direct-acting antivirals for hepatitis C in a large real-world cohort. *Aliment Pharmacol Ther*. 2017;46:432–439.
249. Wang C, Ji D, Chen J, et al. Hepatitis due to reactivation of hepatitis B virus in endemic areas among patients with hepatitis C treated with direct-acting antiviral agents. *Clin Gastroenterol Hepatol*. 2016;5:30370–30376.
250. Terrault NA, Bzowej NH, Chang KM, et al. AASLD guidelines for the treatment of chronic hepatitis B. *Hepatology*. 2016;61:261–283.
251. Kumar M, Chauhan R, Gupta N, et al. Spontaneous increases in alanine aminotransferase levels in asymptomatic chronic hepatitis B virus-infected patients. *Gastroenterology*. 2009;136:1272–1280.
252. Bonino F, Rosina F, Rizzetto M, et al. Chronic hepatitis in HBsAg carriers with serum HBV-DNA and anti-HBe. *Gastroenterology*. 1986;90:1268–1273.
253. Foy MC, Thio CL, Hwang HS, et al. False-negative hepatitis B virus (HBV) surface antigen in a vaccinated dialysis patient with a high level of HBV DNA in the United States. *Clin Vaccine Immunol*. 2012;19:820–822.
254. Pawlotsky JM. Molecular diagnosis of viral hepatitis. *Gastroenterology*. 2002;122:1554–1568.
255. Lok AS, Heathcote EJ, Hoofnagle JH. Management of hepatitis B: 2000—summary of a workshop. *Gastroenterology*. 2001;120:1828–1853.
256. Maasoumy B, Wiegand SB, Jaroszewicz J, et al. Hepatitis B core-related antigen (HBcrAg) levels in the natural history of hepatitis B virus infection in a large European cohort predominantly infected with genotypes A and D. *Clin Microbiol Infect*. 2015;21:606.e1–606.e10.
257. Van Bommel F, Van Bommel A, Krauel A, et al. Serum HBV RNA is an early predictor of HBeAg seroconversion in patients with chronic Hepatitis B (CHB) treated with pegylated interferon alfa-2a. *Hepatology*. 2015;62:336A.
258. Wang J, Shen T, Huang X, et al. Serum hepatitis B virus RNA is encapsidated pregenome RNA that may be associated with persistence of viral infection and rebound. *J Hepatol*. 2016;65:700–710.
259. Hadler SC, Murphy BL, Schable CA, et al. Epidemiological analysis of the significance of low-positive test results for antibody to hepatitis B surface and core antigens. *J Clin Microbiol*. 1984;19:521–525.
260. Sharifi-Mood B, Sanei-Moghaddam E, Khosravi S. Occult hepatitis B Virus infection among anti-HBc only positive individuals in the southeast of Iran in high prevalence of HBV infection region. *IRCMJ*. 2009;11:90–92.
261. Lok AS, Lai CL, Wu PC. Prevalence of isolated antibody to hepatitis B core antigen in an area endemic for hepatitis B virus infection: implications in hepatitis B vaccination programs. *Hepatology*. 1988;8:766–770.
262. Hoofnagle JH, Seef LB, Bales ZB, et al. Type B hepatitis after transfusion with blood containing antibody to hepatitis B core antigen. *N Engl J Med*. 1978;298:1379–1383.
263. Dickson RC, Everhart JE, Lake JR, et al. Transmission of hepatitis B by transplantation of livers from donors positive for antibody to hepatitis B core antigen. The National Institute of Diabetes and Digestive and Kidney Diseases Liver Transplantation Database. *Gastroenterology*. 1997;113:1668–1674.
264. Marcellin P, Heathcote EJ, Buti M, et al. Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med*. 2008;359:2442–2455.
265. Marcellin P, Lau GK, Bonino F, et al. Peginterferon alfa-2a alone, lamivudine alone, and the two in combination in patients with HBeAg-negative chronic hepatitis B. *N Engl J Med*. 2004;351:1206–1217.
266. Kim WR, Loomba R, Berg T, et al. Impact of long-term tenofovir disoproxil fumarate on incidence of hepatocellular carcinoma in patients with chronic hepatitis B. *Cancer*. 2015;121:3631–3638.
267. Dienstag JL, Goldin RD, Heathcote EJ, et al. Histological outcome during long-term lamivudine therapy. *Gastroenterology*. 2003;124:105–117.
268. Malekzadeh R, Mohamadnejad M, Rakhshani N, et al. Reversibility of cirrhosis in chronic hepatitis B. *Clin Gastroenterol Hepatol*. 2004;2:344–347.
269. van Zonneveld M, Honkoop P, Hansen BE, et al. Long-term follow-up of alpha-interferon treatment of patients with chronic hepatitis B. *Hepatology*. 2004;39:804–810.
270. Papatheodoridis GV, Dimou E, Dimakopoulos K, et al. Outcome of hepatitis B e antigen-negative chronic hepatitis B on long-term nucleos(t)ide analog therapy starting with lamivudine. *Hepatology*. 2005;42:121–129.
271. Papatheodoridis GV, Manesis E, Hadziyannis SJ. The long-term outcome of interferon-alpha treated and untreated patients with HBeAg-negative chronic hepatitis B. *J Hepatol*. 2001;34:306–313.
272. Niederau C, Heintges T, Lange S, et al. Long-term follow-up of HBeAg-positive patients treated with interferon alfa for chronic hepatitis B. *N Engl J Med*. 1996;334:1422–1427.
273. Liaw YF, Sung JJ, Chow WC, et al. Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med*. 2004;351:1521–1531.
274. Kumar M, Satapathy S, Monga R, et al. A randomized controlled trial of lamivudine to treat acute hepatitis B. *Hepatology*. 2007;45:97–101.
275. Jochum C, Maischack F, Anastasiou OE, et al. Treatment of fulminant acute hepatitis B with nucleos(t)ide analogues is safe and does not lead to secondary chonification of hepatitis B. *Z Gastroenterol*. 2016;54:1306–1311.
276. Tillmann HL, Hadem J, Leifeld L, et al. Safety and efficacy of lamivudine in patients with severe acute or fulminant hepatitis B, a multicenter experience. *J Viral Hepat*. 2006;13:256–263.
277. Belloni L, Allweiss L, Guerrieri F, et al. IFN-alpha inhibits HBV transcription and replication in cell culture and in humanized mice by targeting the epigenetic regulation of the nuclear cccDNA minichromosome. *J Clin Invest*. 2012;122:529–537.
278. Janssen HL, van Zonneveld M, Senturk H, et al. Pegylated interferon alfa-2b alone or in combination with lamivudine for HBeAg-positive chronic hepatitis B: a randomised trial. *Lancet*. 2005;365:123–129.
279. Sonneveld MJ, Wong VW, Woltman AM, et al. Polymorphisms near IL28B and serologic response to peginterferon in HBeAg-positive patients with chronic hepatitis B. *Gastroenterology*. 2012;142:513–520.
280. Torre D, Tambini R. Interferon-alpha therapy for chronic hepatitis B in children: a meta-analysis. *Clin Infect Dis*. 1996;23:131–137.
281. Lampertico P, Del Ninno E, Viganò M, et al. Long-term suppression of hepatitis B e antigen-negative chronic hepatitis B by 24-month interferon therapy. *Hepatology*. 2003;37:756–763.
282. Sonneveld MJ, Hansen BE, Piratvisuth T, et al. Response-guided peginterferon therapy in hepatitis B e antigen-positive chronic hepatitis B using serum hepatitis B surface antigen levels. *Hepatology*. 2013;58:872–880.
283. Buster EH, Flink HJ, Kakaloglu Y, et al. Sustained HBeAg and HBsAg loss after long-term follow-up of HBeAg-positive patients treated with peginterferon alfa-2b. *Gastroenterology*. 2008;135:459–467.
284. Deutsch M, Dourakis S, Manesis EK, et al. Thyroid abnormalities in chronic viral hepatitis and their relationship to interferon alfa therapy. *Hepatology*. 1997;26:206–210.
285. Hoofnagle JH, Di Bisceglie AM, Waggoner JG, et al. Interferon alfa for patients with clinically apparent cirrhosis due to chronic hepatitis B. *Gastroenterology*. 1993;104:1116–1121.
286. Villeneuve JP, Condreay LD, Willems B, et al. Lamivudine treatment for decompensated cirrhosis resulting from chronic hepatitis B. *Hepatology*. 2000;31:207–210.
287. Fontana RJ, Keeffe EB, Carey W, et al. Effect of lamivudine treatment on survival of 309 North American patients awaiting liver transplantation for chronic hepatitis B. *Liver Transpl*. 2002;8:433–439.
288. Jang JW, Choi JY, Kim YS, et al. Long-term effect of antiviral therapy on disease course after decompensation in patients with hepatitis B virus-related cirrhosis. *Hepatology*. 2015;61:1809–1820.
289. Dienstag JL, Schiff ER, Wright TL, et al. Lamivudine as initial treatment for chronic hepatitis B in the United States. *N Engl J Med*. 1999;341:1256–1263.
290. Lai C-L, Chien R-N, Leung NWY, et al. A one-year trial of lamivudine for chronic hepatitis B. *N Engl J Med*. 1998;339:61–68.
291. Liaw YF, Leung NW, Chang TT, et al. Effects of extended lamivudine therapy in Asian patients with chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *Gastroenterology*. 2000;119:172–180.
292. Leung NW, Lai CL, Chang TT, et al. Extended lamivudine treatment in patients with chronic hepatitis B enhances hepatitis B e antigen seroconversion rates: results after 3 years of therapy. *Hepatology*. 2001;33:1527–1532.
293. Chien RN, Liaw YF, Atkins M. Pretherapy alanine transaminase level as a determinant for hepatitis B e antigen seroconversion during lamivudine therapy in patients with chronic hepatitis B. Asian Hepatitis Lamivudine Trial Group. *Hepatology*. 1999;30:770–774.
294. Hadziyannis SJ, Papatheodoridis GV, Dimou E, et al. Efficacy of long-term lamivudine monotherapy in patients with hepatitis B e antigen-negative chronic hepatitis B. *Hepatology*. 2000;32:847–851.
295. Lau DT, Khokhar MF, Doo E, et al. Long-term therapy of chronic hepatitis B with lamivudine. *Hepatology*. 2000;32:828–834.
296. Tassopoulos NC, Volpes R, Pastore G, et al. Efficacy of lamivudine in patients with hepatitis B e antigen-negative/hepatitis B virus DNA-positive (precore mutant) chronic hepatitis B. Lamivudine Precore Mutant Study Group. *Hepatology*. 1999;29:889–896.
297. Schiff ER. Lamivudine for hepatitis B in clinical practice. *J Med Virol*. 2000;61:386–391.
298. Perrillo R, Schiff E, Yoshida E, et al. Adefovir dipivoxil for the treatment of lamivudine-resistant hepatitis B mutants. *Hepatology*. 2000;32:129–134.

299. Marcellin P, Chang TT, Lim SG, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *N Engl J Med*. 2003;348:808–816.
300. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med*. 2003;348:800–807.
301. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B for up to 5 years. *Gastroenterology*. 2006;131:1743–1751.
302. Schiff E, Lai CL, Hadziyannis S, et al. Adefovir dipivoxil for wait-listed and post-liver transplantation patients with lamivudine-resistant hepatitis B: final long-term results. *Liver Transpl*. 2007;13:349–360.
303. Chang TT, Gish RG, de Man R, et al. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2006;354:1001–1010.
304. Gish RG, Lok AS, Chang TT, et al. Entecavir therapy for up to 96 weeks in patients with HBeAg-positive chronic hepatitis B. *Gastroenterology*. 2007;133:1437–1444.
305. Lam YF, Seto WK, Wong D, et al. Seven-year treatment outcome of entecavir in a real-world cohort: effects on clinical parameters, HBsAg and HBeAg levels. *Clin Transl Gastroenterol*. 2017;8:e125.
306. McMahon MA, Jilek BL, Brennan TP, et al. The HBV drug entecavir—effects on HIV-1 replication and resistance. *N Engl J Med*. 2007;356:2614–2621.
307. Lai CL, Gane E, Liaw YF, Globe Study Group, et al. Telbivudine versus lamivudine in patients with chronic hepatitis B. *N Engl J Med*. 2007;357:2576–2588.
308. Buti M, Tsai N, Petersen J, et al. Seven-year efficacy and safety of treatment with tenofovir disoproxil fumarate for chronic hepatitis B virus infection. *Dig Dis Sci*. 2015;60:1457–1464.
309. Gallant J, Brunetta J, Crofoot G, et al. Efficacy and safety of switching to simpler single tablet regimen of elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (E/C/F/TAF) in HIV-1/hepatitis B co-infected adults. *J Acquir Immune Defic Syndr*. 2016;73:294–298.
310. Buti M, Gane E, Seto WK, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of patients with HBeAg-negative chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol*. 2016;1:196–206.
311. Chan HL, Fund S, Seto WK, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of HBeAg-positive chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol*. 2016;1:185–195.
312. Brouwer WP, Xie Q, Sonneveld MJ, et al. Adding pegylated interferon to entecavir for hepatitis B e antigen-positive chronic hepatitis B: a multicenter randomized trial (ARES study). *Hepatology*. 2015;61:1512–1522.
313. Jun DW, Sohn JH, Kim SG, et al. A randomized, multicenter, phase IIIb open-label study to evaluate the efficacy of sequential therapy of peginterferon alfa-2a following entecavir compared with peginterferon alfa-2a monotherapy in patients with HBeAg positive chronic hepatitis B: interim analysis of POTENT study. Abstract 377. Presented at the 63rd Annual Meeting of the American Association for the Study of Liver Diseases. Boston, 2012.
314. Xie Q, Zhou H, Bai X, et al. A randomized, open-label clinical study of combined pegylated interferon Alfa-2a (40 KD) and entecavir treatment for hepatitis B “e” antigen-positive chronic hepatitis B. *Clin Infect Dis*. 2014;59:1714–1723.
315. Marcellin P, Ahn SH, Ma X, et al. Combination of tenofovir disoproxil fumarate and peginterferon  $\alpha$ -2a increases loss of hepatitis b surface antigen in patients with chronic hepatitis B. *Gastroenterology*. 2016;150:134–144.e10.
316. Chi H, Hanson BE, Guo S, et al. Pegylated interferon alfa-2b Add-on treatment in hepatitis B virus envelope antigen-positive chronic hepatitis B patients treated with nucleos(t)ide analogue: a randomized, controlled trial (PEGON). *J Infect Dis*. 2017;215:1085–1093.
317. Lok AS, Trinh H, Carosi G, et al. Efficacy of entecavir with or without tenofovir disoproxil fumarate for nucleos(t)ide-naïve patients with chronic hepatitis B. *Gastroenterology*. 2012;143:619–628.
318. Das K, Xiong X, Yang H, et al. Molecular modeling and biochemical characterization reveal the mechanism of hepatitis B virus polymerase resistance to lamivudine (3TC) and emtricitabine (FTC). *J Virol*. 2001;75:4771–4779.
319. Bozdayi AM, Uzunalioglu O, Turkyilmaz AR, et al. YSDD: a novel mutation in HBV DNA polymerase confers clinical resistance to lamivudine. *J Viral Hepat*. 2003;10:256–265.
320. Ono SK, Kato N, Shiratori Y, et al. The polymerase L528M mutation cooperates with nucleotide binding-site mutations, increasing hepatitis B virus replication and drug resistance. *J Clin Invest*. 2001;107:449–455.
321. Lok AS, Hussain M, Cursano C, et al. Evolution of hepatitis B virus polymerase gene mutations in hepatitis B e antigen-negative patients receiving lamivudine therapy. *Hepatology*. 2000;32:1145–1153.
322. Liaw YF, Gane E, Leung N, et al. 2-Year GLOBE trial results: telbivudine is superior to lamivudine in patients with chronic hepatitis B. *Gastroenterology*. 2009;136:486–495.
323. Seifer M, Patty A, Serra I, et al. Telbivudine, a nucleoside analog inhibitor of HBV polymerase, has a different in vitro cross-resistance profile than the nucleotide analog inhibitors adefovir and tenofovir. *Antiviral Res*. 2009;81:147–155.
324. Tenney DJ, Rose RE, Baldick CJ, et al. Long-term monitoring shows hepatitis B virus resistance to entecavir in nucleoside-naïve patients is rare through 5 years of therapy. *Hepatology*. 2009;49:1503–1514.
325. Chen CH, Wang JH, Lu SN, et al. Characteristics of adefovir resistance in patients with or without lamivudine-resistant hepatitis B virus treated with adefovir: a 4-year experience. *Liver Int*. 2011;31:206–214.
326. Buti M, Fung S, Gane E, et al. Long-term clinical outcomes in cirrhotic chronic hepatitis B patients treated with tenofovir disoproxil fumarate for up to 5 years. *Hepatol Int*. 2015;9:243.
327. Villet S, Pichoud C, Billioud G, et al. Impact of hepatitis B virus rtA181V/T mutants on hepatitis B treatment failure. *J Hepatol*. 2008;48:747–755.
328. Liaw YF. Acute exacerbation and superinfection in patients with chronic viral hepatitis. *J Formos Med Assoc*. 1995;94:521–528.
329. Park JY, Kim CW, Bae SH, et al. Entecavir plus tenofovir combination therapy in patients with multidrug-resistant chronic hepatitis B: results of a multicentre, prospective study. *Liver Int*. 2016;36:1108.
330. Lim YS, Yoo BC, Byun KS, et al. Tenofovir monotherapy versus tenofovir and entecavir combination therapy in adefovir-resistant chronic hepatitis B patients with multiple drug failure: results of a randomised trial. *Gut*. 2016;65:1042–1051.
331. Torresi J, Earnest Silveira L, Deliyannis G, et al. Reduced anti-genicity of the hepatitis B virus HBsAg protein arising as a consequence of sequence changes in the overlapping polymerase gene that are selected by lamivudine therapy. *Virology*. 2005;293:305–313.
332. Lee KM, Cho SW, Kim SW, et al. Effect of virological response on post-treatment durability of lamivudine-induced HBeAg seroconversion. *J Viral Hepat*. 2002;9:208–212.
333. Chaung KT, Ha NB, Trinh HN, et al. High frequency of recurrent viremia after hepatitis B e antigen seroconversion and consolidation therapy. *J Clin Gastroenterol*. 2012;46:865–870.
334. Lee HA, Seo YS, Park SW, et al. Hepatitis B surface antigen titer is a good indicator of durable viral response after entecavir off-treatment for chronic hepatitis B. *Clin Mol Hepatol*. 2016;22:382–389.
335. Berg T, Simon KG, Mauss S, et al. Long-term response after stopping tenofovir disoproxil fumarate in non-cirrhotic HBeAg-negative patients—FINITE study. *J Hepatol*. 2017;67:918.
336. Lok AS. Prevention of recurrent hepatitis B post-liver transplantation. *Liver Transpl*. 2002;8(10 suppl 1):S67–S73.
337. Loomba R, Rowley AK, Wesley R, et al. Hepatitis B immuno-globulin and lamivudine improve hepatitis B-related outcomes after liver transplantation: meta-analysis. *Clin Gastroenterol Hepatol*. 2008;6:696–700.
338. Bock CT, Tillmann HL, Torresi J, et al. Selection of hepatitis B virus polymerase mutants with enhanced replication by lamivudine treatment after liver transplantation. *Gastroenterology*. 2002;122:264–273.
339. Mutimer D, Ferasz-Neto BH, Harrison R, et al. Acute liver graft failure due to emergence of lamivudine resistant hepatitis B virus: rapid resolution during treatment with adefovir. *Dig*. 2001;49:860–863.
340. Fung J, Wong T, Chok K, et al. Long-term outcomes of entecavir therapy for chronic Hepatitis B after liver transplantation: results up to 8 years. *Hepatology*. 2017;66:1036–1044.
341. Teperman LW, Poordad F, Bzowej N, et al. A randomized trial of emtricitabine/tenofovir disoproxil fumarate after hepatitis B immunoglobulin withdrawal post-liver transplant. *Liver Transpl*. 2013;19:594–601.
342. Fox AN, Terrault NA. The option of HBIG-free prophylaxis against recurrent HBV. *J Hepatol*. 2012;56:1189–1197.
343. Sanchez-Fueyo A, Rimola A, Grande L, et al. Hepatitis B immunoglobulin discontinuation followed by hepatitis B virus vaccination: a new strategy in the prophylaxis of hepatitis B virus recurrence after liver transplantation. *Hepatology*. 2000;31:496–501.
344. Angelico M, Di Paolo D, Trinito MO, et al. Failure of a reinforced triple course of hepatitis B vaccination in patients transplanted for HBV-related cirrhosis. *Hepatology*. 2002;35:176–181.
345. Rosenau J, Hooman N, Hadem J, et al. Failure of hepatitis B vaccination with conventional HBsAg vaccine in patients with continuous HBIG prophylaxis after liver transplantation. *Liver Transpl*. 2007;13:367–373.
346. Bienze U, Gunther M, Neuhaus R, et al. Immunization with an adjuvant hepatitis B vaccine after liver transplantation for hepatitis B-related disease. *Hepatology*. 2003;38:811–819.
347. Rosenau J, Hooman N, Rifai K, et al. Hepatitis B virus immunization with an adjuvant containing vaccine after liver transplantation for hepatitis B-related disease: failure of humoral and cellular immune response. *Transpl Int*. 2006;19:828–833.
348. Peters MG, Andersen J, Lynch P, et al. Randomized controlled study of tenofovir and adefovir in chronic hepatitis B virus and HIV infection: ACTG A5127. *Hepatology*. 2006;44:1110–1116.
349. Matthews GV, Bartholomeusz A, Locarnini S, et al. Characteristics of drug resistant HBV in an international collaborative study of HIV-HBV-infected individuals on extended lamivudine therapy. *AIDS*. 2006;20:863–870.
350. Matthews GV, Avihingsanon A, Lewin SR, et al. A randomized trial of combination hepatitis B therapy in HIV/HBV coinfected antiretroviral naive individuals in Thailand. *Hepatology*. 2008;48:1062–1069.
351. Bani-Sadr F, Palmer P, Scieux C, et al. Ninety-six-week efficacy of combination therapy with lamivudine and tenofovir in patients coinfected with HIV-1 and wild-type hepatitis B virus. *Clin Infect Dis*. 2004;39:1062–1064.
352. de Vries-Sluijs TE, Reijnders JG, Hansen BE, et al. Long-term therapy with tenofovir is effective for patients co-infected with human immunodeficiency virus and hepatitis B virus. *Gastroenterology*. 2010;139:1934–1941.
353. Jain MK, Comanor L, White C, et al. Treatment of hepatitis B with lamivudine and tenofovir in HIV/HBV-coinfected patients: factors associated with response. *J Viral Hepat*. 2007;14:176–182.
354. Department of Health and Human Services, Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents. <http://aidsinfo.nih.gov/contentfiles/vguidelines/adultandadolescentgl.pdf>. Accessed December 12, 2017.
355. Perrillo RP, Gish R, Falck-Ytter YT. American Gastroenterological Association. Institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology*. 2015;148:221–244.e3.
356. Evans AM, Jovanovic BD, Su YC, et al. Rituximab-associated hepatitis B virus (HBV) reactivation in lymphoproliferative diseases: meta-analysis and examination of FDA safety reports. *Ann Oncol*. 2011;22:1170–1180.
357. Yeo W, Chan TC, Leung NW, et al. Hepatitis B virus reactivation in lymphoma patients with prior resolved hepatitis B undergoing anticancer therapy with or without rituximab. *J Clin Oncol*. 2009;27:605–611.
- 357a. Mitka M. FDA: increased HBV reactivation risk with ofatumumab or rituximab. *JAMA*. 2013;310:1664.
358. Lau GK, Liang R, Chiu EK, et al. Hepatic events after bone marrow transplantation in patients with hepatitis B infection: a case controlled study. *Bone Marrow Transplant*. 1997;19:795–799.
359. Buti M, Manzano ML, Morillas RM, et al. Randomized prospective study evaluating tenofovir disoproxil fumarate prophylaxis against hepatitis B virus reactivation in anti-HBe-positive patients with rituximab-based regimens to treat hematologic malignancies: The Preblin study. *PLoS ONE*. 2017;12:e0184550.
360. Zhou K, Terrault N. Management of hepatitis B in special populations. *Best Pract Res Clin Gastroenterol*. 2017;31:311–320.
361. Huang H, Li X, Zhu J, et al. Entecavir vs lamivudine for prevention of hepatitis B virus reactivation among patients with untreated diffuse large B-cell lymphoma receiving R-CHOP chemotherapy: a randomized clinical trial. *JAMA*. 2014;312:2521–2530.
362. Lau GK, Liang R, Lee CK, et al. Clearance of persistent hepatitis B virus infection in Chinese bone marrow transplant recipients whose donors were anti-hepatitis B core- and anti-hepatitis B surface antibody-positive. *J Infect Dis*. 1998;178:1585–1591.



363. Brown RS Jr, Verna EC, Pereira MR, et al. Hepatitis B virus and human immunodeficiency virus drugs in pregnancy: findings from the Antiretroviral Pregnancy Registry. *J Hepatol*. 2012;57:953–959.
364. Brown RS Jr, McMahon BJ, Lok AS, et al. Antiviral therapy in chronic hepatitis B viral infection during pregnancy: a systematic review and meta-analysis. *Hepatology*. 2016;63:319–333.
365. Kim DK, Riley LE, Harriman KH, et al. Advisory Committee on Immunization Practices Recommended Immunization Schedule for Adults Aged 19 Years or Older—United States, 2017. *MMWR Morb Mortal Wkly Rep*. 2017;66:136–138.
366. Keefe EB. Is hepatitis A more severe in patients with chronic hepatitis B and other chronic liver diseases? *Am J Gastroenterol*. 1995;90:201–205.
367. Vento S, Garofano T, Renzini C, et al. Fulminant hepatitis associated with hepatitis B virus superinfection in patients with chronic hepatitis C. *N Engl J Med*. 1998;338:286–290.
368. Corden S, Ballard AL, Ijaz S, et al. HBV DNA levels and transmission of hepatitis B by health care workers. *J Clin Virol*. 2003;27:52–58.
369. Incident Investigation Teams and Others. Transmission of hepatitis B to patients from four infected surgeons without hepatitis B e antigen. *N Engl J Med*. 1997;336:178–184.
370. Spijkerman IJ, van Doorn LJ, Janssen MH, et al. Transmission of hepatitis B virus from a surgeon to his patients during high-risk and low-risk surgical procedures during 4 years. *Infect Control Hosp Epidemiol*. 2002;23:306–312.
371. Centers for Disease Control and Prevention. Updated CDC recommendations for the management of hepatitis B virus-infected health-care providers and students. *MMWR Recomm Rep*. 2012;61(RR-3):1–12.
372. Yang HJ, Lu SN, Liaw YF, et al. Hepatitis B e antigen and the risk of hepatocellular carcinoma. *N Engl J Med*. 2002;347:168–174.
373. Yang HJ, Sherman M, Su J, et al. Nomograms for risk of hepatocellular carcinoma in patients with chronic hepatitis B virus infection. *J Clin Oncol*. 2010;28:2437–2444.
374. Fattovich G, Stroffolini T, Zagni I, et al. Hepatocellular carcinoma in cirrhosis: incidence and risk factors. *Gastroenterology*. 2004;127(5 suppl 1):S35–S50.
375. Zhang BH, Zhang BH, Tang ZY. Randomized controlled trial of screening for hepatocellular carcinoma. *J Cancer Res Clin Oncol*. 2004;130:417–422.
376. Zhang W, Wang X, Wang Y, et al. Effective viral suppression is necessary to reduce hepatocellular carcinoma development in cirrhotic patients with chronic hepatitis B: results of a 10-year follow up. *Medicine (Baltimore)*. 2017;96:e8454.
377. Huang LM, Lu CY, Chen DS, et al. Hepatitis B virus infection, its sequelae, and prevention by vaccination. *Curr Opin Immunol*. 2011;23:237–243.
378. Chen DS. Hepatitis B vaccination: the key towards elimination and eradication of hepatitis B. *J Hepatol*. 2009;50:805–816.
379. Koff RS. Immunogenicity of hepatitis B vaccines: implications of immune memory. *Vaccine*. 2002;20:3695–3701.
380. Gwack J, Park SK, Lee EH, et al. Hepatitis B vaccination and liver cancer mortality reduction in Korean children and adolescents. *Asian Pac J Cancer Prev*. 2011;12:2205–2208.
381. Chang MH, Chen CJ, Lai MS, et al. Universal hepatitis B vaccination in Taiwan and the incidence of hepatocellular carcinoma in children. Taiwan Childhood Hepatoma Study Group. *N Engl J Med*. 1997;336:1855–1859.
382. Halperin SA, Ward B, Cooper C, et al. Comparison of safety and immunogenicity of two doses of investigational hepatitis B virus surface antigen co-administered with an immunostimulatory phosphorothioate oligodeoxynucleotide and three doses of a licensed hepatitis B vaccine in healthy adults 18–55 years of age. *Vaccine*. 2012;30:2556–2563.
383. Heyward WL, Kyle M, Blumenau J, et al. Immunogenicity and safety of an investigational hepatitis B vaccine with a Toll-like receptor 9 agonist adjuvant (HBsAg-1018) compared to a licensed hepatitis B vaccine in healthy adults 40–70 years of age. *Vaccine*. 2013;31:5300–5305.
384. Jackson S, Lentino J, Kopp J, et al. Immunogenicity of a two-dose investigational hepatitis B vaccine, HBsAg-1018, using a toll-like receptor 9 agonist adjuvant compared with a licensed hepatitis B vaccine in adults. *Vaccine*. 2018;36:668–674. <https://doi.org/10.1016/j.vaccine.2017.12.038>.
385. Janssen J, Jackson S, Heyward WL, et al. Immunogenicity of an investigational hepatitis B vaccine with a toll-like receptor 9 agonist adjuvant HBsAg-1018 compared with a licensed hepatitis B vaccine in subpopulations of healthy adults 18–70 years of age. *Vaccine*. 2015;33:833–837.
386. Abara WE, Qaseem A, Schillie S, et al. Hepatitis B vaccination, screening and linkage to care: best practice advice from the American College of Physicians and Centers for Disease Control and Prevention. *Ann Intern Med*. 2017;167:794–804.
387. Schillie S, Vellozzi C, Reingold A, et al. Prevention of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices. *MMWR Recomm Rep*. 2018;67:1–31.
388. World Health Organization. Global Hepatitis Report, Geneva, 2017. <http://apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf?ua=1>. Accessed May 2017.
389. Hill HA, Elam-Evans LD, Yankey D, Centers for Disease Control and Prevention, et al. National, state, and urban area vaccination coverage among children aged 19–35 months—United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2015;64:889–896.
390. American Academy of Pediatrics Policy Statement. Elimination of Perinatal Hepatitis B: Providing the First Vaccine Dose Within 24 Hours of Birth. *Pediatrics*. 2017;140:<http://pediatrics.aappublications.org/content/140/3/e20171870>.
391. Centers for Disease Control and Prevention. Hepatitis B FAQs for Health Professionals. <https://www.cdc.gov/hepatitis/hbv/hbvfaq.htm#vaccFAQ>.
392. Centers for Disease Control and Prevention. Alternate two dose hepatitis B vaccination schedule for adolescents aged 11 to 15 years. *JAMA*. 2000;283:2100.
393. Lindsay KL, Herbert DA, Gitnick GL. Hepatitis B vaccine: low postvaccination immunity in hospital personnel given gluteal injections. *Hepatology*. 1985;5:1088–1090.
394. Kao JH, Chen DS. Hepatitis B vaccination: to boost or not to boost? *Lancet*. 2005;366:1337–1338.
395. Schillie S, Murphy TV, Fenlon N, et al. Update: shortened interval for postvaccination serologic testing of infants born to hepatitis B-infected mothers. *MMWR Morb Mortal Wkly Rep*. 2015;64:1118–1120.
396. Prather AA, Hall M, Fury JM, et al. Sleep and antibody response to hepatitis B vaccination. *Sleep*. 2012;35:1063–1069.
397. Yang S, Tian G, Cui Y, et al. Factors influencing immunologic response to hepatitis B vaccine in adults. *Sci Rep*. 2016;6:27251.
398. Tedaldi EM, Baker RK, Moorman AC, et al. Hepatitis A and B vaccination practices for ambulatory patients infected with HIV. *Clin Infect Dis*. 2004;38:1478–1484.
399. Pettit NN, DePestel DD, Malani PN, et al. Factors associated with seroconversion after standard dose hepatitis B vaccination and high-dose revaccination among HIV-infected patients. *HIV Clin Trials*. 2010;11:332–339.
400. de Vries-Sluijs TE, Hansen BE, van Doornum GJ, et al. A randomized controlled study of accelerated versus standard hepatitis B vaccination in HIV-positive patients. *J Infect Dis*. 2011;203:984–991.
401. Wong EK, Bodsworth NJ, Slade MA, et al. Response to hepatitis B vaccination in a primary care setting: influence of HIV infection, CD4+ lymphocyte count and vaccination schedule. *Int J STD AIDS*. 1996;7:490–494.
402. Collier AC, Corey L, Murphy VL, et al. Antibody to human immunodeficiency virus (HIV) and suboptimal response to hepatitis B vaccination. *Ann Intern Med*. 1988;109:101–105.
403. Hadler SC, Judson FN, O'Malley PM, et al. Outcome of hepatitis B virus infection in homosexual men and its relation to prior human immunodeficiency virus infection. *J Infect Dis*. 1991;163:454–459.
404. Launay O, van der Vliet D, Rosenberg AR, et al. Safety and immunogenicity of 4 intramuscular double doses and 4 intradermal low doses vs. standard hepatitis B vaccine regimen in adults with HIV-1. *JAMA*. 2011;305:1432–1440.
405. Valdez H, Mitsuyasu R, Landay A, et al. Interleukin-2 increases CD4+ lymphocyte numbers but does not enhance responses to immunization: results of A5046s. *J Infect Dis*. 2003;187:320–325.
406. Sasaki Md, Focaccia R, de Messias-Reason JJ. Efficacy of granulocyte-macrophage colony-stimulating factor (GM-CSF) as a vaccine adjuvant for hepatitis B virus in patients with HIV infection. *Vaccine*. 2003;21:4545–4549.
407. Overton ET, Kang M, Peters MG, et al. Immune response to hepatitis B vaccine in HIV-infected subjects using granulocyte-macrophage colony-stimulating factor (GM-CSF) as a vaccine adjuvant: ACTG study 5220. *Vaccine*. 2010;28:5597–5604.
408. Rey D, Piroth L, Wendling MJ, et al. Safety and immunogenicity of double-dose versus standard-dose hepatitis B revaccination in non-responding adults with HIV-1 (ANRS HB04 B-BOOST): a multicentre, open-label, randomised controlled trial. *Lancet Infect Dis*. 2015;15:1283–1291.
409. Propst T, Propst A, Lhotka K, et al. Reinforced intradermal hepatitis B vaccination in hemodialysis patients is superior in antibody response to intramuscular or subcutaneous vaccination. *Am J Kidney Dis*. 1998;32:1041–1045.
410. Bruguera M, Rodicio JL, Alcazar JM, et al. Effects of different dose levels and vaccination schedules on immune response to a recombinant DNA hepatitis B vaccine in haemodialysis patients. *Vaccine*. 1990;8(suppl):S47–S49.
411. Waite NM, Thomson LG, Goldstein MB. Successful vaccination with intradermal hepatitis B vaccine in hemodialysis patients previously nonresponsive to intramuscular hepatitis B vaccine. *J Am Soc Nephrol*. 1995;5:1930–1934.
412. Yousaf F, Gandham S, Galler M, et al. Systematic review of the efficacy and safety of intradermal versus intramuscular hepatitis B vaccination in end-stage renal disease population unresponsive to primary vaccination series. *Ren Fail*. 2015;37:1080–1088.
413. Centers for Disease Control and Prevention. Guidelines for Vaccinating Kidney Dialysis Patients and Patients with Chronic Kidney Disease. <https://www.cdc.gov/vaccines/pubs/downloads/dialysis-guide-2012.pdf>. Accessed August 2, 2016.
414. McMahon BJ, Dentinger CM, Bruden D, et al. Antibody levels and protection after hepatitis B vaccine: results of a 22-year follow-up study and response to a booster dose. *J Infect Dis*. 2009;200:1390–1396.
415. Bruce MG, Bruden D, Hurlburt D, et al. Antibody levels and protection after hepatitis B vaccine: results of a 30-year follow-up study and response to a booster dose. *J Infect Dis*. 2016;214:16–22.
416. Simons BC, Spradling PR, Bruden DJ, et al. A Longitudinal hepatitis B vaccine cohort demonstrates long-lasting hepatitis B virus (HBV) cellular immunity despite loss of antibody against HBV surface antigen. *J Infect Dis*. 2016;214:273–280.
417. Van Damme P. Long-term protection after hepatitis B vaccine. *J Infect Dis*. 2016;214:1–3.
418. Ni YH, Huang LM, Chang MH, et al. Two decades of universal hepatitis B vaccination in Taiwan: impact and implication for future strategies. *Gastroenterology*. 2007;132:1287–1293.
419. Lopes VB, Hassing RJ, de Vries-Sluijs TE, et al. Long-term response rates of successful hepatitis B vaccination in HIV-infected patients. *Vaccine*. 2012;31:1040–1044.
420. U.S. Public Health Service. Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV, and HIV and Recommendations for Postexposure Prophylaxis. *MMWR Recomm Rep*. 2001;50(RR-11):1–42.
421. Borgia G, Carleo MA, Gaeta GB, et al. Hepatitis B in pregnancy. *World J Gastroenterol*. 2012;18:4677–4683.
422. Stevens CE, Taylor PE, Tong MJ, et al. Yeast-recombinant hepatitis B vaccine: efficacy with hepatitis B immune globulin in prevention of perinatal hepatitis B virus transmission. *JAMA*. 1987;257:2612–2616.
423. Stevens CE, Toy PT, Tong MJ, et al. Perinatal hepatitis B virus transmission in the United States: prevention by passive-active immunization. *JAMA*. 1985;253:1740–1745.
424. Wong VC, Ip HM, Reesink HW, et al. Prevention of the HBsAg carrier state in newborn infants of mothers who are chronic carriers of HBsAg and HBeAg by administration of hepatitis-B vaccine and hepatitis-B immunoglobulin: double-blind randomised placebo-controlled study. *Lancet*. 1984;1:921–926.
425. Lee C, Gong Y, Brok J, et al. Effect of hepatitis B immunization in newborn infants of mothers positive for hepatitis B surface antigen: systematic review and meta-analysis. *BMJ*. 2006;332:328–336.
426. Machaira M, Papaevangelou V, Voulomanou EK, et al. Hepatitis B vaccine alone or with hepatitis B immunoglobulin in neonates of HBsAg+/HBeAg-mothers: a systematic review and meta-analysis. *J Antimicrob Chemother*. 2015;70:396–404.
427. Lu Y, Liang XF, Wang FZ, et al. Hepatitis B vaccine alone may be enough for preventing hepatitis B virus transmission in neonates of HBsAg (+)/HBeAg (-) mothers. *Vaccine*. 2017;35:40–45.
428. World Health Organization, Media Centre. Immunization Coverage. <http://www.who.int/mediacentre/factsheets/fs378/en/>. Accessed December 5, 2017.
429. World Health Organization. Immunization Coverage. [http://www.who.int/immunization/monitoring\\_surveillance/data/en](http://www.who.int/immunization/monitoring_surveillance/data/en). Accessed April 2017. (revised).



430. Lin J, Liao S, Guo GZ, et al. Significance of hepatitis B virus DNA quantity in transmission between mothers and infants. *Chinese J Ped.* 2002;40:84–87.
431. Patton H, Tran TT. Management of hepatitis B during pregnancy. *Nat Rev Gastroenterol Hepatol.* 2014;11:402–409.
432. Zou H, et al. Virologic factors associated with failure to passive-active immunoprophylaxis in infants born to HBsAg-positive mothers. *J Viral Hepat.* 2012;19:e18–e25.
433. Pan CQ, Duan Z, Dai E, et al. China Study Group for the Mother-to-Child Transmission of Hepatitis B. Tenofovir to prevent hepatitis B transmission in mothers with high viral load. *N Engl J Med.* 2016;374:2324–2334.
434. Greenup AJ, Tan PK, Nguyen V, et al. Efficacy and safety of tenofovir disoproxil fumarate in pregnancy to prevent perinatal transmission of hepatitis B virus. *J Hepatol.* 2014;61:502–507.
435. Chen HL, Lee CN, Chang CH, et al. Taiwan Study Group for the Prevention of Mother-to-Infant Transmission of HBV (PreMIT Study); Taiwan Study Group for the Prevention of Mother-to-Infant Transmission of HBV PreMIT Study. Efficacy of maternal tenofovir disoproxil fumarate in interrupting mother-to-infant transmission of hepatitis B virus. *Hepatology.* 2015;62:375–386.
436. Wang L, Kourtis AP, Ellington S, et al. Safety of tenofovir during pregnancy for the mother and fetus: a systematic review. *Clin Infect Dis.* 2013;57:1773–1781.
437. Ayres A, Yuen L, Jackson KM, et al. Short duration of lamivudine for the prevention of hepatitis B virus transmission in pregnancy: lack of potency and selection of resistance mutations. *J Viral Hepat.* 2014;21:809–817.
438. Han GR, Cao MK, Zhao W, et al. A prospective and open-label study for the efficacy and safety of telbivudine in pregnancy for the prevention of perinatal transmission of hepatitis B virus infection. *J Hepatol.* 2011;55:1215–1221.
439. Wang J, Zhu Q, Zhang X. Effect of delivery mode on maternal-infant transmission of hepatitis B virus by immunoprophylaxis. *Chin Med J.* 2002;115:1510–1512.
440. Beasley RP, Stevens CE, Shiao IS, et al. Evidence against breastfeeding as a mechanism for vertical transmission of hepatitis B. *Lancet.* 1975;2:740–741.
441. Carman WF, Zanetti AR, Karayiannis P, et al. Vaccine-induced escape mutant of hepatitis B virus. *Lancet.* 1990;336:325–329.
442. Hsu HY, Chang MH, Liaw SH, et al. Changes of hepatitis B surface antigen variants in carrier children before and after universal vaccination in Taiwan. *Hepatology.* 1999;30:1312–1317.
443. He C, Nomura F, Itoga S, et al. Prevalence of vaccine-induced escape mutants of hepatitis B virus in the adult population in China: a prospective study in 176 restaurant employees. *J Gastroenterol Hepatol.* 2001;16:1373–1377.
444. Carman WF, Trautwein C, van Deursen FJ, et al. Hepatitis B virus envelope variation after transplantation with and without hepatitis B immune globulin prophylaxis. *Hepatology.* 1996;24:489–493.
445. Ando T, Kojima K, Isoda H, et al. Reactivation of resolved infection with the hepatitis B virus immune escape mutant G145R during dasatinib treatment for chronic myeloid leukemia. *Int J Hematol.* 2015;102:379–382.
446. Aghakhani A, Mohraz M, Aghasadeghi MR, et al. Occult hepatitis B virus infection and S gene escape mutants in HIV-infected patients after hepatitis B virus vaccination. *Int J STD AIDS.* 2016;27:967–972.
447. Amponsah-Dacosta E, Lebelo RL, Rakgole JN, et al. Hepatitis B virus infection in post-vaccination South Africa: occult HBV infection and circulating surface gene variants. *J Clin Virol.* 2015;63:12–17.
448. Ogata N, Cote PJ, Zanetti AR, et al. Licensed recombinant hepatitis B vaccines protect chimpanzees against infection with the prototype surface gene mutant of hepatitis B virus. *Hepatology.* 1999;30:779–786.

## SHORT VIEW SUMMARY

**Definition**

- Hepatitis delta virus (HDV) occurs as a coinfection or superinfection with HBV, establishing a chronic infection in hepatocytes.
- About 5% of people with chronic hepatitis B globally have evidence of exposure to HDV.
- HDV prevalence generally mirrors hepatitis B virus (HBV) prevalence, although there are geographic differences.
- HDV is primarily transmitted via the parenteral route, but there is some sexual transmission. Perinatal transmission is uncommon.

**Microbiology**

- HDV is a small, defective RNA virus that relies on host cell machinery for replication.

- HDV requires the envelope of HBV for viral assembly and transmission.

**Clinical Presentation and Diagnosis**

- Clinical presentation of HDV is variable, but both chronic and superinfection can be present as acute severe hepatitis.
- Superinfection of HDV in individuals with chronic hepatitis B results in chronic infection in greater than 90% of cases. The risk of cirrhosis and hepatocellular carcinoma is significantly elevated in these patients.
- Laboratory diagnosis of HDV is with enzyme immunoassay for HDV antibodies and HDV RNA by real-time polymerase chain reaction.

**Therapy**

- HDV is treated with pegylated interferon- $\alpha$  weekly for a minimum of 48 weeks.
- Newer therapies are currently under development.

**Prevention**

- Vaccination against hepatitis B (see Chapter 145) can be preventative because HDV needs HBV to replicate.

**HEPATITIS DELTA VIRUS**

Delta agent was identified by Mario Rizzetto in 1977 as a nuclear antigen distinct from hepatitis B surface antigen (HBsAg), hepatitis B core antigen (HBcAg), and hepatitis B e antigen (HBeAg) in hepatocytes of some HBsAg carriers in Italy. It soon became clear that this passenger virus, termed *hepatitis D virus* (HDV), accompanied hepatitis B virus (HBV) infection and required the HBsAg for transmission. This unique HDV RNA genome is circular and resembles plant pathogens, including viroids and virusoids.<sup>1</sup> HDV is the only member of the genus *Deltavirus*.<sup>2</sup> In nature HDV is only found in patients who are also infected with HBV.

HDV is an enveloped virus of 36 nm that is distinct from 22-nm HBsAg or 42-nm HBV particles. Because HDV derives its viral envelope from HBV, the outer coat consists of the three forms (L, M, and S) of HBsAg and host lipids that surround the nucleocapsid, which includes the HDV genome in a complex with hepatitis delta antigen (HDAg). The genome is a small, single-stranded, circular RNA genome of approximately 1680 nucleotides, which has a high guanine-cytosine content, allowing it to fold into a rodlike structure.<sup>3</sup> Similar to other RNA viruses, there is no proofreading ability of the RNA polymerases, so the mutation rate is high, and the virus exists as a swarm of closely related quasispecies in an individual. The C-terminal region of the large-HDAg (L-HDAg) is the most variable.

**Hepatitis Delta Antigen**

The HDAg is the only known protein encoded by HDV and consists of two isoforms: small S-HDAg (24 kilodalton [kDa]) and large L-HDAg (27 kDa) (Fig. 146.1). Both are initiated from the same adenine-thymine-guanine (ATG) codon on the antigenomic RNA and are identical except that the L-HDAg has an additional 19 amino acids at the end of the C-terminus due to a unique RNA editing event during replication via the cellular enzyme adenosine deaminase acting on RNA 1 (ADAR1).<sup>4-6</sup> Each is translated from a distinct species of RNA. Both L- and S-HDAgs are phosphorylated and contain a nuclear localization signal that imports them into the nucleus.<sup>7,8</sup> L-HDAg acts as a dominant-negative inhibitor of replication and is necessary for binding HBsAg for virion assembly.<sup>9</sup> Binding of the L-HDAg to HBsAg is facilitated by prenylation of the

C-terminal cysteine residue, which is a current target for new therapeutics (see “Treatment”). S-HDAg is involved in the initiation of RNA replication and is believed to have RNA chaperone activity.<sup>10</sup> Posttranslational modification of the S-HDAg affects the synthesis of genomic RNA and is a potential novel therapeutic target. Of note, L-HDAg and S-HDAg downregulate HBV replication through repression of two HBV enhancer regions and by activating the myxovirus resistance A (MxA) gene, which reduces export of HBV messenger (m)RNA from the nucleus.<sup>11</sup>


**Hepatitis D Virus Life Cycle Hepatocyte Entry**

HDV infects only hepatocytes with no evidence of extrahepatic sites of its replication. The mechanism of HDV entry is similar to HBV because it requires L-HBsAg for entry through binding the sodium taurocholate cotransporting polypeptide receptor, followed by internalization through endocytosis (see “Attachment, Entry, and Hepatotropism” in Chapter 145). Once in the hepatocytes, the HDV nucleoprotein is released and translocated by HDAg to the nucleus via a nuclear localization signal.<sup>3</sup> The host range of HDV is limited to those species that can support the replication of hepadnaviruses and supply, in *trans*, the HBsAg envelope. For instance, HDV can infect woodchucks and can be packaged with HBsAg derived from woodchuck hepatitis virus. Chimpanzees are susceptible to HDV infection, and the infection is similar to that in humans.<sup>12</sup>


**RNA Genome and Replication**

The HDV RNA genome does not encode protein. Instead, the complementary antigenomic HDV RNA codes for HDAg and contains a sequence of approximately 85 nucleotides with an intrinsic ribozyme activity.<sup>13,14</sup> It is estimated that about 300,000 copies of genomic RNAs can be found in an infected hepatocyte during replication and 10-fold-less antigenomic RNA. S-HDAg is believed to be involved in RNA replication and may serve as an RNA chaperone.<sup>15</sup>

HDV replication involves the transcription of HDAg-encoding RNA from genomic RNA, the synthesis of antigenomic RNA from genomic RNA, and the synthesis of genomic RNA from antigenomic template via a double rolling circle mechanism (Fig. 146.2). Because HDV does



**FIG. 146.1** Schematic structures of hepatitis D virus genomic and antigenomic RNA and hepatitis D antigen (HDAg)-encoding messenger RNA (mRNA). The minimum ribozymes are indicated by light boxes. The italicized numbers for HDAg are amino-acid residues. All other numbers are nucleotide positions on the genomic-sense RNA. The L-HDAg and S-HDAg are encoded by the antigenomic strand indicated by the open reading frame (ORF). (Modified from MacNaughton TB, Lai MC. *The molecular biology of hepatitis delta virus*. In: Ou JE, ed. *Hepatitis Viruses*. Norwell, MA: Kluwer; 2002:109–128.)



**FIG. 146.2** Proposed model of hepatitis D virus RNA replication. The hepatitis D antigen-encoding messenger RNA (mRNA) is synthesized by polymerase II (Pol II) from the genomic RNA template independently of RNA replication. RNA Pol I synthesizes antigenomic RNA, and RNA Pol II synthesizes the genomic RNA. AG, Antigenomic strand; G, genomic strand. (Modified from MacNaughton TB, Lai MC. *The molecular biology of hepatitis delta virus*. In: Ou JE, ed. *Hepatitis Viruses*. Norwell, MA: Kluwer; 2002:109–128.)

not encode a polymerase, it recruits the host RNA polymerase II (Pol II) for transcription of HDV mRNA and for genomic RNA synthesis and likely uses host Pol II, and possibly also host Pol I, for antigenomic RNA synthesis.<sup>10,16</sup> Because of the rolling circle mechanism of RNA synthesis, multimeric linear transcripts are produced that are cleaved into monomers by the ribozyme activity. The linear product is ligated to form the antigenomic molecule, which then serves as a template for the synthesis of genomic RNAs through a rolling circle mechanism; HDV then interacts with the clamp of RNA Pol II, a cellular structure that holds RNA and DNA in place. HDAg loosens the clamp, which facilitates forward translocation of RNA Pol II while sacrificing fidelity.<sup>17</sup> HDV envelopment follows the scheme of virion assembly similar to

HBV. Because HDV assembly can also occur in an L-HBsAg-independent fashion,<sup>18</sup> noninfectious HDV virions can be produced in abundance (≈90%). These particles are noninfectious because of the absence of pre-S1 domain, which is required for infectivity.

## Pathogenesis

HDAg is the only viral protein known to be expressed during HDV infection. A high titer of immunoglobulin (Ig)M anti-HDV is strongly associated with elevated hepatitis D viremia and the severity of liver injury and alanine aminotransferase (ALT)/aspartate aminotransferase levels, whereas a more favorable course to HDV infection is found in individuals with IgG anti-HDV.<sup>19</sup> Although these antibody responses were present during acute and chronic infection, there was no association of IgM anti-HDV levels with HDV RNA, suggesting there is no protective role of anti-HDV antibodies. Woodchucks immunized with recombinant HDAg are only partially protected from subsequent challenge with HDV, which suggests that other mechanisms are responsible for immunity.<sup>20</sup>

The mechanisms of liver damage in HDV infection are unclear. There is conflicting evidence regarding whether HDV has a direct viral cytopathic effect that contributes to hepatocyte injury. In cell lines, expression of HDAg led to cytotoxic changes<sup>21</sup>; however, in transgenic mice, HDAg expression had no cytopathic effect.<sup>22</sup> HDV RNA levels have not been associated with severity of liver disease in 80 individuals with genotype 1 disease, supporting the concept that HDV does not directly lead to hepatocyte injury.<sup>23</sup> It is likely that the adaptive immune response plays a role in liver injury; however, the data are also conflicting. In one study HDV-specific T-cell responses correlated with lower ALT levels, suggesting that immune control of viral replication leads to lesser degrees of liver injury.<sup>24</sup> However, in another study histologic assessment demonstrated that the degree of cellular infiltration in the portal tracts and lobules correlates with the degree of staining for HDAg in the liver, suggesting that the immune response contributes to hepatocellular injury.<sup>25</sup> Thus the etiology of liver injury in HDV infection is unclear.

The adaptive immune response plays an important role in control of HDV infection, but the precise mechanisms are not understood. HDV-specific CD8 T-cell responses were detectable in those with previous infection but not in chronic infection.<sup>9,26</sup> Another study demonstrated that chronic HDV was associated with a Th-2 response and secretion of interleukin-10. Taken together, these suggest that control of an HDV infection require Th1 CD4 and CD8 T-cell responses.

## Epidemiology

An estimated 15 to 20 million people have serologic evidence of exposures to HDV, with an overall global prevalence of approximately 5% of HBV-infected persons.<sup>27</sup> Since the 1980s the prevalence of HDV has declined substantially as a result of better control of HBV due to more widespread vaccination practices and improved socioeconomic conditions. Infection with HDV has a worldwide distribution, although there are considerable geographic differences that do not entirely mirror the prevalence of HBV infection. Prevalence in the United States and Northern Europe are about 5% to 15% in HBsAg-positive carriers.<sup>28</sup> Immigrants from the Middle East, Africa, and Eastern Europe, and older patients from the Mediterranean Basin (most acquired parenterally), as well as young injection drug users account for the largest proportion of cases.<sup>29–31</sup> In the developing world, where HBV is highly endemic, such as Asia and sub-Saharan Africa, as many as two-thirds of the HBV-infected population is infected with HDV, although data in several countries are lacking. Some areas with the highest prevalence include parts of Central and West Africa (33% in Mauritania, 67% in Gabon), parts of the Amazon Basin (42%), the Middle East (33%–47%), India, Pakistan (24%–89%), and Northern Vietnam (15%–43%). In Mongolia greater than 60% to 88% of people with chronic hepatitis B have evidence of chronic HDV.<sup>32</sup> Higher HDV prevalence is also seen in patients with HIV-HBV coinfection compared with HBV monoinfection.<sup>33</sup> It should be noted that many of the estimated prevalence data are based on anti-HDV alone because commercial assays for HDV RNA are not readily available in many countries. In a recent study from Tanzania the prevalence of anti-HDV positivity was 5%; however, HDV RNA was not detectable in any of the anti-HDV-positive patients, and a



second serologic test (HDV-Ab kit; DiaPro, Milan, Italy) was negative in all the samples.<sup>34</sup>

There are eight HDV genotypes that have been identified by phylogenetic analyses. Divergence between genotypes ranges from 20% to 40%. Some of the genotypes are more aggressive than others.<sup>35</sup> HDV-1 strain is the most common worldwide, present in almost all continents. Four HDV-1 subgenotypes have been identified, and most have been associated with severe disease, including cirrhosis, fulminant hepatitis, and death. HDV-2 and HDV-4 strains are found in Asia and are associated with milder disease. HDV-3 is found in northern South America, especially the Amazon Basin, and is associated with the most severe form of HDV.<sup>36,37</sup> HDV-5 to HDV-8 are present only in Africa. HDV-5 is the next most common genotype after HDV-1 and originates from western and sub-Saharan Africa<sup>35</sup> (Fig. 146.3).

## Diagnosis

HDV is diagnosed with detection of anti-HDV IgG, and because this remains positive with HDV clearance, chronic infection can be established using real-time polymerase chain reaction.<sup>38</sup> HDV IgM antibodies have a role if the HDV RNA is negative, and there is still a high suspicion for chronic HDV because viremia can fluctuate over time, and the variability of the HDV sequence can produce false-negative results.

## Transmission and Clinical Manifestations


Similar to HBV, HDV is spread by blood, blood products, and bodily secretions and only requires a small inoculum to transmit.<sup>9</sup> Unlike HBV, perinatal transmission is uncommon. As a result of universal blood screening for HBsAg and widespread HBV vaccination, HDV has virtually disappeared in multiply transfused subjects and hemophiliacs in developed countries.<sup>39</sup> After exposure there is a short incubation period of 3 to 7 weeks. Because HDV requires the surface antigen from HBV, disease may occur as an acute coinfection with HBV or as a superinfection of a chronic HBV infection.

As with acute hepatitis B mono-infection, the clinical presentation and natural history of acute coinfection with HBV and HDV are highly variable. Acute coinfection usually occurs after an incubation period of 45 to 160 days. The incubation period during coinfection may display a biphasic pattern of ALT levels because the titers of the infecting inoculum for the two viruses may be different or because the two viruses

are expressed sequentially.<sup>12,40</sup> In animal models the incubation period of each virus has been shown to be inversely proportional to the dose of the virus.<sup>41</sup> Usually the first episode is due to hepatitis B replication and immune response, followed by that of hepatitis D. Clinically, coinfection presents as an acute self-limited hepatitis with complete recovery with only 4% to 5% progressing to chronic HBV and HDV coinfection.<sup>42–44</sup> In rare instances a severe or fulminant hepatitis can occur.<sup>45</sup>

Superinfection of HDV in individuals with chronic hepatitis B generally results in acute severe hepatitis with a relatively short incubation period (2–8 weeks), followed by chronic hepatitis D in 90% of the cases. Of interest, when HDV clearance occurs after superinfection, there is increased HBsAg loss compared with HBV mono-infection.<sup>46</sup> Superinfection with HDV can be associated with fulminant hepatitis and chronic active hepatitis with cirrhosis. Fulminant hepatitis, a severe form of acute hepatitis, is 10 times more common in HBV-HDV coinfection than in chronic HBV infection alone. The prognosis with fulminant hepatitis is very poor. Massive liver necrosis can occur with mortality approaching 80% in the absence of liver transplantation. In patients with chronic hepatitis D, HDV is the dominant virus because the L-HDAg suppresses HBV replication (see “Hepatitis Delta Antigen”). Thus most HBV-HDV coinfecting patients have low serum HBV DNA levels, and most are HBeAg negative.

Once chronic HDV infection is established, the clinical course of hepatitis is accelerated compared with HBV mono-infection. Cirrhosis occurs in 60% to 80% of chronic hepatitis D patients within 5 to 10 years, and the risk of hepatocellular carcinoma (HCC) is about threefold higher than in HBV mono-infection.<sup>47–49</sup> In one study where the HDV seroprevalence was 6%, the relative risk of cirrhosis and HCC was 2.58 and 2.87, respectively, in patients with both infections compared with those with HBV alone.<sup>50</sup> The high risk of HCC may reflect the higher rate of cirrhosis. Once cirrhosis is established, the risk for clinical decompensation is twofold higher than in HBV mono-infection.<sup>47</sup> Splenomegaly along with elevated aminotransferases and high levels of viremia are common in chronic hepatitis D.<sup>47</sup> However, unlike HBV, the level of HDV viremia does not seem to correlate with severity of disease.<sup>48,51</sup> The clinical course is affected by both the HDV genotype (see “Epidemiology”) and the HBV genotype.<sup>47,52</sup> Liver disease is known to be accelerated in HIV-infected patients with triple coinfection (HIV/



**FIG. 146.3** Worldwide distribution of hepatitis D virus (HDV) genotypes and subgenotypes. (From Le Gal F, Brichler S, Drugan T, et al. Genetic diversity and worldwide distribution of the Deltavirus genus: a study of 2,152 clinical strains. *Hepatology*. 2017;66:1826–1841.)

HBV/HDV), and the incidence of HCC and mortality is higher than in patients with HIV alone.<sup>53,54</sup>

Up to 30% of HBV-HDV coinfecting individuals also have hepatitis C virus (HCV) antibodies; however, in one study 84% of them had undetectable HCV RNA, suggesting that HDV is the dominant virus in this triple infection.<sup>55</sup>

## Treatment

Interferon- $\alpha$  (IFN- $\alpha$ ) and pegylated (PEG) IFN- $\alpha$  have been shown to have antiviral activity against HDV. However, the efficacy of IFN- $\alpha$  in the treatment of HDV is limited unless high doses (9 million units three times a week) are used as in the reported studies. In one of the largest trials to date, treatment with IFN- $\alpha$  was only successful in achieving sustained HDV virologic response in 25% to 30% of individual patients.<sup>56</sup> With PEG IFN- $\alpha$ , sustained virologic response has been variable, occurring in 17% to 43% of patients in clinical trials. The current recommended treatment for chronic hepatitis D is weekly PEG IFN- $\alpha$  for a minimum of 48 weeks. The ultimate duration of therapy for HDV is still not known (also see Chapter 47), and late relapses have been reported.<sup>57</sup> In the long term, biochemical and histologic improvements with IFN- $\alpha$  can occur, even among those who relapse. In a previous study, improvement in liver histology was maintained 10 years post-treatment among the patients who received high-dose IFN- $\alpha$ .<sup>58</sup> Nucleoside reverse-transcriptase inhibitors with activity against HBV, including lamivudine and entecavir, have been shown to be ineffective at controlling HDV replication, which is not surprising because HDV does not have its own polymerase.<sup>59</sup> HDV RNA declines, and virologic suppression have been reported in HIV/HBV/HDV coinfecting patients receiving long-term tenofovir DF, correlating in one study with declines in HBsAg and reduction in fibrosis.<sup>60</sup> This decline may have been related to the longer duration of therapy that led to decrease in HBsAg.

Newer therapies are under development and fall into three categories: entry inhibitors, farnesyltransferase inhibitors, and nucleic acid polymers.

Blocking HDV entry with Myrcludex B  $\pm$  PEG IFN- $\alpha$  has shown promise in phase II clinical trials.<sup>61</sup> Farnesyltransferase inhibitors (lonafarnib) prevent virion assembly, and in early phase II clinical trials, a reduction in HDV RNA was observed over 28 days.<sup>62</sup> Longer studies up to 24 weeks with and without ritonavir as a boosting agent and/or PEG IFN- $\alpha$  have led to HDV RNA suppression at the end of treatment but not necessarily off treatment. Gastrointestinal toxicity is a dose-limiting side effect of lonafarnib. The mechanism of action of nucleic acid polymers is believed to be decreasing viral secretion from hepatocytes. One example is intravenous REP 2139 (Replicor; Montréal, Quebec, Canada), which was given in combination with PEG IFN- $\alpha$  to 12 HBV-HDV coinfecting patients in a phase II clinical trial.<sup>63</sup> Patients received 30 weeks of REP 2139 and 48 weeks of PEG IFN- $\alpha$  with a 15-week overlap of the two drugs. One year after the end of treatment, 7 were HDV RNA negative and 8 were HBV DNA negative. Four individuals had a serious adverse event, which was attributed to the PEG IFN- $\alpha$ , but only 1 discontinued therapy. Other treatments under investigation include PEG IFN- $\gamma$ , which is being tested at two doses for 48 weeks in 33 individuals. Interim 24-week results from 10 of the 33 participants showed that five had a  $\geq 2$ -log decline in HDV RNA, and 4 became HDV RNA negative.<sup>64</sup> Further studies with these and other agents in development are needed to determine if they are superior to PEG IFN- $\alpha$ .

## Transplantation

Liver transplantation is reserved for patients with end-stage liver disease from HBV-HDV coinfection. Reinfection of the graft is prevented with administration of hepatitis B immunoglobulin because prevention of HBV reinfection stops propagation of HDV. There is no evidence for latency of HDV outside the liver. The 5-year survival of liver transplantation for HDV is  $>80\%$ .

## Key References

The complete reference list is available online at Expert Consult.

3. Botelho-Souza LF, Vasconcelos MPA, Dos Santos AO, et al. Hepatitis delta: virological and clinical aspects. *Virology*. 2017;14:177.
16. Sureau C, Negro F. The hepatitis delta virus: replication and pathogenesis. *J Hepatol*. 2016;64:S102–S116.
27. Noureddin M, Gish R. Hepatitis delta: epidemiology, diagnosis and management 36 years after discovery. *Curr Gastroenterol Rep*. 2014;16:365.
29. Rizzetto M, Ciancio A. Epidemiology of hepatitis D. *Semin Liver Dis*. 2012;32:211–219.
33. Calle Serrano B, Manns MP, Wedemeyer H. Hepatitis delta and HIV infection. *Semin Liver Dis*. 2012;32:120–129.
35. Le Gal F, Brichler S, Drugan T, et al. Genetic diversity and worldwide distribution of the *Deltavirus* genus: a study of 2,152 clinical strains. *Hepatology*. 2017;66:1826–1841.
52. Yurdaydin C. Recent advances in managing hepatitis D. *F1000Res*. 2017;6:1596.
58. Farci P, Mandas A, Coiana A, et al. Treatment of chronic hepatitis D with interferon alfa-2a. *N Engl J Med*. 1994;330:88–94.
62. Koh C, Canini L, Dahari H, et al. Oral prenylation inhibition with lonafarnib in chronic hepatitis D infection: a proof-of-concept randomised, double-blind, placebo-controlled phase 2A trial. *Lancet Infect Dis*. 2015;15:1167–1174.
63. Bazinet M, Pântea V, Cebotarescu V, et al. Safety and efficacy of REP 2139 and pegylated interferon alfa-2a for treatment-naïve patients with chronic hepatitis B virus and hepatitis D virus co-infection (REP 301 and REP 301-LTF): a non-randomised, open-label, phase 2 trial. *Lancet Gastroenterol Hepatol*. 2017;12:877.