

at the maximal tolerated doses of 1 to 1.5 mg/kg/day.¹⁸⁹ Unfortunately, high doses of conventional amphotericin B are usually not tolerated for more than several days before renal function deteriorates, especially in patients with diabetes or receiving concomitant nephrotoxic therapies.

Lipid Amphotericin B Formulations

Lipid formulations of amphotericin B are safer than ABD for long-term administration and, in our opinion, are the preferred first-line treatment for severe mucormycosis.^{187,190} Most experience in the treatment of mucormycosis has been with either liposomal amphotericin B (L-AMB) or amphotericin B lipid complex (ABLC). Although no comparative studies have been performed, outcomes with the use of the two lipid amphotericin B formulations are similar to those historically reported for conventional ABD, albeit with lower rates of nephrotoxicity.¹⁸⁹

In one of the few published case series, 24 patients with mucormycosis and diabetes as the predominant underlying risk factors were treated with ABLC after failure or intolerance of the conventional ABD formulation.^{190,191} The overall response rate (improvement or cure of infection) was 71% (17/24 patients), with few reported toxic effects, even in patients with preexisting renal dysfunction. Several case series have reported the successful treatment of mucormycosis with the liposomal formulation of amphotericin B, sometimes administered at high doses (i.e., 10 mg/kg/day) for prolonged treatment courses.^{192,193}

The optimal dosing approach for L-AMB in patients with mucormycosis is frequently debated. Preclinical pharmacokinetics/pharmacodynamics studies in murine models of pulmonary mucormycosis that simulated human dosing have suggested that ABLC at 5 mg/kg/day or L-AMB at 10 mg/kg/day results in rapid antifungal accumulation in the lung, reduced fungal burden, and improved survival.¹⁹⁴ However, these animal models focus on short treatment periods and do not routinely assess nephrotoxicity of higher-dose L-AMB regimens. Both clinical and observational studies have suggested an early “window” of 10 to 14 days when nephrotoxicity risk is lowest with L-AMB, which may be shortened in patients receiving concomitant nephrotoxic agents or aggressive diuresis.^{195–197}

In one of the few randomized trials that compared standard doses of L-AMB (3 mg/kg/day) to a higher dose-regimen (10 mg/kg/day for 14 days, then 3 mg/kg/day) for invasive aspergillosis, patients randomized to the higher-dosed L-AMB regimen failed to achieve higher response rates but experienced significantly higher rates of nephrotoxicity and severe hypokalemia.¹⁹⁸

The feasibility of high-dose (10 mg/kg/day) L-AMB treatment for the initial treatment of mucormycosis was explored in a multicentric prospective French study of 40 patients with invasive mucormycosis.¹⁹⁹ The planned treatment of 10 mg/kg/day was administered as an infusion of at least 2 hours for 4 weeks. In patients in whom there was doubling of serum creatinine levels, compared with baseline, it was recommended that the L-AMB dose be reduced to 7.5 mg/kg/day. If creatinine values did not improve within 3 days, L-AMB dosing was further reduced to 5 mg/kg/day. Among the 33 patients who could be analyzed at 4 weeks after starting therapy or end of therapy, a favorable response was documented in 12 of 33 (36%) at week 4 and 14 of 31 (45%) by week 12. Most patients (24/31, 71%) underwent at least one surgical procedure. Of importance, although 4 weeks of high-dose L-AMB was planned, the average treatment duration at this dose was 13.5 days (range, 0–28), with 40% of patients experiencing a doubling in the baseline serum creatinine and severe hypokalemia (serum potassium <3 mmol/L). Therefore considerations for using doses of L-AMB higher than 5 mg/kg/day should take into account that a sizable proportion of patients will develop renal injury requiring dosage reduction or possibly a switch to triazole therapy after the first 1 to 2 weeks of therapy. Daily infusions of a liter of saline are used in some centers to reduce L-AMB nephrotoxicity in adults who can tolerate an increased sodium load.

Triazoles

The prospects for using posaconazole in the treatment of mucormycosis has improved with the introduction of an extended-release tablet formulation with improved bioavailability compared with the older suspension, and an IV formulation solubilized in sulfobutyl ether β -cyclodextrin. These newer formulations are dosed differently than the oral suspension,

which has to be administered two to four times daily in smaller doses to maximize absorption. The extended-release tablet and IV formulation of posaconazole are the preferred formulations of posaconazole for most patients due to their more reliable pharmacokinetics properties.²⁰⁰

Most of the published clinical experience concerning mucormycosis treatment with posaconazole has used the older suspension formulation administered at 800 mg/day in divided doses. In an open-label study evaluating posaconazole as salvage therapy (not initial therapy), the overall success rate of posaconazole (800 mg/day) was 70% in 24 patients, and it was well tolerated with only minimal GI side effects.^{201,202} Similarly, a retrospective survey of posaconazole-based salvage therapy in 91 patients with refractory mucormycosis indicated an overall success rate of 61%, including 65% in patients with pulmonary mucormycosis.²⁰³ Among patients who were not categorized as treatment successes, 21% had evidence of stable disease after 12 weeks of treatment. At present the US Food and Drug Administration (FDA) has not approved posaconazole for primary or salvage therapy of mucormycosis, indicating the need for further studies.

The extended-release tablet formulation of posaconazole is administered at a dose of 300 mg (three 100-mg delayed-release tablets) twice a day on the first day, then 300 mg daily. Although the absorption of the tablets is improved with food, the tablets have adequate bioavailability even in patients with poor dietary intake or receiving acid-suppression therapy, such as a proton pump inhibitor.²⁰⁴ The IV formulation of posaconazole is dosed at 300 mg IV twice daily on day 1, followed by 300 mg daily thereafter, and rapidly achieves plasma concentration exposures of greater than 500 ng/mL in 95% of patients, which often took more than a week in patients receiving posaconazole suspension.²⁰⁵ Although therapeutic drug monitoring has been recommended for posaconazole in *Aspergillus* treatment guidelines to ensure serum trough levels greater than 1.5 mg/L during the treatment of infection,^{206,207} a similar relationship between serum trough levels of posaconazole and outcome of mucormycosis has not been reported.

In 2015 the FDA approved isavuconazole for treatment of invasive mucormycosis based on an open observational report by the manufacturer²⁰⁸ of 21 patients who received isavuconazole as primary therapy for proven or probable disease; 7 had the drug discontinued, 2 were still on therapy when reported, 6 died during isavuconazole treatment, and 6 were alive when isavuconazole treatment was considered completed. Results with salvage therapy in 16 patients were also considered favorable. The small numbers and diverse results make comparisons with amphotericin B and posaconazole difficult.²⁰⁹ Isavuconazole may be a possible alternative to posaconazole or L-AMB, particularly for longer-term therapy. An advantage of isavuconazole is that it is administered as a prodrug formulation (isavuconazonium) in either an IV and oral formulation with excellent bioavailability that is not affected by dietary intake or acid-suppression therapy. Isavuconazole is dosed with a 372-mg loading dose administered every 8 hours for 6 doses (48 hours), then a 372-mg maintenance dose daily. Isavuconazole is generally well tolerated, making it a possible alternative for patients who are clinically stable and have limiting drug interactions with posaconazole or cannot tolerate the nephrotoxic effects of L-AMB. However, some species, such as *M. circinelloides*, exhibit MICs greater than achievable serum levels of patients with currently recommended isavuconazole dosing.¹⁷² We have also observed cases of breakthrough mucormycosis during isavuconazole treatment in patients with prolonged neutropenia and relapsed hematologic malignancies.

Combination Therapy

Successful treatment of mucormycosis with combinations of amphotericin B, terbinafine, rifampicin, L-AMB, posaconazole, and echinocandins has also been described in small case series and case reports.^{183,210} Interpretation of these studies is confounded by small numbers, retrospective nature, and confounding factors such as neutrophil recovery and surgery.^{210,211} One of the largest case-control studies evaluating the possible benefits of combination therapy for invasive mucormycosis was performed at MD Anderson Cancer Center,²¹² which identified 47 patients who received early L-AMB monotherapy and 59 patients who received combination treatment. The most common combination regimens were L-AMB plus an echinocandin (46%), L-AMB plus

posaconazole (27%), or triple combination therapy (27%). A propensity score analysis for combination therapy was used to attempt to adjust for confounding variables associated with prescription of combination therapy. The investigators could not identify any survival benefit for combination therapy over timely administration of L-AMB, even after propensity score adjustment (odds ratio, 0.8; 95% confidence interval, 0.3 to 2.4; $P = .69$) or if patients were stratified into groups for low versus high risk for mortality.²¹²

Treatment Duration

The duration of treatment required for mucormycosis is highly individualized to the patient. Near normalization of radiographic imaging, negative biopsy specimens, and cultures from the affected site and recovery from immunosuppression are indicators that a patient is a candidate for stopping antifungal therapy. Late relapses of mucormycosis after successful treatment have been reported several years after discontinuation of secondary posaconazole prophylaxis or onset of new immunosuppression.²¹³ In one case relapsing disease was accompanied by increased ¹⁸F-fluorodeoxyglucose uptake in lesions detected by positron emission tomography–CT.²¹³ Therefore continued follow-up of patients is critical in any patient who discontinues treatment for mucormycosis.

Prophylaxis

Because mucormycosis is a relatively rare infection, primary prophylaxis is generally not recommended. Secondary prophylaxis is often desired in patients requiring further immunosuppression after treatment for mucormycosis. Posaconazole, and possibly isavuconazole, appears to be a safe option for patients who require continuous, oral long-term antifungal therapy because they remain at high risk for relapsing infection.²¹³

Surgical Management

In rhinosinusitis surgical débridement of infected tissue is a crucial component of therapy and should be urgently performed to limit the aggressive spread of infection to contiguous structures. Repeated removal of necrotic tissue or radical surgical resection (e.g., exenteration of the orbit) with subsequent reconstructive surgeries may be required for lifesaving control of rapidly evolving infection.¹¹⁸ However, rhinosinusitis has been treated successfully in select patients without radical resection, particularly when multimodality treatment options are used with careful follow-up.²¹⁴ Extension to the brain usually portends a fatal outcome. Decisions regarding the extent of surgical débridement are highly individualized to the patient. Intraoperative frozen sections and MRI can help determine the extent of involved tissue and tissue margins. Conditions such as low platelet counts and other bleeding problems must be corrected with sufficient transfusions before surgical intervention. Unfortunately, bleeding risks may limit surgical options in some patients with profound thrombocytopenia.

In patients with pulmonary mucormycosis, surgical treatment in conjunction with systemic antifungal therapy has been shown to significantly improve survival compared with antifungal therapy alone.^{118,215} One large case series reported a mortality rate of 55% in patients who received systemic antifungal therapy alone compared with 27% in patients who received antifungal therapy plus surgical intervention.¹¹⁰ Although the reported outcomes may reflect a selection bias in offering surgery to less severely ill patients with unifocal disease, removal of infected or devitalized tissue early, when the infection is localized, provides the greatest benefit. Cavitation of lesions near the hilum has been followed by fatal hemoptysis, providing an incentive for resection in selected patients. Repeated surgeries are also necessary in some cases. The benefit of pulmonary resection of a dominant lung lesion (debulking) is unknown in patients with multifocal or disseminated mucormycosis.

Adjunctive Therapies

Patients with profound neutropenia and progressive mucormycosis despite optimal therapy may be candidates for neutrophil transfusion. With currently available techniques, more than 10^{10} granulocytes can be infused, resulting in an immediate postinfusion absolute neutrophil count usually exceeding 1000/ μ L. Repeated infusions are needed every

few days and can lead to diminishing response in the patient's neutrophil count due to allosensitization. Success of this bridging procedure in treating mucormycosis is still uncertain but likely depends ultimately on the patient's marrow recovery.

Several other adjunctive measures have been explored for improving tissue viability, impeding fungal proliferation, and improving host immunity. Hyperbaric oxygen therapy was reported to be a beneficial adjunct to standard surgical and antifungal therapy for mucormycosis, particularly for diabetic patients with rhinocerebral disease.^{216–218} Although this is not one of the approved uses of hyperbaric oxygen (see Chapter 50), the increased oxygen pressure achieved with hyperbaric oxygen may improve neutrophil activity and the putative oxidative killing effects of polyene antifungals.²¹⁹ High oxygen concentrations have also been reported to inhibit growth of Mucorales in vitro and to improve the rate of wound healing by increasing the release of tissue growth factors.²²⁰ Lack of convincing clinical evidence supporting the benefit of hyperbaric oxygen therapy, however, limits its recommendation for routine clinical use for this expensive, controversial, and logistically difficult intervention for mucormycosis.⁷¹

Multiple immune-augmentation strategies have been proposed for mucormycosis, including administration of cytokines that enhance phagocytic activity, such as granulocyte colony-stimulating factor, granulocyte-macrophage colony-stimulating factor, or IFN- γ alone or in combination with granulocyte transfusions. Various combinations of these approaches have been reported with some favorable outcomes in case reports.^{219,221} However, many of these immune-augmentation strategies carry some risk for enhancing inflammatory lung injury^{222,223}; therefore the relative benefits of such adjunctive strategies must be balanced against the risk for increased harm to the patient.

Prognosis

The site of infection and underlying host factors are the key prognostic determinants of mucormycosis outcome. Active hematologic malignancy, allogeneic HSCT, and disseminated infection are associated with poor outcome.¹⁶⁶ In our experience, most patients with hematologic cancer who develop mucormycosis die within 12 weeks of diagnosis.^{121,212,224} However, earlier diagnosis of the disease and aggressive treatment have been associated with improved survival rates in recent series.^{166,212} Correction of underlying immune impairment (e.g., rapid tapering of glucocorticoids), combined with aggressive multimodality treatment approaches, offer the best chance for patient survival.

ENTOMOPHTHORAMYCOSIS

Infections caused by fungi in the subphylum Entomophthoromycotina, called *entomophthoromycosis*, include both conidiobolomycosis and basidiobolomycosis. These are rare infections of the paranasal sinus and subcutaneous tissues, principally encountered in the tropics, that rarely affect other tissues.²²⁵ In contrast to mucormycosis, entomophthoromycosis is usually a chronic, nonangioinvasive infection in relatively immunocompetent individuals.^{226,227} Although rare, case clusters of invasive disease have been reported in both immunocompromised and immunocompetent patients.^{228,229} Conidiobolomycosis affects primarily the head and face, whereas basidiobolomycosis is often localized to the subcutaneous tissues of the trunk and arms or the GI tract.^{226,227} The infections are characterized by slow-growing, tumor-like masses in infected tissues that can remain indolent for years. Both fungi are common inhabitants of the soil throughout the world, including the United States. However, most cases of conidiobolomycosis are found in tropical Africa, South America, Central America, and Asia.²²⁶ Similarly, reports of cutaneous basidiobolomycosis are primarily concentrated in tropical areas of Africa and Southeast Asia and in the tropical and subtropical regions of Asia, Australia, and South America.²²⁷ Entomophthoromycosis also demonstrates some age specificity: conidiobolomycosis is uncommon in children, but 88% of basidiobolomycosis cases occur in patients younger than 20 years.²³⁰

Conidiobolomycosis

Subcutaneous rhinofacial conidiobolomycosis is the most common manifestation of infection caused by *Conidiobolus coronatus*. Symptoms typically begin with nasal discharge, epistaxis, unilateral nasal obstruction,

sinus tenderness, and extensive and persistent facial swelling that may result in disfigurement. The infection slowly progresses with granulomatous inflammation in the subcutaneous tissue without bone involvement or ulceration of the skin.²³¹ Systemic symptoms are rare, but disseminated conidiobolomycosis has been observed.²²⁶ Infections with *Conidiobolus incongruus* are extremely rare but often more aggressive.¹⁵⁹

Basidiobolomycosis

Infections caused by *Basidiobolus ranarum* often begin as a nodular subcutaneous lesion on the trunk, arms, or buttocks. The mode of transmission for *B. ranarum* is assumed to be through minor trauma and insect bites. Fungal spores are found in the bristles of mites and are probably carried by other insects. *B. ranarum* has been theorized to be present on "toilet leaves" used for skin cleaning after a bowel movement, resulting in direct inoculation of the perineum.²³² The predominance of lesions in the buttocks, thighs, and perineum would appear to support this theory. The subcutaneous lesions elicited by *B. ranarum* are typically firm but not painful, with edema around the involved sites. Deeper invasion of muscle underlying the subcutaneous tissue has been reported.²²⁷ Several cases have been reported of otherwise healthy persons with GI tract basidiobolomycosis, most often in the colon,²³³ that may present as obstruction or mimic the presentation of Crohn disease.^{232,234,235} Clinical features of GI basidiobolomycosis include abdominal pain, nausea, vomiting, diarrhea, or abdominal mass.²³⁵ Peripheral blood eosinophilia and high serum immunoglobulin E concentrations are usual. Most cases are slowly progressive locally and may be mistaken for Crohn disease or form large abdominal masses mimicking malignancy. Chronic fibrotic-appearing lesions, including angioinvasive disease reminiscent of mucormycosis seen in diabetic and immunocompromised patients, have also been described.²²⁹

Diagnosis

In areas where entomophthoromycosis is common the diagnosis is often suspected from the clinical appearance of the patient and characteristic lesions or swelling.^{232,235} Definitive diagnosis requires biopsy of the involved site, with characteristic findings of broad, sparsely septated

hyphae surrounded by eosinophilic granular material (Splendore-Hoeppli phenomena).^{232,236,237} Tissue eosinophilia and granulomatous inflammation are usual. Peripheral eosinophilia may also be present, but cultures from the infected site are often negative. PCR assay of DNA extracted from tissue or from the isolated fungus, followed by sequencing, may be the most useful confirmatory diagnostic test.^{13,233}

Differential Diagnosis

Eumycetomas have some clinical similarities to entomophthoromycosis, presenting as a chronic granulomatous fungal disease that affects the limbs, most commonly the feet (mycetoma pedis or Madura foot), and on occasion the abdomen, chest, and head, typically with draining sinuses. Eumycetomas can be differentiated by the presence of grains surrounded by neutrophils on biopsy and by culture. Pythiosis is usually a disease of horses, cattle, dogs, and cats but is a rare cause of subcutaneous disease in humans that can resemble entomophthoromycosis or can cause GI or disseminated infection resembling mucormycosis. Although the histologic appearance of the hyphae is similar, presence of vascular invasion and the appearance of motile zoospores on water culture of the causative organism, *Pythium insidiosum*, can distinguish the entities.¹³

Therapy and Prevention

Conidiobolus spp. are generally more resistant to systemic antifungals than *Basidiobolus* spp.²³⁸ In the older literature agents used to treat entomophthoromycosis included potassium iodide, trimethoprim-sulfamethoxazole, ketoconazole, itraconazole, and amphotericin B with varying success and clinical outcome.^{226,227,239–241} At present the preferred drugs for rhinofacial conidiobolomycosis appear to be a combination of a saturated solution of potassium iodide and itraconazole.²⁴¹ Surgical removal and reconstructive surgery for grossly swollen or disfigured tissues, combined with medical therapy, often provides the best chance for complete recovery. Subcutaneous basidiobolomycosis has been reported to respond to itraconazole.²⁴² Response of GI basidiobolomycosis to mold-active azoles has generally been favorable in small numbers of cases, including itraconazole, posaconazole, or voriconazole.^{241,243–245}

Key References

The complete reference list is available online at Expert Consult.

- Sivagnanam S, Sengupta DJ, Hoogstraal D, et al. Seasonal clustering of sinopulmonary mucormycosis in patients with hematologic malignancies at a large comprehensive cancer center. *Antimicrob Resist Infect Control*. 2017;6:123.
- Tribble DR, Rodriguez CJ, Weintrob AC, et al. Environmental factors related to fungal wound contamination after combat trauma in Afghanistan, 2009–2011. *Emerg Infect Dis*. 2015;21:1759–1769.
- Chibucos MC, Soliman S, Gebremariam T, et al. An integrated genomic and transcriptomic survey of mucormycosis-causing fungi. *Nat Commun*. 2016;7:12218.
- Farmakiotis D, Kontoyiannis DP. Mucormycoses. *Infect Dis Clin North Am*. 2016;30:143–163.
- Pana ZD, Seidel D, Skiada A, et al. Invasive mucormycosis in children: an epidemiologic study in European and non-European countries based on two registries. *BMC Infect Dis*. 2016;16:667.
- Davoudi S, Graviss LS, Kontoyiannis DP. Healthcare-associated outbreaks due to Mucorales and other uncommon fungi. *Eur J Clin Invest*. 2015;45:767–773.
- Cheng VCC, Chen JHK, Wong SCY, et al. Hospital outbreak of pulmonary and cutaneous zygomycosis due to contaminated linen items from substandard laundry. *Clin Infect Dis*. 2016;62:714–721.
- Kontoyiannis DP, Yang H, Song J, et al. Prevalence, clinical and economic burden of mucormycosis-related hospitalizations in the United States: a retrospective study. *BMC Infect Dis*. 2016;16:730.
- Lamoth F, Chung SJ, Damonti L, et al. Changing epidemiology of invasive mold infections in patients receiving azole prophylaxis. *Clin Infect Dis*. 2017;64:1619–1621.
- Ibrahim AS, Voelz K. The mucormycete-host interface. *Curr Opin Microbiol*. 2017;40:40–45.
- Gebremariam T, Lin L, Liu M, et al. Bicarbonate correction of ketoacidosis alters host-pathogen interactions and alleviates mucormycosis. *J Clin Invest*. 2016;126:2280–2294.
- Baldin C, Ibrahim AS. Molecular mechanisms of mucormycosis—the bitter and the sweet. *PLoS Pathog*. 2017;13:e1006408.
- Lass-Flörl C, Aigner M, Nachbaur D, et al. Diagnosing filamentous fungal infections in immunocompromised patients applying computed tomography-guided percutaneous lung biopsies: a 12-year experience. *Infection*. 2017;45:867–875.
- Caillot D, Valot S, Lafon I, et al. Is it time to include CT “reverse halo sign” and qPCR targeting Mucorales in serum to EORTC-MSG criteria for the diagnosis of pulmonary mucormycosis in leukemia patients? *Open Forum Infect Dis*. 2016;3:ofw190.
- Kontoyiannis DP, Lewis RE. Treatment principles for the management of mold infections. *Cold Spring Harb Perspect Med*. 2015;5:a019737.
- Nam BD, Kim TJ, Lee KS, et al. Pulmonary mucormycosis: serial morphologic changes on computed tomography correlate with clinical and pathologic findings. *Eur Radiol*. 2018;28:788–795.
- Rodriguez-Lobato E, Ramirez-Hobak L, Aquino-Matus JE, et al. Primary cutaneous mucormycosis caused by *Rhizopus oryzae*: a case report and review of literature. *Mycopathologia*. 2017;182:387–392.
- Taj-Aldeen SJ, Gamaletsou MN, Rammaert B, et al. Bone and joint infections caused by Mucormycetes: a challenging osteoarticular mycosis of the twenty-first century. *Med Mycol*. 2017;55:691–704.
- Normand AC, Cassagne C, Gautier M, et al. Decision criteria for MALDI-TOF MS-based identification of filamentous fungi using commercial and in-house reference databases. *BMC Microbiol*. 2017;17:25.
- Cassagne C, Normand AC, Lollivier C, et al. Performance of MALDI-TOF MS platforms for fungal identification. *Mycoses*. 2016;59:678–690.
- Springer J, Goldenberger D, Schmidt F, et al. Development and application of two independent real-time PCR assays to detect clinically relevant Mucorales species. *J Med Microbiol*. 2016;65:227–234.
- Springer J, Lackner M, Ensinger C, et al. Clinical evaluation of a Mucorales-specific real-time PCR assay in tissue and serum samples. *J Med Microbiol*. 2016;65:1414–1421.
- Arendrup MC, Jensen RH, Meletiadis J. In vitro activity of isavuconazole and comparators against clinical isolates of the Mucorales order. *Antimicrob Agents Chemother*. 2015;59:7735–7742.
- Chowdhary A, Singh PK, Kathuria S, et al. Comparison of the EUCAST and CLSI broth microdilution methods for testing isavuconazole, posaconazole, and amphotericin B against molecularly identified Mucorales species. *Antimicrob Agents Chemother*. 2015;59:7882–7887.
- Dannaoui E. Antifungal resistance in Mucorales. *Int J Antimicrob Agents*. 2017;50:617–621.
- Espinel-Ingroff A, Alvarez-Fernandez M, Cantón E, et al. Multicenter study of epidemiological cutoff values and detection of resistance in *Candida* spp. to anidulafungin, caspofungin, and micafungin using the Sensititre YeastOne colorimetric method. *Antimicrob Agents Chemother*. 2015;59:6725–6732.
- Lamoth F, Damonti L, Alexander BD. Role of antifungal susceptibility testing in non-*Aspergillus* invasive mold infections. *J Clin Microbiol*. 2016;54:1638–1640.
- Gebremariam T, Alkharzi S, Baldin C, et al. Prophylaxis with isavuconazole or posaconazole protect immunosuppressed mice from pulmonary mucormycosis. *Antimicrob Agents Chemother*. 2017;61:e02589–16.
- Gebremariam T, Wiederhold NP, Alqarhi A, et al. Monotherapy or combination therapy of isavuconazole and micafungin for treating murine mucormycosis. *J Antimicrob Chemother*. 2017;72:462–466.
- Stanzani M, Vianelli N, Cavo M, et al. Liposomal amphotericin B nephrotoxicity in patients with hematological malignancies: a retrospective cohort analysis. *Antimicrob Agents Chemother*. 2017;61:e02651–16.
- Lanternier E, Poiree S, Elie C, et al. Prospective pilot study of high-dose (10 mg/kg/day) liposomal amphotericin B (L-AMB) for the initial treatment of mucormycosis. *J Antimicrob Chemother*. 2015;70:3116–3123.

200. Dekkers BGJ, Bakker M, van der Elst KCM, et al. Therapeutic drug monitoring of posaconazole: an update. *Curr Fungal Infect Rep*. 2016;10:51–61.
206. Patterson TF, Thompson GR 3rd, Denning DW, et al. Practice guidelines for the diagnosis and management of aspergillosis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;63:e1–e60.
208. Marty FM, Ostrosky-Zeichner L, Cornely OA, et al. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect Dis*. 2016;16:828–837.
209. Spellberg B, Brass E. The VITAL study: case control studies are hypothesis-generating. *Lancet Infect Dis*. 2016;16:886.
212. Kyvernitakis A, Torres HA, Jiang Y, et al. Initial use of combination treatment does not impact survival of 106 patients with haematologic malignancies and mucormycosis: a propensity score analysis. *Clin Microbiol Infect*. 2016;22:811.e811–811.e818.
241. Gupta M, Narang T, Kaur RJ, et al. A prospective case series evaluating efficacy and safety of combination of itraconazole and potassium iodide in rhinofacial conidiobolomycosis. *Int J Dermatol*. 2016;55:208–214.
242. Kumaravel S, Bharath K, Rajesh NG, et al. Delay and misdiagnosis of basidiobolomycosis in tropical South India: case series and review of the literature. *Paediatr Int Child Health*. 2016;36:52–57.
243. Mandhan P, Hassan KO, Samaan SM, et al. Visceral basidiobolomycosis: an overlooked infection in immunocompetent children. *Afr J Paediatr Surg*. 2015;12:193–196.

References

- Paltauf A. Mucormycosis: ein Beitrag zur Kenntnis der menschlichen Fadenpilz-krankungen. *Virchows Arch Pathol Anat.* 1885;102:543–564.
- Kwon-Chung KJ. Taxonomy of fungi causing mucormycosis and entomophthoromycosis (zygomycosis) and nomenclature of the disease: molecular mycologic perspectives. *Clin Infect Dis.* 2012;54 Suppl 1:S8–S15.
- Roden MM, Zaoutis TE, Buchanan WL, et al. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. *Clin Infect Dis.* 2005;41:634–653.
- Skiada A, Pagano L, Groll A, et al. Zygomycosis in Europe: analysis of 230 cases accrued by the registry of the European Confederation of Medical Mycology (ECMM) Working Group on Zygomycosis. *Clin Microbiol Infect.* 2011;17:1859–1867.
- Petrikos G, Skiada A, Drogari-Apiranthitou M. Epidemiology of mucormycosis in Europe. *Clin Microbiol Infect.* 2014;20 Suppl 6:67–73.
- Ribes JA, Vanover-Sams CL, Baker DJ. Zygomycetes in human disease. *Clin Microbiol Rev.* 2000;13:236–301.
- Sivagnanam S, Sengupta DJ, Hoogstraal D, et al. Seasonal clustering of sinopulmonary mucormycosis in patients with hematologic malignancies at a large comprehensive cancer center. *Antimicrob Resist Infect Control.* 2017;6:123.
- Shpitzer T, Keller N, Wolf M, et al. Seasonal variations in rhino-cerebral *Mucor* infection. *Ann Otol Rhinol Laryngol.* 2005;114:695–698.
- Neblett Fanfair R, Benedict K, Bos J, et al. Necrotizing cutaneous mucormycosis after a tornado in Joplin, Missouri, in 2011. *N Engl J Med.* 2012;367:2214–2225.
- Andresen D, Donaldson A, Choo L, et al. Multifocal cutaneous mucormycosis complicating polymicrobial wound infections in a tsunami survivor from Sri Lanka. *Lancet.* 2005;365:876–878.
- Tribble DR, Rodriguez CJ, Weintrob AC, et al. Environmental factors related to fungal wound contamination after combat trauma in Afghanistan, 2009–2011. *Emerg Infect Dis.* 2015;21:1759–1769.
- Ma L-J, Ibrahim AS, Skory C, et al. Genomic analysis of the basal lineage fungus *Rhizopus oryzae* reveals a whole-genome duplication. *PLoS Genet.* 2009;5:e1000549.
- Mendoza L, Vilela R, Voelz K, et al. Human fungal pathogens of Mucorales and Entomophthorales. *Cold Spring Harb Perspect Med.* 2014;5.
- Vitale RG, de Hoog GS, Schwarz P, et al. Antifungal susceptibility and phylogeny of opportunistic members of the order Mucorales. *J Clin Microbiol.* 2012;50:66–75.
- Lewis RE, Lortholary O, Spellberg B, et al. How does antifungal pharmacology differ for mucormycosis versus aspergillosis? *Clin Infect Dis.* 2012;54 Suppl 1(suppl 1):S67–S72.
- Chibucos MC, Soliman S, Gebremariam T, et al. An integrated genomic and transcriptomic survey of mucormycosis-causing fungi. *Nat Commun.* 2016;7:12218.
- Farmakiotis D, Kontoyiannis DP. Mucormycoses. *Infect Dis Clin North Am.* 2016;30:143–163.
- Gioverti MV, Cawcutt KA, Abidi M, et al. Gastrointestinal mucormycosis in immunocompromised hosts. *Mycoses.* 2015;58:714–718.
- Spellberg B. Gastrointestinal mucormycosis: an evolving disease. *Gastroenterol Hepatol (N Y).* 2012;8:140–142.
- Lee SC, Billmyre RB, Li A, et al. Analysis of a food-borne fungal pathogen outbreak: virulence and genome of a *Mucor circinelloides* isolate from yogurt. *MBio.* 2014;5:e01390–01314.
- Pana ZD, Seidel D, Skiada A, et al. Invasive mucormycosis in children: an epidemiologic study in European and non-European countries based on two registries. *BMC Infect Dis.* 2016;16:667.
- Davoudi S, Graviss LS, Kontoyiannis DP. Healthcare-associated outbreaks due to Mucorales and other uncommon fungi. *Eur J Clin Invest.* 2015;45:767–773.
- Cheng VCC, Chen JHK, Wong SCY, et al. Hospital outbreak of pulmonary and cutaneous zygomycosis due to contaminated linen items from substandard laundry. *Clin Infect Dis.* 2016;62:714–721.
- Gartenberg G, Bottoni EJ, Keusch GT, et al. Hospital-acquired mucormycosis (*Rhizopus rhizopodiformis*) of skin and subcutaneous tissue: epidemiology, mycology and treatment. *N Engl J Med.* 1978;299:1115–1118.
- Mead JH, Lupton GP, Dillavou CL, et al. Cutaneous *Rhizopus* infection. Occurrence as a postoperative complication associated with an elasticized adhesive dressing. *JAMA.* 1979;242:272–274.
- Chakrabarti A, Kumar P, Padhye AA, et al. Primary cutaneous zygomycosis due to *Saksenaia vasiformis* and *Apophysomyces elegans*. *Clin Infect Dis.* 1997;24:580–583.
- Jain JK, Markowitz A, Khilani PV, et al. Localized mucormycosis following intramuscular corticosteroid. Case report and review of the literature. *Am J Med Sci.* 1978;275:209–216.
- Verweij PE, Voss A, Donnelly JP, et al. Wooden sticks as the source of a pseudoepidemic of infection with *Rhizopus microsporus* var. *rhizopodiformis* among immunocompromised patients. *J Clin Microbiol.* 1997;35:2422–2423.
- Paydas S, Yavuz S, Disel U, et al. Mucormycosis of the tongue in a patient with acute lymphoblastic leukemia: a possible relation with use of a tongue depressor. *Am J Med.* 2003;114:618–620.
- Duffy J, Harris J, Gade L, et al. Mucormycosis outbreak associated with hospital linens. *Pediatr Infect Dis J.* 2014;33:472–476.
- Baker RD. Mucormycosis; a new disease? *J Am Med Assoc.* 1957;163:805–808.
- Baker RD. Pulmonary mucormycosis. *Am J Pathol.* 1956;32:287–313.
- Artis WM, Fountain JA, Delcher HK, et al. A mechanism of susceptibility to mucormycosis in diabetic ketoacidosis: transferrin and iron availability. *Diabetes.* 1982;31:1109–1114.
- Boelaert JR, de Locht M, Van Cutsem J, et al. Mucormycosis during deferoxamine therapy is a siderophore-mediated infection. In vitro and in vivo animal studies. *J Clin Invest.* 1993;91:1979–1986.
- Boelaert JR, van Roost GF, Vergauwe PL, et al. The role of desferrioxamine in dialysis-associated mucormycosis: report of three cases and review of the literature. *Clin Nephrol.* 1988;29:261–266.
- Boelaert JR, Vergauwe PL, Vandepitte JM. Mucormycosis infection in dialysis patients. *Ann Intern Med.* 1987;107:782–783.
- Sims CR, Ostrosky-Zeichner L. Contemporary treatment and outcomes of zygomycosis in a non-oncologic tertiary care center. *Arch Med Res.* 2007;38:90–93.
- Hospital DR, Chung KK, Laird K, et al. *Saksenaia erythrospora* infection following combat trauma. *J Clin Microbiol.* 2011;49:3707–3709.
- Fahimzad A, Chavoshzadeh Z, Abdollahpour H, et al. Necrosis of nasal cartilage due to mucormycosis in a patient with severe congenital neutropenia due to HAX1 deficiency. *J Investig Allergol Clin Immunol.* 2008;18:469–472.
- Kumar N, Hanks ME, Chandrasekaran P, et al. Gain-of-function signal transducer and activator of transcription 1 (STAT1) mutation-related primary immunodeficiency is associated with disseminated mucormycosis. *J Allergy Clin Immunol.* 2014;134:236–239.
- Kontoyiannis DP, Yang H, Song J, et al. Prevalence, clinical and economic burden of mucormycosis-related hospitalizations in the United States: a retrospective study. *BMC Infect Dis.* 2016;16:730.
- Petrikos G, Skiada A, Lortholary O, et al. Epidemiology and clinical manifestations of mucormycosis. *Clin Infect Dis.* 2012;54(suppl 1):S23–S34.
- Lanternier F, Dannaoui E, Morizot G, et al. A global analysis of mucormycosis in France: the RetroZygo Study (2005–2007). *Clin Infect Dis.* 2012;54 Suppl 1:S35–S43.
- Kontoyiannis DP, Marr KA, Park BJ, et al. Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001–2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis.* 2010;50:1091–1100.
- Pappas PG, Alexander BD, Andes DR, et al. Invasive fungal infections among organ transplant recipients: results of the Transplant-Associated Infection Surveillance Network (TRANSNET). *Clin Infect Dis.* 2010;50:1101–1111.
- Kontoyiannis DP, Azie N, Franks B, et al. Prospective antifungal therapy (PATH) alliance: focus on mucormycosis. *Mycoses.* 2014;57:240–246.
- Parker VJ, Jergens AE, Whitley EM, et al. Isolation of *Cokeromyces recurvatus* from the gastrointestinal tract in a dog with protein-losing enteropathy. *J Vet Diagn Invest.* 2011;23:1014–1016.
- Kontoyiannis DP, Marr KA, Park BJ, et al. Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001–2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis.* 2010;50:1091–1100.
- Marty FM, Cosimi LA, Baden LR. Breakthrough zygomycosis after voriconazole treatment in recipients of hematopoietic stem-cell transplants. *N Engl J Med.* 2004;350:950–952.
- Kontoyiannis DP, Lionakis MS, Lewis RE, et al. Zygomycosis in a tertiary-care cancer center in the era of *Aspergillus*-active antifungal therapy: a case-control observational study of 27 recent cases. *J Infect Dis.* 2005;191:1350–1360.
- Lamoth F, Chung SJ, Damonti L, et al. Changing epidemiology of invasive mold infections in patients receiving azole prophylaxis. *Clin Infect Dis.* 2017;64:1619–1621.
- England AC 3rd, Weinstein M, Ellner JJ, et al. Two cases of rhinocerebral zygomycosis (mucormycosis) with common epidemiologic and environmental features. *Am Rev Respir Dis.* 1981;124:497–498.
- Lueg EA, Ballagh RH, Forte V. Analysis of the recent cluster of invasive fungal sinusitis at the Toronto Hospital for Sick Children. *J Otolaryngol.* 1996;25:366–370.
- Prevoo RL, Starink TM, de Haan P. Primary cutaneous mucormycosis in a healthy young girl. Report of a case caused by *Mucor hiemalis* Wehmer. *J Am Acad Dermatol.* 1991;24(5 Pt 2):882–885.
- Ibrahim AS, Voelz K. The mucormycete-host interface. *Curr Opin Microbiol.* 2017;40:40–45.
- Bouchard JR, Oumeziane NA, Lissitzky JC, et al. Attachment of spores of the human pathogenic fungus *Rhizopus oryzae* to extracellular matrix components. *Eur J Cell Biol.* 1996;70:76–83.
- Ibrahim AS, Spellberg B, Avanesian V, et al. *Rhizopus oryzae* adheres to, is phagocytosed by, and damages endothelial cells in vitro. *Infect Immun.* 2005;73:778–783.
- Liu M, Spellberg B, Phan QT, et al. The endothelial cell receptor GRP78 is required for mucormycosis pathogenesis in diabetic mice. *J Clin Invest.* 2010;120:1914–1924.
- Gebremariam T, Lin L, Liu M, et al. Bicarbonate correction of ketoacidosis alters host-pathogen interactions and alleviates mucormycosis. *J Clin Invest.* 2016;126:2280–2294.
- Gebremariam T, Liu M, Luo G, et al. CoH3 mediates fungal invasion of host cells during mucormycosis. *J Clin Invest.* 2014;124:237–250.
- Baldin C, Ibrahim AS. Molecular mechanisms of mucormycosis—the bitter and the sweet. *PLoS Pathog.* 2017;13:e1006408.
- de Locht M, Boelaert JR, Schneider YJ. Iron uptake from ferrioxamine and from ferrirrhizoferrin by germinating spores of *Rhizopus microsporus*. *Biochem Pharmacol.* 1994;47:1843–1850.
- Ibrahim AS, Gebremariam T, Lin L, et al. The high affinity iron permease is a key virulence factor required for *Rhizopus oryzae* pathogenesis. *Mol Microbiol.* 2010;77:587–604.
- Alexander J, Limaye AP, Ko CW, et al. Association of hepatic iron overload with invasive fungal infection in liver transplant recipients. *Liver Transpl.* 2006;12:1799–1804.
- Artis WM, Patrusky E, Rastinejad F, et al. Fungistatic mechanism of human transferrin for *Rhizopus oryzae* and *Trichophyton mentagrophytes*: alternative to simple iron deprivation. *Infect Immun.* 1983;41:1269–1278.
- Boelaert JR, Van Cutsem J, de Locht M, et al. Deferoxamine augments growth and pathogenicity of *Rhizopus*, while hydroxyypyridinone chelators have no effect. *Kidney Int.* 1994;45:667–671.
- Boelaert JR, de Locht M, Schneider YJ. The effect of deferoxamine on different zygomycetes. *J Infect Dis.* 1994;169:231–232.
- Ibrahim AS, Gebremariam T, Fu Y, et al. The iron chelator deferasirox protects mice from mucormycosis through iron starvation. *J Clin Invest.* 2007;117:2649–2657.
- Reed C, Ibrahim A, Edwards JE Jr, et al. Deferasirox, an iron-chelating agent, as salvage therapy for rhinocerebral mucormycosis. *Antimicrob Agents Chemother.* 2006;50:3968–3969.
- Spellberg B, Ibrahim AS, Chin-Hong PV, et al. The Deferasirox-AmBisome Therapy for Mucormycosis (DEFEAT Mucor) study: a randomized, double-blinded, placebo-controlled trial. *J Antimicrob Chemother.* 2012;67:715–722.
- Cornely OA, Arikian-Akdag S, Dannaoui E, et al. ESCMID and ECMM joint clinical guidelines for the diagnosis and management of mucormycosis 2013. *Clin Microbiol Infect.* 2014;20 Suppl 3:5–26.
- Waldorf AR, Levitz SM, Diamond RD. In vivo bronchoalveolar macrophage defense against *Rhizopus oryzae* and *Aspergillus fumigatus*. *J Infect Dis.* 1984;150:752–760.
- Waldorf AR, Ruderman N, Diamond RD. Specific susceptibility to mucormycosis in murine diabetes and bronchoalveolar macrophage defense against *Rhizopus*. *J Clin Invest.* 1984;74:150–160.
- Schmidt S, Tramsen L, Perkhof S, et al. *Rhizopus oryzae* hyphae are damaged by human natural killer (NK) cells, but suppress NK cell mediated immunity. *Immunobiology.* 2013;218:939–944.
- Chamilos G, Lewis RE, Lamas G, et al. Zygomycetes hyphae trigger an early, robust proinflammatory response in human polymorphonuclear neutrophils through

- Toll-like receptor 2 induction but display relative resistance to oxidative damage. *Antimicrob Agents Chemother*. 2008;52:722–724.
76. Diamond RD, Haudenschild CC, Erickson NF 3rd. Monocyte-mediated damage to *Rhizopus oryzae* hyphae in vitro. *Infect Immun*. 1982;38:292–297.
77. Perkhof S, Kainzner B, Kehrel BE, et al. Potential antifungal effects of human platelets against *Zygomycetes* in vitro. *J Infect Dis*. 2009;200:1176–1179.
78. Lionakis MS, Kontoyiannis DP. Glucocorticoids and invasive fungal infections. *Lancet*. 2003;362:1828–1838.
79. Chinn RY, Diamond RD. Generation of chemotactic factors by *Rhizopus oryzae* in the presence and absence of serum: relationship to hyphal damage mediated by human neutrophils and effects of hyperglycemia and ketoacidosis. *Infect Immun*. 1982;38:1123–1129.
80. Potenza L, Vallerini D, Barozzi P, et al. Mucorales-specific T cells emerge in the course of invasive mucormycosis and may be used as a surrogate diagnostic marker in high-risk patients. *Blood*. 2011;118:5416–5419.
81. Chamilos G, Ganguly D, Lande R, et al. Generation of IL-23 producing dendritic cells (DCs) by airborne fungi regulates fungal pathogenicity via the induction of T(H)-17 responses. *PLoS ONE*. 2010;5:e12955.
82. Guo D, Jaber BL, Lee S, et al. Impact of iron dextran on polymorphonuclear cell function among hemodialysis patients. *Clin Nephrol*. 2002;58:134–142.
83. Omara FO, Blakley BR. The effects of iron deficiency and iron overload on cell-mediated immunity in the mouse. *Br J Nutr*. 1994;72:899–909.
84. Omara FO, Blakley BR, Huang HS. Effect of iron status on endotoxin-induced mortality, phagocytosis and interleukin-1 alpha and tumor necrosis factor-alpha production. *Vet Hum Toxicol*. 1994;36:423–428.
85. Cantiniaux B, Janssens A, Boelaert JR, et al. Ferritin-associated iron induces neutrophil dysfunction in hemodieresis. *J Lab Clin Med*. 1999;133:353–361.
86. Sheldon WH, Bauer H. Activation of quiescent mucormycotic granuloma in rabbits by induction of acute alloxan diabetes. *J Exp Med*. 1958;108:171–178.
87. Chandler FW, Kaplan W, Ajello L A colour atlas and textbook of the histopathology of mycotic diseases; 1980.
88. Ben-Ami R, Luna M, Lewis RE, et al. A clinicopathological study of pulmonary mucormycosis in cancer patients: extensive angioinvasion but limited inflammatory response. *J Infect*. 2009;59:134–138.
89. Stergiopoulou T, Meletiadis J, Roilides E, et al. Host-dependent patterns of tissue injury in invasive pulmonary aspergillosis. *Am J Clin Pathol*. 2007;127:349–355.
90. Mallis A, Mastronikolis SN, Naxakis SS, Papadas AT. Rhinocerebral mucormycosis: an update. *Eur Rev Med Pharmacol Sci*. 2010;14:987–992.
91. Yohai RA, Bullock JD, Aziz AA, et al. Survival factors in rhino-orbital-cerebral mucormycosis. *Surv Ophthalmol*. 1994;39:3–22.
92. Ochiai H, Iseda T, Miyahara S, et al. Rhinocerebral mucormycosis—case report. *Neurol Med Chir (Tokyo)*. 1993;33:373–376.
93. Frater JL, Hall GS, Procop GW. Histologic features of zygomycosis: emphasis on perineural invasion and fungal morphology. *Arch Pathol Lab Med*. 2001;125:375–378.
94. Ferguson BJ. Is your case report really an invasive fungal rhinosinusitis? *Laryngoscope*. 2005;115:560, author reply 560–562.
95. Demirag A, Elkhannas EA, Henry ML, et al. Pulmonary *Rhizopus* infection in a diabetic renal transplant recipient. *Clin Transplant*. 2000;14:8–10.
96. Tsai TC, Hou CC, Chou MS, et al. Rhinosino-orbital mucormycosis causing cavernous sinus thrombosis and internal carotid artery occlusion: radiological findings in a patient with treatment failure. *Kaohsiung J Med Sci*. 1999;15:556–561.
97. Sehgal A, Raghavendran M, Kumar D, et al. Rhinocerebral mucormycosis causing basilar artery aneurysm with concomitant fungal colonic perforation in renal allograft recipient: a case report. *Transplantation*. 2004;78:949–950.
98. Delbrouck C, Jacobs F, Fernandez Aguilar S, et al. Carotid artery occlusion due to fulminant rhinocerebral mucormycosis. *Acta Otorhinolaryngol Belg*. 2004;58:135–140.
99. Howells RC, Ramadan HH. Usefulness of computed tomography and magnetic resonance in fulminant invasive fungal rhinosinusitis. *Am J Rhinol*. 2001;15:255–261.
100. Fatterpekar G, Mukherji S, Arbeale A, et al. Fungal diseases of the paranasal sinuses. *Semin Ultrasound CT MR*. 1999;20:391–401.
101. Greenberg MR, Lippman SM, Grinnell VS, et al. Computed tomographic findings in orbital *Mucor*. *West J Med*. 1985;143:102–103.
102. Greenberg RN, Scott LJ, Vaughn HH, et al. Zygomycosis (mucormycosis): emerging clinical importance and new treatments. *Curr Opin Infect Dis*. 2004;17:517–525.
103. Chamilos G, Marom EM, Lewis RE, et al. Predictors of pulmonary zygomycosis versus invasive pulmonary aspergillosis in patients with cancer. *Clin Infect Dis*. 2005;41:60–66.
104. Lass-Flörl C, Aigner M, Nachbaur D, et al. Diagnosing filamentous fungal infections in immunocompromised patients applying computed tomography-guided percutaneous lung biopsies: a 12-year experience. *Infection*. 2017;45:867–875.
105. Lass-Flörl C, Resch G, Nachbaur D, et al. The value of computed tomography-guided percutaneous lung biopsy for diagnosis of invasive fungal infection in immunocompromised patients. *Clin Infect Dis*. 2007;45:e101–e104.
106. Chamilos G, Lewis RE, Kontoyiannis DP. Delaying Amphotericin B-based frontline therapy significantly increases mortality among patients with hematologic malignancy who have zygomycosis. *Clin Infect Dis*. 2008;47:503–509.
107. Caillot D, Valot S, Lafon I, et al. Is it time to include CT “reverse halo sign” and qPCR targeting Mucorales in serum to EORTC-MSG criteria for the diagnosis of pulmonary mucormycosis in leukemia patients? *Open Forum Infect Dis*. 2016;3:ofw190.
108. Legouge C, Caillot D, Chretien ML, et al. The reversed halo sign: pathognomonic pattern of pulmonary mucormycosis in leukemic patients with neutropenia? *Clin Infect Dis*. 2014;58:672–678.
109. Gupta N, Kumar R, Prasad R. Reversed halo sign. *Indian J Chest Dis Allied Sci*. 2014;56:247–248.
- 109a. Thomas R, Madan R, Gooptu M, et al. Significance of the reverse halo sign in immunocompromised patients. *AJR Am J Roentgenol*. 2019;30:1–6.
110. Lee FY, Mossad SB, Adal KA. Pulmonary mucormycosis: the last 30 years. *Arch Intern Med*. 1999;159:1301–1309.
111. Gupta KL, Khullar DK, Behera D, et al. Pulmonary mucormycosis presenting as fatal massive haemoptysis in a renal transplant recipient. *Nephrol Dial Transplant*. 1998;13:3258–3260.
112. Funada H, Matsuda T. Pulmonary mucormycosis in a hematology ward. *Intern Med*. 1996;35:540–544.
113. Kontoyiannis DP, Lewis RE. Treatment principles for the management of mold infections. *Cold Spring Harb Perspect Med*. 2015;5:a019737.
114. Lanternier F, Sun H-Y, Ribaud P, et al. Mucormycosis in organ and stem cell transplant recipients. *Clin Infect Dis*. 2012;54:1629–1636.
115. Trifilio SM, Bennett CL, Yarnold PR, et al. Breakthrough zygomycosis after voriconazole administration among patients with hematologic malignancies who receive hematopoietic stem-cell transplants or intensive chemotherapy. *Bone Marrow Transplant*. 2007;39:425–429.
116. Nam BD, Kim TJ, Lee KS, et al. Pulmonary mucormycosis: serial morphologic changes on computed tomography correlate with clinical and pathologic findings. *Eur Radiol*. 2018;28:788–795.
117. McAdams HP, Rosado de Christenson M, Strollo DC, et al. Pulmonary mucormycosis: radiologic findings in 32 cases. *AJR Am J Roentgenol*. 1997;168:1541–1548.
118. Tedder M, Spratt JA, Anstadt MP, et al. Pulmonary mucormycosis: results of medical and surgical therapy. *Ann Thorac Surg*. 1994;57:1044–1050.
119. Wahba H, Truong MT, Lei X, et al. Reversed halo sign in invasive pulmonary fungal infections. *Clin Infect Dis*. 2008;46:1733–1737.
120. Georgiadou SP, Sipsas NV, Marom EM, et al. The diagnostic value of halo and reversed halo signs for invasive mold infections in compromised hosts. *Clin Infect Dis*. 2011;52:1144–1155.
121. Kontoyiannis DP, Wessel VC, Bodey GP, et al. Zygomycosis in the 1990s in a tertiary-care cancer center. *Clin Infect Dis*. 2000;30:851–856.
122. O’Connell MA, Pluss JL, Schkade P, et al. *Rhizopus*-induced hypersensitivity pneumonitis in a tractor driver. *J Allergy Clin Immunol*. 1995;95:779–780.
123. Hedenstierna G, Alexandersson R, Belin L, et al. Lung function and *Rhizopus* antibodies in wood trimmers. A cross-sectional and longitudinal study. *Int Arch Occup Environ Health*. 1986;58:167–177.
124. Bearer EA, Nelson PR, Chowes MY, et al. Cutaneous zygomycosis caused by *Saksenaea vasiformis* in a diabetic patient. *J Clin Microbiol*. 1994;32:1823–1824.
125. Rodriguez-Lobato E, Ramirez-Hobak L, Aquino-Matus JE, et al. Primary cutaneous mucormycosis caused by *Rhizopus oryzae*: a case report and review of literature. *Mycopathologia*. 2017;182:387–392.
126. De Decker K, Van Poucke S, Wojciechowski M, et al. Successful use of posaconazole in a pediatric case of fungal necrotizing fasciitis. *Pediatr Crit Care Med*. 2006;7:482–485.
127. Devi SC, Kanungo R, Barreto E, et al. Favorable outcome of amphotericin B treatment of zygomycotic necrotizing fasciitis caused by *Apophysomyces elegans*. *Int J Dermatol*. 2008;47:407–409.
128. Jain D, Kumar Y, Vasishta RK, et al. Zygomycotic necrotizing fasciitis in immunocompetent patients: a series of 18 cases. *Mod Pathol*. 2006;19:1221–1226.
129. Kordy FN, Al-Mohsen IZ, Hashem F, et al. Successful treatment of a child with posttraumatic necrotizing fasciitis caused by *Apophysomyces elegans*: case report and review of literature. *Pediatr Infect Dis J*. 2004;23:877–879.
130. Lakshmi V, Rani TS, Sharma S, et al. Zygomycotic necrotizing fasciitis caused by *Apophysomyces elegans*. *J Clin Microbiol*. 1993;31:1368–1369.
131. Thami GP, Kaur S, Bawa AS, et al. Post-surgical zygomycotic necrotizing subcutaneous infection caused by *Absidia corymbifera*. *Clin Exp Dermatol*. 2003;28:251–253.
132. Nouri-Majalan N, Moghimi M. Skin mucormycosis presenting as an erythema-nodosum-like rash in a renal transplant recipient: a case report. *J Med Case Rep*. 2008;2:112.
133. Vernon SE, Dave SP. Cutaneous zygomycosis associated with urate panniculitis. *Am J Dermatopathol*. 2006;28:327–330.
134. Farmakiotis D, Ciurea AM, Cahuayme-Zuniga L, et al. The diagnostic yield of skin biopsy in patients with leukemia and suspected infection. *J Infect*. 2013;67:265–272.
135. Taj-Aldeen SJ, Gamaletsou MN, Rammaert B, et al. Bone and joint infections caused by Mucormycetes: a challenging osteoarticular mycosis of the twenty-first century. *Med Mycol*. 2017;55:691–704.
136. Bittencourt AL, Ayala MA, Ramos EA. A new form of abdominal zygomycosis different from mucormycosis: report of two cases and review of the literature. *Am J Trop Med Hyg*. 1979;28:564–569.
137. Park YS, Lee JD, Kim TH, et al. Gastric mucormycosis. *Gastrointest Endosc*. 2002;56:904–905.
138. Oliver MR, Van Voorhis WC, Boeckh M, et al. Hepatic mucormycosis in a bone marrow transplant recipient who ingested naturopathic medicine. *Clin Infect Dis*. 1996;22:521–524.
139. Yoshita S, Kitano K, Nagaya T, et al. Zygomycosis presenting as acute myocardial infarction during hematological malignancies. *Intern Med*. 2008;47:839–842.
140. Naumann R, Kerkmann ML, Schuler U, et al. *Cunninghamella bertholletiae* infection mimicking myocardial infarction. *Clin Infect Dis*. 1999;29:1580–1581.
141. Benbow EW, Bonshek RE, Stoddart RW. Endobronchial zygomycosis. *Thorax*. 1987;42:553–554.
142. Polo JR, Luno J, Menarguez C, et al. Peritoneal mucormycosis in a patient receiving continuous ambulatory peritoneal dialysis. *Am J Kidney Dis*. 1989;13:237–239.
143. Nakamura M, Weil WB Jr, Kaufman DB. Fatal fungal peritonitis in an adolescent on continuous ambulatory peritoneal dialysis: association with deferoxamine. *Pediatr Nephrol*. 1989;3:80–82.
144. Branton MH, Johnson SC, Brooke JD, et al. Peritonitis due to *Rhizopus* in a patient undergoing continuous ambulatory peritoneal dialysis. *Rev Infect Dis*. 1991;13:19–21.
145. Schwartz JR, Nagle MG, Elkins RC, et al. Mucormycosis of the trachea: an unusual cause of acute upper airway obstruction. *Chest*. 1982;81:653–654.
146. Connor BA, Anderson RJ, Smith JW. Mucor mediastinitis. *Chest*. 1979;75:525–526.
147. Echols RM, Selinger DS, Halliwell C, et al. *Rhizopus* osteomyelitis. A case report and review. *Am J Med*. 1979;66:141–145.
148. Mishra B, Mandal A, Kumar N. Mycotic prosthetic-valve endocarditis. *J Hosp Infect*. 1992;20:122–125.
149. Roy TM, Anderson KC, Farrow JR. Cardiac mucormycosis complicating diabetes mellitus. *J Diabet Complications*. 1990;4:132–135.
150. Lussier N, Laverdiere M, Weiss K, et al. Primary renal mucormycosis. *Urology*. 1998;52:900–903.
151. Tierney MR, Baker AS. Infections of the head and neck in diabetes mellitus. *Infect Dis Clin North Am*. 1995;9:195–216.
152. Bosken CH, Szporn AH, Kleinerman J. Superior vena cava syndrome due to mucormycosis in a patient with lymphoma. *Mt Sinai J Med*. 1987;54:508–511.
153. Air EL, Vagal AA, Kendler A, et al. Isolated cerebellar mucormycosis, slowly progressive over 1 year in an immunocompetent patient. *Surg Neurol Int*. 2010;1:81.
154. Verma A, Brozman B, Pettit CK. Isolated cerebral mucormycosis: report of a case and review of the literature. *J Neurol Sci*. 2006;240:65–69.
155. Cuadrado LM. Cerebral mucormycosis associated with intravenous drug use. *Clin Infect Dis*. 1996;22:198.

156. Hopkins RJ, Rothman M, Fiore A, et al. Cerebral mucormycosis associated with intravenous drug use: three case reports and review. *Clin Infect Dis*. 1994;19:1133–1137.
157. Leen CL, Brettell RP. Fungal infections in drug users. *J Antimicrob Chemother*. 1991;28 Suppl A:83–96.
158. Glazer M, Nusair S, Breuer R, et al. The role of BAL in the diagnosis of pulmonary mucormycosis. *Chest*. 2000;117:279–282.
159. Chayakulkeeree M, Ghannoum MA, Perfect JR. Zygomycosis: the re-emerging fungal infection. *Eur J Clin Microbiol Infect Dis*. 2006;25:215–229.
160. Ruchel R, Schaffrinski M, Seshan KR, et al. Vital staining of fungal elements in deep-seated mycotic lesions during experimental murine mycoses using the parentally applied optical brightener Blankophor. *Med Mycol*. 2000;38:231–237.
161. Hammond SP, Bialek R, Milner DA, et al. Molecular methods to improve diagnosis and identification of mucormycosis. *J Clin Microbiol*. 2011;49:2151–2153.
162. Normand AC, Cassagne C, Gautier M, et al. Decision criteria for MALDI-TOF MS-based identification of filamentous fungi using commercial and in-house reference databases. *BMC Microbiol*. 2017;17:25.
163. Cassagne C, Normand AC, L'Olivier C, et al. Performance of MALDI-TOF MS platforms for fungal identification. *Mycoses*. 2016;59:678–690.
164. Kontoyiannis DP, Chamliou G, Hassan SA, et al. Increased culture recovery of Zygomycetes under physiologic temperature conditions. *Am J Clin Pathol*. 2007;127:208–212.
165. Tarrand JJ, Lichterfeld M, Warraich I, et al. Diagnosis of invasive septate mold infections. A correlation of microbiological culture and histologic or cytologic examination. *Am J Clin Pathol*. 2003;119:854–858.
166. Hammond SP, Baden LR, Marty FM. Mortality in hematologic malignancy and hematopoietic stem cell transplant patients with mucormycosis, 2001 to 2009. *Antimicrob Agents Chemother*. 2011;55:5018–5021.
167. Springer J, Goldenberger D, Schmidt F, et al. Development and application of two independent real-time PCR assays to detect clinically relevant Mucorales species. *J Med Microbiol*. 2016;65:227–234.
168. Springer J, Lackner M, Ensinger C, et al. Clinical evaluation of a Mucorales-specific real-time PCR assay in tissue and serum samples. *J Med Microbiol*. 2016;65:1414–1421.
169. Kobayashi M, Togitani K, Machida H, et al. Molecular polymerase chain reaction diagnosis of pulmonary mucormycosis caused by *Cunninghamella bertholletiae*. *Respirology*. 2004;9:397–401.
170. Bialek R, Konrad F, Kern J, et al. PCR based identification and discrimination of agents of mucormycosis and aspergillosis in paraffin wax embedded tissue. *J Clin Pathol*. 2005;58:1180–1184.
171. Rickerts V, Atta J, Herrmann S, et al. Successful treatment of disseminated mucormycosis with a combination of liposomal amphotericin B and posaconazole in a patient with acute myeloid leukaemia. *Mycoses*. 2006;49 Suppl 1:27–30.
172. Arendrup MC, Jensen RH, Meletiadis J. In vitro activity of isavuconazole and comparators against clinical isolates of the Mucorales order. *Antimicrob Agents Chemother*. 2015;59:7735–7742.
173. Chowdhary A, Singh PK, Kathuria S, et al. Comparison of the EUCAST and CLSI broth microdilution methods for testing isavuconazole, posaconazole, and amphotericin B against molecularly identified Mucorales species. *Antimicrob Agents Chemother*. 2015;59:7882–7887.
174. Dannaoui E. Antifungal resistance in Mucorales. *Int J Antimicrob Agents*. 2017;50:617–621.
175. Espinel-Ingroff A, Bartlett M, Chaturvedi V, et al. Optimal susceptibility testing conditions for detection of azole resistance in *Aspergillus* spp.: NCCLS collaborative evaluation. National Committee for Clinical Laboratory Standards. *Antimicrob Agents Chemother*. 2001;45:1828–1835.
176. Espinel-Ingroff A, Alvarez-Fernandez M, Cantón E, et al. Multicenter study of epidemiological cutoff values and detection of resistance in *Candida* spp. to anidulafungin, caspofungin, and micafungin using the Sensititre YeastOne colorimetric method. *Antimicrob Agents Chemother*. 2015;59:6725–6732.
177. Lamoth F, Damonti L, Alexander BD. Role of antifungal susceptibility testing in non-*Aspergillus* invasive mold infections. *J Clin Microbiol*. 2016;54:1638–1640.
178. Dannaoui E, Rijs AJ, Verweij PE. Invasive infections due to *Apophysomyces elegans*. *Mayo Clin Proc*. 2003;78:252–253.
179. Lewis RE, Albert ND, Kontoyiannis DP. Comparative pharmacodynamics of posaconazole in neutropenic murine models of invasive pulmonary aspergillosis and mucormycosis. *Antimicrob Agents Chemother*. 2014;58:6767–6772.
180. Ibrahim AS, Bowman JC, Avanesian V, et al. Caspofungin inhibits *Rhizopus oryzae* 1,3-beta-D-glucan synthase, lowers burden in brain measured by quantitative PCR, and improves survival at a low but not a high dose during murine disseminated zygomycosis. *Antimicrob Agents Chemother*. 2005;49:721–727.
181. Spellberg B, Fu Y, Edwards JE Jr, et al. Combination therapy with amphotericin B lipid complex and caspofungin acetate of disseminated zygomycosis in diabetic ketoacidotic mice. *Antimicrob Agents Chemother*. 2005;49:830–832.
182. Lewis RE, Leventakos K, Liao G, et al. Efficacy of caspofungin in neutropenic and corticosteroid-immunosuppressed murine models of invasive pulmonary mucormycosis. *Antimicrob Agents Chemother*. 2011;55:3584–3587.
183. Spellberg B, Ibrahim A, Roilides E, et al. Combination therapy for mucormycosis: why, what, and how? *Clin Infect Dis*. 2012;54 Suppl 1:S73–S78.
184. Ibrahim AS, Gebremariam T, Fu Y, et al. Combination echinocandin-polyene treatment of murine mucormycosis. *Antimicrob Agents Chemother*. 2008;52:1556–1558.
185. Gebremariam T, Alkhazraji S, Baldin C, et al. Prophylaxis with isavuconazole or posaconazole protect immunosuppressed mice from pulmonary mucormycosis. *Antimicrob Agents Chemother*. 2017;61:e02589-16.
186. Gebremariam T, Wiederhold NP, Alqarihi A, et al. Monotherapy or combination therapy of isavuconazole and micafungin for treating murine mucormycosis. *J Antimicrob Chemother*. 2017;72:462–466.
187. Kontoyiannis DP, Lewis RE. How I treat mucormycosis. *Blood*. 2011;118:1216–1224.
188. Ghadiali MT, Deckard NA, Farooq U, et al. Frozen-section biopsy analysis for acute invasive fungal rhinosinusitis. *Otolaryngol Head Neck Surg*. 2007;136:714–719.
189. Gleissner B, Schilling A, Anagnostopoulou I, et al. Improved outcome of zygomycosis in patients with hematological diseases? *Leuk Lymphoma*. 2004;45:1351–1360.
190. Perfect JR. Treatment of non-*Aspergillus* moulds in immunocompromised patients, with amphotericin B lipid complex. *Clin Infect Dis*. 2005;40 Suppl 6:S401–S408.
191. Walsh TJ, Hiemenz JW, Seibel NL, et al. Amphotericin B lipid complex for invasive fungal infections: analysis of safety and efficacy in 556 cases. *Clin Infect Dis*. 1998;26:1383–1396.
192. Garbino J, Adam A. Use of high-dose liposomal amphotericin B: efficacy and tolerance. *Acta Biomed*. 2006;77 Suppl 4:19–22.
193. Revankar SG, Hasan MS, Smith JW. Cure of disseminated zygomycosis with cerebral involvement using high dose liposomal amphotericin B and surgery. *Med Mycol*. 2007;45:183–185.
194. Lewis RE, Albert ND, Liao G, et al. Comparative pharmacodynamics of amphotericin B lipid complex and liposomal amphotericin B in a murine model of pulmonary mucormycosis. *Antimicrob Agents Chemother*. 2010;54:1298–1304.
195. Stanzani M, Vianelli N, Cavo M, et al. Liposomal amphotericin B nephrotoxicity in patients with hematological malignancies: a retrospective cohort analysis. *Antimicrob Agents Chemother*. 2017;61:e02651-16.
196. Prentice HG, Hann IM, Herbrecht R, et al. A randomized comparison of liposomal versus conventional amphotericin B for the treatment of pyrexia of unknown origin in neutropenic patients. *Br J Haematol*. 1997.
197. Safdar A, Ma J, Saliba F, et al. Drug-induced nephrotoxicity caused by amphotericin B lipid complex and liposomal amphotericin B: a review and meta-analysis. *Medicine (Baltimore)*. 2010;89:236–244.
198. Cornely OA, Maertens J, Bresnik M, et al. Liposomal amphotericin B as initial therapy for invasive mold infection: a randomized trial comparing a high-loading dose regimen with standard dosing (AmBiload trial). *Clin Infect Dis*. 2007;44:1289–1297.
199. Lanternier F, Poiree S, Elie C, et al. Prospective pilot study of high-dose (10 mg/kg/day) liposomal amphotericin B (L-AMB) for the initial treatment of mucormycosis. *J Antimicrob Chemother*. 2015;70:3116–3123.
200. Dekkers BGJ, Bakker M, van der Elst KCM, et al. Therapeutic drug monitoring of posaconazole: an update. *Curr Fungal Infect Rep*. 2016;10:51–61.
201. Greenberg RN, Mullane K, van Burik JA, et al. Posaconazole as salvage therapy for zygomycosis. *Antimicrob Agents Chemother*. 2006;50:126–133.
202. Raad II, Graybill JR, Bustamante AB, et al. Safety of long-term oral posaconazole use in the treatment of refractory invasive fungal infections. *Clin Infect Dis*. 2006;42:1726–1734.
203. van Burik J-AH, Hare RS, Solomon HF, et al. Posaconazole is effective as salvage therapy in zygomycosis: a retrospective summary of 91 cases. *Clin Infect Dis*. 2006;42:e61–e65.
204. Kraft WK, Chang PS, van Iersel MLPS, et al. Posaconazole tablet pharmacokinetics: lack of effect of concomitant medications altering gastric pH and gastric motility in healthy subjects. *Antimicrob Agents Chemother*. 2014;58:4020–4025.
205. Cornely OA, Haider S, Grigg A, et al. Phase 3 Pharmacokinetic (PK) and Safety Study of Posaconazole (POS) IV in Patients (Pts) at Risk for Invasive Fungal Infection (IFI). 53rd Interscience Conference on Antimicrobial Agents and Chemotherapy; 2013. <https://www.aspergillus.org.uk/content/phase-3-pharmacokinetics-pk-and-safety-study-posaconazole-pos-iv-patients-pts-risk-invasive>.
206. Patterson TF, Thompson GR 3rd, Denning DW, et al. Practice guidelines for the diagnosis and management of aspergillosis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2016;63:e1–e60.
207. Walsh TJ, Raad I, Patterson TF, et al. Treatment of invasive aspergillosis with posaconazole in patients who are refractory to or intolerant of conventional therapy: an externally controlled trial. *Clin Infect Dis*. 2007;44:2–12.
208. Marty FM, Ostrosky-Zeichner L, Cornely OA, et al. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect Dis*. 2016;16:828–837.
209. Spellberg B, Brass E. The VITAL study: case control studies are hypothesis-generating. *Lancet Infect Dis*. 2016;16:886.
210. Reed C, Bryant R, Ibrahim AS, et al. Combination polyene-caspofungin treatment of rhino-orbital-cerebral mucormycosis. *Clin Infect Dis*. 2008;47:364–371.
211. Pagano L, Cornely OA, Busca A, et al. Combined antifungal approach for the treatment of invasive mucormycosis in patients with hematologic diseases: a report from the SEIFEM and FUNGISCOPE registries. *Haematologica*. 2013;98:e127–e130.
212. Kyvernitakis A, Torres HA, Jiang Y, et al. Initial use of combination treatment does not impact survival of 106 patients with haematologic malignancies and mucormycosis: a propensity score analysis. *Clin Microbiol Infect*. 2016;22:811.e811–811.e818.
213. Davoudi S, Anderlini P, Fuller GN, et al. A long-term survivor of disseminated *Aspergillus* and Mucorales infection: an instructive case. *Mycopathologia*. 2014.
214. Hamilton JF, Bartkowski HB, Rock JP. Management of CNS mucormycosis in the pediatric patient. *Pediatr Neurol*. 2003;38:212–215.
215. Raj P, Vella EJ, Bickerton RC. Successful treatment of rhinocerebral mucormycosis by a combination of aggressive surgical debridement and the use of systemic liposomal amphotericin B and local therapy with nebulized amphotericin—a case report. *J Laryngol Otol*. 1998;112:367–370.
216. Bitterman H. Hyperbaric oxygen for invasive fungal infections. *Isr Med Assoc J*. 2007;9:387–388.
217. Ferguson BJ, Mitchell TG, Moon R, et al. Adjunctive hyperbaric oxygen for treatment of rhinocerebral mucormycosis. *Rev Infect Dis*. 1988;10:551–559.
218. John BV, Chamliou G, Kontoyiannis DP. Hyperbaric oxygen as an adjunctive treatment for zygomycosis. *Clin Microbiol Infect*. 2005;11:515–517.
219. Gil-Lamagnere C, Simitsopoulou M, Roilides E, et al. Interferon-gamma and granulocyte-macrophage colony-stimulating factor augment the activity of polymorphonuclear leukocytes against medically important Zygomycetes. *J Infect Dis*. 2005;191:1180–1187.
220. Barratt DM, Van Meter K, Asmar P, et al. Hyperbaric oxygen as an adjunct in zygomycosis: randomized controlled trial in a murine model. *Antimicrob Agents Chemother*. 2001;45:3601–3602.
221. Abzug MJ, Walsh TJ. Interferon-gamma and colony-stimulating factors as adjuvant therapy for refractory fungal infections in children. *Pediatr Infect Dis J*. 2004;23:769–773.
222. Hubel K, Dale DC, Liles WC. Therapeutic use of cytokines to modulate phagocyte function for the treatment of infectious diseases: current status of granulocyte colony-stimulating factor, granulocyte-macrophage colony-stimulating factor, macrophage colony-stimulating factor, and interferon-gamma. *J Infect Dis*. 2002;185:1490–1501.
223. Hubel K, Dale DC, Engert A, et al. Current status of granulocyte (neutrophil) transfusion therapy for infectious diseases. *J Infect Dis*. 2001;183:321–328.
224. Lewis RE, Georgiadou SP, Sampsonas F, et al. Risk factors for early mortality in haematological malignancy patients with pulmonary mucormycosis. *Mycoses*. 2014;57:49–55.
225. El-Shabrawi MH, Arnaout H, Madkour I, et al. Entomophthoromycosis: a challenging emerging disease. *Mycoses*. 2014;57 Suppl 3:132–137.

226. Gugnani HC. Entomophthoromycosis due to *Conidiobolus*. *Eur J Epidemiol*. 1992;8:391–396.
227. Gugnani HC. A review of zygomycosis due to *Basidiobolus ranarum*. *Eur J Epidemiol*. 1999;15:923–929.
228. Walker SD, Clark RV, King CT, et al. Fatal disseminated *Conidiobolus coronatus* infection in a renal transplant patient. *Am J Clin Pathol*. 1992;98:559–564.
229. Bigliazzi C, Poletti V, Dell'Amore D, et al. Disseminated basidiobolomycosis in an immunocompetent woman. *J Clin Microbiol*. 2004;42:1367–1369.
230. Mugerwa JW. Subcutaneous phycomycosis in Uganda. *Br J Dermatol*. 1976;94:539–544.
231. Martinson FD. Clinical, epidemiological and therapeutic aspects of entomophthoromycosis. *Ann Soc Belg Med Trop*. 1972;52:329–342.
232. Khan ZU, Khoursheed M, Makar R, et al. *Basidiobolus ranarum* as an etiologic agent of gastrointestinal zygomycosis. *J Clin Microbiol*. 2001;39:2360–2363.
233. El-Shabrawi MH, Kamal NM, Kaerger K, et al. Diagnosis of gastrointestinal basidiobolomycosis: a mini-review. *Mycoses*. 2014;57 Suppl 3:138–143.
234. Khan ZU, Prakash B, Kapoor MM, et al. Basidiobolomycosis of the rectum masquerading as Crohn's disease: case report and review. *Clin Infect Dis*. 1998;26:521–523.
235. Lyon GM, Smilack JD, Komatsu KK, et al. Gastrointestinal basidiobolomycosis in Arizona: clinical and epidemiological characteristics and review of the literature. *Clin Infect Dis*. 2001;32:1448–1455.
236. Yousef OM, Smilack JD, Kerr DM, et al. Gastrointestinal basidiobolomycosis. Morphologic findings in a cluster of six cases. *Am J Clin Pathol*. 1999;112:610–616.
237. Kimura M, Furuta T, Maekura S, et al. Eumycotic mycetoma of the lower leg. *Rinsho Byori*. 1997;45:801–804.
238. Guarro J, Aguilar C, Pujol I. In-vitro antifungal susceptibilities of *Basidiobolus* and *Conidiobolus* spp. strains. *J Antimicrob Chemother*. 1999;44:557–560.
239. Dworzack DL, Pollock AS, Hodges GR, et al. Zygomycosis of the maxillary sinus and palate caused by *Basidiobolus haptosporus*. *Arch Intern Med*. 1978;138:1274–1276.
240. Foss NT, Rocha MR, Lima VT, et al. Entomophthoromycosis: therapeutic success by using amphotericin B and terbinafine. *Dermatology*. 1996;193:258–260.
241. Gupta M, Narang T, Kaur RJ, et al. A prospective case series evaluating efficacy and safety of combination of itraconazole and potassium iodide in rhinofacial conidiobolomycosis. *Int J Dermatol*. 2016;55:208–214.
242. Kumaravel S, Bharath K, Rajesh NG, et al. Delay and misdiagnosis of basidiobolomycosis in tropical South India: case series and review of the literature. *Paediatr Int Child Health*. 2016;36:52–57.
243. Mandhan P, Hassan KO, Samaan SM, et al. Visceral basidiobolomycosis: an overlooked infection in immunocompetent children. *Afr J Paediatr Surg*. 2015;12:193–196.
244. Zabolinejad N, Naseri A, Davoudi Y, et al. Aelami MH. Colonic basidiobolomycosis in a child: report of a culture-proven case. *Int J Infect Dis*. 2014;22:41–43.
245. Rose SR, Lindsley MD, Hurst SF, et al. Gastrointestinal basidiobolomycosis treated with posaconazole. *Med Mycol Case Rep*. 2012;2:11–14.

SHORT VIEW SUMMARY

Microbiology and Epidemiology

- *Sporothrix schenckii* sensu lato, a group of closely related species, are dimorphic fungi that cause sporotrichosis with budding yeast in tissue and a mold in culture. Fungal cells in tissue may be oval, spherical, or elongate (cigar-shaped), with variable size in the same lesion.
- Worldwide distribution, especially in tropical and subtropical regions.
- Acquisition is associated with exposure to soil, plants, plant products (hay, straw, sphagnum moss), and a variety of animals (especially cats).
- Grows slowly on fungal culture from involved tissues.

Diagnosis

- Lymphocutaneous disease: presentation with an indolent papulonodular lesion, sometimes ulcerating, should suggest the diagnosis. Secondary lesions along lymphangitic channels are characteristic. Biopsy will show pseudoepitheliomatous hyperplasia with

pyogranulomas and microabscesses in the dermis. Yeastlike cells can be seen with Gomori methenamine silver stain in one-third of cases but are not seen with hematoxylin and eosin stain. Culture will be positive.

- The most common extracutaneous form is osteoarticular. Pulmonary disease (usually cavitary) and meningeal disease are also well described. All are diagnosed by culture. Organisms can be seen in tissue with Gomori methenamine silver stain but may not be sufficiently distinctive to permit diagnosis of sporotrichosis.
- Multifocal dissemination may also occur, in immunocompromised subjects: cultures of skin lesions and joints are usually positive, whereas blood, spinal fluid, and bone marrow cultures are only occasionally positive.

Therapy

- Itraconazole at 100 to 200 mg/day (or 65 mg to 130 mg/day if using the enhanced bioavailability SUBA-itraconazole formulation;

see Chapter 40B) orally is the treatment of choice for lymphocutaneous disease, with 3 to 6 months of therapy typically needed for complete resolution. Terbinafine 250 mg daily has also been successful in some cases of cutaneous disease.

- Therapy with a saturated solution of potassium iodide is also effective for lymphocutaneous disease, although it is associated with many side effects.
- Itraconazole is also active in extracutaneous disease, but extended therapy may be required. In difficult settings, amphotericin B is also employed. Success has also been reported in a few cases with terbinafine. Improvement of immune status (e.g., introduction of antiretroviral therapy in subjects with human immunodeficiency virus coinfection, reduction of immunosuppressive agents) is useful.

MYCOLOGY

Sporothrix schenckii sensu lato (meaning in the broadest sense) is a dimorphic fungus that exists in a hyphal form in vitro at temperatures less than 37°C. Colonies are initially white but gradually become brown to black due to the production of pigmented conidia. In vivo or at 37°C on rich media such as brain heart infusion agar, the organism converts to an oval- or cigar-shaped budding yeast. Along with the characteristic morphology of the sporulating mold, identification is based on demonstration of this conversion to a yeast form. DNA sequencing has shown that *S. schenckii* sensu stricto (meaning in a narrow sense) is a cluster of several closely related species.¹⁻³ Identification to the species level is currently done by partial sequencing of the calmodulin gene. Identification of species by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry appears promising.⁶ There are small morphologic and biochemical differences between some species, but these are not sufficient for identification. As an exception, *Sporothrix luriei* has been rarely isolated from humans and differs by producing a variety of unusual shapes in vivo.⁷ The most common species is *S. schenckii* sensu stricto, which is found in America, Asia, and Africa. *Sporothrix brasiliensis* has so far been found only in Brazil, where it has often caused disseminated infection. Erythema nodosum or erythema multiforme has accompanied skin lesions in some of the cutaneous *S. brasiliensis* cases.⁸ *Sporothrix mexicana* has been isolated from a few environmental sources and infections in Mexico, Brazil, Italy, and Portugal. *Sporothrix pallida*, which may be nonpathogenic, and *Sporothrix globosa* have been more broadly distributed geographically.⁵ Because most of the information about sporotrichosis has not distinguished the cryptic species within the *Sporothrix schenckii* complex, the term *S. schenckii* is understood for the remainder of this chapter to include these closely related cryptic species.⁴

EPIDEMIOLOGY

Sporotrichosis has been reported from locations around the globe, but most case reports come from the tropical and subtropical regions of the Americas.^{2,9} Most of the cases in the United States so far have been identified as *S. schenckii*.¹⁰ Regions of hyperendemicity are known.^{9,11} *Sporothrix schenckii* is most often isolated from soil, plants, or plant products such as straw, wood, sphagnum moss, and thorny plants, though the fungus is not a plant pathogen. Scratches on exposed skin of florists, rose gardeners, horticulturalists, farmers, miners, and armadillo hunters have increased risk of infection.^{2,12,13} Because most cases appear to be due to occupational or avocational exposure to these materials, typically in the form of gardening or farming, patients with suggestive syndromes should be asked about these activities. Cases of animal-to-human transmission involving squirrels, horses, dogs, cats, pigs, mules, insects, and birds have been described.^{14,15} An ongoing epidemic of sporotrichosis in Brazil has its origin in thousands of urban cats infected with *Sporothrix brasiliensis*, transmitting infection to thousands of humans and dogs.^{11,16} Finally, sporotrichosis was apparently transmitted from the infected cheek of a mother to her infant¹⁷ and to a lung transplant recipient via the transplanted lung.¹⁸

CLINICAL SYNDROMES

Infections due to *S. schenckii* can be divided into several syndromes. The lymphocutaneous forms are the most common.

Lymphocutaneous Sporotrichosis

Cutaneous disease arises at sites of minor trauma and inoculation of the fungus into the skin. The initial lesion is most often on a distal extremity, but almost any site may be involved, including such central locations as the nose and the ocular adnexa.^{2,19,20} This preference



FIG. 259.1 Sporotrichosis of the fifth finger in a gardener. Three nodular lesions are visible on the hand and arm.

for cooler parts of the body corresponds to the known intolerance of some strains of *S. schenckii* to growth at 37°C.²¹ Initial lesions are papulonodular, often erythematous, and range in size from a few millimeters to 2 to 4 cm. The lesions may be smooth or verrucous, and they often ulcerate and develop raised erythematous borders.¹⁹ Lesions often develop proximally along lymphatic channels—these secondary lesions evolve in the same fashion as the primary lesion (Fig. 259.1). Secondary lesions do not usually involve a lymph node, although lymphadenopathy may develop. The lesions are typically painless, even after they ulcerate. The fixed, or plaque, form of sporotrichosis differs by not demonstrating any tendency to spread locally. Although spontaneous resolution of fixed sporotrichosis has been described,¹⁹ the lesions of sporotrichosis usually wax and wane over months to years. The patient will not have systemic symptoms, and laboratory examinations will be normal.

The indolent progression and physical examination features suggesting both lymphocutaneous and fixed sporotrichosis are also produced by a number of other organisms (Table 259.1). Cultures of the drainage from skin lesions are occasionally helpful, but culture of biopsy material is preferred and is diagnostic when positive. Microscopic examination will reveal pseudoepitheliomatous hyperplasia in the epidermis with pyogranulomas and granulomas, often with areas of liquefactive necrosis in the mid and upper dermis.²² Gomori methenamine silver stain will detect yeast cells in about one-third of cases, particularly if multiple sections are examined.²³

Extracutaneous Sporotrichosis

Osteoarticular involvement is the most common form of extracutaneous sporotrichosis.^{2,24} Involvement is of the major joints of the extremities—wrist (Fig. 259.2), elbow, ankle, and knee (Fig. 259.3); the hip, shoulder, and spine are not involved.²⁵ Most patients present with involvement of a single joint without previously having had sporotrichosis at any other site. The joint is swollen and painful on motion, an effusion is present, and a sinus tract may develop. The overlying skin may or may not be erythematous. Systemic symptoms are minimal and, other than elevation of the erythrocyte sedimentation rate, laboratory examinations are unrevealing. Untreated, other joints may become involved. Tenosynovitis associated with carpal tunnel syndrome or nerve entrapment has been reported.²⁶ The radiologic changes of osteomyelitis develop slowly and include loss of articular cartilage, periosteal reaction, and periarticular osteopenia and cystic changes. Failure to consider the diagnosis has resulted in an average 25-month delay before diagnosis.²⁷ Repeated culture of fluid from joint aspiration, as well as culture and microscopic examination of tissue from synovial biopsies, is often required to make the diagnosis. Differential considerations include pigmented villonodular synovitis, tuberculosis, gout, osteoarthritis, and rheumatoid arthritis.



FIG. 259.2 Sporotrichosis of the bones of the wrist.

TABLE 259.1 Differential Diagnosis of Sporotrichoid Lesions

Lesions Resembling Lymphocutaneous Sporotrichosis

(papulonodular lesions with or without central ulceration and with one or more nodules in the proximal skin along paths of presumed lymphatic spread)

Nocardiosis due to *Nocardia brasiliensis*
 Cutaneous leishmaniasis
 Mycobacterial infection due to:
 M. tuberculosis (tuberculosis cutis verrucosa)
 M. marinum
 M. chelonae
 M. kansasii
 M. fortuitum

Infections that may rarely resemble lymphocutaneous sporotrichosis:

Mycobacterium leprae
 Cowpox
 Staphylococcus aureus
 Streptococcus pyogenes
 Francisella tularensis
 Scedosporium apiospermum
 Fusarium species

Lesions Resembling Plaque Sporotrichosis

(chronic, indurated hyperkeratotic plaques)

Infections—as for lymphocutaneous disease and also:

 Blastomycosis
 Paracoccidioidomycosis
 Chromoblastomycosis
 Lobomycosis
 Neoplasms
 Squamous carcinoma
 Basal cell carcinoma
 Mycosis fungoides

Other

 Psoriasis
 Lupus vulgaris
 Pyoderma gangrenosum

Modified from Kostman JR, DiNubile MJ. Nodular lymphangitis: a distinctive but often unrecognized syndrome. *Ann Intern Med.* 1993;118:883–888; and Smego RA Jr, Castiglia M, Asperilla MO. Lymphocutaneous syndrome—a review of non-sporothrix causes. *Medicine.* 1999;78:38–63.

Pulmonary sporotrichosis is well described and may present with multifocal noncavitary disease as part of the syndrome of multifocal extracutaneous infection (see later) or primary cavitary disease following fungal inhalation.^{28,29} The typical patient with primary cavitary disease is a 30- to 60-year-old-male. Approximately one-third of the patients are alcoholic; one-third have a concomitant medical illness such as pulmonary tuberculosis, diabetes mellitus, sarcoidosis, and steroid use; and one-third are apparently normal. Patients are occasionally asymptomatic but will usually have a productive cough, low-grade fever, or weight loss. Other than elevation of the erythrocyte sedimentation rate, laboratory abnormalities are minimal. The chest radiograph reveals unilateral or bilateral cavitary lesions, usually with an associated parenchymal infiltrate



FIG. 259.3 Sporotrichosis of the knee with formation of a Baker cyst. (From Kwon-Chung KJ, Bennett J. *Medical Mycology*. Philadelphia: Lee & Febiger; 1992:712.)



FIG. 259.5 Hematogenously disseminated skin lesions scattered over the skin of a previously normal male farmer who also had joint lesions.



FIG. 259.4 Chest roentgenogram demonstrating extensive bilateral cavitation due to sporotrichosis.

(Fig. 259.4). Pleural effusions and hilar lymphadenopathy are occasionally noted. Gram stain or cytologic examination of sputum or bronchial washings will sometimes reveal elongated budding yeast,³⁰ and sputum culture will usually yield the organism. With some patients, however, repeated cultures and long-term follow-up are necessary in order to make the diagnosis.³¹ Untreated, the cavities of pulmonary sporotrichosis gradually enlarge and produce progressive pulmonary dysfunction. A single case of spontaneous resolution of noncavitary infection has been reported.³² The differential diagnosis includes mycobacterial infections (due to both *Mycobacterium tuberculosis* and other mycobacteria), histoplasmosis, and coccidioidomycosis.

An indolent meningitis due to *S. schenckii* has been described, with skin lesions or other extra-central nervous system foci noted in about half of cases.^{33–36} Cerebrospinal fluid (CSF) analysis demonstrates a lymphocytic pleocytosis, an elevated protein, and hypoglycorrhachia. Cultures of the CSF are positive, although repeated cultures of large volumes of CSF or serologic studies may be required to make the diagnosis. Infections of a variety of other sites have been reported but are uncommon. Involvement of the ocular adnexa, sometimes with spread to the eye, has been described.^{20,37} Endophthalmitis may even occur without prior trauma or other evidence of sporotrichosis.³⁸ Cases of isolated involvement of the sinuses, kidney, testes, and epididymis have also been reported.^{24,39}

Multifocal Extracutaneous Sporotrichosis

In otherwise normal patients with extracutaneous sporotrichosis, the lesions are generally restricted to a single site and are only locally progressive. Occasionally a patient with osteoarticular sporotrichosis will have involvement of several joints, but the presentation is otherwise identical to that in patients with involvement of only a single joint. A much smaller group of patients, on the other hand, present with weight loss or variable low-grade fever and often have several widely scattered cutaneous lesions without necessarily showing a single primary distal extremity lesion with the pattern of lymphangitic spread (Fig. 259.5). Mild anemia, leukocytosis, and elevation of the erythrocyte sedimentation rate may be present. Osteolytic bone lesions and arthritis are common, and spread to the palate, eyes, and central nervous system may develop.^{24,40} Noncavitary lung lesions may also be seen.⁴¹ Untreated infection is ultimately fatal. Patients with this form of sporotrichosis almost always have some form of immunosuppression, commonly hematologic malignancy,^{24,40} or human immunodeficiency virus (HIV) infection (see later). Cultures of skin lesions and joints are usually positive, whereas blood and bone marrow cultures are occasionally positive. Immunosuppressed patients with what appears to be simple cutaneous sporotrichosis should be examined for other sites of infection, and a technetium pyrophosphate bone scan or positron

emission tomography/computed tomography (PET/CT) should be performed.^{41a}

CLINICAL MANIFESTATIONS OF SPOROTRICHOSIS IN THE HIV-INFECTED PATIENT

Infection with HIV predisposes to invasive, atypical, or disseminated manifestations of sporotrichosis. When CD4⁺ counts are relatively well preserved, localized infection may follow direct cutaneous inoculation in a pattern analogous to that in immunocompetent patients.⁴² However, widespread lymphocutaneous sporotrichosis or multifocal extracutaneous disease may be seen in patients with more advanced HIV infection. In a review of patients with acquired immunodeficiency syndrome (AIDS) with disseminated sporotrichosis,⁴³ almost all had fewer than 100 CD4⁺ T cells/ μ L. Multiple ulcerative skin lesions are usually present. Sporotrichosis may also present as multifocal tenosynovitis and arthritis with or without overt cutaneous disease or systemic dissemination and thus may resemble disseminated gonococcal infection or the seronegative spondyloarthropathies such as reactive arthritis or psoriatic arthritis (Fig. 259.6), which are seen with a higher frequency in the setting of AIDS.⁴⁴ Widespread visceral dissemination also occurs, as evidenced by reports of meningitis with parenchymal brain lesions^{35,36}; lung abscess, liver, and spleen involvement⁴⁵; endophthalmitis⁴⁶; and fungemia with spread to esophagus, colon, testes, bone marrow, and lymph nodes.⁴⁶ Dissemination may manifest as part of the immune reconstitution inflammatory syndrome.⁴⁷ Sinusitis with invasion of the contiguous bone and soft tissues has also been described,⁴⁸ and emphasizes the potential for the respiratory tract as the initial focus of infection in HIV-infected patients.

DIAGNOSIS

Diagnosis is best made by culture of the affected site, though repeated attempts at culture may have to be made. A positive culture from any site is ordinarily diagnostic of infection, although a case of saprophytic involvement of the respiratory tract has been described.⁴⁹ A positive blood culture strongly suggests the multifocal form of sporotrichosis seen in immunocompromised hosts, and the lysis-centrifugation system may be more sensitive in detecting fungemia in nonimmunocompromised patients.⁵⁰ Serodiagnosis has been described⁵¹ but has often been complicated by the presence of antibodies in individuals without evidence

of sporotrichosis⁵² and by cross-reactivity with other fungi.² Molecular diagnostic techniques such as polymerase chain reaction have also been described.⁵³

Examination of biopsy specimens reveals a pyogranulomatous response and is diagnostic if characteristic 1- to 3- μ m \times 3- to 10- μ m cigar-shaped yeast forms are seen. Unfortunately, the yeast may be difficult to detect unless multiple sections are examined,²³ although lesions from immunocompromised hosts may contain numerous yeasts (Fig. 259.7). In addition, *S. schenckii* often assumes a more rounded tissue form, making the biopsy suggestive but not diagnostic. The organisms may be surrounded by a stellate, periodic acid–Schiff–positive, eosinophilic material known as an *asteroid body*. In the brain or eye, a capsule has sometimes been demonstrable around the yeastlike cells.

As with other immunosuppressed patients, individuals with advanced AIDS may have a high fungal load that results in positive smears and cultures.³⁰ Skin biopsy may reveal fungal elements with a limited inflammatory response, and this should prompt the clinician to initiate a search for a systemic immunodeficiency.⁵⁴

THERAPY

Treatment guidelines for sporotrichosis have been proposed.⁵⁵ Due to its convenience and consistent efficacy,⁵⁶ itraconazole at 100 to 200 mg/day (or 65 mg to 130 mg/day if using the enhanced bioavailability SUBA-itraconazole formulation; see Chapter 40B) has become the therapy of choice for cutaneous sporotrichosis.⁵⁵ Therapy with itraconazole is given for 2 to 4 weeks beyond complete resolution of all lesions and usually requires 3 to 6 months to effect a clinical cure. Relapse has been observed on occasion after cessation of therapy. Should relapse develop, a higher dose of itraconazole (200 mg twice daily), or terbinafine or iodide, may be tried. Terbinafine has been effective in adults at doses of 250 to 1000 mg/day for 12 to 24 weeks,^{37,57} but direct comparative data do not exist and the limited data suggest more frequent and more durable cures with higher doses. Further, the maximum US Food and Drug Administration–approved terbinafine regimen is 250 mg daily for 12 weeks; the safety of extended higher-dose therapy has not yet been extensively validated. Iodides are an effective and inexpensive but poorly tolerated therapy for cutaneous sporotrichosis⁵⁸ and have been effective in cases of therapeutic failure of itraconazole.⁵⁹ Iodide therapy is prescribed as a saturated solution of potassium iodide (SSKI), with therapy begun at 5 to 10 drops taken orally three times per day. The dose is gradually advanced to 25 to 40 drops three times daily (for children) or 40 to 50 drops three times daily (for adults). SSKI has a bitter taste and is made more palatable by taking it in milk, juice, or a carbonated beverage. Side effects include nausea, anorexia, diarrhea, parotid or lacrimal gland enlargement, and an acneiform rash. These side effects will remit with reduction of the dose of SSKI or temporary



FIG. 259.6 Extensive multifocal sporotrichosis with tenosynovitis of the toes, arthritis of the ankles, and associated lymphedema in a patient with advanced acquired immunodeficiency syndrome. This patient also had tenosynovitis of the wrists and hands along with arthritis of the wrists and knees.




FIG. 259.7 Numerous yeasts of sporotrichosis in a cutaneous lesion from an immunosuppressed patient. In the normal host, organisms are usually difficult to locate. Although a single cigar-shaped form is present (arrowhead), most of the yeasts have a rounded form that is consistent with, but not diagnostic of, sporotrichosis. (Courtesy Dr. Ronald Rapini, Houston, TX.)

cessation of therapy. For both terbinafine and iodide, therapy should be continued until 2 to 4 weeks after the cutaneous lesions have resolved, a process that usually takes 3 to 6 months. Some patients are allergic to iodides, and in others cutaneous disease may respond slowly to iodide therapy or rarely fail to respond at all.

Ketoconazole has not proven to be effective, and amphotericin B is too toxic to be used in this setting. Consistent with its limited in vitro activity, fluconazole has only modest clinical activity and should be used only if other therapies are not tolerated.^{60–62} Clinical data with the newer azoles are sparse,⁶³ and elevated minimum inhibitory concentrations have been reported with both voriconazole and posaconazole.^{60–62,64} Due to the temperature sensitivity of this organism, local application of heat is a useful adjunct therapy and on occasion has been curative.⁶⁵ Given the toxicity of both the azoles and the iodides in pregnant women (skeletal bone deformities and goiter, respectively),⁶⁶ use of heat may be especially valuable during pregnancy.^{55,67}

Therapy of extracutaneous sporotrichosis is often difficult. Osteoarticular sporotrichosis has been treated with intravenous amphotericin B (a lipid-associated preparation is recommended due to the improved safety profile of these formulations),⁵⁵ but itraconazole at 200 mg twice daily for at least 1 year is the preferred approach. Ketoconazole (400–800 mg/day) and fluconazole (200–400 mg/day) appear less efficacious, and ketoconazole is hepatotoxic.^{55,68} As noted previously, very few clinical data are available with the newer azoles (voriconazole and posaconazole), and the echinocandins do not appear consistently active.^{2,60,61,69} Intraarticular amphotericin B is sometimes given, although its role has not been clearly defined. Surgical débridement is often employed, but its utility is also uncertain. SSKI has rarely been reported to be effective but usually is not.

If diagnosed prior to the development of cavities, pulmonary sporotrichosis may be treated with itraconazole, SSKI, or amphotericin B.^{28,29} Cure of cavitary disease typically requires pulmonary resection plus a perioperative course of itraconazole or amphotericin B.⁵⁵ Treatment failure is often associated with incomplete resection.⁴

Sporothrix schenckii meningitis does not consistently respond to amphotericin B, and the addition of 5-fluorocytosine may be warranted. Limited data suggest that itraconazole might be useful as suppressive or step-down therapy.⁵⁵ The number of reported cases of involvement of other specific sites is too limited to permit generalization. Extracutaneous sporotrichosis in the immunocompromised host usually responds at least partially to either amphotericin B or itraconazole, although relapse is common.

Therapy of Patients With Acquired Immunodeficiency Syndrome

Therapy of sporotrichosis in AIDS should be tailored to the presenting syndrome. Itraconazole appears to be the drug of choice, and individuals with limited cutaneous disease can be treated with 200 mg twice daily. Amphotericin B should be used as initial therapy of disseminated disease, with lipid-associated formulations preferred over amphotericin B deoxycholate.⁵⁵ Based on the observation of frequent relapse and dissemination, chronic suppressive therapy with itraconazole appears warranted following initial control of infection. Although unsupported by direct data, it appears reasonable to discontinue therapy in individuals who have been treated for itraconazole for at least 1 year and whose CD4⁺ cell counts have been about 200 cells/μL for at least a year.⁵⁵

Both anecdotal and published experience suggest that multifocal extracutaneous disease in HIV-infected patients may respond poorly, if at all, to current therapies.^{70,71} Therapy should be initiated with amphotericin B followed by lifetime suppression with itraconazole. Progression may occur despite amphotericin therapy. Because disseminated disease has been reported to develop despite ongoing fluconazole being given for other indications,⁷² fluconazole is not a first-line choice. As has been demonstrated for other opportunistic pathogens, the use of highly active antiretroviral therapies may also assist in clearing the infection.⁷³

Itraconazole Blood Level Monitoring

When itraconazole is used for noncutaneous infection or in HIV-infected subjects, confirmation of adequate blood levels is recommended (see Chapter 40B). In HIV in particular, levels may be reduced by achlorhydria, malabsorption, or diarrhea due to other opportunistic pathogens. As for other fungal infections, target levels of the parent (unmetabolized) itraconazole molecule of at least 500 ng/mL by high-performance liquid chromatography generally appear adequate. The increased bioavailability of itraconazole cyclodextrin suspension is helpful in achieving such blood levels.

PROGNOSIS

Cutaneous sporotrichosis responds well to therapy and has an excellent prognosis. Osteoarticular sporotrichosis may require prolonged therapy but is not life threatening. Other forms of extracutaneous sporotrichosis can be difficult to treat and may have substantial morbidity and mortality.

Key References

The complete reference list is available online at Expert Consult.

- Marimon R, Cano J, Gene J, et al. *Sporothrix brasiliensis*, *S. globosa*, and *S. mexicana*, three new *Sporothrix* species of clinical interest. *J Clin Microbiol*. 2007;45:3198–3206.
- Barros MB, de Almeida Paes R, Schubach AO. *Sporothrix schenckii* and sporotrichosis. *Clin Microbiol Rev*. 2011;24:633–654.
- Zhang Y, Hagen F, Stielow B, et al. Phylogeography and evolutionary patterns in *Sporothrix* spanning more than 14000 human and animal case reports. *Persoonia*. 2015;35:1–20.
- Rangel-Gamboa L, Martinez-Hernandez F, Maravilla P, et al. Update of phylogenetic and genetic diversity of *Sporothrix schenckii* sensu lato. *Med Mycol*. 2016;54:248–255.
- Camacho E, Leon-Navarro I, Rodriguez-Brito S, et al. Molecular epidemiology of human sporotrichosis in Venezuela reveals high frequency of *Sporothrix globosa*. *BMC Infect Dis*. 2015;15:94.
- Ajello L, Kaplan W. A new variant of *Sporothrix schenckii*. *Mykosen*. 1969;12:633–644.
- Almeida-Paes R, de Oliveira MM, Freitas DE, et al. Sporotrichosis in Rio de Janeiro, Brazil: *Sporothrix brasiliensis* is associated with atypical clinical presentations. *PLoS Negl Trop Dis*. 2014;8:e3094.
- Chakrabarti A, Bonifaz A, Gutierrez-Galhardo MC, et al. Global epidemiology of sporotrichosis. *Med Mycol*. 2015;53:3–14.
- Schubach A, Barros MBD, Wanke B. Epidemic sporotrichosis. *Curr Opin Infect Dis*. 2008;21:129–133.
- Feeney KT, Arthur IH, Whittle AJ, et al. Outbreak of sporotrichosis, Western Australia. *Emerg Infect Dis*. 2007;13:1228–1231.
- Alves SH, Boettcher CS, Oliveira DC, et al. *Sporothrix schenckii* associated with armadillo hunting in Southern Brazil: epidemiological and antifungal susceptibility profiles. *Rev Soc Bras Med Trop*. 2010;43:523–525.
- Saravanakumar PS, Eslami P, Zar FA. Lymphocutaneous sporotrichosis associated with a squirrel bite: case report and review. *Clin Infect Dis*. 1996;23:647–648.
- Reed KD, Moore FM, Geiger GE, et al. Zoonotic transmission of sporotrichosis: case report and review. *Clin Infect Dis*. 1993;16:384–387.
- Sanchoene KO, Madrid IM, Klafke GB, et al. *Sporothrix brasiliensis* outbreaks and the rapid emergence of feline sporotrichosis. *Mycoses*. 2015;58:652–658.
- Smith LM. Sporotrichosis: report of four clinically atypical cases. *South Med J*. 1945;38:505–515.
- Bahr NC, Janssen K, Billings J, et al. Respiratory failure due to possible donor-derived *Sporothrix schenckii* infection in a lung transplant recipient. *Case Rep Infect Dis*. 2015;2015:925718.
- Bargman HB. Sporotrichosis of the nose with spontaneous cure. *Can Med Assoc J*. 1981;124:1027.
- Gordon D. Ocular sporotrichosis. *Arch Ophthalmol*. 1947;37:56–72.
- Kwon-Chung KJ. Comparison of isolates of *Sporothrix schenckii* obtained from fixed cutaneous lesions with isolates from other types of lesions. *J Infect Dis*. 1979;139:424–431.
- Bullpitt P, Weedon D. Sporotrichosis: a review of 39 cases. *Pathology*. 1978;10:249–256.
- Wilson DE, Mann JJ, Bennett JE, et al. Clinical features of extracutaneous sporotrichosis. *Medicine (Baltimore)*. 1967;46:265–279.
- Janex PC, Mann RJ. Extracutaneous sporotrichosis. *J Hand Surg Am*. 1987;12A:441–445.
- Stratton CW, Lichtenstein KA, Lowenstein SR, et al. Granulomatous tenosynovitis and carpal tunnel syndrome caused by *Sporothrix schenckii*. *Am J Med*. 1981;71:161–164.
- Crout JE, Brewer NS, Tompkins RB. Sporotrichosis arthritis: clinical features in seven patients. *Ann Intern Med*. 1977;86:294–297.
- Aung AK, Spelman DW, Thompson PJ. Pulmonary sporotrichosis: an evolving clinical paradigm. *Semin Respir Crit Care Med*. 2015;36:756–766.
- Pluss JL, Opal SM. Pulmonary sporotrichosis: review of treatment and outcome. *Medicine (Baltimore)*. 1986;65:143–153.
- Gori S, Lupetti A, Moscato G, et al. Pulmonary sporotrichosis with hyphae in a human immunodeficiency virus-infected patient. A case report. *Acta Cytol*. 1997;41:519–521.
- Khan FA, Guarneri JJ, Sierra MF. Primary pulmonary sporotrichosis complicated by perirectal abscess. *Am Rev Respir Dis*. 1975;112:119–123.
- Pueringer RJ, Iber C, Deike MA, et al. Spontaneous remission of extensive pulmonary sporotrichosis. *Ann Intern Med*. 1986;104:366–367.
- Scott EN, Kaufman L, Brown AC, et al. Serologic studies in the diagnosis and management of meningitis due to *Sporothrix schenckii*. *N Engl J Med*. 1987;317:935–940.
- Hessler C, Kauffman CA, Chow FC. The upside of bias: a case of chronic meningitis due to *Sporothrix schenckii* in an immunocompetent host. *Neurohospitalist*. 2017;7:30–34.
- Galhardo MCG, Silva MTT, Lima MA, et al. *Sporothrix schenckii* meningitis in AIDS during immune reconstitution syndrome. *J Neurol Neurosurg Psychiatry*. 2010;81:696–699.

36. Silva-Vergara ML, Maneira FRZ, de Oliveira RM, et al. Multifocal sporotrichosis with meningeal involvement in a patient with AIDS. *Med Mycol*. 2005;43:187–190.
37. Fan B, Wang JF, Zheng B, et al. Clinical features of 10 cases of eyelid sporotrichosis in Jilin Province (Northeast China). *Can J Ophthalmol*. 2016;51:297–301.
38. Font RL, Jakobiec FA. Granulomatous necrotizing retinochoroiditis caused by *Sporotrichum schenckii*. *Arch Ophthalmol*. 1976;94:1513–1519.
39. Friedman SJ, Doyle JA. Extracutaneous sporotrichosis. *Int J Dermatol*. 1983;22:171–173.
40. Lynch PJ, Voorhees JJ, Harrell ER. Systemic sporotrichosis. *Ann Intern Med*. 1970;73:23–30.
41. Aung AK, Teh BM, McGrath C, et al. Pulmonary sporotrichosis: case series and systematic analysis of literature on clinico-radiological patterns and management outcomes. *Med Mycol*. 2013;51:534–544.
- 41a. Bassetti MA, Carnelutti D, Muser E, et al. 18F-Fluorodeoxyglucose positron emission tomography and infectious diseases: current applications and future perspectives. *Curr Opin Infect Dis*. 2017;30:192–200.
42. Keiser P, Whittle D. Sporotrichosis in human immunodeficiency virus-infected patients: report of a case. *Rev Infect Dis*. 1991;13:1027–1028.
43. Al-Tawfiq JA, Wools KK. Disseminated sporotrichosis and *Sporothrix schenckii* fungemia as the initial presentation of human immunodeficiency virus infection. *Clin Infect Dis*. 1998;26:1403–1406.
44. Oscherwitz SL, Rinaldi MG. Disseminated sporotrichosis in a patient infected with human immunodeficiency virus. *Clin Infect Dis*. 1992;15:568–569.
45. Lipstein-Kresch E, Isenberg HD, Singer C, et al. Disseminated *Sporothrix schenckii* infection with arthritis in a patient with acquired immunodeficiency syndrome. *J Rheumatol*. 1985;12:805–808.
46. Heller HM, Fuhrer J. Disseminated sporotrichosis in patients with AIDS: case report and review of the literature. *AIDS*. 1991;5:1243–1246.
47. Gutierrez-Galhardo MC, do Valle ACF, Fraga BLB, et al. Disseminated sporotrichosis as a manifestation of immune reconstitution inflammatory syndrome. *Mycoses*. 2010;53:78–80.
48. Morgan M, Reves R. Invasive sinusitis due to *Sporothrix schenckii* in a patient with AIDS. *Clin Infect Dis*. 1996;23:1319–1320.
49. Lowenstein M, Markowitz SM, Nottebart HC, et al. Existence of *Sporothrix schenckii* as a pulmonary saprophyte. *Chest*. 1978;73:419–421.
50. Kosinski RM, Axelrod P, Rex JH, et al. Wreiole A. *Sporothrix schenckii* fungemia without disseminated sporotrichosis. *J Clin Microbiol*. 1992;30:501–503.
51. Bernardes-Engemann AR, de Lima Barros M, Zeitone T, et al. Validation of a serodiagnostic test for sporotrichosis: a follow-up study of patients related to the Rio de Janeiro zoonotic outbreak. *Med Mycol*. 2015;53:28–33.
52. Scott EN, Muchmore HG. Immunoblot analysis of antibody responses to *Sporothrix schenckii*. *J Clin Microbiol*. 1989;27:300–304.
53. Rodrigues AM, de Hoog GS, de Camargo ZP. Molecular diagnosis of pathogenic *Sporothrix* species. *PLoS Negl Trop Dis*. 2015;9:e0004190.
54. Fitzpatrick JE, Eubanks S. Acquired immunodeficiency syndrome presenting as disseminated cutaneous sporotrichosis. *Int J Dermatol*. 1988;27:406–407.
55. Kauffman CA, Bustamante B, Chapman SW, et al. Clinical practice guidelines for the management of sporotrichosis: 2007 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2007;45:1255–1265.
56. Sharkey-Mathis PK, Kauffman CA, Graybill JR, et al. Treatment of sporotrichosis with itraconazole. *Am J Med*. 1993;95:279–285.
57. Chapman SW, Pappas P, Kauffmann C, et al. Comparative evaluation of the efficacy and safety of two doses of terbinafine (500 and 1000 mg day⁻¹) in the treatment of cutaneous or lymphocutaneous sporotrichosis. *Mycoses*. 2004;47:62–68.
58. Yamada K, Zaitz C, Framil VM, et al. Cutaneous sporotrichosis treatment with potassium iodide: a 24-year experience in Sao Paulo State, Brazil. *Rev Inst Med Trop Sao Paulo*. 2011;53:89–93.
59. Fischman Gompertz O, Rodrigues AM, Fernandes GF, et al. Atypical clinical presentation of sporotrichosis caused by *Sporothrix globosa* resistant to itraconazole. *Am J Trop Med Hyg*. 2016;94:1218–1222.
60. Brilhante RS, Rodrigues AM, Sidrim JJ, et al. In vitro susceptibility of antifungal drugs against *Sporothrix brasiliensis* recovered from cats with sporotrichosis in Brazil. *Med Mycol*. 2016;54:275–279.
61. Marimon R, Serena C, Gen J, et al. In vitro antifungal susceptibilities of five species of *Sporothrix*. *Antimicrob Agents Chemother*. 2008;52:732–734.
62. Espinel-Ingroff A, Boyle K, Sheehan DJ. In vitro antifungal activities of voriconazole and reference agents as determined by NCCLS methods: review of the literature. *Mycopathologia*. 2001;150:101–115.
63. Paixao AG, Galhardo MCG, Almeida-Paes R, et al. The difficult management of disseminated *Sporothrix brasiliensis* in a patient with advanced AIDS. *AIDS Res Ther*. 2015;12:16.
64. Almeida-Paes R, Oliveira MME, Freitas DFS, et al. Refractory sporotrichosis due to *Sporothrix brasiliensis* in humans appears to be unrelated to in vivo resistance. *Med Mycol*. 2017;55:507–517.
65. Galiana J, Conti-Diaz IA. Healing effects of heat and a rubefacient on nine cases of sporotrichosis. *Sabouraudia*. 1963;3:64–71.
66. Sobel JD. Use of antifungal drugs in pregnancy: a focus on safety. *Drug Saf*. 2000;1:77–85.
67. Vanderveen EE, Messenger AL, Voorhees JJ. Sporotrichosis in pregnancy. *Cutis*. 1982;30:761–763.
68. Calhoun DL, Washkin H, White MP, et al. Treatment of systemic sporotrichosis with ketoconazole. *Rev Infect Dis*. 1991;13:47–51.
69. Nakai T, Uno J, Ikeda F, et al. In vitro antifungal activity of micafungin (FK463) against dimorphic fungi: comparison of yeast-like and mycelial forms. *Antimicrob Agents Chemother*. 2003;47:1376–1381.
70. Kauffman CA. Old and new therapies for sporotrichosis. *Clin Infect Dis*. 1995;21:981–985.
71. Donabedian H, O'Donnell E, Olszewski C, et al. Disseminated cutaneous and meningeal sporotrichosis in an AIDS patient. *Diagn Microbiol Infect Dis*. 1994;18:111–115.
72. Goldani LZ, Aquino VR, Dargel AA. Disseminated cutaneous sporotrichosis in an AIDS patient receiving maintenance therapy with fluconazole for previous cryptococcal meningitis. *Clin Infect Dis*. 1999;28:1337–1338.
73. Carr A, Marriott D, Field A, et al. Treatment of HIV-1-associated microsporidiosis and cryptosporidiosis with combination antiretroviral therapy. *Lancet*. 1998;351:256–261.

References

- Marimon R, Cano J, Gene J, et al. *Sporothrix brasiliensis*, *S. globosa*, and *S. mexicana*, three new *Sporothrix* species of clinical interest. *J Clin Microbiol*. 2007;45:3198–3206.
- Barros MB, de Almeida Paes R, Schubach AO. *Sporothrix schenckii* and sporotrichosis. *Clin Microbiol Rev*. 2011;24:633–654.
- Zhang Y, Hagen F, Stielow B, et al. Phylogeography and evolutionary patterns in *Sporothrix* spanning more than 14 000 human and animal case reports. *Persoonia*. 2015;35:1–20.
- Rangel-Gamboa L, Martinez-Hernandez F, Maravilla P, et al. Update of phylogenetic and genetic diversity of *Sporothrix schenckii* sensu lato. *Med Mycol*. 2016;54:248–255.
- Camacho E, Leon-Navarro I, Rodriguez-Brito S, et al. Molecular epidemiology of human sporotrichosis in Venezuela reveals high frequency of *Sporothrix globosa*. *BMC Infect Dis*. 2015;15:94.
- Oliveira MMA, Santos C, Sampaio P, et al. Development and optimization of a new MALDI-TOF protocol for identification of the *Sporothrix* species complex. *Res Microbiol*. 2015;166:102–110.
- Ajello L, Kaplan W. A new variant of *Sporothrix schenckii*. *Mykosen*. 1969;12:633–644.
- Almeida-Paes R, de Oliveira MM, Freitas DF, et al. Sporotrichosis in Rio de Janeiro, Brazil: *Sporothrix brasiliensis* is associated with atypical clinical presentations. *PLoS Negl Trop Dis*. 2014;8:e3094.
- Chakrabarti A, Bonifaz A, Gutierrez-Galhardo MC, et al. Global epidemiology of sporotrichosis. *Med Mycol*. 2015;53:3–14.
- Rodrigues AM, de Hoog S, de Camargo ZP. Emergence of pathogenicity in the *Sporothrix schenckii* complex. *Med Mycol*. 2012.
- Schubach A, Barros MBD, Wanke B. Epidemic sporotrichosis. *Curr Opin Infect Dis*. 2008;21:129–133.
- Feeney KT, Arthur IH, Whittle AJ, et al. Outbreak of sporotrichosis, Western Australia. *Emerg Infect Dis*. 2007;13:1228–1231.
- Alves SH, Boettcher CS, Oliveira DC, et al. *Sporothrix schenckii* associated with armadillo hunting in Southern Brazil: epidemiological and antifungal susceptibility profiles. *Rev Soc Bras Med Trop*. 2010;43:523–525.
- Saravanakumar PS, Eslami P, Zar FA. Lymphocutaneous sporotrichosis associated with a squirrel bite: case report and review. *Clin Infect Dis*. 1996;23:647–648.
- Reed KD, Moore FM, Geiger GE, et al. Zoonotic transmission of sporotrichosis: case report and review. *Clin Infect Dis*. 1993;16:384–387.
- Sanchothene KO, Madrid IM, Klafke GB, et al. *Sporothrix brasiliensis* outbreaks and the rapid emergence of feline sporotrichosis. *Mycoses*. 2015;58:652–658.
- Smith LM. Sporotrichosis: report of four clinically atypical cases. *South Med J*. 1945;38:505–515.
- Bahr NC, Janssen K, Billings J, et al. Respiratory failure due to possible donor-derived *Sporothrix schenckii* infection in a lung transplant recipient. *Case Rep Infect Dis*. 2015;2015:925718.
- Bargman HB. Sporotrichosis of the nose with spontaneous cure. *Can Med Assoc J*. 1981;124:1027.
- Gordon D. Ocular sporotrichosis. *Arch Ophthalmol*. 1947;37:56–72.
- Kwon-Chung KJ. Comparison of isolates of *Sporothrix schenckii* obtained from fixed cutaneous lesions with isolates from other types of lesions. *J Infect Dis*. 1979;139:424–431.
- Quintella LP, Passos SLR, do Vale ACF, et al. Histopathology of cutaneous sporotrichosis in Rio de Janeiro: a series of 119 consecutive cases. *J Cutan Pathol*. 2011;38:25–32.
- Bullpitt P, Weedon D. Sporotrichosis: a review of 39 cases. *Pathology*. 1978;10:249–256.
- Wilson DE, Mann JJ, Bennett JE, et al. Clinical features of extracutaneous sporotrichosis. *Medicine (Baltimore)*. 1967;46:265–279.
- James PC, Mann RJ. Extracutaneous sporotrichosis. *J Hand Surg Am*. 1987;12A:441–445.
- Stratton CW, Lichtenstein KA, Lowenstein SR, et al. Granulomatous tenosynovitis and carpal tunnel syndrome caused by *Sporothrix schenckii*. *Am J Med*. 1981;71:161–164.
- Crout JE, Brewer NS, Tompkins RB. Sporotrichosis arthritis: clinical features in seven patients. *Ann Intern Med*. 1977;86:294–297.
- Aung AK, Spelman DW, Thompson PJ. Pulmonary sporotrichosis: an evolving clinical paradigm. *Semin Respir Crit Care Med*. 2015;36:756–766.
- Pluss JL, Opal SM. Pulmonary sporotrichosis: review of treatment and outcome. *Medicine (Baltimore)*. 1986;65:143–153.
- Gori S, Lupetti A, Moscato G, et al. Pulmonary sporotrichosis with hyphae in a human immunodeficiency virus-infected patient. A case report. *Acta Cytol*. 1997;41:519–521.
- Khan FA, Guarneri JJ, Sierra MF. Primary pulmonary sporotrichosis complicated by perirectal abscess. *Am Rev Respir Dis*. 1975;112:119–123.
- Pueringer RJ, Iber C, Deike MA, et al. Spontaneous remission of extensive pulmonary sporotrichosis. *Ann Intern Med*. 1986;104:366–367.
- Scott EN, Kaufman L, Brown AC, et al. Serologic studies in the diagnosis and management of meningitis due to *Sporothrix schenckii*. *N Engl J Med*. 1987;317:935–940.
- Hessler C, Kauffman CA, Chow FC. The upside of bias: a case of chronic meningitis due to *Sporothrix schenckii* in an immunocompetent host. *Neurohospitalist*. 2017;7:30–34.
- Galhardo MCG, Silva MTT, Lima MA, et al. *Sporothrix schenckii* meningitis in AIDS during immune reconstitution syndrome. *J Neurol Neurosurg Psychiatry*. 2010;81:696–699.
- Silva-Vergara ML, Maneira FRZ, de Oliveira RM, et al. Multifocal sporotrichosis with meningeal involvement in a patient with AIDS. *Med Mycol*. 2005;43:187–190.
- Fan B, Wang JF, Zheng B, et al. Clinical features of 10 cases of eyelid sporotrichosis in Jilin Province (Northeast China). *Can J Ophthalmol*. 2016;51:297–301.
- Font RL, Jakobiec FA. Granulomatous necrotizing retinochoroiditis caused by *Sporotrichum schenckii*. *Arch Ophthalmol*. 1976;94:1513–1519.
- Friedman SJ, Doyle JA. Extracutaneous sporotrichosis. *Int J Dermatol*. 1983;22:171–173.
- Lynch PJ, Voorhees JJ, Harrell ER. Systemic sporotrichosis. *Ann Intern Med*. 1970;73:23–30.
- Aung AK, Teh BM, McGrath C, et al. Pulmonary sporotrichosis: case series and systematic analysis of literature on clinico-radiological patterns and management outcomes. *Med Mycol*. 2013;51:534–544.
- Bassetti MA, Carnelutti D, Muser E, et al. 18F-Fluorodeoxyglucose positron emission tomography and infectious diseases: current applications and future perspectives. *Curr Opin Infect Dis*. 2017;30:192–200.
- Keiser P, Whittle D. Sporotrichosis in human immunodeficiency virus-infected patients: report of a case. *Rev Infect Dis*. 1991;13:1027–1028.
- Al-Tawfiq JA, Wools KK. Disseminated sporotrichosis and *Sporothrix schenckii* fungemia as the initial presentation of human immunodeficiency virus infection. *Clin Infect Dis*. 1998;26:1403–1406.
- Oscherwitz SL, Rinaldi MG. Disseminated sporotrichosis in a patient infected with human immunodeficiency virus. *Clin Infect Dis*. 1992;15:568–569.
- Lipstein-Kresch E, Isenberg HD, Singer C, et al. Disseminated *Sporothrix schenckii* infection with arthritis in a patient with acquired immunodeficiency syndrome. *J Rheumatol*. 1985;12:805–808.
- Heller HM, Fuhrer J. Disseminated sporotrichosis in patients with AIDS: case report and review of the literature. *AIDS*. 1991;5:1243–1246.
- Gutierrez-Galhardo MC, do Valle ACF, Fraga BLB, et al. Disseminated sporotrichosis as a manifestation of immune reconstitution inflammatory syndrome. *Mycoses*. 2010;53:78–80.
- Morgan M, Reeves R. Invasive sinusitis due to *Sporothrix schenckii* in a patient with AIDS. *Clin Infect Dis*. 1996;23:1319–1320.
- Lowenstein M, Markowitz SM, Nottebart HC, et al. Existence of *Sporothrix schenckii* as a pulmonary saprophyte. *Chest*. 1978;73:419–421.
- Kosinski RM, Axelrod P, Rex JH, et al. *Sporothrix schenckii* fungemia without disseminated sporotrichosis. *J Clin Microbiol*. 1992;30:501–503.
- Bernardes-Engemann AR, de Lima Barros M, Zeitune T, et al. Validation of a serodiagnostic test for sporotrichosis: a follow-up study of patients related to the Rio de Janeiro zoonotic outbreak. *Med Mycol*. 2015;53:28–33.
- Scott EN, Muchmore HG. Immunoblot analysis of antibody responses to *Sporothrix schenckii*. *J Clin Microbiol*. 1989;27:300–304.
- Rodrigues AM, de Hoog GS, de Camargo ZP. Molecular diagnosis of pathogenic *Sporothrix* species. *PLoS Negl Trop Dis*. 2015;9:e0004190.
- Fitzpatrick JE, Eubanks S. Acquired immunodeficiency syndrome presenting as disseminated sporotrichosis. *Int J Dermatol*. 1988;27:406–407.
- Kauffman CA, Bustamante B, Chapman SW, et al. Clinical practice guidelines for the management of sporotrichosis: 2007 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2007;45:1255–1265.
- Sharkey-Mathis PK, Kauffman CA, Graybill JR, et al. Treatment of sporotrichosis with itraconazole. *Am J Med*. 1993;95:279–285.
- Chapman SW, Pappas P, Kauffmann C, et al. Comparative evaluation of the efficacy and safety of two doses of terbinafine (500 and 1000 mg day⁻¹) in the treatment of cutaneous or lymphocutaneous sporotrichosis. *Mycoses*. 2004;47:62–68.
- Yamada K, Zaitz C, Framil VM, et al. Cutaneous sporotrichosis treatment with potassium iodide: a 24-year experience in Sao Paulo State, Brazil. *Rev Inst Med Trop Sao Paulo*. 2011;53:89–93.
- Fischman Gompertz O, Rodrigues AM, Fernandes GF, et al. Atypical clinical presentation of sporotrichosis caused by *Sporothrix globosa* resistant to itraconazole. *Am J Trop Med Hyg*. 2016;94:1218–1222.
- Brilhante RS, Rodrigues AM, Sidrim JJ, et al. In vitro susceptibility of antifungal drugs against *Sporothrix brasiliensis* recovered from cats with sporotrichosis in Brazil. *Med Mycol*. 2016;54:275–279.
- Marimon R, Serena C, Gen J, et al. In vitro antifungal susceptibilities of five species of *Sporothrix*. *Antimicrob Agents Chemother*. 2008;52:732–734.
- Espinel-Ingroff A, Boyle K, Sheehan DJ. In vitro antifungal activities of voriconazole and reference agents as determined by NCCLS methods: review of the literature. *Mycopathologia*. 2001;150:101–115.
- Paixao AG, Galhardo MCG, Almeida-Paes R, et al. The difficult management of disseminated *Sporothrix brasiliensis* in a patient with advanced AIDS. *AIDS Res Ther*. 2015;12:16.
- Almeida-Paes R, Oliveira MME, Freitas DFS, et al. Refractory sporotrichosis due to *Sporothrix brasiliensis* in humans appears to be unrelated to in vivo resistance. *Med Mycol*. 2017;55:507–517.
- Galiana J, Conti-Diaz IA. Healing effects of heat and a rubefacient on nine cases of sporotrichosis. *Sabouraudia*. 1963;3:64–71.
- Sobel JD. Use of antifungal drugs in pregnancy: a focus on safety. *Drug Saf*. 2000;1:77–85.
- Vanderveen EE, Messenger AL, Voorhees JJ. Sporotrichosis in pregnancy. *Cutis*. 1982;30:761–763.
- Calhoun DL, Washkin H, White MP, et al. Treatment of systemic sporotrichosis with ketoconazole. *Rev Infect Dis*. 1991;13:47–51.
- Nakai T, Uno J, Ikeda F, et al. In vitro antifungal activity of micafungin (FK463) against dimorphic fungi: comparison of yeast-like and mycelial forms. *Antimicrob Agents Chemother*. 2003;47:1376–1381.
- Kauffman CA. Old and new therapies for sporotrichosis. *Clin Infect Dis*. 1995;21:981–985.
- Donabedian H, O'Donnell E, Olszewski C, et al. Disseminated cutaneous and meningeal sporotrichosis in an AIDS patient. *Diagn Microbiol Infect Dis*. 1994;18:111–115.
- Goldani LZ, Aquino VR, Dargel AA. Disseminated cutaneous sporotrichosis in an AIDS patient receiving maintenance therapy with fluconazole for previous cryptococcal meningitis. *Clin Infect Dis*. 1999;28:1337–1338.
- Carr A, Marriott D, Field A, et al. Treatment of HIV-1-associated microsporidiosis and cryptosporidiosis with combination antiretroviral therapy. *Lancet*. 1998;351:256–261.

SHORT VIEW SUMMARY

Definition

- Chromoblastomycosis (chromomycosis) is a chronic fungal infection limited to the skin and subcutaneous tissue.
- It produces scaly nodular, tumorous, verrucous, plaque or cicatricial lesions, typically affecting the lower extremities.
- It is defined by the microscopic presence of muriform cells.

Epidemiology

- Chromoblastomycosis has a worldwide distribution, with most cases occurring in tropical and subtropical regions.
- The largest numbers of cases are reported from Madagascar, Brazil, Mexico, Venezuela, and Costa Rica.
- The disease has a male predominance and is more prevalent in those aged 40 to 69. It is commonly associated with outdoor activities such as farming or woodcutting and with absence of footwear.

Microbiology

- Infection is most commonly caused by species of *Fonsecaea* (*F. pedrosoi*, *F. monophora*, *F. nubica*) or *Cladophialophora* (*C. carrionii*).
- Less commonly it is caused by *Exophiala*, *Phialophora*, or *Rhinocladiella* species.
- Other dark-walled fungi are rarely reported as causative agents.

Diagnosis

- The presence of muriform cells (also called sclerotic, "copper penny," or Medlar bodies) is pathognomonic.
- Muriform cells are observed microscopically in skin scrapings prepared with potassium hydroxide or in routinely stained skin biopsy specimens.
- Culture on standard mycologic media is possible but not always necessary.

Therapy

- No proven treatment has been identified.
- Small lesions may be treated successfully with surgical excision, liquid nitrogen or topical heat application, or photocoagulation. Use of curettage and electrocautery is discouraged because of reports of disease spread after their use.
- Oral itraconazole 200 to 400 mg daily or terbinafine 250 to 500 mg daily in combination with local liquid nitrogen is the most effective therapy.
- Posaconazole has been used successfully in a small number of patients with disease that is refractory to itraconazole or terbinafine.

Prevention

- No vaccine is available.
- The need for footwear and proper protective clothing should be stressed.

Chromoblastomycosis (chromomycosis) is a chronic, localized fungal infection of the skin and subcutaneous tissue that produces raised scaly lesions, usually of the lower extremities. The lesions of chromoblastomycosis are frequently warty or cauliflower-like in appearance, with pathognomonic muriform cells (also called "copper penny" or sclerotic bodies) found at histologic examination. This disease of tropical and subtropical distribution is produced by inoculation of the infecting fungi in association with minor trauma. Alexandrino Pedroso, for whom the major etiologic agent is named, first noted the disease in 1911, although the first publication to describe what was likely chromoblastomycosis appeared in 1914, authored by Max Rudolph.^{1,2} The first reports to include identification of the fungal cause of this disease were published 1 year later by Medlar and Lane, who described a patient with disease acquired not in the tropics but in the United States.

ETIOLOGIC AGENTS

Infection is caused by one of several dark-walled (dematiaceous) fungi found in the soil and in association with cacti, thorny plants, and other live or decaying vegetation. *Fonsecaea pedrosoi* is the most common cause of chromoblastomycosis. Other *Fonsecaea* species (*F. monophora* and *F. nubica*) and *Cladophialophora carrionii* are also common etiologic agents. *Phialophora verrucosa* and *Rhinocladiella aquaspersa* are less commonly reported. In the largest reports from Brazil,³ Mexico,⁴ Sri Lanka,⁵ India,⁶ and Japan,⁷ *F. pedrosoi* has been responsible for 66% to 96% of all infections. With the more recent recognition of *F. monophora* and *F. nubica* as new species, identification of these organisms as the cause of both new cases and old cases (with reidentification of isolates previously identified as *F. pedrosoi*) of chromoblastomycosis continues to increase.⁸⁻¹² *Fonsecaea compacta* is currently believed to be a variant

of *F. pedrosoi* and not a distinct species. Rare or isolated reports of disease caused by *Botryomyces caespitosus*, *Chaetomium funicola*, *Cladophialophora arxii*, *Cladophialophora boppii*, *Cyphellophora ludovicensis*, *Exophiala* (*Wangiella*) *dermatitidis*, *Exophiala jeanselmei*, *Exophiala spinifera*, *Fonsecaea pugnacius*, *Phaeosclera dermatioides*, *Rhinocladiella phaeophora*, *Rhinocladiella similis*, *Rhinocladiella tropicalis*, and *Rhytidhysterion* species have been published.^{1,2,13-19}

EPIDEMIOLOGY

Chromoblastomycosis has been described to occur throughout the world, although most cases arise in tropical and subtropical regions, especially those with high annual rainfall. Large numbers of cases have been described from Madagascar, Brazil, Mexico, Venezuela, and Costa Rica. Disease is more prevalent in males (4:1 ratio), in those aged 40 to 69,³ in association with outdoor activities such as farming and woodcutting, and in the absence of footwear. In Madagascar, a unique epidemiology has been described in what is probably the largest focus of endemic disease.²⁰ Madagascar has two distinct foci of infection, with disease secondary to *F. pedrosoi* occurring in the humid, rainy, northern evergreen forest region and disease secondary to *C. carrionii* found in the arid southern desert region. In a study of 1343 cases of disease over 40 years in that country, the prevalence of 1 case per 1920 inhabitants in the southern desert region has been described, with an incredible 1 in 910 prevalence in a single district of that region.

PATHOLOGY AND PATHOGENESIS

Traumatic inoculation of these fungi results in a mixed chronic suppurative and granulomatous host response.²¹ The epidermis typically becomes thickened in a process called pseudoepitheliomatous hyperplasia,



FIG. 260.1 Sclerotic bodies of chromoblastomycosis. (From Beneke ES, Rogers AL. *Medical Mycology and Human Mycoses*. Belmont, CA: Star Publishing; 1996.)

a histologic morphologic appearance that may be misidentified as malignancy by more inexperienced microscopists. Foci of polymorphonuclear cells and microabscesses are seen in both the epidermis and dermis. In the dermis, granulomas that include multinucleated giant cells and epithelioid cells are present, along with varying amounts of fibrosis. Fibrosis is increased in older lesions and can extend into the subcutaneous tissue, although disease rarely extends deep into the subcutaneous tissue. The hallmark of chromoblastomycosis, the muriform cells—also called sclerotic, copper penny, or Medlar bodies—may be found intracellularly in macrophages or extracellularly in abscesses. These are darkly pigmented (brown-golden), thick-walled, rounded cells, 4 to 12 μm in diameter, with cross walls in one or two planes (Fig. 260.1). Hyphae may also occasionally be seen, usually in the epidermis. The host response to these structures results in a process termed *transepithelial elimination*, in which fungi and damaged tissue are expelled through the epidermis, a process similar to that seen in calcinosis cutis.²² The immunologic response of the host in this mycosis is still not well understood. Disease appears to be associated with an ineffective immune response to the organism, with chronic inflammation produced in response to persistence of the fungi in tissue. The cell-mediated immune response appears to play a central role. High interleukin-10 (IL-10) levels, low interferon- γ levels, and inefficient T-cell proliferation have been noted in patients with severe disease, with the opposite seen in those with milder chromoblastomycosis.²³ This ineffective response has been linked to lack of recognition of these fungi by Toll-like receptors (TLRs) in a murine model.²⁴ Antibody responses have shown association with disease chronicity and extent but do not appear to provide any degree of protection in this infection.²⁵

CLINICAL MANIFESTATIONS

Weeks to months after inoculation of the causative organisms through minor trauma, subjects typically develop a small scaly papule on the lower extremity at the site of the trauma (Figs. 260.2–260.4). This lesion slowly develops into a superficial nodule, commonly with an irregular friable surface. Frequently, these nodules later spread out to become purplish, irregular, raised plaques. In descriptions by Carrion,²⁶ lesions of chromoblastomycosis were categorized into five types: (1) early nodular lesions, described as soft and pink-violaceous, with smooth, verrucous, or scaly surfaces; (2) tumorous lesions—large, papillomatous, often lobulated masses with crusting, sometimes described as cauliflower-like; (3) verrucous lesions with prominent hyperkeratosis; (4) plaque lesions; and (5) cicatricial lesions. Most lesions have black dots associated with their outer surface that are composed of fungi and necrotic debris, the products of transepithelial elimination. Although not typically painful, lesions may be associated with pruritus, are easily traumatized, and bleed readily. Ulceration is generally limited to those lesions with bacterial superinfection. Large lesions may become hyperkeratotic, and limb distortion, including elephantiasis, can occur as a result of blockage of normal lymphatic drainage.



FIG. 260.2 Chromoblastomycosis of the foot. (From Beneke ES, Rogers AL. *Medical Mycology and Human Mycoses*. Belmont, CA: Star Publishing; 1996.)



FIG. 260.3 Chromoblastomycosis with multiple verrucous nodules. (From McGinnis MR, Chandler FW. *Chromoblastomycosis*. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. *Pathology of Infectious Diseases*. Norwalk, CT: Appleton & Lange; 1997.)



FIG. 260.4 Chromoblastomycosis of the lower leg with lobulated, confluent nodules with focal ulcerations. (From McGinnis MR, Chandler FW. *Chromoblastomycosis*. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. *Pathology of Infectious Diseases*. Norwalk, CT: Appleton & Lange; 1997.)

Although lesions have been described to occur chiefly on the lower extremities (80%–85%) in most regions of the world,^{3,27} an exception to this pattern is reported from Japan. Evaluation of 290 lesions from that country found that chromoblastomycosis occurs most commonly on the upper extremities of male subjects and on the face or neck of females.⁷

Persistence of the lesions of chromoblastomycosis for 30 years has been reported, and delays in diagnosis of 1 to 3 years are not unusual. Although most lesions remain localized without spread to deeper structures, localized dissemination may occur via autoinoculation or via the lymphatics. Hematogenous spread has only rarely been described but does include reports of dissemination to the central nervous system.⁷ The occurrence of secondary bacterial infection has been reported to affect as many as 63% of persons with chromoblastomycosis.⁴ Although apparently rare, at least 14 cases of squamous cell carcinomas and 1 case of melanoma have been reported; all occurred after 8 or more years of disease.^{28,29}

The differential diagnosis of chromoblastomycosis includes psoriasis, other mycoses (e.g., blastomycosis, coccidioidomycosis, lobomycosis, mycetoma, paracoccidioidomycosis, cutaneous phaeohyphomycosis, sporotrichosis, tinea), cutaneous tuberculosis, leprosy, leishmaniasis, protothecosis, keratoacanthoma, squamous cell carcinoma, and sarcoidosis.

DIAGNOSIS

Chromoblastomycosis should be suspected in persons with chronic scaly or friable lesions of the extremities, especially in rural tropical climates. Microscopic examination of skin scrapings can provide a rapid diagnosis of chromoblastomycosis because the characteristic muriform cells can be seen in potassium hydroxide preparations, especially those containing black dots (see Fig. 260.1). These unique structures may also be readily observed with standard staining of skin punch biopsy specimens with hematoxylin and eosin (Fig. 260.5). Although not absolutely necessary, culture can be performed to identify the specific cause of infection. Standard mycologic media (Sabouraud glucose agar), with and without cycloheximide, should be used and cultures incubated for at least 4 weeks. In culture, the fungal agents of chromoblastomycosis appear as dark molds. Under standard culture conditions, these fungi may be identified by the microscopic appearance of hyphae and reproductive structures. The muriform structures seen in tissue have been produced *in vitro* with low pH and the addition of propranolol, but this is not necessary for clinical diagnosis.³⁰ Exoantigen testing of the fungus has been developed to aid in identification, although this is not commonly used.³¹ Serologic and skin tests have also been developed, but their use in this rare disease is limited to specialized centers in more endemic regions of the world.

THERAPY

Although spontaneous resolution has been reported,³² this is only a rare occurrence, and most chromoblastomycosis is a chronic indolent infection. When caused by its most common agent, *F. pedrosoi*, it is difficult to eradicate, even with prolonged therapy. Multiple modalities




FIG. 260.5 Dermal abscess with quadruple cluster of organisms in a mixture of neutrophils, macrophages, eosinophils, and a giant cell (hematoxylin and eosin). (From McGinnis MR, Chandler FW. Chromoblastomycosis. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997.)

have been used to treat patients with chromoblastomycosis, including surgery, local (physical) treatments, and antifungal agents.³³ Unfortunately, no single reproducibly successful treatment strategy has been identified. Surgical removal of small lesions appears to be effective, as does local application of liquid nitrogen, topical heat, and photocoagulation. Local curettage or electrocautery has been reported sometimes to result in disease spread and is to be discouraged. Heat therapy (42°C–46°C) with pocket warmers and other devices providing prolonged daily warmth directly to the lesions has been described as effective with 2 to 12 months of treatment.³⁴ Cryotherapy with liquid nitrogen sprays or applied with soaked cotton swabs or balls has been successful in the cure of small early lesions and may be used effectively in combination with antifungal medications on larger lesions. Topical imiquimod has been used successfully in four cases of disease secondary to *F. pedrosoi*.³⁵ The use of topical ajoene (a garlic extract) and 5-fluorouracil has been reported to be effective against disease secondary to *C. carrionii*.³⁶

Currently, the best therapy appears to be itraconazole or terbinafine, perhaps with adjunctive cryotherapy with liquid nitrogen or other local treatments.^{37,38} Other antifungal agents, including amphotericin B (intravenous or intralesional), 5-fluorocytosine, ketoconazole, and fluconazole, have been used with poor to mixed success, alone or in combination.

Itraconazole has been reported to be effective in many patients in uncontrolled, nonrandomized studies. Early study with lower doses (100 to 200 mg daily) of itraconazole documented high response rates with this azole antifungal agent, but the numbers of cures were small (3 of 10 patients treated for 12 to 24 months).³⁹ Treatment of chromoblastomycosis caused by *C. carrionii* with itraconazole has met with much greater success than treatment of that caused by other agents. One study reported cure in 2 of 5 patients with disease secondary to *F. pedrosoi* and 8 of 9 patients with *C. carrionii*, all given 100 to 400 mg of the drug daily for 4 to 8 months.⁴⁰ Queiroz-Telles and colleagues have reported the cure of 42% of patients (8 of 19) treated with 200 to 400 mg of itraconazole for a median of 7 months.⁴¹ Another group described the therapy for 10 patients with disease secondary to *F. pedrosoi*, 4 of whom had experienced failure of prior therapy with ketoconazole.⁴² All patients received 200 to 400 mg of itraconazole daily, and 8 also received monthly cryotherapy with liquid nitrogen. Nine patients were cured with 3 to 12 months of therapy; 2 of these responded with sustained cures after only 3 months of itraconazole at the lower dose of 200 mg daily. The remaining patient had marked improvement without cure. Recurrence was noted in a single patient who had been cured with a 6-month course of itraconazole (400 mg daily) and cryotherapy. Decreased *in vitro* susceptibility to itraconazole has been reported in one study of sequential clinical isolates of *F. pedrosoi*, potentially accounting for treatment failures.⁴³ Bonifaz and associates described good success with smaller lesions treated with itraconazole or cryosurgery and in larger lesions treated with itraconazole followed by cryosurgery.⁴⁴ In that study, which included 4 patients in each group and dosage of itraconazole at 100 mg three times daily for 5 to 14 months, 8 of 12 patients (67%) were cured and the remaining 4 showed improvement. In addition to daily therapy, success with itraconazole given as pulse therapy, 400 mg daily for 7 days/month for 6 to 12 months, has also been reported.⁴⁵ If the better-absorbed SUBA-itraconazole formulation is used, 65-mg capsules would be equivalent to 100-mg capsules of the older formulation.

The allylamine antifungal agent terbinafine has produced excellent results in the treatment of chromoblastomycosis. In the largest study to date, terbinafine, 500 mg daily, was given to 35 patients for up to 12 months.⁴⁶ In that study, 16 patients had experienced failure of thiabendazole in the past and almost half had lesions of longer than 10 years' duration. Improvement, defined as lack of bacterial superinfection and resolution of edema, was seen after 2 to 4 months of therapy and, after 12 months, 86% obtained mycologic cures (72% with clinical cures).²⁷ Patients with *C. carrionii* infections were noted to respond more quickly than those with *F. pedrosoi* infection in that study. Unexpectedly, partial reversal of fibrosis of the lesions of chromoblastomycosis has also been reported to occur with terbinafine therapy. This reversal has been suggested to be independent of mycologic cure of infection in those receiving terbinafine.^{47,48} Terbinafine used in alternate-week or

combination therapy with itraconazole to treat 4 patients with resistant infections successfully has also been reported.⁴⁹

The newer broad-spectrum azole antifungal agents may potentially be useful in this disease. In vitro testing has shown that the minimum inhibitory concentrations of voriconazole for *F. pedrosoi* and *F. compacta* are lower than those seen with itraconazole.⁵⁰ *F. pedrosoi* has also been shown to be susceptible to the echinocandin caspofungin in one small

in vitro study.⁵¹ Posaconazole (800 mg/day of the suspension in divided doses for 6 to 12 months) has been used to successfully treat a small number of patients (five of six) with disease refractory to itraconazole and terbinafine.⁵²

No vaccinations exist to prevent chromoblastomycosis. Proper protective clothing, especially footwear, and early treatment of the lesions are the only available preventive measures against this disease.

Key References

The complete reference list is available online at Expert Consult.

1. Kwon-Chung KJ, Bennett JE. Chromoblastomycosis. In: *Medical Mycology*. Philadelphia: Lea & Febiger; 1992:337–355.
2. Queiroz-Telles F, de Hoog S, Santos DW, et al. Chromoblastomycosis. *Clin Microbiol Rev*. 2017;30:233–276.
3. Minotto R, Bernardi CD, Mallmann LF, et al. Chromoblastomycosis: a review of 100 cases in the state of Rio Grande do Sul, Brazil. *J Am Acad Dermatol*. 2001;44:585–592.
4. Bonifaz A, Carrasco-Gerard E, Saul A. Chromoblastomycosis: clinical and mycologic experience of 51 cases. *Mycoses*. 2001;44:1–7.
5. Attapattu MC. Chromoblastomycosis—a clinical and mycological study of 71 cases from Sri Lanka. *Mycopathologia*. 1997;137:145–151.
6. Agarwal R, Singh G, Ghosh A, et al. Chromoblastomycosis in India: review of 169 cases. *PLoS Negl Trop Dis*. 2017;11:e0005534.
35. de Sousa M, Belda W Jr, Spina R, et al. Topical application of imiquimod as a treatment for chromoblastomycosis. *Clin Infect Dis*. 2014;58:1734–1737.
38. Queiroz-Telles F, Santos DW. Challenges in the therapy of chromoblastomycosis. *Mycopathologia*. 2013;175:477–488.
39. Restrepo A, Gonzalez A, Gomez I, et al. Treatment of chromoblastomycosis with itraconazole. *Ann N Y Acad Sci*. 1988;544:504–516.
40. Borelli D. A clinical trial of itraconazole in the treatment of deep mycoses and leishmaniasis. *Rev Infect Dis*. 1987;9(suppl 1):S57–S63.
41. Queiroz-Telles F, Purim KS, Filius JN, et al. Itraconazole in the treatment of chromoblastomycosis due to *Fonsecaea pedrosoi*. *Int J Dermatol*. 1992;31:805–812.
42. Kullavanijaya P, Rojanavanich V. Successful treatment of chromoblastomycosis due to *Fonsecaea pedrosoi* by the combination of itraconazole and cryotherapy. *Int J Dermatol*. 1995;34:804–807.
43. Andrade TS, Castro LGM, Nunes RS, et al. Susceptibility of sequential *Fonsecaea pedrosoi* isolates from chromoblastomycosis patients to antifungal agents. *Mycoses*. 2004;47:216–221.
44. Bonifaz A, Martinez-Soto E, Carrasco-Gerard E, et al. Treatment of chromoblastomycosis with itraconazole, cryosurgery, and a combination of both. *Int J Dermatol*. 1997;36:542–547.
45. Ungpakorn R, Reangchainam S. Pulse itraconazole 400 mg daily in the treatment of chromoblastomycosis. *Clin Exp Dermatol*. 2006;31:245–247.
46. Esterre P, Inzan CK, Ratsioharana M, et al. A multicentre trial of terbinafine in patients with chromoblastomycosis: effect on clinical and biologic criteria. *J Dermatol Treat*. 1998;9(suppl 1):S29–S34.
49. Gupta AK, Taborda PR, Sanzovo AD. Alternate week and combination itraconazole and terbinafine therapy for chromoblastomycosis caused by *Fonsecaea pedrosoi* in Brazil. *Med Mycol*. 2002;40:529–534.
50. Radford SA, Johnson EM, Warnock DW. In vitro studies of activity of voriconazole (UK-109,496), a new triazole antifungal agent, against emerging and less common mold pathogens. *Antimicrob Agents Chemother*. 1997;41:841–843.
51. Del Poeta M, Schell WA, Perfect JR. In vitro antifungal activity of pneumocandin L-743,872 against a variety of clinically important molds. *Antimicrob Agents Chemother*. 1997;41:1835–1836.
52. Negróni R, Tobón A, Bustamante B, et al. Posaconazole treatment of refractory eumycetoma and chromoblastomycosis. *Rev Inst Med Trop Sao Paulo*. 2005;47:339–346.

References

- Kwon-Chung KJ, Bennett JE. Chromoblastomycosis. In: *Medical Mycology*. Philadelphia: Lea & Febiger; 1992:337–355.
- Queiroz-Telles F, de Hoog S, Santos DW, et al. Chromoblastomycosis: a review of 100 cases in the state of Rio Grande do Sul, Brazil. *J Am Acad Dermatol*. 2001;44:585–592.
- Bonifaz A, Carrasco-Gerard E, Saul A. Chromoblastomycosis: clinical and mycologic experience of 51 cases. *Mycoses*. 2001;44:1–7.
- Attapattu MC. Chromoblastomycosis—a clinical and mycological study of 71 cases from Sri Lanka. *Mycopathologia*. 1997;137:145–151.
- Agarwal R, Singh G, Ghosh A, et al. Chromoblastomycosis in India: review of 169 cases. *PLoS Negl Trop Dis*. 2017;11:e0005534.
- Fukushiro R. Chromomycosis in Japan. *Int J Dermatol*. 1983;22:221–229.
- de Hoog GS, Attili-Angelis D, Vicente VA, et al. Molecular ecology and pathogenic potential of *Fonsecaea* species. *Med Mycol*. 2004;42:405–416.
- Najafzadeh MJ, Gueidan C, Badali H, et al. Genetic diversity and species delimitation in the opportunistic genus *Fonsecaea*. *Med Mycol*. 2009;47(special issue):17–25.
- Xi L, Sun J, Lu C, et al. Molecular diversity of *Fonsecaea* (Chaetothyriales) causing chromoblastomycosis in southern China. *Med Mycol*. 2009;47(special issue):27–33.
- Najafzadeh MJ, Sun J, Vicente V, et al. *Fonsecaea nubica* sp. nov., a new agent of human chromoblastomycosis revealed using molecular data. *Med Mycol*. 2010;48:800–806.
- Fransisca C, He Y, Chen Z, et al. Molecular identification of chromoblastomycosis clinical isolates in Guangdong. *Med Mycol*. 2017;55:851–858.
- Barba-Gomez JF, Mayorga J, McGinnis MR, et al. Chromoblastomycosis caused by *Exophiala spinifera*. *J Am Acad Dermatol*. 1992;26:367–370.
- Chowdhary A, Guarro J, Randhawa HS, et al. A rare case of chromoblastomycosis in a renal transplant recipient caused by a non-sporulating species of *Rhytidhysterion*. *Med Mycol*. 2008;46:163–166.
- Piepenbring M, Mendez OA, Espinoza AA, et al. Chromoblastomycosis caused by *Chaetomium funicola*: a case report from Western Panama. *Br J Dermatol*. 2007;157:1025–1029.
- Gomes RR, Vicente VA, de Azevedo CM, et al. Molecular epidemiology of agents of human chromoblastomycosis in Brazil with the description of two novel species. *PLoS Negl Trop Dis*. 2016;10:e0005102.
- de Azevedo CM, Gomes RR, Vicente VA, et al. *Fonsecaea pugnacius*, a novel agent of disseminated chromoblastomycosis. *J Clin Microbiol*. 2015;53:2674–2685.
- Kampirapap K, Reangchainam S, Ornpaew P, et al. Chromoblastomycosis masquerading as dermatophytosis, with the description of a new opportunistic species. *Southeast Asian J Trop Med Public Health*. 2015;46:105–109.
- Heidrich D, Gonzalez GM, Pagani DM, et al. Chromoblastomycosis caused by *Rhinochrysiella similis*: case report. *Med Mycol*. 2017;16:25–27.
- Esterre P, Andriantsimahavandy A, Ramarcel ER, et al. Forty years of chromoblastomycosis in Madagascar: a review. *Am J Trop Med Hyg*. 1996;55:45–47.
- Uribe F, Zuluaga AI, Leon W, et al. Histopathology of chromoblastomycosis. *Mycopathologia*. 1989;105:1–6.
- Batres E, Wolf JE Jr, Rudolph AH, et al. Transepithelial elimination of cutaneous chromomycosis. *Arch Dermatol*. 1978;114:1231–1232.
- Gimenes VMF, Sousa MG, et al. Cytokine and lymphocyte proliferation in patients with different clinical forms of chromoblastomycosis. *Microbes Infect*. 2005;7:708–713.
- Sousa M, Reid DM, Schweighoffer E, et al. Restoration of pattern recognition receptor costimulation to treat chromoblastomycosis, a chronic fungal infection of the skin. *Cell Host Microbe*. 2011;9:436–443.
- Esterre P, Jahevitra M, Andriantsimahavandy A. Humoral immune response in chromoblastomycosis during and after therapy. *Clin Diagn Lab Immunol*. 2000;7:497–500.
- Carrión AL. Chromoblastomycosis. *Ann N Y Acad Sci*. 1950;50:1255–1282.
- Silva JP, de Souza W, Rozental S. Chromoblastomycosis: a retrospective study of 325 cases in Amazonian Region (Brazil). *Mycopathologia*. 1999;143:171–175.
- Gon A, Minelli L. Melanoma in a long-standing lesion of chromoblastomycosis. *Int J Dermatol*. 2006;45:1331–1333.
- Rojas OC, González GM, Moreno-Treviño M, et al. Chromoblastomycosis by *Cladophialophora carrionii* associated with squamous cell carcinoma and review of published reports. *Mycopathologia*. 2015;179:153–157.
- da Silva JP, Alviano DS, Alviano CS, et al. Comparison of *Fonsecaea pedrosoi* sclerotic cells obtained in vivo and in vitro: ultra-structure and antigenicity. *FEMS Immunol Med Microbiol*. 2002;33:63–69.
- Espinell-Ingroff A, Shadomy S, Dixon D, et al. Exoantigen test for *Cladosporium bantianum*, *Fonsecaea pedrosoi*, and *Phialophora verrucosa*. *J Clin Microbiol*. 1986;23:305–310.
- Nishimoto K, Yoshimura S, Honma K. Chromomycosis spontaneously healed. *Int J Dermatol*. 1984;23:408–410.
- Queiroz-Telles F, Esterre P, Perez-Blanco M, et al. Chromoblastomycosis: an overview of clinical manifestations, diagnosis and treatment. *Med Mycol*. 2009;47(special issue):3–15.
- Tagami H, Ginoza M, Imaizumi S, et al. Successful treatment of chromoblastomycosis with topical heat therapy. *J Am Acad Dermatol*. 1984;10:615–619.
- de Sousa M, Belda W Jr, Spina R, et al. Topical application of imiquimod as a treatment for chromoblastomycosis. *Clin Infect Dis*. 2014;58:1734–1737.
- Perez-Blanco M, Valles RH, Zeppenfeldt GF, et al. Ajoene and 5-fluorouracil in the topical treatment of *Cladophialophora carrionii* chromoblastomycosis in humans: a comparative open study. *Med Mycol*. 2003;41:517–520.
- Queiroz-Telles F. Chromoblastomycosis: a neglected tropical disease. *Rev Inst Med Trop Sao Paulo*. 2015;19(suppl 57):46–50.
- Queiroz-Telles F, Santos DW. Challenges in the therapy of chromoblastomycosis. *Mycopathologia*. 2013;175:477–488.
- Restrepo A, Gonzalez A, Gomez I, et al. Treatment of chromoblastomycosis with itraconazole. *Ann N Y Acad Sci*. 1988;544:504–516.
- Borelli D. A clinical trial of itraconazole in the treatment of deep mycoses and leishmaniasis. *Rev Infect Dis*. 1987;9(suppl 1):S57–S63.
- Queiroz-Telles F, Purim KS, Filius JN, et al. Itraconazole in the treatment of chromoblastomycosis due to *Fonsecaea pedrosoi*. *Int J Dermatol*. 1992;31:805–812.
- Kullavanijaya P, Rojanavanich V. Successful treatment of chromoblastomycosis due to *Fonsecaea pedrosoi* by the combination of itraconazole and cryotherapy. *Int J Dermatol*. 1995;34:804–807.
- Andrade TS, Castro LGM, Nunes RS, et al. Susceptibility of sequential *Fonsecaea pedrosoi* isolates from chromoblastomycosis patients to antifungal agents. *Mycoses*. 2004;47:216–221.
- Bonifaz A, Martinez-Soto E, Carrasco-Gerard E, et al. Treatment of chromoblastomycosis with itraconazole, cryosurgery, and a combination of both. *Int J Dermatol*. 1997;36:542–547.
- Ungpakorn R, Reangchainam S. Pulse itraconazole 400 mg daily in the treatment of chromoblastomycosis. *Clin Exp Dermatol*. 2006;31:245–247.
- Esterre P, Inzan CK, Ratsioharana M, et al. A multicentre trial of terbinafine in patients with chromoblastomycosis: effect on clinical and biologic criteria. *J Dermatol Treat*. 1998;9(suppl 1):S29–S34.
- Esterre P, Risteli L, Ricard-Blum S. Immunohistochemical study of type I collagen turnover and of matrix metalloproteinases in chromoblastomycosis before and after treatment by terbinafine. *Pathol Res Pract*. 1998;194:847–853.
- Ricard-Blum S, Hartmann DJ, Esterre P. Monitoring of extracellular matrix metabolism and cross-linking in tissue, serum and urine of patients with chromoblastomycosis, a chronic skin fibrosis. *Eur J Clin Invest*. 1998;28:748–754.
- Gupta AK, Taborda PR, Sanzovo AD. Alternate week and combination itraconazole and terbinafine therapy for chromoblastomycosis caused by *Fonsecaea pedrosoi* in Brazil. *Med Mycol*. 2002;40:529–534.
- Radford SA, Johnson EM, Warnock DW. *In vitro* studies of activity of voriconazole (UK-109,496), a new triazole antifungal agent, against emerging and less common mold pathogens. *Antimicrob Agents Chemother*. 1997;41:841–843.
- Del Poeta M, Schell WA, Perfect JR. *In vitro* antifungal activity of pneumocandin L-743,872 against a variety of clinically important molds. *Antimicrob Agents Chemother*. 1997;41:1835–1836.
- Negroni R, Tobon A, Bustamante B, et al. Posaconazole treatment of refractory eumycetoma and chromoblastomycosis. *Rev Inst Med Trop Sao Paulo*. 2005;47:339–346.

SHORT VIEW SUMMARY

Definition

- Mycetoma is an infection of the skin and subcutaneous tissue characterized by a triad of localized swelling, draining sinuses, and grains (aggregates of infecting organisms). Unlike actinomycosis, which also forms grains in lesions, mycetoma enters the body through the skin.
- It most commonly affects a single site, typically involving the lower extremity and especially the foot.
- Eumycetoma (eumycotic mycetoma) is mycetoma caused by fungi, most commonly *Madurella mycetomatis*.
- Actinomycetoma (actinomycotic mycetoma) is mycetoma caused by bacteria, most commonly *Nocardia brasiliensis*.

Epidemiology

- Distribution is worldwide, with most cases occurring in tropical and subtropical regions.
- Largest numbers of cases are reported from Africa, Latin America, and the Indian subcontinent.
- Mycetoma occurs predominantly in men, ages 20 to 40, typically with occupations that expose them to the environment.

Microbiology

- *Madurella* spp., *M. mycetomatis*, *Trematosphaeria grisea*, and *Scedosporium apiospermum* complex are the most common causes of eumycetoma, although *Falciformispora senegalensis*, *Falciformispora tompkinsii*, *Exophiala jeanselmei*, and many other genera and species of fungi have been reported as etiologic agents.
- *Nocardia brasiliensis*, *Actinomadura madurae*, *Streptomyces somaliensis*, and *Actinomadura pelletieri* are the most common causes of actinomycetoma, although disease secondary to other species of *Actinomadura*, *Nocardia*, and *Streptomyces* has been described.

Diagnosis

- Clinical presentation of the classic triad of chronic, painless soft tissue swelling with draining sinuses that discharge grains is pathognomonic.
- Microscopic examination of the grains can differentiate between eumycetoma and actinomycetoma.
- Culture of causative agent from grains can better direct selection of antimicrobial therapy.
- Radiographic techniques (plain radiographs, ultrasound, computed tomography, and

magnetic resonance imaging) can be used adjunctively in making (or excluding) the diagnosis and determining its extent.

Therapy

- Small lesions may be treated successfully with surgical excision alone.
- Actinomycetoma is typically treated with medical therapy alone.
- Eumycetoma commonly requires combined medical and surgical therapy, but results are poor.
- No single therapy has proved most effective for either form of mycetoma.
- Most actinomycetoma regimens include parenteral aminoglycosides and oral sulfa drugs.
- Less severe actinomycetoma may be treated with 6 to 24 months of trimethoprim-sulfamethoxazole.
- Eumycetoma is typically treated with a regimen of an oral azole antifungal drug for 6 to 24 months, perhaps combined with debulking surgery.

Prevention

- No vaccine is available.
- Use of footwear and proper protective clothing should protect against this infection.

Mycetoma is a chronic progressive granulomatous infection of the skin and subcutaneous tissue most often affecting the lower extremities, typically a single foot. The disease is unique from other cutaneous or subcutaneous diseases in its triad of localized swelling, underlying sinus tracts, and production of grains or granules (composed of aggregations of the causative organism) within the sinus tracts. These infections may be caused by fungi and termed *eumycotic mycetoma* or *eumycetoma*, or by filamentous higher bacteria and termed *actinomycotic mycetoma* or *actinomycetoma*. The term *mycetoma* can also be found in the literature incorrectly referring to a fungus ball found in a preexisting cavity in the lung or within a paranasal sinus, most often caused by *Aspergillus* spp. Grain formation by infecting organisms is restricted to the diseases mycetoma, actinomycosis (see Chapter 254), and botryomycosis. Actinomycosis is a disease produced by the anaerobic and microaerophilic higher bacteria that normally colonize the mouth and gastrointestinal and urogenital tracts. The portal of entry in actinomycosis is from those colonized sites, whereas in mycetoma the portal is the skin and subcutaneous tissue into which the organism was inoculated by minor trauma. Botryomycosis is a chronic bacterial infection of soft tissues in which the causative organism, often *Staphylococcus aureus*, is found in loose clusters among the pus.¹ In a rare form of ringworm called *dermatophyte mycetoma*, there are also loosely compacted clusters of hyphae in subcutaneous pus.² In contrast, mycetoma grains are dense clusters of organisms.

ETIOLOGIC AGENTS

The agents of mycetoma are fungi and aerobic filamentous bacteria that have been found on plants and in the soil.^{3,4} The predominance of bacterial versus fungal causes of mycetoma varies among geographic locations. Eumycotic (true fungal) disease is caused by a variety of fungal organisms. These can be divided into those that form dark grains and those that form pale or white grains (Table 261.1). Color distinctions are made by observing unstained specimens. Among the fungi causing dark-grained mycetoma, the most common are *Madurella mycetomatis*, *Falciformispora* (formerly *Leptosphaeria*) *senegalensis*, and *Trematosphaeria grisea*. Other agents include *Corynespora cassicola*, *Curvularia geniculata*, *Curvularia lunata*, *Emarellia grisea*, *Emarellia paragrisea*, *Exophiala jeanselmei*, *Exophiala oligosperma*, *Falciformispora* (formerly *Leptosphaeria*) *tompkinsii*, *Madurella fahalii*, *Madurella pseudomycetomatis*, *Madurella tropicana*, *Phialophora verrucosa*, *Plenodomas avramii*, *Pseudochaetosphaeronema larense*, *Rhinocladiella atrovirens*, *Medicopsis* (formerly *Pyrenochaeta*) *mackinnonii*, *Biatrispora* spp., *Rousoella* spp., *Rhytidhysterion* spp., and *Medicopsis* (formerly *Pyrenochaeta*) *romeroi*. *Scedosporium apiospermum* complex species are the most common cause of pale-colored grains. Other fungi in that category include *Fusarium* (formerly *Acremonium*) *falciforme*, *Sarocladium* (formerly *Acremonium*) *kiliense*, *Acremonium recifei*, *Aspergillus flavus*, *Aspergillus hollandicus*, *Aspergillus nidulans*, *Phialophora* (formerly *Cylindrocarpon*) *cyaneus*, *Cylindrocarpon destructans*, *Diaporthe phaseolorum*, *Fusarium solani*,

TABLE 261.1 Typical Morphologic Features of Mycetoma Grains

GRAIN COLOR	CAUSATIVE AGENT
Eumycetoma (Eumycotic Mycetoma)^a	
Black grains	<i>Madurella</i> spp., <i>Biatrispora</i> spp., <i>Trematosphaeria</i> spp., <i>Pseudochaetosphaeroma</i> spp., <i>Roussoella</i> spp., <i>Rhytidhysterion</i> spp., <i>Curvularia</i> spp., <i>Exophiala</i> spp., <i>Falciformispora</i> spp., <i>Medicopsis</i> spp., <i>Phaeoacremonium</i> spp., <i>Phialophora verrucosa</i>
Pale grains (white to yellow)	<i>Scedosporium apiospermum</i> complex, <i>Aspergillus</i> spp., <i>Diaporthe phaseolorum</i> , <i>Fusarium</i> spp., <i>Neotestudina rosatii</i> , <i>Pleurostomophora ochracea</i>
Actinomycetoma (Actinomycotic Mycetoma)^b	
Pale grains (white to yellow)	<i>Actinomadura madurae</i> , <i>Nocardia</i> spp.
Yellow to brown grains	<i>Streptomyces somaliensis</i>
Red to pink grains	<i>Actinomadura pelletieri</i>

^aHyphae of 2- to 5-μm diameter are observed within grain.

^bFilaments of 0.5- to 1-μm diameter are observed within grain.

Fusarium moniliforme, *Fusarium keratoplasticum*, *Fusarium pseudensiforme*, *Neotestudina rosatii*, *Phaeoacremonium* spp., *Pleurostomophora ochracea*, and *Polycyrtella hominis*.⁵⁻¹³

Actinomycetoma is caused by members of the order Actinomycetales, most commonly *Nocardia brasiliensis*, *Actinomadura madurae*, *Streptomyces somaliensis*, and *Actinomadura pelletieri*. Cases have been reported that were caused by *Actinomadura latina*, *Nocardia aobensis*, *Nocardia farcinica*, *Nocardia harenae*, *Nocardia otitidiscaviarum* (formerly *N. caviae*), *Nocardia mexicana*, *Nocardia transvalensis*, *Nocardia veterana*, *Nocardia yamanashiensis*, and *Nocardiopsis dassonvillei*.^{4,14-18} Actinomycetoma grains are typically white or pale yellow, except those caused by *Actinomadura pelletieri*, which are red to pink.

Some reports use species names that are not currently recognized, leaving in doubt the identification.¹⁹ The modern use of molecular techniques, including polymerase chain reaction, DNA sequencing, and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF), has both assisted in better identifying and classifying the agents of mycetoma, and made comparisons between current and previous pathogen identification difficult to track.²⁰⁻²²

EPIDEMIOLOGY

The oldest description of this disease appears to date back to the ancient Indian Sanskrit text *Atharva Veda*, in which reference is made to *pada valmikam*, translated to mean “anthill foot.”²³ More modern descriptions from Madras, India, in the 19th century led to this disease initially being called “madura foot,” or *maduromycosis*, a term still used by some today to describe eumycotic mycetoma. Mycetoma is most commonly found in tropical and subtropical climates, with the highest incidence reported from endemic areas in the Indian subcontinent, the Middle East, Africa, and Central and South America. One of the largest current groups of cases is in Sudan. Only scattered reports describe cases originating in the United States, Europe, and Japan. Disease occurs around five times more frequently in males, commonly in the 20- to 40-year-old age range. Disease is more common in agricultural workers and outdoor laborers but is not exclusively seen in rural areas. Disease occurs sporadically throughout most areas of the world, and some postulate that the increased numbers in tropical regions may also result in part from less use of protective clothing, chiefly shoes, in the warmer, poorer endemic regions.

The causative agents of mycetoma vary from region to region and with climate. Worldwide, *M. mycetomatis* is the most common cause of this disease, but *A. madurae*, *M. mycetomatis*, and *S. somaliensis* are more commonly reported from drier regions, whereas *S. apiospermum* complex spp., *Nocardia* spp., and *A. pelletieri* are more common in those areas with higher annual rainfall. In India, *Nocardia* spp. and *T. grisea* are the most common causes of mycetoma; in the Middle East, *M. mycetomatis* and *S. somaliensis*; in West Africa, *F. senegalensis*; and



FIG. 261.1 Mycetoma of the foot. (From Beneke ES, Rogers AL. Medical Mycology and Human Mycoses. Belmont, CA: Star Publishing; 1996.)

in East Africa, *M. mycetomatis* and *S. somaliensis*. In Central and South America, *T. grisea* and *Nocardia* spp. are the common causes of mycetoma, and in the United States, *S. apiospermum* complex spp. are the most commonly recovered causative agent.²³

PATHOLOGY AND PATHOGENESIS

Infection follows inoculation of organisms, frequently through horn punctures, wood splinters, or preexisting abrasions or trauma. After inoculation, these normally nonpathogenic organisms grow and survive through the production of grains (also called granules or sclerotia), structures composed of masses of mycelial fungi or bacterial filaments and a matrix component. The matrix material has been shown to be host derived with some pathogens. In eumycetoma, hyphal elements often have thickened cell walls toward the periphery of grains, potentially conferring protection against the host immune system.²⁴ Grains are seen in histopathology within abscesses containing polymorphonuclear cells. Complement-dependent chemotaxis of polymorphonuclear leukocytes has been shown to be induced by both fungal (*M. mycetomatis* and *S. apiospermum* complex) and actinomycotic (*S. somaliensis*) antigens in vitro.²⁵ Cells of the innate immune system attempt to engulf and inactivate these organisms, but in disease they ultimately fail to accomplish this goal. Abscesses containing grains are seen in association with granulomatous inflammation and fibrosis. *Nocardia brasiliensis* has been shown to be resistant to human neutrophil peptides.²⁶ Three types of immune responses have been described in response to the grains of mycetoma.²⁷ The type 1 response is seen as neutrophils degranulate and adhere to the grain surface, leading to gradual disintegration of the grain. Type 2 response is characterized by the disappearance of neutrophils and arrival of macrophages to clear grains and neutrophil debris. Type 3 response is marked by the formation of epithelioid granuloma. This host response does not appear to be able to control infection but likely accounts for the partial spontaneous healing that is seen in the disease.

It is not clear whether persons who develop mycetoma have predisposing immune deficits.⁴ Disease does not appear to be more common in immunocompromised hosts, and early studies of immune function in persons with mycetoma have not clearly documented a common deficit.^{28,29} Recent work examining genes responsible for innate immune functions has identified polymorphisms that appear to predispose people to this infection, which may be linked with neutrophil function.³⁰ It has been suggested that the greater frequency of disease in men is not completely explained by increased frequency of exposure to soil and plant material. Progesterone has been shown in vitro to inhibit the growth of *M. mycetomatis*, *M. romeroi*, and *N. brasiliensis*.^{31,32} In the study of *N. brasiliensis*, estradiol limited disease produced in animals.³¹

CLINICAL MANIFESTATIONS

More than 75% of persons with mycetoma have a lesion of a lower extremity, most commonly the foot (70%) (Figs. 261.1 and 261.2). Next



FIG. 261.2 Mycetoma of the leg (seen from back of knee). (Courtesy Dr. Glenn W. Wortmann.)



FIG. 261.3 Mycetoma of the arm caused by *Madurella mycetomatis*. (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)

in frequency is disease of the hand (15%), followed by the upper extremities and other areas of the body that may be exposed by carrying firewood or thorny brush, including the upper back and adjacent neck, top of the head, and, rarely, the face (Fig. 261.3). Lesions in more than one anatomic site are extraordinarily rare. Disease begins in most cases as a single, small, painless subcutaneous nodule. This nodule slowly increases in size, becomes fixed to the underlying tissue, and ultimately develops sinus tracts beneath the lesion. These tracts open to the surface and drain purulent material with grains. Grains are several millimeters in diameter and may be seen by close inspection of a gauze bandage covering the sinus tract. Progression to draining sinus tracts can take weeks, months, and even years, occurring more rapidly in actinomycetoma. In a study of patients in India, the average time to presentation with disease from history of probable inciting trauma was 3 years for *N. brasiliensis*, 7 years for *A. madurae*, and 9 years for *T. grisea*.³³

Disease can affect the skin, subcutaneous tissue, and eventually contiguous bone, spreading along fascial planes. Overlying skin appears smooth and shiny and is commonly fixed to the underlying tissue. Skin may be hypopigmented or hyperpigmented, with signs of both old healed and active sinuses, displaying the cycle of spontaneous healing of older sinus tracts and simultaneous spread of infection to new areas that is typical of this disease. Swelling is often firm and nontender, and the overlying skin is not erythematous. Muscle, tendons, and nerves are generally spared direct infection, but extensive local damage may lead to muscle wasting, bone destruction, and limb deformities. Lymphatic spread is rare, although it may follow surgical manipulation. Hematogenous




FIG. 261.4 Eumycetoma grain of *Fusarium (Acremonium) falciforme*. (Gomori methenamine silver and hematoxylin and eosin stains.) (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)




FIG. 261.5 Eumycetoma grain of *Scedosporium apiospermum-Pseudallescheria boydii* complex. (Hematoxylin and eosin stain.) (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)

spread has not been documented. This disease and its effects are generally localized, and thus no signs or symptoms of systemic illness are usually seen in mycetoma unless secondary bacterial infection occurs. When left untreated, disease continues to progress, and bacterial superinfection can lead to increased morbidity from local abscess formation, cellulitis, bacterial osteomyelitis, and, rarely, septic death.

Differential diagnosis includes botryomycosis, chronic bacterial osteomyelitis, tuberculous osteomyelitis, chromoblastomycosis, phaeohyphomycosis, and soft tissue or bone tumor.

DIAGNOSIS

A diagnosis of mycetoma can be made by the classic triad of painless soft tissue swelling, draining sinus tracts, and extrusion of grains. Diagnosis of the causative organism can be made by microscopic observation and culture of a grain. Deep biopsy with histopathology and culture is usually not necessary, although obtaining a deep tissue biopsy avoids the bacterial contamination of surface cultures. Grains may not be seen in any one histopathologic section because they are scattered along the tracts. When a grain is present in the section, its large size and surrounding cluster of neutrophils make it difficult to miss, even without fungal or bacterial stains (Figs. 261.4 through 261.9). Organisms are usually not seen outside the grain. An alternate diagnostic strategy is the ultrasound-guided needle aspiration of grains directly




FIG. 261.6 Eumycetoma grain of *Curvularia geniculata*. (Hematoxylin and eosin stain.) (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)




FIG. 261.7 Eumycetoma grain of *Neotestudina rosatii*. (Gomori methenamine silver and hematoxylin and eosin stains.) (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)




FIG. 261.8 Actinomycetoma grain. (Gridley stain.) (From Beneke ES, Rogers AL. Medical Mycology and Human Mycoses. Belmont, CA: Star Publishing; 1996.)

from an unopened sinus tract for microscopic observation and culture.³⁴ Evaluation of spontaneously extruded grains may not allow diagnosis because these grains are often composed of dead organisms and contaminated with skin surface bacteria that outgrow the mycetomatous agent in culture.




FIG. 261.9 *Nocardia brasiliensis* grain. (Hematoxylin and eosin stain.) (From Chandler FW, Ajello L. Mycetoma. In: Connor DH, Chandler FW, Schwartz DA, et al, eds. Pathology of Infectious Diseases. Norwalk, CT: Appleton & Lange; 1997:1035–1044.)

The grains (or granules or sclerotia) of mycetoma are usually 0.2 to 5 mm in diameter and thus may be observed grossly, without magnification. Microscopic evaluation of crushed grains prepared with potassium hydroxide or stained with Gram stain is useful in differentiating fungal from bacterial causes. On inspection, actinomycetes are recognized by the production of 0.5- to 1- μ m-wide filaments and fungi by 2- to 5- μ m-wide hyphae. Many reports and reviews have detailed the use of grain color, size, and consistency to diagnose the specific cause of mycetoma, but recovery of the causative agents in culture is more accurate and of greater clinical usefulness when resources are available.

Culture of grains recovered from aspirated material or biopsy specimens can be used to diagnose the specific cause of mycetoma. If extruded grains are used, most experts suggest rinsing these in 70% alcohol or with antibiotic-containing saline solutions to decrease bacterial contamination. Specimens should be cultured on mycologic and mycobacteriologic media and held for at least 4 weeks.

The role of radiology in the management of mycetoma is that of adjunctive assessment of disease extent and involvement of bone, and perhaps long-term follow-up of disease regression or progression. Radiographic studies can help define the extent of disease and aid in the differentiation of mycetoma from other disease. Standard radiographic studies can reveal bony involvement such as periosteal erosion secondary to invasion, osteoporosis, and changes consistent with osteomyelitis, including lytic lesions. Ultrasonography has been used successfully in the differentiation of mycetoma from osteomyelitis or tumor. In a study of 100 patients with foot swelling who underwent ultrasonography before surgical excision, these lesions were found to have characteristics that distinguished them from other diseases.³⁵ Eumycetoma were found to produce single or multiple thick-walled cavities, without acoustic enhancement, with grains represented as distinct hyperreflective echoes. Actinomycetoma produced similar results, except grains produced fine echoes that were found at the bottom of the cavities. Magnetic resonance imaging (MRI) and computed tomography have also been evaluated in the management of mycetoma. Both modalities provide accurate assessment of disease extent when compared with surgical findings, especially in the soft tissues.³⁶ When compared directly, computed tomography appears to be more sensitive for detecting early changes consistent with bone involvement. A *dot-in-circle sign* has been described as a potentially specific diagnostic finding seen with MRI.^{37,38} The dots are tiny hypointense foci (believed to be grains) within spherical, high-intensity lesions (the circle) surrounded by low-intensity matrix on T2-weighted imaging, which represent granulomas scattered in areas of fibrosis. T1-weighted, fat-saturated, postgadolinium images may also produce this appearance. An MRI grading system has been developed by the Mycetoma Research Centre (Khartoum, Sudan) for use in the diagnosis and management of mycetoma.³⁹

The use of serology in the diagnosis and long-term management of this disease has been advocated by some authorities. Of the tests

described, counterimmunoelectrophoresis has been the most commonly used. Lack of standardization or widespread availability limits the use of these tests to centers that see a large volume of such patients. In the United States, the infrequency of the diagnosis and the diverse number of pathogens render serology of no practical use.

THERAPY

Treatment of mycetoma has proved to be difficult and typically includes prolonged courses of antimicrobial agents, often with surgical debulking.^{40–42} Surgery alone has a limited role in the treatment of mycetoma.⁴³ In addition to limb amputation, surgical excision of smaller lesions may be successful as monotherapy. Typically, surgery is employed adjunctively in fungal mycetoma to debulk large lesions after weeks to months of azole antifungal therapy has been given. Because chemotherapy varies for actinomycetoma and eumycetoma, at a minimum the clinician must differentiate whether a mycetoma is caused by actinomycetes or fungi. Ideally, recovery of the causative organism can allow identification of species, and perhaps even susceptibility testing, to guide therapy. Treatment regimens are currently based on expert opinion because no randomized controlled trials have been performed. Duration of therapy is also not defined, and most patients require 3 to 24 months of therapy to obtain an adequate response.

The most commonly described regimens for actinomycetoma include parenteral aminoglycosides combined with oral sulfonamide drugs or dapsone. Therapy with 5-week cycles of oral trimethoprim-sulfamethoxazole (TMP-SMX) (2 double-strength tablets twice daily) with parenteral amikacin (15 mg/kg/day divided into 2 daily doses) given during the first 3 weeks has been reported to produce successful results in all treated patients.^{4,14,44} More than half of 56 patients treated with this regimen were cured with one or two cycles; none required more than four cycles, and only one relapsed. The single patient with relapse was treated successfully with three cycles of TMP-SMX combined with netilmicin, an aminoglycoside. Other aminoglycoside-sulfonamide regimens include gentamicin or streptomycin in combination with either TMP-SMX or dapsone.^{40,41,44,45} TMP-SMX alone may be successful in the treatment of smaller lesions. Other regimens that have been used include streptomycin with either sulfadoxine-pyrimethamine

or rifampin; a combination of penicillin, gentamicin, and TMP-SMX followed by TMP-SMX and amoxicillin; and regimens that include amoxicillin-clavulanate, tetracycline, rifampin, fusidic acid, clindamycin, or imipenem-cilastatin.^a Success with linezolid therapy has also been reported in one patient.^{4,14}

Antifungal therapy for eumycetoma most commonly includes the use of azole antifungals because amphotericin B has not been effective in producing long-term cures. The first-line agent in the treatment of this disease is considered to be itraconazole (200–400 mg/day). Cure rates with this drug are quite variable, and it is suggested that all patients be evaluated for surgical debulking after their disease is controlled with azole therapy, typically after a year of therapy.^{41,48,49} In vitro, all three of the newer azoles—isavuconazole,⁵⁰ posaconazole, and voriconazole—have good activity against many of the causative agents of eumycetoma. Case reports of successful therapy with voriconazole have been published,^{51–53} as has a small case series of successful therapy for previously azole-refractory disease that responded to posaconazole.⁵⁴ Because of its susceptibility pattern, mycetoma secondary to the *S. apiospermum* complex should be treated with voriconazole. Successful therapy with terbinafine, an allylamine antifungal, has also been reported. Improvement or cure was seen in 16 of 20 patients who completed 24 to 48 weeks of terbinafine therapy (500 mg twice daily).⁵⁵ There are very little safety data on doses this high. In vitro, terbinafine has limited activity against *M. mycetomatis* compared with that observed with itraconazole, ketoconazole, or posaconazole.⁵⁶

PREVENTION

No preventive vaccine is available against any of the causative agents of mycetoma. Disease prevention is best accomplished by reduction of the incidence of the traumatic inoculation of the causative organisms. Wearing shoes and clothing to protect against splinters and thorn pricks should be stressed. Debilitating disease can be prevented by early identification and treatment of lesions, usually with minor surgery and chemotherapy.

^aReferences 4, 14, 40, 41, 46, 47.

Key References

The complete reference list is available online at Expert Consult.

4. Nenoff P, van de Sande WWJ, Fahal AH, et al. Eumycetoma and actinomycetoma – an update on causative agents, epidemiology, pathogenesis, diagnostics and therapy. *J Eur Acad Dermatol Venereol*. 2015;29:1873–1883.
6. Queiroz-Telles F, Wagner D, Pedrozo C. Fungal infections of implantation (chromoblastomycosis, mycetoma, entomophthoromycosis and lacaziosis). In: Hospenthal DR, Rinaldi MG, eds. *Diagnosis and Treatment of Fungal Infections*. 2nd ed. Switzerland: Springer International Publishing; 2015:261–276.
14. Welsh O, Vera-Cabrera L, Welsh E, et al. Actinomycetoma and advances in its treatment. *Clin Dermatol*. 2012;30:372–381.
40. Welsh O, Al-Abdely HM, Salinas-Carmona MC, et al. Mycetoma medical therapy. *PLoS Negl Trop Dis*. 2014;8:e3218.
41. Ameen M, Arenas R. Developments in the management of mycetomas. *Clin Exp Dermatol*. 2008;34:1–7.
42. Zijlstra EE, van de Sande WW, Welsh O, et al. Mycetoma: a unique neglected tropical disease. *Lancet Infect Dis*. 2016;16:100–112.
43. Suleiman SH, Wadaella ELS, Fahal AH. The surgical treatment of mycetoma. *PLoS Negl Trop Dis*. 2016;10:e0004690.
44. Raman M, Bhat R, Garg T, et al. A modified two-step treatment for actinomycetoma. *Indian J Dermatol Venereol Leprol*. 2007;73:235–239.
45. Fahal AH. Evidence-based guidelines for the management of mycetoma patients. Mycetoma Research Centre, Khartoum, Sudan, October 2002. www.mycetoma.edu.sd.

References

- Padilla-Desgarnes C, Vazquez-Gonzalez D, Bonifaz A. Botryomycosis. *Clin Dermatol*. 2012;30:397–402.
- Berg JC, Hamacher KL, Roberts GD. Pseudomycetoma caused by *Microsporium canis* in an immunosuppressed patient: a case report and review of the literature. *J Cutan Pathol*. 2007;34:431–434.
- Ahmed A, Adelmann D, Fahal A, et al. Environmental occurrence of *Madurella mycetomatis*, the major agent of human eumycetoma in Sudan. *J Clin Microbiol*. 2002;40:1031–1036.
- Nenoff P, van de Sande WWJ, Fahal AH, et al. Eumycetoma and actinomycetoma – an update on causative agents, epidemiology, pathogenesis, diagnostics and therapy. *J Eur Acad Dermatol Venereol*. 2015;29:1873–1883.
- Kwon-Chung KJ, Bennett JE. Mycetoma. In: *Medical Mycology*. Philadelphia: Lea & Febiger; 1992:560–593.
- Queiroz-Telles F, Wagner D, Pedrozo C. Fungal infections of implantation (chromoblastomycosis, mycetoma, entomophthoromycosis and lacaziosis). In: Hospenthal DR, Rinaldi MG, eds. *Diagnosis and Treatment of Fungal Infections*. 2nd ed. Switzerland: Springer International Publishing; 2015:261–276.
- Desnos-Ollivier M, Bretagne S, Dromer F, et al. Molecular identification of black-grain mycetoma agents. *J Clin Microbiol*. 2006;44:3517–3523.
- Iriart X, Binois R, Flor A, et al. Eumycetoma caused by *Diaporthe phaseolorum* (*Phomopsis phaseoli*): a case report and a mini-review of *Diaporthe/Phomopsis* spp. invasive infections in humans. *Clin Microbiol Infect*. 2011;17:1492–1494.
- de Hoog GS, van Diepeningen AD, Mahgoub E, et al. New species of *Madurella*, causative agents of black-grain mycetoma. *J Clin Microbiol*. 2012;50:988–994.
- Hemashettar BM, Siddaramappa B, Munjunathaswamy BS, et al. *Phaeoacremonium kraidenii*, a cause of white grain eumycetoma. *J Clin Microbiol*. 2006;44:4619–4622.
- Mhmoud NA, Ahmed SA, Fahal AH, et al. *Pleurostomophora ochracea*, a novel agent of human eumycetoma with yellow grains. *J Clin Microbiol*. 2012;50:2987–2994.
- Borman AM, Desnos-Ollivier M, Campbell CK, et al. Novel taxa associated with human fungal black-grain mycetomas: *Eumarellia grisea* gen. nov., sp. nov., and *Eumarellia paragrisea* sp. nov. *J Clin Microbiol*. 2016;54:1738–1745.
- Al-Hatmi AMS, Bonifaz A, Tirado-Sanchez A, et al. *Fusarium* species causing eumycetoma: report of two cases and comprehensive review of the literature. *Mycoses*. 2017;60:204–212.
- Welsh O, Vera-Cabrera L, Welsh E, et al. Actinomycetoma and advances in its treatment. *Clin Dermatol*. 2012;30:372–381.
- Kresch-Tronik NS, Carrillo-Casas EM, Arenas R, et al. *Nocardia harenae*, an uncommon causative organism of mycetoma: report on two patients. *J Med Microbiol*. 2012;61:1153–1155.
- Rodriguez-Nava V, Couble A, Molinard C, et al. *Nocardia mexicana* sp. nov., a new pathogen isolated from human mycetomas. *J Clin Microbiol*. 2004;42:4530–4535.
- Mitja O, Hays R, Van Straten C, et al. Case report: mycetoma caused by *Nocardia yamanashiensis*, Papua New Guinea. *Am J Trop Med Hyg*. 2012;86:1043–1045.
- Vongphoumy I, Dance DAB, Ditttrich S, et al. Case report: actinomycetoma caused by *Nocardia aobensis* from Lao PDR with favourable outcome after short-term antibiotic treatment. *PLoS Negl Trop Dis*. 2015;9:e0003729.
- Quintana ET, Wierzbicka K, Mackiewicz P, et al. *Streptomyces sudanensis* sp. nov., a new pathogen isolated from patients with actinomycetoma. *Antonie Van Leeuwenhoek*. 2008;93:305–313.
- Fraser M, Borman AM, Johnson EM. Rapid and robust identification of the agents of black-grain mycetoma by matrix-assisted laser desorption/ionization-time of flight mass spectrometry. *J Clin Microbiol*. 2017;55:2521–2528.
- Ahmed SA, van de Sande WWJ, Stevens DA, et al. Revision of agents of black-grain eumycetoma in the order Pleosporales. *Persoonia*. 2014;33:141–154.
- van de Sande WWJ. Phylogenetic analysis of the complete mitochondrial genome of *Madurella mycetomatis* confirms its taxonomic position within the order Sordariales. *PLoS ONE*. 2012;7:e38654.
- Green WO, Adams TE. Mycetoma in the United States: a review and report of seven additional cases. *Am J Clin Pathol*. 1964;42:75–91.
- Wethered DB, Markey MA, Hay RJ, et al. Ultrastructural and immunogenic changes in the formation of mycetoma grains. *J Med Vet Mycol*. 1986;25:39–46.
- Yousif MA, Hay RJ. Leucocyte chemotaxis to mycetoma agents—the effect of the antifungal drugs griseofulvin and ketoconazole. *Trans R Soc Trop Med Hyg*. 1987;81:319–321.
- Vera-Cabrera L, Salinas-Carmona MC, Waksman N, et al. Host defenses in subcutaneous mycosis. *Clin Dermatol*. 2012;30:382–388.
- Fahal AH, el Toum EA, el Hassan AM, et al. The host tissue reaction to *Madurella mycetomatis*: new classification. *J Med Vet Mycol*. 1995;33:15–17.
- Bendl BJ, Mackey D, Al-Saati F, et al. Mycetoma in Saudi Arabia. *J Trop Med Hyg*. 1987;90:51–59.
- Mahgoub ES, Gumaa SA, El Hassan AM. Immunological status of mycetoma patients. *Bull Soc Pathol Exot Filiales*. 1977;70:48–54.
- van de Sande WWJ, Fahal A, Verbrugh H, et al. Polymorphisms in genes involved in innate immunity predispose toward mycetoma susceptibility. *J Immunol*. 2007;179:3065–3074.
- Hernández-Hernández F, López-Martínez R, Méndez-Tovar LJ, et al. *Nocardia brasiliensis*: in vitro and in vivo growth in response to steroid sex hormones. *Mycopathologia*. 1995;132:79–85.
- Méndez-Tovar LJ, de Biève C, López-Martínez R. Effets des hormones sexuelles humaines sur le développement in vitro des agents d'eumycetomes. *J Mycol Méd*. 1991;1:141–143.
- Maiti PK, Ray A, Bandyopadhyay S. Epidemiological aspects of mycetoma from a retrospective study of 264 cases in West Bengal. *Trop Med Int Health*. 2002;7:788–792.
- Yousif BM, Fahal AH, Shakir MY. A new technique for the diagnosis of mycetoma using fixed blocks of aspirated material. *Trans R Soc Trop Med Hyg*. 2010;104:6–9.
- Fahal AH, Sheik HE, Homeida MM, et al. Ultrasonographic imaging of mycetoma. *Br J Surg*. 1997;84:1120–1122.
- Sharif HS, Clark DC, Aabed MY, et al. Mycetoma: comparison of MR imaging with CT. *Radiology*. 1991;178:865–870.
- Sarris I, Berendt AR, Athanasous N, et al. MRI of mycetoma of the foot: two cases demonstrating the dot-in-circle sign. *Skeletal Radiol*. 2003;32:179–183.
- Laohawiriyakamol T, Tanutit P, Kanjanapradit K, et al. The "dot-in-circle" sign in musculoskeletal mycetoma on the magnetic resonance imaging and ultrasound. *Springerplus*. 2014;3:671.
- El Shamy ME, Fahal AN, Shakir MY, et al. New MRI grading system for the diagnosis and management of mycetoma. *Trans R Soc Trop Med Hyg*. 2012;106:738–742.
- Welsh O, Al-Abdely HM, Salinas-Carmona MC, et al. Mycetoma medical therapy. *PLoS Negl Trop Dis*. 2014;8:e3218.
- Ameen M, Arenas R. Developments in the management of mycetomas. *Clin Exp Dermatol*. 2008;34:1–7.
- Zijlstra EE, van de Sande WW, Welsh O, et al. Mycetoma: a unique neglected tropical disease. *Lancet Infect Dis*. 2016;16:100–112.
- Suleiman SH, Wadaella ELS, Fahal AH. The surgical treatment of mycetoma. *PLoS Negl Trop Dis*. 2016;10:e0004690.
- Raman M, Bhat R, Garg T, et al. A modified two-step treatment for actinomycetoma. *Indian J Dermatol Venereol Leprol*. 2007;73:235–239.
- Fahal AH. Evidence-based guidelines for the management of mycetoma patients. Mycetoma Research Centre, Khartoum, Sudan, October 2002. www.mycetoma.edu.sd.
- Wortman PD. Treatment of a *Nocardia brasiliensis* mycetoma with sulfamethoxazole and trimethoprim, amikacin, and amoxicillin and clavulanate. *Arch Dermatol*. 1993;129:564–567.
- Ameen M, Arenas R, del Mercado EV, et al. Efficacy of imipenem therapy for *Nocardia* actinomycetomas refractory to sulfonamides. *J Am Acad Dermatol*. 2010;62:239–246.
- Zein HAM, Fahal AH, Mahgoub ES, et al. Predictors of cure, amputation and follow-up dropout among patients with mycetoma seen at the Mycetoma Research Centre, University of Khartoum, Sudan. *Trans R Soc Trop Med Hyg*. 2012;106:639–644.
- Fahal AH, Rahman IA, El-Hassan AM, et al. The safety and efficacy of itraconazole for the treatment of patients with eumycetoma due to *Madurella mycetomatis*. *Trans R Soc Trop Med Hyg*. 2011;105:127–132.
- Kloeezen W, Meis JF, Curfs-Breuker I, et al. In vitro antifungal activity of isavuconazole against *Madurella mycetomatis*. *Antimicrob Agents Chemother*. 2012;56:6054–6056.
- Lacroix C, de Kendler E, Morel P, et al. *Madurella mycetomatis* mycetoma treated successfully with oral voriconazole. *Br J Dermatol*. 2005;152:1062–1094.
- Loulergue P, Hot A, Dannaoui E, et al. Short report. Successful treatment of black-grain mycetoma with voriconazole. *Am J Trop Med Hyg*. 2006;75:1106–1107.
- Porte L, Khatibi S, El Hajj L, et al. *Scedosporium apiospermum* mycetoma with bone involvement successfully treated with voriconazole. *Trans R Soc Trop Med Hyg*. 2006;100:891–894.
- Negroni R, Tobon A, Bustamante B, et al. Posaconazole treatment of refractory eumycetoma and chromoblastomycosis. *Rev Inst Med Trop Sao Paulo*. 2005;47:339–346.
- N'Diaye B, Dieng MT, Perez A, et al. Clinical efficacy and safety of oral terbinafine in fungal mycetoma. *Int J Dermatol*. 2006;45:154–157.
- van Belkum A, Fahal AH, van de Sande WWJ. In vitro susceptibility of *Madurella mycetomatis* to posaconazole and terbinafine. *Antimicrob Agents Chemother*. 2011;55:1771–1773.

Cryptococcosis (*Cryptococcus neoformans* and *Cryptococcus gattii*)

John R. Perfect

SHORT VIEW SUMMARY

Life Cycle

- The life cycle represents both asexual (clinical) and sexual (recombination) cycles, with the impact of the sexual cycle on pathogenesis.

Taxonomy

- There are 19 cryptococcal species with two major pathogenic species, *Cryptococcus neoformans* and *Cryptococcus gattii*.
- At present, the following taxonomic divisions have been proposed, but this area continues to evolve in the molecular era: *C. neoformans* var. *grubii* (serotype A), with five genotypes (VNI, VNII, VNBI, VNBII, and the hybrid VNIII); *C. neoformans* var. *neoformans* (serotype D or genotype VNIV); and five other cryptic species: *Cryptococcus gattii*, *Cryptococcus bacillisporus*, *Cryptococcus deuterogattii*, *Cryptococcus tetragattii*, and *Cryptococcus decagattii* (serotypes B/C or VGI–VGV).¹

Identification

- Biochemical tests, including urease production, melanin formation, and appearance of a capsule
- Antigen tests, cultures, DNA-based tests, or matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF)

Ecology

- *C. neoformans* (serotypes A and D) found in pigeon guano or associated with certain trees (mopane trees)
- *C. gattii* found around trees from eucalyptus to coniferous species (landscape of niche may be changing related to climate changes)

Epidemiology

- Not a routine constituent of human microbiota but can colonize humans without evidence of disease
- Widespread asymptomatic infections occur in the exposed populations
- Risk factors for disease: acquired immunodeficiency syndrome (AIDS), corticosteroid treatment, transplantation, cancer, monoclonal antibodies (from anti-tumor necrosis factor to anti-CD52), sarcoidosis, autoantibodies to granulocyte-macrophage colony-stimulating factor, idiopathic CD4 lymphocytopenia, and new tyrosine kinase inhibitors for cancer treatment; identifies primarily cell-mediated immunity defects but humoral perturbations may also be a factor (see Table 262.1)
- Antiretroviral therapy (ART) impact on appearance of clinical cryptococcosis is impressive in patients with AIDS. However, even widespread availability of ART does not

eliminate high numbers of cryptococcosis in some populations.

- A million new cases of cryptococcosis per year, with more than 600,000 deaths worldwide at the peak of the human immunodeficiency virus (HIV) epidemic; now reduced although not yet eliminated with ART availability, with an estimated present census of over 200,000 new cases per year worldwide with over 100,000 deaths per year.

Pathogenicity

- Virulence factors (examples include but not limited to: capsule, melanin, high temperature growth, urease, phospholipase)
- Many understandings of cryptococcal disease at the genetic and molecular level have been acquired.

Host Responses

- Efficient protection requires intact and coordinated cell-mediated, innate, and humoral immunity networks.

Pathogenesis

- Three factors: (1) host defenses, (2) virulence of strains, and (3) size of infectious inocula
- Cryptococcosis is considered primarily a reactivation clinical disease but disease can occur with the initial infection.

Clinical Manifestations (see Table 262.2)

- Major body sites for infection are the central nervous system and lung.
- Other body sites with unique infection considerations are the skin, prostate, peritoneum, and eye.
- Cryptococci have produced infection within any organ of the human body.
- Disease primarily depends on host immunity—either too little or too much (immune reconstitution inflammatory syndrome [IRIS]).

Laboratory Diagnosis

- Microscopy: India ink and histopathologic stains, including a general fungal Gomori methenamine silver stain, or the specific capsular polysaccharide stains (mucicarmine or alcian blue)
- Culture of cerebrospinal fluid (CSF), blood, bronchoalveolar lavage
- Cryptococcal antigen detection in CSF and serum is sensitive and specific as determined by either latex agglutination assay, enzyme-linked immunosorbent assay, or a lateral flow assay.

Management

- The Infectious Diseases Society of American Guidelines in 2010 established road maps for

therapies and strategies (<https://academic.oup.com/cid/article/50/3/291/392360>); most of its principles are still intact but a few updates can be made.

- Direct antifungal drug resistance uncommonly arises during therapy, but relapse/persistent isolates should have in vitro antifungal susceptibility testing performed and compared to original isolate.
- Treatment strategies are outlined in this chapter for (1) cryptococcal meningitis in AIDS patients, transplant recipients, and non-AIDS, nontransplant patients and (2) pulmonary infections.
- Treatment has been best studied for cryptococcal meningitis in HIV-infected patients (2018 US Department of Health and Human Services guidelines at <https://aidsinfo.nih.gov/guidelines/html/4/adult-and-adolescent-oi-prevention-and-treatment-guidelines/333/cryptococcosis>). Treatment is initiated with amphotericin B, preferably with the liposomal formulation at 3 to 4 mg/kg daily, the alternatives being the lipid complex at 5 mg/kg or conventional amphotericin B at 0.7 mg/kg/day. Any one of these can be used plus flucytosine 25 mg/kg every 6 hours (100 mg/kg/day with normal renal function) for at least 2 weeks and until clinically improved. In cryptococcal meningitis, delaying ART during these 2 weeks may prevent IRIS from complicating management and increased early deaths. Patients who respond to induction-phase combination antifungal therapy may then be switched to fluconazole, 400 to 800 mg/day, for 8 to 10 weeks as a consolidation phase. Finally, a suppressive phase is begun with fluconazole 200 mg once daily. Antifungal therapy in HIV-infected patients can be stopped after 1 to 2 years in patients who respond to ART with a CD4 count above 100/μL for at least 3 months, a nondetectable viral load, and a negative or low serum cryptococcal antigen. In non-HIV-infected patients who improve, therapy is also generally extended for 1 year.
- Complications of cryptococcal meningitis requiring special attention are (1) early increased intracranial pressure, which can lead to blindness, permanent dementia, and death; (2) development of hydrocephalus, which may require placement of a ventriculoperitoneal shunt; and (3) IRIS, which may be mistaken for therapeutic failure.