

symptomatic children fail to control an outbreak of diarrhea, consideration can be given to treating all infected children.²³¹ Treatment-based strategies for transmission prevention in endemic settings, however, have not proven beneficial. Although metronidazole exposure within the previous 7 days decreased the likelihood of *Giardia* detection⁶⁰ in separate community-based studies of children in high-transmission

endemic settings, community-based administration of metronidazole, albendazole, or nitazoxanide did not diminish *Giardia* prevalence.^{118,195,232}

Human milk can be protective against *Giardia*. Both human and animal breast milk can contain anti-*Giardia* antibodies; studies have demonstrated protection of breastfeeding infants from symptomatic infection.^{114,233,234}

Key References

The complete reference list is available online at Expert Consult.

- Dawson SC. An insider's guide to the microtubule cytoskeleton of *Giardia*. *Cell Microbiol*. 2010;12:588–598.
- Yu LZ, Birkly CW Jr, Adam RD. The two nuclei of *Giardia* each have complete copies of the genome and are partitioned equationally at cytokinesis. *Eukaryot Cell*. 2002;1:191–199.
- Ankarklev J, Jerlstrom-Hultqvist J, Ringqvist E, et al. Behind the smile: cell biology and disease mechanisms of *Giardia* species. *Nat Rev Microbiol*. 2010;8:413–422.
- Clark CG, Diamond LS. Methods for cultivation of luminal parasitic protists of clinical importance. *Clin Microbiol Rev*. 2002;15:329–341.
- Adam RD. Biology of *Giardia lamblia*. *Clin Micro Rev*. 2001;14:447–475.
- Nash TE, McCutchan T, Keister D, et al. Restriction-endonuclease analysis of DNA from 15 *Giardia* isolates obtained from humans and animals. *J Infect Dis*. 1985;152:64–73.
- Feng Y, Xiao L. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. *Clin Microbiol Rev*. 2011;110–140.
- Caccio SM, Ryan U. Molecular epidemiology of giardiasis. *Mol Biochem Parasitol*. 2008;160:75–80.
- Franzen O, Jerlstrom-Hultqvist J, Castro E, et al. Draft genome sequencing of *Giardia intestinalis* assemblage B isolate GS: is human giardiasis caused by two different species? *PLoS Pathog*. 2009;5:e1000560.
- Nash TE, Herrington DA, Losonsky GA, et al. Experimental human infections with *Giardia lamblia*. *J Infect Dis*. 1987;156:974–984.
- Haque R, Roy S, Kabir M, et al. *Giardia* assemblage A infection and diarrhea in Bangladesh. *J Infect Dis*. 2005;192:2171–2173.
- Morrison HG, McArthur AG, Gillin FD, et al. Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science*. 2007;317:1921–1926.
- Ye Q, Tian H, Chen B, et al. *Giardia*'s primitive GPL biosynthesis pathways with parasitic adaptation 'patches': implications for *Giardia*'s evolutionary history and for finding targets against Giardiasis. *Sci Rep*. 2017;7:9507.
- Pyrhijä J, Pyrihová E, Kolisko M, et al. Minimal cytosolic iron-sulfur cluster assembly machinery of *Giardia intestinalis* is partially associated with mitochondria. *Mol Microbiol*. 2016;102:701–714.
- Adam RD, Aggarwal A, Lal AA, et al. Antigenic variation of a cysteine-rich protein in *Giardia lamblia*. *J Exp Med*. 1988;167:109–118.
- Nash TE, Herrington DA, Levine MM, et al. Antigenic variation of *Giardia lamblia* in experimental human infections. *J Immunol*. 1990;144:4362–4369.
- Nash TE. Surface antigenic variation in *Giardia lamblia*. *Mol Microbiol*. 2002;45:585–590.
- Adam RD, Nigam A, Seshadri V, et al. The *Giardia lamblia* vsp gene repertoire: characteristics, genomic organization, and evolution. *BMC Genomics*. 2010;11:424.
- Prucca CG, Rivero FD, Lujan HD. Regulation of antigenic variation in *Giardia lamblia*. *Annu Rev Microbiol*. 2011;65:611–630.
- Singer SM, Elmendorf HG, Conrad JT, et al. Biological selection of variant-specific surface proteins in *Giardia lamblia*. *J Infect Dis*. 2001;183:119–124.
- Carranza PG, Lujan HD. New insights regarding the biology of *Giardia lamblia*. *Microbes Infect*. 2010;12:71–80.
- Gillin FD, Reiner DS, McCaffery JM. Cell biology of the primitive eukaryote *Giardia lamblia*. *Annu Rev Microbiol*. 1996;50:679–705.
- Touz MC, Zamponi N. Sorting without a Golgi complex. *Traffic*. 2017;18:637–645.
- Svard SG, Hagblom P, Palm JE. *Giardia lamblia*—a model organism for eukaryotic cell differentiation. *FEMS Microbiol Lett*. 2003;218:3–7.
- DuBois KN, Abodeedy M, Sakanari J, et al. Identification of the major cysteine protease of *Giardia* and its role in encystation. *J Biol Chem*. 2008;283:18024–18031.
- Palm D, Weiland M, McArthur AG, et al. Developmental changes in the adhesive disk during *Giardia* differentiation. *Mol Biochem Parasitol*. 2005;141:199–207.
- Roxström-Lindquist K, Palm D, Reiner D, et al. *Giardia* immunity—an update. *Trends Parasitol*. 2006;22:26–31.
- Berkman DS, Lescano AG, Gilman RH, et al. Effects of stunting, diarrhoeal disease, and parasitic infection during infancy on cognition in late childhood: a follow-up study. *Lancet*. 2002;359:564–571.
- Rogawski ET, Bartelt LA, Platts-Mills JA, et al. Determinants and impact of *Giardia* infection in the first 2 years of life in the MAL-ED Birth cohort. *J Pediatr Infect Dis*. 2017;6:153–160.
- Hollm-Delgado MG, Gilman RH, Bern C, et al. Lack of an adverse effect of *Giardia intestinalis* infection on the health of Peruvian children. *Am J Epidemiol*. 2008;168:647–655.
- Craun GF, Brunkard JM, Yoder JS, et al. Causes of outbreaks associated with drinking water in the United States from 1971 to 2006. *Clin Microbiol Rev*. 2010;23:507–528.
- Adam EA, Yoder JS, Gould LH, et al. Giardiasis outbreaks in the United States, 1971–2011. *Epidemiol Infect*. 2016;144:2790–2801.
- Stuart JM, Orr HJ, Warburton FG, et al. Risk factors for sporadic giardiasis: a case-control study in southwestern England. *Emerg Infect Dis*. 2003;9:229–233.
- Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis*. 2011;17:7–15.
- Thompson SC. *Giardia lamblia* in children and the child care setting: a review of the literature. *J Paediatr Child Health*. 1994;30:202–209.
- Swaminathan A, Torresi J, Schlagenhauf P, et al. A global study of pathogens and host risk factors associated with infectious gastrointestinal disease in returned international travellers. *J Infect*. 2009;59:19–27.
- Oberhuber G, Kastner N, Stolte M. Giardiasis: a histologic analysis of 567 cases. *Scand J Infect Dis*. 1997;32:48–51.
- Hjelt K, Pørrregaard A, Krasilnikov PA. Giardiasis causing chronic diarrhoea in suburban Copenhagen: incidence, physical growth, clinical symptoms and small intestinal abnormality. *Acta Paediatr*. 1992;81:881–886.
- Humen MA, Perez PF, Lievin-Le Moal V. Lipid raft-dependent adhesion of *Giardia intestinalis* trophozoites to a cultured human endocyte-like Caco-2/TC7 cell monolayer leads to cytoskeleton-dependent functional injuries. *Cell Microbiol*. 2011;13:1683–1702.
- Maia-Brigagao C, Morgado-Diaz JA, De Souza W. *Giardia* disrupts the arrangement of tight, adherens and desmosomal junction proteins of intestinal cells. *Parasitol Int*. 2012;61:280–287.
- Bartelt LA, Roche J, Kolling G, et al. Persistent *G. lamblia* impairs growth in a murine malnutrition model. *J Clin Invest*. 2013;123:2672–2684.
- Stadelmann B, Merino MC, Persson L, et al. Arginine consumption by the intestinal parasite *Giardia intestinalis* reduces proliferation of intestinal epithelial cells. *PLoS ONE*. 2012;7:e45325.
- Cotton JA, Bhargava A, Ferraz JG, et al. *Giardia duodenalis* cathepsin B proteases degrade intestinal epithelial interleukin-8 and attenuate interleukin-8-induced neutrophil chemotaxis. *Infect Immun*. 2014;82:2772–2787.
- Aggarwal A, Nash TE. Antigenic variation of *Giardia lamblia* in vivo. *Infect Immun*. 1988;56:1420–1423.
- Solaymani-Mohammadi S, Singer SM. Host immunity and pathogen strain contribute to intestinal disaccharidase impairment following gut infection. *J Immunol*. 2011;187:3769–3775.
- Bienz M, Dai WJ, Welle M, et al. Interleukin-6-deficient mice are highly susceptible to *Giardia lamblia* infection but exhibit normal intestinal immunoglobulin A responses against the parasite. *Infect Immun*. 2003;71:1569–1573.
- Zhou P, Li E, Zhu N, et al. Role of interleukin-6 in the control of acute and chronic *Giardia lamblia* infections in mice. *Infect Immun*. 2003;71:1566–1568.
- Solaymani-Mohammadi S, Singer SM. *Giardia duodenalis*: the double-edged sword of immune responses in giardiasis. *Exp Parasitol*. 2010;126:292–297.
- Singer SM, Nash TE. T-cell-dependent control of acute *Giardia lamblia* infections in mice. *Infect Immun*. 2000;68:170–175.
- Istre GR, Dunlop TS, Gaspard B, et al. Waterborne giardiasis at a mountain resort: evidence for acquired immunity. *Am J Public Health*. 1984;74:602–604.
- Gilman RH, Marquis GS, Miranda E, et al. Rapid reinfection by *Giardia lamblia* after treatment in a hyperendemic third world community. *Lancet*. 1988;1:343–345.
- Langford TD, Housley MP, Boes M, et al. Central importance of immunoglobulin A in host defense against *Giardia* spp. *Infect Immun*. 2002;70:11–18.
- Nash TE, Ohi CA, Thomas E, et al. Treatment of patients with refractory giardiasis. *Clin Infect Dis*. 2001;33:22–28.
- Hanevik K, Krisoffersen E, Svard S, et al. Human cellular immune response against *Giardia lamblia* 5 years after acute giardiasis. *J Infect Dis*. 2011;204:1779–1786.
- Rivero FD, Saura A, Prucca CG, et al. Disruption of antigenic variation is crucial for effective parasite vaccine. *Nat Med*. 2010;16:551–557.
- Dann SM, Manthey CR, Le C, et al. IL-17A promotes protective IgA responses and expression of other potential effectors against the lumen-dwelling enteric parasite *Giardia*. *Exp Parasitol*. 2015;156:68–78.
- Ridley MJ, Ridley DS. Serum antibodies and jejunal histology in giardiasis. *J Clin Pathol*. 1976;29:30–34.
- Bartelt LA, Bolick DT, Mayneris-Pexachs J, et al. Cross-modulation of pathogen-specific pathways enhances malnutrition during enteric co-infection with *Giardia lamblia* and enteroaggregative *Escherichia coli*. *PLoS Pathog*. 2017;13:e1006471.
- Singer SM, Nash TE. The role of normal flora in mice. *J Infect Dis*. 2000;181:1510–1512.
- Gerbaba TK, Gupta P, Rioux K, et al. *Giardia duodenalis*-induced alterations of commensal bacteria kill *Caenorhabditis elegans*: a new model to study microbial-microbial interactions in the gut. *Am J Physiol Gastrointest Liver Physiol*. 2015;308:G550–G561.
- Keselman A, Li E, Maloney J, et al. The microbiota contributes to CD8+ T cell activation and nutrient malabsorption following intestinal infection with *Giardia duodenalis*. *Infect Immun*. 2016;84:2853–2860.
- Smith PD, Gillin FD, Spira WM, et al. Chronic giardiasis: studies on drug sensitivity, toxin production, and host immune response. *Gastroenterology*. 1982;83:797–803.
- Cantey PT, Roy S, Lee B, et al. Study of nonoutbreak giardiasis: novel findings and implications for research. *Am J Med*. 2011;124:1175.e1–1175.e8.
- Hanevik K, Wensaas KA, Rortveit G, et al. Irritable bowel syndrome and chronic fatigue 6 years after *Giardia* infection: a controlled prospective cohort study. *Clin Infect Dis*. 2014;59:1394–1400.
- Muhsen K, Levine MM. A systematic review and meta-analysis of the association between *Giardia lamblia* and endemic pediatric diarrhea in developing countries. *Clin Infect Dis*. 2012;55(suppl 4):S271–S293.
- Veenemans J, Mank T, Ottenhof M, et al. Protection against diarrhea associated with *Giardia intestinalis* is lost with multi-nutrient supplementation: a study in Tanzanian children. *PLoS Negl Trop Dis*. 2011;5:e1158.
- Ignatius R, Gahutu JB, Klotz C, et al. High prevalence of *Giardia duodenalis* Assemblage B infection and association with underweight in Rwandan children. *PLoS Negl Trop Dis*. 2012;6:e1677.
- Farthing MJG, Mata L, Urrutia JJ, et al. Natural history of *Giardia* infection of infants and children in rural Guatemala and its impact on physical growth. *Am J Clin Nutr*. 1986;43:395–405.
- Sullivan PS, DuPont HL, Arafat RR, et al. Illness and reservoirs associated with *Giardia lamblia* infection in rural Egypt: the case against treatment in developing world environments of high endemicity. *Am J Epidemiol*. 1988;127:1272–1281.
- Haque R, Mondal D, Karim A, et al. Prospective case-control study of the association between common enteric protozoal parasites and diarrhea in Bangladesh. *Clin Infect Dis*. 2009;48:1191–1197.
- Ish-Horowitz M, Korman SH, Shapiro M, et al. Asymptomatic giardiasis in children. *Pediatr Infect Dis J*. 1989;8:773–779.
- Hiatt RA, Markell EK, Ng E. How many stool examinations are necessary to detect pathogenic intestinal protozoa? *Am J Trop Med Hyg*. 1995;53:36–39.

189. Garcia LS, Shimizu RY. Evaluation of nine immunoassay kits (enzyme immunoassay and direct fluorescence) for detection of *Giardia lamblia* and *Cryptosporidium parvum* in human fecal specimens. *J Clin Microbiol.* 1997;35:1526–1529.
190. Aldeen WE, Carroll K, Robison A, et al. Comparison of nine commercially available enzyme-linked immunosorbent assays for the detection of *Giardia lamblia* in fecal specimens. *J Clin Microbiol.* 1998;36:1338–1340.
192. Haque R, Roy S, Siddique A, et al. Multiplex real-time PCR assay for detection of *Entamoeba histolytica*, *Giardia intestinalis*, and *Cryptosporidium* spp. *Am J Trop Med Hyg.* 2007;76:713–717.
195. Mejia R, Vicuna Y, Broncano N, et al. A novel, multi-parallel, real-time polymerase chain reaction approach for eight gastrointestinal parasites provides improved diagnostic capabilities to resource-limited at-risk populations. *Am J Trop Med Hyg.* 2013;88:1041–1047.
199. Polage CR, Stoddard GJ, Rolfs RT, et al. Physician use of parasite tests in the g from 1997 to 2006 and in a Utah *Cryptosporidium* outbreak in 2007. *J Clin Microbiol.* 2011;49:591–596.
202. Isaac-Renton J, Blatherwick J, Bowie WR, et al. Epidemic and endemic seroprevalence of antibodies to *Cryptosporidium* and *Giardia* in residents of three communities with different drinking water supplies. *Am J Trop Med Hyg.* 1999;60:578–583.
205. Gardner TB, Hill DR. Treatment of giardiasis. *Clin Micro Rev.* 2001;14:114–128.
208. Granados CE, Reveiz L, Uribe LG, et al. Drugs for treating giardiasis. *Cochrane Database Syst Rev.* 2012;(12):CD007787.
209. Lalle M. Giardiasis in the post genomic era: treatment, drug resistance and novel therapeutic perspectives. *Infect Disord Drug Targets.* 2010;10:283–294.
210. Speelman P. Single-dose tinidazole for the treatment of giardiasis. *Antimicrob Agents Chemother.* 1985;27:227–229.
215. Rossignol JF. *Cryptosporidium* and *Giardia*: treatment options and prospects for new drugs. *Exp Parasitol.* 2010;124:45–53.
230. Backer HD. Water disinfection for international travelers. In: Keystone JS, Kozarsky PE, Freedman DO, et al, eds. *Travel Medicine.* 3rd ed. Philadelphia: Elsevier; 2013:37–49.
233. Mahmud MA, Chappell CL, Hossain MM, et al. Impact of breast-feeding on *Giardia lamblia* infections in Bilbeis, Egypt. *Am J Trop Med Hyg.* 2001;65:257–260.

References

- Dawson SC. An insider's guide to the microtubule cytoskeleton of *Giardia*. *Cell Microbiol*. 2010;12:588–598.
- Schwartz CL, Heumann JM, Dawson SC, et al. A detailed, hierarchical study of *Giardia lamblia*'s ventral disc reveals novel microtubule-associated protein complexes. *PLoS ONE*. 2012;7:e43783.
- Hagen KD, Hirakawa MP, House SA, et al. Novel structural components of the ventral disc and lateral crest in *Giardia intestinalis*. *PLoS Negl Trop Dis*. 2011;5:e1442.
- Yu LZ, Birky CW Jr, Adam RD. The two nuclei of *Giardia* each have complete copies of the genome and are partitioned equationally at cytokinesis. *Eukaryot Cell*. 2002;1:191–199.
- Ankarklev J, Jerlstrom-Hultqvist J, Ringqvist E, et al. Behind the smile: cell biology and disease mechanisms of *Giardia* species. *Nat Rev Microbiol*. 2010;8:413–422.
- Carpenter ML, Assaf ZJ, Gourguechon S, et al. Nuclear inheritance and genetic exchange without meiosis in the binucleate parasite *Giardia intestinalis*. *J Cell Sci*. 2012;125(Pt 10):2523–2532.
- Clark CG, Diamond LS. Methods for cultivation of luminal parasitic protists of clinical importance. *Clin Microbiol Rev*. 2002;15:329–341.
- Adam RD. Biology of *Giardia lamblia*. *Clin Micro Rev*. 2001;14:447–475.
- Di Matteo A, Scandurra FM, Testa F, et al. The O2-scavenging flavodiiron protein in the human parasite *Giardia intestinalis*. *J Biol Chem*. 2008;283:4061–4068.
- Nash TE, McCutchan T, Keister D, et al. Restriction-endonuclease analysis of DNA from 15 *Giardia* isolates obtained from humans and animals. *J Infect Dis*. 1985;152:64–73.
- Xiao L, Fayer R. Molecular characterisation of species and genotypes of *Cryptosporidium* and *Giardia* and assessment of zoonotic transmission. *Int J Parasitol*. 2008;38:1239–1255.
- Feng Y, Xiao L. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. *Clin Microbiol Rev*. 2011;24:110–140.
- Caccio SM, Ryan U. Molecular epidemiology of giardiasis. *Mol Biochem Parasitol*. 2008;160:75–80.
- Wielinga CM, Thompson RC. Comparative evaluation of *Giardia duodenalis* sequence data. *Parasitology*. 2007;134(Pt 12):1795–1821.
- Fantiniati M, Bello AR, Fernandes O, et al. Identification of *Giardia lamblia* Assemblage E in humans points to a new anthroponotic cycle. *J Infect Dis*. 2016;214:1256–1259.
- Franzen O, Jerlstrom-Hultqvist J, Castro E, et al. Draft genome sequencing of *Giardia intestinalis* assemblage B isolate GS: is human giardiasis caused by two different species? *PLoS Pathog*. 2009;5:e1000560.
- Nash T. Surface antigen variability and variation in *Giardia lamblia*. *Parasitol Today*. 1992;8:229–234.
- Sprong H, Caccio SM, van der Giessen JW. Identification of zoonotic genotypes of *Giardia duodenalis*. *PLoS Negl Trop Dis*. 2009;3:e558.
- Nash TE, Herrington DA, Losonsky GA, et al. Experimental human infections with *Giardia lamblia*. *J Infect Dis*. 1987;156:974–984.
- Haque R, Roy S, Kabir M, et al. *Giardia* assemblage A infection and diarrhea in Bangladesh. *J Infect Dis*. 2005;192:2171–2173.
- Sahagun J, Clavel A, Goni P, et al. Correlation between the presence of symptoms and the *Giardia duodenalis* genotype. *Eur J Clin Microbiol Infect Dis*. 2008;27:81–83.
- Perez Cordon G, Cordova Paz Soldan O, Vargas Vasquez F, et al. Prevalence of enteroparasites and genotyping of *Giardia lamblia* in Peruvian children. *Parasitol Res*. 2008;103:459–465.
- Morrison HG, McArthur AG, Gillin FD, et al. Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science*. 2007;317:1921–1926.
- Davis-Hayman SR, Nash TE. Genetic manipulation of *Giardia lamblia*. *Mol Biochem Parasitol*. 2002;122:1–7.
- Ye Q, Tian H, Chen B, et al. *Giardia*'s primitive GPL biosynthesis pathways with parasitic adaptation 'patches': implications for *Giardia*'s evolutionary history and for finding targets against Giardiasis. *Sci Rep*. 2017;7:9507.
- Lane S, Lloyd D. Current trends in research into the waterborne parasite *Giardia*. *Crit Rev Microbiol*. 2002;28:123–147.
- Pyrhij J, Pyrhijová E, Kolisko M, et al. Minimal cytosolic iron-sulfur cluster assembly machinery of *Giardia intestinalis* is partially associated with mitochondria. *Mol Microbiol*. 2016;102:701–714.
- Adam RD, Aggarwal A, Lal AA, et al. Antigenic variation of a cysteine-rich protein in *Giardia lamblia*. *J Exp Med*. 1988;167:109–118.
- Nash TE, Herrington DA, Levine MM, et al. Antigenic variation of *Giardia lamblia* in experimental human infections. *J Immunol*. 1990;144:4362–4369.
- Nash TE. Surface antigenic variation in *Giardia lamblia*. *Mol Microbiol*. 2002;45:585–590.
- Adam RD, Nigam A, Seshadri V, et al. The *Giardia lamblia* *Giardia* vsp gene repertoire: characteristics, genomic organization, and evolution. *BMC Genomics*. 2010;11:424.
- Prucsa CG, Rivero FD, Lujan HD. Regulation of antigenic variation in *Giardia lamblia*. *Annu Rev Microbiol*. 2011;65:611–630.
- Upcroft JA, Krauer KG, Upcroft P. Chromosome sequence maps of the *Giardia lamblia* assemblage A isolate WB. *Trends Parasitol*. 2010;26:484–491.
- Saraiya AA, Li W, Wang CC. A microRNA derived from an apparent canonical biogenesis pathway regulates variant surface protein gene expression in *Giardia lamblia*. *RNA*. 2011;17:2152–2164.
- Singer SM, Elmendorf HG, Conrad JT, et al. Biological selection of variant-specific surface proteins in *Giardia lamblia*. *J Infect Dis*. 2001;183:119–124.
- Lauwaet T, Davids BJ, Reiner DS, et al. Encystation of *Giardia lamblia*: a model for other parasites. *Curr Opin Microbiol*. 2007;10:554–559.
- Carranza PG, Lujan HD. New insights regarding the biology of *Giardia lamblia*. *Microbes Infect*. 2010;12:71–80.
- Faso C, Hehl AB. Membrane trafficking and organelle biogenesis in *Giardia lamblia*: use it or lose it. *Int J Parasitol*. 2011;41:471–480.
- Gillin FD, Reiner DS, McCaffery JM. Cell biology of the primitive eukaryote *Giardia lamblia*. *Annu Rev Microbiol*. 1996;50:679–705.
- Touz MC, Zamponi N. Sorting without a Golgi complex. *Traffic*. 2017;18:637–645.
- Svard SG, Hagblom P, Palm JE. *Giardia lamblia*—a model organism for eukaryotic cell differentiation. *FEMS Microbiol Lett*. 2003;218:3–7.
- Morf L, Spycher C, Rehrauer H, et al. The transcriptional response to encystation stimuli in *Giardia lamblia* is restricted to a small set of genes. *Eukaryot Cell*. 2010;9:1566–1576.
- DuBois KN, Abodeely M, Sakanari J, et al. Identification of the major cysteine protease of *Giardia* and its role in encystation. *J Biol Chem*. 2008;283:18024–18031.
- Konrad C, Spycher C, Hehl AB. Selective condensations drives partitioning and sequential secretion of cyst wall proteins in differentiating *Giardia lamblia*. *PLoS Pathog*. 2010;6:e1000835.
- Erlandsen SL, Macechko PT, van Keulen H, et al. Formation of the *Giardia* cyst wall: studies on extracellular assembly using immunogold labeling and high resolution field emission SEM. *J Eukaryot Microbiol*. 1996;43:416–429.
- Chatterjee A, Carpentieri A, Ratner DM, et al. *Giardia* cyst wall protein 1 is a lectin that binds to curled fibrils of the GalNAc homopolymer. *PLoS Pathog*. 2010;6:e1001059.
- Hetsko ML, McCaffery JM, Svard SG, et al. Cellular and transcriptional changes during excystation of *Giardia lamblia* in vitro. *Exp Parasitol*. 1998;88:172–183.
- Palm D, Weiland M, McArthur AG, et al. Developmental changes in the adhesive disk during *Giardia* differentiation. *Mol Biochem Parasitol*. 2005;141:199–207.
- Ward W, Alvarado L, Rawlings ND, et al. A primitive enzyme for a primitive cell: the protease required for excystation of *Giardia*. *Cell*. 1997;89:437–444.
- Reiner DS, Hetsko ML, Meszaros JG, et al. Calcium signaling in excystation of the early diverging eukaryote, *Giardia lamblia*. *J Biol Chem*. 2003;278:2533–2540.
- Roxström-Lindquist K, Palm D, Reiner D, et al. *Giardia* immunity—an update. *Trends Parasitol*. 2006;22:26–31.
- Painter JE, Gargano JW, Collier SA, et al. Giardiasis surveillance – United States, 2011–2012. *MMWR*. 2015;64:15–25.
- Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis*. 2011;17:7–15.
- Thompson SC. *Giardia lamblia* in children and the child care setting: a review of the literature. *J Paediatr Child Health*. 1994;30:202–209.
- Overturf GD. Endemic giardiasis in the United States—role of the daycare center [editorial]. *Clin Infect Dis*. 1994;18:764–765.
- Waldram A, Vivanos R, Harley C, et al. Prevalence of *Giardia* infection in households of *Giardia* cases and risk factors for household transmission. *BMC Infect Dis*. 2017;17:486.
- Swaminathan A, Torresi J, Schlegelhauf P, et al. A global study of pathogens and host risk factors associated with infectious gastrointestinal disease in returned international travellers. *J Infect*. 2009;59:19–27.
- Mark-Carew MP, Wade SE, Chang YF, et al. Prevalence of *Giardia duodenalis* assemblages among dairy herds in the New York City Watershed. *Vet Parasitol*. 2012;185:151–157.
- Minetti C, Tawenan W, Hogg R, et al. Occurrence and diversity of *Giardia duodenalis* assemblages in livestock in the UK. *Transbound Emerg Dis*. 2014;61:e60–e67.
- Isaac-Renton JL, Cordeiro C, Sarafis K, et al. Characterization of *Giardia duodenalis* isolates from a waterborne outbreak. *J Infect Dis*. 1993;167:431–440.
- Moorehead WP, Guasparini R, Donovan CA, et al. Giardiasis outbreak from a chlorinated community water supply. *Can J Public Health*. 1990;81:358–362.
- Rendtorff RC. The experimental transmission of human intestinal protozoan parasites: II. *Giardia lamblia* cysts given in capsules. *Am J Hyg*. 1954;59:209–220.
- Prado MS, Cairncross S, Strina A, et al. Asymptomatic giardiasis and growth in young children: a longitudinal study in Salvador, Brazil. *Parasitology*. 2005;131(Pt 1):51–56.
- Rogawski ET, Bartelt LA, Platts-Mills JA, et al. Determinants and impact of *Giardia* infection in the first 2 years of life in the MAL-ED Birth cohort. *J Ped Infect Dis*. 2017;6:153–160.
- Donowitz JR, Alam M, Kabir M, et al. A prospective longitudinal cohort to investigate the effects of early life giardiasis on growth and all cause diarrhea. *Clin Infect Dis*. 2016;63:792–797.
- Al-Mekhlafi MS, Azlin M, Nor Aini U, et al. Giardiasis as a predictor of childhood malnutrition in Orang Asli children in Malaysia. *Trans R Soc Trop Med Hyg*. 2005;99:686–691.
- Carvalho-Costa FA, Gonçalves AQ, Lassance SL, et al. *Giardia lamblia* and other intestinal parasitic infections and their relationships with nutritional status in children in Brazilian Amazon. *Rev Inst Med Trop São Paulo*. 2007;49:147–153.
- Hollm-Delgado MG, Gilman RH, Bern C, et al. Lack of an adverse effect of *Giardia intestinalis* infection on the health of Peruvian children. *Am J Epidemiol*. 2008;168:647–655.
- Mahmud MA, Chappell C, Hossain MM, et al. Risk factors for development of first symptomatic *Giardia* infection among infants of a birth cohort in rural Egypt. *Am J Trop Med Hyg*. 1995;53:84–88.
- Dib HH, Lu SQ, Wen SF. Prevalence of *Giardia lamblia* with or without diarrhea in South East, South East Asia and the Far East. *Parasitol Res*. 2008;103:239–251.
- Taniuchi M, Sobuz SU, Begum S, et al. Etiology of diarrhea in Bangladeshi infants in the first year of life analyzed using molecular methods. *J Infect Dis*. 2013;208:794–802.
- Craun GF, Brunkard JM, Yoder JS, et al. Causes of outbreaks associated with drinking water in the United States from 1971 to 2006. *Clin Microbiol Rev*. 2010;23:507–528.
- Adam EA, Yoder JS, Gould LH, et al. Giardiasis outbreaks in the United States, 1971–2011. *Epidemiol Infect*. 2016;144:2790–2801.
- Stuart JM, Orr HJ, Warburton FG, et al. Risk factors for sporadic giardiasis: a case-control study in southwestern England. *Emerg Infect Dis*. 2003;9:229–233.
- Daly ER, Roy SJ, Blaney DD, et al. Outbreak of giardiasis associated with a community drinking-water source. *Epidemiol Infect*. 2010;138:491–500.
- Beer KD, Gargano JW, Roberts VA, et al. Outbreaks associated with environmental and undetermined water exposures—United States 2011–2012. *MMWR Morb Mortal Wkly Rep*. 2015;63:849–851.
- Baldursson S, Karanis P. Waterborne transmission of protozoan parasites: review of worldwide outbreaks—an update 2004–2010. *Water Res*. 2011;45:6603–6614.
- Mintz ED, Hudson-Wragg M, Mshar P, et al. Foodborne giardiasis in a corporate office setting. *J Infect Dis*. 1993;167:250–253.
- Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis*. 2011;17:7–15.
- Thompson SC. *Giardia lamblia* in children and the child care setting: a review of the literature. *J Paediatr Child Health*. 1994;30:202–209.
- Overturf GD. Endemic giardiasis in the United States—role of the daycare center [editorial]. *Clin Infect Dis*. 1994;18:764–765.
- Waldram A, Vivanos R, Harley C, et al. Prevalence of *Giardia* infection in households of *Giardia* cases and risk factors for household transmission. *BMC Infect Dis*. 2017;17:486.
- Swaminathan A, Torresi J, Schlegelhauf P, et al. A global study of pathogens and host risk factors associated with infectious gastrointestinal disease in returned international travellers. *J Infect*. 2009;59:19–27.
- Mark-Carew MP, Wade SE, Chang YF, et al. Prevalence of *Giardia duodenalis* assemblages among dairy herds in the New York City Watershed. *Vet Parasitol*. 2012;185:151–157.
- Minetti C, Tawenan W, Hogg R, et al. Occurrence and diversity of *Giardia duodenalis* assemblages in livestock in the UK. *Transbound Emerg Dis*. 2014;61:e60–e67.
- Isaac-Renton JL, Cordeiro C, Sarafis K, et al. Characterization of *Giardia duodenalis* isolates from a waterborne outbreak. *J Infect Dis*. 1993;167:431–440.
- Moorehead WP, Guasparini R, Donovan CA, et al. Giardiasis outbreak from a chlorinated community water supply. *Can J Public Health*. 1990;81:358–362.
- Rendtorff RC. The experimental transmission of human intestinal protozoan parasites: II. *Giardia lamblia* cysts given in capsules. *Am J Hyg*. 1954;59:209–220.

85. Kohli A, Bushen OY, Pinkerton RC, et al. *Giardia duodenalis* assemblage, clinical presentation and markers of intestinal inflammation in Brazilian children. *Trans R Soc Trop Med Hyg.* 2008;102:718–725.
86. Jenkins MC, O'Brien CN, Murphy C, et al. Antibodies to the ventral disc protein delta-giardin prevent in vitro binding of *Giardia lamblia* trophozoites. *J Parasitol.* 2009;95:895–899.
87. House SA, Richter DJ, Pham JK, et al. *Giardia* flagellar motility is not directly required to maintain attachment to surfaces. *PLoS Pathog.* 2011;7:e1002167.
88. Oberhuber G, Kastner N, Stolte M. Giardiasis: a histologic analysis of 567 cases. *Scand J Infect Dis.* 1997;32:48–51.
89. Duncombe VM, Bolin TD, Davis AE, et al. Histopathology in giardiasis: a correlation with diarrhea. *Aust N Z J Med.* 1978;8:392–396.
90. Hanevik K, Hausken T, Morken MH, et al. Persisting symptoms and duodenal inflammation related to *Giardia duodenalis* infection. *J Infect.* 2007;55:524–530.
91. Hjelt K, Poerregaard A, Krasilnikoff PA. Giardiasis causing chronic diarrhoea in suburban Copenhagen: incidence, physical growth, clinical symptoms and small intestinal abnormality. *Acta Paediatr.* 1992;81:881–886.
92. Humen MA, Perez PF, Lievin-Le Moal V. Lipid raft-dependent adhesion of *Giardia intestinalis* trophozoites to a cultured human enterocyte-like Caco-2/TC7 cell monolayer leads to cytoskeleton-dependent functional injuries. *Cell Microbiol.* 2011;13:1683–1702.
93. Chin AC, Teoh DA, Scott KG, et al. Strain-dependent induction of enterocyte apoptosis by *Giardia lamblia* disrupts epithelial barrier function in a caspase-3-dependent manner. *Infect Immun.* 2002;70:3673–3680.
94. Panaro MA, Cianciulli A, Mitolo V, et al. Caspase-dependent apoptosis of the HCT-8 epithelial cell line induced by the parasite *Giardia intestinalis*. *FEMS Immunol Med Microbiol.* 2007;51:302–309.
95. Buret AG. Pathophysiology of enteric infections with *Giardia duodenalis*. *Parasite.* 2008;15:261–265.
96. Maia-Brigao C, Morgado-Diaz JA, De Souza W. *Giardia* disrupts the arrangement of tight, adherens and desmosomal junction proteins of intestinal cells. *Parasitol Int.* 2012;61:280–287.
97. Halliez MCM, Motta J-P, Feener TD, et al. *Giardia duodenalis* induces paracellular bacterial translocation and causes postinfectious visceral hypersensitivity. *Am J Physiol Gastrointest Liver Physiol.* 2016;310:G574–G585.
98. Troeger H, Eppe HJ, Schneider T, et al. Effect of chronic *Giardia lamblia* infection on epithelial transport and barrier function in human duodenum. *Gut.* 2007;56:328–335.
99. Bartelt LA, Roche J, Kolling G, et al. Persistent *G. lamblia* impairs growth in a murine malnutrition model. *J Clin Invest.* 2013;123:2672–2684.
100. Yu LC, Huang CY, Kuo WT, et al. SGLT-1-mediated glucose uptake protects human intestinal epithelial cells against *Giardia duodenalis*-induced apoptosis. *Int J Parasitol.* 2008;38:923–934.
101. Stadelmann B, Merino MC, Persson L, et al. Arginine consumption by the intestinal parasite *Giardia intestinalis* reduces proliferation of intestinal epithelial cells. *PLoS ONE.* 2012;7:e45325.
102. Ringqvist E, Palm JE, Skarin H, et al. Release of metabolic enzymes by *Giardia* in response to interaction with intestinal epithelial cells. *Mol Biochem Parasitol.* 2008;159:85–91.
103. Li E, Zhou P, Singer SM. Neuronal nitric oxide synthase is necessary for elimination of *Giardia lamblia* infections in mice. *J Immunol.* 2006;176:516–521.
104. Cotton JA, Bhargava A, Ferraz JG, et al. *Giardia duodenalis* cathepsin B proteases degrade intestinal epithelial interleukin-8 and attenuate interleukin-8-induced neutrophil chemotaxis. *Infect Immun.* 2014;82:2772–2787.
105. Aggarwal A, Nash TE. Antigenic variation of *Giardia lamblia* in vivo. *Infect Immun.* 1988;56:1420–1423.
106. Solaymani-Mohammadi S, Singer SM. Host immunity and pathogen strain contribute to intestinal disaccharidase impairment following gut infection. *J Immunol.* 2011;187:3769–3775.
107. Tako EA, Hassimi MF, Li E, et al. Transcriptomic analysis of the host response to *Giardia duodenalis* infection reveals redundant mechanisms for parasite control. *MBio.* 2013;4:e00660–13.
108. Bienz M, Dai WJ, Welle M, et al. Interleukin-6-deficient mice are highly susceptible to *Giardia lamblia* infection but exhibit normal intestinal immunoglobulin A responses against the parasite. *Infect Immun.* 2003;71:1569–1573.
109. Zhou P, Li E, Zhu N, et al. Role of interleukin-6 in the control of acute and chronic *Giardia lamblia* infections in mice. *Infect Immun.* 2003;71:1566–1568.
110. Solaymani-Mohammadi S, Singer SM. *Giardia duodenalis*: the double-edged sword of immune responses in giardiasis. *Exp Parasitol.* 2010;126:292–297.
111. Andersen YS, Gillin FD, Eckmann L. Adaptive immunity-dependent intestinal hypermotility contributes to host defense against *Giardia* spp. *Infect Immun.* 2006;74:2473–2476.
112. Li E, Zhao A, Shea-Donohue T, et al. Mast cell-mediated changes in smooth muscle contractility during mouse giardiasis. *Infect Immun.* 2007;75:4514–4518.
113. Kamda JD, Nash TE, Singer SM. *Giardia duodenalis*: dendritic cell defects in IL-6 deficient mice contribute to susceptibility to intestinal infection. *Exp Parasitol.* 2012;130:288–291.
114. Li X, Zhang X, Gong P, et al. TLR2/- mice display decreased severity of giardiasis via enhanced proinflammatory cytokines production dependent on AKT signal pathway. *Front Immunol.* 2017;8:1186.
115. Singer SM, Nash TE. T-cell-dependent control of acute *Giardia lamblia* infections in mice. *Infect Immun.* 2000;68:170–175.
116. Istre GR, Dunlop TS, Gaspard B, et al. Waterborne giardiasis at a mountain resort: evidence for acquired immunity. *Am J Public Health.* 1984;74:602–604.
117. Isaac-Renton JL, Lewis LF, Ong CS, et al. A second community outbreak of waterborne giardiasis in Canada and serological investigation of patients. *Trans R Soc Trop Med Hyg.* 1994;88:395–399.
118. Gilman RH, Marquis GS, Miranda E, et al. Rapid reinfection by *Giardia lamblia* after treatment in a hyperendemic third world community. *Lancet.* 1988;1:343–345.
119. Stäger S, Müller N. *Giardia lamblia* infections in B-cell-deficient transgenic mice. *Infect Immun.* 1997;65:3944–3946.
120. Rosales-Borjas DM, Diaz-Rivadeneira J, Dona-Leyva A, et al. Secretory immune response to membrane antigens during *Giardia lamblia* infection in humans. *Infect Immun.* 1998;66:756–759.
121. Langford TD, Housley MP, Boes M, et al. Central importance of immunoglobulin A in host defense against *Giardia* spp. *Infect Immun.* 2002;70:11–18.
122. Char S, Cevallos AM, Yanson P, et al. Impaired IgA response to *Giardia* heat shock antigen in children with persistent diarrhoea and giardiasis. *Gut.* 1993;34:38–40.
123. Oksenhendler E, Gerard L, Fieschi C, et al. Infections in 252 patients with common variable immunodeficiency. *Clin Infect Dis.* 2008;46:1547–1554.
124. Janoff EN, Smith PD, Blaser MJ. Acute antibody responses to *Giardia lamblia* are depressed in patients with AIDS. *J Infect Dis.* 1988;157:798–804.
125. Nash TE, Ohi CA, Thomas E, et al. Treatment of patients with refractory giardiasis. *Clin Infect Dis.* 2001;33:22–28.
126. Saghaug CS, Sørnes S, Peirasmaki D, et al. Human memory CD4+ T cell immune responses against *Giardia lamblia*. *Clin Vaccine Immunol.* 2015;23:11–18.
127. Hanevik K, Krisoffersen E, Svard S, et al. Human cellular immune response against *Giardia lamblia* 5 Years after acute giardiasis. *J Infect Dis.* 2011;204:1779–1786.
128. Rivero FD, Saura A, Prucca CG, et al. Disruption of antigenic variation is crucial for effective parasite vaccine. *Nat Med.* 2010;16:551–557.
129. Davids BJ, Palm JE, Housley MP, et al. Polymeric immunoglobulin receptor in intestinal immune defense against the lumen-dwelling protozoan parasite *Giardia*. *J Immunol.* 2006;177:6281–6290.
130. Dann SM, Mantey CR, Le C, et al. IL-17A promotes protective IgA responses and expression of other potential effectors against the lumen-dwelling enteric parasite *Giardia*. *Exp Parasitol.* 2015;156:68–78.
131. Paerewijck O, Maertens B, Dreesen L, et al. Interleukin-17 receptor A (IL-17RA) as a central regulator of the protective immune response against *Giardia*. *Sci Rep.* 2017;7:8520.
132. Heyworth MF. Immunology of *Giardia* and *Cryptosporidium* infections. *J Infect Dis.* 1992;166:465–472.
133. Scott KG, Logan MR, Klammer GM, et al. Jejunal brush border microvillous alterations in *Giardia muris*-infected mice: role of T lymphocytes and interleukin-6. *Infect Immun.* 2000;68:3412–3418.
134. Ridley MJ, Ridley DS. Serum antibodies and jejunal histology in giardiasis. *J Clin Pathol.* 1976;29:30–34.
135. Cotton JA, Motta JP, Schenck LP, et al. *Giardia duodenalis* infection reduces granulocyte infiltration in an in vivo model of bacterial toxin-induced colitis and attenuates inflammation in human intestinal tissue. *PLoS ONE.* 2014;9:e109087.
136. Manko A, Motta JP, Cotton JA, et al. *Giardia* co-infection promotes the secretion of antimicrobial peptides beta-defensin 2 and trefoil factor 3 and attenuates attaching and effacing bacteria-induced intestinal disease. *PLoS ONE.* 2017;12:e0178647.
137. Bartelt LA, Bolick DT, Mayneris-Perxachs J, et al. Cross-modulation of pathogen-specific pathways enhances malnutrition during enteric co-infection with *Giardia lamblia* and enteroaggregative *Escherichia coli*. *PLoS Pathog.* 2017;13:e1006471.
138. Singer SM, Nash TE. The role of normal flora in mice. *J Infect Dis.* 2000;181:1510–1512.
139. Travers MA, Sow C, Zirah S, et al. Deconjugated bile salts produced by extracellular bile-salt hydrolase-like activities from the probiotic *Lactobacillus johnsonii* La1 inhibit *Giardia duodenalis* in vitro growth. *Front Microbiol.* 2016;7:1453.
140. Torres MF, Uetanabaro AP, Costa AF, et al. Influence of bacteria from the duodenal microbiota of patients with symptomatic giardiasis on the pathogenicity of *Giardia duodenalis* in gnotoxenic mice. *J Med Microbiol.* 2000;49:209–215.
141. Barash NR, Malnoey JG, Singer SM, et al. *Giardia* alters commensal microbial diversity throughout the murine gut. *Infect Immun.* 2017;85:e00948–16.
142. Gerbaba TK, Gupta P, Rioux K, et al. *Giardia duodenalis*-induced alterations of commensal bacteria kill *Caenorhabditis elegans*: a new model to study microbial-microbial interactions in the gut. *Am J Physiol Gastrointest Liver Physiol.* 2015;308:G550–G561.
143. Beatty JK, Akierman SV, Motta JP, et al. *Giardia duodenalis* induces pathogenic dysbiosis of human intestinal microbiota biofilms. *Int J Parasitol.* 2017;47:311–326.
144. Chen TL, Chen S, Wu HW, et al. Persistent gut barrier damage and commensal bacterial influx following eradication of *Giardia* infection in mice. *Gut Pathog.* 2013;5:26.
145. Keselman A, Li E, Maloney J, et al. The microbiota contributes to CD8+ T Cell activation and nutrient malabsorption following intestinal infection with *Giardia duodenalis*. *Infect Immun.* 2016;84:2853–2860.
146. Smith PD, Gillin FD, Spira WM, et al. Chronic giardiasis: studies on drug sensitivity, toxin production, and host immune response. *Gastroenterology.* 1982;83:797–803.
147. Cantey PT, Roy S, Lee B, et al. Study of nonoutbreak giardiasis: novel findings and implications for research. *Am J Med.* 2011;124:1175.e1–1175.e8.
148. Hill DR. Giardiasis. Issues in diagnosis and management. *Infect Dis Clin North Am.* 1993;7:503–525.
149. Painter JE, et al. Association between *Giardia* and arthritis or joint pain in a large health insurance cohort: could it be reactive arthritis. *Epidemiol Infect.* 2017;145:471–477.
150. Berney DM, Rampton D, van der Walt JD. Giardiasis of the stomach. *Postgrad Med J.* 1994;70:237–238.
151. Ankarklev J, Hestvik E, Lebbad M, et al. Common coinfections of *Giardia intestinalis* and *Helicobacter pylori* in non-symptomatic Ugandan children. *PLoS Negl Trop Dis.* 2012;6:e1780.
152. Lengerich EJ, Addiss DG, Juranek DD. Severe giardiasis in the United States. *Clin Infect Dis.* 1994;18:760–763.
153. Robertson LJ. Severe giardiasis and cryptosporidiosis in Scotland, UK. *Epidemiol Infect.* 1996;117:551–561.
154. Behera B, Mirdha BR, Makharia GK, et al. Parasites in patients with malabsorption syndrome: a clinical study in children and adults. *Dig Dis Sci.* 2008;53:672–679.
155. Solomons NW. Giardiasis: nutritional implications. *Rev Infect Dis.* 1982;4:859–869.
156. Gillon J. Clinical studies in adults presenting with giardiasis to a gastrointestinal unit. *Scot Med J.* 1985;30:89–95.
157. Rana SV, Bhasin DK, Vinayak VK. Lactose hydrogen breath test in *Giardia lamblia*-positive patients. *Dig Dis Sci.* 2005;50:259–261.
158. Litleskare S, Wensasas KA, Eide GE, et al. Perceived food intolerance and irritable bowel syndrome in a population 3 years after a giardiasis-outbreak: a historical cohort study. *BMC Gastroenterol.* 2015;15:164.
159. Hanevik K, Wensasas KA, Rortveit G, et al. Irritable bowel syndrome and chronic fatigue 6 years after giardia infection: a controlled prospective cohort study. *Clin Infect Dis.* 2014;59:1394–1400.
160. Hanevik K, Kristoffersen E, Mørch K, et al. *Giardia*-specific cellular immune responses in post-giardiasis chronic fatigue syndrome. *BMC Immunol.* 2017;18:5.
161. Nako JK, Collier SA, Gargano JW. Giardiasis and subsequent irritable bowel syndrome: a longitudinal cohort study using health insurance data. *J Infect Dis.* 2017;215:798–805.
162. Bartelt LA, Platts-Mills JA. *Curr Opin Infect Dis.* 2016;29:502–507.
163. Muhsen K, Levine MM. A systematic review and meta-analysis of the association between *Giardia lamblia* and endemic pediatric diarrhea in developing countries. *Clin Infect Dis.* 2012;55(suppl 4):S271–S293.
164. Kotloff KL, Nataro JP, Blackwelder WC, et al. Burden and aetiology of diarrhoeal disease in infants and young

- children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet*. 2013;382:209–222.
165. Breurec S, Vanel N, Bata P, et al. Etiology and epidemiology of diarrhea in hospitalized children from low income country: a matched case-control study in Central African Republic. *PLoS Negl Trop Dis*. 2016;10:e0004283.
 166. Liu J, Platts-Mills JA, Juma J, et al. Use of quantitative molecular diagnostic methods to identify cases of diarrhea in children: a reanalysis of the GEMS case-control study. *Lancet*. 2016;388:1291–1301.
 167. Veenemans J, Mank T, Ottenhof M, et al. Protection against diarrhea associated with *Giardia intestinalis* is lost with multi-nutrient supplementation: a study in Tanzanian children. *PLoS Negl Trop Dis*. 2011;5:e1158.
 168. Sackey ME, Weigel MM, Armijos RX. Predictors and nutritional consequences of intestinal parasitic infections in rural Ecuadorian children. *J Trop Pediatr*. 2003;49:17–23.
 169. Goto R, Panter-Brick C, Northrop-Clewes CA, et al. Poor intestinal permeability in mildly stunted Nepali children: associations with weaning practices and *Giardia lamblia* infection. *Br J Nutr*. 2002;88:141–149.
 170. Newman RD, Moore SR, Lima AA, et al. A longitudinal study of *Giardia lamblia* infection in north-east Brazilian children. *Trop Med Int Health*. 2001;6:624–634.
 171. Botero-Garcés JH, García-Montoya GM, Grisales-Patino D, et al. *Giardia intestinalis* and nutritional status in children participating in the complementary nutrition program, Antioquia, Colombia, May to October 2006. *Rev Inst Med Trop São Paulo*. 2009;51:155–162.
 172. Ignatius R, Gahutu JB, Klotz C, et al. High prevalence of *Giardia duodenalis* Assemblage B infection and association with underweight in Rwandan children. *PLoS Negl Trop Dis*. 2012;6:e1677.
 173. Aijampur SS, Koshy B, Venkataramani M, et al. Effect of cryptosporidial and giardial diarrhoea on social maturity, intelligence and physical growth in children in a semi-urban slum in south India. *Ann Trop Paediatr*. 2011;31:205–212.
 174. Al-Mekhlafi HM, Al-Makatar MT, Jani R, et al. Burden of *Giardia duodenalis* infection and its adverse effects on growth of schoolchildren in rural Malaysia. *PLoS Negl Trop Dis*. 2013;7:e2516.
 175. Gupta MC, Urrutia JJ. Effect of periodic antiscaris and anti-*Giardia* treatment on nutritional status of preschool children. *Am J Clin Nutr*. 1982;36:79–86.
 176. Farthing MJG, Mata L, Urrutia JJ, et al. Natural history of *Giardia* infection of infants and children in rural Guatemala and its impact on physical growth. *Am J Clin Nutr*. 1986;43:395–405.
 177. Fraser D, Bilenko N, Deckelbaum RJ, et al. *Giardia lamblia* carriage in Israeli Bedouin infants: risk factors and consequences. *Clin Infect Dis*. 2000;30:419–424.
 178. Sullivan PS, DuPont HL, Arafat RR, et al. Illness and reservoirs associated with *Giardia lamblia* infection in rural Egypt: the case against treatment in developing world environments of high endemicity. *Am J Epidemiol*. 1988;127:1272–1281.
 179. Lunn PG, Erinoso HO, Northrop-Clewes CA, et al. *Giardia intestinalis* is unlikely to be a major cause of the poor growth of rural Gambian infants. *J Nutr*. 1999;129:872–877.
 180. Goto R, Mascie-Taylor CG, Lunn PG. Impact of anti-*Giardia* and anthelmintic treatment on infant growth and intestinal permeability in rural Bangladesh: a randomised double-blind controlled study. *Trans R Soc Trop Med Hyg*. 2009;103:520–529.
 181. Mondal D, Petri WA Jr, Sack RB, et al. *Entamoeba histolytica*-associated diarrheal illness is negatively associated with the growth of preschool children: evidence from a prospective study. *Trans R Soc Trop Med Hyg*. 2006;100:1032–1038.
 182. Haque R, Mondal D, Karim A, et al. Prospective case-control study of the association between common enteric protozoal parasites and diarrhea in Bangladesh. *Clin Infect Dis*. 2009;48:1191–1197.
 183. Ish-Horowitz M, Korman SH, Shapiro M, et al. Asymptomatic giardiasis in children. *Pediatr Infect Dis J*. 1989;8:773–779.
 184. Moya-Camarena SY, Sotelo N, Valencia ME. Effects of asymptomatic *Giardia intestinalis* infection on carbohydrate absorption in well-nourished Mexican children. *Am J Trop Med Hyg*. 2002;66:255–259.
 185. Bhavnani D, Goldstick JE, Cevallos W, et al. Synergistic effects between rotavirus and coinfecting pathogens on diarrheal disease: evidence from a community-based study in northwestern Ecuador. *Am J Epidemiol*. 2012;176:387–395.
 186. Becker-Dreps S, Bucardo F, Vilchez S, et al. Etiology of childhood diarrhea after rotavirus vaccine introduction: a prospective, population-based study in Nicaragua. *Pediatr Infect Dis J*. 2014;33:1156–1163.
 187. Hiatt RA, Markell EK, Ng E. How many stool examinations are necessary to detect pathogenic intestinal protozoa? *Am J Trop Med Hyg*. 1995;53:36–39.
 188. Mank TG, Zaat JO, Deelder AM, et al. Sensitivity of microscopy versus enzyme immunoassay in the laboratory diagnosis of giardiasis. *Eur J Clin Microbiol Infect Dis*. 1997;16:615–619.
 189. García LS, Shimizu RY. Evaluation of nine immunoassay kits (enzyme immunoassay and direct fluorescence) for detection of *Giardia lamblia* and *Cryptosporidium parvum* in human fecal specimens. *J Clin Microbiol*. 1997;35:1526–1529.
 190. Aldeen WE, Carroll K, Robison A, et al. Comparison of nine commercially available enzyme-linked immunosorbent assays for the detection of *Giardia lamblia* in fecal specimens. *J Clin Microbiol*. 1998;36:1338–1340.
 191. Church D, Miller K, Lichtenfeld A, et al. Screening for *Giardia/Cryptosporidium* infections using an enzyme immunoassay in a centralized regional microbiology laboratory. *Arch Pathol Lab Med*. 2005;129:754–759.
 192. Haque R, Roy S, Siddique A, et al. Multiplex real-time PCR assay for detection of *Entamoeba histolytica*, *Giardia intestinalis*, and *Cryptosporidium* spp. *Am J Trop Med Hyg*. 2007;76:713–717.
 193. Calderaro A, Gorrini C, Montecchini S, et al. Evaluation of a real-time polymerase chain reaction assay for the laboratory diagnosis of giardiasis. *Diagn Microbiol Infect Dis*. 2010;66:261–267.
 194. Vanni I, Caccio SM, van Lith L, et al. Detection of *Giardia duodenalis* assemblages A and B in human feces by simple, assemblage-specific PCR assays. *PLoS Negl Trop Dis*. 2012;6:e1776.
 195. Mejia R, Vicuna Y, Broncano N, et al. A novel, multi-parallel, real-time polymerase chain reaction approach for eight gastrointestinal parasites provides improved diagnostic capabilities to resource-limited at-risk populations. *Am J Trop Med Hyg*. 2013;88:1041–1047.
 196. Johnston SP, Ballard MM, Beach MJ, et al. Evaluation of three commercial assays for detection of *Giardia* and *Cryptosporidium* organisms in fecal specimens. *J Clin Microbiol*. 2003;41:623–626.
 197. García LS, García JP. Detection of *Giardia lamblia* antigens in human fecal specimens by a solid-phase qualitative immunochromatographic assay. *J Clin Microbiol*. 2006;44:4587–4588.
 198. Youn S, Kabir M, Haque R, et al. Evaluation of a screening test for detection of *Giardia* and *Cryptosporidium*. *J Clin Microbiol*. 2009;47:451–452.
 199. Polage CR, Stoddard GJ, Rolfs RT, et al. Physician use of parasite tests in the United States from 1997 to 2006 and in a Utah *Cryptosporidium* outbreak in 2007. *J Clin Microbiol*. 2011;49:591–596.
 200. Swierczewski B, Odundo E, Ndonye J, et al. Comparison of the Triage Micro Parasite Panel and microscopy for the detection of *Entamoeba histolytica/Entamoeba dispar*, *Giardia lamblia*, and *Cryptosporidium parvum* in stool samples collected in Kenya. *J Trop Med*. 2012;2012:564721.
 201. Soliman MM, Taghi-Kilani R, Abou-Shady AF, et al. Comparison of serum antibody response to *Giardia lamblia* of symptomatic and asymptomatic patients. *Am J Trop Med Hyg*. 1998;58:232–239.
 202. Isaac-Renton J, Blatherwick J, Bowie WR, et al. Epidemic and endemic seroprevalence of antibodies to *Cryptosporidium* and *Giardia* in residents of three communities with different drinking water supplies. *Am J Trop Med Hyg*. 1999;60:578–583.
 203. Crump JA, Mendoza CE, Priest JW, et al. Comparing serologic response against enteric pathogens with reported diarrhea to assess the impact of improved household drinking water quality. *Am J Trop Med Hyg*. 2007;77:136–141.
 204. Cedillo-Rivera R, Leal YA, Yezpe-Mulia L, et al. Seroepidemiology of giardiasis in Mexico. *Am J Trop Med Hyg*. 2009;80:6–10.
 205. Gardner TB, Hill DR. Treatment of giardiasis. *Clin Micro Rev*. 2001;14:114–128.
 206. Upcroft P, Upcroft JA. Drug targets and mechanisms of resistance in the anaerobic protozoa. *Clin Micro Rev*. 2001;14:150–164.
 207. Nabarro LE, Lever RA, Armstrong M, et al. Increased incidence of nitroimidazole-refractory giardiasis at the hospital for Tropical Diseases London: 2008–2013. *Clin Microbiol Infect*. 2015;21:791–796.
 208. Granados CE, Reveiz L, Uribe LG, et al. Drugs for treating giardiasis. *Cochrane Database Syst Rev*. 2012;(12):CD007787.
 209. Lalle M. Giardiasis in the post genomic era: treatment, drug resistance and novel therapeutic perspectives. *Infect Disord Drug Targets*. 2010;10:283–294.
 210. Speelman P. Single-dose tinidazole for the treatment of giardiasis. *Antimicrob Agents Chemother*. 1985;27:227–229.
 211. Falagas ME, Walker AM, Jick H, et al. Late incidence of cancer after metronidazole use: a matched metronidazole user/nonuser study. *Clin Infect Dis*. 1998;26:384–388.
 212. Samuelson J. Why metronidazole is active against both bacteria and parasites. *Antimicrob Agents Chemother*. 1999;43:1533–1541.
 213. Upcroft J, Upcroft P. My favorite cell: *Giardia*. *Bioessays*. 1998;20:256–263.
 214. Leitsch D, Burgess AG, Dunn LA, et al. Pyruvate:ferredoxin oxidoreductase and thioredoxin reductase are involved in 5-nitroimidazole activation while flavin metabolism is linked to 5-nitroimidazole resistance in *Giardia lamblia*. *J Antimicrob Chemother*. 2011;66:1756–1765.
 215. Rossignol JF. *Cryptosporidium* and *Giardia*: treatment options and prospects for new drugs. *Exp Parasitol*. 2010;124:45–53.
 216. Anderson VR, Curran MP. Nitazoxanide: a review of its use in the treatment of gastrointestinal infections. *Drugs*. 2007;67:1947–1967.
 217. Escobedo AA, Alvarez G, Gonzalez ME, et al. The treatment of giardiasis in children: single-dose tinidazole compared with 3 days of nitazoxanide. *Ann Trop Med Parasitol*. 2008;102:199–207.
 218. Hoffman PS, Sisson G, Croxen MA, et al. Antiparasitic drug nitazoxanide inhibits the pyruvate oxidoreductases of *Helicobacter pylori*, selected anaerobic bacteria and parasites, and *Campylobacter jejuni*. *Antimicrob Agents Chemother*. 2007;51:868–876.
 219. Solaymani-Mohammadi S, Genkinger JM, Loffredo CA, et al. A meta-analysis of the effectiveness of albendazole compared with metronidazole as treatments for infections with *Giardia duodenalis*. *PLoS Negl Trop Dis*. 2010;4:e682.
 220. Abboud P, Leme V, Gargala G, et al. Successful treatment of metronidazole- and albendazole-resistant giardiasis with nitazoxanide in a patient with acquired immunodeficiency syndrome. *Clin Infect Dis*. 2001;32:1792–1794.
 221. Sawatzki M, Peter S, Hess C. Therapy-resistant diarrhea due to *Giardia lamblia* in a patient with common variable immunodeficiency disease. *Digestion*. 2007;75:101–102.
 222. Mørch K, Hanevik K, Robertson LJ, et al. Treatment-ladder and genetic characterisation of parasites in refractory giardiasis after an outbreak in Norway. *J Infect*. 2008;56:268–273.
 223. Besirbellioglu BA, Ulcay A, Can M, et al. *Saccharomyces boulardii* and infection due to *Giardia lamblia*. *Scand J Infect Dis*. 2006;38:479–481.
 224. Rotblatt MD. Giardiasis and amebiasis in pregnancy. *Drug Intell Clin Pharm*. 1983;17:187–188.
 225. Burtin P, Taddio A, Ariburnu O, et al. Safety of metronidazole in pregnancy: a meta-analysis. *Am J Obstet Gynecol*. 1995;172:525–529.
 226. Rosado JL, Caamano MC, Montoya YA, et al. Interaction of zinc or vitamin A supplementation and specific parasite infections on Mexican infants' growth: a randomized clinical trial. *Eur J Clin Nutr*. 2009;63:1176–1184.
 227. Quhui L, Morales GG, Mendez RO, et al. Could giardiasis be a risk factor for low zinc status in schoolchildren from northwestern Mexico? A cross-sectional study with longitudinal follow-up. *BMC Public Health*. 2010;10:85.
 228. Lima AA, Soares AM, Lima NL, et al. Effects of vitamin A supplementation on intestinal barrier function, growth, total parasitic, and specific *Giardia* spp infections in Brazilian children: a prospective randomized, double-blind, placebo-controlled trial. *J Pediatr Gastroenterol Nutr*. 2010;50:309–315.
 229. Morken MH, Valeur J, Norin E, et al. Antibiotic or bacterial therapy in post-giardiasis irritable bowel syndrome. *Scand J Gastroenterol*. 2009;44:1296–1303.
 230. Backer HD. Water disinfection for international travelers. In: Keystone JS, Kozarsky PE, Freedman DO, et al, eds. *Travel Medicine*. 3rd ed. Philadelphia: Elsevier; 2013:37–49.
 231. Bartlett AV, Englander SJ, Jarvis BA, et al. Controlled trial of *Giardia lamblia*: control strategies in day care centers. *Am J Public Health*. 1991;81:1001–1006.
 232. Asai T, Córdova Vidal C, Strauss W, et al. Effect of mass stool examination and mass treatment for decreasing intestinal helminth and protozoan infection rates in Bolivian children: a cross-sectional study. *PLoS Negl Trop Dis*. 2016;10:e0005147.
 233. Mahmud MA, Chappell CL, Hossain MM, et al. Impact of breast-feeding on *Giardia lamblia* infections in Bilbeis, Egypt. *Am J Trop Med Hyg*. 2001;65:257–260.
 234. Bilenko N, Ghosh R, Levy A, et al. Partial breastfeeding protects Bedouin infants from infection and morbidity: prospective cohort study. *Asia Pac J Clin Nutr*. 2008;17:243–249.

SHORT VIEW SUMMARY

Microbiology and Epidemiology

- *Trichomonas vaginalis* is a flagellated parasite.
- Infection is through sexual transmission.
- Susceptibility to human immunodeficiency virus (HIV) infection may be increased in infected women.

Diagnosis

- Infection causes vaginitis in women and may cause urethritis in men; it is often asymptomatic.

- In women, most cases are diagnosed using wet prep microscopy of vaginal fluid as the point-of-care test, but this lacks sensitivity.
- The gold standard for diagnosis is nucleic acid amplification testing.

Therapy

- Metronidazole is the usual therapy.
- Resistance to metronidazole may occur.
- Tinidazole may be more effective.

Prevention

- Sexual partners should be treated.
- Condoms prevent infection.

Trichomonas vaginalis is the causative agent of trichomoniasis, a common cause of vaginitis. In men, it may cause urethritis but is more commonly asymptomatic. Despite being a readily diagnosed and treatable sexually transmitted disease (STD), trichomoniasis is not a reportable infection and its control has received relatively little emphasis from public health STD control programs. More recently, however, recognition of the high rates of disease and of associations of trichomoniasis in women with adverse outcomes of pregnancy and increased risk for human immunodeficiency virus (HIV) infection suggest a need for increased control efforts.

TAXONOMY

Trichomonas vaginalis is a parasitic pear-shaped protozoan, with an average size of $10 \times 7 \mu\text{m}$ (Fig. 280.1). It has four free flagella and one recurrent flagellum, along the outer margin of the undulating membrane; a costa at the base of the undulating membrane; and an axostyle extending through the cell.¹ Unlike most eukaryotes, *T. vaginalis* lacks mitochondria and instead uses the hydrogenosome to accomplish fermentative carbohydrate metabolism, with hydrogen as the electron acceptor. The hydrogenosome appears to have a common ancestry with mitochondria based on similarities in protein import.² However, major differences exist between hydrogenosomes and mitochondria in that hydrogenosomes lack cytochromes, mitochondrial respiratory chain enzymes, and their own DNA.

Trichomonas tenax, found in oral gingival and tracheobronchial sites, and *Pentatrichomonas hominis*, isolated from the intestinal tract, are considered nonpathogenic. Each human species has specific tropism for its site of infection. *Tritrichomonas foetus*, which is found exclusively in cattle, is perhaps the nonhuman trichomonad most similar to *T. vaginalis*. Aside from *T. foetus* having three anterior flagella (vs. four in *T. vaginalis*), there are few morphologic differences between the parasites, and *T. foetus* causes the STD known as bovine trichomoniasis. *Tritrichomonas foetus* can be invasive to the fetus, having been demonstrated in the placenta and the fetal lung, gut, and lymph nodes, and is a known cause of abortion in infected cattle.³

The life cycle of *T. vaginalis* is simple in that the trophozoite is transmitted through coitus and no cyst form is known. The trophozoite divides by binary fission and, in natural infections, gives rise to a population in the lumen and on the mucosal surfaces of the urogenital tracts of humans. In an infected person there are estimated to be 10^1 to 10^5

protozoa/mL of vaginal fluid.⁴ The organisms create microulcerations in the genital mucosa by direct contact, mediated by surface proteins.⁵

Our current understanding of immunity to *T. vaginalis* has come largely from observations of responses in human patients and experimentation using in vitro and animal models of the related species *T. foetus*. Natural infection seems to produce immunity that is only partially protective, because reinfection of patients can be as high as 30% on follow-up.⁶ The presence of *T. vaginalis* in the reproductive tract results in parasite-specific antibodies in the reproductive tract and, in most cases, circulating antibodies in the serum⁷; there is also evidence of lymphocyte priming, as detected by antigen-specific proliferation of peripheral blood mononuclear cells.⁸ Thus natural infection with *T. vaginalis* results in priming of acquired immune responses. A study of patients infected with *T. vaginalis* and HIV indicated no evidence of increased levels or duration of parasite infection in these patients compared with those in patients infected with *T. vaginalis* but not HIV.⁹ These observations may indicate that innate immunity involving chemotaxis and subsequent influx of neutrophils is much more important than acquired immunity in controlling infections with *T. vaginalis*, because neutrophils are often the most numerous leukocytes present in response to infection.¹⁰

The genome of *T. vaginalis* has been sequenced and provides considerable resources that may be used to define genes important in the pathogenesis of human trichomoniasis. The release of the 5X genome sequence data makes the most comprehensive genomic sequence of this parasite available to date.¹¹

EPIDEMIOLOGY

Humans are the only natural host of *T. vaginalis*. Trichomoniasis is an extremely common infection in the United States and worldwide. The prevalence of trichomoniasis in inner-city US STD clinics typically approaches 25% and may be higher in certain populations. In Los Angeles, for example, the prevalence among African-American patients at a public clinic was 38%.¹² Among women ages 14 to 19 years, the overall prevalence of trichomoniasis in the United States was recently found to be 3.1%, but among African Americans the prevalence was 13.3%.¹³ An estimate of the incidence of STDs in the United States yielded an annual incidence of trichomoniasis of 7.4 million new cases.¹⁴ The World Health Organization has estimated that this infection accounts for almost 50% of all curable infections worldwide.¹⁵ Studies of African



FIG. 280.1 *Trichomonas vaginalis*. (From Centers for Disease Control and Prevention, National Center for Infectious Diseases, Division of Parasitic Diseases.)



FIG. 280.2 Vaginal discharge in a patient with trichomoniasis. Note the bubbles, which give the discharge a “frothy” appearance.

populations have reported the prevalence of vaginal trichomoniasis to be between 11% and 25%. Laga and colleagues reported an incidence of 38% during a 4-month exposure interval among HIV-infected women in Zaire.¹⁶

Epidemiologically, *T. vaginalis* infections are commonly associated with other STDs and are a marker of high-risk sexual behavior. Trichomoniasis is frequently seen concomitantly with other STDs, particularly gonorrhea. Unlike other STDs, which have a higher prevalence among adolescents and young adults, the rates of trichomoniasis are more evenly distributed among sexually active women of all age groups, probably as a result of a lack of an organized disease control effort for this infection.¹⁷ Although survival on fomites is documented, the organism is thought to be transmitted almost exclusively by sexual activity.¹⁸ Vertical transmission from mother to infant can occur rarely and cause genital infection or respiratory distress.

The reported prevalence of urethral infection with *T. vaginalis* in males has varied, depending on the population studied and the diagnostic techniques used. In a series of sentinel studies using cultures of urine, urethra, coronal sulcus, and semen, Krieger and associates reported a prevalence of 11% in men attending an STD clinic.¹⁹ Urethritis was present in half of the men with *Trichomonas* as the sole urethral pathogen. In a similar study conducted at an STD clinic in Denver, investigators found a prevalence rate of 2.8% by using urine sediment culture.²⁰ At an STD clinic in Birmingham, Alabama, *T. vaginalis* was detected by polymerase chain reaction assay in 17% of men attending for a new-problem visit or screening. There was no significant difference in the detection of the organism between men with and without urethral symptoms (20% and 14.5%, respectively). Of men with nongonococcal urethritis (NGU), nearly as many had *T. vaginalis* detected as had *Chlamydia* (19.9% and 25.2%, respectively).²¹ Detection of *T. vaginalis* in men is increased if semen is tested in addition to urine and urethral specimens.²²

CLINICAL MANIFESTATIONS

The organisms create microulcerations in the genital mucosa by direct contact, mediated by surface proteins.⁵ In women, it is the squamous epithelium of the vagina that is infected. Interestingly, although 2% to 17% of female neonates may have vaginal colonization if the mother is infected, the infection may not be sustained once the maternal effects of estrogen wear off and the vaginal pH becomes neutral.²³ Neonatal respiratory tract infections with *T. vaginalis* have also been reported.²⁴ Detection of *Trichomonas* infection in older children should raise the suspicion of child abuse.²⁵

TABLE 280.1 Sensitivity of Clinical and Laboratory Findings in Vaginal Trichomoniasis

CLINICAL MANIFESTATION		PERCENT POSITIVE
Symptoms	None	9–56
	Discharge	50–75
	Malodorous	~10
	Irritating, pruritic	23–82
	Dyspareunia	10–50
	Dysuria	30–50
	Lower abdominal discomfort	5–12
Signs	None	~15
	Vulvar erythema	10–20
	Excessive discharge	50–75
	Yellow, green	5–20
	Frothy	10–50
	Vaginal wall inflammation	40–75
	Strawberry cervix (direct visualization)	1–2
	Colpitis macularis (colposcope)	45
Laboratory findings	pH >4.5	66–91
	Positive whiff test	~75
	Excess polymorphonuclear neutrophils on wet mount	~75

Data from Honigberg BM, ed. *Trichomonads Parasitic in Humans*. New York: Springer-Verlag; 1990; Bickley LS, Krisher KK, Punsalang A Jr, et al. Comparison of direct fluorescent antibody, acridine orange, wet mount, and culture for detection of *Trichomonas vaginalis* in women attending a public sexually transmitted disease clinic. *Sex Transm Dis*. 1989;16:127–131; and Rein MF. Uncertainties and controversies in trichomoniasis. In: Sobel JD, ed. *Vulvovaginal Infections: Current Concepts in Diagnosis and Therapy*. New York: Academy Professional Information Services; 1990:73–85.

The incubation period of this infection is unknown; however, human challenge studies done in the 1940s suggested an incubation period of 4 to 28 days in women.²⁶ The clinical manifestations of trichomoniasis in the female are summarized in Table 280.1. Symptoms in women include vaginal discharge, pruritus, and irritation. Signs of infection include vaginal discharge (42%), odor (50%), and edema or erythema (22% to 37%). The discharge is classically described as frothy, but it is actually frothy in only about 10% of patients (Fig. 280.2). The color of the discharge may vary. Colpitis macularis (strawberry cervix) is a specific clinical sign for this infection but is detected with reliability only by colposcopy and rarely during routine examination.¹⁰ Other complaints may include dysuria and lower abdominal pain; the cause

of the latter is unclear. The urethra is also infected in most women.²⁷ Almost 50% of all women with *T. vaginalis* are asymptomatic²⁸; therefore if these women are not screened, the diagnosis will be missed. The extent of the inflammatory response to the parasite may determine the severity of the symptoms. Factors that influence the host inflammatory response are not well understood but may include hormonal levels, coexisting vaginal flora, and strain and relative concentration of the organisms present in the vagina. In men the infection is usually asymptomatic, but some studies have suggested that it is a more common cause of NGU than was previously recognized^{19,21} and should certainly be considered as a diagnostic possibility in a man who does not respond to initial therapy for NGU. Trichomoniasis in men may also rarely cause epididymitis, prostatitis, and superficial penile ulcerations.²⁹ Spontaneous resolution of trichomoniasis in men has been reported.³⁰

DIAGNOSIS

Direct microscopic examination of vaginal fluid for the presence of motile trichomonads is the most common diagnostic method employed for the detection of trichomoniasis. However, even for skilled diagnosticians, the sensitivity of this test varies from 40 to 60%, and may be less in asymptomatic women.³¹ Sensitivity decreases dramatically if specimen transport time is longer than 10 minutes.³² The pH of the vaginal fluid will usually be higher than 4.5 in women with trichomoniasis but can be normal (≤ 4.5). In the latter cases, the trichomonads are often sparser because they prefer a more alkaline pH. Neutrophils as well as altered vaginal bacterial microbiota are often seen. Trichomoniasis is most often associated with intermediate vaginal microbiota as defined by Nugent criteria, raising the possibility that *T. vaginalis* may in some way be able to alter the vaginal microbiota composition to enhance its survival.³³ Culture media are commercially available and until recently have been the gold standard for diagnosis, but the cultured specimen requires incubation for up to 7 days prior to finalization.³⁴ Though not recommended routinely, culture is currently the only acceptable specimen type for susceptibility testing. Additional tests include the Affirm VPIII (Becton Dickinson, Franklin Lakes, NJ), a nucleic acid probe hybridization semiautomated system that also detects candidiasis and bacterial vaginosis and has results available in less than an hour. The OSOM Trichomonas Rapid Test (Sekisui, Lexington, MA), an antigen detection test using immunochromatographic capillary flow dipstick technology, is Clinical Laboratories Improvement Amendments–waived with results available in 10 minutes and sensitivity estimates from 77% to 98%, with specificities of 99% to 100%. Finally, AmpliVue Trichomonas assay and Solana Trichomonas assay (Quidel, San Diego, CA) use an isothermal helicase-dependent amplification-based method to detect trichomonas DNA, with results available within an hour and performance characteristics comparable to culture. All of these tests are currently licensed only for vaginal specimens, can be used as point of care tests, and have sensitivities of about 80%.^{35,36} They may be a good choice if microscopy is not available.

Nucleic acid amplification tests (NAATs) are now commercially available for the diagnosis of trichomoniasis and have replaced culture as the gold standard diagnostic test. Genital (endocervical, provider and patient-collected vaginal swabs), urine, and liquid PAP-based specimens are acceptable, with some variation depending on the specific NAAT test being utilized.³⁷ With sensitivities greater than 90% and specificities from 95% to 100%, current US Food and Drug Association (FDA)–approved NAATs include the Aptima *Trichomonas vaginalis* assay (Hologic, San Diego, CA); the ProbeTec *Trichomonas vaginalis* Q^x assay (BD Diagnostics, Franklin Lakes, NJ); and the Xpert TV test (Cepheid, Sunnyvale, CA),³⁸ which was recently approved by the FDA for use with male urine. Diagnosis in general is much more difficult for males, with the best culture results obtained by combining urethral swabs and urine sediment into one specimen. However, compared with NAATs, this method is highly insensitive for use in men.³⁹ Although most are not approved for men by the FDA, several studies have found NAATs to be highly sensitive and specific in this population and these tests are available from laboratories that have performed the necessary validation studies.³⁹

Lack of suitable screening and diagnostic tests for males has hampered public health control efforts of the disease. The increasing use of NAATs testing in women and men should be of great importance in this regard.

THERAPY

In most cases, the infection is easily treated with a single 2-g oral dose of metronidazole or tinidazole, and because it is an STD, sexual partners should be routinely treated.⁴⁰ The reported cure rates are 90% to 100%. Failure to treat the male sexual partner is likely the most common cause of recurrent disease in women and should be explored before assuming that the woman has a strain of *T. vaginalis* that is resistant to metronidazole. Metronidazole intravaginal gel has limited efficacy and should not be used. Although there continues to be some controversy about the safety of metronidazole in pregnancy, there has never been a documented case of fetal malformation attributed to its use, even when it is used in the first trimester.⁴¹ Tinidazole has a plasma half-life twice that of metronidazole (12–14 hours for tinidazole vs. 6–7 hours for metronidazole)⁴² and appears to be better tolerated, with fewer gastrointestinal side effects.

Resistance has been reported, estimated to be from 2.5% to 10% for metronidazole and less than 1% for tinidazole.⁴³ One mechanism of development of anaerobic resistance to metronidazole is controlled by hydrogenosomes, in that metronidazole competes for hydrogen as an electron acceptor. In metronidazole-resistant *T. vaginalis*, the expression levels of the hydrogenosomal enzymes pyruvate ferredoxin oxidoreductase, ferredoxin, malic enzyme, and hydrogenase are reduced dramatically, which probably eliminates the ability of the parasite to activate metronidazole.⁴⁴ Resistance has also been associated with mutations in nitroreductase genes.⁴⁵ Resistance is relative and can usually be overcome with higher doses of oral metronidazole. However, there does not appear to be a definitive correlation between in vitro and in vivo resistance. Intravenous formulations offer no advantage over the oral drug. Some authorities have recommended higher doses of oral medication in combination with pharmacy-prepared intravaginal preparations.

Tinidazole, with its more favorable pharmacokinetics, may be the drug of choice when resistance is encountered. The Centers for Disease Control and Prevention (CDC) tested 195 metronidazole-resistant *T. vaginalis* clinical isolates submitted for minimal lethal concentration (MLC) testing for both metronidazole and tinidazole. The mean aerobic metronidazole MLC was 400 $\mu\text{g/mL}$ compared with an aerobic tinidazole MLC of 100 $\mu\text{g/mL}$.⁴⁶ Several clinical studies have evaluated various doses of tinidazole for treatment of metronidazole-resistant trichomoniasis. The largest series of patients was reported by Sobel and associates.⁴⁷ In this study, 20 patients with clinically refractory trichomoniasis (failure to respond to therapy with oral metronidazole, at least 500 mg twice daily for 7 days) were treated with high doses of oral and vaginal tinidazole (2–3 g PO, plus 1–1.5 g intravaginally for 14 days). The cure rate was 92% (22 of 24 patients); no patients discontinued therapy because of side effects. Current CDC recommendations for patients failing a single 2-g dose of metronidazole are to treat next with metronidazole, 500 mg PO twice daily for 7 days, or a 2-g dose of tinidazole immediately. Failing that, treatment with tinidazole or metronidazole at 2 g PO for 5 days is suggested.⁴⁰

There are limited anecdotal reports of success with paromomycin cream; however, this therapy may also be associated with a high incidence of local side effects.⁴⁸ A case report of two women with resistant trichomoniasis reported cure with a combination of intravaginal paromomycin cream and high-dose oral tinidazole.⁴⁹ There is also a report of cure with prolonged intravaginal boric acid suppositories.⁵⁰ Women with asymptomatic infection should be treated. If their infection is left untreated, they may later become symptomatic and may continue to transmit the infection while untreated. Occasionally, patients manifest a true allergy to metronidazole. Because there is no effective alternative, desensitization is the only option.⁵¹

COMPLICATIONS

Long considered a minor STD, with few associated complications, infection with *T. vaginalis* has more recently been implicated as a cause of preterm delivery in several studies. In a large multicenter study, after adjusting for demographic, behavioral, and microbiologic variables, *T. vaginalis* was significantly associated with low birth weight, premature rupture of membranes, and preterm delivery (relative risk, 1.4).⁵² Minkoff and coworkers⁵³ also documented a significant correlation between trichomoniasis and premature rupture of membranes. In that study,

the incidence of this complication at term was 27.5% in women with *T. vaginalis* infection versus 12.8% in those without ($P = .03$).

Prospective studies of treatment of trichomoniasis during pregnancy for the prevention of preterm birth have yielded disappointing results. Among women with asymptomatic infection treated with metronidazole during the second and third trimesters of pregnancy, a trend toward increased preterm delivery was seen compared with the placebo group. However, the dose of metronidazole used was four times the recommended dose. In addition, the study was stopped prematurely because of a slow accrual of subjects and the trend for increased risk for preterm delivery in the treatment group.⁵⁴ A second study, conducted in Uganda, also found that treatment of trichomoniasis during pregnancy resulted in an increase in the incidence of preterm birth. However, this study was actually a subgroup analysis of a larger trial and was not properly designed to determine the effect of treatment of *T. vaginalis* during pregnancy on preterm birth⁵⁵; therefore the question remains unanswered. Since the publication of these reports, the CDC has not revised recommendations for treatment during pregnancy. Trichomoniasis has also been associated with vaginal cuff cellulitis after abdominal hysterectomy.⁵⁶

Acquisition of HIV has been associated with trichomoniasis in several African studies, possibly as a result of the local inflammation often caused by the parasite. Leroy and colleagues⁵⁷ found a significant

difference between the prevalence of trichomoniasis in a cohort of HIV-infected and uninfected pregnant women in Rwanda (20.2% and 10.9%, respectively; $P = 0.0007$). In a prospective study by Laga and associates,¹⁶ trichomoniasis positivity was significantly associated with HIV seroconversion (odds ratio, 1.9) in a multivariate analysis of a cohort of women in Zaire. Similar findings were reported from a cohort of women in Kenya.⁵⁸ The associations between HIV and trichomoniasis, as well as other STDs, may relate to the following: (1) increased shedding of HIV as a result of the local inflammation produced by the STD; (2) increased susceptibility to HIV as a result of the macroscopic or microscopic breaks in mucosal barriers caused by the STD; (3) a higher prevalence of STDs among HIV-infected individuals as a result of common risk factors for both infections; and (4) an increased susceptibility to STDs as a result of the immunosuppression associated with HIV infection. Given the higher prevalence and incidence of trichomoniasis than most other treatable STDs in most studies to date, the attributable fraction of HIV acquisitions caused by trichomoniasis may eclipse the relative contribution of other STDs.¹² The RNA concentration of HIV in the seminal fluid of men with urethritis was significantly higher in men with trichomoniasis than in those with symptomatic urethritis with an unidentified cause.⁵⁹ In addition, successful treatment of trichomonal urethritis reduced the levels of HIV RNA so that they were similar to those seen in uninfected control subjects.⁶⁰

Key References

The complete reference list is available online at Expert Consult.

- Dyall SD, Johnson PJ. Origins of hydrogenosomes and mitochondria: evolution and organelle biogenesis. *Curr Opin Microbiol.* 2000;3:404–411.
- Cu-Uvin S, Ko H, Jamieson DJ, et al. HIV Epidemiology Research Study (HERS) Group. Prevalence, incidence, and persistence or recurrence of trichomoniasis among human immunodeficiency virus (HIV)-positive women and among HIV-negative women at high risk for HIV infection. *Clin Infect Dis.* 2002;34:1406–1411.
- Wolner-Hanssen P, Krieger JN, Stevens CE, et al. Clinical manifestations of vaginal trichomoniasis. *JAMA.* 1989;261:571–576.
- Carlton JM, Hirt RP, Silva JC, et al. Draft genome sequence of the sexually transmitted pathogen *Trichomonas vaginalis*. *Science.* 2007;315:207–212.
- Sutton M, Sternberg M, Koumans EH, et al. The prevalence of *Trichomonas vaginalis* infection among reproductive-age women in the United States, 2001–2004. *Clin Infect Dis.* 2007;45:1319–1326.
- Laga M, Manoka A, Kivuvu M, et al. Non-ulcerative sexually transmitted diseases as risk factors for HIV-1 transmission in women: results from a cohort study. *AIDS.* 1993;7:95–102.
- Krieger JN, Verdon M, Siegel N, et al. Natural history of urogenital trichomoniasis in men. *J Urol.* 1993;149:1455–1458.
- Schwebke JR, Hook EW III. High rates of *Trichomonas vaginalis* among men attending a sexually transmitted diseases clinic: implications for screening and urethritis management. *J Infect Dis.* 2003;188:465–468.
- Seña AC, Miller WC, Hobbs MM, et al. *Trichomonas vaginalis* infection in male sexual partners: implications for diagnosis, treatment, and prevention. *Clin Infect Dis.* 2007;44:13–22.
- Rein MF, Muller M. *Trichomonas vaginalis* and trichomoniasis. *Sex Transm Dis.* 1990;481–492.
- Schwebke J, Rompalo A, Taylor S, et al. Re-evaluating the treatment of nongonococcal urethritis: emphasizing emerging pathogens—a randomized clinical trial. *Clin Infect Dis.* 2011;52:163–170.
- Martin DH, Zozaya M, Lillis RA, et al. Unique vaginal microbiota that includes an unknown *Mycoplasma*-like organism is associated with *Trichomonas vaginalis* infection. *J Infect Dis.* 2013;207:1922–1931.
- Kurth A, Whittington WLH, Golden MR, et al. Performance of a new, rapid assay for *Trichomonas vaginalis*. *J Clin Microbiol.* 2004;42:2940–2943.
- Schwebke J, Hobbs MM, Taylor SN, et al. Molecular testing for *Trichomonas vaginalis* in women: results from a prospective US clinical trial. *J Clin Microbiol.* 2011;49:4106–4111.
- Nye MB, Schwebke JR, Body BA. Comparison of APTIMA *Trichomonas vaginalis* transcription-mediated amplification to wet mount microscopy, culture and PCR for diagnosis of trichomoniasis in men and women. *Am J Obstet Gynecol.* 2009;200:188.e1–188.e7.
- Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines. *MMWR Recomm Rep.* 2006;55(RR-11):49–56.
- Paulish-Miller TE, Augostini P, Schuyler JA, et al. *Trichomonas vaginalis* metronidazole resistance is associated with single nucleotide polymorphisms in the nitroreductase genes *ntf4Tv* and *ntf6Tv*. *Antimicrob Agents Chemother.* 2014;58:2938–2943.
- Crowell A, Sanders-Lewis KA, Secor WE. In vitro metronidazole and tinidazole activities against metronidazole-resistant strains of *Trichomonas vaginalis*. *Antimicrob Agents Chemother.* 2003;47:1407–1409.
- Sobel J, Nyirjesy P, Brown W. Tinidazole therapy for metronidazole-resistant vaginal trichomoniasis. *Clin Infect Dis.* 2001;33:1341–1346.
- Sobel J, Nagappan V, Nyirjesy P. Metronidazole-resistant vaginal trichomoniasis—an emerging problem. *N Engl J Med.* 1999;341:292–293.
- Klebanoff M, Carey J, Hauth J, et al. Failure of metronidazole to prevent preterm delivery among pregnant women with asymptomatic *Trichomonas vaginalis* infection. *N Engl J Med.* 2001;345:487–493.
- McClelland R, Sangare L, Hassan W, et al. Infection with *Trichomonas vaginalis* increases the risk of HIV-1 acquisition. *J Infect Dis.* 2007;195:698–702.
- Price M, Zimba D, Hoffman IF, et al. Addition of treatment for trichomoniasis to syndromic management of urethritis in Malawi: a randomized clinical trial. *Sex Transm Dis.* 2003;30:516–522.

References

- Honigberg B, King VM. Structure of *Trichomonas vaginalis* Donne. *J Parasitol*. 1964;50:345–364.
- Dyall SD, Johnson PJ. Origins of hydrogenosomes and mitochondria: evolution and organelle biogenesis. *Curr Opin Microbiol*. 2000;3:404–411.
- Burgess D, Knoblock KF. Identification of *Trichomonas foetus* in sections of bovine placental tissue with monoclonal antibodies. *J Parasitol*. 1989;75:977–980.
- Phillip A, Carter-Scott P, Rogers C. An agar culture technique to quantitate *Trichomonas vaginalis* from women. *J Infect Dis*. 1987;155:304–308.
- Arroyo R, Engbring J, Alderete J. Molecular basis of host epithelial cell recognition by *Trichomonas vaginalis*. *Mol Microbiol*. 1992;6:853–862.
- Niccolai L, Kopicko JJ, Kassie A, et al. Incidence and predictors of reinfection with *Trichomonas vaginalis* in HIV-infected women. *Sex Transm Dis*. 2000;27:284–288.
- Honigberg B, Burgess DE. Trichomonads of importance in human medicine including *Dientamoeba fragilis*. In: Kreier JP, ed. *Parasitic Protozoa*. Vol. 9. New York: Academic Press; 1994:1–109.
- Mason P, Patterson BA. Proliferative response of human lymphocytes to secretory and cellular antigens of *Trichomonas vaginalis*. *J Parasitol*. 1985;71:265–268.
- Cu-Uvin S, Ko H, Jamieson DJ, et al; HIV Epidemiology Research Study (HERS) Group. Prevalence, incidence, and persistence or recurrence of trichomoniasis among human immunodeficiency virus (HIV)-positive women and among HIV-negative women at high risk for HIV infection. *Clin Infect Dis*. 2002;34:1406–1411.
- Wolner-Hanssen P, Krieger JN, Stevens CE, et al. Clinical manifestations of vaginal trichomoniasis. *JAMA*. 1989;261:571–576.
- Carlton JM, Hirt RP, Silva JC, et al. Draft genome sequence of the sexually transmitted pathogen *Trichomonas vaginalis*. *Science*. 2007;315:207–212.
- Sorvillo F, Kerndt P. *Trichomonas vaginalis* and amplification of HIV-1 transmission. *Lancet*. 1998;351:213–214.
- Sutton M, Sternberg M, Koumans EH, et al. The prevalence of *Trichomonas vaginalis* infection among reproductive-age women in the United States, 2001–2004. *Clin Infect Dis*. 2007;45:1319–1326.
- Weinstock H, Berman S, Cates W. Sexually transmitted diseases among American youth: incidence and prevalence estimates, 2000. *Perspect Sex Reprod Health*. 2004;36:6–10.
- Cates W. American Social Health Association Panel. Estimates of the incidence and prevalence of sexually transmitted diseases in the United States. *Sex Transm Dis*. 1999;26:S2–S7.
- Laga M, Manoka A, Kivuvu M, et al. Non-ulcerative sexually transmitted diseases as risk factors for HIV-1 transmission in women: results from a cohort study. *AIDS*. 1993;7:95–102.
- Lossick J. Epidemiology of urogenital trichomoniasis. In: Honigberg BM, ed. *Trichomonads Parasitic in Humans*. New York: Springer-Verlag; 1989:311–341.
- Wilcox R. Epidemiological aspects of human trichomoniasis. *Br J Vener Dis*. 1960;36:167–174.
- Krieger JN, Verdon M, Siegel N, et al. Natural history of urogenital trichomoniasis in men. *J Urol*. 1993;149:1455–1458.
- Joyner J, Douglas J, Ragsdale S, et al. Comparative prevalence of infection with *Trichomonas vaginalis* among men attending a sexually transmitted diseases clinic. *Sex Transm Dis*. 2000;27:236–240.
- Schwebke JR, Hook EW III. High rates of *Trichomonas vaginalis* among men attending a sexually transmitted diseases clinic: implications for screening and urethritis management. *J Infect Dis*. 2003;188:465–468.
- Seña AC, Miller WC, Hobbs MM, et al. *Trichomonas vaginalis* infection in male sexual partners: implications for diagnosis, treatment, and prevention. *Clin Infect Dis*. 2007;44:13–22.
- Danesh IS, Stephen JM, Gorbach J. Neonatal *Trichomonas vaginalis* infection. *J Emerg Med*. 1995;13:51–54.
- al-Salihi FL, Curran JP, Wang J. Neonatal *Trichomonas vaginalis*: report of three cases and review of the literature. *Pediatrics*. 1974;53:196–200.
- Jones JG, Yamaguchi T, Lambert B. *Trichomonas vaginalis* infestations in sexually abused girls. *Am J Dis Child*. 1985;139:846–847.
- Hesseltine H. Experimental human vaginal trichomoniasis. *J Infect Dis*. 1942;71:127.
- Rein MF, Muller M. *Trichomonas vaginalis* and trichomoniasis. *Sex Transm Dis*. 1990;481–492.
- Fouts AC, Kraus SJ. *Trichomonas vaginalis* reevaluation of its clinical presentation and laboratory diagnosis. *J Infect Dis*. 1980;177:167–174.
- Krieger JN. Trichomoniasis in men: old issues and new data. *Sex Transm Dis*. 1995;22:83–96.
- Schwebke J, Rompalo A, Taylor S, et al. Re-evaluating the treatment of nongonococcal urethritis: emphasizing emerging pathogens—a randomized clinical trial. *Clin Infect Dis*. 2011;52:163–170.
- Krieger JN, Tam MR, Stevens CE, et al. Diagnosis of trichomoniasis: comparison of conventional wet-mount examination with cytologic studies, cultures, and monoclonal antibody staining of direct specimens. *JAMA*. 1988;259:1223–1227.
- Kingston MA, Bansal D, Carlin EM. ‘Shelf life’ of *Trichomonas vaginalis*. *Int J STD AIDS*. 2003;14:28–29.
- Martin DH, Zozaya M, Lillis RA, et al. Unique vaginal microbiota that includes an unknown *Mycoplasma*-like organism is associated with *Trichomonas vaginalis* infection. *J Infect Dis*. 2013;207:1922–1931.
- Draper D, Parker R, Patterson E, et al. Detection of *Trichomonas vaginalis* in pregnant women with the InPouch TV system. *J Clin Microbiol*. 1993;31:1016–1018.
- Briselden AM, Hillier SL. Evaluation of Affirm VP microbial identification test for *Gardnerella vaginalis* and *Trichomonas vaginalis*. *J Clin Microbiol*. 1994;32:148–152.
- Kurth A, Whittington WLH, Golden MR, et al. Performance of a new, rapid assay for *Trichomonas vaginalis*. *J Clin Microbiol*. 2004;42:2940–2943.
- Schwebke J, Hobbs MM, Taylor SN, et al. Molecular testing for *Trichomonas vaginalis* in women: results from a prospective US clinical trial. *J Clin Microbiol*. 2011;49:4106–4111.
- Van Der Pol B. Clinical and Laboratory Testing for *Trichomonas vaginalis* Infection. *J Clin Microbiol*. 2016;54:7–12.
- Nye MB, Schwebke JR, Body BA. Comparison of APTIMA *Trichomonas vaginalis* transcription-mediated amplification to wet mount microscopy, culture and PCR for diagnosis of trichomoniasis in men and women. *Am J Obstet Gynecol*. 2009;200:188.e1–188.e7.
- Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines. *MMWR Recomm Rep*. 2006;55(RR-11):49–56.
- Roe F. Safety of nitroimidazoles. *Scand J Infect Dis Suppl*. 1985;46:72–81.
- Wood B, Monroe A. Pharmacokinetics of tinidazole and metronidazole after single large oral doses. *Br J Vener Dis*. 1975;51:51–53.
- Schwebke JR, Barrientes FJ. Prevalence of *Trichomonas vaginalis* isolates with resistance to metronidazole and tinidazole. *Antimicrob Agents Chemother*. 2006;50:4209–4210.
- Land KM, Delgadillo-Correa MG, Tachezy J, et al. Targeted gene replacement of a ferredoxin gene in *Trichomonas vaginalis* does not lead to metronidazole resistance. *Mol Microbiol*. 2003;51:115–122.
- Paulish-Miller TE, Augustini P, Schuyler JA, et al. *Trichomonas vaginalis* metronidazole resistance is associated with single nucleotide polymorphisms in the nitroreductase genes *ntf4Tv* and *ntf6Tv*. *Antimicrob Agents Chemother*. 2014;58:2938–2943.
- Crowell A, Sanders-Lewis KA, Secor WE. In vitro metronidazole and tinidazole activities against metronidazole-resistant strains of *Trichomonas vaginalis*. *Antimicrob Agents Chemother*. 2003;47:1407–1409.
- Sobel J, Nyirjesy P, Brown W. Tinidazole therapy for metronidazole-resistant vaginal trichomoniasis. *Clin Infect Dis*. 2001;33:1341–1346.
- Sobel J, Nagappan V, Nyirjesy P. Metronidazole-resistant vaginal trichomoniasis—an emerging problem. *N Engl J Med*. 1999;341:292–293.
- Nyirjesy P, Gilbert J, Mulcahy LJ. Resistant trichomoniasis: successful treatment with combination therapy. *Sex Transm Dis*. 2011;38:962–963.
- Muzny C, Barnes A, Mena L. Symptomatic *Trichomonas vaginalis* infection in the setting of severe nitroimidazole allergy: successful treatment with boric acid. *Sex Health*. 2012;9:389–391.
- Pearlman M, Yashar C, Ernst S, et al. An incremental dosing protocol for women with severe vaginal trichomoniasis and adverse reactions to metronidazole. *Am J Obstet Gynecol*. 1996;174:934–936.
- Cotch MF, Pastorek JG, Nugent RP, et al. *Trichomonas vaginalis* associated with low birth weight and preterm delivery. *Sex Transm Dis*. 1997;24:361–362.
- Minkoff H, Grunebaum AN, Schwarz RH, et al. Risk factors for prematurity and premature rupture of membranes: a prospective study of the vaginal flora in pregnancy. *Am J Obstet Gynecol*. 1984;150:965–972.
- Klebanoff M, Carey J, Hauth J, et al. Failure of metronidazole to prevent preterm delivery among pregnant women with asymptomatic *Trichomonas vaginalis* infection. *N Engl J Med*. 2001;345:487–493.
- Kigozi G, Brahmabhatt H, Wabwire-Mangen F, et al. Treatment of *Trichomonas* in pregnancy and adverse outcomes of pregnancy: a subanalysis of a randomized trial in Rakai, Uganda. *Am J Obstet Gynecol*. 2003;189:1398–1400.
- Soper DE, Bump RC, Hurt WG. Bacterial vaginosis and trichomoniasis vaginitis are risk factors for cuff cellulitis after abdominal hysterectomy. *Am J Obstet Gynecol*. 1990;163:1016–1023.
- Leroy V, De Clercq A, Ladner J, et al. Should screening of genital infections be part of antenatal care in areas of high HIV prevalence? A prospective cohort study from Kigali, Rwanda, 1992–1993. *Genitourin Med*. 1995;71:207–211.
- McClelland R, Sangare L, Hassan W, et al. Infection with *Trichomonas vaginalis* increases the risk of HIV-1 acquisition. *J Infect Dis*. 2007;195:698–702.
- Hobbs M, Kazembe P, Reed A, et al. *Trichomonas vaginalis* as a cause of urethritis in Malawian men. *Sex Transm Dis*. 1999;26:381–387.
- Price M, Zimba D, Hoffman IF, et al. Addition of treatment for trichomoniasis to syndromic management of urethritis in Malawi: a randomized clinical trial. *Sex Transm Dis*. 2003;30:516–522.

SHORT VIEW SUMMARY**Definition**

- Babesiosis is an emerging infectious disease caused by hemoprotozoan parasites of the genus *Babesia*.

Epidemiology and Microbiology

- Most cases of human babesiosis occur in the United States, particularly the Northeast and the upper Midwest, and are caused by *Babesia microti*. *B. microti* is primarily transmitted during the bite of an *Ixodes scapularis* nymphal tick but also by transfusion of contaminated blood products, predominantly packed red blood cells. *B. microti* has been the pathogen most frequently implicated by investigations of transfused-related illnesses in the United States and accounted for one-fourth of transfusion-related deaths caused by microbial infection from fiscal year 2010 to fiscal year 2016 (see Chapter 304).
- Human babesiosis also occurs in Europe and East Asia, where the major etiologic agents are *Babesia divergens* and *Babesia venatorum*, respectively.
- Risk factors for severe babesiosis include anatomic or functional asplenia, low CD4⁺ T-cell counts associated with human immunodeficiency virus/acquired immunodeficiency syndrome or therapy for transplantation, and impaired antibody production associated with chronic immunosuppressive therapy.

Diagnosis

- Fever is the salient symptom, and often is accompanied by chills, sweats, headache, anorexia, and/or myalgia.
- Anemia is the major laboratory finding. Severe anemia often is preceded by severe thrombocytopenia. Elevated levels of lactate dehydrogenase and total bilirubin are consistent with hemolytic anemia.
- The definitive diagnosis is made on Giemsa-stained thin blood smears. Trophozoites often appear as rings. Tetrads of merozoites are pathognomonic.
- If parasites are not visualized on smear but babesiosis remains suspected, polymerase chain reaction assay–based amplification of parasite DNA should be performed.

Therapy (See Table 281.1)

- Mild *B. microti* illness should be treated with a single 7- to 10-day course of oral atovaquone (750 mg every 12 hours) *plus* oral azithromycin (500 mg on day 1, then 250 mg from day 2 onward). Symptoms typically abate within 48 hours and resolve within 2 weeks.
- Severe *B. microti* illness requires hospital admission, and can be complicated by pulmonary edema and respiratory distress. Other complications include congestive heart failure, renal failure, disseminated intravascular coagulation, shock, and splenic rupture. Initial

therapy should consist of a 7- to 10-day course of oral atovaquone (750 mg every 12 hours) *plus* intravenous azithromycin (500 mg daily). If symptoms do not abate, therapy should be extended as needed. If microbial resistance is suspected, intravenous clindamycin (600 mg every 6 hours) *plus* oral quinine (650 mg every 8 hours) or oral atovaquone is an alternative.

- If the patient is severely immunocompromised, antimicrobial therapy should be administered for at least 6 consecutive weeks, including 2 final weeks during which parasites are no longer detected on blood smear.
- Partial or complete red blood cell exchange transfusion is recommended for cases with high-grade parasitemia ($\geq 10\%$), severe anemia (hemoglobin < 10 g/dL) in the context of parasitemia, or organ (pulmonary or renal) compromise.

Prevention

- No vaccine is available. Individuals at risk of severe babesiosis should avoid tick exposure in endemic areas. Protective measures include covering exposed skin, applying an acaricide to skin, using permethrin-impregnated clothes, and thoroughly examining the skin for ticks.

The first recorded mention of babesiosis is believed to date to biblical times. In the Book of Exodus, the fifth plague is described as “very grievous murrain” that fell “upon thy cattle which is in the field, upon the horses, upon the asses, upon the camels, upon the oxen and upon the sheep.” Bovine babesiosis is still referred to as murrain in the Irish countryside. In 1888, the microbiologist and pathologist Victor Babes attributed the febrile hemoglobinuria and death of cattle in Romania to an intraerythrocytic microorganism.¹ Soon after, Theobald Smith and Frederick Kilborne described a similar piroplasm (*L. pirum*, “pear”) in the erythrocytes of Texas cattle with fever. Initially named *Pyrosoma*, this organism became known as *Babesia bigemina*. Smith and Kilborne also identified the tick *Rhipicephalus annulatus* as its vector, establishing for the first time that hematophagous arthropods can transmit an infectious agent to a vertebrate host. The first well-documented case of human babesiosis was reported in 1957 in a 33-year-old splenectomized herdsman who had been grazing cattle on tick-infested pastures near Zagreb, Croatia. The infection was fulminant and fatal. Originally identified as *Babesia bovis*, the causative agent was later reported to be *Babesia divergens*, a parasite of cattle. In 1969, a 59-year-old woman who had summered on Nantucket Island off the coast of Massachusetts presented with a history of fever, headache, and crampy abdominal pain. This patient had an intact spleen, and the causative agent was

Babesia microti, a parasite of white-footed mice. Andrew Spielman and colleagues identified the vector as the deer tick *Ixodes dammini* (now recognized as *Ixodes scapularis*). As the number of cases on the island grew, the disease became known as “Nantucket fever.” Babesiosis caused by *B. microti* is now recognized as endemic on the mainland of the United States, particularly in the Northeast and the upper Midwest.

EPIDEMIOLOGY**United States: *Babesia microti* Geographic Distribution**

Human babesiosis became a nationally notifiable disease in January 2011.² Notification is made via the National Notifiable Diseases Surveillance System using the standard case definition developed by the Centers for Disease Control and Prevention (CDC) and the Council of State and Territorial Epidemiologists. In 2014, babesiosis was reportable from 31 states, of which 22 notified the CDC of at least one case.³ Most cases (>90%) are reported from the seven states with well-established foci of zoonotic transmission (Connecticut, Massachusetts, Minnesota, New Jersey, New York, Rhode Island, Wisconsin). In the Northeast, areas of high endemicity include islands off the southern coast of New England.⁴ On the mainland, babesiosis is highly endemic in southeastern Massachusetts, including Cape Cod, the coastal counties of western




FIG. 281.1 Geographic distribution of human babesiosis and its tick vectors. Dark colors indicate states or provinces where human babesiosis is endemic or sporadic (defined by ≥ 5 cases), whereas light colors indicate areas where tick vectors are present but human babesiosis is rare (< 5 cases), undocumented, or absent. Circles depict single cases, except in three locations (Mexico, Montenegro, and eastern Poland) where all patients were diagnosed at the same hospital. Colors distinguish the etiologic agents: red for *Babesia microti*, orange for *Babesia duncani*, blue for *Babesia divergens*, green for *Babesia venatorum*, pink for *Babesia crassa*, black for KO1, and yellow for *Babesia* sp. XXB/HangZhou. White circles depict cases caused by *Babesia* isolates that were not characterized at the molecular level. Asymptomatic infections and cases of travel-associated babesiosis are omitted. *Ixodes ricinus* and *Ixodes persulcatus* are sympatric in southern Finland, Estonia, Latvia, and northwestern Russia. The island of Hokkaido in northern Japan is enzootic for both *I. persulcatus* and *Ixodes ovatus*. (From Vannier E, Krause X. In: Ryan ET, Hill DR, Solomon T, et al, eds. Hunter's Tropical Medicine and Emerging Infectious Diseases. 10th ed. Oxford, UK: Elsevier; 2019.)

Rhode Island and eastern Connecticut, the counties east of the Hudson River in the Lower Hudson Valley (New York), and the south central counties of New Jersey. In the upper Midwest, babesiosis caused by *B. microti* is moderately endemic in Minnesota and Wisconsin (Fig. 281.1).

Incidence and Prevalence

The incidence of babesiosis has steadily increased over the past 30 years. In New York State, the first state to mandate the reporting of babesiosis, more than 5800 cases have been notified to the health authorities. About 1100 cases were reported from 1986 to 2005, and more than 3500 cases in the following decade. The sharp increase noted between 2006 and 2015 resulted from the emergence of babesiosis in the Lower Hudson Valley combined with the maintenance of highly endemic foci on eastern Long Island.^{5,6} Along the northeastern seaboard, babesiosis has emerged in areas that were once at the periphery of the “historical heartland,” namely southern Maine and southern New Hampshire to the north and Pennsylvania, Delaware, and Maryland to the south.^{3,4} In the upper Midwest, the number of annual cases has risen steadily since 2005.⁷ A few cases have been reported in Illinois, Indiana, and Michigan.³ Babesiosis caused by *B. microti* remains an emerging infectious disease in the United States.^{1,3,4}

The seroprevalence of antibodies against *B. microti* antigen varies by year, study site, and sample population but has consistently been high in endemic areas, such as Nantucket (7% among blood donors),⁸ Block Island (9% among residents),⁹ southern Connecticut (0.5%–9% among blood donors),^{9,10} Shelter Island (4% in blood donors; 4%–7% among residents),¹¹ and eastern Long Island (16% among residents).¹² In Minnesota, seroprevalence among blood donors has been reported at 2%.¹³ Given that seroprevalence is much higher than prevalence of clinical babesiosis, asymptomatic infection is more common than recognized or reported.

Modes of Transmission

Tick Bite

B. microti is primarily acquired during the blood meal of the tick *I. scapularis*,¹⁴ but only one-half of patients recall a tick bite in the 8 weeks prior to symptom onset.³ The nymphal stage, the primary vector, is most active from late spring to early summer (see “Microbiology” later).¹⁴ Because the latency period typically lasts from 1 to 4 weeks, one-fifth of cases are diagnosed in June, one-half in July, and another one-fifth in August.³ Some cases present as early as late spring. Patients who

present in late summer or early fall likely acquired the infection from an adult female tick.

Blood Transfusion

B. microti can be transmitted via transfusion of blood components prepared from contaminated blood donated by asymptomatic carriers or symptomatic individuals who did not suspect the infection.¹⁵ Implicated blood donors typically have engaged in outdoor activities.¹⁶ Less than one-fifth have experienced fatigue or fever within 24 months prior to index donation.¹⁶ Even fewer (8%) have experienced chills. Consistent with the seasonality of tick-borne babesiosis, three-fourths of transfusion-transmitted babesiosis (TTB) cases are diagnosed from June through November, with one-fifth in August alone.¹⁷ Given that asymptomatic infection can persist in untreated individuals for longer than a year,^{16,18} TTB is diagnosed year-round.¹⁷ Most TTB cases (87%) occur in residents of highly endemic states (see earlier list).¹⁷ Outside endemic areas, TTB typically involves contaminated blood products that are imported from endemic areas or are derived from blood donations by individuals following their return from endemic areas.

More than 250 TTB cases caused by *B. microti* have been identified since the first case was reported in 1979; three-fourths have occurred since 2005.^{16,17,19} Most cases involve red blood cell (RBC) units. A few have been attributed to whole-blood-derived platelets contaminated with residual RBCs. At time of transfusion, the age of liquid-stored RBC units has ranged from 4 to 42 days; one platelet unit was 5 days old.^{16,17,20} In March 2018, when the US Food and Drug Administration (FDA) approved two tests for donor blood screening (see next paragraph), *B. microti* was still the pathogen most frequently implicated by investigations of transfusion-related illnesses.¹⁹ From fiscal year 2010 to fiscal year 2016, *B. microti* accounted for one-fourth of transfusion-related deaths caused by microbial infection.²¹

Using an arrayed fluorescence immunoassay for detection of *B. microti* immunoglobulin G (IgG) and a real-time polymerase chain reaction (PCR) for detection of *B. microti* DNA, the risk posed by blood donations in some highly endemic areas (Massachusetts and Connecticut) has been carefully assessed.¹⁶ Among donations collected from June 2012 through September 2014, 0.3% reacted in one test or both. Three-quarters of the donations were positive for *B. microti* IgG but negative for *B. microti* DNA, whereas one-fifth were positive for both. Antibody-positive donations were obtained throughout the year. Consistent with the seasonality of tick-borne babesiosis, donations that were DNA positive but IgG

negative were obtained from June through September. Among donors who tested positive for *B. microti* DNA at time of index donation, most (86%) were no longer infected a year later. The median time to parasite DNA clearance was 4.7 months. Among those who tested positive for *B. microti* IgG, only 8% had seroreverted a year later. The median time to seroreversion was 17.1 months. None of the 75,000+ donations screened with the two FDA-approved tests were implicated in a TTB case. In contrast, 1 in 18,000 unscreened donations obtained from the same area was implicated in a TTB case. When five highly endemic states (Connecticut, Massachusetts, Minnesota, New Jersey, New York) were analyzed along with two states that recently emerged as endemic (Maine and New Hampshire), the risk for TTB was estimated at 1 case per 100,000 donations for the period from January 1, 2010 to August 31, 2016.

Transplacental Passage

Unequivocal evidence of transplacental transmission came from a case reported in 2012.²² A splenectomized mother was admitted after a 10-day history of fever and rigors. Intraerythrocytic *B. microti* ring forms were identified on a blood smear and in the amniotic fluid. On the second day of hospitalization, the mother underwent cesarean section. Immediately after delivery, blood was obtained from the neonate and intraerythrocytic parasites were noted on a thin blood film.

Probable congenital babesiosis has been reported for at least nine other cases.^{23–26} For seven cases, mothers were asymptomatic throughout pregnancy. In one such case, *B. microti* likely was acquired during the first trimester of pregnancy.²⁵ Two symptomatic mothers received a diagnosis of Lyme disease during the last trimester of pregnancy, but *B. microti* infection was not suspected.²⁴ Symptoms of babesiosis developed in the neonates during the third to the sixth week of life and often consisted of fever accompanied by pallor.²⁴ Laboratory tests revealed hemolytic anemia, thrombocytopenia, and neutropenia. In all cases but one, anemia was so severe that a blood transfusion was required. At diagnosis, parasitemia ranged from 2% to 40%. When tested for *B. microti* antibody, neonates and their mothers were seropositive. One neonate who presented with tachycardia and respiratory distress required an exchange transfusion, which was performed manually.²⁶ This neonate had a twin who was seropositive for *B. microti* but remained healthy. In another case, *B. microti* antibody was detected in a heel stick blood sample obtained on the third day of life, raising the possibility that maternal IgG directed against *B. microti* protects newborns from congenital babesiosis very early in life.²³ In addition to vertical transmission, babesiosis has been diagnosed in neonates following blood transfusion.

Solid Organ Transplantation

Two cases of transplantation-associated babesiosis have been described.²⁷ Both men had received a kidney allograft obtained from the same deceased donor and began experiencing symptoms 5 weeks posttransplantation. The diagnosis of babesiosis was made 8 weeks posttransplantation. Despite immunosuppressive therapy, both men tested seropositive for *B. microti*. Pretransplantation serum samples did not react with *B. microti* antigen, an observation consistent with the denial by these allograft recipients of tick exposure and blood transfusion prior to transplantation. The organ donor, however, had received multiple blood transfusions, including one from a blood donor who tested positive for *B. microti* antibody during the retrospective investigation. It is presumed that some *B. microti* parasites from the blood donor had remained in the vasculature/fluids of the renal allografts at time of transplantation. In support of this explanation, corneas from the deceased organ donor were transplanted but neither recipient presented clinical or serologic evidence of *B. microti* infection.

Risk Factors

Babesiosis caused by *B. microti* often is severe in individuals with one or several of the following risk factors: asplenia,^{28–30} X-linked agammaglobulinemia,²⁹ malignancy,³¹ low CD4⁺ T-cell counts associated with human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS),^{32,33} blockade of tumor necrosis factor- α (TNF- α) for chronic inflammatory disorders such as inflammatory bowel disease,³⁴ and immunosuppressive therapy for transplantation,^{35,36} B-cell lymphoma, or an autoimmune disorder.^{29,32} These risk factors also pertain to the transmission of *B. microti* by blood transfusion. In this setting, given

that most cases have significant comorbidity, risk factors for TTB include conditions that require blood transfusion, such as hematologic disorders, cardiovascular surgery or procedure, gastrointestinal disease, bleeding and surgery, and anemia of prematurity.¹⁷ Two-thirds of TTB patients are older than age 50 years, but two-thirds of those younger than age 40 years have a hereditary RBC disorder.¹⁷

United States: Other *Babesia* Species

Babesia duncani and *B. duncani*-type organisms are the etiologic agents of human babesiosis along the Pacific Coast.¹ The index case (WA1) occurred in a 41-year-old normosplenic man from a forested area of south central Washington State.³⁷ The four cases in California (CA1–CA4) developed in splenectomized men in their 20s to 40s who lived or had outdoor exposure in areas ranging from Sonoma County to San Bernardino County.³⁸ The first case of TTB caused by *B. duncani* involved a 76-year-old normosplenic man who had received RBCs donated by a 34-year-old Washington State resident.³⁹ The second case was diagnosed in a premature male infant who had received two blood transfusions.⁴⁰ The implicated donor lived in the San Francisco Bay area and had vacationed in a rural area of central Oregon 3 months prior to donating. The third case involved a 59-year-old patient who had sickle cell disease.⁴¹ The 67-year-old donor reported a possible tick bite during a hike in the San Francisco Bay area. The winter tick *Dermacentor albipictus* presumably is the vector for transmission of *B. duncani*.^{41a} The prevalence of *B. duncani* infection remains unclear because clinical cases have been too few to validate the serologic test.

B. divergens-like organisms have caused disease in Arkansas, Kentucky, Michigan, Missouri, and Washington State.^{42–45} All six patients were older than 50 years of age, and asplenic. In one case, transfusion is the suspected mode of transmission.⁴³ The patient lived in Arkansas but blood donated in Missouri was the likely culprit, thereby bringing to three the number of *B. divergens*-like infections that originated in Missouri. Isolates from the five cases that originated in the Midwest were identical to piroplasms found in eastern cottontail rabbits, implying that *Ixodes dentatus* may be a vector for zoonotic transmission of *B. divergens*-like organisms. In Tennessee, three cases were uncovered by querying claims data from a managed care organization.⁴⁶ One such case may have been caused by a *B. divergens*-like organism.

Europe: *Babesia divergens*

Approximately 40 cases of babesiosis have been attributed to *B. divergens*, a parasite of cattle that is transmitted by *Ixodes ricinus*.⁴⁷ Most cases occurred in France and Ireland, particularly in regions with extensive cattle farming. Isolated cases have been reported from Portugal, Spain, Norway, Sweden, Finland, Georgia (ex-USSR), Turkey, and Croatia (including the index case). Nearly all patients experienced a severe illness and had been splenectomized,⁴⁷ although one had a rudimentary spleen and another had functional hyposplenism caused by celiac disease.^{48,49} TTB caused by *B. divergens* has not been reported from Europe, although asymptomatic *B. divergens* infection is not uncommon, as indicated by seroprevalence in Slovenia (4.1% of forestry workers), western Austria (2.1% of blood donors), midwestern Germany (4.2% of individuals with clinical or serologic evidence of Lyme borreliosis), and southern Sweden (7.0% of individuals seropositive for *Borrelia burgdorferi* s.l.).^{50–51a}

Europe: Other *Babesia* Species

Cases of *B. venatorum* infection have been reported in Italy, Austria, Germany, and Sweden.^{52–54} The five patients were splenectomized men older than 50 years of age. Roe deer are a reservoir host for *B. venatorum* in Europe, and the bite of an *I. ricinus* tick is the suspected mode of transmission. The prevalence of asymptomatic infection remains unknown because *B. venatorum* sera cross-react with *B. divergens* antigen. Attributed to *B. divergens* based on morphology and seroreactivity, some early cases of babesiosis in Europe may have been caused by *B. venatorum*.

The first case of *B. microti* infection native to Europe was reported in 2007 from Germany.⁵⁵ A 42-year-old woman presented with fever and chest pain 1 week after chemotherapy for acute myeloid leukemia. Intraerythrocytic parasites were observed on Giemsa-stained blood smears and identified as US-type *B. microti*. The platelet concentrate transfused 6 days before the diagnosis of babesiosis had been prepared

from blood donated by an individual who subsequently tested seropositive for *B. microti*. Additional cases were reported in 2016 from northeastern Poland.⁵⁶ All six patients were immunocompetent and had been bitten by a tick within 8 weeks prior to presentation. No parasites were seen on blood smear, but a fragment of the *B. microti* 18S ribosomal RNA (rRNA) gene was amplified and sequenced, establishing for the first time that *B. microti* organisms of the Munich lineage can cause disease in humans. Cases of travel-associated babesiosis have been reported throughout Europe and explained by a recent stay in the northeastern United States.⁴⁷ Asymptomatic *B. microti* infection is prevalent in residents of rural, forested areas across Europe as indicated by serosurveys conducted in northeastern Poland (4.4% of foresters), eastern Switzerland (1.3% of blood donors), midwestern Germany (8.3% of individuals with clinical or serologic evidence of Lyme borreliosis), eastern France (2% of foresters who are seropositive for other tick-borne pathogens), and Belgium (9% of patients presenting within 1 month after a tick bite).^{50,57–60}

Asia: *Babesia venatorum*

The first case of *B. venatorum* infection in Asia was documented in northwestern China.⁶¹ A series of 48 cases was subsequently reported from northeastern China.⁶² All 48 cases reported a tick bite within 2 months prior to seeking medical care; none had received a blood transfusion within 2 years. All had a spleen, but only one-third were older than 50 years. A survey of ticks revealed that the taiga tick *Ixodes persulcatus* is the likely vector for *B. venatorum* in northeastern China.

Asia: Other *Babesia* Species

The first two cases of *B. microti* infection in Asia were documented in Taiwan.⁶³ A case of TTB was subsequently reported from Kobe, Japan.⁶⁴ The isolate has defined the Kobe lineage, one of the four lineages within the *B. microti* species complex.⁶⁵ The asymptomatic blood donor resided on the nearby Awaji Island, where the field mouse *Apodemus speciosus* was identified as a reservoir host for Kobe-type organisms.⁶⁶ The vector is unknown. *B. microti* organisms of the Hobetsu lineage are found in *A. speciosus* mice throughout Japan, and can be transmitted by *Ixodes ovatus* ticks.⁶⁷ Hobetsu-type organisms have infected humans, as revealed by a serosurvey, but no cases of babesiosis have been reported. In mainland China, *B. microti* has caused disease in the southern provinces, particularly in Yunnan along the border with Myanmar.^{68,69} At least two cases were caused by US-type organisms. One case was coinfecting with *Plasmodium falciparum* and another with *Plasmodium vivax*.⁶⁸

In South Korea, a 75-year-old splenectomized woman who presented with fever and severe anemia was infected with a large *Babesia* species (KO1) that is closely related to organisms found in sheep. A series of 31 cases caused by *Babesia crassa*, a species found in sheep, has been documented in northeastern China.⁷⁰ All cases were mild to moderate; none were asplenic.

Rest of the World

A case of US-type *B. microti* has been documented in Australia.¹ On the American continent, a case of *B. microti* infection was identified in Manitoba, Canada, whereas three cases were diagnosed on the Yucatan peninsula, Mexico. Asymptomatic *B. microti* infection has been documented in Bolivia and Brazil. Cases caused by *Babesia* spp. that infect cattle (*B. bigemina*, *B. bovis*) have been reported from Colombia. Asymptomatic infections with these two *Babesia* spp. have been reported from Brazil, Colombia, and Cuba. On the African continent, three cases have been reported from Egypt, of which one was acquired from a pet dog. A case of *B. divergens*-like infection was diagnosed in a splenectomized resident of the Canary Islands. In South Africa, babesiosis was diagnosed in two patients upon their return from Namibia and Zimbabwe. Isolated cases of *B. bovis* infection, of which two were fatal, have been documented in Mozambique.

MICROBIOLOGY

Babesia spp. are protozoan parasites of the phylum Apicomplexa, class Aconoidasida, and order Piroplasmida. More than 100 *Babesia* spp. are known to infect vertebrates, including mammals and birds. In the northern hemisphere, most *Babesia* spp. are maintained in their enzootic cycle by hard-bodied ticks of the genus *Ixodes*. In the United States, *B. microti* is maintained by *I. scapularis* ticks.

The Enzootic Cycle of *Babesia microti* Ticks and Reservoir Hosts

The life cycle of *I. scapularis* has three active stages (larva, nymph, adult) and requires 2 years for completion.^{1,14} In the fall, adult ticks feed primarily on white-tailed deer (*Odocoileus virginianus*). Deer are incompetent hosts for *B. microti* but are essential for the mating of *I. scapularis* and therefore its maintenance in the life cycle. Adult female ticks overwinter in an engorged state and lay eggs in the spring. Because *B. microti* does not reach the ovaries of adult ticks, eggs are free of *B. microti* (no transovarial transmission). Larvae hatch in late July and become infected as they feed in late summer on a wide range of small mammals. The primary reservoir host for *B. microti* is the white-footed mouse (*Peromyscus leucopus*).¹⁴ Other competent reservoirs include short-tailed shrews, eastern chipmunks, and raccoons.⁷¹ Passerines, raptorial birds, and ground-dwelling birds (e.g., wood thrush, veery) are reservoirs but of lesser competence. Fed to repletion, larvae overwinter and molt into nymphs in late spring. If the larva is infected, so can be the nymph (transstadial transmission). Nymphs feed on *P. leucopus* and other vertebrates from May through July. Feeding of nymphs in early summer ensures that *P. leucopus* mice are reservoirs of *B. microti* when larvae feed in late summer. In the fall, nymphs molt into adults.

Parasite Acquisition by Ticks

Soon after attachment of a larva or a nymph, host erythrocytes accumulate in its gut.^{1,14} *Babesia microti* pregametocytes mature into gametocytes. Microtubules accumulate at the anterior end of the parasite to form a raylike structure that contributes to the fusion of gametocytes into a zygote. Using an arrowhead structure, zygotes penetrate the tick gut epithelium. The arrowhead structure dissociates, and zygotes translocate to the basal lamina. In the hemolymph, zygotes become ookinetes that travel to the salivary acini. There, ookinetes hypertrophy into sporoblasts, which remain dormant while the larva or nymph overwinters and, until the next tick stage, feeds on a reservoir host or, incidentally, on a human.

Parasite Transmission to Hosts

Larvae, nymphs, and adult ticks all feed on humans, but the nymph is the primary vector for *B. microti* transmission.^{1,14} The size of the nymph (<2.5 mm; as small as a poppy seed), its gray ground color, and the inconspicuous feeding site explain why only one-half of patients recall a tick bite.³ As nymphs attach to the host, their temperatures rise. Sporogony is initiated, and the sporoblast membrane folds. Micronemes and rhoptries appear. The cytoplasm starts to bud from the parent sporoblast, and nuclear division ensues. During the final hours of feeding, which typically last 72 hours, as many as 100,000 sporozoites are deposited in the dermis of the host. Events that lead sporozoites toward the bloodstream are not well understood. *Babesia* sporozoites do not undergo merogony in an exoerythrocytic compartment but readily invade erythrocytes. Surface antigens, particularly glycosylphosphatidylinositol-anchored proteins encoded by genes of the *bmn* family, are thought to mediate initial attachment.^{72,73} Micronemes also contribute to host cell invasion, as evidenced by the role of the apical membrane antigen 1 (AMA1),⁷⁴ a micronemal protein that anchors in the plasma membrane of the parasite at its apex. A role for AMA1 is consistent with a role for the rhoptry neck protein (RON) macromolecular complex that is projected onto the erythrocyte and anchors in its plasma membrane.⁷⁵ As with *Plasmodium* species, the erythrocyte membrane is thought to invaginate as the parasite glides using the moving junction formed by nascent AMA1-RON2 interactions. Once the invasion process is completed, the parasitophorous vacuole gradually disintegrates. The organism differentiates into a trophozoite that moves freely in the RBC cytoplasm and eventually undergoes binary fission, resulting in two to four merozoites (Fig. 281.2). Merozoites egress the erythrocyte, thereby triggering its lysis, and soon invade other erythrocytes.

Classification of *Babesia* Species Single-Gene Analysis

Using full-length 18S rRNA gene sequences, phylogenetic analysis of the order Piroplasmida has revealed five clades.⁷⁶ The first clade contains the *B. microti* species complex as well as *Babesia rodhaini*, a parasite of rodents. The second clade consists of piroplasmids found in the western United States and may constitute a species complex because *B. duncani*




FIG. 281.2 Giemsa stain of a thin blood film. The patient was diagnosed with babesiosis caused by *Babesia microti*. Trophozoites often are seen as ring forms, some with two chromatin dots. Ring forms have a peripheral location, as with *Plasmodium falciparum*, but their large clear central vacuole and the absence of brown pigment (hemozoin) are characteristic of *B. microti*. A single trophozoite can divide by merogony to generate four merozoites arranged in a tetrad, also known as Maltese cross (arrow). Tetrads are rare but pathognomonic of small *Babesia* spp. such as *B. microti* and *B. duncani*. Tetrads also can be seen in human red blood cells infected with *B. divergens* or *B. venatorum*. (Courtesy Dr. J. Vyas.)

(WA1, CA5) and *B. duncani*-type (WA2, CA6) organisms cluster separately from other patient isolates (CA1–CA4) and those of wildlife roaming in the same habitat.⁷⁷ The third clade contains *Theileria* spp. only. The fourth clade contains *B. divergens* isolates from cattle in Europe, but also *B. divergens*-like organisms from patients in the United States; *B. odocoilei*, which infects white-tailed deer in the United States; and *B. venatorum* isolates from patients in Europe and China.^{43,54,62} The fifth clade contains *Babesia* spp. that infect ungulates but rarely humans. These include *B. bovis* and *B. bigemina*, parasites of cattle, as well as *B. crassa* and *B. caballi*, which infect sheep and horses, respectively. Piroplasms of the fourth and fifth clades are *Babesia* spp. sensu stricto.

Long regarded as a single species, *B. microti* is a species complex. The internal topology of this complex was refined by analysis of the gene encoding the η subunit of the chaperonin-containing *t*-complex polypeptide 1 (CCT η).⁶⁵ The number, position, and length of introns, as well as the nucleotide-based and amino-acid-based phylogenies, all concur to indicate that the *B. microti* species complex consists of four lineages (US-type, Munich, Kobe, and Hobetsu) and that these four lineages correspond to four separate species.

Whole-Genome Analysis

The genomes of *B. microti*, *B. divergens*, and *B. bovis* have been fully sequenced.^{72,78,79} Each genome consists of four nuclear chromosomes, a linear mitochondrial genome, and a circular apicoplast genome. The nuclear genome of *B. microti* is the smallest (6.7 Mb vs. approximately 8 Mb for *B. bovis* and approximately 10 Mb for *B. divergens*). It is notable for the absence of a large family of *ves*-like genes that, in *B. bovis*, encode variant erythrocyte surface antigens (VESAs). VESAs mediate the sequestration of infected erythrocytes in the vasculature. The four *ves*-like genes present in the *B. microti* genome are not arranged in tandem and most likely are not expressed, hence the lack of cerebral manifestations in babesiosis caused by *B. microti*. *Babesia divergens*, a species with numerous *ves*-like genes, does not evoke cerebral manifestations because the encoded VESA proteins lack the cysteine-rich domains critical for cytoadhesion of infected erythrocytes.

Genome-wide analysis of single nucleotide polymorphisms among *B. microti* isolates obtained from ticks and humans in the United States revealed strong population structure with deep divergence between the Midwest and the Northeast.^{80,81} Within the Northeast, three clusters were identified. The first cluster is found on Nantucket, an island 30

miles off Cape Cod. The second cluster encompasses the Northeast mainland except Cape Cod. The third cluster stretches from Cape Cod to the nearby Martha's Vineyard island. The first and second clusters are closely related, whereas the third cluster emerged earlier. Despite the close geographic proximity of these clusters, there is little evidence of genetic recombination.⁸⁰ This strong population structure is consistent with the low dispersal capability of *B. microti* that is imposed by the short-range migration of *P. leucopus*, the primary reservoir host. Unlike the Nantucket cluster, the Northeast mainland population presents genetic evidence of recent demographic expansion, possibly from west to east.^{80,81} The cluster that spans from Martha's Vineyard to Cape Cod may correspond to a relictual (surviving) population that was maintained in its enzootic cycle by *Ixodes muris* or *Ixodes angustus* prior to the arrival of *I. scapularis*.⁸⁰ Whether these distinct *B. microti* populations differ by the number or severity of symptoms they evoke, or both, is unknown.

CLINICAL MANIFESTATIONS

Babesia microti Infection

Symptoms typically appear 1 to 4 weeks after the bite of an infected tick, but 3 to 7 weeks when the infection is transmitted through blood transfusion (median interval, 37 days; range, 11–176 days).^{1,17} Most patients experience a gradual onset of fatigue, weakness, or malaise or a combination of these, followed within days by fever and one or more of the following: chills, sweats, headache, myalgia, and anorexia.¹ Fever is intermittent or persistent and has reached 40.9°C (105.6°F). Less frequent symptoms include arthralgia, neck stiffness, nonproductive cough, sore throat, nausea/vomiting, weight loss, and emotional lability. Photophobia, conjunctivitis, joint swelling, diarrhea, and crampy abdominal pain are rare. Dark urine raises the suspicion of severe hemolytic anemia, and may be accompanied by shortness of breath.

On physical examination, fever is the salient feature.¹ The skin may be pale or yellowish. A local red rash can mark the tick bite site; an erythema migrans rash is diagnostic of intercurrent Lyme disease.⁴ Ecchymoses and petechiae are rare. Examination of the mouth is unremarkable except when a slight pharyngeal erythema is noted. Scleral icterus is consistent with severe hemolytic anemia. Retinopathy with splinter hemorrhages and retinal infarcts have been reported. Tenderness of the upper left quadrant suggests splenomegaly and may be accompanied by hepatomegaly. Abdominal pain or unexplained hypotension with increased heart rate raises the suspicion of splenic rupture and hemoperitoneum. Splenic rupture with or without palpable splenomegaly has been reported.^{82,83} Splenic infarction and subcapsular hematoma may develop in the absence of splenic rupture. Splenic infarction and splenic rupture are confirmed by computed tomography.

Severe babesiosis requires hospital admission.^{5,28–30} Of the cases reported to the CDC from 2011 to 2014, one-half were admitted at least overnight.³ Of those for which the length of hospitalization was documented, one-half had a hospital stay of 3 to 5 days (range, 1–39 days). In smaller case series, the median length of stay typically has ranged from 5 to 9 days, with one patient remaining in the hospital for 100 days.^{5,7,28–30,32} Two-thirds of cases are diagnosed in persons 50 to 79 years of age, with the largest number occurring in people 60 to 69 years of age.³ Patients who are admitted to the hospital are older (median age, 68 years) than those who are not admitted (median age, 58 years).² Although age is a risk factor for hospitalization, it is not a risk factor for severe babesiosis.^{9,29,30} This finding is best explained by the fact that comorbidities and immunosenescence are not strict correlates of chronologic age. Asplenia and autoimmune disorders predispose to severe babesiosis and therefore hospital admission.²⁹ Among clinical features, nausea/vomiting and diarrhea are strong predictors of hospitalization.²⁹

Severe babesiosis can lead to complications. Pulmonary edema and acute respiratory distress syndrome are most common, followed by congestive heart failure and renal failure.^{5,7,28–30,32} Disseminated intravascular coagulation, shock, and splenic rupture are rare. In earlier case series, when clindamycin plus quinine was the mainstay of therapy, death occurred in 6% to 9% of hospitalized patients.^{28,30} Of the 10,305 Medicare recipients who received a diagnosis of babesiosis between 2006 and 2013, 1% died within 30 days.⁸⁴ Among inpatients, the 30-day mortality rate was 3%. In this series, atovaquone plus azithromycin was

the mainstay of therapy, whereas clindamycin plus quinine was used to treat one-fourth of patients. Death is more frequent (approximately 20%) among immunocompromised patients.³² Of the TTB cases caused by *B. microti* until fiscal year 2016, one-tenth ended in death.^{16,17,21}

Babesia duncani Infection

The eight documented cases caused by *B. duncani* were moderate to severe.^{37–41} All but one had one or several comorbidities. Clinical manifestations were similar to those of *B. microti*. Of the five patients who presumably acquired the infection via tick bite, two had a complicated course.³⁸ One died of cardiopulmonary arrest despite antimicrobial therapy, transfusion, and hemodialysis. The other experienced disseminated intravascular coagulation, pulmonary edema, and renal insufficiency but recovered following antimicrobial therapy and exchange transfusion. The three cases acquired via blood transfusion were severe. One was so anemic that packed RBCs were transfused before diagnosis.³⁹ Another, a neonate, developed severe respiratory distress.⁴⁰ The third developed acute renal failure requiring hemodialysis.⁴¹ All three patients recovered following standard antimicrobial therapy. The implicated blood donors were unaware of their babesial infection but recalled experiencing nausea, fatigue, or a mild flulike illness.

Babesia divergens Infection

Most cases caused by *B. divergens* infection occur in asplenic patients, and are severe.⁴⁷ Symptoms tend to appear abruptly 1 to 3 weeks following a tick bite, and consist of persistent high fever (40°C–41°C [104°F–105.8°F]), headache, shaking chills, drenching sweats, myalgia, and lumbar and abdominal pain. Dark urine and jaundice are consistent with severe hemolytic anemia. Without immediate treatment, a shocklike syndrome can develop, with renal failure and pulmonary edema. Once a fatal disease, symptomatic *B. divergens* infection is now successfully managed with exchange transfusion as an adjunct to antimicrobial therapy. *B. divergens* rarely causes disease in spleen-intact individuals. One such case was mild and resolved without adequate antimicrobial therapy.⁸⁵ Another case was severe and relapsed despite standard antimicrobial therapy, but eventually resolved after extended therapy with an alternative regimen.⁸⁶ Cases of *B. divergens*-like infection in the United States have been severe. All six patients had been splenectomized and three died despite medical intervention.^{42–45}

Babesia venatorum Infection

B. venatorum causes mild to severe illness. All five cases in Europe occurred in splenectomized patients.^{52–54,86a} One patient was markedly fatigued and produced dark urine.⁵⁴ Another patient experienced fever (39°C), chills, headache, jaundice, and dark urine. He recently had begun chemotherapy for diffuse large B-cell lymphoma. The third patient had been treated with rituximab for relapsing Hodgkin disease, and presented with weakness, shortness of breath, and lethargy.⁵³ He was hospitalized for severe anemia and tested positive in the direct Coombs test, as sometimes seen in babesiosis. He was initiated on prednisolone for presumptive autoimmune hemolytic anemia, which likely worsened the illness. The fourth patient was admitted for recurrent fever, hemolysis, and acute renal failure.⁵² He too had been treated with prednisolone for suspected autoimmune hemolytic anemia. The fifth patient was admitted for fever, myalgia, and dark urine. His history included idiopathic thrombocytopenic purpura, bouts of hemolytic anemia, and administration of rituximab 2 years earlier. He was treated with cyclosporine, prednisolone, human immunoglobulin, and a series of antimicrobial agents, including co-trimoxazole (trimethoprim-sulfamethoxazole). The diagnosis of babesiosis was made 2 months later, during a third hospital stay for fever and rigors. Co-trimoxazole, which was curative in a mild case of *B. divergens* infection, may have prevented the progression into severe disease. Despite a large spectrum of clinical severity, symptoms in the five European patients resolved following standard or extended antimicrobial therapy without exchange transfusion. A report from northeastern China indicated that *B. venatorum* causes disease in two-thirds of spleen-intact individuals.⁶² Symptoms were those evoked by *B. microti*, although fever, fatigue, chills, and myalgia were experienced by fewer patients. Of the 32 symptomatic individuals, 7 were admitted

for irregular fever as high as 40°C. The median length of stay was 16 days (range, 11–20 days). Only 4 patients received appropriate antibiotic therapy (although clindamycin monotherapy), but all 32 patients recovered.

PATHOGENESIS

Erythrocyte Clearance and Rupture

Continuous clearance of *Babesia*-infected RBCs and uninfected RBCs by phagocytes that reside in the red pulp of the spleen contributes to splenomegaly and splenic rupture,⁸⁷ and can lead to hemophagocytic lymphohistiocytosis, a fatal condition.⁸⁸ As the immune response unfolds, inflammatory cytokines are produced. By suppressing erythropoiesis, interferon- γ (IFN- γ) and interleukin-6 likely contribute to severe anemia. Persistent anemia, despite parasite clearance, has been attributed to autoantibodies that target RBCs.⁸⁹ Egress of merozoites is accompanied by erythrocyte rupture and hemoglobin release into the bloodstream. Hemoglobin is rapidly complexed by haptoglobin; minute amounts of free hemoglobin, however, are sufficient to promote systemic inflammation.⁹⁰

Inflammatory and Immune Response

The spleen is the immunodominant organ in babesiosis.¹ This feature, along with the clearance of parasitized RBCs, explains why asplenia is a major risk factor for severe babesiosis.²⁹ Another key risk factor for severe babesiosis is a low CD4⁺ T-cell count, as seen in HIV/AIDS patients and allograft recipients.^{32,33,35,36} In mice infected with *B. microti*, depletion of CD4⁺ T cells, a major source of IFN- γ , or neutralization of IFN- γ is sufficient to break host resistance and prevent parasite clearance.^{91,92} While central to host immunity, cytokines likely contribute to clinical manifestations. For example, circulating TNF- α and interleukin-6 are elevated during acute illness with *B. microti*,⁹³ and likely fuel the triad of symptoms consisting of fever, chills, and sweats. Cytokines may also contribute to complications. Mice infected with *B. duncani* die from pulmonary edema associated with intravascular margination of leukocytes. TNF- α is localized to the alveolar septa, whereas IFN- γ is expressed by cells around and within pulmonary vessels.⁹⁴ In this model, TNF- α blockade promotes survival, whereas IFN- γ receptor blockade promotes death.^{95,96} Although nearly every immunocompetent patient infected with *B. microti* has a positive serology at diagnosis, a role for antibodies is uncertain, as demonstrated in mice.⁹¹ Patients who are or were recently treated with rituximab are prone to persistent or relapsing babesiosis or both,^{32,97–99} indicating that B cells, and most likely antibodies, are key to clearing *B. microti* parasites in some immunocompromised hosts.

DIAGNOSIS

A diagnosis of babesiosis should be considered for patients who experience typical symptoms of babesiosis and may have been exposed to ticks in an endemic area between late spring and early fall or who have received blood components, particularly packed RBCs, in the past 6 months.¹ Given that *I. scapularis* transmits *B. microti*, *B. burgdorferi*, and *Anaplasma phagocytophilum* in the Northeast and upper Midwest of the United States, babesiosis should always be suspected in patients diagnosed with Lyme disease or human granulocytic anaplasmosis in these regions, particularly if symptoms worsen or do not abate within days to weeks of appropriate antibiotic therapy.^{4,100}

Routine Laboratory Findings

Routine laboratory tests provide clues. Anemia is evident on a complete blood count (low RBC count, low hematocrit, and low hemoglobin), and often accompanied by an elevated reticulocyte count.¹ The white blood cell (WBC) count is usually normal to slightly decreased, but sometimes elevated.^{29,30} A differential may reveal neutropenia as seen in one-third of adults and three-quarters of neonates.¹⁰¹ A severe thrombocytopenia often precedes severe anemia. Elevated blood levels of lactate dehydrogenase and total bilirubin are consistent with hemolytic anemia, findings that are substantiated by a depressed haptoglobin level. Elevated levels of aspartate aminotransferase, alanine aminotransferase, and alkaline phosphatase indicate hepatocyte injury, whereas elevated

aspartate aminotransferase levels without elevated alanine aminotransferase levels are consistent with hemolysis. Severe hemolysis is associated with excess urobilinogen, but rarely with hemoglobinuria. Elevated levels of blood urea nitrogen and serum creatinine denote renal compromise. The direct Coombs test may be positive on admission or in follow-up or both,^{52,53,89} implicating autoantibodies as contributors to the anemia. Autoantibodies against platelets can be induced, leading to immune thrombocytopenia.¹⁰² In asplenic patients, babesiosis has reactivated or provoked an Evans syndrome.¹⁰³

A study of 34 patients admitted to a hospital on Long Island, NY, reported that severe anemia (hemoglobin <10 g/dL) is a strong risk factor for complications as a whole, and that high-grade parasitemia ($\geq 10\%$) is marginally associated with disseminated intravascular coagulation.²⁸ A review of 139 cases that occurred in New York State between 1982 and 1993 indicated that elevated alkaline phosphatase levels (>125 U/L) and elevated WBC counts ($>5 \times 10^3/\mu\text{L}$) are strong predictors of severe outcome, defined as hospitalization longer than 2 weeks, stay in an intensive care unit longer than 2 days, or death.³⁰ A review of 128 cases diagnosed at Marshfield Clinic in Wisconsin between 1999 and 2015 identified total bilirubin levels greater than 1.9 mg/dL as a strong predictor of severe disease, defined as admission to the intensive care unit; an illness complicated by heart failure, shock, or splenic rupture; or the need for intubation or exchange transfusion.²⁹ In this study, high-grade parasitemia ($>10\%$), elevated creatinine levels (>1.2 mg/dL), and elevated WBC counts ($>10 \times 10^3/\mu\text{L}$) were moderate predictors of severe illness, whereas low WBC counts ($<5 \times 10^3/\mu\text{L}$) were highly predictive of severe disease.²⁹

Microscopy

Microscopic examination of Giemsa-stained thin blood smears has long been used to make a definitive diagnosis of babesiosis.^{1,47,54,77} *B. microti* trophozoites appear round, oval, or ameboid (see Fig. 281.2). The ring form is most common, with one or two red chromatic dots and a light blue cytoplasm. Tetrads of merozoites arranged in a Maltese cross are rare, but pathognomonic of *B. microti* and *B. duncani*. *B. divergens* and *B. venatorum* merozoites appear as paired piriforms but may form tetrads in human RBCs. The analytic sensitivity of microscopy (approximately 0.01% parasitemia as evaluated by highly experienced laboratory personnel) is sufficient in most cases because initial parasitemia typically ranges from 0.05% to 10% in spleen-intact patients, and even higher in asplenic patients (30%–50%, with one case at 85%). When parasitemia is high, extracellular *B. microti* merozoites, either single or in clumps, can be observed. *Babesia* rings resemble early-stage *P. falciparum* trophozoites but lack the brownish deposit (hemozoin) typical of older rings of *P. falciparum*. Gametocytes and schizonts are not observed in babesiosis.

Nucleic Acid Amplification

When *Babesia* organisms are not identified on smear but babesiosis is suspected, amplification of the parasite 18S rRNA gene is recommended. PCR assays that use a fluorescent probe can detect as few as 1 to 5 parasites per microliter of blood, that is, as low as 0.0001% parasitemia.^{104–106} Often qualitative, PCR testing provides little help in predicting the course of disease. By use of species-specific primers or probes, real-time PCR assays are well-suited for speciation of the causative organism. The CDC now uses a real-time PCR assay to identify *B. microti* in human blood.¹⁰⁷ This assay, although less sensitive than others, can detect *B. microti* isolates from all four lineages (US-type, Munich, Kobe, and Hobetsu) of the species complex. In patients who experience acute babesiosis caused by *B. microti* in the United States, the persistence of babesial DNA correlates with the persistence of symptoms and is shortened by antimicrobial therapy.³¹

Serology

The caveat of serologic testing is that IgG titers can be low or negative at time of presentation, and typically persist beyond parasite clearance.¹ An indirect fluorescent antibody (IFA) test to detect *B. microti*-specific IgG in serum or plasma can be ordered from commercial laboratories. A titer of 1:64 or greater is considered positive. Titers of 1:1024 or greater indicate active or recent infection. Following acute illness, titers

slowly wane over 6 to 12 months.³¹ Serology is not practical for *B. divergens* infection because the incubation period is short and the infection fulminant.⁴⁷ Sera that react with *B. microti* antigen do not cross react with *B. divergens* or *B. duncani* antigen and vice versa.

THERAPY

Asymptomatic *Babesia microti* Infection

Asymptomatic carriers need not be treated because they typically are immunocompetent and clear the infection within 1 year.¹⁶ Should they become at risk of severe babesiosis during this period (see “Risk Factors” earlier), treatment is recommended. At present, asymptomatic carriers often are identified during investigational product-release testing at selected blood drives. Their numbers will significantly increase once a systematic screening of the blood supply is instituted.

Mild *Babesia microti* Infection

Mild babesiosis caused by *B. microti* is treated with oral atovaquone plus oral azithromycin for 7 to 10 days.¹⁰⁸ Clindamycin plus quinine was the first regimen to demonstrate efficacy against *B. microti*; quinine, however, is poorly tolerated and often interrupted. A prospective, nonblinded, randomized clinical trial demonstrated that atovaquone plus azithromycin is as effective as clindamycin plus quinine in resolving symptoms of babesiosis and clearing *B. microti* parasites.¹⁰⁹ Importantly, untoward drug reactions were noted in fewer of the patients treated with atovaquone plus azithromycin (15% vs. 72% for clindamycin plus quinine).¹⁰⁹ Regimens for adults and children are provided in Table 281.1. Symptoms usually abate within 48 hours following initiation of antimicrobial therapy, and resolve within 1 to 2 weeks. Fatigue may persist for a few months. Low-grade parasitemia can persist for as long as 3 months, but follow-up PCR testing is not recommended because immunocompetent patients eventually clear the infection. Symptoms can recrudesce should a malignancy develop or an immunosuppressive regimen be administered before complete parasite clearance.³¹

Severe *Babesia microti* Infection First-Line Antimicrobial Therapy

Severe babesiosis caused by *B. microti* should be treated with oral atovaquone plus intravenous azithromycin. Based on cumulative clinical experience, it has long been recommended that severe babesiosis be treated with intravenous clindamycin plus oral quinine.¹⁰⁸ Quinine, however, has frequent and often severe side effects (cinchonism) and cardiotoxicity. No prospective study has compared the two regimens for severe *B. microti* illness, but a retrospective study of 40 patients admitted for babesiosis, including 11 admitted to intensive care, found that all patients but one resolved the infection while on atovaquone plus azithromycin.¹¹⁰ Regimens for adults and children are provided in Table 281.1. Intravenous azithromycin should be initiated at 500 mg/day.¹⁰⁰ Hematocrit, platelet count, parasitemia, liver enzymes, and renal function should be monitored daily until symptoms abate and parasitemia is reduced ($<4\%$). Thereafter, for patients who are immunocompetent and have a functional spleen, azithromycin can be prescribed orally and the dosage reduced to 250 mg/day.¹⁰⁰ In such patients, atovaquone plus azithromycin typically is administered for a total of 7 to 10 days. Duration, however, should be extended if symptoms persist. The approach to severely immunocompromised patients and asplenic patients is discussed separately (see later).

Adjunctive Exchange Transfusion

Partial or complete RBC exchange transfusion should be considered for patients with high-grade parasitemia ($\geq 10\%$) or severe anemia (hemoglobin <10 g/dL) in the context of parasitemia.¹⁰⁸ RBC exchange transfusion reduces parasitemia, corrects anemia, and removes inflammatory mediators and toxic by-products of RBC lysis (e.g., hemoglobin). Prompt use of RBC exchange transfusion is associated with favorable outcome.¹¹¹ A 90% reduction in parasitemia has been recommended as a target of apheresis and can be achieved by transfusion of a volume of allogeneic RBCs that amounts to 2.5 times the estimated volume of recipient RBCs.¹¹¹ When parasitemia is less than 10% and hemoglobin is 10 g/dL or greater, one should take into account the degree of systemic illness; acute respiratory distress syndrome and a systemic inflammatory

TABLE 281.1 Treatment of Human Babesiosis

SPECIES	ILLNESS	HOST ^a	FIRST-LINE REGIMEN ^b	ALTERNATIVE REGIMEN ^b
<i>Babesia microti</i>	Mild		Atovaquone 750 mg q12h PO plus azithromycin 500 mg PO on day 1 and 250 mg/d PO from day 2 on; for 7–10 d ^d	
	Severe ^c	Immunocompetent	Atovaquone 750 mg q12h PO plus azithromycin 500 mg/d IV ^{e,f} ; for 7–10 d ^d	Clindamycin 600 mg q6h IV plus quinine 650 mg q6–8h PO ^{g,h,i} ; for same duration
		Immunocompromised and/or asplenic	Atovaquone 750 mg q12h PO plus azithromycin 500 mg/d IV; for at least 6 consecutive wk, including 2 final wk during which parasites are no longer detected on blood smear ^{k,l}	Atovaquone 750 mg q12h PO plus clindamycin 600 mg q6h IV with or without azithromycin 500 mg/d IV ^{m,n} ; for same duration
<i>Babesia divergens</i>	Mild	Immunocompetent	Clindamycin 600 mg q8h PO plus quinine 650 mg q6–8h PO; for 7–10 d ^h	Atovaquone 750 mg q12h PO plus azithromycin 500 mg PO on day 1 and 250 mg/d PO from day 2 on; for 7–10 d ^{d,n}
	Severe	Immunocompromised and/or asplenic	Immediate complete RBC exchange transfusion combined with clindamycin 600 mg q6h IV plus quinine 650 mg q6–8h PO; for 7–10 d ^{d,h}	

^aDosages are provided for treatment of adults. For pediatric cases, consider using atovaquone 20 mg/kg q12h PO (up to 750 mg/dose) plus azithromycin 5–10 mg/kg PO (up to 250–500 mg/dose), or clindamycin 7–10 mg/kg q6–8h IV or PO (up to 600 mg/dose) plus quinine 8 mg/kg q8h PO (up to 650 mg/dose).

^bAtovaquone is available in a suspension form, and should be taken with dietary fat to increase absorption.

^cConsider partial or complete RBC exchange transfusion in cases of high-grade parasitemia ($\geq 10\%$), severe anemia (hemoglobin < 10 g/dL), or organ (pulmonary or renal) compromise.

^dDuration may be extended if symptoms other than fatigue linger.

^eIntravenous azithromycin may be replaced with oral azithromycin (250–500 mg/day) once the patient has improved.

^fRecommended for the treatment of severe babesiosis in hospitalized patients.

^gIntravenous clindamycin may be replaced with oral clindamycin (600 mg q8h) once the patient has improved.

^hBecause quinine carries a risk of QT segment prolongation and torsade de pointes, patients should be monitored by electrocardiography. In the setting of hepatic or renal disease, quinine serum levels should be monitored.

ⁱIf cardiac toxicity or untoward adverse effects occur, quinine can be replaced with oral atovaquone (750 mg q12h).

^jIntravenous azithromycin can be replaced with oral azithromycin (500 mg/d) once the patient has improved. Higher doses of oral azithromycin (600–1000 mg/d) accelerate symptom resolution and parasite clearance.

^kBlood smear is recommended to monitor resolution of infection in severely immunocompromised patients, but recent case reports support the use of a real-time polymerase chain reaction assay to ensure complete parasite clearance.

^lDuration may vary. In a patient who had been treated with rituximab, parasite clearance was achieved after 27 months of uninterrupted antimicrobial therapy. If asplenia is the only comorbidity, antimicrobial therapy may be shortened as long as clinical cure and parasite clearance are achieved in less than 4 weeks.

^mProlonged administration of clindamycin places the patient at risk of *Clostridioides difficile* (formerly *Clostridium difficile*) infection.

ⁿAtovaquone can be replaced with atovaquone-proguanil. Atovaquone-proguanil should be taken with dietary fat, but should not be given to patients with severe renal impairment.

IV, Intravenously; PO, orally; RBC, red blood cell.

response-like syndrome are indications for apheresis.¹¹² RBC exchange transfusion also is recommended for patients with renal compromise.¹⁰⁸

Alternative Antimicrobial Regimens

In patients who do not improve clinically within 48 hours of treatment with atovaquone plus azithromycin, intravenous clindamycin plus oral quinine should be considered.^{100,108} An alternative regimen can consist of intravenous clindamycin plus oral atovaquone with or without intravenous azithromycin, a regimen favored by the authors.¹¹² Once the patient has improved, antibiotics can be administered orally.^{5,97} Atovaquone-proguanil has been added to azithromycin or various two- or three-drug regimens, particularly when symptoms persisted or relapsed.^{36,97–99,112} In an AIDS patient, high-grade parasitemia resolved following addition of atovaquone-proguanil to a four-drug regimen that consisted of azithromycin, atovaquone, clindamycin, and quinine; an asymptomatic recurrence was treated with atovaquone-proguanil monotherapy.³³ Doxycycline is prescribed for intercurrent Lyme disease but has yet to be shown effective against *B. microti*.

Severely Immunocompromised Patients

A retrospective case-control study conducted more than 10 years ago provided the first evidence that severely immunocompromised patients, particularly those who are or were recently treated with rituximab for B-cell lymphoma, are at risk of relapsing babesiosis and should be treated with antimicrobial therapy for at least 6 consecutive weeks, including 2 final weeks during which parasites are no longer seen on blood smear.³² Given that no particular drug regimen is superior to another,³² the first-line regimen can consist of oral atovaquone plus intravenous azithromycin. Regimens for adults and children are provided in Table 281.1. Intravenous azithromycin should be initiated at 500 mg/day.¹⁰⁰ Hematocrit, platelet count, parasitemia, liver enzymes, and renal function should be monitored daily until symptoms abate and parasitemia is reduced ($< 4\%$). Thereafter, azithromycin can be administered orally but the dosage should be no less than 500 mg/day because lower dosages may

promote antibiotic resistance.^{99,100} Higher dosages of oral azithromycin (600–1000 mg/day) can be considered because they accelerate resolution of fever and clearance of parasites,¹¹³ thereby minimizing the risk of microbial relapse. Once the patient is no longer critically ill, laboratory testing can be performed every 2 to 3 days until antimicrobial therapy is completed. Close clinical follow-up is recommended.

The approach to treatment outlined for B-cell lymphoma patients extends to any patient who is or was recently treated with rituximab for autoimmune disorders, including rheumatoid arthritis, Evans syndrome, and confirmed or presumptive autoimmune hemolytic anemia.^{29,32,97,98} Time from last dose of rituximab to seroconversion and parasite clearance appears to be approximately 18 months.⁹⁷ This finding is consistent with the observation that patients who ended rituximab therapy as far as 18 months prior to a diagnosis of babesiosis are at risk of relapsing babesiosis should a standard course of antimicrobial therapy be administered.³² A prolonged antimicrobial therapy is also recommended for patients with X-linked agammaglobulinemia or low CD4⁺ T-cell counts due to HIV/AIDS, as well as patients who receive chronic immunosuppressive medication for malignancy, stem cell or solid organ transplantation, or autoimmune disorders.^{29,32,100} Reduction or interruption of such immunosuppressive regimens, when possible, is desirable.

Blood smear is recommended to monitor parasite clearance in severely immunocompromised patients.³² Recent case reports, however, stress the benefit of using real-time PCR to detect low-grade parasitemia.^{97,98} A premature interruption of antimicrobial therapy due to apparent cure has led to the emergence of microbial resistance to atovaquone and azithromycin.⁹⁹ Such resistance has been associated with missense mutations in the mitochondrial cytochrome b gene (*cytb*) and the apicoplast-encoded ribosomal protein subunit L4 gene (*rpl4*), respectively.⁸¹

Asplenic Patients

Asplenic patients account for at least one-half of reported cases of relapsing babesiosis.^{29,32} Asplenic patients, particularly when presenting with comorbidities that impair the immune response to *B. microti*

parasites, should be treated with prolonged antimicrobial therapy.³² If asplenia is the only comorbidity, therapy may be shortened to less than 6 weeks as long as clinical cure and parasite clearance are achieved in less than 4 weeks. Given the importance of the spleen for host resistance to *Babesia* parasites, a nonsurgical management of splenic rupture is preferred when the patient is at risk of tick exposure in areas endemic for babesiosis.^{82,87}

Infection With Other *Babesia* Species

Symptomatic *B. duncani* infection typically has been treated with intravenous clindamycin plus oral quinine.^{37–41} One patient developed a diffuse urticarial rash attributed to quinine, and was cured while receiving intravenous clindamycin monotherapy.³⁷

In Europe, symptomatic *B. divergens* infection in asplenic patients is managed with immediate, complete RBC exchange transfusion followed by intravenous clindamycin plus oral quinine.⁴⁷ Although used empirically, quinine may provide little to no benefit as demonstrated by its lack of activity in vitro against *B. divergens*¹¹⁴ and the full recovery by two asplenic patients despite premature interruption of quinine. A splenectomized individual who had a low CD4⁺ T-cell count due to HIV infection resolved a severe *B. divergens* infection while treated with atovaquone plus azithromycin.¹¹⁵ A relapse of babesiosis in a spleen-intact individual was treated successfully with atovaquone-proguanil plus azithromycin.⁸⁶ In the United States, cases of *B. divergens*-like infections typically have been treated with intravenous clindamycin plus oral quinine or quinidine.^{42–45} Quinidine is no longer available in the United States.

Two of the European cases of *B. venatorum* infection were successfully treated with intravenous clindamycin, alone or in combination with oral quinine.⁵⁴ A third case was treated with oral clindamycin plus oral quinine but the latter had to be interrupted.⁵³ Four weeks after clindamycin monotherapy ended, symptoms and parasitemia relapsed. Symptoms quickly resolved following initiation of oral atovaquone plus oral azithromycin, but low-level parasitemia persisted despite 11 weeks of therapy. To avoid relapse, atovaquone monotherapy was prescribed for

6 months until seroconversion and parasite clearance were noted. A pediatric case of mild *B. venatorum* infection in China was successfully treated with oral atovaquone plus oral azithromycin.⁶¹ Symptoms resolved in 3 days, and parasites cleared within 30 days.

PREVENTION

Despite the rise of transfusion-transmitted babesiosis, protection of the blood supply has relied unsatisfactorily on the permanent deferral of blood donors with a history of symptomatic babesiosis.^{15,19,21} In March 2018, the FDA approved a nucleic acid test and an arrayed fluorescent immunoassay for the detection of *B. microti* DNA and *B. microti* antibody in human whole blood and human plasma, respectively. These two tests, when combined, hold promise of dramatically reducing the incidence of TTB caused by *B. microti* in endemic areas, and possibly across the United States.¹⁶ In January 2019, the FDA approved the use of another nucleic acid test that is highly sensitive as it detects transcripts of the *B. microti* 18S rRNA gene (rather the gene itself).

Individuals at risk of severe babesiosis (e.g., asplenic and immunocompromised individuals) should avoid areas endemic for *Babesia* spp. from May through September when ticks seek blood.¹ Strategies for tick avoidance and tick removal, and a review of various acaricides are found at the CDC website (<http://www.cdc.gov/ticks/avoid/index.html>). No vaccine is available to protect humans from any of the zoonotic *Babesia* spp.

Given the increasing confluence of *Ixodes* ticks, *Babesia* spp., and individuals at risk of severe babesiosis in ever-enlarging areas, clinicians should be increasingly alert to the possibility of this previously arcane disease.

ACKNOWLEDGMENTS

The authors express their sincere gratitude to Dr. Tristram C. Dammin for inspiring discussions and unwavering support. They also acknowledge the support of the Dorothy Harrison Egan Foundation (EV) and the Edmund C. Lynch Jr. Research Fund (JAG).

Key References

The complete reference list is available online at Expert Consult.

- Vannier E, Krause PJ. Human babesiosis. *N Engl J Med*. 2012;366:2397–2407.
- Herwaldt B, Montgomery S, Woodhall D, et al. Babesiosis surveillance - 18 states, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:505–509.
- Diuk-Wasser MA, Vannier E, Krause PJ. Coinfection by *Ixodes* tick-borne pathogens: ecological, epidemiological, and clinical consequences. *Trends Parasitol*. 2016;32:40–42.
- Joseph JT, Roy SS, Shams N, et al. Babesiosis in Lower Hudson Valley, New York, USA. *Emerg Infect Dis*. 2011;17:843–847.
- Kowalski TJ, Jobe DA, Dolan EC, et al. The emergence of clinically relevant babesiosis in Southwestern Wisconsin. *WJMJ*. 2015;114:152–157.
- Leiby DA. Transfusion-transmitted *Babesia* spp.: bull's-eye on *Babesia microti*. *Clin Microbiol Rev*. 2011;24:14–28.
- Moritz ED, Winton CS, Tonnetti L, et al. Screening for *Babesia microti* in the U.S. blood supply. *N Engl J Med*. 2016;375:2236–2245.
- Herwaldt BL, Linden JV, Bosserman E, et al. Transfusion-associated babesiosis in the United States: a description of cases. *Ann Intern Med*. 2011;155:509–519.
- Saetre K, Godhwani N, Maria M, et al. Congenital babesiosis after maternal infection with *Borrelia burgdorferi* and *Babesia microti*. *J Pediatric Infect Dis Soc*. 2018;7:e1–e5.
- Yager PH, Luginbuhl LM, Dekker JP. Case records of the Massachusetts General Hospital. Case 6-2014. A 35-day-old boy with fever, vomiting, mottled skin, and severe anemia. *N Engl J Med*. 2014;370:753–762.
- Brennan MB, Herwaldt BL, Kazmierczak JJ, et al. Transmission of *Babesia microti* Parasites by Solid Organ Transplantation. *Emerg Infect Dis*. 2016;22.
- Hatcher JC, Greenberg PD, Antique J, et al. Severe babesiosis in Long Island: review of 34 cases and their complications. *Clin Infect Dis*. 2001;32:1117–1125.
- Mareedu N, Schotthoefer AM, Tompkins J, et al. Risk factors for severe infection, hospitalization, and prolonged antimicrobial therapy in patients with babesiosis. *Am J Trop Med Hyg*. 2017;97:1218–1225.
- White DJ, Talarico J, Chang HG, et al. Human babesiosis in New York State: review of 139 hospitalized cases and analysis of prognostic factors. *Arch Intern Med*. 1998;158:2149–2154.
- Krause PJ, Spielman A, Telford SR 3rd, et al. Persistent parasitemia after acute babesiosis. *N Engl J Med*. 1998;339:160–165.
- Krause PJ, Gewurz BE, Hill D, et al. Persistent and relapsing babesiosis in immunocompromised patients. *Clin Infect Dis*. 2008;46:370–376.
- Vyas JM, Telford SR, Robbins GK. Treatment of refractory *Babesia microti* infection with atovaquone-proguanil in an HIV-infected patient: case report. *Clin Infect Dis*. 2007;45:1588–1590.
- Ather I, Pourafshar N, Schain D, et al. Babesiosis: an unusual cause of sepsis after kidney transplantation and review of the literature. *Transpl Infect Dis*. 2017;19.
- Lubin AS, Snyderman DR, Miller KB. Persistent babesiosis in a stem cell transplant recipient. *Leuk Res*. 2011;35:e77–e78.
- Persing DH, Herwaldt BL, Glaser C, et al. Infection with a *Babesia*-like organism in northern California. *N Engl J Med*. 1995;332:298–303.
- Swei A, O'Connor KE, Couper LI, et al. Evidence for transmission of the zoonotic apicomplexan parasite *Babesia duncani* by the tick *Dermacentor albipictus*. *Int J Parasitol*. 2019;49:95–103.
- Burgess MJ, Rosenbaum ER, Pritt BS, et al. Possible transfusion-transmitted *Babesia divergens*-like/MO-1 infection in an Arkansas patient. *Clin Infect Dis*. 2017;64:1622–1625.
- Hildebrandt A, Gray JS, Hunfeld KP. Human babesiosis in Europe: what clinicians need to know. *Infection*. 2013;41:1057–1072.
- Svensson J, Hunfeld K-P, Persson KEM. High seroprevalence of *Babesia* antibodies amongst *Borrelia burgdorferi*-infected humans in Sweden. *Ticks Tick Borne Dis*. 2019;10:186–190.
- Herwaldt BL, Caccio S, Gherlinzoni F, et al. Molecular characterization of a non-*Babesia divergens* organism causing zoonotic babesiosis in Europe. *Emerg Infect Dis*. 2003;9:942–948.
- Sun Y, Li SG, Jiang JF, et al. *Babesia venatorum* infection in child, China. *Emerg Infect Dis*. 2014;20:896–897.
- Jiang JF, Zheng YC, Jiang RR, et al. Epidemiological, clinical, and laboratory characteristics of 48 cases of “*Babesia venatorum*” infection in China: a descriptive study. *Lancet Infect Dis*. 2015;15:196–203.
- Wei Q, Tsuji M, Zamoto A, et al. Human babesiosis in Japan: isolation of *Babesia microti*-like parasites from an asymptomatic transfusion donor and from a rodent from an area where babesiosis is endemic. *J Clin Microbiol*. 2001;39:2178–2183.
- Zhou X, Li SG, Chen SB, et al. Co-infections with *Babesia microti* and *Plasmodium* parasites along the China-Myanmar border. *Infect Dis Poverty*. 2013;2:24.
- Jia N, Zheng YC, Jiang JF, et al. Human babesiosis caused by a *Babesia crassa*-like pathogen: a case series. *Clin Infect Dis*. 2018;67:1110–1119.
- Cornillot E, Dassouli A, Garg A, et al. Whole genome mapping and re-organization of the nuclear and mitochondrial genomes of *Babesia microti* isolates. *PLoS ONE*. 2013;8:e72657.
- Conrad PA, Kjemtrup AM, Carreno RA, et al. Description of *Babesia duncani* n.sp. (Apicomplexa: babesidae) from humans and its differentiation from other piroplasms. *Int J Parasitol*. 2006;36:779–789.
- Carpi G, Walter KS, Mamoun CB, et al. *Babesia microti* from humans and ticks hold a genomic signature of strong population structure in the United States. *BMC Genomics*. 2016;17:888.
- Lemieux JE, Tran AD, Freimark L, et al. A global map of genetic diversity in *Babesia microti* reveals strong population structure and identifies variants associated with clinical relapse. *Nat Microbiol*. 2016;1:16079.
- El Khoury MY, Gandhi R, Dandache P, et al. Non-surgical management of spontaneous splenic rupture due to *Babesia microti* infection. *Ticks Tick Borne Dis*. 2011;2:235–238.
- Menis M, Forshee RA, Kumar S, et al. Babesiosis occurrence among the elderly in the United States, as recorded in large Medicare databases during 2006–2013. *PLoS ONE*. 2015;10:e0140332.
- Gonzalez LM, Rojo S, Gonzalez-Camacho F, et al. Severe babesiosis in immunocompetent man, Spain, 2011. *Emerg Infect Dis*. 2014;20:724–726.
- Blackberg J, Lazarevic VL, Hunfeld K-P, et al. Low-virulent *Babesia venatorum* infection masquerading as

- hemophagocytic syndrome. *Ann Hematol.* 2018;97:731–733.
87. Li S, Goyal B, Cooper JD, et al. Splenic rupture from babesiosis, an emerging concern? A systematic review of current literature. *Ticks Tick Borne Dis.* 2018;9:1377–1382.
 88. Akel T, Mobarakai N. Hematologic manifestations of babesiosis. *Ann Clin Microbiol Antimicrob.* 2017;16:6.
 89. Woolley AE, Montgomery MW, Savage WJ, et al. Post-babesiosis warm autoimmune hemolytic anemia. *N Engl J Med.* 2017;376:939–946.
 97. Raffalli J, Wormser GP. Persistence of babesiosis for >2 years in a patient on rituximab for rheumatoid arthritis. *Diagn Microbiol Infect Dis.* 2016;85:231–232.
 98. Simon MS, Westblade LF, Dziedzic A, et al. Clinical and Molecular Evidence of Atovaquone and Azithromycin Resistance in Relapsed *Babesia microti* infection associated with Rituximab and Chronic Lymphocytic Leukemia. *Clin Infect Dis.* 2017.
 99. Wormser GP, Prasad A, Neuhaus E, et al. Emergence of resistance to azithromycin-atovaquone in immunocompromised patients with *Babesia microti* infection. *Clin Infect Dis.* 2010;50:381–386.
 100. Sanchez E, Vannier E, Wormser GP, et al. Diagnosis, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: a review. *JAMA.* 2016;315:1767–1777.
 103. Shatzel JJ, Donohoe K, Chu NQ, et al. Profound autoimmune hemolysis and Evans syndrome in two asplenic patients with babesiosis. *Transfusion.* 2015;55:661–665.
 107. Souza SS, Bishop HS, Sprinkle P, et al. Comparison of *Babesia microti* real-time polymerase chain reaction assays for confirmatory diagnosis of babesiosis. *Am J Trop Med Hyg.* 2016;95:1413–1416.
 108. Wormser GP, Dattwyler RJ, Shapiro ED, et al. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis.* 2006;43:1089–1134.
 109. Krause PJ, Lepore T, Sikand VK, et al. Atovaquone and azithromycin for the treatment of babesiosis. *N Engl J Med.* 2000;343:1454–1458.
 110. Kletsova EA, Spitzer ED, Fries BC, et al. Babesiosis in Long Island: review of 62 cases focusing on treatment with azithromycin and atovaquone. *Ann Clin Microbiol Antimicrob.* 2017;16:26.
 111. Spaete J, Patrozou E, Rich JD, et al. Red cell exchange transfusion for babesiosis in Rhode Island. *J Clin Apher.* 2009;24:97–105.
 112. Stowell CP, Gelfand JA, Shepard JA, et al. Case records of the Massachusetts General Hospital. Case 17-2007. A 25-year-old woman with relapsing fevers and recent onset of dyspnea. *N Engl J Med.* 2007;356:2313–2319.
 113. Weiss LM, Wittner M, Tanowitz HB. The treatment of babesiosis. *N Engl J Med.* 2001;344:773.
 115. Gonzalez LM, Castro E, Lobo CA, et al. First report of *Babesia divergens* infection in an HIV patient. *Int J Infect Dis.* 2015;33:202–204.

References

- Vannier E, Krause PJ. Human babesiosis. *N Engl J Med*. 2012;366:2397–2407.
- Herwaldt B, Montgomery S, Woodhall D, et al. Babesiosis surveillance - 18 states, 2011. *MMWR Morb Mortal Wkly Rep*. 2012;61:505–509.
- Gray EB, Herwaldt BL. Surveillance for babesiosis - United States, 2014. Centers for Disease Control and Prevention. https://www.cdc.gov/parasites/babesiosis/resources/babesiosis_surveillance_summary_2016.pdf.
- Diuk-Wasser MA, Vannier E, Krause PJ. Coinfection by *Ixodes* tick-borne pathogens: ecological, epidemiological, and clinical consequences. *Trends Parasitol*. 2016;32:30–42.
- Joseph JT, Roy SS, Shams N, et al. Babesiosis in Lower Hudson Valley, New York, USA. *Emerg Infect Dis*. 2011;17:843–847.
- Linden JV, Prusinski MA, Crowder LA, et al. Transfusion-transmitted and community-acquired babesiosis in New York, 2004 to 2015. *Transfusion*. 2018;58:660–668.
- Kowalski TJ, Jobe DA, Dolan EC, et al. The emergence of clinically relevant babesiosis in Southwestern Wisconsin. *WMJ*. 2015;114:152–157.
- Johnson ST, Cable RG, Tonnetti L, et al. Seroprevalence of *Babesia microti* in blood donors from *Babesia*-endemic areas of the northeastern United States: 2000 through 2007. *Transfusion*. 2009;49:2574–2582.
- Krause PJ, McKay K, Gadbaw J, et al. Increasing health burden of human babesiosis in endemic sites. *Am J Trop Med Hyg*. 2003;68:431–436.
- Leiby DA, Chung AP, Gill JE, et al. Demonstrate parasitemia among Connecticut blood donors with antibodies to *Babesia microti*. *Transfusion*. 2005;45:1804–1810.
- Linden JV, Wong SJ, Chu FK, et al. Transfusion-associated transmission of babesiosis in New York State. *Transfusion*. 2000;40:285–289.
- Hilton E, DeVoti J, Benach JL, et al. Seroprevalence and seroconversion for tick-borne diseases in a high-risk population in the northeast United States. *Am J Med*. 1999;106:404–409.
- Tonnetti L, Thorp AM, Deisting B, et al. *Babesia microti* seroprevalence in Minnesota blood donors. *Transfusion*. 2013;53:1698–1705.
- Spielman A, Wilson ML, Levine JF, et al. Ecology of *Ixodes dammini*-borne human babesiosis and Lyme disease. *Annu Rev Entomol*. 1985;30:439–460.
- Leiby DA. Transfusion-transmitted *Babesia* spp.: bull's-eye on *Babesia microti*. *Clin Microbiol Rev*. 2011;24:14–28.
- Moritz ED, Winton CS, Tonnetti L, et al. Screening for *Babesia microti* in the U.S. blood supply. *N Engl J Med*. 2016;375:2236–2245.
- Herwaldt BL, Linden JV, Bosserman E, et al. Transfusion-associated babesiosis in the United States: a description of cases. *Ann Intern Med*. 2011;155:509–519.
- Bloch EM, Levin AE, Williamson PC, et al. A prospective evaluation of chronic *Babesia microti* infection in seroreactive blood donors. *Transfusion*. 2016;56:1875–1882.
- Biological Product and HCT/P Deviation Reports - Annual Summary for Fiscal Year 2016. Silver Spring, MD: Food and Drug Administration: <https://www.fda.gov/downloads/BiologicsBloodVaccines/SafetyAvailability/ReportaProblem/BiologicalProductDeviations/UCM564872.pdf>.
- Johnson ST, Cable RG, Leiby DA. Lookback investigations of *Babesia microti*-seropositive blood donors: seven-year experience in a *Babesia*-endemic area. *Transfusion*. 2012;52:1509–1516.
- Fatalities Reported to FDA Following Blood Collection and Transfusion - Annual Summary for FY2016. Silver Spring, MD: Food and Drug Administration: <https://www.fda.gov/downloads/BiologicsBloodVaccines/SafetyAvailability/ReportaProblem/TransfusionDonationFatalities/UCM598243.pdf>.
- Cornett JK, Malhotra A, Hart D. Vertical transmission of babesiosis from a pregnant, splenectomized mother to her neonate. *Infect Dis Clin Pract*. 2012;20:408–410.
- Joseph JT, Purtil K, Wong SJ, et al. Vertical transmission of *Babesia microti*, United States. *Emerg Infect Dis*. 2012;18:1318–1321.
- Saetre K, Godhwani N, Maria M, et al. Congenital babesiosis after maternal infection with *Borrelia burgdorferi* and *Babesia microti*. *J Pediatric Infect Dis Soc*. 2018;7:e1–e5.
- Surra ND, Jesus JE. The anemic and thrombocytopenic febrile neonate. *J Emerg Med*. 2015;48:675–678.
- Yager PH, Luginbuhl LM, Dekker JP. Case records of the Massachusetts General Hospital. Case 6-2014. A 35-day-old boy with fever, vomiting, mottled skin, and severe anemia. *N Engl J Med*. 2014;370:753–762.
- Brennan MB, Herwaldt BL, Kazmierczak JJ, et al. Transmission of *Babesia microti* Parasites by Solid Organ Transplantation. *Emerg Infect Dis*. 2016;22.
- Hatcher JC, Greenberg PD, Antique J, et al. Severe babesiosis in Long Island: review of 34 cases and their complications. *Clin Infect Dis*. 2001;32:1117–1125.
- Mareedu N, Schotthoef AM, Tompkins J, et al. Risk factors for severe infection, hospitalization, and prolonged antimicrobial therapy in patients with babesiosis. *Am J Trop Med Hyg*. 2017;97:1218–1225.
- White DJ, Talarico J, Chang HG, et al. Human babesiosis in New York State: review of 139 hospitalized cases and analysis of prognostic factors. *Arch Intern Med*. 1998;158:2149–2154.
- Krause PJ, Spielman A, Telford SR 3rd, et al. Persistent parasitemia after acute babesiosis. *N Engl J Med*. 1998;339:160–165.
- Krause PJ, Gewurz BE, Hill D, et al. Persistent and relapsing babesiosis in immunocompromised patients. *Clin Infect Dis*. 2008;46:370–376.
- Vyas JM, Telford SR, Robbins GK. Treatment of refractory *Babesia microti* infection with atovaquone-proguanil in an HIV-infected patient: case report. *Clin Infect Dis*. 2007;45:1588–1590.
- Cullen G, Sands BE, Yajnik V. Babesiosis in a patient on infliximab for Crohn's disease. *Inflamm Bowel Dis*. 2010;16:1269–1270.
- Ather I, Pourafshar N, Schain D, et al. Babesiosis: an unusual cause of sepsis after kidney transplantation and review of the literature. *Transp Infect Dis*. 2017;19.
- Lubin AS, Snyderman DR, Miller KB. Persistent babesiosis in a stem cell transplant recipient. *Leuk Res*. 2011;35:e77–e78.
- Quick RE, Herwaldt BL, Thomford JW, et al. Babesiosis in Washington State: a new species of *Babesia*? *Ann Intern Med*. 1993;119:284–290.
- Persing DH, Herwaldt BL, Glaser C, et al. Infection with a *Babesia*-like organism in northern California. *N Engl J Med*. 1995;332:298–303.
- Herwaldt BL, Kjemtrup AM, Conrad PA, et al. Transfusion-transmitted babesiosis in Washington State: first reported case caused by a WA1-type parasite. *J Infect Dis*. 1997;175:1259–1262.
- Kjemtrup AM, Lee B, Fritz CL, et al. Investigation of transfusion transmission of a WA1-type babesial parasite to a premature infant in California. *Transfusion*. 2002;42:1482–1487.
- Bloch EM, Herwaldt BL, Leiby DA, et al. The third described case of transfusion-transmitted *Babesia duncani*. *Transfusion*. 2012;52:1517–1522.
- Swei A, O'Connor KE, Couper LI, et al. Evidence for transmission of the zoonotic complex parasite *Babesia duncani* by the tick *Dermacentor albipictus*. *Int J Parasitol*. 2019;49:95–103.
- Beattie JF, Michelson ML, Holman PJ. Acute babesiosis caused by *Babesia divergens* in a resident of Kentucky. *N Engl J Med*. 2002;347:697–698.
- Burgess MJ, Rosenbaum ER, Pritt BS, et al. Possible transfusion-transmitted *Babesia divergens*-like/MO-1 infection in an Arkansas patient. *Clin Infect Dis*. 2017;64:1622–1625.
- Herwaldt B, Persing DH, Precigout EA, et al. A fatal case of babesiosis in Missouri: identification of another piroplasm that infects humans. *Ann Intern Med*. 1996;124:643–650.
- Herwaldt BL, de Bruyn G, Pieniazek NJ, et al. *Babesia divergens*-like infection, Washington State. *Emerg Infect Dis*. 2004;10:622–629.
- Jones SG, Coulter S, Conner W. Using administrative medical claims data to supplement state disease registry systems for reporting zoonotic infections. *J Am Med Inform Assoc*. 2012;20:193–198.
- Hildebrandt A, Gray JS, Hunfeld KP. Human babesiosis in Europe: what clinicians need to know. *Infection*. 2013;41:1057–1072.
- Haapasalo K, Suomalainen P, Sukura A, et al. Fatal babesiosis in man, Finland, 2004. *Emerg Infect Dis*. 2010;16:1116–1118.
- O'Connell S, Lyons C, Abdou M, et al. Splenic dysfunction from celiac disease resulting in severe babesiosis. *Ticks Tick Borne Dis*. 2017;8:537–539.
- Hunfeld KP, Lambert A, Kampen H, et al. Seroprevalence of *Babesia* infections in humans exposed to ticks in midwestern Germany. *J Clin Microbiol*. 2002;40:2431–2436.
- Sonnleitner ST, Fritz J, Bednarska M, et al. Risk assessment of transfusion-associated babesiosis in Tyrol: appraisal by seroepidemiology and polymerase chain reaction. *Transfusion*. 2014;54:1725–1732.
- Svensson J, Hunfeld K-P, Persson KEM. High seroprevalence of *Babesia* antibodies amongst *Borrelia burgdorferi*-infected humans in Sweden. *Ticks Tick Borne Dis*. 2019;10:186–190.
- Blum S, Gattringer R, Haschke E, et al. The case: hemolysis and acute renal failure. Babesiosis. *Kidney Int*. 2011;80:681–683.
- Haselbarth K, Tenter AM, Brade V, et al. First case of human babesiosis in Germany - Clinical presentation and molecular characterisation of the pathogen. *Int J Med Microbiol*. 2007;297:197–204.
- Herwaldt BL, Caccio S, Gherlinzoni F, et al. Molecular characterization of a non-*Babesia divergens* organism causing zoonotic babesiosis in Europe. *Emerg Infect Dis*. 2003;9:942–948.
- Hildebrandt A, Hunfeld KP, Baier M, et al. First confirmed autochthonous case of human *Babesia microti* infection in Europe. *Eur J Clin Microbiol Infect Dis*. 2007;26:595–601.
- Moniuszko-Malinowska A, Swiecicka I, Dunaj J, et al. Infection with *Babesia microti* in humans with non-specific symptoms in North East Poland. *Infect Dis (Lond)*. 2016;48:537–543.
- Foppa IM, Krause PJ, Spielman A, et al. Entomologic and serologic evidence of zoonotic transmission of *Babesia microti*, eastern Switzerland. *Emerg Infect Dis*. 2002;8:722–726.
- Lempereur L, Shiels B, Heyman P, et al. A retrospective serological survey on human babesiosis in Belgium. *Clin Microbiol Infect*. 2015;21:96 e91–96 e97.
- Pancewicz S, Moniuszko A, Bieniarz E, et al. Anti-*Babesia microti* antibodies in foresters highly exposed to tick bites in Poland. *Scand J Infect Dis*. 2011;43:197–201.
- Rigaud E, Jaulhac B, Garcia-Bonnet N, et al. Seroprevalence of seven pathogens transmitted by the *Ixodes ricinus* tick in forestry workers in France. *Clin Microbiol Infect*. 2016;22:735 e731–735 e739.
- Sun Y, Li SG, Jiang JF, et al. *Babesia venatorum* infection in child, China. *Emerg Infect Dis*. 2014;20:896–897.
- Jiang JF, Zheng YC, Jiang RR, et al. Epidemiological, clinical, and laboratory characteristics of 48 cases of “*Babesia venatorum*” infection in China: a descriptive study. *Lancet Infect Dis*. 2015;15:196–203.
- Zhou X, Xia S, Huang JL, et al. Human babesiosis, an emerging tick-borne disease in the People's Republic of China. *Parasit Vectors*. 2014;7:509.
- Wei Q, Tsuji M, Zamoto A, et al. Human babesiosis in Japan: isolation of *Babesia microti*-like parasites from an asymptomatic transfusion donor and from a rodent from an area where babesiosis is endemic. *J Clin Microbiol*. 2001;39:2178–2183.
- Nakajima R, Tsuji M, Oda K, et al. *Babesia microti*-group parasites compared phylogenetically by complete sequencing of the *CCteta* gene in 36 isolates. *J Vet Med Sci*. 2009;71:55–68.
- Tsuji M, Wei Q, Zamoto A, et al. Human babesiosis in Japan: epizootologic survey of rodent reservoir and isolation of new type of *Babesia microti*-like parasite. *J Clin Microbiol*. 2001;39:4316–4322.
- Zamoto-Niikura A, Tsuji M, Qiang W, et al. Detection of two zoonotic *Babesia microti* lineages, the Hobetsu and U.S. lineages, in two sympatric tick species, *Ixodes ovatus* and *Ixodes persulcatus*, respectively, in Japan. *Appl Environ Microbiol*. 2012;78:3424–3430.
- Zhou X, Li SG, Chen SB, et al. Co-infections with *Babesia microti* and *Plasmodium* parasites along the China-Myanmar border. *Infect Dis Poverty*. 2013;2:24.
- Zhou X, Li SG, Wang JZ, et al. Emergence of human babesiosis along the border of China with Myanmar: detection by PCR and confirmation by sequencing. *Emerg Microbes Infect*. 2014;3:e55.
- Jia N, Zheng YC, Jiang JF, et al. Human babesiosis caused by a *Babesia crassa*-like pathogen: a case series. *Clin Infect Dis*. 2018;67:1110–1119.
- Hersh MH, Tibbetts M, Strauss M, et al. Reservoir competence of wildlife host species for *Babesia microti*. *Emerg Infect Dis*. 2012;18:1951–1957.
- Cornillot E, Dassouli A, Garg A, et al. Whole genome mapping and re-organization of the nuclear and mitochondrial genomes of *Babesia microti* isolates. *PLoS ONE*. 2013;8:e72657.
- Silva JC, Cornillot E, McCracken C, et al. Genome-wide diversity and gene expression profiling of *Babesia microti* isolates identify polymorphic genes that mediate host-pathogen interactions. *Sci Rep*. 2016;6:35284.
- Moitra P, Zheng H, Anantharaman V, et al. Expression, purification, and biological characterization of *Babesia microti* apical membrane antigen 1. *Infect Immun*. 2015;83:3890–3901.
- Wang G, Efstratiou A, Adjou Moumouni PF, et al. Expression of truncated *Babesia microti* apical membrane protein 1 and rhoptry neck protein 2 and evaluation of their protective efficacy. *Exp Parasitol*. 2017;172:5–11.
- Criado-Fornelio A, Martinez-Marcos A, Buling-Sarana A, et al. Molecular studies on *Babesia*, *Theileria* and

- Hepatozoon* in southern Europe. Part II. Phylogenetic analysis and evolutionary history. *Vet Parasitol*. 2003;114:173–194.
77. Conrad PA, Kjemtrup AM, Carreno RA, et al. Description of *Babesia duncani* n.sp. (Apicomplexa: babesidae) from humans and its differentiation from other piroplasms. *Int J Parasitol*. 2006;36:779–789.
 78. Brayton KA, Lau AO, Herndon DR, et al. Genome sequence of *Babesia bovis* and comparative analysis of apicomplexan hemoprotozoa. *PLoS Pathog*. 2007;3:1401–1413.
 79. Jackson AP, Otto TD, Darby A, et al. The evolutionary dynamics of variant antigen genes in *Babesia* reveal a history of genomic innovation underlying host-parasite interaction. *Nucleic Acids Res*. 2014;42:7113–7131.
 80. Carpi G, Walter KS, Mamoun CB, et al. *Babesia microti* from humans and ticks hold a genomic signature of strong population structure in the United States. *BMC Genomics*. 2016;17:888.
 81. Lemieux JE, Tran AD, Freimark L, et al. A global map of genetic diversity in *Babesia microti* reveals strong population structure and identifies variants associated with clinical relapse. *Nat Microbiol*. 2016;1:16079.
 82. El Khoury MY, Gandhi R, Dandache P, et al. Non-surgical management of spontaneous splenic rupture due to *Babesia microti* infection. *Ticks Tick Borne Dis*. 2011;2:235–238.
 83. Kuwayama DP, Briones RJ. Spontaneous splenic rupture caused by *Babesia microti* infection. *Clin Infect Dis*. 2008;46:e92–e95.
 84. Menis M, Forshee RA, Kumar S, et al. Babesiosis occurrence among the elderly in the United States, as recorded in large Medicare databases during 2006–2013. *PLoS ONE*. 2015;10:e0140332.
 85. Martinot M, Zadeh MM, Hansmann Y, et al. Babesiosis in immunocompetent patients, Europe. *Emerg Infect Dis*. 2011;17:114–116.
 86. Gonzalez LM, Rojo S, Gonzalez-Camacho F, et al. Severe babesiosis in immunocompetent man, Spain, 2011. *Emerg Infect Dis*. 2014;20:724–726.
 - 86a. Blackberg J, Lazarevic VL, Hunfeld K-P, et al. Low-virulent *Babesia venatorum* infection masquerading as hemophagocytic syndrome. *Ann Hematol*. 2018;97:731–733.
 87. Li S, Goyal B, Cooper JD, et al. Splenic rupture from babesiosis, an emerging concern? A systematic review of current literature. *Ticks Tick Borne Dis*. 2018;9:1377–1382.
 88. Akel T, Mobarakai N. Hematologic manifestations of babesiosis. *Ann Clin Microbiol Antimicrob*. 2017;16:6.
 89. Woolley AE, Montgomery MW, Savage WJ, et al. Post-babesiosis warm autoimmune hemolytic anemia. *N Engl J Med*. 2017;376:939–946.
 90. Larsen R, Gozzelino R, Jeney V, et al. A central role for free heme in the pathogenesis of severe sepsis. *Sci Transl Med*. 2010;2:51ra71.
 91. Clawson ML, Paciorekowski N, Rajan TV, et al. Cellular immunity, but not gamma interferon, is essential for resolution of *Babesia microti* infection in BALB/c mice. *Infect Immun*. 2002;70:5304–5306.
 92. Igarashi I, Waki S, Ito M, et al. Role of CD4+ T cells in the control of primary infection with *Babesia microti* in mice. *J Protozool Res*. 1994;4:164–171.
 93. Shaio MF, Lin PR. A case study of cytokine profiles in acute human babesiosis. *Am J Trop Med Hyg*. 1998;58:335–337.
 94. Hemmer RM, Ferrick DA, Conrad PA. Up-regulation of tumor necrosis factor-alpha and interferon-gamma expression in the spleen and lungs of mice infected with the human *Babesia* isolate WA1. *Parasitol Res*. 2000;86:121–128.
 95. Aguilar-Delfin I, Wettstein PJ, Persing DH. Resistance to acute babesiosis is associated with interleukin-12- and gamma interferon-mediated responses and requires macrophages and natural killer cells. *Infect Immun*. 2003;71:2002–2008.
 96. Hemmer RM, Ferrick DA, Conrad PA. Role of T cells and cytokines in fatal and resolving experimental babesiosis: protection in TNFRp55-/- mice infected with the human *Babesia* WA1 parasite. *J Parasitol*. 2000;86:736–742.
 97. Raffalli J, Wormser GP. Persistence of babesiosis for >2 years in a patient on rituximab for rheumatoid arthritis. *Diagn Microbiol Infect Dis*. 2016;85:231–232.
 98. Simon MS, Westblade LF, Dziedzic A, et al. Clinical and Molecular Evidence of Atovaquone and Azithromycin Resistance in Relapsed *Babesia microti* infection associated with Rituximab and Chronic Lymphocytic Leukemia. *Clin Infect Dis*. 2017.
 99. Wormser GP, Prasad A, Neuhaus E, et al. Emergence of resistance to azithromycin-atovaquone in immunocompromised patients with *Babesia microti* infection. *Clin Infect Dis*. 2010;50:381–386.
 100. Sanchez E, Vannier E, Wormser GP, et al. Diagnosis, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: a review. *JAMA*. 2016;315:1767–1777.
 101. Wormser GP, Villafuerte P, Nolan SM, et al. Neutropenia in Congenital and Adult Babesiosis. *Am J Clin Pathol*. 2015;144:94–96.
 102. Narurkar R, Mamorska-Dyga A, Agarwal A, et al. Babesiosis-associated immune thrombocytopenia. *Stem Cell Investig*. 2017;4:1.
 103. Shatzel JJ, Donohoe K, Chu NQ, et al. Profound autoimmune hemolysis and Evans syndrome in two asplenic patients with babesiosis. *Transfusion*. 2015;55:661–665.
 104. Rollend L, Bent SJ, Krause PJ, et al. Quantitative PCR for detection of *Babesia microti* in Ixodes scapularis ticks and in human blood. *Vector Borne Zoonotic Dis*. 2013;13:784–790.
 105. Teal AE, Habura A, Ennis J, et al. A new real-time PCR assay for improved detection of the parasite *Babesia microti*. *J Clin Microbiol*. 2012;50:903–908.
 106. Wang G, Wormser GP, Zhuge J, et al. Utilization of a real-time PCR assay for diagnosis of *Babesia microti* infection in clinical practice. *Ticks Tick Borne Dis*. 2015;6:376–382.
 107. Souza SS, Bishop HS, Sprinkle P, et al. Comparison of *Babesia microti* real-time polymerase chain reaction assays for confirmatory diagnosis of babesiosis. *Am J Trop Med Hyg*. 2016;95:1413–1416.
 108. Wormser GP, Dattwyler RJ, Shapiro ED, et al. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis*. 2006;43:1089–1134.
 109. Krause PJ, Lepore T, Sikand VK, et al. Atovaquone and azithromycin for the treatment of babesiosis. *N Engl J Med*. 2000;343:1454–1458.
 110. Kletsova EA, Spitzer ED, Fries BC, et al. Babesiosis in Long Island: review of 62 cases focusing on treatment with azithromycin and atovaquone. *Ann Clin Microbiol Antimicrob*. 2017;16:26.
 111. Spaete J, Patrozou E, Rich JD, et al. Red cell exchange transfusion for babesiosis in Rhode Island. *J Clin Apher*. 2009;24:97–105.
 112. Stowell CP, Gelfand JA, Shepard JA, et al. Case records of the Massachusetts General Hospital. Case 17-2007. A 25-year-old woman with relapsing fevers and recent onset of dyspnea. *N Engl J Med*. 2007;356:2313–2319.
 113. Weiss LM, Wittner M, Tanowitz HB. The treatment of babesiosis. *N Engl J Med*. 2001;344:773.
 114. Brasseur P, Lecoublet S, Kapel N, et al. Quinine in the treatment of *Babesia divergens* infections in humans. *Eur J Clin Microbiol Infect Dis*. 1996;15:840–841.
 115. Gonzalez LM, Castro E, Lobo CA, et al. First report of *Babesia divergens* infection in an HIV patient. *Int J Infect Dis*. 2015;33:202–204.