



FIG. 237.12 Congenital syphilis. (A) Scaling annular rash of congenital syphilis. (B) Ulceration of perineal region in newborn with congenital syphilis. (C) Face of the same infant reveals perioral papules, ulcerous lesions, and labial fissures. (D) Dusky macular rash extending over the anterior trunk of a newborn infant with congenital syphilis. (Courtesy Dr. Leon Chameides, Hartford, CT.)

TABLE 237.3 Clinical Signs of Congenital Syphilis

Early Congenital Syphilis

Osteochondritis, periostitis
 Snuffles, hemorrhagic rhinitis
 Bullous lesions, palmar/plantar exanthem
 Hemolytic anemia
 Hepatosplenomegaly
 Jaundice
 Central nervous system involvement: elevated cell count or elevated protein in cerebrospinal fluid
 Generalized lymphadenopathy
 Pneumonitis
 Nephrotic syndrome
 Nonimmune hydrops fetalis
 Intrauterine growth retardation

Late Congenital Syphilis

Frontal bosses
 Short maxillas
 Saddle nose
 Protruding mandible
 Interstitial keratitis
 Eighth nerve deafness
 High palatal arch
 Hutchinson incisors
 Mulberry molars
 Sternoclavicular thickening (Higoumenaki sign)
 Clutton joints (bilateral painless swelling of knees)
 Saber shins
 Flaring scapulas

Modified from Kampmeier RH. Essentials of Syphilology. 3rd ed. Philadelphia: JB Lippincott; 1943; Stokes JH, Beerman H, Ingraham NR. Modern Clinical Syphilology: Diagnosis, Treatment: Case Study. 3rd ed. Philadelphia: Saunders; 1945; and Patton ME, Su JR, Nelson R, et al; Centers for Disease Control and Prevention. Primary and secondary syphilis—United States, 2005–2013. MMWR Morb Mortal Wkly Rep. 2014;63:402–406.

neonates will manifest both direct and indirect hyperbilirubinemia, as well as elevated transaminases. The major hematologic features of CS are anemia, leukocytosis or leukopenia, in some cases monocytosis, and thrombocytopenia.^{462,482} Renal involvement with an immune complex glomerulonephritis, evidenced by the appearance of nephrotic syndrome, may develop and usually occurs at about the fourth month of life.⁴⁸³ A generalized osteochondritis and periostitis or periostitis may affect the architecture of all bones of the skeletal system but are most prominent in the long bones and can be easily discerned on radiographs.^{484,485} CNS involvement is common, occurring in at least 22% of infected neonates,⁴⁸⁶ ranging from asymptomatic invasion by *T. pallidum* to acute syphilitic leptomeningitis. When untreated, neurosyphilis in the infant can lead to a chronic meningovascular process, which results in hydrocephalus, cranial nerve palsies, and even cerebral infarction. Asymptomatic CNS involvement is inferred from the presence of abnormal CSF studies, such as pleocytosis (defined as >25 white blood cells/ μ L for infants <1 month), high CSF protein levels (defined as >150 mg/dL in term infants <1 month of age and >170 mg/dL in preterm infants <1 month of age, although some experts use a threshold of >40 mg), a reactive CSF VDRL test, or a combination of these, in a healthy-appearing infant. Asymptomatic CNS syphilis occurs in approximately 40% of infants who have clinical, laboratory, or radiographic abnormalities of CS but is infrequent in infants without such manifestations. Because CSF studies are of low sensitivity to detect spread of spirochetes into the brain, in some cases asymptomatic disease can be diagnosed only by serum or CSF IgM immunoblotting or by PCR detection of spirochetal DNA.⁴⁸⁶ Hypopituitarism manifesting with persistent neonatal hypoglycemia has been observed among some infants with CS.⁴⁸⁷ Although rare, chorioretinitis, glaucoma, and uveitis have also been described in newborns with early CS.⁶⁰

With a few exceptions, the untreated child who survives the first 6 to 12 months of life will develop manifestations of late CS.⁶⁰ The generalized osteochondritis, perichondritis, and periostitis may result in deformities of the nose (saddle nose)⁴⁸⁸ and the metaphyses of the lower extremities (anterior bowing, or saber shin).⁶⁰ Cardiovascular syphilis is rare, whereas interstitial keratitis is common.⁴⁸⁹ Photophobia, pain, circumcorneal inflammation, and superficial and deep vascularization of the cornea may occur at any time between the ages of 5 and 30 years.⁴⁸⁹ Asymptomatic late congenital neurosyphilis occurs in up to one-third of perinatally infected children, whereas symptomatic disease is rare and fits the adult patterns of tabes dorsalis, syphilitic encephalitis, and local gummas. Eighth nerve compromise with resulting sensorineural deafness is particularly common.⁴⁸⁴ Other late characteristic stigmata include recurrent arthropathy and bilateral knee effusions (Clutton joints)⁴²⁵; centrally notched, widely spread, peg-shaped upper central incisors (Hutchinson teeth)⁴⁹⁰; frontal bossing; and poorly developed maxillas.⁴⁹¹

A complete evaluation of the neonate suspected of having CS should include a complete physical examination and DF microscopic examination or DFA staining of suspicious lesions or body fluids (e.g., nasal discharge). Conventional blood tests, including a complete blood count and liver and renal function tests, along with HIV testing and long bone radiographs, should be done in all infants. Nontreponemal tests (VDRL or RPR) and, if available, IgM immunoblotting are recommended on serum and CSF (VDRL only on CSF). The CDC no longer recommends obtaining umbilical cord blood because of the potential for contamination with maternal blood and because RPR testing of umbilical cord blood is prone to false positives owing to contamination with Wharton jelly.⁴⁹² The nontreponemal test that is performed on the infant should be the same as that done on the mother so that the maternal and infant titers can be directly compared. Other tests should be performed as clinically indicated (i.e., chest radiography, abdominal ultrasonography, ophthalmologic examination, auditory brainstem response, and neuroimaging studies). It is important to note that a diagnosis of acquired syphilis in an older child is suggestive of sexual abuse, although some cases might be attributable to progression of CS.⁴⁹³

To maximize treatment of children potentially infected with *T. pallidum*, including those who are asymptomatic at the time of initial presentation, in 1996 the CDC issued a revised case definition for CS (see Table 237.2).⁴⁹⁴ The criteria include all infants with clinical evidence of active syphilis, as well as normal-appearing neonates and stillbirths delivered to women with untreated or inadequately treated syphilis. According to these guidelines, a confirmed diagnosis of CS requires laboratory demonstration of *T. pallidum* in tissues or lesions of infected infants (see Table 237.2). Amniotic fluid, placental tissue, cytobrushings or sections of umbilical cord, nasal fluid, scrapings from a rash, tissue impressions, and lymph node aspirates are examples of specimens that can be examined for spirochetes.^{473,495–497} DF microscopy, DFA-TP, immunohistochemistry, and silver staining can all be used to visualize *T. pallidum*. PCR can reveal small amounts of *T. pallidum* DNA in lesion exudates or tissues but cannot be used to confirm the presence of living organisms.^{497–499} Because most of these detection methods are difficult to perform in most clinical settings, a confirmed diagnosis is achieved only in a small proportion of newborns, with most affected children receiving a diagnosis of presumptive CS. CS is highly probable if the neonate has any of the following: abnormal physical examination findings consistent with CS; a serum quantitative nontreponemal serologic titer that is greater than or equal to fourfold (two dilutions) the corresponding maternal titer; a reactive serum VDRL/RPR test and abnormal CSF (see earlier); and reactive treponemal antibody test after 15 months of age.

As per Table 237.2, the diagnosis of presumptive CS is made on the basis of (1) identification of syphilis in the mother; (2) adequacy of maternal treatment; (3) presence of clinical, laboratory, or radiographic evidence of syphilis in the infant; and (4) comparison of maternal (at delivery) and infant nontreponemal serologic titers obtained using the same test and preferably performed in the same laboratory.⁵⁰⁰ Infants younger than 1 month who have normal physical examination findings and serum VDRL or RPR titers less than fourfold the maternal titer are presumed to have CS if their mother was not treated or received

inadequate or suboptimal therapy. Some experts would also presume infants to have CS if their mothers had contact with a person with primary or secondary syphilis within 90 days before delivery and were not treated or were inadequately treated, even if the mother had nonreactive serology. A diagnosis of CS is unlikely if the infant has normal physical examination findings and a nonreactive VDRL/RPR test and the mother was adequately treated during pregnancy and had an appropriate serologic response, or if the neonate has normal physical examination findings and serum VDRL or RPR titer the same or less than fourfold the maternal titer and the mother was adequately treated before pregnancy and her titers remained low (VDRL <1:2; RPR <1:4) and stable before and during pregnancy and at delivery. Because at least one-third of the mothers who give birth to syphilitic children have not had prenatal care, and about half have had a nonreactive serologic test during the first trimester of pregnancy, serologic testing of the mother is always warranted at delivery, especially in high-risk patients. No infant or mother should leave the hospital unless the maternal serologic status has been documented at least once during pregnancy, and at delivery in communities and populations in which the risk for CS is high.

Parenteral penicillin is the drug of choice for the treatment of CS (see Table 237.2).^{60,501} Infants 4 weeks of age or younger who have confirmed or presumptive disease should be treated with either aqueous crystalline penicillin G, 50,000 units/kg IV every 12 hours for the first 7 days of life and every 8 hours beyond 1 week of age for a total of 10 days, or aqueous procaine penicillin G, 50,000 units/kg IM once daily for 10 days (see Table 237.2). The levels of penicillin that are achieved in the CSF after intramuscular administration of procaine penicillin are lower than those with intravenous administration of aqueous penicillin.⁵⁰² However, the clinical significance of this observation is unclear because there have been no treatment failures reported after treatment with procaine penicillin.⁴⁶¹ If more than 1 day of penicillin therapy is missed, the entire course should be restarted. Single-dose therapy with intramuscular injection of BPG (50,000 units/kg) can be considered in the infant with normal physical examination findings, normal CSF, and normal radiographic and laboratory studies and whose nontreponemal test titer is the same or less than the maternal titer, under the following circumstances: (1) The mother was treated with the appropriate regimen for 30 or fewer days before delivery; (2) the mother received her recommended penicillin therapy for the stage of infection during the pregnancy but the nontreponemal titer did not decrease fourfold at the time of delivery; and (3) the mother has untreated syphilis or her treatment status is undocumented. Children who are diagnosed with CS after 1 month of age (including those with late CS) and children with acquired syphilis should be treated with aqueous penicillin G (50,000 units/kg IV every 4 to 6 hours for 10 days). In addition, for children with CS or findings compatible with CNS involvement, some experts suggest that the 10-day course of aqueous penicillin be followed with a single dose of intramuscular BPG (50,000 units/kg).

Follow-up of children treated for CS after the newborn period should be conducted as is recommended for neonates. The risk of infection for the infant is minimal if the mother has received adequate penicillin treatment during pregnancy. Nevertheless, the child must be examined monthly after delivery and until the nontreponemal test or PCR assay result becomes negative. Penicillin treatment should never be withheld to prove the diagnosis, and every neonate born to a syphilitic mother should be promptly treated unless adequate, serologically proven effective treatment with penicillin can be documented more than 1 month before delivery. Because giving penicillin to the neonate is almost risk free, all neonates born to syphilitic mothers should be treated, regardless of whether the mother was treated during pregnancy. Adequate treatment of the mother usually but not always ensures that the fetus will not be infected. Infants born to mothers who are coinfecting with syphilis and HIV should receive the same evaluation and treatment as those whose mothers do not have HIV infection.⁵⁰¹ There is insufficient evidence to determine whether such infants require different evaluation, treatment, or follow-up.

IMMUNITY

For nearly 200 years, syphilologists have intensely pondered and debated the existence, timing, and nature of protective immunity in acquired

syphilis.^{169,503,504} In the 19th century, it was believed that individuals became immune to reinfection during primary syphilis, a notion referred to by the term “chance immunity.” This flawed concept was based largely on “autoinoculation” experiments in which material taken from presumptive chancres failed to cause lesions when inoculated elsewhere on the same patient. Using his primate model, Neisser concluded that chancre immunity is dependent on the presence of active infection and that cure of the disease leads to the disappearance of immunity.¹⁶⁹ Chesney,⁵⁰⁵ in contrast, correctly concluded from data generated using the rabbit model that syphilitic infection does indeed elicit lasting immunity, provided the initial infection is of sufficient duration. Important to note, he found substantially less cross-immunity to challenge with heterologous strains of *T. pallidum*, findings confirmed years later, also in rabbits, by Turner and Hollander.^{132,505} In a meticulously executed study involving nearly 300 rabbits, which took advantage of the recent availability of penicillin, Magnuson and Rosenau⁵⁰⁶ showed definitively that immunity to homologous challenge in the animal model was dose and time dependent and could be achieved in the absence of concurrent infection. To a significant extent, Magnuson and coworkers¹⁴⁰ replicated these results in humans several years later in a remarkable (and unrepeatable) series of experiments conducted at Sing Sing Prison. Although these investigators went to great length to explain that all participants were volunteers, the authors of this chapter must state unequivocally that they do not condone experimentation in incarcerated populations even if it was considered acceptable in the era in which it was performed. In any event, the Magnuson group showed that patients with treated primary and secondary syphilis were susceptible to challenge with the Nichols strain of *T. pallidum*, whereas those with treated or untreated late latent disease were resistant. Interpretation of these experiments is clouded by the extraordinarily high challenge inoculum used (1×10^5), more than 2000-fold greater than the median infective dose (ID₅₀) determined in the same study, compounded by the likely heterologous challenge in the majority of “volunteers.” However, there is also abundant clinical experience that patients successfully treated for primary and secondary syphilis can be reinfectd.^{508,509} It seems reasonable to conclude, therefore, that the immunity that develops during early syphilis is limited, even more so against heterologous infection, and that it dissipates rapidly after treatment. In a controversial paper, Grassly and coworkers³⁵ used mathematical models derived from epidemiologic data to argue that the oscillatory pattern of syphilis epidemics over an approximate 11-year cycle is due to the buildup of immunity in untreated populations, followed by the waning of immunity

after large-scale treatment. Their hypothesis deserves reconsideration given recent technical advances in sequencing genomes of syphilis spirochetes in lesions^{33,180} and our much improved knowledge of *T. pallidum*’s OMP repertoire,^{34,73} the presumptive targets of protective antibodies.

The lack of an inbred animal model has greatly hindered efforts to dissect the components of the immune response that confer protection. Because *T. pallidum* is an extracellular pathogen, one would expect passive administration of antibodies from immune animals to protect against homologous challenge. Such experiments have repeatedly shown that administration of large amounts of serum or purified immunoglobulins yields only modest lesion attenuation in rabbits challenged intradermally and that dissemination of spirochetes is not prevented.¹⁶⁹ In other words, passive immunization is at best only partially protective. The paucity of antigenic targets on the surface of the syphilis spirochete, coupled with other mechanisms for antibody avoidance (e.g., limited production of antibodies against surface-exposed epitopes, antigenic variation, and differential expression of OMPs; see earlier), is an attractive explanation for these results.⁷³

The inability of public health measures to curtail the explosive increase in syphilis cases worldwide, with particular concern for the global burden of maternal syphilis and CS, has called renewed attention to the need for an effective vaccine.^{5,434,503,504,510} The feasibility of eliciting a protective immune response against *T. pallidum* was demonstrated nearly 50 years ago. In a landmark study, Miller⁵⁰⁷ reported complete and long-lasting (1 year) protection of rabbits immunized over a 37-week period with *T. pallidum* rendered noninfectious by gamma irradiation. Important to note, this regimen did not protect against challenge with the Haiti B strain, thought at the time to be subspecies *pertenue* but now known to be subspecies *pallidum*. Until recently, efforts to replicate this stunning result by immunization with recombinant *T. pallidum* proteins believed to contain surface-exposed epitopes yielded either no or only partial degrees of protection. In what appears to be a major advance, Lithgow and colleagues⁵¹¹ showed that immunization of rabbits with TP0751 reduced spirochete dissemination. The high degree of conservation of TP0751 in *T. pallidum* strains suggests that a vaccine containing this lipoprotein could be useful on a global scale.

ACKNOWLEDGMENTS

We are deeply indebted to Ms. Carson Karanian and Ms. Morgan Ledoyt for their assistance with myriad aspects of manuscript production, referencing, and figure generation.

Key References

The complete reference list is available online at Expert Consult.

4. Oriel JD. *The Scars of Venus*. London: Springer-Verlag; 1994.
6. Larsen SA, Steiner BM, Rudolph AH. Laboratory diagnosis and interpretation of tests for syphilis. *Clin Microbiol Rev*. 1995;8:1–21.
9. Ghanem KG, Workowski KA. Management of adult syphilis. *Clin Infect Dis*. 2011;53(suppl 3):S110–S128.
10. Smajs D, Norris SJ, Weinstock GM. Genetic diversity in *Treponema pallidum*: implications for pathogenesis, evolution and molecular diagnostics of syphilis and yaws. *Infect Genet Evol*. 2012;12:191–202.
16. Harper KN, Zuckerman MK, Harper ML, et al. The origin and antiquity of syphilis revisited: an appraisal of Old World pre-Columbian evidence for treponemal infection. *Am J Phys Anthropol*. 2011;146(suppl 53):99–133.
38. Ghanem KG. Neurosyphilis: a historical perspective and review. *CNS Neurosci Ther*. 2010;16:e157–e168.
45. MacDonald CJ. Communications: the contribution of the Tuskegee study to medical knowledge. *J Natl Med Assoc*. 1974;66:1–7.
60. Shafii T, Radolf JD, Sanchez PJ, et al. Congenital syphilis. In: Holmes KK, Sparling PF, Stamm WE, et al, eds. *Sexually Transmitted Diseases*. New York: McGraw Hill; 2008:1577–1612.
79. Salazar JC, Hazlett KR, Radolf JD. The immune response to infection with *Treponema pallidum*, the stealth pathogen. *Microbes Infect*. 2002;4:1133–1140.
82. Magnuson HJ, Eagle H, Fleischman R. The minimal infectious inoculum of *Spirochaeta pallida* (Nichols strain) and a consideration of its rate of multiplication in vivo. *Am J Syph Gonorrhea Vener Dis*. 1948;32:1–18.
86. Peng RR, Wang AL, Li J, et al. Molecular typing of *Treponema pallidum*: a systematic review and meta-analysis. *PLoS Negl Trop Dis*. 2011;5:e1273.
96. Zetola NM, Klausner JD. Syphilis and HIV infection: an update. *Clin Infect Dis*. 2007;44:1222–1228.
116. Peterman TA, Su J, Bernstein KT, et al. Syphilis in the United States: on the rise? *Expert Rev Anti Infect Ther*. 2015;13:161–168.
117. Cohen SE, Chew Ng RA, Katz KA, et al. Repeat syphilis among men who have sex with men in California, 2002–2006: implications for syphilis elimination efforts. *Am J Public Health*. 2012;102:e1–e8.
118. Peterman TA, Newman DR, Maddox L, et al. High risk for HIV following syphilis diagnosis among men in Florida, 2000–2011. *Public Health Rep*. 2014;129:164–169.
120. Wasserheit JN. Epidemiological synergy. Interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis*. 1992;19:61–77.
134. Cruz AR, Ramirez LG, Zuluaga AV, et al. Immune evasion and recognition of the syphilis spirochete in blood and skin of secondary syphilis patients: two immunologically distinct compartments. *PLoS Negl Trop Dis*. 2012;6:e1717.
140. Magnuson HJ, Thomas EW, Olansky S, et al. Inoculation syphilis in human volunteers. *Medicine (Baltimore)*. 1956;35:33–82.
150. Lukehart SA, Hook EW 3rd, Baker-Zander SA, et al. Invasion of the central nervous system by *Treponema pallidum*: implications for diagnosis and treatment. *Ann Intern Med*. 1988;109:855–862.
151. Rolfs RT, Joesoef MR, Hendershot EF, et al. A randomized trial of enhanced therapy for early syphilis in patients with and without human immunodeficiency virus infection. The Syphilis and HIV Study Group. *N Engl J Med*. 1997;337:307–314.
168. Lafond RE, Lukehart SA. Biological basis for syphilis. *Clin Microbiol Rev*. 2006;19:29–49.
187. Jordaan HF. Secondary syphilis. A clinicopathological study. *Am J Dermatopathol*. 1988;10:399–409.
194. Jackman JD Jr, Radolf JD. Cardiovascular syphilis. *Am J Med*. 1989;87:425–433.
203. Clark EG, Danbolt N. The Oslo study of the natural history of untreated syphilis: an epidemiologic investigation based on a restudy of the Boeck-Bruusgaard material; a review and appraisal. *J Chron Dis*. 1955;2:311–344.
204. White RM. Unraveling the Tuskegee study of untreated syphilis. *Arch Intern Med*. 2000;160:585–598.
217. Musher DM, Hamill RJ, Baughn RE. Effect of human immunodeficiency virus (HIV) infection on the course of syphilis and on the response to treatment. *Ann Intern Med*. 1990;113:872–881.
219. Golden MR, Marra CM, Holmes KK. Update on syphilis: resurgence of an old problem. *JAMA*. 2003;290:1510–1514.
235. Chapel TA. The signs and symptoms of secondary syphilis. *Sex Transm Dis*. 1980;7:161–164.
257. Chahine LM, Khoriaty RN, Tomford WJ, et al. The changing face of neurosyphilis. *Int J Stroke*. 2011;6:136–143.
262. Nagappa M, Sinha S, Taly AB, et al. Neurosyphilis: MRI features and their phenotypic correlation in a cohort of 35 patients from a tertiary care university hospital. *Neuroradiology*. 2013;55:379–388.

287. Szilak I, Marty F, Helft J, et al. Neurosyphilis presenting as herpes simplex encephalitis. *Clin Infect Dis*. 2001;32:1108–1109.
300. Tong SY, Haqqani H, Street AC. A pox on the heart: five cases of cardiovascular syphilis. *Med J Aust*. 2006;184:241–243.
316. Wheeler HL, Agarwal S, Goh BT. Dark ground microscopy and treponemal serological tests in the diagnosis of early syphilis. *Sex Transm Infect*. 2004;80:411–414.
318. Creagan L, Bauer HM, Samuel MC, et al. An evaluation of the relative sensitivities of the Venereal Disease Research Laboratory test and the *Treponema pallidum* particle agglutination test among patients diagnosed with primary syphilis. *Sex Transm Dis*. 2007;34:1016–1018.
329. Gayet-Ageron A, Lautenschlager S, Ninet B, et al. Sensitivity, specificity and likelihood ratios of PCR in the diagnosis of syphilis: a systematic review and meta-analysis. *Sex Transm Infect*. 2013;89:251–256.
334. Gayet-Ageron A, Sednaoui P, Lautenschlager S, et al. Use of *Treponema pallidum* PCR in testing of ulcers for diagnosis of primary syphilis. *Emerg Infect Dis*. 2015;21:127–129.
337. Wu BR, Tsai MS, Yang CJ, et al. Spirochetemia due to *Treponema pallidum* using polymerase-chain-reaction assays in patients with early syphilis: prevalence, associated factors and treatment response. *Clin Microbiol Infect*. 2014;20:O524–O527.
346. Tong ML, Lin LR, Liu LL, et al. Analysis of 3 algorithms for syphilis serodiagnosis and implications for clinical management. *Clin Infect Dis*. 2014;58:1116–1124.
350. Nandwani R, Evans DT. Are you sure it's syphilis? A review of false positive serology. *Int J STD AIDS*. 1995;6:241–248.
362. Sena A, Pillay A, Cox DL, et al. *Treponema* and *Brachyspira*, human host-associated spirochetes. In: Jorgensen JH, Pfaller MA, Carroll KC, et al, eds. *Manual of Clinical Microbiology*. 11th ed. Washington, DC: ASM Press; 2015.
363. Sena AC, White BL, Sparling PF. Novel *Treponema pallidum* serologic tests: a paradigm shift in syphilis screening for the 21st century. *Clin Infect Dis*. 2010;51:700–708.
365. Morshed MG, Singh AE. Recent trends in the serologic diagnosis of syphilis. *Clin Vaccine Immunol*. 2015;22:137–147.
374. Tucker JD, Bien CH, Peeling RW. Point-of-care testing for sexually transmitted infections: recent advances and implications for disease control. *Curr Opin Infect Dis*. 2013;26:73–79.
388. Clement ME, Okeke NL, Hicks CB. Treatment of syphilis: a systematic review. *JAMA*. 2014;312:1905–1917.
394. Idsoe O, Guthe T, Willcox RR. Penicillin in the treatment of syphilis. The experience of three decades. *Bull World Health Organ*. 1972;47(suppl):1–68.
402. Rolfs RT. Treatment of syphilis, 1993. *Clin Infect Dis*. 1995;20(suppl 1):S23–S38.
410. Riedner G, Rusizoka M, Todd J, et al. Single-dose azithromycin versus penicillin G benzathine for the treatment of early syphilis. *N Engl J Med*. 2005;353:1236–1244.
417. Ghanem KG, Erbelding EJ, Cheng WW, et al. Doxycycline compared with benzathine penicillin for the treatment of early syphilis. *Clin Infect Dis*. 2006;42:e45–e49.
427. Lukehart SA, Godornes C, Molini BJ, et al. Macrolide resistance in *Treponema pallidum* in the United States and Ireland. *N Engl J Med*. 2004;351:154–158.
432. Wendel GD Jr, Sheffield JS, Hollier LM, et al. Treatment of syphilis in pregnancy and prevention of congenital syphilis. *Clin Infect Dis*. 2002;35:S200–S209.
447. Romanowski B, Sutherland R, Fick GH, et al. Serologic response to treatment of infectious syphilis. *Ann Intern Med*. 1991;114:1005–1009.
453. Schmid GP, Stoner BP, Hawkes S, et al. The need and plan for global elimination of congenital syphilis. *Sex Transm Dis*. 2007;34:S5–S10.
462. Hollier LM, Harstad TW, Sanchez PJ, et al. Fetal syphilis: clinical and laboratory characteristics. *Obstet Gynecol*. 2001;97:947–953.
463. Casal C, Araujo Eda C, Corvelo TC. Risk factors and pregnancy outcomes in women with syphilis diagnosed using a molecular approach. *Sex Transm Infect*. 2013;89:257–261.
484. Obladen M. Curse on two generations: a history of congenital syphilis. *Neonatology*. 2013;103:274–280.
491. Sparling PF, Musher MNSD, Healy BP. Clinical manifestations of syphilis. In: Holmes KK, Sparling PF, Stamm WE, et al, eds. *Sexually Transmitted Diseases*. 4th ed. New York: McGraw Hill Medical; 2008:662–684.
506. Magnuson HJ, Rosenau BJ. The rate of development and degree of acquired immunity in experimental syphilis. *Am J Syph Gonorrhea Vener Dis*. 1948;32:418–436.

References

- Stokes JH, Beerman H, Ingraham NR. *Modern Clinical Syphilology*. 3rd ed. Philadelphia: Saunders; 1944.
- Wright AD. Venereal disease and the great. *Br J Vener Dis*. 1971;47:295–306.
- Tramont EC. The impact of syphilis on humankind. *Infect Dis Clin North Am*. 2004;18:101–110.
- Oriel JD. *The Scars of Venus*. London: Springer-Verlag; 1994.
- Peeling RW, Mabey D, Kamb ML, et al. Syphilis. *Nat Rev Dis Primers*. 2017;3:17073.
- Larsen SA, Steiner BM, Rudolph AH. Laboratory diagnosis and interpretation of tests for syphilis. *Clin Microbiol Rev*. 1995;8:1–21.
- Sena AC, Pillay A, Cox DL, et al. *Treponema* and *Brachyspira*, human host-associated spirochetes. In: Jorgensen JH, Pfaller MA, Carroll KC, et al, eds. *Manual of Clinical Microbiology*. 11th ed. Washington, D.C.: ASM Press; 2015:1055–1081.
- Dowell D, Polgreen PM, Beekmann SE, et al. Dilemmas in the management of syphilis: a survey of infectious diseases experts. *Clin Infect Dis*. 2009;49:1526–1529.
- Ghanem KG, Workowski KA. Management of adult syphilis. *Clin Infect Dis*. 2011;53(suppl 3):S110–S128.
- Smajs D, Norris SJ, Weinstock GM. Genetic diversity in *Treponema pallidum*: implications for pathogenesis, evolution and molecular diagnostics of syphilis and yaws. *Infect Genet Evol*. 2012;12:191–202.
- Staudova B, Strouhal M, Zbanikova M, et al. Whole genome sequence of the *Treponema pallidum* subsp. *endemicum* strain Bosnia A: the genome is related to yaws treponemes but contains few loci similar to syphilis treponemes. *PLoS Negl Trop Dis*. 2014;8:e3261.
- Nechvatal L, Petrosova H, Grillova L, et al. Syphilis-causing strains belong to separate SS14-like or Nichols-like groups as defined by multilocus analysis of 19 *Treponema pallidum* strains. *Int J Med Microbiol*. 2014;304:645–653.
- Hudson EH. Treponematoses in perspective. *Bull World Health Organ*. 1965;32:735–748.
- Hackett CJ. On the origin of the human treponematoses (pinta, yaws, endemic syphilis and venereal syphilis). *Bull World Health Organ*. 1963;29:7–41.
- McGough LJ. Syphilis in history: a response to 2 articles. *Clin Infect Dis*. 2005;41:573–575, author reply 5.
- Harper KN, Zuckerman MK, Harper ML, et al. The origin and antiquity of syphilis revisited: an appraisal of Old World pre-Columbian evidence for treponemal infection. *Am J Phys Anthropol*. 2011;146(suppl 53):99–133.
- Rothschild BM. History of syphilis. *Clin Infect Dis*. 2005;40:1454–1463.
- Mays S, Crane-Kramer G, Bayliss A. Two probable cases of treponemal disease of Medieval date from England. *Am J Phys Anthropol*. 2003;120:133–143.
- von Hunnius TE, Roberts CA, Boylston A, et al. Histological identification of syphilis in pre-Columbian England. *Am J Phys Anthropol*. 2006;129:559–566.
- Barnes I, Thomas MG. Evaluating bacterial pathogen DNA preservation in museum osteological collections. *Proc Biol Sci*. 2006;273:645–653.
- von Hunnius TE, Yang D, Eng B, et al. Digging deeper into the limits of ancient DNA research on syphilis. *J Archaeol Sci*. 2007;34:2091–2100.
- Bouwman AS, Brown TA. The limits of biomolecular paleopathology: ancient DNA cannot be used to study venereal syphilis. *J Archaeol Sci*. 2005;32:703–713.
- Bouwman AS, Brown TA. The limits of biomolecular paleopathology: ancient DNA cannot be used to study venereal syphilis. *J Archaeol Sci*. 2005;32:703–713.
- Gogarten JF, Dux A, Schuenemann VJ, et al. Tools for opening new chapters in the book of *Treponema pallidum* evolutionary history. *Clin Microbiol Infect*. 2016;22:916–921.
- Harper KN, Ocampo PS, Steiner BM, et al. On the origin of the treponematoses: a phylogenetic approach. *PLoS Negl Trop Dis*. 2008;2:e148.
- Mulligan CJ, Norris SJ, Lukehart SA. Molecular studies in *Treponema pallidum* evolution: toward clarity? *PLoS Negl Trop Dis*. 2008;2:e184.
- de Melo FL, de Mello JC, Fraga AM, et al. Syphilis at the crossroad of phylogenetics and paleopathology. *PLoS Negl Trop Dis*. 2010;4:e575.
- Strouhal M, Mikalova L, Havlickova P, et al. Complete genome sequences of two strains of *Treponema pallidum* subsp. *pertenue* from Ghana, Africa: identical genome sequences in samples isolated more than 7 years apart. *PLoS Negl Trop Dis*. 2017;11:e005894.
- Harper KN, Fyumagwa RD, Hoare R, et al. *Treponema pallidum* infection in the wild baboons of East Africa: distribution and genetic characterization of the strains responsible. *PLoS ONE*. 2012;7:e50882.
- Fribourg-Blanc A, Mollaret HH. Natural treponematoses of the African primate. *Primates Med*. 1969;3:113–121.
- Zobanikova M, Strouhal M, Mikalova L, et al. Whole genome sequence of the *Treponema* Fribourg-Blanc: undescribed simian isolate is highly similar to the yaws subspecies. *PLoS Negl Trop Dis*. 2013;7:e2172.
- McGough LJ, Erbeling E. Historical Evidence of syphilis and other treponemes. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006.
- Arora N, Schuenemann VJ, Jager G, et al. Origin of modern syphilis and emergence of a pandemic *Treponema pallidum* cluster. *Nat Microbiol*. 2016;2:16245.
- Radolf JD, Deka RK, Anand A, et al. *Treponema pallidum*, the syphilis spirochete: making a living as a stealth pathogen. *Nat Rev Microbiol*. 2016;14:744–759.
- Grassly NC, Fraser C, Garnett GP. Host immunity and synchronized epidemics of syphilis across the United States. *Nature*. 2005;433:417–421.
- Grassly NC, Fraser C, Garnett GP. Host immunity and synchronized epidemics of syphilis across the United States. *Nature*. 2005;433:417–421.
- Nichols HJ, Hough HH. Demonstration of *Spirochaeta pallida* in the cerebrospinal fluid. From a patient with nervous relapse following the use of salvarsan. *JAMA*. 1913;40:108–110.
- Ghanem KG. Neurosyphilis: a historical perspective and review. *CNS Neurol Ther*. 2010;16:e157–e168.
- Fraser CM, Norris SJ, Weinstock GM, et al. Complete genome sequence of *Treponema pallidum*, the syphilis spirochete. *Science*. 1998;281:375–388.
- Stamm LV, Stapleton JT, Bassford PJ Jr. *In vitro* assay to demonstrate high-level erythromycin resistance of a clinical isolate of *Treponema pallidum*. *Antimicrob Agents Chemother*. 1988;32:164–169.
- Parran T. *Shadow on the Land*. New York: Reynal & Hitchcock; 1937.
- Mahoney JF, Arnold RC, Stermer BL, et al. Penicillin treatment of early syphilis: II. *JAMA*. 1944;126:63–67.
- Rolfs RT, Goldberg M, Sharrar RG. Risk factors for syphilis: cocaine use and prostitution. *Am J Public Health*. 1990;80:853–857.
- Peterman TA, Heffelfinger JD, Swint EB, et al. The changing epidemiology of syphilis. *Sex Transm Dis*. 2005;32:S4–S10.
- MacDonald CJ. Communications: the contribution of the Tuskegee study to medical knowledge. *J Natl Med Assoc*. 1974;66:1–7.
- White RM. Unraveling the Tuskegee study of untreated syphilis. *Arch Intern Med*. 2000;160:585–598.
- Fairchild AL, Bayer R. Uses and abuses of Tuskegee. *Science*. 1999;284:919–921.
- Zenilman J. The Guatemala sexually transmitted disease studies: what happened. *Sex Transm Dis*. 2013;40:277–279.
- Hook EW 3rd. Remembering Thomas Parran, his contributions and missteps going forward: history informs us. *Sex Transm Dis*. 2013;40:281–282.
- Semeniuk I, Reverby S. A shocking discovery. *Nature*. 2010;467:645.
- Obrigton-Tito AJ, Tito RY, Metcalf J, et al. Subsistence strategies in traditional societies distinguish gut microbiomes. *Nat Commun*. 2015;6:6505.
- Gomez A, Petzelkova KJ, Burns MB, et al. Gut Microbiome of coexisting BaAka pygmies and Bantu reflects gradients of traditional subsistence patterns. *Cell Rep*. 2016;14:2142–2153.
- Wilcox RR, Guthrie T. *Treponema pallidum*. A bibliographic review of the morphology, culture, and survival of *T. pallidum* and associated organisms. *Bull WHO*. 1966;35:1–69.
- Miao R, Fieldsteel AH. Genetics of *Treponema*: relationship between *Treponema pallidum* and five cultivable treponemes. *J Bacteriol*. 1978;133:101–107.
- Miao RM, Fieldsteel AH. Genetic relationship between *Treponema pallidum* and *Treponema pertenue*, two noncultivable human pathogens. *J Bacteriol*. 1980;141:427–429.
- Norris SJ, Weinstock GM. Comparative genomics of spirochetes. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006:19–38.
- Seshadri R, Myers GS, Tettelin H, et al. Comparison of the genome of the oral pathogen *Treponema denticola* with other spirochete genomes. *Proc Natl Acad Sci USA*. 2004;101:5646–5651.
- Giacani L, Lukehart SA. The endemic treponematoses. *Clin Microbiol Rev*. 2014;27:89–115.
- Radolf JD, Hazlett KRO, Lukehart SA. Pathogenesis of syphilis. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponemes: Cellular and Molecular Biology*. Norfolk, UK: Caister Academic Press; 2006:197–236.
- Shafit T, Radolf JD, Sanchez PJ, et al. Congenital syphilis. In: Holmes KK, Sparling PF, Stamm WE, et al, eds. *Sexually Transmitted Diseases*. New York: McGraw Hill; 2008:1577–1612.
- Smajs D, Norris SJ, Weinstock GM. Genetic diversity in *Treponema pallidum*: implications for pathogenesis, evolution and molecular diagnostics of syphilis and yaws. *Infect Genet Evol*. 2012;12:191–202.
- Cejkova D, Zobanikova M, Chen L, et al. Whole genome sequences of three *Treponema pallidum* ssp. *pertenue* strains: yaws and syphilis treponemes differ in less than 0.2% of the genome sequence. *PLoS Negl Trop Dis*. 2012;6:e1471.
- Knauf S, Batamuzi EK, Mlengeya T, et al. *Treponema* infection associated with genital ulceration in wild baboons. *Vet Pathol*. 2012;49:292–303.
- Cox DL. Culture of *Treponema pallidum*. *Methods Enzymol*. 1994;236:390–405.
- Edmondson DG, Hu B, Norris SJ. Long-term *In vitro* culture of the syphilis spirochete *Treponema pallidum* subsp. *pallidum*. *MBio*. 2018;9.
- Lukehart SA, Marra CM. Isolation and laboratory maintenance of *Treponema pallidum*. *Curr Protoc Microbiol*. 2007.
- Turner TB, Hardy PH, Newman B. Infectivity tests in syphilis. *Br J Vener Dis*. 1969;45:183–195.
- Izard J, Renken C, Hsieh CE, et al. Cryo-electron tomography elucidates the molecular architecture of *Treponema pallidum*, the syphilis spirochete. *J Bacteriol*. 2009;191:7566–7580.
- Liu J, Howell JK, Bradley SD, et al. Cellular architecture of *Treponema pallidum*: novel flagellum, periplasmic cone, and cell envelope as revealed by cryo electron tomography. *J Mol Biol*. 2010;403:546–561.
- Thomas DD, Navab M, Haake DA, et al. *Treponema pallidum* invades intercellular junctions of endothelial cell monolayers. *Proc Natl Acad Sci USA*. 1988;85:3608–3612.
- Harman M, Vig DK, Radolf JD, et al. Viscous dynamics of Lyme disease and syphilis spirochetes reveal flagellar torque and drag. *Biophys J*. 2013;105:2273–2280.
- Paster BJ, Dewhirst FE. The phylogenetic diversity of the genus *Treponema*. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006:9–18.
- Radolf JD, Kumar S. The *Treponema pallidum* outer membrane. *Curr Top Microbiol Immunol*. 2017.
- Radolf JD, Norgard MV, Schulz WW. Outer membrane ultrastructure explains the limited antigenicity of virulent *Treponema pallidum*. *Proc Natl Acad Sci USA*. 1989;86:2051–2055.
- Walker EM, Zampighi GA, Blanco DR, et al. Demonstration of rare protein in the outer membrane of *Treponema pallidum* subsp. *pallidum* by freeze-fracture analysis. *J Bacteriol*. 1989;171:5005–5011.
- Liu J, Howell JK, Bradley SD, et al. Cellular architecture of *Treponema pallidum*: novel flagellum, periplasmic cone, and cell envelope as revealed by cryo electron tomography. *J Mol Biol*. 2010;403:546–561.
- Cox DL, Luthra A, Dunham-Ems S, et al. Surface immunolabeling and consensus computational framework to identify candidate rare outer membrane proteins of *Treponema pallidum*. *Infect Immun*. 2010;78:5178–5194.
- Sell S, Norris SJ. The biology, pathology, and immunology of syphilis. *Int Rev Exp Pathol*. 1983;24:203–276.
- Salazar JC, Hazlett KR, Radolf JD. The immune response to infection with *Treponema pallidum*, the stealth pathogen. *Microbes Infect*. 2002;4:1133–1140.
- Seshadri R, Myers GS, Tettelin H, et al. Comparison of the genome of the oral pathogen *Treponema denticola* with other spirochete genomes. *Proc Natl Acad Sci USA*. 2004;101:5646–5651.
- Correia FF, Plummer AR, Paster BJ, et al. Genome size of human oral *Treponema* species by pulsed-field gel electrophoresis. *Oral Microbiol Immunol*. 2004;19:129–131.
- Magnuson HJ, Eagle H, Fleischman R. The minimal infectious inoculum of *Spirochaeta pallida* (Nichols strain) and a consideration of its rate of multiplication *in vivo*. *Am J Syph Gonorrhea Vener Dis*. 1948;32:1–18.
- Norris SJ, Cox DL, Weinstock GM. Biology of *Treponema pallidum*: correlation of functional activities with genome sequence data. *J Mol Microbiol Biotechnol*. 2001;3:37–62.
- Pillay A, Liu H, Ebrahim S, et al. Molecular typing of *Treponema pallidum* in South Africa: cross-sectional studies. *J Clin Microbiol*. 2002;40:256–258.
- Marra C, Sahi S, Tantalo L, et al. Enhanced molecular typing of *Treponema pallidum*: geographical distribution of strain types and association with neurosyphilis. *J Infect Dis*. 2010;202:1380–1388.
- Peng RR, Wang AL, Li J, et al. Molecular typing of *Treponema pallidum*: a systematic review and meta-analysis. *PLoS Negl Trop Dis*. 2011;5:e1273.

87. Kumar S, Caimano MJ, Anand A, et al. Sequence variation of rare outer membrane protein beta-barrel domains in clinical strains provides insights into the evolution of *Treponema pallidum* subsp. *pallidum*, the syphilis spirochete. *MBio*. 2018;9.
88. Smolak A, Rowley J, Nagelkerke N, et al. Trends and predictors of syphilis prevalence in the general population: global pooled analyses of 1103 prevalence measures including 136 million syphilis tests. *Clin Infect Dis*. 2017.
89. Fenton KA, Breban R, Vardavas R, et al. Infectious syphilis in high-income settings in the 21st century. *Lancet Infect Dis*. 2008;8:244–253.
90. Centers for Disease Control and Prevention. 2011 Sexually Transmitted Disease Surveillance. <http://www.cdc.gov/STD/stats11/default.htm>. Accessed April 1, 2014.
91. Mayer KH, Mimiaga MJ. Resurgent syphilis in the United States: urgent need to address an evolving epidemic. *Ann Intern Med*. 2011;155:192–193.
92. de Voux A, Kidd S, Grey JA, et al. State-specific rates of primary and secondary syphilis among men who have sex with men - United States, 2015. *MMWR Morb Mortal Wkly Rep*. 2017;66:349–354.
93. French P, Gomberg M, Janier M, et al. IUSTI: 2008 European guidelines on the management of syphilis. *Int J STD AIDS*. 2009;20:300–309.
94. Centers for Disease Control and Prevention. National notifiable diseases surveillance system (NNDSS). <https://www.cdc.gov/nndss/conditions/syphilis/>. Accessed December 13, 2018.
95. Jones DL, Irwin KL, Inciardi J, et al. The high-risk sexual practices of crack-smoking sex workers recruited from the streets of three American cities. The Multicenter Crack Cocaine and HIV Infection Study Team. *Sex Transm Dis*. 1998;25:187–193.
96. Zetola NM, Klausner JD. Syphilis and HIV infection: an update. *Clin Infect Dis*. 2007;44:1222–1228.
97. European Centre for Disease Prevention and Control (ECDC). *Sexually Transmitted Infections in Europe, 1990-2010*. Stockholm: ECDC; 2012.
98. European Centre for Disease Prevention and Control. Annual Epidemiological Report 2016 – Syphilis. <http://ecdc.europa.eu/en/healthtopics/Syphilis/Pages/Annuairepidemiologicalreport.aspx>. Accessed June 2, 2018.
99. Scott CM, Flint SR. Oral syphilis—re-emergence of an old disease with oral manifestations. *Int J Oral Maxillofac Surg*. 2005;34:58–63.
100. Herbert LJ, Middleton SL. An estimate of syphilis incidence in Eastern Europe. *J Glob Health*. 2012;2:10402.
101. Bailey H, Turkova A, Thorne C. Syphilis, hepatitis C and HIV in Eastern Europe. *Curr Opin Infect Dis*. 2017;30:93–100.
102. Tucker JD, Yin YP, Wang B, et al. An expanding syphilis epidemic in China: epidemiology, behavioural risk and control strategies with a focus on low-tier female sex workers and men who have sex with men. *Sex Transm Infect*. 2011;87(suppl 2):ii16–ii18.
103. Tucker JD, Cohen MS. China's syphilis epidemic: epidemiology, proximate determinants of spread, and control responses. *Curr Opin Infect Dis*. 2011;24:50–55.
104. Sanchez J, Volquez C, Totten PA, et al. The etiology and management of genital ulcers in the Dominican Republic and Peru. *Sex Transm Dis*. 2002;29:559–567.
105. Lopes F, Latorre MR, Campos Pignatari AC, et al. HIV, HPV, and syphilis prevalence in a women's penitentiary in the city of Sao Paulo, 1997–1998. *Cad Saude Publica*. 2001;17:1473–1480.
- 105a. Pan American Health Organization. <http://new.paho.org>. Accessed January 2018.
106. Mejia A, Bautista CT, Leal L, et al. Syphilis infection among female sex workers in Colombia. *J Immigr Minor Health*. 2009;11:92–98.
107. Cruz AR, Pillay A, Zuluaga AV, et al. Secondary syphilis in Cali, Colombia: new concepts in disease pathogenesis. *PLoS Negl Trop Dis*. 2010;4:e690.
108. Zoni AC, Gonzalez MA, Sjogren HW. Syphilis in the most at-risk populations in Latin America and the Caribbean: a systematic review. *Int J Infect Dis*. 2013;17:e84–e92.
109. Chow JY, Konda KA, Calvo GM, et al. Demographics, behaviors, and sexual health characteristics of high risk men who have sex with men and transgender women who use social media to meet sex partners in Lima, Peru. *Sex Transm Dis*. 2017;44:143–148.
110. Kojima N, Park H, Konda KA, et al. The PICASSO Cohort: baseline characteristics of a cohort of men who have sex with men and male-to-female transgender women at high risk for syphilis infection in Lima, Peru. *BMC Infect Dis*. 2017;17:255.
111. Jaffe HW, Choi K, Thomas PA, et al. National case-control study of Kaposi's sarcoma and *Pneumocystis carinii* pneumonia in homosexual men: part 1. Epidemiologic results. *Ann Intern Med*. 1983;99:145–151.
112. Hutchinson CM, Rompalo AM, Reichart CA, et al. Characteristics of patients with syphilis attending Baltimore STD clinics. Multiple high-risk subgroups and interactions with human immunodeficiency virus infection. *Arch Intern Med*. 1991;151:511–516.
113. Ansell DA, Hu TC, Straus M, et al. HIV and syphilis seroprevalence among clients with sexually transmitted diseases attending a walk-in clinic at Cook County Hospital. *Sex Transm Dis*. 1994;21:93–96.
114. Katz KA, Pillay A, Ahrens K, et al. Molecular epidemiology of syphilis—San Francisco, 2004–2007. *Sex Transm Dis*. 2010;37:660–663.
115. Page-Shafer K, Shiboski CH, Osmond DH, et al. Risk of HIV infection attributable to oral sex among men who have sex with men and in the population of men who have sex with men. *AIDS*. 2002;16:2350–2352.
116. Peterman TA, Su J, Bernstein KT, et al. Syphilis in the United States: on the rise? *Expert Rev Anti Infect Ther*. 2015;13:161–168.
117. Cohen SE, Chew Ng RA, Katz KA, et al. Repeat syphilis among men who have sex with men in California, 2002–2006: implications for syphilis elimination efforts. *Am J Public Health*. 2012;102:e1–e8.
118. Peterman TA, Newman DR, Maddox L, et al. High risk for HIV following syphilis diagnosis among men in Florida, 2000–2011. *Public Health Rep*. 2014;129:164–169.
119. Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Infect*. 1999;75:3–17.
120. Wasserheit JN. Epidemiological synergy. Interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis*. 1992;19:61–77.
121. Tosca A, Lehou J, Hatjivasiliou M, et al. Infiltrate of syphilitic lesions before and after treatment. *Genitourin Med*. 1988;64:289–293.
122. Van Voorhis WC, Barrett LK, Koelle DM, et al. Primary and secondary syphilis lesions contain mRNA for Th1 cytokines. *J Infect Dis*. 1996;173:491–495.
123. McBroom RL, Styles AR, Chiu MJ, et al. Secondary syphilis in persons infected with and not infected with HIV-1: a comparative immunohistologic study. *Am J Dermatopathol*. 1999;21:432–441.
124. Salazar JC, Cruz AR, Pope CD, et al. *Treponema pallidum* elicits innate and adaptive cellular immune responses in skin and blood during secondary syphilis: a flow-cytometric analysis. *J Infect Dis*. 2007;195:879–887.
125. Sheffield JS, Wendel GD Jr, McIntire DD, et al. Effect of genital ulcer disease on HIV-1 coreceptor expression in the female genital tract. *J Infect Dis*. 2007;196:1509–1516.
126. Theus SA, Harrich DA, Gaynor R, et al. *Treponema pallidum*, lipoproteins, and synthetic lipoprotein analogues induce human immunodeficiency virus type 1 gene expression in monocytes via NF- κ B activation. *J Infect Dis*. 1998;177:941–950.
127. Jarzebowski W, Caumes E, Dupin N, et al. Effect of early syphilis infection on plasma viral load and CD4 cell count in human immunodeficiency virus-infected men: results from the FHDH-ANRS CO4 cohort. *Arch Intern Med*. 2012;172:1237–1243.
128. Newman L, Kamb M, Hawkes S, et al. Global estimates of syphilis in pregnancy and associated adverse outcomes: analysis of multinational antenatal surveillance data. *PLoS Med*. 2013;10:e1001396.
129. Centers for Disease Control and Prevention. 2016 Sexually transmitted diseases surveillance. Updated 2017. <https://www.cdc.gov/std/stats16/figures/44.htm>.
130. Gueff B, Rosahn PD. Experimental mouse syphilis, a critical review of the literature. *Am J Syphilology*. 1948;32:59–88.
131. Silver AC, Dunne DW, Zeiss CJ, et al. MyD88 deficiency markedly worsens tissue inflammation and bacterial clearance in mice infected with *Treponema pallidum*, the agent of syphilis. *PLoS ONE*. 2013;8:e71388.
132. Turner TB, Hollander DH. *Biology of the Treponematoses*. Geneva: World Health Organization; 1957.
133. Tantaló LC, Lukehart SA, Marra CM. *Treponema pallidum* strain-specific differences in neuroinvasion and clinical phenotype in a rabbit model. *J Infect Dis*. 2005;191:75–80.
134. Cruz AR, Ramirez LG, Zuluaga AV, et al. Immune evasion and recognition of the syphilis spirochete in blood and skin of secondary syphilis patients: two immunologically distinct compartments. *PLoS Negl Trop Dis*. 2012;6:e1717.
135. Lafont RE, Lukehart SA. Biological basis for syphilis. *Clin Microbiol Rev*. 2006;19:29–49.
136. Cameron CE. The *T. pallidum* outer membrane and outer membrane proteins. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006:237–266.
137. Radolf JD. *Treponema pallidum* and the quest for outer membrane proteins. *Mol Microbiol*. 1995;16:1067–1073.
138. Smajs D, McKevitt M, Howell JK, et al. Transcriptome of *Treponema pallidum*: gene expression profile during experimental rabbit infection. *J Bacteriol*. 2005;187:1866–1874.
139. Osbak KK, Houston S, Lithgow KV, et al. Characterizing the syphilis-causing *Treponema pallidum* ssp. *pallidum* proteome using complementary mass spectrometry. *PLoS Negl Trop Dis*. 2016;10:e0004988.
140. Magnuson HJ, Thomas EW, Olansky S, et al. Inoculation syphilis in human volunteers. *Medicine (Baltimore)*. 1956;35:33–82.
141. Quatresooz P, Pierard GE. Skin homing of *Treponema pallidum* in early syphilis: an immunohistochemical study. *Appl Immunohistochem Mol Morphol*. 2009;17:47–50.
142. Schmidt PJ. Syphilis, a disease of direct transfusion. *Transfusion*. 2001;41:1069–1071.
143. Adegoke AO, Akanni O, Dirisu J. Risk of transfusion-transmitted syphilis in a tertiary hospital in Nigeria. *N Am J Med Sci*. 2011;3:78–81.
144. Mahoney JF, Bryant KK. Time element in penetration of genital mucosa by *Treponema pallidum*. *J Vener Dis Inf*. 1934;15:1–5.
145. Stokes JH, Beerman H, Ingraham NR. *Modern Clinical Syphilology*. 3rd ed. Philadelphia: W.B. Saunders; 1944.
146. Brown WH, Pearce L. A note on the dissemination of *Spirochaeta pallida* from the primary focus of infection. *Arch Derm Syphilol*. 1920;2:470–472.
147. Salazar JC, Rathi A, Michael NL, et al. Assessment of the kinetics of *Treponema pallidum* dissemination into blood and tissues in experimental syphilis by real-time quantitative PCR. *Infect Immun*. 2007;75:2954–2958.
148. Raiziss GW, Severac M. Rapidity with which *Spirochaeta pallida* invade the bloodstream. *Arch Derm Syphilol*. 1937;35:1101–1109.
149. Merritt HH, Adams RD, Solomon HC. *Neurosyphilis*. New York: Oxford University Press; 1946.
150. Lukehart SA, Hook EW 3rd, Baker-Zander SA, et al. Invasion of the central nervous system by *Treponema pallidum*: implications for diagnosis and treatment. *Ann Intern Med*. 1988;109:855–862.
151. Rolfs RT, Joesoef MR, Hendershot EF, et al. A randomized trial of enhanced therapy for early syphilis in patients with and without human immunodeficiency virus infection. The Syphilis and HIV Study Group. *N Engl J Med*. 1997;337:307–314.
152. Cameron CE, Brouwer NL, Tisch LM, et al. Defining the interaction of the *Treponema pallidum* adhesin Tp0751 with laminin. *Infect Immun*. 2005;73:7485–7494.
153. Radolf JD, Hazlett KRO, Lukehart SA. Pathogenesis of Syphilis. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponemes: Cellular and Molecular Biology*. Norfolk, UK: Caister Academic Press; 2006:197–236.
154. Harman MW, Dunham-Emms SM, Caimano MJ, et al. The heterogeneous motility of the Lyme disease spirochete in gelatin mimics dissemination through tissue. *Proc Natl Acad Sci USA*. 2012;109:3059–3064.
155. Thomas DD, Baseman JB, Alderete JF. Fibronectin tetrapeptide is target for syphilis spirochete cytoadherence. *J Exp Med*. 1985;162:1715–1719.
156. Cameron CE. Identification of a *Treponema pallidum* laminin-binding protein. *Infect Immun*. 2003;71:2525–2533.
157. Cameron CE, Brown EL, Kuroiwa JM, et al. *Treponema pallidum* fibronectin-binding proteins. *J Bacteriol*. 2004;186:7019–7022.
158. Brinkman MB, McGill MA, Pettersson J, et al. A novel *Treponema pallidum* antigen, TP0136, is an outer membrane protein that binds human fibronectin. *Infect Immun*. 2008;76:1848–1857.
159. Ke W, Molini BJ, Lukehart SA, et al. *Treponema pallidum* subsp. *pallidum* TP0136 protein is heterogeneous among isolates and binds cellular and plasma fibronectin via its NH2-terminal end. *PLoS Negl Trop Dis*. 2015;9:e0003662.
160. Chan K, Nasereddin T, Alter L, et al. *Treponema pallidum* lipoprotein TP0435 expressed in *Borrelia burgdorferi* produces multiple surface/periplasmic isoforms and mediates adherence. *Sci Rep*. 2016;6:25593.
161. Parker ML, Houston S, Petrosova H, et al. The structure of *Treponema pallidum* TP0751 (Pallilysin) reveals a non-canonical lipocalin fold that mediates adhesion to extracellular matrix components and interactions with host cells. *PLoS Pathog*. 2016;12:e100519.
162. Brautigam CA, Deka RK, Liu WZ, et al. Insights into the potential function and membrane organization of the TP0435 (Tp17) lipoprotein from *Treponema pallidum* derived from structural and biophysical analyses. *Protein Sci*. 2015;24:11–19.

163. Hayes NS, Muse KE, Collier AM, et al. Parasitism by virulent *Treponema pallidum* of host cell surfaces. *Infect Immun*. 1977;17:174–186.
164. Charon NW, Li C, Goldstein SE. The beguiling motility of the genus *Treponema*. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006:127–170.
165. Kao WA, Petrosova H, Ebady R, et al. Identification of Tp0751 (Pallilysin) as a *Treponema pallidum* vascular adhesion by heterologous expression in the Lyme disease spirochete. *Sci Rep*. 2017;7:1538.
166. Sykes JA, Miller JN, Kalan AJ. *Treponema pallidum* within cells of a primary chancre from a human female. *Br J Vener Dis*. 1974;50:40–44.
167. Sykes JA, Kalan J. Intracellular *Treponema pallidum* in cells of a syphilitic lesion of the uterine cervix. *Am J Obstet Gynecol*. 1975;122:361–367.
168. Lafond RE, Lukehart SA. Biological basis for syphilis. *Clin Microbiol Rev*. 2006;19:29–49.
169. Radolf JD, Lukehart SA. Immunology of syphilis. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponemes: Cellular and Molecular Biology*. Norfolk, UK: Caister Academic Press; 2006:285–322.
170. Hawley KL, Cruz AR, Benjamin SJ, et al. IFN γ enhances CD64-potentiated phagocytosis of *Treponema pallidum* opsonized with human syphilitic serum by human macrophages. *Front Immunol*. 2017;8:1227.
171. Salazar JC, Hazlett KR, Radolf JD. The immune response to infection with *Treponema pallidum*, the stealth pathogen. *Microbes Infect*. 2002;4:1133–1140.
172. Moore MW, Cruz AR, LaVake CJ, et al. Phagocytosis of *Borrelia burgdorferi* and *Treponema pallidum* potentiates innate immune activation and induces gamma interferon production. *Infect Immun*. 2007;75:2046–2062.
173. Lukehart SA. Scientific monogamy: thirty years dancing with the same bug: 2007 Thomas Parran Award Lecture. *Sex Transm Dis*. 2008;35:2–7.
174. Stary G, Klein I, Bruggen MC, et al. Host defense mechanisms in secondary syphilitic lesions: a role for IFN γ /IL-17-producing CD8 $^{+}$ T cells? *Am J Pathol*. 2010;177:2421–2432.
175. Cox DL, Akins DR, Porcella SE, et al. *Treponema pallidum* in gel microdroplets: a novel strategy for investigation of treponemal molecular architecture. *Mol Microbiol*. 1995;15:1151–1164.
176. Myint M, Bashiri H, Harrington RD, et al. Relapse of secondary syphilis after benzathine penicillin G: molecular analysis. *Sex Transm Dis*. 2004;31:196–199.
177. Centurion-Lara A. Antigenic variation in *Treponema pallidum*. In: Radolf JD, Lukehart SA, eds. *Pathogenic Treponema: Molecular and Cellular Biology*. Norwich, UK: Caister Academic Press; 2006:267–284.
178. Ho EL, Lukehart SA. Syphilis: using modern approaches to understand an old disease. *J Clin Invest*. 2011;121:4584–4592.
179. Giacani L, Brandt SL, Ke W, et al. Transcription of TP0126, *Treponema pallidum* putative OmpW homolog, is regulated by the length of a homopolymeric guanosine repeat. *Infect Immun*. 2015;83:2275–2289.
180. Pinto M, Borges V, Antelo M, et al. Genome-scale analysis of the non-cultivable *Treponema pallidum* reveals extensive within-patient genetic variation. *Nat Microbiol*. 2016;2:16190.
181. Lukehart SA, Shaffer JM, Baker-Zander SA. A subpopulation of *Treponema pallidum* is resistant to phagocytosis: possible mechanism of persistence. *J Infect Dis*. 1992;166:1449–1453.
182. Crissey JT, Denenholz DA. Syphilis. *Clin Dermatol*. 1984;2:1–166.
183. Martin-Ezquerro G, Fernandez-Casado A, Barco D, et al. *Treponema pallidum* distribution patterns in mucocutaneous lesions of primary and secondary syphilis: an immunohistochemical and ultrastructural study. *Hum Pathol*. 2009;40:624–630.
184. Quatresooz P, Pierard GE. Skin homing of *Treponema pallidum* in early syphilis: an immunohistochemical study. *Appl Immunohistochem Mol Morphol*. 2009;17:47–50.
185. Merritt HH, Adams RD, Solomon HC. *Neurosyphilis*. New York: Oxford University Press; 1946.
186. Abell E, Marks R, Jones EW. Secondary syphilis: a clinico-pathological review. *Br J Dermatol*. 1975;93:53–61.
187. Jordaan HF. Secondary syphilis. A clinicopathological study. *Am J Dermatopathol*. 1988;10:399–409.
188. Jeerapaet P, Ackerman AB. Histologic patterns of secondary syphilis. *Arch Dermatol*. 1973;107:373–377.
189. Turner DR. Lymphadenopathy in secondary syphilis. *J Pathol*. 1971;104:x.
190. Jorizzo JL, McNeely MC, Baughn RE, et al. Role of circulating immune complexes in human secondary syphilis. *J Infect Dis*. 1986;153:1014–1022.
191. Bhorade MS, Carag HB, Lee HJ, et al. Nephropathy of secondary syphilis. A clinical and pathological spectrum. *JAMA*. 1971;216:1159–1166.
192. Gamble CN, Reardan JB. Immunopathogenesis of syphilitic glomerulonephritis. Elution of antitreponemal antibody from glomerular immune-complex deposits. *N Engl J Med*. 1975;292:449–454.
193. Fargen KM, Alvernia JE, Lin CS, et al. Cerebral syphilitic gummatous: a case presentation and analysis of 156 reported cases. *Neurosurgery*. 2009;64:568–575, discussion 575–76.
194. Jackman JD Jr, Radolf JD. Cardiovascular syphilis. *Am J Med*. 1989;87:425–433.
195. Roberts WC, Ko JM, Vowels TJ. Natural history of syphilitic aortitis. *Am J Cardiol*. 2009;104:1578–1587.
196. Guyure KA. Infections. In: Prayson RA, ed. *Neuropathology*. Philadelphia: Churchill Livingstone; 2005:287–338.
197. Brown E, Gray F. Bacterial infections. In: Love S, Louis DN, Ellison DW, eds. *Greenfield's Neuropathology*. London: Hodder Arnold; 2008:1391–1445.
198. Wenhai L, Jianzhong Z, Cao Y. Detection of *Treponema pallidum* in skin lesions of secondary syphilis and characterization of the inflammatory infiltrate. *Dermatology*. 2004;208:94–97.
199. Leader BT, Godornes C, VanVoorhis WC, et al. CD4 $^{+}$ lymphocytes and γ interferon predominate in local immune responses in early experimental syphilis. *Infect Immun*. 2007;75:3021–3026.
200. van Voorhis WC, Barrett LK, Nasio JM, et al. Lesions of primary and secondary syphilis contain activated cytolytic T cells. *Infect Immun*. 1996;64:1048–1050.
201. Stary G, Klein I, Bruggen MC, et al. Host defense mechanisms in secondary syphilitic lesions: a role for IFN γ -IL-17-producing CD8 $^{+}$ T cells? *Am J Pathol*. 2010;177:2421–2432.
202. Babolin C, Amedei A, Ozolins D, et al. T β F1 from *Treponema pallidum* activates inflammasome and promotes the development of regulatory T cells. *J Immunol*. 2011;187:1377–1384.
203. Clark EG, Danbolt N. The Oslo study of the natural history of untreated syphilis: an epidemiologic investigation based on a restudy of the Boeck-Bruusgaard material: a review and appraisal. *J Chron Dis*. 1955;2:311–344.
204. White RM. Unraveling the Tuskegee study of untreated syphilis. *Arch Intern Med*. 2000;160:585–598.
205. Peters JJ, Peers JH, Olansky S, et al. Untreated syphilis in the male Negro: pathologic findings in syphilitic and nonsyphilitic patients. *J Chronic Dis*. 1955;1:127–148.
206. Rockwell DH, Yobs AR, Moore MB Jr. The Tuskegee study of untreated syphilis: the 30th year of observation. *Arch Intern Med*. 1964;114:792–798.
207. Rosahn PD. Studies in syphilis: the end results of untreated syphilis. *J Vener Dis Inf*. 1946;27:293–301.
208. Sena AC, Zhang XH, Li T, et al. A systematic review of syphilis serological treatment outcomes in HIV-infected and HIV-uninfected persons: rethinking the significance of serological non-responsiveness and the serofast state after therapy. *BMC Infect Dis*. 2015;15:479.
209. Golden MR, Marra CM, Holmes KK. Update on syphilis: resurgence of an old problem. *JAMA*. 2003;290:1510–1514.
210. Sparling PF, Swartz MN, Musher DM, et al. Clinical manifestations of syphilis. In: Holmes KK, Sparling PF, Stamm WE, et al, eds. *Sexually Transmitted Diseases*. 4th ed. New York: McGraw-Hill; 2008:661–684.
211. Hook EWR. Syphilis. *Lancet*. 2017;389:1550–1557.
212. Garnett GP, Aral SO, Hoyle DV, et al. The natural history of syphilis. Implications for the transmission dynamics and control of infection. *Sex Transm Dis*. 1997;24:185–200.
213. Merritt HH, Moore M. Acute syphilitic meningitis. *Medicine (Baltimore)*. 1935;14:119–183.
214. Simon RP. Neurosyphilis. *Arch Neurol*. 1985;42:606–613.
215. Collis TK, Celum CL. The clinical manifestations and treatment of sexually transmitted diseases in human immunodeficiency virus-positive men. *Clin Infect Dis*. 2001;32:611–622.
216. Musher DM. Syphilis, neurosyphilis, penicillin, and AIDS. *J Infect Dis*. 1991;163:1201–1206.
217. Musher DM, Hamill RJ, Baughn RE. Effect of human immunodeficiency virus (HIV) infection on the course of syphilis and on the response to treatment. *Ann Intern Med*. 1990;113:872–881.
218. Rompalo AM, Joeseof MR, O'Donnell JA, et al. Clinical manifestations of early syphilis by HIV status and gender: results of the syphilis and HIV study. *Sex Transm Dis*. 2001;28:158–165.
219. Golden MR, Marra CM, Holmes KK. Update on syphilis: resurgence of an old problem. *JAMA*. 2003;290:1510–1514.
220. Farhi D, Dupin N. Management of syphilis in the HIV-infected patient: facts and controversies. *Clin Dermatol*. 2010;28:539–545.
221. Ghanem KG, Workowski KA. Management of adult syphilis. *Clin Infect Dis*. 2011;53(suppl 3):S110–S128.
222. Palacios R, Jimenez-Onate F, Aguilar M, et al. Impact of syphilis infection on HIV viral load and CD4 cell counts in HIV-infected patients. *J Acquir Immune Defic Syndr*. 2007;44:356–359.
223. Kofoed K, Gerstoft J, Mathiesen LR, et al. Syphilis and human immunodeficiency virus (HIV)-1 coinfection: influence on CD4 T-cell count, HIV-1 viral load, and treatment response. *Sex Transm Dis*. 2006;33:143–148.
224. Buchacz K, Patel P, Taylor M, et al. Syphilis increases HIV viral load and decreases CD4 cell counts in HIV-infected patients with new syphilis infections. *AIDS*. 2004;18:2075–2079.
225. Jarzabowski W, Caumes E, Dupin N, et al. Effect of early syphilis infection on plasma viral load and CD4 cell count in human immunodeficiency virus-infected men: results from the FHDH-ANRS CO4 cohort. *Arch Intern Med*. 2012;172:1237–1243.
226. Sadiq ST, McSorley J, Copas AJ, et al. The effects of early syphilis on CD4 counts and HIV-1 RNA viral loads in blood and semen. *Sex Transm Infect*. 2005;81:380–385.
227. Manfredi R, Sabbatani S, Pocaterra D, et al. Syphilis does not seem to involve virological and immunological course of concurrent HIV disease. *AIDS*. 2006;20:305–306.
228. Cohen SE, Klausner JD, Engelman J, et al. Syphilis in the modern era: an update for physicians. *Infect Dis Clin North Am*. 2013;27:705–722.
229. Crissey JT, Denenholz DA. Syphilis. *Clin Dermatol*. 1984;2:1–166.
230. Chapel TA. The variability of syphilitic chancres. *Sex Transm Dis*. 1978;5:68–70.
231. Fiumara NJ. Treatment of primary and secondary syphilis. Serological response. *JAMA*. 1980;243:2500–2502.
232. Flynn TR, Hunter GJ, Johnson MM. Case records of the Massachusetts General Hospital. Case 6-2010. A 37-year-old man with a lesion on the tongue. *N Engl J Med*. 2010;362:740–748.
233. Mindel A, Tovey SJ, Timmins DJ, et al. Primary and secondary syphilis, 20 years' experience. 2. Clinical features. *Genitourin Med*. 1989;65:1–3.
234. Cruz AR, Pillay A, Zuluaga AV, et al. Secondary syphilis in cali, Colombia: new concepts in disease pathogenesis. *PLoS Negl Trop Dis*. 2010;4:e690.
235. Chapel TA. The signs and symptoms of secondary syphilis. *Sex Transm Dis*. 1980;7:161–164.
236. Fiumara NJ. Treatment of secondary syphilis: an evaluation of 204 patients. *Sex Transm Dis*. 1977;4:96–99.
237. Tucker JD, Shah S, Jarell AD, et al. Lues maligna in early HIV infection case report and review of the literature. *Sex Transm Dis*. 2009;36:512–514.
238. Pleimes M, Hartschuh W, Kutzner H, et al. Malignant syphilis with ocular involvement and organism-depleted lesions. *Clin Infect Dis*. 2009;48:83–85.
239. Cid PM, Cudos ES, Zamora Vargas FX, et al. Pathologically confirmed malignant syphilis using immunohistochemical staining: report of 3 cases and review of the literature. *Sex Transm Dis*. 2014;41:94–97.
240. Jalili A, Mosleh M, Grabmeier-Pfistershammer K, et al. Malignant syphilis in a HIV infected patient. *Sex Transm Dis*. 2015;42:223–225.
241. Ridruejo E, Mordoh A, Herrera F, et al. Severe cholestatic hepatitis as the first symptom of secondary syphilis. *Dig Dis Sci*. 2004;49:1401–1404.
242. Mylona EE, Baraboutis IG, Papastamopoulos V, et al. Gastric syphilis: a systematic review of published cases of the last 50 years. *Sex Transm Dis*. 2010;37:177–183.
243. Chen CY, Chi KH, George RW, et al. Diagnosis of gastric syphilis by direct immunofluorescence staining and real-time PCR testing. *J Clin Microbiol*. 2006;44:3452–3456.
244. Aldave AJ, King JA, Cunningham ET Jr. Ocular syphilis. *Curr Opin Ophthalmol*. 2001;12:433–441.
245. Pless ML, Kroschinsky D, LaRocque RC, et al. Case records of the Massachusetts General Hospital. Case 26-2010. A 54-year-old man with loss of vision and a rash. *N Engl J Med*. 2010;363:865–874.
246. Moore JE. Syphilitic Iritis. A study of 249 patients. *Am J Ophthalmol*. 1931;14:110–122.
247. Dutta Majumder P, Chen EJ, Shah J, et al. Ocular syphilis: an update. *Ocul Immunol Inflamm*. 2017;1–9.
248. Radolf JD, Kaplan RP. Unusual manifestations of secondary syphilis and abnormal humoral immune response to *Treponema pallidum* antigens in a homosexual man with asymptomatic human immunodeficiency virus infection. *J Am Acad Dermatol*. 1988;18:423–428.
249. Yimtae K, Srirompotong S, Lertsukprasert K. Otopharyngitis: a review of 85 cases. *Otolaryngol Head Neck Surg*. 2007;136:67–71.
250. Rademacher SE, Radolf JD. Prominent osseous and unusual dermatologic manifestations of early syphilis in two patients with discordant serological statuses for human immunodeficiency virus infection. *Clin Infect Dis*. 1996;23:462–467.

251. Dismukes WE, Delgado DG, Mallernee SV, et al. Destructive bone disease in early syphilis. *JAMA*. 1976;236:2646–2648.
252. Reginaldo AJ, Schumacher HR, Jimenez S, et al. Synovitis in secondary syphilis. Clinical, light, and electron microscopic studies. *Arthritis Rheum*. 1979;22:170–176.
253. David G, Perpoint T, Bobieux A, et al. Secondary pulmonary syphilis: report of a likely case and literature review. *Clin Infect Dis*. 2006;42:e11–e15.
254. Crouzy F, Alvarez V, Gex G, et al. Unusual presentations and pitfalls of secondary syphilis: osteitis, pneumonia and malignancy. *BMJ Case Rep*. 2015;2015.
255. Jeny F, Fargelat A, Laurent-Roussel S, et al. Pulmonary consolidations due to secondary syphilis with positive bronchial washing immunohistochemistry. *Am J Respir Crit Care Med*. 2016;193:1061–1062.
256. Hooshmand H, Escobar MR, Kopf SW. Neurosyphilis. A study of 241 patients. *JAMA*. 1972;219:726–729.
257. Chahine LM, Khoriaty RN, Tomford WJ, et al. The changing face of neurosyphilis. *Int J Stroke*. 2011;6:136–143.
258. Ghanem KG. REVIEW: neurosyphilis: a historical perspective and review. *CNS Neurosci Ther*. 2010;16:e157–e168.
259. Marra C, Sahi S, Tantal L, et al. Enhanced molecular typing of *Treponema pallidum*: geographical distribution of strain types and association with neurosyphilis. *J Infect Dis*. 2010;202:1380–1388.
260. Marra CM, Sahi SK, Tantal LC, et al. Toll-like receptor polymorphisms are associated with increased neurosyphilis risk. *Sex Transm Dis*. 2014;41:440–446.
261. Workowski KA, Bolan GA, Centers for Disease C, Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recomm Rep*. 2015;64:1–137.
262. Nagappa M, Sinha S, Taly AB, et al. Neurosyphilis: MRI features and their phenotypic correlation in a cohort of 35 patients from a tertiary care university hospital. *Neuroradiology*. 2013;55:379–388.
263. Chesney AM, Kemp JE. Incidence of *Spirochaeta pallidum* in cerebrospinal fluid during early stages of syphilis. *JAMA*. 1924;22:1725–1728.
264. Hahn RD, Clark EG. Asymptomatic neurosyphilis: a review of the literature. *Am J Syph Gonorrhea Vener Dis*. 1946;30:305–316.
265. Moore JE. Asymptomatic neurosyphilis: a comparison of early and late asymptomatic neurosyphilis. *Arch Derm Syphilol*. 1928;18:99–108.
266. Hahn RD, Clark EG, et al. Asymptomatic neurosyphilis: prognosis. *Am J Syph Gonorrhea Vener Dis*. 1946;30:513–548.
267. Hahn RD, Clark EG. Asymptomatic neurosyphilis: a review of the literature. *Am J Syph Gonorrhea Vener Dis*. 1946;30:305–316.
268. Sparling PF. Natural history of syphilis. In: Holmes KK, Sparling PF, Mardh PA, et al, eds. *Sexually Transmitted Diseases*. New York: McGraw-Hill; 1999:473–478.
269. Marra CM. Update on neurosyphilis. *Curr Infect Dis Rep*. 2009;11:127–134.
270. Marra CM, Maxwell CL, Smith SL, et al. Cerebrospinal fluid abnormalities in patients with syphilis: association with clinical and laboratory features. *J Infect Dis*. 2004;189:369–376.
271. Flood JM, Weinstock HS, Guroy ME, et al. Neurosyphilis during the AIDS epidemic, San Francisco, 1985–1992. *J Infect Dis*. 1998;177:931–940.
272. Ghanem KG, Moore RD, Rompalo AM, et al. Lumbar puncture in HIV-infected patients with syphilis and no neurologic symptoms. *Clin Infect Dis*. 2009;48:816–821.
273. Feng W, Caplan M, Matheus MG, et al. Meningovascular syphilis with fatal vertebral occlusion. *Am J Med Sci*. 2009;338:169–171.
274. Bauerle J, Zitzmann A, Egger K, et al. The great imitator—still today! A case of meningovascular syphilis affecting the posterior circulation. *J Stroke Cerebrovasc Dis*. 2015;24:e1–e3.
275. Lair L, Naidech AM. Modern neuropsychiatric presentation of neurosyphilis. *Neurology*. 2004;63:1331–1333.
276. Kodama K, Okada S, Komatsu N, et al. Relationship between MRI findings and prognosis for patients with general paresis. *J Neuropsychiatry Clin Neurosci*. 2000;12:246–250.
277. Berbel-Garcia A, Porta-Etessam J, Martinez-Salio A, et al. Magnetic resonance image-reversible findings in a patient with general paresis. *Sex Transm Dis*. 2004;31:350–352.
278. Luo W, Ouyang Z, Xu H, et al. The clinical analysis of general paresis with 5 cases. *J Neuropsychiatry Clin Neurosci*. 2008;20:490–493.
279. Agayeva N, Karli-Oguz K, Saka E. Teaching neuroImages: a neurosyphilis case presenting with atypical neuroradiologic findings. *Neurology*. 2013;80:e119.
280. Dewhurst K. The neurosyphilitic psychoses today. A survey of 91 cases. *Br J Psychiatry*. 1969;115:31–38.
281. Osman C, Clark TW. Tabes dorsalis and Argyll Robertson pupils. *N Engl J Med*. 2016;375:e40.
282. Russouw HG, Roberts MC, Emsley RA, et al. Psychiatric manifestations and magnetic resonance imaging in HIV-negative neurosyphilis. *Biol Psychiatry*. 1997;41:467–473.
283. Mehrabian S, Raycheva MR, Petrova EP, et al. Neurosyphilis presenting with dementia, chronic chorioretinitis and adverse reactions to treatment: a case report. *Cases J*. 2009;2:8334.
284. Abdelrahman KT, Santamaria DD, Rakocevic G. Pearls and oysters: neurosyphilis presenting as mesial temporal encephalitis. *Neurology*. 2012;79:e206–e208.
285. Denays R, Collier A, Rubinstein M, et al. A 51-year-old woman with disorientation and amnesia. *Lancet*. 1999;354:1786.
286. Bash S, Hathout GM, Cohen S. Mesiotemporal T2-weighted hyperintensity: neurosyphilis mimicking herpes encephalitis. *AJNR Am J Neuroradiol*. 2001;22:314–316.
287. Szilak I, Marty F, Helft J, et al. Neurosyphilis presenting as herpes simplex encephalitis. *Clin Infect Dis*. 2001;32:1108–1109.
288. Hahn RD. Tabes dorsalis with special reference to primary optic atrophy. *Br J Vener Dis*. 1957;33:139–148.
289. Cushing H. *Intracranial Tumors: Notes Upon a Series of Two Thousand Verified Cases With Surgical-Mortality Percentages Pertaining Thereto*. Springfield: Charles C. Thomas; 1932.
290. Horowitz HW, Valsamis MP, Wicher V, et al. Brief report: cerebral syphilitic gumma confirmed by the polymerase chain reaction in a man with human immunodeficiency virus infection. *N Engl J Med*. 1994;331:1488–1491.
291. Seeley WW, Venna N. Neurosyphilis presenting with gummatous oculomotor nerve palsy. *J Neurol Neurosurg Psychiatry*. 2004;75:789.
292. Aldave AJ, King JA, Cunningham ET Jr. Ocular syphilis. *Curr Opin Ophthalmol*. 2001;12:433–441.
293. Yimtae K, Srirompotong S, Lertsukprasert K. Otsyphilis: a review of 85 cases. *Otolaryngol Head Neck Surg*. 2007;136:67–71.
294. Smith MM, Anderson JC. Neurosyphilis as a cause of facial and vestibulocochlear nerve dysfunction: MR imaging features. *AJNR Am J Neuroradiol*. 2000;21:1673–1675.
295. Becker GD. Late syphilitic hearing loss: a diagnostic and therapeutic dilemma. *Laryngoscope*. 1979;89:1273–1288.
296. Zoller M, Wilson WR, Nadol JB Jr, et al. Detection of syphilitic hearing loss. *Arch Otolaryngol*. 1978;104:63–65.
297. Pletcher SD, Cheung SW. Syphilis and otolaryngology. *Otolaryngol Clin North Am*. 2003;36:595–605, vi.
298. Moore JE. Cardiovascular syphilis: a summary of recent information with special reference to treatment with penicillin. *Am J Syph Gonorrhea Vener Dis*. 1949;33:43–55.
299. Roberts WC, Lensing FD, Kourlis H Jr, et al. Full blown cardiovascular syphilis with aneurysm of the innominate artery. *Am J Cardiol*. 2009;104:1595–1600.
300. Tong SY, Haqqani H, Street AC. A pox on the heart: five cases of cardiovascular syphilis. *Med J Aust*. 2006;184:241–243.
301. Kennedy JL, Barnard JJ, Prahlow JA. Syphilitic coronary artery ostial stenosis resulting in acute myocardial infarction and death. *Cardiology*. 2006;105:25–29.
302. Tanaka M, Okamoto M, Murayama T. A case of acute myocardial infarction due to cardiovascular syphilis with aortic regurgitation and bilateral coronary ostial stenosis. *Surg Case Rep*. 2016;2:138.
303. Roberts WC, Bose R, Ko JM, et al. Identifying cardiovascular syphilis at operation. *Am J Cardiol*. 2009;104:1588–1594.
304. Tadavarthy SM, Castaneda-Zuniga WR, Klugman J, et al. Syphilitic aneurysms of the innominate artery. *Radiology*. 1981;139:31–34.
305. De Rango P, De Socio GV, Silvestri V, et al. An unusual case of epigastric and back pain: expanding descending thoracic aneurysm resulting from tertiary syphilis diagnosed with positron emission tomography. *Circ Cardiovasc Imaging*. 2013;6:1120–1121.
306. Parnis R. Gumma of the liver. *Br J Surg*. 1975;62:236.
307. Chudomirova K, Chapkanov A, Abadjieva T, et al. Gummatous cutaneous syphilis. *Sex Transm Dis*. 2009;36:239–240.
308. Foucar FH. Syphilis of the gastro-intestinal tract: report of a case of gumma of the transverse colon with review of literature. *Am J Pathol*. 1937;13:65–80, 5.
309. King AJ, Catterall RD. Syphilis of bones. *Br J Vener Dis*. 1959;35:116–128.
310. Francis HC, Kampmeier RH. The bone lesions in acquired tertiary syphilis. *South Med J*. 1943;36:556–559.
311. Johns D. Syphilitic disorders of the spine. Report of two cases. *J Bone Joint Surg Br*. 1970;52:724–731.
312. Sena AC, White BL, Sparling PF. Novel *Treponema pallidum* serologic tests: a paradigm shift in syphilis screening for the 21st century. *Clin Infect Dis*. 2010;51:700–708.
313. Hoover KW, Radolf JD. Serodiagnosis of syphilis in the recombinant era: reversal of fortune. *J Infect Dis*. 2011;204:1295–1296.
314. Association of Public Health Laboratories. Laboratory Diagnostic Testing for *Treponema pallidum*: Expert Consultation Meeting Summary Report; January 13–15, 2009. https://www.aphl.org/programs/infectious_disease/std/Documents/ID_2009Jan_Laboratory-Guidelines-Treponema-pallidum-Meeting-Report.pdf. Accessed April 3, 2019.
315. Larsen SA, Steiner BM, Rudolph AH. Laboratory diagnosis and interpretation of tests for syphilis. *Clin Microbiol Rev*. 1995;8:1–21.
316. Wheeler HL, Agarwal S, Goh BT. Dark ground microscopy and treponemal serological tests in the diagnosis of early syphilis. *Sex Transm Infect*. 2004;80:411–414.
317. Trinh TT, Kamb ML, Luu M, et al. Syphilis testing practices in the Americas. *Trop Med Int Health*. 2017;22:1196–1203.
318. Creagan L, Bauer HM, Samuel MC, et al. An evaluation of the relative sensitivities of the Venereal Disease Research Laboratory test and the *Treponema pallidum* particle agglutination test among patients diagnosed with primary syphilis. *Sex Transm Dis*. 2007;34:1016–1018.
319. Hook EW 3rd, Roddy RE, Lukehart SA, et al. Detection of *Treponema pallidum* in lesion exudate with a pathogen-specific monoclonal antibody. *J Clin Microbiol*. 1985;22:241–244.
320. Janier M, Hegyi V, Dupin N, et al. 2014 European guideline on the management of syphilis. *J Eur Acad Dermatol Venerol*. 2014;28:1581–1593.
321. Behrhorf W, Springer E, Brauninger W, et al. PCR testing for *Treponema pallidum* in paraffin-embedded skin biopsy specimens: test design and impact on the diagnosis of syphilis. *J Clin Pathol*. 2008;61:390–395.
322. Steere AC, Strle F, Wormser GP, et al. Lyme borreliosis. *Nat Rev Dis Primers*. 2016;2:16090.
323. Ito F, Hunter EF, George RW, et al. Specific immunofluorescent staining of pathogenic treponemes with a monoclonal antibody. *J Clin Microbiol*. 1992;30:831–838.
324. Buffet M, Grange PA, Gerhardt P, et al. Diagnosing *Treponema pallidum* in secondary syphilis by PCR and immunohistochemistry. *J Invest Dermatol*. 2007;127:2345–2350.
325. Buffet M, Grange PA, Gerhardt P, et al. Diagnosing *Treponema pallidum* in secondary syphilis by PCR and immunohistochemistry. *J Invest Dermatol*. 2007;127:2345–2350.
326. Cruz AR, Ramirez LG, Zuluaga AV, et al. Immune evasion and recognition of the syphilis spirochete in blood and skin of secondary syphilis patients: two immunologically distinct compartments. *PLoS Negl Trop Dis*. 2012;6:e1717.
327. Hay PE, Clarke JR, Taylor-Robinson D, et al. Detection of treponemal DNA in the CSF of patients with syphilis and HIV infection using the polymerase chain reaction. *Genitourin Med*. 1990;66:428–432.
328. Burstein JM, Grimpel E, Lukehart SA, et al. Sensitive detection of *Treponema pallidum* by using the polymerase chain reaction. *J Clin Microbiol*. 1991;29:62–69.
329. Gayet-Ageron A, Lautenschlager S, Ninet B, et al. Sensitivity, specificity and likelihood ratios of PCR in the diagnosis of syphilis: a systematic review and meta-analysis. *Sex Transm Infect*. 2013;89:251–256.
330. Martin IE, Tsang RS, Sutherland K, et al. Molecular characterization of syphilis in patients in Canada: azithromycin resistance and detection of *Treponema pallidum* DNA in whole-blood samples versus ulcerative swabs. *J Clin Microbiol*. 2009;47:1668–1673.
331. Orle KA, Gates CA, Martin DH, et al. Simultaneous PCR detection of *Haemophilus ducreyi*, *Treponema pallidum*, and herpes simplex virus types 1 and 2 from genital ulcers. *J Clin Microbiol*. 1996;34:49–54.
332. Pope V, Fox K, Liu H, et al. Molecular subtyping of *Treponema pallidum* from North and South Carolina. *J Clin Microbiol*. 2005;43:3743–3746.
333. Heymans R, van der Helm JJ, de Vries HJ, et al. Clinical value of *Treponema pallidum* real-time PCR for diagnosis of syphilis. *J Clin Microbiol*. 2010;48:497–502.
334. Gayet-Ageron A, Sednaoui P, Lautenschlager S, et al. Use of *Treponema pallidum* PCR in testing of ulcers for diagnosis of primary syphilis. *Emerg Infect Dis*. 2015;21:127–129.
335. Gayet-Ageron A, Sednaoui P, Lautenschlager S, et al. Use of *Treponema pallidum* PCR in testing of ulcers for diagnosis of primary syphilis. *Emerg Infect Dis*. 2015;21:127–129.

336. Grange PA, Gressier L, Dion PL, et al. Evaluation of a PCR test for detection of *Treponema pallidum* in swabs and blood. *J Clin Microbiol*. 2012;50:546–552.
337. Wu BR, Tsai MS, Yang CJ, et al. Spirochetemia due to *Treponema pallidum* using polymerase-chain-reaction assays in patients with early syphilis: prevalence, associated factors and treatment response. *Clin Microbiol Infect*. 2014;20:O524–O527.
338. Vanhaecke C, Grange P, Benhaddou N, et al. Clinical and biological characteristics of 40 Patients With neurosyphilis and evaluation of *Treponema pallidum* nested polymerase chain reaction in cerebrospinal fluid samples. *Clin Infect Dis*. 2016;63:1180–1186.
339. Musher DM. Editorial commentary: polymerase chain reaction for the tpp47 Gene: a new test for neurosyphilis. *Clin Infect Dis*. 2016;63:1187–1188.
340. Larsen SA, Pope V, Johnson RE, et al. *A Manual of Tests for Syphilis*. Washington, DC: American Public Health Association; 1998.
341. Belisle JT, Brandt ME, Radolf JD, et al. Fatty acids of *Treponema pallidum* and *Borrelia burgdorferi* lipoproteins. *J Bacteriol*. 1994;176:2151–2157.
342. Larsen SA, Pope V, Johnson RE, et al. *A Manual of Tests for Syphilis*. Washington, DC: American Public Health Association; 1998.
343. Pettit DE, Larsen SA, Harbec PS, et al. Toluidine red unheated serum test, a nontreponemal test for syphilis. *J Clin Microbiol*. 1983;18:1141–1145.
344. US Preventive Services Task Force, Bibbins-Domingo K, Grossman DC, et al. Screening for syphilis infection in nonpregnant adults and adolescents: US Preventive Services Task Force recommendation statement. *JAMA*. 2016;315:2321–2327.
345. Smith G, Holman RP. The prozone phenomenon with syphilis and HIV-1 co-infection. *South Med J*. 2004;97:379–382.
346. Tong ML, Lin LR, Liu LL, et al. Analysis of 3 algorithms for syphilis serodiagnosis and implications for clinical management. *Clin Infect Dis*. 2014;58:1116–1124.
347. Discordant results from reverse sequence syphilis screening—five laboratories, United States, 2006–2010. *MMWR Morb Mortal Wkly Rep*. 2011;60:133–137.
348. Hicks CB, Benson PM, Lupton GP, et al. Seronegative secondary syphilis in a patient infected with the human immunodeficiency virus (HIV) with Kaposi sarcoma. A diagnostic dilemma. *Ann Intern Med*. 1987;107:492–495.
349. Tikjob G, Russel M, Petersen CS, et al. Seronegative secondary syphilis in a patient with AIDS: identification of *Treponema pallidum* in biopsy specimen. *J Am Acad Dermatol*. 1991;24:506–508.
350. Nandwani R, Evans DT. Are you sure it's syphilis? A review of false positive serology. *Int J STD AIDS*. 1995;6:241–248.
351. Monath TP, Frey SE. Possible autoimmune reactions following smallpox vaccination: the biologic false positive test for syphilis. *Vaccine*. 2009;27:1645–1650.
352. Zhu WF, Lei SY, Li LJ. Hepatitis C virus infection and biological false-positive syphilis test: a single-center experience. *Hepatobiliary Pancreat Dis Int*. 2011;10:399–402.
353. Henriksen R, Sogaard PE, Grennert L, et al. Autoimmune antibodies and pregnancy outcome in women with false-positive syphilis test results. A retrospective controlled investigation of women from 5170 deliveries. *Acta Obstet Gynecol Scand*. 1989;68:537–540.
354. Smikle MF, James OB, Prabhakar P. Biological false positive serological tests for syphilis in the Jamaican population. *Genitourin Med*. 1990;66:76–78.
355. Koike T. Antiphospholipid syndrome: 30 years and our contribution. *Int J Rheum Dis*. 2015;18:233–241.
356. Asherson RA, Cervera R. Antiphospholipid antibodies and infections. *Ann Rheum Dis*. 2003;62:388–393.
357. Brown ST, Zaidi A, Larsen SA, et al. Serological response to syphilis treatment. A new analysis of old data. *JAMA*. 1985;253:1296–1299.
358. Cai SN, Long J, Chen C, et al. Incidence of asymptomatic neurosyphilis in serofast Chinese syphilis patients. *Sci Rep*. 2017;7:15456.
359. Hook EW 3rd, Behets F, Van Damme K, et al. A phase III equivalence trial of azithromycin versus benzathine penicillin for treatment of early syphilis. *J Infect Dis*. 2010;201:1729–1735.
360. Sena AC, Wolff M, Behets F, et al. Rate of decline in nontreponemal antibody titers and seroreversion after treatment of early syphilis. *Sex Transm Dis*. 2017;44:6–10.
361. Discordant results from reverse sequence syphilis screening—five laboratories, United States, 2006–2010. *MMWR Morb Mortal Wkly Rep*. 2011;60:133–137.
362. Sena A, Pillay A, Cox DL, et al. *Treponema* and *Brachyspira*, human host-associated spirochetes. In: Jorgensen JH, Pfaller MA, Carroll KC, et al, eds. *Manual of Clinical Microbiology*. 11th ed. Washington, DC: ASM Press; 2015.
363. Sena AC, White BL, Sparling PF. Novel *Treponema pallidum* serologic tests: a paradigm shift in syphilis screening for the 21st century. *Clin Infect Dis*. 2010;51:700–708.
364. Park IU, Chow JM, Bolan G, et al. Screening for syphilis with the treponemal immunoassay: analysis of discordant serology results and implications for clinical management. *J Infect Dis*. 2011;204:1297–1304.
365. Morshed MG, Singh AE. Recent trends in the serologic diagnosis of syphilis. *Clin Vaccine Immunol*. 2015;22:137–147.
366. Rourke AR, Nolte FS, Litwin CM. Performance characteristics of the reverse syphilis screening algorithm in a population with a moderately high prevalence of syphilis. *Am J Clin Pathol*. 2016;146:572–577.
367. Mmeje O, Chow JM, Davidson L, et al. Discordant syphilis immunoassays in pregnancy: perinatal outcomes and implications for clinical management. *Clin Infect Dis*. 2015;61:1049–1053.
368. Hernandez-Aguado I, Bolumar F, Moreno R, et al. False-positive tests for syphilis associated with human immunodeficiency virus and hepatitis B virus infection among intravenous drug abusers. Valencian Study Group on HIV Epidemiology. *Eur J Clin Microbiol Infect Dis*. 1998;17:784–787.
369. Augenbraun MH, DeHovitz JA, Feldman J, et al. Biological false-positive syphilis test results for women infected with human immunodeficiency virus. *Clin Infect Dis*. 1994;19:1040–1044.
370. Geisler WM. The prozone phenomenon in syphilis testing. *South Med J*. 2004;97:327–328.
371. Ghanem KG, Erbeling EJ, Wiener ZS, et al. Serological response to syphilis treatment in HIV-positive and HIV-negative patients attending sexually transmitted diseases clinics. *Sex Transm Infect*. 2007;83:97–101.
372. Kingston AA, Vujecich J, Shapiro M, et al. Seronegative secondary syphilis in 2 patients coinfected with human immunodeficiency virus. *Arch Dermatol*. 2005;141:431–433.
373. Peeling RW, Mabey D. Point-of-care tests for diagnosing infections in the developing world. *Clin Microbiol Infect*. 2010;16:1062–1069.
374. Tucker JD, Bien CH, Peeling RW. Point-of-care testing for sexually transmitted infections: recent advances and implications for disease control. *Curr Opin Infect Dis*. 2013;26:73–79.
375. Peeling RW, Mabey D, Herring A, et al. Why do we need quality-assured diagnostic tests for sexually transmitted infections? *Nat Rev Microbiol*. 2006;4:57–S19.
376. Jafari Y, Peeling RW, Shivkumar S, et al. Are *Treponema pallidum* specific rapid and point-of-care tests for syphilis accurate enough for screening in resource limited settings? Evidence from a meta-analysis. *PLoS ONE*. 2013;8:e54695.
377. Castro AR, Esfandiari J, Kumar S, et al. Novel point-of-care test for simultaneous detection of nontreponemal and treponemal antibodies in patients with syphilis. *J Clin Microbiol*. 2010;48:4615–4619.
378. Yin YP, Chen XS, Wei WH, et al. A dual point-of-care test shows good performance in simultaneously detecting nontreponemal and treponemal antibodies in patients with syphilis: a multisite evaluation study in China. *Clin Infect Dis*. 2013;56:659–665.
379. Gliddon HD, Peeling RW, Kamb ML, et al. A systematic review and meta-analysis of studies evaluating the performance and operational characteristics of dual point-of-care tests for HIV and syphilis. *Sex Transm Infect*. 2017;93:S3–S15.
380. Andrade R, Rodriguez-Barradas MC, Yasukawa K, et al. Single dose versus 3 doses of intramuscular benzathine penicillin for early syphilis in HIV: a randomized clinical trial. *Clin Infect Dis*. 2017;64:759–764.
381. Polisel R, Vidal JE, Penalva De Oliveira AC, et al. Neurosyphilis in HIV-infected patients: clinical manifestations, serum Venereal Disease Research Laboratory titers, and associated factors to symptomatic neurosyphilis. *Sex Transm Dis*. 2008;35:425–429.
382. Libois A, De Wit S, Poll B, et al. HIV and syphilis: when to perform a lumbar puncture. *Sex Transm Dis*. 2007;34:141–144.
383. Marra CM, Tantal LC, Maxwell CL, et al. The rapid plasma reagin test cannot replace the Venereal Disease Research Laboratory test for neurosyphilis diagnosis. *Sex Transm Dis*. 2012;39:453–457.
384. Castro R, Prieto ES, da Luz Martins Pereira F. Nontreponemal tests in the diagnosis of neurosyphilis: an evaluation of the Venereal Disease Research Laboratory (VDRL) and the rapid plasma reagin (RPR) tests. *J Clin Lab Anal*. 2008;22:257–261.
385. Jaffe HW, Larsen SA, Peters M, et al. Tests for treponemal antibody in CSF. *Arch Intern Med*. 1978;138:252–255.
386. Muller F, Moskopidis M. Estimation of the local production of antibodies to *Treponema pallidum* in the central nervous system of patients with neurosyphilis. *Br J Vener Dis*. 1983;59:80–84.
387. Prange HW, Moskopidis M, Schipper HI, et al. Relationship between neurological features and intrathecal synthesis of IgG antibodies to *Treponema pallidum* in untreated and treated human neurosyphilis. *J Neurol*. 1983;230:241–252.
388. Clement ME, Okeke NL, Hicks CB. Treatment of syphilis: a systematic review. *JAMA*. 2014;312:1905–1917.
389. Norris SJ, Edmondson DG. In vitro culture system to determine MICs and MBCs of antimicrobial agents against *Treponema pallidum* subsp. *pallidum* (Nichols strain). *Antimicrob Agents Chemother*. 1988;32:68–74.
390. Lukehart SA, Baker-Zander SA, Holmes KK. Efficacy of aztreonam in treatment of experimental syphilis in rabbits. *Antimicrob Agents Chemother*. 1984;25:390–391.
391. Magnuson HJ, Rosenau BJ. The rate of development and degree of acquired immunity in experimental syphilis. *Am J Syph Gonorrhea Vener Dis*. 1948;32:418–436.
392. Lukehart SA, Fohn MJ, Baker-Zander SA. Efficacy of azithromycin for therapy of active syphilis in the rabbit model. *J Antimicrob Chemother*. 1990;25(supplA):91–99.
393. Molini BJ, Tantal LC, Sahi SK, et al. Macrolide resistance in *Treponema pallidum* correlates With 23S rDNA mutations in recently isolated clinical strains. *Sex Transm Dis*. 2016;43:579–583.
394. Idsoe O, Guthe T, Willcox RR. Penicillin in the treatment of syphilis. The experience of three decades. *Bull World Health Organ*. 1972;47(suppl):1–68.
395. Rein MF. Biopharmacology of syphilotherapy. *J Am Vener Dis Assoc*. 1976;3:109–127.
396. Eagle H, Fleishman R, Musselman AD. The effective concentrations of penicillin in vitro and in vivo for streptococci, pneumococci, and *Treponema pallidum*. *J Bacteriol*. 1950;59:625–643.
397. Radolf JD, Moomaw C, Slaughter CA, et al. Penicillin-binding proteins and peptidoglycan of *Treponema pallidum* subsp. *pallidum*. *Infect Immun*. 1989;57:1248–1254.
398. Smith CA, Kamp M, Olansky S, et al. Benzathine penicillin G in the treatment of syphilis. *Bull World Health Organ*. 1956;15:1087–1096.
399. Eagle H, Magnuson HJ, Fleishman R. Relation of the size of the inoculum and the age of the infection to the curative dose of penicillin in experimental syphilis, with particular reference to the feasibility of its prophylactic use. *J Exp Med*. 1947;85:423–440.
400. Zenker PN, Rolfs RT. Treatment of syphilis, 1989. *Rev Infect Dis*. 1990;12(suppl 6):S590–S609.
401. Schroeter AL, Lucas JB, Price EV, et al. Treatment for early syphilis and reactivity of serologic tests. *JAMA*. 1972;221:471–476.
402. Rolfs RT. Treatment of syphilis, 1993. *Clin Infect Dis*. 1995;20(suppl 1):S23–S38.
403. Jefferis FJ, Willcox RR. Treatment of early syphilis with penicillin alone. *Br J Vener Dis*. 1963;39:143–148.
404. Fernando WL. Cerebrospinal fluid findings after treatment of early syphilis with penicillin. *Br J Vener Dis*. 1965;41:168–169.
405. Fernando WL. Cerebrospinal fluid findings after treatment of early syphilis with penicillin. A further series of 80 cases. *Br J Vener Dis*. 1968;44:134–135.
406. Mohr JA, Griffiths W, Jackson R, et al. Neurosyphilis and penicillin levels in cerebrospinal fluid. *JAMA*. 1976;236:2208–2209.
407. Tramont EC. Persistence of *Treponema pallidum* following penicillin G therapy. Report of two cases. *JAMA*. 1976;236:2206–2207.
408. Ghanem KG. Evaluation and management of syphilis in the HIV-infected patient. *Curr Infect Dis Rep*. 2010;12:140–146.
409. Blank LJ, Rompalo AM, Erbeling EJ, et al. Treatment of syphilis in HIV-infected subjects: a systematic review of the literature. *Sex Transm Infect*. 2011;87:9–16.
410. Riedner G, Rusizoka M, Todd J, et al. Single-dose azithromycin versus penicillin G benzathine for the treatment of early syphilis. *N Engl J Med*. 2005;353:1236–1244.
411. Ganesan A, Mesner O, Okulicz JF, et al. A single dose of benzathine penicillin G is as effective as multiple doses of benzathine penicillin G for the treatment of HIV-infected persons with early syphilis. *Clin Infect Dis*. 2015;60:653–660.
412. Clement ME, Okeke NL, Hicks CB. Treatment of syphilis: a systematic review. *JAMA*. 2014;312:1905–1917.
413. Jaffe HW. Treatment of latent syphilis. *J Am Vener Dis Assoc*. 1976;3:143–145.
414. Fiumara NJ. Serologic responses to treatment of 128 patients with late latent syphilis. *Sex Transm Dis*. 1979;6:243–246.
415. Andrade P, Mariano A, Figueiredo A. Solitary frontal ulcer: a syphilitic gumma. *Dermatol Online J*. 2010;16:5.
416. Milligan R, Dent B, Hoare T, et al. Gummatous rectal syphilis imitating rectal carcinoma. *Colorectal Dis*. 2010;12:1269–1270.

417. Ghanem KG, Erbelding EJ, Cheng WW, et al. Doxycycline compared with benzathine penicillin for the treatment of early syphilis. *Clin Infect Dis*. 2006;42:e45–e49.
418. Wong T, Singh AE, De P. Primary syphilis: serological treatment response to doxycycline/tetracycline versus benzathine penicillin. *Am J Med*. 2008;121:903–908.
419. Spornraft-Ragaller P, Abraham S, Lueck C, et al. Response of HIV-infected patients with syphilis to therapy with penicillin or intravenous ceftriaxone. *Eur J Med Res*. 2011;16:47–51.
420. Long CM, Klausner JD, Leon S, et al. Syphilis treatment and HIV infection in a population-based study of persons at high risk for sexually transmitted disease/HIV infection in Lima, Peru. *Sex Transm Dis*. 2006;33:151–155.
421. Shann S, Wilson J. Treatment of neurosyphilis with ceftriaxone. *Sex Transm Infect*. 2003;79:415–416.
422. Marra CM, Boutin P, McArthur JC, et al. A pilot study evaluating ceftriaxone and penicillin G as treatment agents for neurosyphilis in human immunodeficiency virus-infected individuals. *Clin Infect Dis*. 2000;30:540–544.
423. Hook EW 3rd, Roddy RE, Handsfield HH. Ceftriaxone therapy for incubating and early syphilis. *J Infect Dis*. 1988;158:881–884.
424. Romano A, Gueant-Rodriguez RM, Viola M, et al. Cross-reactivity and tolerability of cephalosporins in patients with immediate hypersensitivity to penicillins. *Ann Intern Med*. 2004;141:16–22.
425. Pichichero ME. Cephalosporins can be prescribed safely for penicillin-allergic patients. *J Fam Pract*. 2006;55:106–112.
426. Matejkova P, Flasarova M, Zakoucka H, et al. Macrolide treatment failure in a case of secondary syphilis: a novel A2059G mutation in the 23S rRNA gene of *Treponema pallidum* subsp. *pallidum*. *J Med Microbiol*. 2009;58:832–836.
427. Lukehart SA, Godornes C, Molini BJ, et al. Macrolide resistance in *Treponema pallidum* in the United States and Ireland. *N Engl J Med*. 2004;351:154–158.
428. Grimes M, Sahi SK, Godornes BC, et al. Two mutations associated with macrolide resistance in *Treponema pallidum*: increasing prevalence and correlation with molecular strain type in Seattle, Washington. *Sex Transm Dis*. 2012;39:954–958.
429. Mitchell SJ, Engelman J, Kent CK, et al. Azithromycin-resistant syphilis infection: San Francisco, California, 2000–2004. *Clin Infect Dis*. 2006;42:337–345.
430. Van Damme K, Behets F, Ravelomanana N, et al. Evaluation of azithromycin resistance in *Treponema pallidum* specimens from Madagascar. *Sex Transm Dis*. 2009;36:775–776.
431. Chen XS, Yin YP, Wei WH, et al. High prevalence of azithromycin resistance to *Treponema pallidum* in geographically different areas in China. *Clin Microbiol Infect*. 2013;19:975–979.
432. Wendel GD Jr, Sheffield JS, Hollier LM, et al. Treatment of syphilis in pregnancy and prevention of congenital syphilis. *Clin Infect Dis*. 2002;35:S200–S209.
433. Genc M, Ledger WJ. Syphilis in pregnancy. *Sex Transm Infect*. 2000;76:73–79.
434. Wijesooriya NS, Rochat RW, Kamb ML, et al. Global burden of maternal and congenital syphilis in 2008 and 2012: a health systems modelling study. *Lancet Glob Health*. 2016;4:e252–e533.
435. Patel SJ, Klinger EJ, O'Toole D, et al. Missed opportunities for preventing congenital syphilis infection in New York City. *Obstet Gynecol*. 2012;120:882–888.
436. Alexander JM, Sheffield JS, Sanchez PJ, et al. Efficacy of treatment for syphilis in pregnancy. *Obstet Gynecol*. 1999;93:5–8.
437. Nathan L, Bawdon RE, Sidawi JE, et al. Penicillin levels following the administration of benzathine penicillin G in pregnancy. *Obstet Gynecol*. 1993;82:338–342.
438. Putkonen T, Salo OP, Mustakallio KK. Febrile Herxheimer reaction in different phases of primary and secondary syphilis. *Br J Vener Dis*. 1966;42:181–184.
439. Brown ST. Adverse reactions in syphilis therapy. *J Am Vener Dis Assoc*. 1976;3:172–176.
440. Fekade D, Knox K, Hussein K, et al. Prevention of Jarisch-Herxheimer reactions by treatment with antibodies against tumor necrosis factor alpha. *N Engl J Med*. 1996;335:311–315.
441. Negussie Y, Remick DG, DeForge LE, et al. Detection of plasma tumor necrosis factor, interleukins 6, and 8 during the Jarisch-Herxheimer reaction of relapsing fever. *J Exp Med*. 1992;175:1207–1212.
442. Fathilah J, Choo MM. The Jarisch-Herxheimer reaction in ocular syphilis. *Med J Malaysia*. 2003;58:437–439.
443. Kobayashi J, Nakagawa Y, Tobisawa S, et al. Deterioration of MRI findings related to Jarisch-Herxheimer reaction in a patient with neurosyphilis. *J Neurol*. 2011;258:699–701.
444. Fiumara NJ. Treatment of seropositive primary syphilis: an evaluation of 196 patients. *Sex Transm Dis*. 1977;4:92–95.
445. Fiumara NJ. Treatment of secondary syphilis: an evaluation of 204 patients. *Sex Transm Dis*. 1977;4:96–99.
446. Guinan ME. Treatment of primary and secondary syphilis: defining failure at three- and six-month follow-up. *JAMA*. 1987;257:359–360.
447. Romanowski B, Sutherland R, Fick GH, et al. Serologic response to treatment of infectious syphilis. *Ann Intern Med*. 1991;114:1005–1009.
448. Sena AC, Wolff M, Martin DH, et al. Predictors of serological cure and serofast state after treatment in HIV-negative persons with early syphilis. *Clin Infect Dis*. 2011;53:1092–1099.
449. Sena AC, Wolff M, Behets F, et al. Response to therapy following retreatment of serofast early syphilis patients with benzathine penicillin. *Clin Infect Dis*. 2013;56:420–422.
450. Holman KM, Hook EW 3rd. Clinical management of early syphilis. *Expert Rev Anti Infect Ther*. 2013;11:839–843.
451. Lukehart SA. Serologic testing after therapy for syphilis: is there a test for cure? *Ann Intern Med*. 1991;114:1057–1058.
452. Kamb ML, Newman LM, Riley PL, et al. A road map for the global elimination of congenital syphilis. *Obstet Gynecol Int*. 2010;2010:312798.
453. Schmid GP, Stoner BP, Hawkes S, et al. The need and plan for global elimination of congenital syphilis. *Sex Transm Dis*. 2007;34:S5–S10.
454. Kruger C, Malleyeck I. Congenital syphilis: still a serious, under-diagnosed threat for children in resource-poor countries. *World J Pediatr*. 2010;6:125–131.
455. Walker GJ, Walker DG. Congenital syphilis: a continuing but neglected problem. *Semin Fetal Neonatal Med*. 2007;12:198–206.
456. Hsu KK. Congenital syphilis: time for a national prevention program. *Sex Transm Dis*. 2017;44:503–504.
457. Cruz AR, Castrillon MA, Minotta AY, et al. Gestational and congenital syphilis epidemic in the Colombian Pacific Coast. *Sex Transm Dis*. 2013;40:813–818.
458. Lomotey CJ, Lewis J, Gebrian B, et al. Maternal and congenital syphilis in rural Haiti. *Rev Panam Salud Publica*. 2009;26:197–202.
459. Dorfman DH, Glaser JH. Congenital syphilis presenting in infants after the newborn period. *N Engl J Med*. 1990;323:1299–1302.
460. Fiumara NJ, Fleming WL, Downing JG, et al. The incidence of prenatal syphilis at the Boston City Hospital. *N Engl J Med*. 1952;247:48–52.
461. Sanchez PJ, Wendel GD Jr, Grimpel E, et al. Evaluation of molecular methodologies and rabbit infectivity testing for the diagnosis of congenital syphilis and neonatal central nervous system invasion by *Treponema pallidum*. *J Infect Dis*. 1993;167:148–157.
462. Hollier LM, Harstad TW, Sanchez PJ, et al. Fetal syphilis: clinical and laboratory characteristics. *Obstet Gynecol*. 2001;97:947–953.
463. Casal C, Araujo Eda C, Corvelo TC. Risk factors and pregnancy outcomes in women with syphilis diagnosed using a molecular approach. *Sex Transm Infect*. 2013;89:257–261.
464. Harter C, Benirschke K. Fetal syphilis in the first trimester. *Am J Obstet Gynecol*. 1976;124:705–711.
465. Lomotey CJ, Lewis J, Gebrian B, et al. Maternal and congenital syphilis in rural Haiti. *Rev Panam Salud Publica*. 2009;26:197–202.
466. Gust DA, Levine WC, St Louis ME, et al. Mortality associated with congenital syphilis in the United States, 1992–1998. *Pediatrics*. 2002;109:E79.
467. Judge DM, Tafari N, Naeye RL, et al. Congenital syphilis and perinatal mortality. *Pediatr Pathol*. 1986;5:411–420.
468. Qureshi F, Jacques SM, Reyes MP. Placental histopathology in syphilis. *Hum Pathol*. 1993;24:779–784.
469. Malan AF, Woods DL, Van der Elst CW, et al. Relative placental weight in congenital syphilis. *Placenta*. 1990;11:3–6.
470. Russell P, Altschuler G. Placental abnormalities of congenital syphilis. A neglected aid to diagnosis. *Am J Dis Child*. 1974;128:160–163.
471. Sheffield JS, Sanchez PJ, Wendel GD Jr, et al. Placental histopathology of congenital syphilis. *Obstet Gynecol*. 2002;100:126–133.
472. Walter P, Blot P, Ivanoff B. The placental lesions in congenital syphilis. A study of six cases. *Virchows Arch A Pathol Anat Histol*. 1982;397:313–326.
473. Genest DR, Choi-Hong SR, Tate JE, et al. Diagnosis of congenital syphilis from placental examination: comparison of histopathology, Steiner stain, and polymerase chain reaction for *Treponema pallidum* DNA. *Hum Pathol*. 1996;27:366–372.
474. Budell JW. Treatment of congenital syphilis. *J Am Vener Dis Assoc*. 1976;3:168–171.
475. Reyes MP, Hunt N, Ostrea EM Jr, et al. Maternal/congenital syphilis in a large tertiary-care urban hospital. *Clin Infect Dis*. 1993;17:1041–1046.
476. Stoll BJ, Lee FK, Larsen S, et al. Clinical and serologic evaluation of neonates for congenital syphilis: a continuing diagnostic dilemma. *J Infect Dis*. 1993;167:1093–1099.
477. Fojaco RM, Hensley GT, Moskowitz L. Congenital syphilis and necrotizing funisitis. *JAMA*. 1989;261:1788–1790.
478. Jacques SM, Qureshi F. Necrotizing funisitis: a study of 45 cases. *Hum Pathol*. 1992;23:1278–1283.
479. Khetarpal S, Kempf E, Mostow E. Congenital syphilis: early- and late-stage findings of rhagades and dental anomalies. *Pediatr Dermatol*. 2011;28:401–403.
480. Tiliakos N, Shammaa JM, Nasrallah SM. Syphilitic hepatitis. *Am J Gastroenterol*. 1980;73:60–61.
481. Long WA, Ulshen MH, Lawson EE. Clinical manifestations of congenital syphilitic hepatitis: implications for pathogenesis. *J Pediatr Gastroenterol Nutr*. 1984;3:551–555.
482. Juhlin L. Early congenital syphilis and thrombocytopenia. *Acta Derm Venereol*. 1968;48:166–169.
483. O'Regan S, Fong JS, de Chadarevian JP, et al. Treponemal antigens in congenital and acquired syphilitic nephritis: demonstration by immunofluorescence studies. *Ann Intern Med*. 1976;85:325–327.
484. Obladen M. Curse on two generations: a history of congenital syphilis. *Neonatology*. 2013;103:274–280.
485. Moyer VA, Schneider V, Yetman R, et al. Contribution of long-bone radiographs to the management of congenital syphilis in the newborn infant. *Arch Pediatr Adolesc Med*. 1998;152:353–357.
486. Michelow IC, Wendel GD Jr, Norgard MV, et al. Central nervous system infection in congenital syphilis. *N Engl J Med*. 2002;346:1792–1798.
487. Daaboul JJ, Kartchner W, Jones KL. Neonatal hypoglycemia caused by hypopituitarism in infants with congenital syphilis. *J Pediatr*. 1993;123:983–985.
488. Laird SM. Late congenital syphilis: an analysis of 115 cases. *Br J Vener Dis*. 1950;26:143–145.
489. Hariprasad SM, Moon SJ, Allen RC, et al. Keratopathy from congenital syphilis. *Cornea*. 2002;21:608–609.
490. Hillson S, Grigson C, Bond S. Dental defects of congenital syphilis. *Am J Phys Anthropol*. 1998;107:25–40.
491. Sparling PF, Musher MN, Healy BP. Clinical manifestations of syphilis. In: Holmes KK, Sparling PF, Stamm WE, et al, eds. *Sexually Transmitted Diseases*. 4th ed. New York: McGraw Hill Medical; 2008:662–684.
492. Ingall D, Sanchez PJ, Musher D. Syphilis. In: Remington JS, Klein JD, eds. *Infectious Diseases of the Fetus and Newborn Infant*. Philadelphia: Saunders; 1995.
493. Woods CR. Syphilis in children: congenital and acquired. *Semin Pediatr Infect Dis*. 2005;16:245–257.
494. Glaser JH. Centers for Disease Control and Prevention guidelines for congenital syphilis. *J Pediatr*. 1996;129:488–490.
495. Sanchez PJ, McCracken GH Jr, Wendel GD, et al. Molecular analysis of the fetal IgM response to *Treponema pallidum* antigens: implications for improved serodiagnosis of congenital syphilis. *J Infect Dis*. 1989;159:508–517.
496. Wendel GD Jr, Sanchez PJ, Peters MT, et al. Identification of *Treponema pallidum* in amniotic fluid and fetal blood from pregnancies complicated by congenital syphilis. *Obstet Gynecol*. 1991;78:890–895.
497. Rawstron SA, Vetrano J, Tannis G, et al. Congenital syphilis: detection of *Treponema pallidum* in stillborns. *Clin Infect Dis*. 1997;24:24–27.
498. Wu CC, Tsai CN, Wong WR, et al. Early congenital syphilis and erythema multiforme-like bullous targetoid lesions in a 1-day-old newborn: detection of *Treponema pallidum* genomic DNA from the targetoid plaque using nested polymerase chain reaction. *J Am Acad Dermatol*. 2006;55:S11–S15.
499. Woznicova V, Smajs D, Wechsler D, et al. Detection of *Treponema pallidum* subsp. *pallidum* from skin lesions, serum, and cerebrospinal fluid in an infant with congenital syphilis after clindamycin treatment of the mother during pregnancy. *J Clin Microbiol*. 2007;45:659–661.
500. Grimpel E, Sanchez PJ, Wendel GD, et al. Use of polymerase chain reaction and rabbit infectivity testing to detect *Treponema pallidum* in amniotic fluid, fetal and neonatal sera, and cerebrospinal fluid. *J Clin Microbiol*. 1991;29:1711–1718.
501. Workowski KA. Sexually transmitted infections and HIV: diagnosis and treatment. *Top Antivir Med*. 2012;20:11–16.
502. Azimi PH, Janner D, Berne P, et al. Concentrations of procaine and aqueous penicillin in the cerebrospinal fluid of infants treated for congenital syphilis. *J Pediatr*. 1994;124:649–653.

503. Cameron CE, Lukehart SA. Current status of syphilis vaccine development: need, challenges, prospects. *Vaccine*. 2014;32:1602–1609.
504. Lithgow KV, Cameron CE. Vaccine development for syphilis. *Expert Rev Vaccines*. 2017;16:37–44.
505. Chesney AM. Immunity in syphilis. *Medicine (Baltimore)*. 1926;5:463–547.
506. Magnuson HJ, Rosenau BJ. The rate of development and degree of acquired immunity in experimental syphilis. *Am J Syph Gonorrhea Vener Dis*. 1948;32:418–436.
507. Miller JN. Immunity in experimental syphilis. VI. Successful vaccination of rabbits with *Treponema pallidum*, Nichols strain, attenuated by γ -irradiation. *J Immunol*. 1973;110:1206–1215.
508. Katz KA, Lee MA, Gray T, et al. Repeat syphilis among men who have sex with men—San Diego County, 2004–2009. *Sex Transm Dis*. 2011;38:349–352.
509. Fiumara NJ. Reinfection primary, secondary, and latent syphilis: the serologic response after treatment. *Sex Transm Dis*. 1980;7:111–115.
510. Gottlieb SL, Deal CD, Giersing B, et al. The global roadmap for advancing development of vaccines against sexually transmitted infections: update and next steps. *Vaccine*. 2016;34:2939–2947.
511. Lithgow KV, Hof R, Wetherell C, et al. A defined syphilis vaccine candidate inhibits dissemination of *Treponema pallidum* subspecies *pallidum*. *Nat Commun*. 2017;8:14273.

SHORT VIEW SUMMARY

Definition

- The endemic treponematoses include yaws, endemic syphilis, and pinta. All are cutaneous diseases caused by treponemes closely related to *Treponema pallidum*, the causative agent of syphilis.

Microbiology

- The treponemal species that cause the endemic treponematoses cannot be easily cultured for diagnosis. Microbiologic diagnosis is most often based on nontreponemal and treponemal serologic reactivity in tests designed for syphilis diagnosis.

Epidemiology

- The endemic treponematoses are generally spread only from human to human by direct contact. After global eradication campaigns in the 1950s and 1960s, yaws rates have again increased worldwide. Endemic syphilis and

pinta remain uncommon, but the prevalence is unknown because of the lack of surveillance data.

Clinical Manifestations

- Similar to syphilis, initial lesions of endemic treponematoses resolve spontaneously; recurrent secondary lesions are typically cutaneous or involve bone or cartilage.
- A latent (asymptomatic) stage may occur in individuals infected with yaws or endemic syphilis, and a minority of untreated patients progress to bone, soft tissue, and cartilaginous destruction.

Diagnosis

- Diagnosis of endemic treponematoses is based on clinical recognition of compatible skin lesions and confirmed by serologic testing.
- The development of a point-of-care immunochromatographic lateral flow assay

that simultaneously tests for treponemal and nontreponemal antibodies facilitates rapid diagnosis and treatment.

Therapy

- Benzathine penicillin G was the mainstay of therapy of the endemic treponematoses. More recently, oral azithromycin has been proven to be an easy-to-administer alternative to penicillin injections for therapy of yaws.

Prevention

- Yaws appears to be amenable to mass treatment as a control and prevention measure.
- In 2012 the World Health Organization announced plans to embark on mass treatment initiatives designed to eradicate yaws by 2020.

The endemic treponematoses comprise yaws, endemic syphilis, and pinta and are caused by *Treponema pallidum* subsp. *pertenue*, *Treponema pallidum* subsp. *endemicum*, and *Treponema carateum*. Nearly eliminated in the 1960s by an eradication campaign led by the World Health Organization (WHO), these relatively uncommon diseases are now often seen among children living in low-income and middle-income countries. The bacteria that cause the endemic treponematoses are morphologically and serologically indistinguishable from *T. pallidum* subsp. *pallidum*, the causative agent of sexually transmitted syphilis, and important parallels are found between the natural history of these diseases and syphilis. Similarly, the tools for management are adapted almost entirely from tools historically used as part of syphilis control efforts. Nonetheless, the endemic treponematoses differ from syphilis in terms of epidemiology, clinical manifestations, and at the level of the bacterial genome.^{1–3,4,5}

Disease rates have again increased more recently.^{1,6} Based on the successful eradication of yaws in India and the potential for recently described oral azithromycin therapy to facilitate control and prevention, the WHO announced a new effort to eradicate yaws.^{7,8}

MICROBIOLOGY

Within the genus *Treponema*, four bacteria are presently recognized as human pathogens: *T. pallidum* subsp. *pallidum*, *T. pallidum* subsp. *pertenue*, *T. pallidum* subsp. *endemicum*, and *T. carateum*. Much remains to be learned about these organisms. They cannot be cultured in vitro, and current understanding of their biology is based on the careful study of relatively few clinical isolates, most often propagated in laboratory animals (e.g., rabbits, hamsters, and guinea pigs).^{4,5} Although sequencing of the *T. pallidum* genome showed more than 99% DNA homology,³ several genetic loci have been identified that may permit differentiation of *T. pallidum* subsp. *pallidum* from the other subspecies (*T. pertenue* and *T. endemicum*).^{4,5} *T. carateum* isolates are not available for study at the present time, and any molecular similarities and differences have

yet to be compared with the *T. pallidum* subspecies.⁹ In the future, application of evolving molecular diagnostic methods to the study of human treponemes promises to provide new insights into the biology of these organisms.

Morphology

T. pallidum and the treponemes causing endemic treponematoses are long, thin (8–13 × 0.15 μm), motile bacteria that cannot be seen with Gram stain and are best seen in specimens of a lesion exudate with a Warthin-Starry silver stain, darkfield microscopic examination, or fluorescent antibody techniques. Their regular, spiral morphology and characteristic corkscrew motility are helpful for recognition in clinical specimens.

Antibiotic Sensitivity

Based on clinical and serologic response to therapy as well as studies performed in experimental animals, these treponemes are sensitive to penicillins and tetracyclines. Because the organisms cannot be propagated in vitro, data on minimal inhibitory and bactericidal concentrations are unavailable. Sulfa drugs and fluoroquinolone antimicrobials have not been found to be active against *T. pallidum* subsp. *pallidum*.^{10,11} Clinical and laboratory resistance (mutations of 23S ribosomal RNA gene) to erythromycin and other macrolide antibiotics has been shown for multiple isolates of *T. pallidum* subsp. *pallidum* from North America and Western Europe,¹² but similar resistance among the endemic treponematoses was not documented until 2018.¹³ With the adoption by the WHO of azithromycin as the preferred treatment for yaws,¹⁴ close monitoring for macrolide resistance will be important.

EPIDEMIOLOGY

The endemic treponematoses are diseases of low-income and middle-income countries, with varied geographic distribution. However, the true scope of disease is unknown because endemic treponematoses are

not notifiable diseases in most countries. Yaws appears to have worldwide distribution (Central and South America, Asia, Africa, and Pacific Islands) and is found in warm, humid climates with heavy rainfall including tropical regions and rainforests. Of 13 endemic countries that reported to the WHO between 1990 and 2014, prevalence of active yaws disease ranged from 0.31% to 14.54%, and prevalence of latent yaws ranged from 2.45% to 31.05% with a higher burden of infection in the Pacific Islands.¹⁵ Endemic syphilis is less common than yaws and occurs primarily in dry, arid areas among nomadic and seminomadic rural populations. The disease has been most common in North Africa, Southwest Asia, and the eastern Mediterranean region. In contrast to the global distribution of yaws and endemic syphilis, pinta is limited to the Western Hemisphere. The disease has been described in Mexico, Caribbean islands (including Cuba), and Central and South American countries.^{6,9} The prevalence of pinta is unknown at the present time because of the lack of surveillance data. However, substantial numbers of cases including a case series of more than 200 patients have been described in rural Brazil.¹⁶ Within these regions, each of the endemic treponematoses disproportionately impacts impoverished people living in remote areas. Limited access to hygienic facilities has been associated with increased infection rates.

Transmission

Similar to syphilis, the endemic treponematoses are acquired and spread through direct contact with infectious exudates from patients with active infection or who have relapsed after latent infection (infected but asymptomatic). In the case of endemic syphilis, transmission is possible via fomites (see “Endemic Syphilis [Bejel]”). However, in contrast to syphilis, little evidence exists that transmission of endemic treponematoses occurs via blood, blood products, or transplacentally from untreated pregnant women to unborn children. Protective immunity induced by infection has been shown only in rabbit experimental models conducted in the early 20th century; however, a later study in hamsters concluded that immunity was not absolute.¹

CLINICAL MANIFESTATIONS

The endemic treponematoses can be characterized by several common clinical characteristics as well as characteristics that distinguish them from one another.¹⁷ Although there are several important differences, there are numerous parallels between the well-described course of untreated syphilis (*T. pallidum* subsp. *pallidum* infection) and the natural history of the endemic treponematoses. Similar to syphilis, each is a chronic bacterial infection that progresses through a series of well-described clinical stages, predictably progressing from an early, localized stage to a later, more widespread stage. Early in the course of infection, even before primary lesions develop, the treponemes spread hematogenously throughout the body; without treatment, they subsequently give rise to secondary and late manifestations of infection. Once primary lesions develop, regional lymphadenopathy follows. As in the case of syphilis, primary lesions of yaws and pinta resolve spontaneously without treatment; then in the secondary stage, skin lesions may recur up to 5 years, with the frequency of these recurrences declining with time.¹⁴ Recurrent secondary manifestations of early endemic treponematoses are primarily dermatologic; however, a substantial proportion of persons also develop osteitis or other bony or cartilaginous lesions. Primary and secondary lesions may resolve, and a patient's infection may become latent, where the patient harbors infection without exhibiting signs and symptoms.¹³

Late Complications and Sequelae

In a minority of untreated patients with yaws and endemic syphilis, late complications of infection occur. The lesions of late infections tend to be more destructive than the early lesions and most commonly manifest either as ulcerative or hyperkeratotic cutaneous lesions or as bone and joint involvement. A distinguishing element of the endemic treponematoses is that, in contrast to syphilis, little evidence exists of cardiovascular or neurologic sequelae in late or untreated infection.

Differential Diagnosis

The differential diagnosis varies between the endemic treponematoses and their stages. As the clinical presentation of endemic treponematoses

can be mistaken for sexually transmitted syphilis, it is difficult to distinguish between the two diseases especially in countries where both are endemic. Early yaws may present similar to impetigo, scabies, molluscum contagiosum, lichen planus, and cutaneous leishmaniasis. The differential diagnosis for endemic syphilis includes orofacial herpes simplex and noninfectious aphthous ulcers. Early lesions of pinta may be confused with eczema, psoriasis, and leprosy.

Yaws

Yaws, one of the 20 neglected tropical diseases according to the WHO, is a chronic infection caused by *T. pallidum* subsp. *pertenue* and is the most common of the endemic treponematoses. After years of relatively low levels of prevalence (see “Prevention”), there has been a resurgence of disease.^{1,6,7} Yaws is most common in children 2 to 15 years of age. Human-to-human transmission of yaws is thought to be facilitated by skin breaks (e.g., lacerations, insect bites), crowding, and a relative lack of hygiene. These circumstances are generally more common in children than in adults and may explain why most initial infections occur in children and adolescents. Notably, yaws-like strains have been identified from skin lesions of African monkeys, baboons, and gorillas suggesting that *T. pallidum* subsp. *pertenue* may also cause the same disease in nonhuman primates.² To date, there has been no evidence of transmission from nonhuman primates to humans, but it is possible that nonhuman primates may be a potential reservoir for the bacteria.

A primary papule forms at the site of inoculation (commonly the legs, feet, or buttocks) after an incubation period of 9 to 90 days (mean, 21 days). Over a period of 3 to 6 months, the primary papule of yaws may increase in size and then heal spontaneously. Primary lesions are typically pruritic, facilitating autoinoculation; tender regional lymphadenopathy, fever, and constitutional symptoms may also develop. At about the time the primary papule heals, secondary lesions near or distant from the initial lesion often occur. These lesions may also be papular and are thought to be a consequence of local (autoinoculation) and hematogenous or lymphatic spread of infection (Fig. 238.1). In the secondary stage, lesions of yaws predominately involve the skin near the nose and mouth and bones or cartilage of the extremities and, even if untreated, generally heal without scarring (Fig. 238.2). Similar to the primary lesions, lesions of secondary yaws resolve spontaneously without therapy, and patients enter a latent stage that, if untreated, may last a lifetime.⁴ Occasionally, secondary infection or ulceration of cutaneous yaws lesions occurs and results in more pronounced lesions and scarring.

About 10% of untreated patients develop late lesions of yaws characterized by hyperkeratotic plaques, destructive bony lesions, or gummata. In the 1820s, Spaniards first coined the term *gangosa* (nasal voice) to describe these disfiguring chronic manifestations, as they destroy bone,



FIG. 238.1 Initial papillomatous yaws lesion on upper thigh (also called primary frambesioma, mother yaw, *chancre pia nique*). Initial lesion usually commences as a papule on lower extremities and slowly enlarges to form a raspberry-like lesion. (From Ferine PL, Hopkins DR, Niemei PLA, et al. Handbook of Endemic Treponematoses. Geneva: World Health Organization; 1984.)



FIG. 238.2 Early ulceropapillomatous yaws on the leg (also called *ulcère post-chancereux*). (From Ferine PL, Hopkins DR, Niemel PLA, et al. Handbook of Endemic Treponematoses. Geneva: World Health Organization; 1984.)



FIG. 238.3 Gangosa. (From Ferine PL, Hopkins DR, Niemel PLA, et al. Handbook of Endemic Treponematoses. Geneva: World Health Organization; 1984.)

cartilage, and soft tissue of the mouth and nose (Fig. 238.3). Bowing of the anterior tibiae (saber shins) may likewise be a late manifestation of yaws arising from infectious periostitis.

Healing is promoted with therapeutic administration of azithromycin or penicillin. Complete healing may occur without sequelae in patients receiving early treatment. Yaws infections may be prevented with prophylactic administration of azithromycin in exposed patients (see “Therapy” and “Prevention”).

Endemic Syphilis (Bejel)

Endemic syphilis is caused by *T. pallidum* subsp. *endemicum*.^{1,6} Similar to yaws, endemic syphilis is a disease of childhood and adolescence, with most cases occurring in individuals 2 to 15 years of age. Not only does transmission of endemic syphilis occur via direct contact, but

transmission via fomites (i.e., on shared eating or drinking utensils) also may occur because early lesions are often mucosal. The primary lesions of endemic syphilis are typically mucous patches on oral pharyngeal mucosa or lesions at the angles of the lips (angular stomatitis) and may go unobserved. These painless lesions may resolve without therapy, but secondary lesions may appear 3 to 6 months later. Secondary lesions of endemic syphilis manifest as rashes, as mucosal lesions, or with bony and cartilaginous involvement. Similar to secondary syphilis, endemic syphilis rashes in the secondary stage may display a wide variety of morphologies including disseminated papular lesions and condylomata lata in moist areas of the skin accompanied by regional lymphadenopathy.

After a period of latency shorter than for yaws (6 months to several years), late complications of endemic syphilis are relatively common.⁴ These complications may manifest as gummatous lesions or chronic ulcerative skin lesions in 25% to 50% of patients. In contrast to early lesions, late lesions tend to be destructive, chronic progressive lesions, some of which progress to deforming bony and cartilaginous facial lesions referred to as *gangosa* (see “Yaws”). In addition to the ulcerative lesions of endemic syphilis, bony involvement may occur and is manifested as osteoperiostitis of the long bones and hands causing disability and deformity.

Pinta

Pinta is the endemic treponematoses caused by a unique treponemal species, *T. carateum*.⁶ The peak age prevalence for pinta is older than for yaws or endemic syphilis, with most individuals 15 to 30 years of age. Pinta is also thought to spread between individuals through direct lesion contact. An important difference between pinta and the other endemic treponematoses is that without treatment the lesions tend to persist. The classic initial lesion of pinta is a papule or erythematous epithelial plaque. These lesions tend to occur on parts of the body that are not typically clothed, most often the leg, foot, forearm, or back of the hands where skin to skin contact is common. These lesions gradually enlarge through local extension to form hyperkeratotic pigmented lesions. The lesions of pinta are also accompanied by regional lymphadenopathy. Disseminated, small lesions (pintids) may occur distal to the initial lesion 3 to 9 months after infection and slowly enlarge and coalesce. Over time, the lesions of pinta become pigmented taking on a darker color described as slate blue. Late pinta is characterized further by additional pigmentary cutaneous changes; lesions may include dyschromic treponeme-containing lesions and achromic treponeme-free lesions. The depigmentation process of pinta occurs at different rates in the same lesion, giving the lesions a mottled appearance. No other disability or late complications of pinta have been described.

Attenuated Disease

A mild presentation of both yaws and endemic syphilis has been described in regions of lower prevalence. In highly endemic areas the papillomatous stage of yaws tends to last months to years, but in areas of low endemicity the stage is shortened and characterized by fewer, dry crops of lesions in the intertriginous regions.^{1,6} Similarly in areas of lower prevalence, endemic syphilis has been characterized by fewer early and late lesions lasting a shorter period of time. Multiple factors are thought to lead to attenuated disease including improved hygiene, nutrition, and access to treatment.⁶

DIAGNOSIS

Diagnosis of individual cases of the endemic treponematoses in large part depends on recognition of appropriate clinical findings and is confirmed with serologic testing with tests for syphilis.⁶

Direct Visualization

Although not widely available, demonstration of characteristic treponemes with silver staining or darkfield microscopy of lesion exudates or immunofluorescent antibody stains for treponemes can provide immediate and highly specific diagnosis of the endemic treponematoses. However, facilities capable of performing these methods are limited in low-income and middle-income countries where endemic treponematoses are common. Therefore these methods are rarely used for diagnosis.

Serology

No serologic tests have been specifically developed for diagnosis of endemic treponematoses; however, because the humoral antibody response to these diseases is indistinguishable from the response to syphilis, serologic tests for syphilis are important for confirmation of clinically suspected infections. Serologic tests for syphilis and the endemic treponematoses are divided into nontreponemal and treponemal tests.¹⁸ The nontreponemal assays are based on cross-reactivity of cardiolipin-cholesterol-lecithin antigens with antibodies to *T. pallidum* and include tests such as the rapid plasma reagin (RPR) card test and Venereal Disease Research Laboratory (VDRL) tube tests. These tests provide quantifiable results that are useful not only for screening for infection but also for evaluation of response to therapy after treatment. In contrast, the treponemal assays are based on the reactivity of antibodies to either recombinant *T. pallidum* antigens or antigens from *T. pallidum* propagated in laboratory animals. The treponemal tests are available in a variety of formats including fluorescent treponemal antibody absorption, *T. pallidum* hemagglutination, or several different commercial enzyme-linked immunosorbent assays. False-positive results sometimes occur with both nontreponemal and treponemal tests. Because treponemal and nontreponemal tests are unrelated, use of an unrelated test to confirm an initial test result (i.e., confirmatory testing of reactive RPR or VDRL test results with a treponemal test) greatly increases the specificity of test results.

Six months after effective treatment, most patients with active endemic treponematoses have a fourfold (two dilutions) or greater decline in RPR or VDRL titers. Reinfection or relapse may be indicated with a VDRL or RPR titer that increases two or more dilutions. Reversion of treponemal serologic test results to nonreactive is considerably less common than for nontreponemal tests. Early in the disease, patients may have negative serologic tests.¹⁶ Patients with latent infections will have positive serologic tests. Patients with advanced disease may have nontreponemal tests that have low titers or become nonreactive.²

Point-of-Care Syphilis and Endemic Treponematoses Test

A simple, point-of-care test based on simultaneous detection of both treponemal and nontreponemal antibodies has been developed for rapid diagnosis and treatment of syphilis and endemic treponematoses.¹⁹ This immunochromatographic lateral flow assay detects both immunoglobulins (IgM and IgG) against a recombinant *T. pallidum* antigen (T1) and a nontreponemal antigen (T2). This assay may be used for testing serum, plasma, or whole blood; has a 1-year shelf life at room temperature; distinguishes between past and active infection; confirms the diagnosis in real time for patients with skin lesions; and can be performed outside of clinics, providing remote communities access to diagnostic testing. In addition, the test is useful for estimation of population prevalence of endemic treponematoses and detection of continued transmission after control efforts (e.g., mass treatment programs). A meta-analysis of 7267 tests showed good performance, but sensitivity was higher in patients with high-titer (RPR $\geq 1:16$) infection compared with low-titer (RPR $< 1:16$) infection for both the T1 component (98.2% vs. 90.1%, $P < .0001$) and the T2 component (98.2% vs. 80.6%, $P < .0001$).²⁰ As the number of active cases declines after a mass treatment campaign, a diagnostic test with higher sensitivity or repeat testing may be needed to confirm that all cases have been identified and treated.

Polymerase Chain Reaction

The use of polymerase chain reaction (PCR) for surveillance has been proposed to improve diagnostic accuracy of skin ulcers, especially in settings where other causative pathogens may be coendemic and cause similar lesions.²¹ Treponemal DNA is detected with use of a PCR assay alone, which applies a qualitative or quantitative amplification-based method to detect single genes or with reverse-transcriptase PCR followed by the use of labeled primers to target amplicons from *T. pallidum* subsp. *pallidum* 16S ribosomal RNA. Specific areas in the *T. pallidum* genome that distinguish it from other bacterial organisms have been identified including *tpf-1*, *bmp*, *tmpA*, 47-kDa lipoprotein, and *polA*.⁴ However, PCR testing is expensive, requiring electricity and trained laboratorians at equipped laboratory facilities to perform the test (which

may not be available in field conditions); access and delivery of specimens from remote areas may also be difficult. The only reliable method at the present time to distinguish *T. pallidum* subspecies (*pallidum*, *pertenue*, and *endemicum*) is the combined use of PCR and DNA sequencing, which identifies molecular differences between subspecies (e.g., *gpd*, *tpnC*, *tpnI*, *arp*, IGR19), but availability is limited to a few reference laboratories.⁵

THERAPY

Benzathine penicillin G had been the preferred drug for treatment of the endemic treponematoses.^{1,2,6} The need for refrigerated storage of penicillin, the inconvenience of injections for patients, the potential for allergic reactions to β -lactam antibiotics, and the risk of needlestick injuries to health care providers are drawbacks for many countries. A randomized trial in 2012 demonstrated that a single oral dose of 30 mg/kg of azithromycin (up to a maximal dose of 2 g) was noninferior to benzathine penicillin G for treatment of early yaws. Cure rates are similar with recommended doses of azithromycin or benzathine penicillin G.¹⁴ Based on these landmark findings, the WHO now recommends azithromycin as the preferred treatment for endemic treponematoses.⁸

Observational studies in the Pacific region where endemic treponematoses and trachoma occur concurrently suggest that lower doses of azithromycin (20 mg/kg, up to maximal dose of 1 g) typically used to treat trachoma may also be effective against endemic treponematoses.^{22,23} Studies comparing efficacy of the two doses (30 mg/kg and 20 mg/kg) of azithromycin for treatment are ongoing. If the infected individual is intolerant to azithromycin, WHO recommends a single dose of benzathine penicillin G, 600,000 U for children younger than 10 years of age or 1.2 million U for individuals 10 years of age or older, may be administered for treatment. Despite the lack of formal studies, extrapolation from experience with sexually transmitted syphilis suggests that tetracycline or doxycycline given for 14 days may also be effective.

Response to therapy may be determined through resolution of early lesions or, in patients with late infection, arrest of progression. Serologic response to therapy may also be documented by declines in RPR or VDRL titers of two or more dilutions 6 months following treatment (as with sexually transmitted syphilis). After treatment, however, titers may not revert to seronegativity.

PREVENTION

In the 1950s it was estimated that more than 50 to 160 million people worldwide were infected with yaws, of which 40 million may have had permanent disfigurement of the face, joints, bones, and soft tissues. Because of the large geographic distribution and disease burden, the WHO led a global control program from 1952 to 1964 using active serologic screening and staged treatment in 46 countries.^{6,7} In this effort, about 300 million examinations were performed, and more than 50 million people were treated. Prevalence of endemic treponematoses was reduced by more than 95%. However, as disease prevalence waned, so did resources and commitment for continued surveillance and control efforts.^{2,24} Infection rates of yaws once again rebounded^{6,7} with global prevalence of disease estimated at about 460,000 cases in the late 1990s; however, this is likely an underestimate.¹

In the past decade, progress has been made to eliminate yaws. The potential effectiveness of community-based control efforts has been demonstrated in India, where reimplementing the control measures reduced the number of reported new cases from 3571 in 1996 to 0 cases in 2004 followed by negative serologic surveys among children younger than 5 years old for 3 consecutive years. In 2006 yaws was eliminated from India and formally declared eradicated by the WHO in 2016.²⁴ Ecuador also reported elimination of yaws through a mass treatment program.²⁵

Based on the confirmed success in India and the potential to simplify provision of therapy for entire communities, in 2012 the WHO announced a new, aggressive roadmap that targets global elimination of yaws by 2020.^{7,8} In contrast to the staged treatment strategy of the 1950s, two new treatment strategies will simplify case detection and treatment of active and latent disease. Total community treatment is the treatment of an entire community regardless of active cases. This strategy first aims for 100% treatment coverage of a confirmed endemic village or

community followed by surveillance and, if necessary, incorporates mop ups with additional treatment of identified cases. Total targeted treatment, the treatment of all active clinical cases and contacts, is used every 3 to 6 months to detect and treat remaining active cases.⁸ Pilot total community treatment programs with azithromycin in some endemic countries (Solomon Islands, Papua New Guinea, and Ghana) demonstrated the potential for eradication of endemic treponematoses.^{22,26,27} A longitudinal study followed the 16,092 inhabitants of a Papua New Guinea island (Lihir Island) in which nearly 84% received a dose of azithromycin. Although prevalence of yaws initially declined, it began steadily increasing 2 years later, likely due to relapse of untreated latent infections and reintroduction of yaws through in-migration. This study is the first to identify a point mutation in a yaws strain that conferred

resistance to macrolides, previously unreported among endemic treponematoses. For successful eradication, the WHO now recommends treatment coverage rates should exceed 90%, and if rates are less, another round of mass treatment should be considered along with close monitoring for macrolide resistance.²⁸

Scientific advances including development of point-of-care and molecular diagnostics will aid in achieving the 2020 goal. However, further studies are needed to continually improve these eradication strategies. The future of endemic treponemal research lies in improving surveillance and geographic mapping, integrating point-of-care testing, determining etiology of nontreponematoses skin ulcers by using advanced molecular diagnostics, monitoring for macrolide resistance, and looking for *T. pallidum* infections in nonhuman primates.²⁵

Key References

The complete reference list is available online at Expert Consult.

3. Cejkova D, Zbanikova M, Chen L, et al. Whole genome sequences of three *Treponema pallidum* ssp. *pertenue* strains: yaws and syphilis treponemes differ in less than 0.2% of the genome sequence. *PLoS Negl Trop Dis*. 2012;6:e1471.
4. Giacani L, Lukehart SA. The endemic treponematoses. *Clin Microbiol Rev*. 2014;27:89–115.
8. Eradication of yaws—the Morges strategy. *Wkly Epidemiol Rec*. 2012;87:189–194.
14. Mitja O, Hays R, Ipai A, et al. Single-dose azithromycin versus benzathine benzylpenicillin for treatment of yaws in children in Papua New Guinea: an open-label, non-inferiority, randomised trial. *Lancet*. 2012;379:342–347.
19. Castro AR, Esfandiari J, Kumar S, et al. Novel point-of-care test for simultaneous detection of nontreponemal and treponemal antibodies in patients with syphilis. *J Clin Microbiol*. 2010;48:4615–4619.
21. Mitja O, Lukehart SA, Pokowas G, et al. *Haemophilus ducreyi* as a cause of skin ulcers in children from a yaws-endemic area of Papua New Guinea: a prospective cohort study. *Lancet Glob Health*. 2014;2:e235–e241.
22. Marks M, Vahi V, Sokana O, et al. Impact of community mass treatment with azithromycin for trachoma elimination on the prevalence of yaws. *PLoS Negl Trop Dis*. 2015;9:e0003988.

References

1. Antal GM, Lukehart SA, Meheus AZ. The endemic treponematoses. *Microbes Infect.* 2002;4:83–94.
2. Mitja O, Asiedu K, Mabey D. Yaws. *Lancet.* 2013;381:763–773.
3. Cejkova D, Zobanikova M, Chen L, et al. Whole genome sequences of three *Treponema pallidum* ssp. pertenue strains: yaws and syphilis treponemes differ in less than 0.2% of the genome sequence. *PLoS Negl Trop Dis.* 2012;6:e1471.
4. Giacani L, Lukehart SA. The endemic treponematoses. *Clin Microbiol Rev.* 2014;27:89–115.
5. Mitja O, Smajs D, Bassat Q. Advances in the diagnosis of endemic treponematoses: yaws, bejel, and pinta. *PLoS Negl Trop Dis.* 2013;7:e2283.
6. Farnsworth N, Rosen T. Endemic treponematoses: review and update. *Clin Dermatol.* 2006;24:181–190.
7. Asiedu K, Fitzpatrick C, Jannin J. Eradication of yaws: historical efforts and achieving WHO's 2020 target. *PLoS Negl Trop Dis.* 2014;8:e3016.
8. Eradication of yaws—the Morges strategy. *Wkly Epidemiol Rec.* 2012;87:189–194.
9. Stamm LV. Pinta: Latin America's forgotten disease? *Am J Trop Med Hyg.* 2015;93:901–903.
10. Veller-Fornasa C, Tarantello M, Cipriani R, et al. Effect of ofloxacin on *Treponema pallidum* in incubating experimental syphilis. *Genitourin Med.* 1987;63:214.
11. Svindland HB. Treatment of gonorrhoea with sulphamethoxazole-trimethoprim. Lack of effect on concomitant syphilis. *Br J Vener Dis.* 1973;49:50–53.
12. Lukehart SA, Godornes C, Molini BJ, et al. Macrolide resistance in *Treponema pallidum* in the United States and Ireland. *N Engl J Med.* 2004;351:154–158.
13. Mitja O, Codornes C, Houine W, et al. Re-emergence of yaws after single mass azithromycin treatment followed by targeted treatment: a longitudinal study. *Lancet.* 2018;391:1599–1607.
14. Mitja O, Hays R, Ipai A, et al. Single-dose azithromycin versus benzathine benzylpenicillin for treatment of yaws in children in Papua New Guinea: an open-label, non-inferiority, randomised trial. *Lancet.* 2012;379:342–347.
15. Mitja O, Marks M, Konan D, et al. Global epidemiology of yaws: a systematic review. *Lancet Glob Health.* 2015;3:e324–e331.
16. Castro LG. Nonvenereal treponematoses. *J Am Acad Dermatol.* 1994;31:1075–1076.
17. Perine PL, Hopkins DR, Niemei PLA, et al. *Handbook of Endemic Treponematoses.* Geneva: World Health Organization; 1984.
18. Larsen SA, Pope V, Johnson RE, et al, eds. *A manual of tests for syphilis.* 9th ed. Washington DC: American Public Health Association; 1998.
19. Castro AR, Esfandiari J, Kumar S, et al. Novel point-of-care test for simultaneous detection of nontreponemal and treponemal antibodies in patients with syphilis. *J Clin Microbiol.* 2010;48:4615–4619.
20. Marks M, Yin YP, Chen XS, et al. Metaanalysis of the performance of a combined treponemal and nontreponemal rapid diagnostic test for syphilis and yaws. *Clin Infect Dis.* 2016;63:627–633.
21. Mitja O, Lukehart SA, Pokowas G, et al. *Haemophilus ducreyi* as a cause of skin ulcers in children from a yaws-endemic area of Papua New Guinea: a prospective cohort study. *Lancet Glob Health.* 2014;2:e235–e241.
22. Marks M, Vahi V, Sokana O, et al. Impact of community mass treatment with azithromycin for trachoma elimination on the prevalence of yaws. *PLoS Negl Trop Dis.* 2015;9:e0003988.
23. Ghinai R, El-Duah P, Chi KH, et al. A cross-sectional study of 'yaws' in districts of Ghana which have previously undertaken azithromycin mass drug administration for trachoma control. *PLoS Negl Trop Dis.* 2015;9:e0003496.
24. World Health Organization. Eradication of yaws in India. *Wkly Epidemiol Rec.* 2015;90:161–168.
25. Marks M, Mitja O, Vestergaard LS, et al. Challenges and key research questions for yaws eradication. *Lancet Infect Dis.* 2015;15:1220–1225.
26. Mitja O, Houine W, Moses P, et al. Mass treatment with single-dose azithromycin for yaws. *N Engl J Med.* 2015;372:703–710.
27. Abdulai AA, Agana-Nsiire P, Biney F, et al. Community-based mass treatment with azithromycin for the elimination of yaws in Ghana—results of a pilot study. *PLoS Negl Trop Dis.* 2018;12:e0006303.
28. Yaws: strict compliance with WHO's Morges strategy critical to achieve eradication; February 13, 2018. http://www.who.int/neglected_diseases/news/strict_compliance_with_WHO_Morges_strategy/en/. Accessed May 25, 2018.

SHORT VIEW SUMMARY

Definition

- Leptospirosis is caused by infection with pathogenic spirochetes of the genus *Leptospira*.

Epidemiology

- Leptospirosis is a zoonosis of global distribution.
- Leptospirosis is maintained in nature by chronic renal infection of carrier animals, especially rodents.
- Infections typically occur after occupational or recreational exposure to water or soil contaminated with rodent urine.
- In developing countries with poor housing standards, leptospirosis outbreaks occur regularly in urban settings after heavy rainfall and flooding.

Microbiology

- Twenty-three named leptospiral species have been described, of which 10 are known to be pathogenic (see Table 239.1).

Diagnosis

- Clinical diagnosis requires a high index of suspicion based on epidemiologic exposure. The signs and symptoms of early leptospirosis are nonspecific.
- Serologic tests are helpful when positive but may have poor sensitivity during the first week of illness. Polymerase chain reaction assay of serum and urine appears to be a promising diagnostic technique. Culture can be done using blood early in the course and urine later on but requires special media.

Therapy

- Antibiotic therapy (see Table 239.4) should be initiated as early in the course of the disease as suspicion allows. For mild disease, doxycycline, amoxicillin, or oral ampicillin is recommended. For more severe disease, intravenous ceftriaxone, ampicillin, or penicillin is used.

- Patients with early renal disease with high-output renal dysfunction and hypokalemia should receive aggressive volume repletion and potassium supplementation.
- Patients who progress to oliguric renal failure should undergo rapid initiation of hemodialysis or peritoneal dialysis, which is typically required on only a short-term basis.

Prevention

- The most effective preventive strategy is to reduce direct contact with potentially infected animals and indirect contact with urine-contaminated soil and water.
- Vaccination of agricultural and companion animals is practiced, and vaccines for humans are used in France, Cuba, and parts of Asia.

HISTORY

A syndrome of severe multisystem disease, presenting with profound jaundice and renal function impairment, was described by Weil in Heidelberg in 1886. Other descriptions of disease that probably represent leptospirosis were made earlier, but the etiology cannot be definitively ascribed to leptospiral infection.² Leptospire were first visualized in autopsy specimens from a patient thought to have had yellow fever³ but were not isolated until several years later, almost simultaneously in Germany and Japan.⁴ Diagnostic confusion between severe icteric leptospirosis and yellow fever continued, with prominent researchers such as Stokes and Noguchi dying in their attempts to discover the etiologic agent.⁴ Several authoritative reviews have been published.^{2,4,5,6,7,8}

ETIOLOGY

"Leptospira" derives from the Greek *leptos* (thin) and Latin *spira* (coiled). Aptly named, the leptospire are a mere 0.1 μm in diameter by 6 to 20 μm in length. The cells have pointed ends, one or both of which is usually bent into a characteristic hook (Fig. 239.1). Motility is conferred by the rotation of two axial flagella underlying the membrane sheath, which are inserted at opposite ends of the cell and extend toward the central region.⁹ Because of their small diameter, leptospire are best visualized by darkfield microscopy, appearing as actively motile spirochetes (Fig. 239.2). Leptospire are readily cultured in polysorbate-albumin media if specimens are obtained before initiation of antibiotic therapy.¹⁰

Historically, the genus *Leptospira* was classified into two species, *L. interrogans* and *L. biflexa*, composed of pathogenic and nonpathogenic strains, respectively. Within each species, large numbers of serovars

were differentiated using agglutinating antibodies. Serovar specificity is conferred by lipopolysaccharide (LPS) O-antigens.¹¹ More than 250 serovars of pathogenic leptospire have been described; because of the large number of serovars, antigenically related serovars were grouped into serogroups for convenience in serologic testing.

Leptospire are now classified into a number of species defined by their degree of genetic relatedness, determined by DNA reassociation.^{12,13} There are currently 23 named species, including pathogens (e.g., *L. interrogans*), nonpathogenic saprophytes (e.g., *L. biflexa*), and species of indeterminate pathogenicity (e.g., *L. inadai*) (Table 239.1).¹⁴ This classification is supported by 16S RNA gene sequencing (Fig. 239.3)¹⁵ but is quite distinct from the former serologic classification.⁵

The system of serogroup nomenclature has no taxonomic standing but is retained because presumptive serogroup determination by serologic testing has some epidemiologic value. However, serologic responses may bear surprisingly little relationship to the infecting serovar in individual patients.¹⁶

The genome sequences of several *Leptospira* spp. and strains have been determined,^{17,18,19,20,21} and sequencing of another 200 strains is under way through the *Leptospira* Genomics and Human Health Project. The availability of these genome sequences has already led to better understanding of leptospiral pathogenesis.²²

EPIDEMIOLOGY AND TRANSMISSION

Globally, leptospirosis is a leading zoonotic cause of morbidity and mortality that disproportionately affects vulnerable populations, such as rural subsistence farmers and urban slum dwellers. Human infections




FIG. 239.1 Scanning electron micrograph of cells *leptospira interrogans* showing helical structure and curved (hooked) ends (original magnification $\times 60,000$). (Courtesy Rob Weyant, Centers for Disease Control and Prevention.)




FIG. 239.2 Leptospire viewed by darkfield microscopy (original magnification $\times 100$). (Courtesy Mildred Galton, Public Health Image Library, Centers for Disease Control and Prevention.)

are endemic in most regions, and the peak incidence occurs in the rainy season in tropical regions and the late summer to early fall in temperate regions.²³ In developing countries with poor urban housing standards, leptospirosis outbreaks occur regularly after heavy rainfall and flooding.²⁴ In the United States the highest incidence is found in Hawaii; active surveillance in 1992 detected an annual incidence of approximately 128 per 100,000.²⁵ Based on a systematic review of 80 published studies and databases from 34 countries, it is estimated that there are 1.03 million cases and 58,900 deaths each year.²⁶ Morbidity and mortality are greatest in the poorest regions of the world and in areas where surveillance is not routinely performed. It is likely that these figures still underestimated the burden because leptospirosis patients are commonly misdiagnosed with dengue, malaria, and other infections. More important than incidence, estimates suggest that globally ~ 2.90 million of disability adjusted life-years (DALYs) are lost per annum from the approximately 1.03 million cases.²⁷ This estimate of DALYs from leptospirosis is more than four times greater than the ~ 0.83 million DALYs due to dengue.²⁸

Leptospirosis is maintained in nature by chronic renal infection of carrier animals. The most important reservoirs are rodents and other small mammals, but livestock and companion animals are also significant sources of human infection. Infection of carrier animals usually occurs during infancy, and once infected, animals may shed leptospire in their urine intermittently or continuously throughout life.

Infection occurs through direct or indirect contact with urine or tissues of infected animals. Direct contact is important in transmission to veterinarians, workers in milking sheds on dairy farms, abattoir




FIG. 239.3 Unrooted phylogenetic tree based on 16S RNA gene sequences of the leptospiraceae obtained from genbank. Species composed of pathogens (see Table 239.1) cluster separately from saprophytes (nonpathogenic species) and species whose pathogenicity is intermediate.

TABLE 239.1 Species of *Leptospira* and Some Pathogenic Serovars

SPECIES	SELECTED PATHOGENIC SEROVARS
<i>L. interrogans</i>	Icterohaemorrhagiae, Copenhageni, Canicola, Pomona, Australis, Autumnalis, Pyrogenes, Bratislava, Lai
<i>L. kirschneri</i>	Bim, Bulgarica, Grippotyphosa, Cynopteri
<i>L. borgpetersenii</i>	Ballum, Hardjo, Javanica
<i>L. mayottensis</i>	Kenya, Mini
<i>L. santarosai</i>	Bataviae
<i>L. noguchii</i>	Panama, Pomona
<i>L. weilii</i>	Celledoni, Sarmin
<i>L. alexanderi</i>	Manhao 3
<i>L. alstonii</i>	Sichuan
<i>L. kmetyi</i>	Manilae
<i>L. inadai</i>	Intermediate
<i>L. fainei</i>	Intermediate
<i>L. broomii</i>	Intermediate
<i>L. wolffii</i>	Intermediate
<i>L. licerasiae</i>	Intermediate
<i>L. venezuelensis</i>	Intermediate
<i>L. wolbachii</i>	Nonpathogen
<i>L. biflexa</i>	Nonpathogen
<i>L. vanthiellii</i>	Nonpathogen
<i>L. meyeri</i>	Nonpathogen
<i>L. terpstrae</i>	Nonpathogen
<i>L. yanagawae</i>	Nonpathogen
<i>L. idonii</i>	Nonpathogen

workers, butchers, hunters, and animal handlers; transmission to children handling puppies and to dog handlers has been reported. Males are predominantly affected, with approximately 80% of the total burden.²⁷ Indirect contact is more common and is responsible for disease after exposure to wet soil or water. The great majority of cases are acquired by this route in the tropics, either through occupational exposure to water, as in rice or taro farming, or through exposure to damp soil and water during avocational activities. Overseas operations place the US military, including military dogs, at risk of leptospirosis. This is particularly true of Special Operations Forces, who work closely with local military.²⁹

Recreational exposures have become relatively more important, often in association with adventure tourism to tropical endemic areas. Several large point-source waterborne outbreaks involving attack rates as high as 42% have occurred recently after athletic events.^{30,31} In recent years there has been an increase in leptospirosis cases among dogs in the eastern regions of North America and in the Midwest,³² associated with a shift in the predominant serovars causing disease.^{33,34} Veterinary vaccine manufacturers have responded to this problem by licensing new canine vaccines containing these emerging serovars.

PATHOGENESIS

Leptospire enter the body through cuts and abrasions, mucous membranes or conjunctivae, and aerosol inhalation of microscopic droplets. Swallowing contaminated lake water was the only behavioral risk factor identified in a case-control study of a large leptospirosis outbreak at the 1998 Springfield Triathlon.³⁰ However, the oral mucosae are probably a more important route of entry after ingestion than the intestinal tract. On entering the body there is widespread hematogenous dissemination, with levels as high as 10^6 leptospire per milliliter of blood documented by quantitative polymerase chain reaction (PCR).³⁵ Organisms readily penetrate tissue barriers, resulting in invasion of the central nervous system and aqueous humor of the eye. Transendothelial migration of spirochetes is facilitated by a systemic vasculitis, accounting for a broad spectrum of clinical illness. Severe vascular injury can ensue, leading to pulmonary hemorrhage, ischemia of the renal cortex and tubular-epithelial cell necrosis, and destruction of the hepatic architecture, resulting in jaundice and liver cell injury with or without necrosis.³⁶

The mechanisms by which leptospire cause disease are not clearly understood. Potential virulence factors include immune mechanisms, toxin production, adhesins, and other surface proteins. Human susceptibility to leptospirosis may be related to poor recognition of leptospiral LPS by the innate immune system.^{37,38} Human Toll-like receptor 4 (TLR4), which responds to extremely low concentrations of gram-negative LPS (endotoxin), appears to be unable to bind leptospiral LPS,^{38,39} perhaps because of the unique methylated phosphate residue of its lipid A.⁴⁰ In contrast to human TLR4, mouse TLR4 can recognize leptospiral LPS and has been shown to play an important role in preventing fatal leptospirosis infection in mice.^{39,41} Direct tissue damage may also be due to production of hemolytic toxins, which may act as sphingomyelinases, phospholipases, or pore-forming proteins, and collagenases.⁴²⁻⁴⁴

Immune-mediated mechanisms have been postulated as one factor influencing the severity of symptoms.⁴⁵ Investigation of the same triathlon outbreak mentioned earlier identified the human leukocyte antigen (HLA) DQ6 as an independent risk factor for leptospirosis.⁴⁶ The structural location of HLA-DQ6 polymorphisms associated with disease suggested that leptospire produce a superantigen that can cause nonspecific T-cell activation in susceptible individuals. Other immune mechanisms, including circulating immune complexes, anticardiolipin antibodies, and antiplatelet antibodies, have been proposed, but their significance is unproven. Although immune or autoimmune mechanisms may play a role in some cases of leptospiral uveitis,⁴⁷ a high percentage of such patients have leptospiral DNA in the aqueous humor by PCR.⁴⁸

Much recent work has focused on the roles of surface lipoproteins in leptospiral pathogenesis.⁴⁹ The major outer membrane lipoprotein LipL32 is highly conserved among pathogenic serovars.⁵⁰ LipL32 is a major target of the human immune response⁵¹ and appears to be involved in pathogenesis of tubulointerstitial nephritis.⁵² Virulent leptospire respond to the increased osmolarity of host tissues by inducing expression of the multifunctional Lig surface proteins that mediate interactions

with a variety of host proteins, including extracellular matrix factors.⁵³ The Lig proteins are early antigens; IgM antibodies to their immunoglobulin-like repeats develop early in infection, offering an approach to improved detection of acute infection.⁵⁴ The endostatin-like LenA protein binds the complement regulatory protein, factor H, and pathogenic strains exhibit proteolytic activity that cleaves complement, suggesting an important role in serum resistance.^{55,56}

CLINICAL MANIFESTATIONS

Leptospiral infection is associated with a broad spectrum of severity, ranging from subclinical illness, followed by seroconversion to two clinically recognizable syndromes: a self-limited, systemic illness seen in roughly 90% of infections and a severe, potentially fatal illness accompanied by any combination of renal failure, liver failure, and pneumonitis with hemorrhagic diathesis.^{2,6,7} In some patients the disease has two distinct phases: an initial septicemic stage that is followed by a temporary decline in fever, followed by an immune phase in which the severe symptoms occur. However, in many severe cases the distinction between these two phases is not apparent; in addition, many patients present only with the onset of the second phase of the illness.

The mean incubation period is 10 days, ranging from 5 to 14 days; determination of precise exposures may be difficult, leading to significant imprecision in estimated incubation times. The acute, septicemic phase of illness begins abruptly with high, remittent fever (38°–40°C) and headache, chills, rigors, and myalgias; conjunctival suffusion (redness without exudate); abdominal pain; anorexia, nausea, and vomiting; diarrhea; and cough and pharyngitis. A pretibial maculopapular cutaneous eruption occurs rarely (Table 239.2). Conjunctival suffusion and muscle tenderness, most notable in the calf and lumbar areas, are the most characteristic physical findings but may occur in a minority of cases (see Table 239.2). Other less common signs include lymphadenopathy, splenomegaly, and hepatomegaly. The acute phase lasts from 5 to 7 days. Routine laboratory tests are nonspecific but indicative of a bacterial infection. Leptospire can be recovered from blood and cerebrospinal fluid (CSF) during the acute phase of illness, but meningeal signs are not prominent in this phase. Leptospire may also be recovered from urine, beginning about 5 to 7 days after the onset of symptoms (Fig. 239.4). Urinalysis reveals mild proteinuria and pyuria, with or without hematuria and hyaline or granular casts. Death is rare in the acute phase of illness.

The immune phase of illness generally lasts from 4 to 30 days (see Fig. 239.4). The disappearance of leptospire from the blood and CSF coincides with the appearance of immunoglobulin M (IgM) antibodies.^{7,57} The organisms can be detected in almost all tissues and organs and in urine for several weeks, depending on the severity of the disease. In addition to the acute-phase symptoms described in the preceding paragraph, the immune phase may be characterized by any or all of the following signs and symptoms: jaundice, renal failure, cardiac arrhythmias, pulmonary symptoms, aseptic meningitis, conjunctival suffusion with or without hemorrhage; photophobia; eye pain; muscle tenderness; adenopathy; and hepatosplenomegaly (see Table 239.2). Abdominal pain is not uncommon and may be an indication of pancreatitis.

Aseptic meningitis, with or without symptoms, is characteristic of the immune phase of illness, occurring in up to 80% of cases. In endemic areas a significant proportion of all aseptic meningitis cases may be caused by leptospiral infection.⁵⁸ Symptomatic patients present with an intense, bimodal and frontal throbbing headache with or without delirium. A lymphocytic pleocytosis occurs, with total cell counts generally less than 500/mm.³ CSF protein levels are modestly elevated, between 50 and 100 mg/mL; the CSF glucose concentration is normal. Severe neurologic complications, such as meningoencephalitis, hemiplegia, transverse myelitis, cerebral venous thrombosis, or Guillain-Barré syndrome, occur only rarely.^{5,59} Uveitis is reported in horses and humans and is associated with the second phase of illness, perhaps accounting for underreporting.^{60,61} An unusual case of chronic meningoencephalitis in an immunocompromised child was diagnosed only by next-generation sequencing.⁶²

The most distinctive form of severe illness that may develop after the acute phase of illness is Weil disease, characterized by impaired hepatic and renal function. More severe cases may progress directly

TABLE 239.2 Signs and Symptoms on Admission in Patients With Leptospirosis in Large Case Series

PERCENT WITH:	PUERTO RICO, 1963 ¹⁵⁰ (N = 208)	CHINA, 1965 ¹⁵¹ (N = 168)	VIETNAM, 1973 ¹⁵² (N = 93)	KOREA, 1987 ¹⁵³ (N = 150)	BARBADOS, 1990 ⁶³ (N = 88)	SEYCHELLES, 1998 ¹⁵⁴ (N = 75)	BRAZIL, 1999 ²⁴ (N = 93)	HAWAII, 2001 ¹⁵⁵ (N = 353)	INDIA, 2002 ¹⁵⁶ (N = 74)
Jaundice	49	0	1.5	16	95	27	93	39	34
Anorexia	—	46	—	80	85	—	—	82	—
Headache	91	90	98	70	76	80	75	89	92
Conjunctival suffusion	99	57	42	58	54	—	28.5	28	35
Vomiting	69	18	33	32	50	40	—	73	—
Myalgia	97	64	79	40	49	63	94	91	68
Arthralgia	—	36	—	—	—	31	—	59	12
Abdominal pain	—	26	28	40	43	41	—	51	—
Nausea	75	29	41	46	37	—	—	77	—
Dehydration	—	—	—	—	37	—	—	—	—
Cough	24	57	20	45	32	39	—	—	—
Hemoptysis	9	51	—	40	—	13	20	—	35
Hepatomegaly	69	28	15	17	27	—	—	16	—
Lymphadenopathy	24	49	21	—	21	—	—	—	15
Diarrhea	27	20	29	36	14	11	—	53	—
Rash	6	—	7	—	2	—	—	8	12




FIG. 239.4 Biphasic nature of leptospirosis and relevant investigations at different stages of disease. Specimens 1 and 2 for serology are acute-phase specimens; 3 is a convalescent-phase sample, which may facilitate detection of a delayed immune response; and 4 and 5 are follow-up samples that can provide epidemiologic information, such as the presumptive infecting serogroup. CSF, Cerebrospinal fluid. (Modified from Turner LH. *Leptospirosis*. Br Med J. 1969;1:231–235; and Levett PN. *Leptospirosis*. Clin Microbiol Rev. 2001;14:296–326.)

from the acute phase without the characteristic brief improvement in symptoms to a fulminant illness, with fever greater than 40°C and the rapid onset of liver failure, acute renal failure, hemorrhagic pneumonitis, cardiac arrhythmia, or circulatory collapse.⁷ Mortality rates in patients developing severe disease have ranged from 5% to 40%.^{2,5,6,63} In a study

of 840 hospitalized patients with severe leptospirosis (14% case-fatality rate), risk of death was found to increase with age, especially in adults 40 years of age or older.^{24,64} Altered mental status has been found to be the strongest predictor of death.^{24,65} Other poor prognostic signs include acute renal failure (oliguria, hyperkalemia, serum creatinine >3 mg/

dL), respiratory insufficiency (dyspnea, pulmonary rales, chest radiograph infiltrates), hypotension, and arrhythmias.²⁴ In jaundiced patients disturbance of liver function is out of proportion to the rather mild and nonspecific pathologic findings. Conjugated serum bilirubin levels may rise to 80 mg/dL, accompanied by more modest elevations of serum aminotransferases, alanine aminotransferase and aspartate aminotransferase, which rarely exceed 200 IU/L.⁶⁶ This is in marked contrast to viral hepatitis. Jaundice is slow to resolve, but death due to liver failure almost never occurs in the absence of renal failure. At autopsy, degenerative changes are seen in hepatocytes, Kupffer cells may be hypertrophied, cholestasis is evident, and erythrophagocytosis and mononuclear cell infiltrates are observed.⁶⁷ Hepatocellular necrosis is absent.

Kidney involvement is initially characterized by a unique nonoliguric hypokalemic form of renal insufficiency. Hallmarks are impaired sodium reabsorption, increased distal sodium delivery, and potassium wasting. The impairment in sodium reabsorption appears to be due to selective loss of the epithelial sodium channel in the proximal tubular epithelium. The blood urea nitrogen level is usually less than 100 mg/dL, and the serum creatinine level is usually less than 2 to 8 mg/dL during the acute phase of illness.⁶⁸ Thrombocytopenia occurs in the absence of disseminated intravascular coagulation and may accompany progressive renal dysfunction.⁶⁹ Renal biopsy reveals acute interstitial nephritis; immune-complex glomerulonephritis may also be present.⁷⁰ If electrolyte and volume losses are not replaced, patients progress to oliguric renal failure. In fatal cases the kidneys are swollen and yellow, with prominent cortical blood vessels.³⁶ Histologic findings include a diffuse, mixed tubulointerstitial inflammatory cell infiltrate of lymphocytes, plasma cells, macrophages, and polymorphonuclear leukocytes, accompanied by focal areas of tubular necrosis.⁶⁷ Acute kidney injury in leptospirosis is associated with elevated markers of endothelial injury.⁷¹

Severe pulmonary hemorrhage syndrome (SPHS) can be a prominent manifestation of infection and may occur in the absence of hepatic and renal failure.⁷² Frank hemoptysis can arise simultaneously with the onset of cough during the acute phase of illness.⁷³ However, hemorrhage is often inapparent until patients are intubated; clinicians should suspect SPHS in patients with signs of respiratory distress whether or not they have hemoptysis. With progressive pulmonary involvement, radiographic abnormalities seen most frequently in the lower lobes evolve from small

nodular densities ("snowflake-like") to patchy alveolar infiltrates; confluent consolidation is uncommon but may occur.⁷⁴ The pathophysiology of SPHS is consistent with acute respiratory distress syndrome (ARDS) with diffuse lung injury, impaired gas exchange, and hemodynamic changes indicative of septic shock.⁷⁵ At autopsy, the lungs appear grossly congested and demonstrate focal areas of hemorrhage.⁶⁷ Histologically, damage to the capillary endothelium leads to congestion with foci of interstitial and intraalveolar hemorrhage, diffuse alveolar damage, and severe airspace disorganization.⁷⁶ Although inflammatory infiltrates are usually absent, a community-acquired pneumonia has been described in Indonesia, an area of high prevalence, from four patients who presented with fever, dry cough, dyspnea, and conjunctival suffusion or scleral icterus and crackles on physical examination. Infiltrates on chest radiograph were present in all cases, and the diagnosis of leptospiral pneumonia was supported by real-time PCR from multiple specimens and serologic confirmation.⁷⁷

Congestive heart failure occurs rarely. However, nonspecific electrocardiographic changes are common.⁷⁸ In more than half of patients receiving continuous cardiac monitoring, cardiac arrhythmias may occur, including atrial fibrillation; flutter and tachycardia; and cardiac irritability, including premature ventricular contractions and ventricular tachycardia.⁷⁸ Atrial fibrillation is associated with more severe disease.⁷⁹ Cardiovascular collapse with shock can develop abruptly and in the absence of aggressive supportive care can be fatal. At autopsy, interstitial myocarditis with inflammatory involvement of the conduction system is seen⁸⁰; acute coronary arteritis and aortitis are also common at postmortem examination.⁸¹

LABORATORY DIAGNOSIS

Direct Detection Methods

Direct visualization of leptospires in blood or urine by darkfield microscopic examination has been used for diagnosis (Table 239.3). However, artifacts are commonly mistaken for leptospires, and the method has both low sensitivity (40.2%) and specificity (61.5%).⁸² A range of staining methods has been applied to direct detection, including immunofluorescence staining, immunoperoxidase staining, and silver staining. These methods are not widely used because of the lack of commercially available reagents and their relatively low sensitivity.

TABLE 239.3 Common Diagnostic Tests for Leptospirosis

TEST	SENSITIVITY ^{a,b}	SPECIFICITY ^{a,b}	COSTS ^c	ADVANTAGES	DISADVANTAGES ^d	REFERENCE
Culture	5%–50%	100%	++	Provides definitive evidence. Applicable to human and veterinary diagnosis.	Slow and requires significant expertise	Levett; WHO ¹¹³
Darkfield microscopy (DFM)	10 ⁴ bacteria/mL	Low; confusion with protein fibers	+	Quick and early diagnosis. Applicable to human and veterinary diagnosis.	Unreliable, requires confirmation	Levett, 2001 ⁵ ; WHO ¹¹³
Microscopic agglutination test (MAT)	90%	>90%	+++	Gold standard. ^e Applicable to human and veterinary diagnosis.	Requires a panel of live antigens, difficult (expertise), laborious. Problems with seronegative carrier animals.	Levett, 2001 ⁵ ; WHO ¹¹³
IgM–enzyme-linked immunosorbent assay (ELISA)	>90%	88%–95%	++	Cost effective and relatively rapid (1–2 h)	Serology, needs confirmation by MAT	WHO ¹¹³
Latex agglutination test	82%	95%	++	Easy, quick (30 s), cost effective	Serology, needs confirmation by MAT	WHO ¹¹³
Lateral flow test	81%	96%	++	Easy, quick (10 min), finger prick blood, cost effective	Serology, needs confirmation by MAT	WHO ¹¹³
Real-time polymerase chain reaction (PCR)	100%	93%	+++	Early diagnosis. Applicable to human and veterinary diagnosis.	Few tests validated, sophisticated, expensive equipment, requires expertise	Ahmed et al, 2011 ⁹⁰

^aSensitivity and specificity largely depend on a number of factors: stage of illness, type and producer of test, and panel of clinical materials used for testing.^{5,157}

^bSensitivity of culture depends mainly on the route and transport time to the laboratory. MAT is the gold standard and should be 100%. However, when comparing with culture, this gold standard appears not optimal. Sensitivities of other serologic tests are compared with MAT.

^cExcluding costs for equipment.

^dSerology has the disadvantage that it detects antibodies 7–10 days after the onset of the disease. This is too late for antibiotic treatment, which should start within the first 4 days.

^eSerologic confirmation requires testing of both acute and convalescent serum samples for seroconversion or significant titer rise.

Modified from Hartskeerl RA, Collares-Pereira M, Ellis WA. Emergence, control and re-emerging leptospirosis: dynamics of infection in the changing world. Clin Microbiol Infect. 2011;17:494–501.




FIG. 239.5 Kidney sections stained by silver staining (A) and immunohistochemical staining (B), showing presence of multiple leptospires in tubules. (A from Barnett JK, Barnett D, Bolin CA, et al. *Infect Immun.* 1999;67:853–861.¹⁴⁹ B from Haake DA, Chao G, Zuerner RL, et al. *Infect Immun.* 2000;68:2276–2285.)

Detection of leptospiral antigen in blood or urine has been attempted but without significant success. Many PCR assays^{83–85} and several loop-mediated isothermal amplification assays^{86–88} have been developed for the detection of leptospires, but relatively few have been evaluated in prospective clinical studies,^{88,89} and there have been no multicenter studies of multiple molecular diagnostic methods. The chief advantage of PCR is the prospect of confirming the diagnosis during the early acute (leptospiremic) stage of the illness, before the appearance of IgM antibodies, when treatment is likely to have the greatest benefit.⁹⁰ For early diagnosis, serum or plasma might be the optimal specimen, but urine has been often positive, even early in the course.⁹¹ In fulminating cases, in which death occurs before seroconversion, PCR may be of great diagnostic value.^{83,92} Histologic diagnosis (Fig. 239.5) traditionally relied on silver impregnation staining,³ but immunohistochemical staining offers greater sensitivity and specificity.^{93,94}

Isolation and Identification

Leptospires can be isolated from blood, CSF, and peritoneal dialysate fluids during the first 10 days of illness. Specimens should be collected while the patient is febrile and before antibiotic therapy is initiated. One or two drops of blood should be inoculated directly into culture medium at the bedside. Survival of leptospires in commercial blood culture media or refrigerated blood samples for several days has been reported.^{95,96} Urine can be cultured after the first week of illness. Specimens should be collected aseptically into sterile containers without preservatives and must be processed within a short time of collection; best results are obtained when the delay is less than 1 hour because leptospires do not survive well in acidic environments.¹⁰

Cultures are performed in albumin-polysorbate media such as EMJH (Ellinghausen-McCullough-Johnson-Harris),⁹⁷ which is available

commercially. Primary cultures are performed in semisolid (0.2% agar) medium, to which 5-fluorouracil is usually added as a selective agent. Cultures are incubated at 30°C for several weeks because initial growth may be slow.

Isolated leptospires are identified to serovar level by traditional serologic methods.^{98,99} These techniques are limited in availability to a few reference laboratories. However, molecular methods such as pulsed-field gel electrophoresis show promise for rapid identification of most isolates.¹⁰⁰ Identification to species level is accomplished by sequencing the 16S ribosomal RNA gene of isolates¹⁵ or by sequencing the products of PCR directly.^{90,101} Powerful molecular techniques, such as multiple locus sequence typing and multilocus variable-number tandem-repeat analysis, are being applied to the epidemiologic analysis of leptospirosis.^{102–105} As more genomic data become available, these approaches are becoming more standardized.^{106–109} In addition to molecular typing, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry has been investigated for identification of *Leptospira* isolates.¹¹⁰

Indirect Detection Methods

The majority of leptospirosis cases are diagnosed by serology. The reference standard assay is the microscopical agglutination test (MAT), in which live antigens representing different serogroups of leptospires are reacted with serum samples and then examined by darkfield microscopy for agglutination.¹⁰ Due to regional variations in infecting strains, it has been considered important to include local serovars in a laboratory's MAT panel. Recently, this view has been challenged.¹¹¹ The MAT is a complex test to maintain, perform, and interpret, and participation of the laboratory in a proficiency testing program is essential to maintaining accuracy.¹¹² Use of the MAT is restricted to a few reference laboratories.

A serologically confirmed case of leptospirosis is defined by a fourfold rise in MAT titer to one or more serovars between acute-phase and convalescent serum specimens run in parallel.¹¹³ A single titer of at least 1:800 in the presence of compatible symptoms is strong evidence of recent or current infection.¹¹⁴ Suggestive evidence for recent or current infection includes a single titer of at least 1:200 obtained after the onset of symptoms.¹¹⁵ Delayed seroconversions are common, with up to 10% of patients failing to seroconvert within 30 days of the clinical onset. Cross-reactive antibodies may be associated with syphilis, relapsing fever, Lyme disease, viral hepatitis, human immunodeficiency virus infection, legionellosis, and autoimmune diseases.¹¹⁶

The interpretation of the MAT is complicated by cross-reaction between different serogroups, especially in acute-phase samples.⁵ Cross-reactivity in acute samples is attributable to IgM antibodies, which may persist for several years.¹¹⁷ The MAT is a serogroup-specific assay and should not be used to infer the identity of the infecting serovar.¹⁶ However, knowledge of the presumptive serogroup may be of epidemiologic value in determining potential exposures to animal reservoirs.

Diagnostic application of the MAT is limited by the relatively low sensitivity when acute serum samples are tested.¹¹⁸ Other agglutination assays that detect total immunoglobulins, such as the indirect hemagglutination assay, suffer from similarly low sensitivities on acute specimens but have high case sensitivities when acute and convalescent specimens are tested.¹¹⁹ IgM antibodies are detectable after about the fifth day of illness, and IgM-detection assays are available in several formats.^{119–122} Use of these assays as screening tests offers the potential to enhance the diagnostic capacity of many laboratories, particularly in developing countries, where most cases of leptospirosis occur.¹²³ The modified Faine criteria scoring system, which combines serologic results with clinical history and epidemiologic factors, may be useful in some settings.¹²⁴ The persistently unsatisfactory state of diagnostics, especially in early disease, is summarized in a recent study and a recent review, which highlight the general problem of lacking a gold-standard diagnostic assay by which to assess the utility of tests that are affordable and require less extensive laboratory capacity.^{125,126}

THERAPY

Antibiotic therapy should be initiated as early in the course of the disease as suspicion allows. There have been few randomized or placebo-controlled trials,^{127,128,129,130} and these have produced conflicting results.

TABLE 239.4 Antimicrobial Agents Recommended for Treatment and Chemoprophylaxis of Leptospirosis

INDICATION	COMPOUND	DOSAGE
Chemoprophylaxis	Doxycycline	200 mg PO orally once per week
Treatment of mild leptospirosis	Doxycycline Ampicillin Amoxicillin	100 mg bid PO 500–750 mg q6h PO 500 mg q6h PO
Treatment of moderate-to-severe leptospirosis	Penicillin Ceftriaxone Ampicillin	1.5 MU IV q6h 1 g IV q24h 0.5–1 g IV q6h

bid, Twice daily; IV, intravenously; MU, million units; PO, orally.

Therapeutic benefits of antibiotics may be difficult to demonstrate in populations where patients present for medical care with late or severe disease, or both. Nevertheless, severe disease is usually treated with intravenous penicillin and mild disease with oral doxycycline (Table 239.4). Once-daily ceftriaxone has been shown to be as effective as penicillin.¹³¹ Jarisch-Herxheimer reactions have been reported in patients treated with penicillin.¹³² Patients receiving penicillin should be monitored because of the increased morbidity and mortality of such reactions.

Supportive therapy is essential for hospitalized patients. Patients with early renal disease with high-output renal dysfunction and hypokalemia should receive aggressive volume repletion and potassium supplementation to avoid severe dehydration and acute tubular necrosis.⁶⁸ In patients who progress to oliguric renal failure, rapid initiation of daily hemodialysis reduces mortality, particularly in critically ill patients, and is typically required on only a short-term basis. Renal dysfunction caused by leptospirosis is typically completely reversible.^{133,134} Patients requiring intubation for SPHS have decreased pulmonary compliance and should be managed as cases of ARDS. Protective ventilation strategies involving low tidal volumes (<6 mL/kg) to avoid alveolar injury due to high ventilation pressures have been shown to dramatically improve survival rates.¹³⁵

PREVENTION

Prevention of leptospirosis may be achieved by avoidance of high-risk exposures, adoption of protective measures, immunization, and use of chemoprophylaxis, in varying combinations depending on the environmental circumstances and the degree of human activity.

High-risk exposures include immersion in fresh water, as in swimming, and contact with animals and their body fluids.² Removal of leptospires from the environment is impractical, but reducing direct contact with potentially infected animals and indirect contact with urine-contaminated soil and water remains the most effective preventive strategy available. Consistent application of rodent control measures is important in limiting the extent of contamination. Appropriate protective measures depend on the activity but include wearing boots, goggles, overalls, and rubber gloves. In tropical environments walking barefoot is a common risk factor.¹³⁶ In an urban area of Thailand with high environmental contamination and unavoidable exposure to flood waters, a single dose of doxycycline, 200 g, was considered to have protective efficacy, particularly for flood victims with lacerations, in whom efficacy was estimated to have been 92%.¹³⁷

Immunization of agricultural and companion animals with killed vaccines is widely practiced both to prevent disease and to reduce human exposure. Vaccination of animals at an early age before exposure is crucial in achieving optimum reduction of shedding.¹³⁸ Periodic (usually annual) boosters are required to maintain immunity.¹³⁹ Current bovine and porcine vaccines used in the United States contain serovars Icterohaemorrhagiae, Canicola, Grippotyphosa, Pomona, and Hardjo, whereas canine vaccines contain all except serovar Hardjo. New vaccines for bovine use stimulate a type 1 cell-mediated immune response against serovar Hardjo^{140,141} and appear to protect against renal colonization and urinary shedding.¹⁴²

Human immunization is not widely practiced. A vaccine containing serovar Icterohaemorrhagiae is available in France for workers in high-risk occupations,¹⁴³ and a vaccine has been developed for human use in Cuba.¹⁴⁴ Immunization has been more widely used in Asian countries to prevent large-scale epidemics in agricultural laborers.

For individuals who will be unavoidably exposed to leptospires in endemic environments, chemoprophylaxis is recommended (see Table 239.4). Weekly doxycycline (200 mg) has been shown to be effective in military personnel without previous exposure who underwent jungle training.¹⁴⁵ The use of doxycycline prophylaxis after excess rainfall in local populations in endemic areas has been studied.^{146,147} Symptomatic disease was significantly reduced in one study, but serologic evidence of infection was found equally in subjects and controls.¹⁴⁶ Limitations of doxycycline are its photosensitivity, high frequency of gastrointestinal side effects, dietary calcium restrictions, and contraindication in pregnant women and children. In vitro susceptibility of leptospires to azithromycin¹⁴⁸ and its longer serum half-life suggest that this agent would be a reasonable alternative to doxycycline; however, clinical trials are necessary to validate this approach.

Key References

The complete reference list is available online at Expert Consult.

- Levett PN. Leptospirosis. *Clin Microbiol Rev*. 2001;14:296–326.
- Edwards GA, Domm BM. Human leptospirosis. *Medicine (Baltimore)*. 1960;39:117–156.
- Feigin RD, Anderson DC. Human leptospirosis. *CRC Crit Rev Clin Lab Sci*. 1975;5:413–467.
- Levett PN. *Leptospira*. In: Versalovic J, et al, eds. *Manual of Clinical Microbiology*. 10th ed. Washington, DC: American Society for Microbiology Press; 2011:916–923.
- Levett PN. Usefulness of serologic analysis as a predictor of the infecting serovar in patients with severe leptospirosis. *Clin Infect Dis*. 2003;36:447–452.
- Ren SX, Fu G, Jiang XG, et al. Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole-genome sequencing. *Nature*. 2003;422:888–893.
- Nascimento AL, Ko AI, Martins EA, et al. Comparative genomics of two *Leptospira interrogans* serovars reveals novel insights into physiology and pathogenesis. *J Bacteriol*. 2004;186:2164–2172.
- Bulach DM, Zuerne RL, Wilson P, et al. Genome reduction in *Leptospira borgpetersenii* reflects limited transmission potential. *Proc Natl Acad Sci USA*. 2006;103:14560–14565.
- Trejevo RT, Rigau-Perez JG, Ashford DA, et al. Epidemic leptospirosis associated with pulmonary hemorrhage-Nicaragua, 1995. *J Infect Dis*. 1998;178:1457–1463.
- Ko AI, Galvao Reese M, Ribeiro Durado CM, et al. Urban epidemic of severe leptospirosis in Brazil. *Lancet*. 1999;354:820–825.
- Costa F, Hagan JE, Calcagno J, et al. Global morbidity and mortality of leptospirosis: a systematic review. *PLoS Negl Trop Dis*. 2015;9:e0003898.
- Torgerson PR, Hagan JE, Costa F, et al. Global burden of leptospirosis: estimated in terms of disability adjusted life years. *PLoS Negl Trop Dis*. 2015;9:e0004122.
- Grambusch D, Hoedebecke KL. Unforeseen risk: leptospirosis and the U.S. Special operations community. *J Spec Oper Med*. 2012;12:36–42.
- Morgan J, Bornstein SL, Karpati AM, et al. Outbreak of leptospirosis among triathlon participants and community residents in Springfield, Illinois, 1998. *Clin Infect Dis*. 2002;34:1593–1599.
- Sejvar J, Bancroft E, Winthrop K, et al. Leptospirosis in “Eco-Challenge” athletes, Malaysian borneo, 2000. *Emerg Infect Dis*. 2003;9:702–707.
- Agampodi SB, Matthias MA, Moreno AC, et al. Utility of quantitative polymerase chain reaction in leptospirosis diagnosis: association of level of leptospiemia and clinical manifestations in Sri Lanka. *Clin Infect Dis*. 2012;54:1249–1255.
- Areal VM. The pathologic anatomy and pathogenesis of fatal human leptospirosis (Weil’s disease). *Am J Pathol*. 1962;40:393–423.
- Kasheghe K, Hu W, Ojcius DM, et al. Identification of collagenase as a critical virulence factor for invasiveness and transmission of pathogenic *Leptospira* species. *J Infect Dis*. 2014;209:1105–1115.
- Abdulkader RC, Daher EF, Camargo ED, et al. Leptospirosis severity may be associated with the intensity of humoral immune response. *Rev Inst Med Trop São Paulo*. 2002;44:79–83.
- Ingappa J, Kuffner T, Tappero J, et al. HLA-DQ6 and ingestion of contaminated water: possible gene-environment interaction in an outbreak of leptospirosis. *Genes Immun*. 2004;5:197–202.
- Chu KM, Rathinam R, Namperumalsamy P, et al. Identification of *Leptospira* species in the pathogenesis of uveitis and determination of clinical ocular characteristics in south India. *J Infect Dis*. 1998;177:1314–1321.
- Haake DA. Spirochaetal lipoproteins and pathogenesis. *Microbiology*. 2000;146:1491–1504.
- Haake DA, Suchard MA, Kelley MM, et al. Molecular evolution and mosaicism of leptospiral outer membrane proteins involves horizontal DNA transfer. *J Bacteriol*. 2004;186:2818–2828.
- Choy HA, Kelley MM, Chen TL, et al. Physiological osmotic induction of *Leptospira interrogans* adhesion: LigA and LigB bind extracellular matrix proteins and fibrinogen. *Infect Immun*. 2007;75:2441–2450.
- Croda J, Ramos JG, Matsunaga J, et al. *Leptospira* immunoglobulin-like proteins as a serodiagnostic marker for acute leptospirosis. *J Clin Microbiol*. 2007;45:1528–1534.
- Fraga TR, Courrol Ddos S, Castiblanco-Valencia MM, et al. Immune evasion by pathogenic *Leptospira* strains: the secretion of proteases that directly cleave complement proteins. *J Infect Dis*. 2014;209:876–886.