



Guideline

American Society of Transplantation and Cellular Therapy Series, 1: Enterobacterales Infection Prevention and Management after Hematopoietic Cell Transplantation



Michael J. Satlin^{1,*}, Scott J. Weissman², Paul A. Carpenter^{2,3}, Susan K. Seo⁴, Samuel A. Shelburne⁵

¹ Division of Infectious Diseases, Department of Medicine, Weill Cornell Medicine, New York, New York

² Division of Infectious Diseases, Seattle Children's Hospital, Department of Pediatrics, University of Washington School of Medicine, Seattle, Washington

³ Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington

⁴ Infectious Diseases Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York

⁵ Department of Infectious Diseases, MD Anderson Cancer Center, Houston, Texas

Article history:

Received 6 October 2020

Accepted 7 October 2020

Key Words:

Enterobacterales
Guidelines
Epidemiology
Treatment
Prevention
Hematopoietic
Cell transplant

ABSTRACT

The Practice Guidelines Committee of the American Society of Transplantation and Cellular Therapy partnered with its Transplant Infectious Disease Special Interest Group to update its 2009 compendium-style infectious diseases guidelines for hematopoietic cell transplantation (HCT). A completely fresh approach was taken, with the goal of better serving clinical providers by publishing each stand-alone topic in the infectious diseases series in a concise format of frequently asked questions (FAQs), tables, and figures [1]. Adult and pediatric infectious diseases and HCT content experts developed and then answered FAQs, and then finalized topics with harmonized recommendations that were made by assigning a strength of recommendation ranging from A to E paired with a level of supporting evidence graded I to III. The first topic in the series focuses on potentially life-threatening infections in HCT caused by Enterobacterales, relevant infection risk factors, and practical considerations regarding prevention and treatment of these infections in the setting of emerging multidrug resistance.

© 2020 The American Society for Transplantation and Cellular Therapy. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

INTRODUCTION

Hematopoietic cell transplantation (HCT) recipients are at increased risk for potentially life-threatening bloodstream infections caused by Enterobacterales. This guideline is in the form of frequently asked questions (FAQs) and focuses on relevant risk factors for infection and multidrug resistance that are prevalent in many geographic regions, as well as on strategies to prevent and treat these common gram-negative infections.

EPIDEMIOLOGY

FAQ1: Which organisms compose the order Enterobacterales, and which are the most common pathogens after HCT?

Enterobacterales comprises multiple bacterial families, including the family Enterobacteriaceae that harbors the genera *Escherichia*, *Klebsiella*, and *Enterobacter* [2]. *Escherichia coli* and *Klebsiella pneumoniae* are the most common causes of post-transplantation gram-negative bacteremia [3–5].

Financial disclosure: See Acknowledgments on page 111.

*Correspondence and reprint requests: Michael Satlin, MD, Weill Cornell Medicine, 1300 York Avenue, A-421, New York, NY 10021.

E-mail address: mjs9012@med.cornell.edu (M.J. Satlin).

FAQ2: What are risk factors for developing an Enterobacterales infection after HCT?

Enterobacterales bacteremias most commonly occur before neutrophil engraftment. Risk factors include older age, receipt of a cord blood or mismatched donor graft, prolonged neutropenia, gastrointestinal (GI) mucositis, and lack of antibacterial prophylaxis [6–8]. Postengraftment, GI graft-versus-host disease (GVHD) is another risk factor due to bacterial translocation across a damaged colon [9,10].

FAQ3: What are the most problematic types of multidrug-resistant (MDR) Enterobacterales, and how frequently do they occur during HCT? (Table 1)

These can be dichotomized into third-generation cephalosporin-resistant/carbapenem-susceptible Enterobacterales, which include extended-spectrum β -lactamase (ESBL)-producing and AmpC-producing bacteria, as well as the category of carbapenem-resistant Enterobacterales (CRE). The former are most commonly *E coli* and *K pneumoniae* that have ESBL enzymes that hydrolyze penicillins, extended-spectrum cephalosporins, and aztreonam, but not carbapenems, thereby conferring resistance to many first-line agents for fever and neutropenia [11,12]. Depending on geographic region, 10% to

Table 1

ESBL-Producing, AmpC-Producing, and Carbapenem-Resistant Enterobacterales: Epidemiology, Resistance Phenotypes, and Treatment Considerations

Parameter	Third-Generation Cephalosporin-Resistant/Carbapenem-Susceptible		
	ESBLs	AmpCs	CREs
Common species	<i>Escherichia coli</i> <i>Klebsiella pneumoniae</i> <i>Proteus mirabilis</i>	<i>Enterobacter</i> spp* <i>Klebsiella aerogenes</i> <i>Serratia marcescens</i> <i>Proteus vulgaris</i> <i>Morganella morganii</i> <i>Providencia</i> spp <i>Citrobacter freundii</i>	<i>Klebsiella pneumoniae</i> <i>Enterobacter</i> spp <i>Escherichia coli</i>
Common phenotypic susceptibility profile [†]	Ceftriaxone-resistant Carbapenem- and cefoxitin-susceptible	Ampicillin-, cefazolin-, and cefoxitin-resistant [‡] Carbapenem- and cefepime-susceptible	Carbapenem-, cefepime-, and piperacillin-tazobactam-resistant • KPC: susceptible to new BL-BLIs [§] • MBL (eg, NDM): resistant to new BL-BLIs • OXA-48: susceptible to ceftazidime-avibactam, but often resistant to other new BL-BLIs Carbapenemase production (eg, KPC, NDM, OXA-48) or ESBL or AmpC production combined with loss or dysfunction of outer membrane porins
Resistance mechanisms	Production of plasmid-encoded CTX-M and mutated TEM and SHV β -lactamases	Production of chromosomally encoded β -lactamases that can be derepressed on exposure to certain β -lactam agents	
Preferred therapies	Carbapenems (A-I)	First-line: Carbapenems (B-II) Alternative: Cefepime (B-II)	• KPC: New BL-BLIs (B-II) • MBL: Ceftazidime-avibactam + aztreonam or polymyxin or aminoglycoside-based combination therapy • OXA-48: Ceftazidime-avibactam (B-III) • Noncarbapenemase producers: case-by-case depending on susceptibility testing results

* These organisms are frequently referred to as the “SPICE” organisms.

† Susceptibilities of ESBL- and AmpC-producing Enterobacterales to piperacillin-tazobactam and ceftazidime are variable.

‡ Hyperproduction of AmpC β -lactamase typically also leads to resistance to third-generation cephalosporins and aztreonam.

§ Newly approved BL-BLIs with activity against KPC-producing organisms include ceftazidime-avibactam, meropenem-vaborbactam, and imipenem-relebactam.

|| Plasmid-encoded AmpC β -lactamases have also been identified, but are less common than chromosomally-encoded enzymes.

40% of *E. coli* and *K. pneumoniae* bloodstream isolates in HCT recipients are ESBL producers [13].

Other Enterobacterales harbor *ampC* genes not normally expressed at sufficient levels to confer resistance to third-generation cephalosporins. However, after exposure to certain β -lactam agents, including third-generation cephalosporins, *ampC* gene expression is induced, leading to sufficient AmpC β -lactamase production to hydrolyze third-generation cephalosporins and aztreonam [14]. Thus, isolates that initially appear susceptible to third-generation cephalosporins can quickly develop resistance on therapy [15]. *Enterobacter* spp and *Klebsiella* (*Enterobacter*) *aerogenes* most commonly harbor these genes, followed by other “SPICE” organisms (*Serratia marcescens*, *Proteus vulgaris*, *Morganella morganii*, *Providencia stuartii*, and *Citrobacter freundii*).

CRE bacteremia after HCT is being increasingly reported, particularly in certain endemic areas outside of the United States [13,16,17]. It occurs from either bacterial carbapenemases alone or ESBL/AmpC β -lactamases combined with changes in outer membrane porins that limit antimicrobial access to penicillin-binding proteins. Carbapenemase-producing Enterobacterales are often resistant to all first-line agents for fever and neutropenia, with *K. pneumoniae* carbapenemase (KPC) the most common [18]. Other major carbapenemases include OXA-48 β -lactamases and New Delhi metallo- β -lactamases (NDM).

FAQ4: What are risk factors for infection due to an Enterobacterales that is resistant to standard agents for fever and neutropenia in HCT recipients?

Risk factors include residence in a location with a high prevalence of MDR Enterobacterales, colonization with MDR

Enterobacterales, prolonged hospitalization, receipt of allogeneic HCT, and previous antibacterial therapy [5,19]

FAQ5: Are clinical outcomes attributed to MDR Enterobacterales infections worse than those due to more susceptible pathogens?

Studies of neutropenic patients have generally demonstrated increased mortality after bacteremias caused by ESBL-producing Enterobacterales (ESBL-E) versus non-ESBL-E, in part because ESBL-E-infected patients frequently receive inadequate initial therapy [20–22]. CRE bacteremia during HCT has devastating consequences, with mortality >50% [13,16,17]. No firm conclusions can be established about the outcomes of bacteremias caused by AmpC-hyperproducing Enterobacterales in this population.

TREATMENT

FAQ6: Which antimicrobial agents are best to treat infections caused by ESBL-E and AmpC-producing Enterobacterales during HCT? (Table 1)

Carbapenems are recommended for treating invasive ESBL-E infections despite frequent in vitro susceptibility of ESBL-E to piperacillin-tazobactam (A-I). This is because mortality was increased by 3-fold when piperacillin-tazobactam was compared with a carbapenem in a randomized trial of ESBL-E bacteremia therapy in the general population [23]. The majority of ESBL-E are resistant to cefepime (fourth-generation cephalosporin) and even patients with a cefepime-susceptible ESBL-E infection have worse outcomes when treated with cefepime compared with carbapenems [24,25]. The newer agents ceftolozane-tazobactam and ceftazidime-avibactam are active against ESBL-E in vitro [24]. Although limited data suggest that

they have comparable clinical efficacy to carbapenems for ESBL-E infections in the general patient population [24,26–28], clinical data on the efficacy of these new agents in HCT recipients are needed.

Carbapenem therapy is also recommended for invasive *Enterobacter* spp and *K aerogenes* infections, because carbapenems are not readily hydrolyzed by their AmpC β -lactamases (B-II) [14]. Third-generation cephalosporins (eg, ceftriaxone) should not be used to treat invasive *Enterobacter* infections, because resistance can develop on therapy (D-II) [15]. However, increasing observational data suggest that cefepime may be just as effective as carbapenems and thus can serve as an alternative agent for these infections (B-II) [14,29–31]. Depending on susceptibility results, fluoroquinolones and trimethoprim-sulfamethoxazole are potential oral step-down therapies after clinical improvement for both ESBL-E and AmpC-producing Enterobacterales infections (B-III) [32]. The optimal treatment of infections due to SPICE organisms other than *Enterobacter* (eg, *S marcescens*) is less clear, and treatment decisions can be based on in vitro susceptibility results.

FAQ7: Which antimicrobial agents are best for treating CRE infections during HCT?

This depends on the mechanism of carbapenem resistance (Table 1). Infections caused by KPC-producing CRE are best treated with novel β -lactam/ β -lactamase inhibitors (BL-BLIs), such as ceftazidime-avibactam, meropenem-vaborbactam, or imipenem-relebactam, because of improved clinical outcomes and less toxicity compared with polymyxin-based regimens (B-II) [33–35]. CRE with OXA-48-like carbapenemases are frequently resistant to meropenem-vaborbactam and imipenem-relebactam but are typically susceptible to ceftazidime-avibactam, which is recommended for these infections (B-III) [36–38]. Metallo- β -lactamase (MBL)-producing CRE (eg, NDM) are typically resistant to new BL-BLIs, and although optimal therapies for these infections remain undefined, potential options include ceftazidime-avibactam plus aztreonam and polymyxin- or aminoglycoside-based regimens. Other novel agents, such as eravacycline, plazomicin, or cefiderocol, also may be considered [39,40]. Consultation with an infectious diseases expert is recommended for the treatment of CRE infections [41].

FAQ8: For how long should Enterobacterales bacteremias be treated after HCT?

Although data on outcomes by treatment duration are limited, 10 to 14 days of appropriate antimicrobial therapy is generally recommended (C-III) [42]. One randomized trial of patients with uncomplicated gram-negative bacteremia demonstrated that 7 days was noninferior to 14 days of therapy, but immunocompromised patients were excluded from the study [43]. In uncomplicated central venous catheter (CVC)-related Enterobacterales bacteremia, a course as short as 7 days may be reasonable if management includes CVC removal, but other factors may warrant up to 14 days of therapy (C-III) [44].

HCT recipients who receive 14 days of therapy and remain neutropenic but without signs or symptoms of ongoing infection can often resume antibacterial prophylaxis or have antibacterial therapy discontinued [12]. This strategy is supported by a randomized trial that showed discontinuing antibacterial therapy is safe during neutropenia if patients are afebrile without signs or symptoms of infection [45]. Although we recommend initial i.v. therapy, those who improve clinically and do not have GI pathology that would impair systemic absorption

can be transitioned to highly bioavailable oral therapies, such as fluoroquinolones or trimethoprim-sulfamethoxazole, depending on relevant susceptibility profiles (B-III). Observational study results showing therapeutic equivalence for i.v.-to-oral step-down therapy versus continued i.v. therapy support this approach [32].

FAQ9: Should a CVC be removed for Enterobacterales bacteremia during HCT?

It depends. Enterobacterales bacteremia in HCT is more commonly due to gut translocation during chemotherapy-induced neutropenia or GI GVHD than from direct infection of a CVC [19,46–49]. In this setting of mucosal barrier injury, CVC removal is not associated with improved outcomes and is not routinely recommended (D-III) [46].

Conversely, if the Enterobacterales bacteremia develops when there is no mucosal barrier injury and no clear alternate source of infection, a central line-associated bloodstream infection (CLABSI) is more likely. CVC removal is associated with improved outcomes in this setting and should be considered (B-III) [44].

PREVENTION/EMPIRICAL THERAPY

FAQ10: Does antibacterial prophylaxis during neutropenia decrease the risk of Enterobacterales infections during HCT?

Yes in adults, but not clearly in children. Levofloxacin should be considered for adults during neutropenia after HCT to lower the risk of Enterobacterales bacteremia (B-I) [50]. A randomized, placebo-controlled trial in adult patients with cancer who were expected to have >7 days of neutropenia, of whom one-half were undergoing autologous HCT (allogeneic HCT recipients were excluded), demonstrated that levofloxacin prophylaxis significantly decreased the risk of gram-negative bacteremia [51]. Observational studies, primarily in autologous HCT recipients, reached similar conclusions [52–56]. A recent randomized trial in children undergoing autologous or allogeneic HCT showed a trend toward reduction in all-cause BSI with levofloxacin prophylaxis, but was not powered or designed to specifically assess Enterobacterales BSIs [57].

The degree to which increasing fluoroquinolone resistance among the most common Enterobacterales decreases the effectiveness of fluoroquinolone prophylaxis is unclear [58]. The decision whether to routinely administer levofloxacin prophylaxis after HCT should carefully weigh the benefit of potentially lowering the risk of bacteremia and fever and neutropenia against the risks of increasing antimicrobial resistance and levofloxacin-related adverse effects [59,60]. Levofloxacin prophylaxis may not only increase the risk of fluoroquinolone resistance among breakthrough Enterobacterales infections, but because of co-occurrence of fluoroquinolone and β -lactam resistance in prominent strains, may also increase the risk of ESBL-E bacteremia [19,55]. Thus, careful monitoring for increased rates of ESBL-E infections is warranted when using fluoroquinolone prophylaxis.

FAQ11: Should screening be used to identify HCT recipients colonized with MDR Enterobacterales to guide initial empirical therapy?

No definitive recommendations can be provided due to variable colonization rates. One multicenter study demonstrated that in HCT recipients colonized with carbapenem-resistant *K pneumoniae* (CRKp), rates of post-transplantation CRKp infection were 26% in the autologous setting and 39% in the allogeneic setting [16]. Thus, in transplantation centers where CRE

are prominent pathogens, screening for CRE colonization may have a role in guiding empirical therapy during post-transplantation neutropenia. When CRE are uncommon, limiting screening to patients referred from CRE-endemic areas is another reasonable strategy.

At one center where levofloxacin prophylaxis was administered, one-third of ESBL-E-colonized patients developed ESBL-E bacteremia during post-transplantation neutropenia [19]. Thus, screening for ESBL-E could identify HCT recipients who should be treated empirically with a carbapenem for fever and neutropenia. By contrast, other studies in hematologic malignancies in which antibacterial prophylaxis was not routinely used have reported lower rates of ESBL-E bacteremia in colonized patients [61–63].

FAQ12: Instead of a single approach for prophylaxis and empirical therapy, can an individualized approach be pursued during HCT?

One approach to tailoring prophylaxis and empirical antibiotic therapy in HCT recipients is to use an individualized antibiotic plan (IAP) [64]. Before transplantation, patients would undergo rectal screening for colonization with MDR bacteria. Patients not colonized by MDR bacteria would be assigned a standard 3-tiered IAP to guide their antibiotic management from conditioning through engraftment (eg, levofloxacin for prophylaxis; ceftazidime for fever, stable; and ceftazidime + vancomycin + gentamicin for fever, unstable). Patients colonized by MDR bacteria would be assigned a modified IAP regimen by an algorithmic approach according to the susceptibility phenotype of the isolate. The IAP would be entered into the electronic medical record before transplantation, readily visible and accessible to all providers, updated in the event of new microbiologic data, and inactivated at 100 days post-transplantation.

FAQ13: Should HCT recipients infected or colonized with MDR Enterobacterales be placed on contact precautions?

Contact precautions with gown and gloves are recommended in patients infected or colonized with CRE [65] and should be considered for patients with ESBL-E (B-III). The rationale for this is that contact precautions may decrease the risk of transmission of these organisms on a transplantation unit, although data do not consistently support efficacy in preventing ESBL-E transmission [66,67].

FAQ14: Are there interventions to mitigate the risk of MDR Enterobacterales infection in HCT recipients who are colonized with these organisms?

Using oral antibiotics to decolonize HCT recipients with ESBL-E or CRE gut colonization is not routinely recommended (D-III). Some randomized clinical trials have demonstrated marginal effectiveness in temporarily suppressing gut colonization with these organisms, but benefits are transient and major concerns include emergence of resistance to agents used for decolonization and negative effects on the gut microbiome [68–72].

Fecal microbiota transplantation (FMT) is another potential strategy to suppress or eradicate ESBL-E or CRE gut colonization. However, additional data are needed to evaluate the merits of FMT for this indication, particularly in immunocompromised

HCT recipients, given the concern for transmission of infectious agents [73].

ADDITIONAL CONSIDERATIONS

FAQ15: Is the epidemiology of HCT infections caused by Enterobacterales different in children?

There are some differences. Risk factors for Enterobacterales infections in children include allogeneic HCT or myeloablative conditioning, prolonged neutropenia, and acute GI GVHD [74–76]. *E coli* was the most common cause of CLABSI reported from adult and pediatric oncology centers [77,78], but whereas *Klebsiella* spp were the second most common cause of CLABSI in pediatric oncology units, they were the fourth most common cause in adult oncology units.

Rates of resistance to extended-spectrum cephalosporins and carbapenems among CLABSI isolates of *E coli*, *Klebsiella* spp, and *Enterobacter* spp were similar between adult and pediatric oncology patients, but rates of fluoroquinolone resistance were lower among pediatric *E coli* (38%) than adult *E coli* (65%) [77,79]. In a study conducted primarily in Europe, gram-negative bloodstream isolates were less frequently resistant to anti-pseudomonal BL-BLIs and fluoroquinolones among pediatric allogeneic HCT recipients compared with adult allogeneic HCT recipients [5].

FUTURE DIRECTIONS

FAQ16: How can rapid diagnostics be incorporated into treating Enterobacterales bacteremias during HCT?

Multiplexed PCR/microarray-based platforms can diagnose the type of gram-negative infection and presence of selected resistance genes within 2 hours of blood culture bottles signaling positive growth [79,80]. When combined with antimicrobial stewardship interventions, the use of these assays optimizes pathogen-directed antimicrobial therapy and may decrease mortality [79,81]. These assays also rapidly detect certain carbapenemases, which can guide treatment decisions given that therapy choices for carbapenemase-producing infections depends on carbapenemase type (Table 1).

A limitation of these multiplexed PCR/microarray-based platforms is that they detect only a small number of genetic resistance determinants. In contrast, a new rapid phenotypic susceptibility testing platform identifies the organism and more comprehensive antimicrobial susceptibility results within 7 hours of blood cultures signaling positive [82]. An additional novel diagnostic test combines magnetic resonance and PCR-based amplification to detect 5 common bloodstream pathogens, including *E coli* and *K pneumoniae*, directly from whole blood within 6 hours of blood culture collection [83]. However, this test does not detect Enterobacterales species other than *E coli* and *K pneumoniae* and does not provide antimicrobial susceptibility information. Additional data are needed to better understand how to leverage these assays to improve outcomes of HCT recipients with Enterobacterales bacteremia.

ACKNOWLEDGMENTS

Financial disclosure: There is no funding to report.

Conflict of interest statement: MJS reports receiving consulting fees from Shionogi and Achaogen and grant support from Allergan, Merck, and BioFire Diagnostics. SJW reports receiving consulting fees and having equity in MicrobiomX. The other authors have no conflicts of interest to report.

APPENDIX: GRADING STRENGTH OF RECOMMENDATION AND LEVEL OF EVIDENCE [1]

Recommendation	Grade
Which antimicrobial agents are best to treat infections caused by ESBL-E and AmpC-producing Enterobacterales during HCT?	
1. Carbapenems are recommended to treat invasive ESBL-E infections despite frequent in vitro susceptibility of ESBL-E to other β -lactam agents [23,25].	A-I
2. Carbapenems are recommended for invasive <i>Enterobacter</i> spp and <i>Klebsiella</i> (<i>Enterobacter</i>) <i>aerogenes</i> infections because carbapenems are not hydrolyzed by their AmpC β -lactamases [14].	B-II
3. Third-generation cephalosporins (eg, ceftriaxone) should not be used to treat invasive <i>Enterobacter</i> infections, because resistance can develop on therapy [15].	D-II
4. Cefepime may be an alternative option for invasive <i>Enterobacter</i> infections because of its stability against AmpC β -lactamases [14,29–31].	B-II
5. Depending on susceptibility results, fluoroquinolones and trimethoprim-sulfamethoxazole are potential oral step-down therapies for ESBL-E and AmpC-producing Enterobacterales after clinical improvement [32].	B-III
Which antimicrobial agents are best to treat CRE infections during HCT?	
6. Infections caused by KPC-producing CRE are best treated with novel BL-BLIs, such as ceftazidime-avibactam, meropenem-vaborbactam, or imipenem-relebactam [33–35].	B-II
7. Ceftazidime-avibactam is recommended for the treatment of infections caused by CRE with OXA-48-like carbapenemases, which are frequently resistant to meropenem-vaborbactam and imipenem-relebactam [36–38].	B-III
For how long should Enterobacterales bacteremias be treated after HCT?	
8. In the absence of complications, 10 to 14 days of appropriate antimicrobial therapy is generally recommended [42].	C-III
9. In uncomplicated CVC-related Enterobacterales bacteremia, a course as short as 7 days may be reasonable if management includes CVC removal, but other factors may warrant up to 14 days of therapy [44].	C-III
Should a CVC be removed for Enterobacterales bacteremia during HCT?	
10. In the setting of Enterobacterales bacteremia during chemotherapy-induced neutropenia or GI GVHD, CVC removal is not routinely recommended, unless there are clear indications that the CVC is the source of the infection [46].	D-III
11. If Enterobacterales bacteremia develops without chemotherapy-induced neutropenia or GI GVHD, CVC removal should be considered in the absence of an alternate source of infection [44].	B-III
Should antibacterial prophylaxis be administered during neutropenia to decrease the risk of Enterobacterales infections during HCT?	
12. Levofloxacin prophylaxis should be considered for adults during neutropenia after HCT to lower the risk of Enterobacterales bacteremia [50–56].	B-I
Should HCT recipients infected or colonized with MDR Enterobacterales be placed on contact precautions?	
13. Contact precautions with gown and gloves are recommended in patients infected or colonized with CRE and should be considered for patients with ESBL-E [65,66].	B-III
Should oral antibiotics be used to decolonize HCT recipients who are colonized with MDR Enterobacterales?	
14. Using oral antibiotics to decolonize HCT recipients with ESBL-E or CRE gut colonization should not be performed routinely [68–72].	D-III

REFERENCES

- Carpenter PA, Papanicolaou, Chemaly RF, Boeckh M, Savani BN. American Society for Transplantation and Cellular Therapy Infectious Diseases Guidelines: Preface to the Series. *Biol Blood Marrow Transplant*. 2020. [Manuscript under review].
- Adeolu M, Alnajjar S, Naushad S, Gupta RS. Genome-based phylogeny and taxonomy of the “Enterobacterales”: proposal for Enterobacterales ord. nov., divided into the families Enterobacteriaceae, Erwiniaceae fam. nov., Pectobacteriaceae fam. nov., Yersiniaceae fam. nov., Hafniaceae fam. nov., Morganellaceae fam. nov., and Budviciaceae fam. nov. *Int J Syst Evol Microbiol*. 2016;66:5575–5599.
- Seo SK, Xiao K, Huang YT, et al. Impact of peri-transplant vancomycin and fluoroquinolone administration on rates of bacteremia in allogeneic hematopoietic stem cell transplant (HSCT) recipients: a 12-year single institution study. *J Infect*. 2014;69:341–351.
- Miles-Jay A, Butler-Wu S, Rowhani-Rahbar A, Pergam SA. Incidence rate of fluoroquinolone-resistant gram-negative rod bacteremia among allogeneic hematopoietic cell transplantation patients during an era of levofloxacin prophylaxis. *Biol Blood Marrow Transplant*. 2015;21:539–545.
- Averbuch D, Tridello G, Hoek J, et al. Antimicrobial resistance in gram-negative rods causing bacteremia in hematopoietic stem cell transplant recipients: Intercontinental Prospective Study of the Infectious Diseases Working Group Party of the European Bone Marrow Group. *Clin Infect Dis*. 2017;65:1819–1828.
- Blennow O, Ljungman P, Sparrelid E, Mattsson J, Remberger M. Incidence, risk factors, and outcome of bloodstream infections during the pre-engraftment phase in 521 allogeneic hematopoietic stem cell transplantations. *Transpl Infect Dis*. 2014;16:106–114.
- Girmenia C, Bertina A, Piciocchi A, et al. Incidence, risk factors and outcome of pre-engraftment Gram-negative bacteremia after allogeneic and autologous hematopoietic stem cell transplantation: an Italian prospective multicenter survey. *Clin Infect Dis*. 2017;65:1884–1896.
- Yan CH, Wang Y, Mo XD, et al. Incidence, risk factors, microbiology and outcomes of pre-engraftment bloodstream infection after haploidentical hematopoietic stem cell transplantation and comparison with HLA-identical sibling transplantation. *Clin Infect Dis*. 2018;67(suppl_2):S162–S173.
- Mori Y, Yoshimoto G, Nishida R, et al. Gastrointestinal graft-versus-host disease is a risk factor for postengraftment bloodstream infection in allogeneic hematopoietic stem cell transplant recipients. *Biol Blood Marrow Transplant*. 2018;24:2302–2309.
- Petersen J, Lindner C, Hakki M. Incidence and outcomes of bacterial bloodstream infections during acute graft-versus-host disease involving the gastrointestinal tract after hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2019;25:1648–1653.
- Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev*. 2005;18:657–686.
- Freifeld AG, Bow EJ, Sepkowitz KA, et al. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2011;52:e56–e93.
- Satlin MJ, Walsh TJ. Multidrug-resistant Enterobacteriaceae, *Pseudomonas aeruginosa*, and vancomycin-resistant *Enterococcus*: three major threats to hematopoietic stem cell transplant recipients. *Transpl Infect Dis*. 2017;19. <https://doi.org/10.1111/tid.12762>.
- Tamma PD, Doi Y, Bonomo RA, Johnson JK, Simner PJ. Antibacterial Resistance Leadership Group. A primer on AmpC β -lactamases: necessary knowledge for an increasingly multidrug-resistant world. *Clin Infect Dis*. 2019;69:1446–1455.
- Chow JW, Fine MJ, Shales DM, et al. *Enterobacter* bacteremia: clinical features and emergence of antibiotic resistance during therapy. *Ann Intern Med*. 1991;115:585–590.
- Girmenia C, Rossolini GM, Piciocchi A, et al. Infections caused by carbapenem-resistant *Klebsiella pneumoniae* in SCT recipients: a nationwide retrospective study from Italy. *Bone Marrow Transplant*. 2015;50:282–288.
- Pouch SM, Satlin MJ. Carbapenem-resistant Enterobacteriaceae in special populations: solid organ transplant recipients, stem cell transplant recipients, and patients with hematologic malignancies. *Virulence*. 2017;8:391–402.
- Woodworth KR, Walters MS, Weiner LM, et al. Vital signs: Containment of novel multidrug-resistant organisms and resistance mechanisms—United States, 2006–2017. *MMWR Morb Mortal Wkly Rep*. 2018;67:396–401.
- Satlin MJ, Chavda KD, Baker TM, et al. Colonization with levofloxacin-resistant extended-spectrum β -lactamase-producing Enterobacteriaceae

- and risk of bacteremia in hematopoietic stem cell transplant recipients. *Clin Infect Dis*. 2018;67:1720–1728.
20. Gudiol C, Calatayud L, Garcia-Vidal C, et al. Bacteraemia due to extended-spectrum beta-lactamase-producing *Escherichia coli* (ESBL-EC) in cancer patients: clinical features, risk factors, molecular epidemiology and outcome. *J Antimicrob Chemother*. 2010;65:333–341.
 21. Kang CI, Chung DR, Ko KS, Peck KR, Song JH, Korean Network for Study of Infectious Diseases. Risk factors for infection and treatment outcome of extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* bacteremia in patients with hematologic malignancy. *Ann Hematol*. 2012;91:115–121.
 22. Cornejo-Juárez P, Pérez-Jiménez C, Silva-Sánchez J, et al. Molecular analysis and risk factors for *Escherichia coli* producing extended-spectrum β -lactamase bloodstream infection in hematological malignancies. *PLoS One*. 2012;7:e35780.
 23. Harris PNA, Tambyah PA, Lye DC, et al. Effect of piperacillin-tazobactam vs meropenem on 30-day mortality for patients with *E. coli* or *Klebsiella pneumoniae* bloodstream infection and ceftriaxone resistance: a randomized clinical trial. *JAMA*. 2018;320:984–994.
 24. Castanheira M, Doyle TB, Mendes RE, Sader HS. Comparative activities of ceftazidime-avibactam and ceftolozane-tazobactam against Enterobacteriaceae isolates producing extended-spectrum β -lactamases from United States hospitals. *Antimicrob Agents Chemother*. 2019;63. e00160-19.
 25. Lee NY, Lee CC, Huang WH, Tsui KC, Hsueh PR, Ko WC. Cefepime therapy for monomicrobial bacteremia caused by cefepime-susceptible extended-spectrum beta-lactamase-producing Enterobacteriaceae: MIC matters. *Clin Infect Dis*. 2013;56:488–495.
 26. Mendes RE, Castanheira M, Woosley LN, Stone GG, Bradford PA, Flamm RK. Molecular β -lactamase characterization of aerobic Gram-negative pathogens recovered from patients enrolled in the ceftazidime-avibactam phase 3 trials for complicated intra-abdominal infections, with efficacies analyzed against susceptible and resistant subsets. *Antimicrob Agents Chemother*. 2017;61. e02447-16.
 27. Mendes RE, Castanheira M, Woosley LN, Stone GG, Bradford PA, Flamm RK. Characterization of β -lactamase content of ceftazidime-resistant pathogens recovered during the pathogen-directed phase 3 REPRIME trial for ceftazidime-avibactam: correlation of efficacy against β -lactamase producers. *Antimicrob Agents Chemother*. 2019;63. e02655-18.
 28. Kollef MH, Nováček M, Kivistik Ü, et al. Ceftolozane-tazobactam versus meropenem for treatment of nosocomial pneumonia (ASPECT-NP): a randomized, controlled double-blind, phase 3, non-inferiority trial. *Lancet Infect Dis*. 2019;19:1299–1311.
 29. Lee NY, Lee CC, Li CW, et al. Cefepime therapy for monomicrobial *Enterobacter cloacae* bacteremia: unfavorable outcomes in patients infected by cefepime-susceptible dose-dependent isolates. *Antimicrob Agents Chemother*. 2015;59:7558–7563.
 30. McKamey L, Venugopalan V, Cherabuddi K, et al. Assessing antimicrobial stewardship initiatives: clinical evaluation of cefepime or piperacillin/tazobactam in patients with bloodstream infections secondary to AmpC-producing organisms. *Int J Antimicrob Agents*. 2018;52:719–723.
 31. Tan SH, Ng TM, Chew KL, et al. Outcomes of treating AmpC-producing Enterobacteriaceae bacteremia with carbapenems vs. non-carbapenems. *Int J Antimicrob Agents*. 2020;55: 105860.
 32. Tamma PD, Conley AT, Cosgrove SE, et al. Association of 30-day mortality with oral step-down vs continued intravenous therapy in patients hospitalized with Enterobacteriaceae bacteremia. *JAMA Intern Med*. 2019;179:316–323.
 33. van Duin D, Lok JJ, Earley M, et al. Colistin versus ceftazidime-avibactam in the treatment of infections due to carbapenem-resistant Enterobacteriaceae. *Clin Infect Dis*. 2018;66:163–171.
 34. Wunderink RG, Giamarellos-Bourboulis EJ, Rahav G, et al. Effect and safety of meropenem-vaborbactam versus best-available therapy in patients with carbapenem-resistant Enterobacteriaceae infections: the TANGO II randomized clinical trial. *Infect Dis Ther*. 2018;7:439–455.
 35. Motsch J, Murta de Oliveira C, Stus V, et al. RESTORE-IMI 1: A multicenter, randomized, double-blind trial comparing efficacy and safety of imipenem/relebactam vs colistin plus imipenem in patients with imipenem-nonsusceptible bacterial infections. *Clin Infect Dis*. 2020;70:1799–1808.
 36. Castanheira M, Huband MD, Mendes RE, Flamm RK. Meropenem-vaborbactam tested against contemporary Gram-negative isolates collected worldwide during 2014, including carbapenem-resistant, KPC-producing, multidrug-resistant, and extensively drug-resistant Enterobacteriaceae. *Antimicrob Agents Chemother*. 2017;61. e00567-17.
 37. Canver MC, Satlin MJ, Westblade LF, et al. Activity of imipenem-relebactam and comparator agents against genetically characterized isolates of carbapenem-resistant Enterobacteriaceae. *Antimicrob Agents Chemother*. 2019;63. e00672-19.
 38. Sousa A, Pérez-Rodríguez MT, Soto A, et al. Effectiveness of ceftazidime/avibactam as salvage therapy for treatment of infections due to OXA-48 carbapenemase-producing Enterobacteriaceae. *J Antimicrob Chemother*. 2018;73:3170–3175.
 39. Tumbarello M, Losito AR, Giamarellou H. Optimizing therapy in carbapenem-resistant Enterobacteriaceae infections. *Curr Opin Infect Dis*. 2018;31:566–577.
 40. Wu JY, Srinivas P, Pogue JM. Cefiderocol: a novel agent for the management of multidrug-resistant Gram-negative organisms. *Infect Dis Ther*. 2020;9:17–40.
 41. Burnham JP, Olsen MA, Stawley D, Kwon JH, Babcock HM, Kollef MH. Infectious diseases consultation reduces 30-day and 1-year all-cause mortality for multidrug-resistant organism infections. *Open Forum Infect Dis*. 2018;5. ofy026.
 42. Giannella M, Pascale R, Toschi A, et al. Treatment duration for *Escherichia coli* bloodstream infection and outcomes: retrospective single-centre study. *Clin Microbiol Infect*. 2018;24:1077–1083.
 43. Yahav D, Franceschini E, Koppel F, et al. Seven versus 14 days of antibiotic therapy for uncomplicated Gram-negative bacteremia: a noninferiority randomized controlled trial. *Clin Infect Dis*. 2019;69:1091–1098.
 44. Mermel LA, Allon M, Bouza E, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;49:1–45.
 45. Aguilar-Guisado M, Espigado I, Martín-Peña A, et al. Optimisation of empirical antimicrobial therapy in patients with hematological malignancies and febrile neutropenia (How Long study): an open-label, randomized, controlled phase 4 trial. *Lancet Hematol*. 2017;4:e573–e583.
 46. Fares J, Khalil M, Chaftari AM, Jiang Y, Hachem RY, Raad II. Impact of catheter management on the clinical outcome in adult cancer patients with Gram-negative bacteremia. *Open Forum Infect Dis*. 2019;6(suppl 2):S419.
 47. Taur Y, Xavier JB, Lipuma L, et al. Intestinal domination and the risk of bacteremia in patients undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis*. 2012;55:905–914.
 48. Chaftari AM, Jordan M, Hachem R, et al. A clinical practical approach to the surveillance definition of central line-associated bloodstream infection in cancer patients with mucosal barrier injury. *Am J Infect Control*. 2016;44:931–934.
 49. Kelly MS, Ward DV, Severn CJ, et al. Gut colonization preceding mucosal barrier injury bloodstream infection in pediatric hematopoietic stem cell transplant recipients. *Biol Blood Marrow Transplant*. 2019;25:2274–2280.
 50. Taplitz RA, Kennedy EB, Bow EJ, et al. Antimicrobial prophylaxis for adult patients with cancer-related immunosuppression: ASCO and IDSA clinical practice guideline update. *J Clin Oncol*. 2018;36:3043–3054.
 51. Bucaneve G, Micozzi A, Menichetti F, et al. Levofloxacin to prevent bacterial infection in patients with cancer and neutropenia. *N Engl J Med*. 2005;353:977–987.
 52. Kern WV, Klose K, Jellen-Ritter AS, et al. Fluoroquinolone resistance of *Escherichia coli* at a cancer center: epidemiologic evolution and effects of discontinuing prophylactic fluoroquinolone use in neutropenic patients with leukemia. *Eur J Clin Microbiol Infect Dis*. 2005;24:111–118.
 53. Satlin MJ, Vardhana S, Soave R, et al. Impact of prophylactic levofloxacin on rates of bloodstream infection and fever in neutropenic patients with multiple myeloma undergoing autologous hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2015;21:1808–1814.
 54. Yeshurun M, Vaxman I, Shargian L, et al. Antibacterial prophylaxis with ciprofloxacin for patients with multiple myeloma and lymphoma undergoing autologous hematopoietic cell transplantation: a quasi-experimental single-centre before-after study. *Clin Microbiol Infect*. 2018;24:749–754.
 55. Kern WV, Weber S, Dettenkofer M, et al. Impact of fluoroquinolone prophylaxis during neutropenia on bloodstream infection: data from a surveillance program in 8755 patients receiving high-dose chemotherapy for haematologic malignancies between 2009 to 2014. *J Infect*. 2018;77:68–74.
 56. Signorelli J, Zimmer A, Liewer S, Shostrom VK, Freifeld A. Incidence of febrile neutropenia in autologous hematopoietic stem cell transplant (HSCT) recipients on levofloxacin prophylaxis. *Transpl Infect Dis*. 2020;22: e13225.
 57. Alexander S, Fisher BT, Gaur AH, et al. Effect of levofloxacin prophylaxis on bacteremia in children with acute leukemia or undergoing hematopoietic stem cell transplantation: a randomized clinical trial. *JAMA*. 2018;320:995–1004.
 58. Mikulska M, Averbuch D, Tissot F, et al. Fluoroquinolone prophylaxis in hematological cancer patients with neutropenia: ECIL critical appraisal of previous guidelines. *J Infect*. 2018;76:20–37.
 59. Pépin J, Saheb N, Coulombe MA, et al. Emergence of fluoroquinolones as the predominant risk factor for *Clostridium difficile*-associated diarrhea: a cohort study during an epidemic in Quebec. *Clin Infect Dis*. 2005;41:1254–1260.
 60. Melhorn AJ, Brown DA. Safety concerns with fluoroquinolones. *Ann Pharmacother*. 2007;41:1859–1866.
 61. Arnan M, Gudiol C, Calatayud L, et al. Risk factors for, and clinical relevance of, faecal extended-spectrum β -lactamase producing *Escherichia coli* (ESBL-EC) carriage in neutropenic patients with haematological malignancies. *Eur J Clin Microbiol Infect Dis*. 2011;30:355–360.
 62. Vehreschild MJ, Hamprecht A, Peterson L, et al. A multicentre cohort study on colonization and infection with ESBL-producing Enterobacteriaceae in high-risk patients with haematological malignancies. *J Antimicrob Chemother*. 2014;69:3387–3392.

63. Cornejo-Juárez P, Suárez-Cuenca JA, Volkow-Fernández P, et al. Fecal ESBL *Escherichia coli* carriage as a risk factor for bacteremia in patients with hematological malignancies. *Support Care Cancer*. 2016;24:253–259.
64. Brothers AW, Ratliff SM, Bluestone H, et al. Individualized antibiotic plans: An antimicrobial stewardship intervention to reduce meropenem use on the bone marrow transplant service. Poster presented at: 9th Annual International Pediatric Antimicrobial Stewardship Conference; May 31 - June 1. St. Louis, MO; 2018.
65. Centers for Disease Control and Prevention. Facility guidance for control of carbapenem-resistant Enterobacteriaceae (CRE), November 2015 update—CRE toolkit. Available at: <https://www.cdc.gov/hai/pdfs/cre/CRE-guidance-508.pdf>. Accessed September 14, 2020.
66. Tschudin-Sutter S, Lucet JC, Muters NT, Tacconelli E, Zahar JR, Harbarth S. Contact precautions for preventing nosocomial transmission of extended-spectrum β -lactamase-producing *Escherichia coli*: a point/counterpoint review. *Clin Infect Dis*. 2017;65:342–347.
67. Biehl LM, Higgins P, Wille T, et al. Impact of single-room contact precautions on hospital-acquisition and transmission of multidrug-resistant *Escherichia coli*: a prospective multicenter cohort study in hematological and oncological wards. *Clin Microbiol Infect*. 2019;25:1013–1020.
68. Saidel-Odes L, Polachek H, Peled N, et al. A randomized, double-blind, placebo-controlled trial of selective digestive decontamination using oral gentamicin and oral polymyxin E for eradication of carbapenem-resistant *Klebsiella pneumoniae* carriage. *Infect Control Hosp Epidemiol*. 2012;33:14–19.
69. Huttner B, Hausteiner T, Uçkay I, et al. Decolonization of intestinal carriage of extended-spectrum β -lactamase-producing Enterobacteriaceae with oral colistin and neomycin: a randomized, double-blind, placebo-controlled trial. *J Antimicrob Chemother*. 2013;68:2375–2382.
70. Bar-Yoseph H, Hussein K, Braun E, Paul M. Natural history and decolonization strategies for ESBL/carbapenem-resistant Enterobacteriaceae carriage: systematic review and meta-analysis. *J Antimicrob Chemother*. 2016;71:2729–2739.
71. Stoma I, Karpov I, Iskrov I, et al. Decolonization of intestinal carriage of MDR/XDR Gram-negative bacteria with oral colistin in patients with hematologic malignancies: results of a randomized controlled trial. *Mediterr J Hematol Infect Dis*. 2018;10: e2018030.
72. Catho G, Huttner BD. Strategies for the eradication of extended-spectrum beta-lactamase or carbapenemase-producing Enterobacteriaceae intestinal carriage. *Expert Rev Anti Infect Ther*. 2019;17:557–569.
73. DeFilipp Z, Bloom PP, Torres Soto M, et al. Drug-resistant *E. coli* bacteremia transmitted by fecal microbiota transplant. *N Engl J Med*. 2019;381:2043–2050.
74. Satwani P, Freedman JL, Chaudhury S, et al. A multicenter study of bacterial blood stream infections in pediatric allogeneic hematopoietic cell transplantation recipients: the role of acute gastrointestinal graft-versus-host disease. *Biol Blood Marrow Transplant*. 2017;23:642–647.
75. Levinson A, Pinkney K, Jin Z, et al. Acute gastrointestinal graft-vs-host disease is associated with increased enteric bacterial bloodstream infection density in pediatric allogeneic hematopoietic cell transplant recipients. *Clin Infect Dis*. 2015;61:350–357.
76. Youssef A, Hafez H, Madney Y, et al. Incidence, risk factors, and outcome of blood stream infections during the first 100 days post-pediatric allogeneic and autologous hematopoietic stem cell transplantations. *Pediatr Transplant*. 2020;24:e13610.
77. Weiner-Lastinger LM, Abner S, Edwards JR, et al. Antimicrobial-resistant pathogens associated with adult healthcare-associated infections: summary of data reported to the National Healthcare Safety Network, 2015–2017. *Infect Control Hosp Epidemiol*. 2020;41:1–18.
78. Weiner-Lastinger LM, Abner S, Benin AL, et al. Antimicrobial-resistant pathogens associated with pediatric healthcare-associated infections: summary of data reported to the National Healthcare Safety Network, 2015–2017. *Infect Control Hosp Epidemiol*. 2020;41:19–30.
79. Banerjee R, Teng CB, Cunningham SA, et al. Randomized trial of rapid multiplex polymerase chain reaction-based culture identification and susceptibility testing. *Clin Infect Dis*. 2015;61:1071–1080.
80. Ledebner NA, Lopansri BK, Dhiman N, et al. Identification of Gram-negative bacteria and genetic resistance determinants from positive blood culture broths by use of the Verigene gram-negative blood culture multiplex microarray-based molecular assay. *J Clin Microbiol*. 2015;53:2460–2472.
81. Timbrook TT, Morton JB, McConeghy KW, Caffrey AR, Mylonakis E, LaPlante KL. The effect of molecular rapid diagnostic testing on clinical outcomes in bloodstream infections: a systematic review and meta-analysis. *Clin Infect Dis*. 2017;64:15–23.
82. Pancholi P, Carroll KC, Buchan BW, et al. Multicenter evaluation of the Accelerate PhenoTest BC Kit for rapid identification and phenotypic antimicrobial susceptibility testing using morphokinetic cellular analysis. *J Clin Microbiol*. 2018;56: e01329–17.
83. Nguyen MH, Clancy CJ, Pascual AW, et al. Performance of the T2Bacterial Panel for diagnosing bloodstream infections: a diagnostic accuracy study. *Ann Intern Med*. 2019;170:845–852.