

**FIG. 278.9** Diagnostic approach and management algorithm of the newborn whose mother has been suspected or confirmed to have acquired toxoplasmosis during gestation. CSF, Cerebrospinal fluid; CT, computed tomography; IgG, IgM, and IgA, immunoglobulins G, M, and A, respectively; ISAGA, immunosorbent agglutination assay; PCR, polymerase chain reaction. <sup>a</sup>Consider consultation with a physician expert in management of toxoplasmosis during pregnancy (e.g., in the United States, Palo Alto Medical Foundation–*Toxoplasma* Serology Laboratory [PAMF-TSL], [www.pamf.org/serology/](http://www.pamf.org/serology/); 650-853-4828; e-mail, [toxolab@pamf.org](mailto:toxolab@pamf.org) or US [Chicago] National Collaborative Treatment Trial Study, 773-834-4152). <sup>b</sup>Consider sending serum sample to a reference laboratory (e.g., PAMF-TSL). <sup>c</sup>Testing CSF assumes that lumbar puncture is clinically indicated, deemed safe, and feasible. <sup>d</sup>In an attempt to confirm the diagnosis of congenital toxoplasmosis, CSF should be sent for cell count and differential (congenital toxoplasmosis is one of the few causes of eosinophilic meningitis), protein (congenital toxoplasmosis is one of the few causes of extreme elevation of CSF protein), glucose, and *T. gondii* PCR. <sup>e</sup>The recommended regimen is pyrimethamine plus sulfadiazine plus folinic acid (see text). <sup>f</sup>Major clinical signs are referred here: chorioretinitis, brain calcifications, and hydrocephalus. <sup>g</sup>Maternally transferred IgG antibodies usually decline and disappear within 6 to 12 months of life.

obtained from peripheral blood (and not from the umbilical cords) are preferred. In a retrospective study of 164 untreated infants age 0 to 180 days, in the United States, *T. gondii*-specific IgM, IgA, and IgE antibodies were demonstrable in 86.6%, 77.4%, and 40.2% of the infants, respectively.<sup>338</sup> Testing for IgM and IgA antibodies increased the sensitivity of making the diagnosis of congenital toxoplasmosis to 93%, compared

with testing for IgM or IgA alone. The sensitivity of the IgM and IgA has been reported to be somewhat lower in cohorts of treated (prenatal treatment) infants: 64% to 70% for IgM, 53% to 65% for IgA, and 66% to 81% for both.<sup>458</sup>

If IgG antibodies are detected but serologic tests for IgM and IgA antibodies are negative and *T. gondii* is not isolated, follow-up serologic

testing in suspect cases is indicated to attempt to establish the diagnosis. Maternally transferred antibodies usually decline and disappear within 6 to 9 months.

Studies using the Western blot technique have shown that maternal and infant sera may recognize different *T. gondii* antigens when the infant is congenitally infected.<sup>459,460</sup> Combining Western blot with conventional serologic analysis (i.e., IgG, IgM, and IgA tests) has been reported to be more sensitive for the diagnosis of congenital toxoplasmosis at birth and within the first 3 months of life than either test alone.<sup>460–462</sup> Recently, three IgM-bands at 75, 90, and 100 kilodaltons (also known as the “IgM triplet”) have been reported to increase the sensitivity of the diagnosis of congenital toxoplasmosis to 95.8% when combined with prenatal and serologic neonatal tests.<sup>463</sup>

Additional diagnostic methods that have been used successfully to diagnose the infection in infants are direct demonstration of the organism by isolation in mice or cell culture (e.g., placental tissue, body fluid) and PCR in body fluids (e.g., CSF, blood, and urine).<sup>407,464–468</sup> In a study of 161 consecutive infants, CSF PCR was positive in 27 of the 58 (46.5%) congenitally infected infants and was negative in each of the 103 infants without congenital toxoplasmosis.<sup>468</sup> Evaluation of infants with suspected congenital toxoplasmosis should always include ophthalmologic examination, radiologic studies (particularly to detect the presence of cerebral calcifications; if feasible, CT is preferred over ultrasound), and examination of CSF. A more detailed discussion of diagnostic procedures in congenitally infected infants is available from Peyron and colleagues.<sup>9</sup>

In the absence of a systematic screening program for pregnant women, a secondary prevention program that consists of serologic testing of all newborns for IgM antibodies against *T. gondii* has been implemented in Massachusetts.<sup>469,470</sup> Using routine screening of all newborns, congenital infection was confirmed in approximately 1 in 12,000 infants. Greater than 90% of these were identified only through neonatal screening and not through initial clinical examination.

In infants with congenital toxoplasmosis or congenital infection, a rebound in IgG and IgM antibody titers is frequently observed after discontinuation of therapy. In our experience such a serologic rebound has not been shown to be clinically significant.<sup>9</sup>

## THERAPY

Currently recommended drugs against *T. gondii* act primarily against the tachyzoite form and thus do not eradicate the encysted form (bradyzoite). Pyrimethamine is considered to be the most effective anti-*Toxoplasma* agent, and it should, if possible, always be included in a combination drug regimen used against the parasite. Pyrimethamine is a folic acid antagonist. The most common side effect is dose-related suppression of the bone marrow, which may be decreased by concomitant administration of folinic acid (calcium leucovorin). It is not well established how often a blood count should be obtained; a reasonable strategy would be to check a peripheral blood cell and platelet count twice weekly until hematologic parameters have stabilized in a nontoxic range, and then every 2 to 4 weeks. Folinic acid should be administered concomitantly to avoid bone marrow suppression. The parenteral form of folinic acid is well absorbed orally, and 10 to 20 mg/day of folinic acid (up to 50 mg/day is used in AIDS patients) may be given orally (e.g., with orange juice at the same time as the pyrimethamine). Whereas folinic acid does not inhibit the action of pyrimethamine on tachyzoites, folinic acid does and should not be used in patients being treated with pyrimethamine. Less serious side effects of pyrimethamine include GI distress, rash, headaches, and a bad taste in the mouth.

Unless there are circumstances that preclude the use of more than one drug, there is no role for monotherapy in the treatment of toxoplasmosis. A second drug, such as sulfadiazine or clindamycin, should be added. Sulfadiazine acts synergistically with pyrimethamine; most other sulfonamides have inferior activity. The patient must maintain a good urine output to prevent crystalluria and oliguria. The most common side effects associated with sulfadiazine are skin rashes, which may be life threatening,<sup>471</sup> and crystal-induced nephrotoxicity.<sup>472</sup> Worsening encephalopathy, hallucinations, or a new onset of psychiatric symptoms in patients with AIDS may be sulfadiazine induced and must be

considered in the patient who is nonresponsive to otherwise appropriate anti-*Toxoplasma* treatment.<sup>473</sup> A drug rash with sulfonamide therapy does not necessarily preclude its use because successful desensitization protocols have been reported.<sup>474,475</sup> Clindamycin appears to act by targeting translation in the apicoplast of *T. gondii*.<sup>40</sup> Adverse reactions to clindamycin include rash, nausea, vomiting, and diarrhea, which may be associated with *Clostridioides difficile* (formerly *Clostridium difficile*) infection. Myopathy with electromyographic abnormalities and elevated serum creatine phosphokinase levels have been described.<sup>476</sup>

Although clinical experience with TMP-SMX is more limited, this drug combination, which targets folate metabolism in a manner similar to pyrimethamine plus sulfadiazine, has documented activity and can be used in patients requiring parenteral therapy or those in whom cost of pyrimethamine becomes prohibitive.<sup>477</sup> Atovaquone combined with pyrimethamine or sulfadiazine, or in unusual circumstances as a single agent, is another less-well-studied alternative for patients who can be treated with oral therapy. The role of other drugs, including azithromycin, clarithromycin, and dapsone, is less clear; they should only be used as alternatives when the regimens described earlier cannot be used and should be used in combination with pyrimethamine whenever possible.

Spiramycin has been used in pregnant women to reduce transplacental transmission to the fetus; it has not been shown to be effective for acute therapy, maintenance therapy, or primary prophylaxis of TE in AIDS patients. There is no evidence that spiramycin is teratogenic.

Although drugs used to treat toxoplasmosis in the setting of different clinical entities are basically the same, careful attention should be given to the dose and dosing regimen. Recommended doses in immunocompromised patients are usually higher than those in immunocompetent patients. For instance, the recommended dose of pyrimethamine for patients with TE is 50 to 75 mg/day after a loading dose of 200 mg, whereas the dose to treat fetal infection during pregnancy is 25 to 50 mg/day after a loading dose of 100 mg/day for 2 days in the mother.

## Therapy Regimens in Specific Clinical Entities

### Toxoplasmosis in the Immunocompetent Patient

Treatment of immunocompetent adults with the lymphadenopathic form is rarely indicated; this form is usually self-limited. One small randomized trial demonstrated efficacy of a 1-month course of TMP-SMX in reducing lymphadenopathy (65% response vs. 22% for placebo).<sup>478</sup> If visceral disease is clinically overt or symptoms are severe or persistent, treatment may be indicated for 2 to 4 weeks, followed by reassessment of the patient's condition (Table 278.2). Infections acquired by laboratory accident or transfusion of blood products are potentially more severe, and patients who have been infected in these ways probably should be treated (see Table 278.2).

### Toxoplasmosis in the Immunodeficient Patient

Because experience with treatment of toxoplasmosis in immunodeficient patients has been most extensively studied in patients with AIDS, this section focuses primarily on this group of patients. However, information on treatment in AIDS patients likely can, in large part and in the absence of data, be extrapolated directly to other immunodeficient patients (see Table 278.2).

If left untreated, toxoplasmosis in immunodeficient patients is often lethal. Treatment is recommended for 4 to 6 weeks after the resolution of all signs and symptoms (often for 6 months or longer). At one medical center, 80% of non-AIDS immunodeficient patients with toxoplasmosis improved with specific therapy.<sup>479</sup> This rate of improvement is similar to that observed in appropriately treated AIDS patients with TE.<sup>435,480</sup> Chronic (latent) asymptomatic infection in immunodeficient patients is not treated.

The exact dosing schedule for the treatment of toxoplasmosis in non-AIDS immunocompromised patients has not been defined. However, useful information in this regard has resulted from studies performed in AIDS patients with toxoplasmosis.

Therapy for toxoplasmosis in AIDS patients includes acute (primary or induction) treatment, chronic maintenance treatment (secondary

**TABLE 278.2 Treatment Regimens for Toxoplasmosis in Immunocompetent and Immunocompromised Patients**

	IMMUNOCOMPETENT PATIENTS <sup>b</sup>	IMMUNOCOMPROMISED PATIENTS, INCLUDING THOSE WITH AIDS, TRANSPLANTS, CANCER, OR TAKING IMMUNOSUPPRESSOR DRUGS <sup>c</sup>
<b>Preferred Regimen</b>		
Pyrimethamine (PO)	50 mg q12h for 2 days, followed by 25–50 mg daily	200-mg loading dose, followed by 50 (<60 kg)–75 mg/day (≥60 kg)
<i>plus</i>		
Folinic acid <sup>d</sup> (PO)	10–20 mg daily (during and 1 wk after therapy with pyrimethamine)	10–20 mg daily (up to 50 mg/day) (during and 1 wk after therapy with pyrimethamine)
<i>plus</i>		
Sulfadiazine (PO)	1 g q6h	1000 (<60 kg)–1500 mg (≥60 kg) q6h
<b>Preferred Alternative Regimens</b>		
Pyrimethamine–folinic acid	Same doses as above	Same doses as above
<i>plus</i>		
Clindamycin (PO or IV)	300 mg q6h	600 mg q6h (up to 1200 mg q6h)
<i>or</i>		
Atovaquone (PO)	1500 mg PO twice daily	1500 mg PO twice daily
Trimethoprim-sulfamethoxazole (PO or IV)	10 mg/kg/day (trimethoprim component) divided in two to three doses	10 mg/kg/day (trimethoprim component) divided in two to three doses (doses as high as 15–20 mg/kg/day have been used)
<b>Alternative Regimens With Limited Supportive Data<sup>e</sup></b>		
Pyrimethamine–folinic acid	Same doses as above	Same doses as above
<i>plus</i>		
Clarithromycin (PO)	500 mg q12h	500 mg q12h
<i>or</i>		
Dapsone (PO)	100 mg/day	100 mg/day
<i>or</i>		
Azithromycin (PO)	900–1200 mg/day	900–1200 mg/day
Sulfadiazine (PO)	Same doses as above	Same doses as above
<i>plus</i>		
Atovaquone (PO)	Same doses as above	Same doses as above

<sup>a</sup>Assistance is available for the diagnosis and management of toxoplasmosis at the Palo Alto Medical Foundation–*Toxoplasma* Serology Laboratory; Palo Alto, CA; [www.pamf.org/serology/](http://www.pamf.org/serology/); 650-853-4828; [toxolab@pamf.org](mailto:toxolab@pamf.org).

<sup>b</sup>Indicated in patients with (1) ocular disease associated with primary infection or reactivation of latent infection; (2) severe disease, including myocarditis, myositis, hepatitis, pneumonia, brain, or skin lesions and lymphadenopathy; (3) persisting symptoms; and (4) laboratory accidents.

<sup>c</sup>After the successful use of a combination regimen during the acute/primary therapy phase, same agents at half doses are usually used for maintenance or secondary prophylaxis; for pyrimethamine, use 25 mg (<60 kg) to 50 mg/day (>60 kg); for clindamycin, use 600 mg PO every 8 hours.

<sup>d</sup>Folinic acid (leucovorin); should always be given with pyrimethamine to minimize toxicity; folic acid must not be used as a substitute for folinic acid.

<sup>e</sup>These agents have been used in clinical studies with small numbers of patients and have response rates lower than the standard regimens (see text for references). They should be used only in patients who are intolerant of the standard regimens.

AIDS, Acquired immunodeficiency syndrome; IV, intravenously; PO, orally.

prophylaxis), and primary prophylaxis. There are no convincing data from prospective, carefully designed trials to allow the recommendation of monotherapy for induction, maintenance, or primary prophylaxis. Because relapse occurs in up to 80% of cases<sup>471</sup> after the discontinuation of primary therapy, maintenance therapy is recommended for all patients until the CD4 count rises above 200 cells/mm<sup>3</sup> for at least 6 months in response to ART (plasma HIV viral load will often be below detection limits during this period).

Acute therapy should be for at least 3 weeks,<sup>100</sup> and up to 6 weeks or more may be required for more severely ill patients who have not achieved a complete response. Pyrimethamine combined with sulfadiazine and folinic acid is the therapy of choice for AIDS patients with toxoplasmosis and is the standard to which experimental regimens should be compared. This regimen is associated with clinical response in 68% to 95% of patients with TE.<sup>480,481</sup> Unfortunately, up to 40% of patients develop side effects from one or more of the drugs, often requiring discontinuation of one or both agents.<sup>232,426</sup> Pyrimethamine-clindamycin and folinic acid appear comparable in efficacy to pyrimethamine-sulfadiazine,<sup>480,482</sup> but this combination also has substantial toxicity. TMP-SMX<sup>483–485</sup>

(at 10 mg/kg/day of the TMP component, divided in two doses) has shown efficacy similar to the pyrimethamine-sulfadiazine regimen (with a more rapid radiologic response in the TMP-SMX group) in a randomized pilot trial in 77 patients with AIDS<sup>477</sup>; this provides an alternative regimen for situations in which parenteral therapy is required or when pyrimethamine is unavailable or unaffordable. An international, noncomparative study of atovaquone<sup>486,487</sup> (administered orally as a suspension), combined with either pyrimethamine or sulfadiazine as treatment for acute disease, demonstrated 6-week response rates of 75% (21/28 patients) for atovaquone-pyrimethamine and 82% (9/11) for atovaquone-sulfadiazine.<sup>488</sup> Serum levels of atovaquone may predict response but are not commercially available.<sup>487</sup> Doses of atovaquone lower than currently recommended for treatment (750 mg twice daily vs. 1500 mg twice daily) can often achieve therapeutic levels if not coadministered with efavirenz, which can reduce atovaquone levels by approximately 45%.<sup>489</sup> TMP-SMX and atovaquone both are active against *Pneumocystis*, and thus anti-*Pneumocystis* prophylaxis does not need to be administered when these drugs are used. Because clindamycin-pyrimethamine has no demonstrated anti-*Pneumocystis*

activity, additional prophylaxis (e.g., aerosolized pentamidine) needs to be coadministered if patients are at risk for developing *Pneumocystis pneumonia*.

In a 13-patient pilot study of TE using the combination of pyrimethamine, 75 mg/day, and clarithromycin, 1 g every 12 hours, 62% of patients had a complete and 23% a partial clinical response; 15% of patients died by week 3 of therapy,<sup>490</sup> and adverse events resulted in discontinuation of therapy in 27%. No long-term follow-up data are available, and no additional studies with this drug combination have been published. Doses of clarithromycin higher than 500 mg twice daily have been associated with increased mortality in HIV-infected patients receiving therapy for *Mycobacterium avium* infection and should not be used.<sup>491,492</sup>

Dapsone in combination with pyrimethamine has been reported anecdotally to be effective for the treatment of TE when used in an oral dose of 100 mg/day with 25 mg/day of oral pyrimethamine.<sup>493</sup> Doxycycline has had success in the treatment of TE in a few patients when used at 300 mg/day IV in three divided doses.<sup>494</sup> A dosage of 100 mg twice a day was given to six patients intolerant to pyrimethamine-sulfadiazine, but five had associated neurologic and radiologic recurrences while receiving the drug.<sup>495</sup> Further studies are needed to compare the relative efficacy and toxicity of these alternative regimens. Although azithromycin plus pyrimethamine is effective for the treatment of some cases of TE in AIDS patients, its use should be limited, based on results of a recent study demonstrating an inferior response rate, especially during maintenance therapy.<sup>496,497</sup> Alternative regimens used for acute therapy and their dosage schedules are listed in Table 278.2.

Corticosteroids are often given to patients with TE for the reduction of cerebral edema and raised intracranial pressure. The clinical response and survival in patients with TE who received corticosteroids in addition to antimicrobial therapy have been reported to be no different from that of those who received antimicrobial agents alone.<sup>435</sup> The use of these agents may complicate the interpretation of empirical therapy of TE because partial clinical and radiologic improvement may be seen solely resulting from a reduction in cerebral edema and inflammation or a response of CNS lymphoma; moreover, they may further compromise the immune systems of these already very immunodeficient patients. Their use thus should be limited to situations where clinically significant edema or a mass effect is present.

Seizures occur in up to 35% of patients with TE.<sup>426</sup> One retrospective study demonstrated a poorer outcome in those patients who received anticonvulsant therapy, compared with those who did not.<sup>471</sup> Whether this result represents a true drug effect or a selection bias, given that those receiving anticonvulsant therapy are likely to be more severely ill, is unclear. Anticonvulsant agents may be responsible for numerous side effects and drug interactions; for instance, potentially serious interactions can occur between agents such as carbamazepine, phenobarbital, or phenytoin and other drugs used to treat HIV infection, such as protease inhibitors. Anticonvulsant therapy is probably best administered only when seizures have occurred.<sup>471</sup>

The time to clinical response in AIDS patients with TE who were receiving appropriate anti-*Toxoplasma* therapy has been evaluated in a study that included an objective, graded neurologic examination.<sup>435</sup> Of those with a response, 91% improved with respect to at least half of their baseline abnormalities by day 14.<sup>435</sup> AIDS patients with presumed TE had some degree of improvement within 7 to 10 days of the initiation of appropriate anti-*Toxoplasma* therapy. By contrast, a significant number of patients with an alternative diagnosis, including lymphoma, exhibited signs of clinical deterioration as early as 3 to 5 days after the initiation of the empirical regimen for presumed TE.<sup>435</sup> Headaches and seizures were insensitive indicators for a response to therapy. In some cases toxoplasmosis has progressed to death despite the use of appropriate drug regimens.<sup>480,482</sup>

After successful primary therapy, drug dosages are generally decreased for maintenance therapy. No single maintenance regimen that is efficacious with an acceptable adverse reaction profile has yet been identified. Relapse of TE occurs in approximately 20% to 30% of patients who are receiving maintenance therapy, in part because of nonadherence to and patient intolerance of the prescribed regimen.<sup>232</sup> Pyrimethamine, 25 mg/day, plus sulfadiazine, 500 mg four times daily, has been associated with

the lowest relapse rate<sup>480</sup> and is recommended unless there are contraindications or cost considerations preventing its use. When daily therapy with pyrimethamine-sulfadiazine was compared with a twice-weekly regimen for the prevention of recurrence of TE, the latter was found to be less effective.<sup>498,499</sup> Although a subsequent trial by the same group found that three-times weekly therapy was equivalent to daily therapy, the relapse rates for both groups ( $\approx 14.5/100$  patient-years) was higher than was seen with daily therapy ( $4.4/100$  patient-years) in the earlier study.<sup>499,500</sup> Patients receiving the pyrimethamine-sulfadiazine combination do not require another regimen for PCP prophylaxis. Whereas 25% of patients receiving pyrimethamine-clindamycin subsequently developed PCP,<sup>501</sup> no patient receiving pyrimethamine-sulfadiazine developed PCP.<sup>499,501</sup> One 17-patient study demonstrated that TMP-SMX could be safely substituted for pyrimethamine-sulfadiazine after a median of 24-months' maintenance therapy in patients also receiving ART; the major benefit was in decreasing pill burden.<sup>502</sup>

Because of drug toxicity, many patients are unable to continue taking the pyrimethamine-sulfadiazine combination for maintenance therapy. A higher relapse rate has been reported with the use of pyrimethamine-clindamycin compared with pyrimethamine-sulfadiazine for chronic maintenance of TE; hence it is recommended that the clindamycin dose be 1800 mg/day if tolerated.<sup>480,503,504</sup> Encouraging results have been reported with other drug combinations. These include Fansidar (pyrimethamine-sulfadoxine), which has been used in a dose of one tablet twice weekly,<sup>505</sup> and pyrimethamine-dapsone administered on an intermittent schedule (two to three times a week).<sup>506-508</sup> The long half-life of these agents allowed the longer dosing interval, but Fansidar (currently not available in the United States) may have an increased risk of severe cutaneous reactions, and the longer half-life results in slower drug clearance after discontinuation of therapy.<sup>509</sup> When pyrimethamine was used alone as maintenance therapy at 50 mg/day<sup>510,511</sup> and 100 mg/day,<sup>510</sup> the relapse rates were 10% to 28% and 5%, respectively. Atovaquone alone or in combination regimens also appears to have activity based on uncontrolled trials; combination therapy (with sulfadiazine or pyrimethamine) should be used whenever possible.<sup>486-488,512</sup>

The optimal time to initiate ART in HIV-infected patients with TE has not been well defined. In one randomized trial of 282 patients with opportunistic infections other than tuberculosis (only 5% with TE), which compared early (median, 12 days after initiation of opportunistic infection therapy) to deferred (median, 45 days) initiation of ART, the secondary end point of AIDS progression or death occurred significantly less frequently in the early initiation group.<sup>513</sup> Thus, in the absence of contraindications, ART can reasonably be started 2 to 3 weeks after diagnosis of TE.

Primary prophylaxis against *T. gondii* in patients with AIDS has been shown to be effective in preventing acute TE.<sup>514-517</sup> In addition, the use of combination ART in HIV-infected patients has had a profound effect in decreasing the incidence of TE in these patients.<sup>101,103</sup> Primary prophylaxis is recommended for patients who have detectable *Toxoplasma* IgG antibodies and whose lowest CD4<sup>+</sup> count has been less than 100/mm<sup>3</sup> (many experts use <200/mm<sup>3</sup> as the cutoff rather than 100/mm<sup>3</sup>), regardless of the HIV RNA viral load.<sup>504</sup> TMP-SMX (1 double-strength or single-strength tablet/day), dapsone (50 mg/day) plus pyrimethamine (50 mg/wk), and Fansidar (twice weekly; not available in the United States) have been reported to be effective in preventing the first episode of TE.<sup>504,514,517,518</sup> Atovaquone (1500 mg daily) alone or combined with pyrimethamine (25 mg daily) can also be used, although failures in HSCT patients have been reported.<sup>519</sup>

Studies have demonstrated that it is safe to discontinue primary anti-*Toxoplasma* prophylaxis or chronic maintenance therapy when the recovery of a CD4<sup>+</sup> count greater than 200 cells/mm<sup>3</sup> is achieved and sustained in patients on ART.<sup>300-302,520,521</sup> It is important to recognize that in most of these studies, the median CD4<sup>+</sup> count was greater than 300 cells/mm<sup>3</sup> at enrollment, and viral loads were below detection limits or reasonably controlled in most patients. More recent data from a European collaboration after multiple cohorts demonstrated a low risk of developing TE in seropositive patients with CD4<sup>+</sup> counts of 100 to 200 cells/mm<sup>3</sup> and HIV plasma viral loads below detection limits even if they are not receiving primary prophylaxis; similar benefit was not, however, seen in patients receiving chronic maintenance therapy for



prior TE.<sup>522</sup> Current recommendations are to discontinue prophylaxis in patients receiving cART when the CD4 count has risen above 200 cells/mm<sup>3</sup> for at least 3 months (primary prophylaxis) or 6 months (chronic maintenance), and to consider discontinuing primary prophylaxis in patients with CD4 counts of 100 to 200 cells/mm<sup>3</sup> if HIV plasma viral loads are below detection limits for 3 to 6 months.<sup>504</sup>

At present, TMP-SMX is used by most transplant teams as prophylaxis against *Pneumocystis pneumonia*. Its use has also been shown to protect against toxoplasmosis. However, TMP-SMX is not protective in every case, and some patients are not able to tolerate the drug combination. In addition, in some patients sulfonamides may be contraindicated. Thus use of TMP-SMX alone may be sufficient for prevention of toxoplasmosis in patients who are seronegative for *T. gondii* antibodies and who receive heart transplants from seropositive donors (i.e., D<sup>+</sup>/R<sup>-</sup> patients).<sup>523</sup> The optimal schedule for administration of TMP-SMX in heart transplant patients has not been defined. Physicians must decide whether a schedule of daily administration or administration three times each week is to be used. We routinely recommend daily use of single-strength TMP-SMX whenever feasible.<sup>15</sup>

Roxithromycin, administered at 900 mg once a week and may be given in three divided doses,<sup>524</sup> has been reported in a small randomized trial to be effective for primary prophylaxis; roxithromycin is unavailable in the United States. Based on a number of clinical trials, pyrimethamine alone cannot be recommended for primary prophylaxis.<sup>98,525,526</sup> Clarithromycin<sup>527</sup> and spiramycin<sup>528</sup> have been ineffective for primary prophylaxis when they were used alone. In a randomized, placebo-controlled, primary prophylaxis trial, clindamycin (600 mg/day) was associated with an unacceptably high rate of associated GI disease, in particular diarrhea.<sup>529</sup>

Data on the outcome of treatment of AIDS patients with toxoplasmosis outside the CNS are limited; available information on the therapy of ocular<sup>291,530,531</sup> and pulmonary involvement<sup>285,532</sup> indicates that these forms of toxoplasmosis are also responsive to treatment. Therapy was successful in 50% to 77% of patients with pulmonary toxoplasmosis.<sup>285,532</sup>

### Ocular Toxoplasmosis

A recent report from the American Academy of Ophthalmology highlighted the limited data that are available from randomized controlled trials with well-defined end points demonstrating the benefits of therapy.<sup>533</sup> Treatment is most likely indicated in the following settings: any decrease in visual acuity, macular or peripapillary lesions, lesions greater than one optic disk diameter, lesions associated with a moderate-to-severe vitreous inflammatory reaction, the presence of multiple active lesions, the persistence of active disease for more than 1 month, and any ocular lesions associated with recently acquired infection. Because the disease can be self-limited in immunocompetent patients, many clinicians may not treat small, peripheral retinal lesions that are not immediately vision threatening.<sup>14,316,534–536</sup>

The reported benefits of medical therapy are related primarily to the clinical presentation.<sup>316,534</sup> Because there is so much variation in the clinical manifestations of the retinal disease, and because the disease may be self-limited even without treatment, the response to therapy is difficult to interpret. The combination of pyrimethamine (100-mg loading dose given over 24 hours for 2 days, followed by 25–50 mg daily) and sulfadiazine (1 g given four times daily for 4–6 weeks), depending on the clinical response, which is considered “classic” therapy for ocular toxoplasmosis, is the most common drug combination used (see Table 278.2).<sup>534</sup> TMP-SMX showed responses similar to pyrimethamine-sulfadiazine in a recent randomized, single-blind trial, although the latter regimen was used at lower-than-standard doses.<sup>537</sup> Two recent, open, randomized trials found no differences in response rates to intravitreal clindamycin plus dexamethasone compared with an oral regimen combining pyrimethamine, sulfadiazine, leucovorin, and prednisone/prednisolone.<sup>538,539</sup>

Clindamycin (300 mg orally every 6 hours for a minimum of 3 weeks) has also been used with favorable clinical results.<sup>534</sup> Intravitreal clindamycin combined with dexamethasone has been shown to have similar benefits to the oral combination regimen of pyrimethamine and sulfadiazine.<sup>464</sup> Intravitreal TMP-SMX combined with dexamethasone has shown benefit in small uncontrolled studies.<sup>540,541</sup> Other drugs that

may have activity but have been inadequately studied include atovaquone and pyrimethamine plus azithromycin.<sup>542,543</sup> A small randomized trial comparing oral azithromycin to TMP-SMX found no difference in clinical responses.<sup>544</sup>

Systemic corticosteroids are indicated when lesions involve the macula, optic nerve head, or papillomacular bundle, although no randomized trials have demonstrated their benefit compared with antimicrobial therapy alone.<sup>545</sup> Corticosteroid therapy alone may be associated with progressive disease.<sup>546</sup> Photocoagulation has been used both for the treatment of active lesions and for prophylaxis against the spread of lesions because new lesions appear contiguous to old lesions.<sup>534</sup> In some patients vitrectomy and lens removal may be necessary.

Given the high relapse rate seen in some patients with ocular toxoplasmosis, prevention of recurrences would be highly desirable. A randomized, open-label trial of 124 patients in Brazil found that TMP-SMX (1 double-strength tablet every 3 days) was effective in decreasing the frequency of recurrences from 24% to 7% in a population at high risk for recurrences.<sup>214</sup> Although a 10-year follow-up report suggested that efficacy was lost once drug was stopped,<sup>547</sup> a subsequent randomized, placebo-controlled Brazilian trial confirmed the efficacy of prophylactic TMP-SMX (1 double-strength tablet every 2 days) and suggested sustained benefit for 2 years after discontinuing therapy.<sup>548</sup> Such a regimen should be used in patients with frequent or severe recurrences.<sup>549</sup>

For the approach to ocular toxoplasmosis during pregnancy see “Acute Acquired *Toxoplasma* Infection in Pregnant Women.”

### Acute Acquired *Toxoplasma* Infection in Pregnant Women

Treatment of the acutely infected pregnant woman does not completely eliminate but does decrease the incidence and severity of fetal infection. Because there is usually a delay between the acquisition of acute maternal infection, infection of the placenta, and subsequent infection of the fetus, identification of acute maternal infection necessitates immediate institution of treatment of the mother. Most experience of maternal treatment to prevent transmission to the fetus has been with spiramycin (3 g/day, obtainable in the United States from the FDA, 301-796-1400; after hours, 301-796-8210) (Table 278.3). Spiramycin has been accepted by most investigators as being effective in reducing the frequency of maternal transmission of *T. gondii* to the fetus by approximately 60%.<sup>331,382</sup> Spiramycin is indicated for patients confirmed or suspected to have been infected before 14 weeks of gestation. It appears that its maximal efficacy is best achieved when given within 8 weeks of seroconversion.<sup>334</sup> Spiramycin should be continued until delivery, even if results of the amniotic fluid PCR are negative and ultrasound examinations are normal. A retrospective study suggested that the combination of spiramycin and TMP-SMX (after the 14th week) is more effective than spiramycin alone in reducing transmission during the second trimester.<sup>550</sup> If spiramycin cannot be used or is not available, it may be replaced by sulfadiazine (with appropriate precautions at term) or clindamycin alone. However, there are no data on the efficacy of sulfonamides, including sulfadiazine, or clindamycin when these drugs are used for this purpose.

Because spiramycin does not reliably cross the placenta,<sup>551</sup> if fetal infection is documented or highly suspected or maternal infection is confirmed or highly suspected of having been acquired at 14 weeks of gestation or later, the recommended therapeutic regimen is the combination of sulfadiazine (initial dose 75 mg/kg, followed by 50 mg/kg every 12 hours; maximum, 4 g/day), pyrimethamine (50 mg every 12 hours for 2 days, followed by 50 mg daily), and folinic acid (10–20 mg daily, during, and 1 week after completion of pyrimethamine therapy) (see Table 278.3).<sup>334a,334b</sup> Such treatment might be an alternative to the termination of pregnancy when abortion is not allowed by law or for women who desire to continue their pregnancy. Pyrimethamine should not be used in the first 14 weeks of pregnancy because of a concern for teratogenicity. In this circumstance, if indicated, sulfadiazine plus clindamycin can be considered, although there are no data on its efficacy in this situation.<sup>9</sup> In addition, pyrimethamine-sulfadiazine is also recommended for pregnant women in whom a recently acquired acute infection is highly suspected or confirmed during the late second or third trimesters; this is due to the high rates of vertical transmission observed in those

**TABLE 278.3 Treatment Regimens for Toxoplasmosis During Pregnancy and for Congenital Disease**

INDICATION	DRUG REGIMEN
<b>Pregnant Women</b>	
Pregnant women suspected or confirmed of having acquired infection before 14 weeks of gestation. Not recommended during pregnancy if the fetus has documented or suspected infection (see below)	<b>Spiramycin (oral)</b> <sup>a</sup> : 1 g (3 million units) q8h (for a total of 3 g or 9 million units/day) Spiramycin should be continued until delivery, even if fetal infection is not detected (e.g., negative amniotic fluid PCR test and negative follow-up ultrasound)
Pregnant women at ≥14 weeks of gestation and (1) in whom it is suspected or confirmed that acute infection was acquired at or after 14 weeks of gestation or (2) who have a positive amniotic fluid PCR test or an abnormal ultrasound suggestive of congenital toxoplasmosis	<b>Pyrimethamine</b> : 50 mg q12h for 2 days, followed by 50 mg daily <i>plus</i> <b>Sulfadiazine</b> : 75 mg/kg (first dose), followed by 50 mg/kg q12h (max., 4 g/day) <i>plus</i> <b>Folinic acid</b> <sup>b</sup> : 10–20 mg daily during and for 1 wk after pyrimethamine therapy Pyrimethamine is teratogenic and should not be used during pregnancy before week 18 (in some centers in Europe, it is used as early as week 14). Sulfadiazine should not be used alone; consider clindamycin plus sulfadiazine
<b>Infants</b>	
Congenital infection; treatment regimen is usually recommended for 1 yr	<b>Pyrimethamine</b> : 1 mg/kg q12h for 2 days, followed by 1 mg/kg/day for 2 or 6 mo, followed by 1 mg/kg/day every Monday, Wednesday, and Friday <i>plus</i> <b>Sulfadiazine</b> : 50 mg/kg q12h <i>plus</i> <b>Folinic acid</b> <sup>c</sup> : 10 mg three times weekly <b>Prednisone (if CSF protein ≥1 g/dL or severe chorioretinitis)</b> : 0.5 mg/kg q12h (until CSF protein <1 g/dL or resolution of severe chorioretinitis)
<b>Older Children</b>	
Active disease; treatment is usually continued for 1–2 wk beyond resolution of clinical manifestations	<b>Pyrimethamine</b> : 1 mg/kg q12h (max., 50 mg) for 2 days, followed by 1 mg/kg/day (max., 25 mg) <i>plus</i> <b>Sulfadiazine</b> : 75 mg/kg (first dose), followed by 50 mg/kg q12h (max., 4 g/day) <i>plus</i> <b>Folinic acid</b> <sup>d</sup> : 10–20 mg three times weekly <b>Prednisone (severe chorioretinitis)</b> : 1 mg/kg/day in two divided doses; max., 40 mg/day, rapid taper

<sup>a</sup>Assistance is available for the diagnosis and management of toxoplasmosis at the Palo Alto Medical Foundation–*Toxoplasma* Serology Laboratory; Palo Alto, CA; [www.pamf.org/serology/](http://www.pamf.org/serology/); 650-853-4828; [toxolab@pamf.org](mailto:toxolab@pamf.org); or US (Chicago) National Collaborative Treatment Trial Study; 773-834-4152.

<sup>b</sup>Spiramycin is not teratogenic, and it is available in the United States through the investigational new drug process at the US Food and Drug Administration (301-796-1600). Prior consultation with medical consultants is required.

<sup>c</sup>Folinic acid (leucovorin) should always be given with pyrimethamine to minimize toxicity; folic acid must not be used as a substitute for folinic acid.

<sup>d</sup>CSF, Cerebrospinal fluid; max., maximum; PCR, polymerase chain reaction.

stages of gestation and should be recommended even though fetal infection may not yet have been confirmed.

A group of European investigators reported between 1999 and 2007 that, in their studies, a significant effect of prenatal treatment on the risk of vertical transmission and clinical signs of congenital toxoplasmosis was not detected.<sup>552–555</sup> These results are not surprising because the studies included very few untreated women in their analysis, most untreated women were infected during the third trimester, and severe cases were excluded.<sup>556,557</sup> More recently, several studies from Europe have consistently reported evidence for an association between early treatment and reduced risk in the incidence and severity of congenital toxoplasmosis.<sup>334,336,337,558,559</sup> Given that recent studies have consistently

reported a clinically significant benefit<sup>556</sup> and that a significant benefit was never ruled out in previous studies, most authorities continue to recommend spiramycin or pyrimethamine-sulfadiazine for women with suspected or confirmed acute *T. gondii* infection acquired during gestation.

Pregnant women with toxoplasmic chorioretinitis as a result of reactivation of chronic disease do not have a higher risk to transmit the parasite to their offspring than do pregnant women who have been infected before pregnancy and do not have ocular disease.<sup>560</sup> Their eye disease should be treated according to the indications discussed in the section “Ocular Toxoplasmosis.” Pregnant women with toxoplasmic chorioretinitis thought to be a manifestation of recently acquired infection should be treated because of both the eye disease and the risk of transmission of the infection to their fetus.

## Congenital Infection

Detailed information on and recommendations for the postnatal treatment of congenital toxoplasmosis are reviewed elsewhere,<sup>9,561</sup> but we favor continuous sulfadiazine (50 mg/kg every 12 hours), pyrimethamine (loading dose, 1 mg/kg every 12 hours for 2 days; then beginning on day 3, 1 mg/kg per day for 2 or 6 months; then this dose every Monday, Wednesday, and Friday), and folinic acid (10 mg three times weekly during and for 1 week after pyrimethamine therapy) for a minimum of 12 months (see Table 278.3).<sup>557,562</sup> Serial follow-up to gauge the response of the infant to therapy should include neuroradiology, ophthalmologic examinations, and CSF analysis if indicated.<sup>9</sup> Fansidar (unavailable in the United States) has also been used and reported to be well tolerated,<sup>563,564</sup> although safety concerns and availability of alternative regimens preclude recommending its use.

Studies have shown that outcomes are substantially better for most, but not all, infants treated from the neonatal period for 12 months with pyrimethamine-sulfadiazine and leucovorin, compared with historical control subjects receiving no or short-course therapy.<sup>562,565–567</sup> Improvement in intellectual function, regression of retinal lesions, reduction in anticonvulsant drug requirements, and prevention of auditory sequelae appear to be the major benefits of such treatment, which was combined with CSF shunting if required.<sup>562</sup> Signs of active infection resolved within weeks of initiation of treatment. In a significant number of treated children, cerebral calcifications diminished in size or resolved.<sup>567</sup> They also reported that new central chorioretinal lesions were uncommon (14%) in children with congenital toxoplasmosis who have been treated during their first year of life, compared with historical reports of much higher rates (≥82%) for untreated children or those treated for only 1 month near birth.<sup>349</sup>

## PREVENTION AND PROPHYLAXIS

### General Methods

Prevention is most important in seronegative pregnant women and immunodeficient patients. It is most readily accomplished through education of these patients by their personal physicians (Table 278.4). The goal is to avoid the ingestion of and contact with tissue cysts or sporulated oocysts. Tissue cysts in meat are made noninfectious by heating the meat to 67°C/153°F (i.e., should be cooked to “well done” with no pink meat visible in the center) or by freezing it to –20°C/–4°F (which is not attainable in most home freezers) for at least 48 hours. Meat that is smoked, cured in brine, or dried may still be infectious. Oysters, clams, or mussels should not be eaten raw. Hands should be washed thoroughly after handling raw meat or vegetables, eggs should not be eaten raw, and unpasteurized milk (particularly milk from goats) should be avoided. Vectors such as flies and cockroaches should be controlled. Areas contaminated with cat feces should be avoided altogether. Disposable gloves should be worn while disposing of cat litter material, working in the garden, or cleaning a child’s sandbox. Oocysts are killed if the cat litter pan is soaked in nearly boiling water for 5 minutes. If the litter pan is cleaned every day, oocysts will not have a chance to sporulate. Serologic testing of cats is unwarranted because testing does not demonstrate whether the infected cat is excreting oocysts. Untreated water has been shown to be an effective vehicle for the transmission of the parasite, and drinking water sources potentially contaminated with oocytes should be avoided.

### Serologic Screening and Prophylaxis Acute *Toxoplasma gondii* Infection and Toxoplasmosis in the Immunodeficient Patient

Transmission of *T. gondii* and death caused by the infection have resulted from the transfusion of leukocyte-rich blood products and by organ transplantation in immunodeficient patients. Transmission of infection by these routes may occur frequently enough to warrant screening for antibody to *T. gondii* in leukocyte-rich blood product donors and possibly to exclude seropositive people as organ donors to seronegative potential recipients whenever feasible.

Primary prophylaxis can prevent toxoplasmosis in patients dually infected with HIV and *T. gondii* (see “*Toxoplasmosis in the Immunodeficient Patient*” under “Therapy”). Prophylactic treatment (pyrimethamine, 25 mg orally every day for 6 weeks after transplantation) has been used with apparent success in seronegative recipients of hearts transplanted from seropositive donors.<sup>568</sup> However, TMP-SMX used for PCP prophylaxis in solid-organ transplant patients is also effective as primary prophylaxis against *T. gondii* and can be used without addition of pyrimethamine. Primary prophylaxis in BMT and HSCT patients is particularly challenging because TMP-SMX cannot safely be used early (i.e., before engraftment), whereas in patients with all other transplant organs, it can be used immediately after transplantation. Atovaquone is an alternative, but its efficacy has not been evaluated in controlled

trials, and breakthroughs have been reported.<sup>569</sup> A preemptive approach using *Toxoplasma* PCR in whole blood, similar to the strategy for preventing CMV disease using serum PCR, should be entertained for immunocompromised patients in whom prophylaxis is not feasible during periods of high risk for development of toxoplasmosis.<sup>281</sup>

### Congenital *Toxoplasma gondii* Infection and Toxoplasmosis

Congenital toxoplasmosis is a preventable disease. It is therefore the responsibility of physicians who care for pregnant women to educate them on how they can prevent themselves from becoming infected and thereby not place their fetus at risk. A lack of adoption of a systematic serologic screening program in the United States leaves education as the principal means of preventing this tragic disease. If physicians choose to screen their patients serologically, the appropriate tests must be used, the laboratory performing the tests must be competent, and the test results must be interpreted correctly. Nonreference laboratories can effectively accomplish the initial screening testing by simultaneously performing IgG and IgM antibody tests. Women with positive results in the initial IgG antibody test should have a test for IgM antibody performed on the same serum. Only IgM-positive test results need to be sent to a reference laboratory for confirmatory testing (e.g., PAMF-TSL).<sup>359,379,381</sup> Clinical decisions should not be made based on positive IgM test results alone. In patients with IgG antibodies at any titer, a negative IgM antibody test result in the first trimester, and no clinical signs of acute toxoplasmosis, no further testing would be necessary because the probability of acute acquired infection in these women is extremely low. Given the same circumstances in the second trimester of pregnancy, a negative IgM test result rules out, for practical purposes, recent acquisition of acute infection. A negative IgM test in the third trimester may occur in a patient who acquired the infection earlier in gestation.

In some countries (e.g., France, Austria, Italy, Slovenia, Lithuania, and Uruguay), initially seronegative pregnant women are retested at various intervals during gestation to detect seroconversion and institute treatment.<sup>570</sup> Monthly screening of seronegative pregnant women was recently reported to significantly decrease the risk of vertical transmission and of clinical signs at 3 years of age.<sup>558</sup> The appropriate use of prenatal diagnosis, followed by the optimal use of spiramycin and/or pyrimethamine-sulfadiazine-folinic acid, can markedly reduce the incidence of clinically significant congenital toxoplasmosis.<sup>4</sup> Its absence is often associated with poor outcomes.<sup>338</sup> Results of a study in France<sup>333</sup> on the incidence of *T. gondii* infection in fetuses of women whose date of acquiring the infection during the gestation was known and who were treated with spiramycin are shown in Table 278.5.

<sup>a</sup>References 333, 336, 337, 558, 571, 572.

**TABLE 278.4 Measures to Prevent Primary  
*Toxoplasma gondii* Infection**

- Avoid contact with materials potentially contaminated with cat feces, especially handling of cat litter and gardening. Gloves are advised when these activities are necessary. Because oocysts require 1 to 2 days to mature, dispose of all cat feces daily.
- Disinfect cat litter box with near-boiling water for 5 minutes before handling.
- Avoid mucous membrane contact when handling raw meat.
- Wash hands thoroughly after contact with raw meat.
- Kitchen surfaces and utensils that have come in contact with raw meat should be washed.
- Freeze meat to –20°C (–4°F) for at least 48 hours.
- Cook meat to 67°C (153°F) or “well done” (meat should not be “pink” in the center).
- Avoid ingestion of dried, smoked, or cured meat because they may be infectious.
- Wash fruits and vegetables before consumption.
- Refrain from skinning animals.
- Avoid drinking untreated water potentially contaminated with oocysts.
- Avoid drinking unpasteurized goat’s milk.
- Avoid eating raw oysters, clams, and mussels.

**TABLE 278.5 Rates of Congenital Transmission in 270 Women and the Sensitivity and Negative Predictive Value of Amniotic Fluid PCR for Prenatal Diagnosis of Congenital Toxoplasmosis, According to Gestational Age at Which Maternal Infection Was Acquired**

GESTATIONAL AGE AT MATERNAL INFECTION <sup>a</sup> (wk)	NO. OF INFECTED FETUSES <sup>b</sup> /TOTAL (%)	AMNIOTIC FLUID PCR	
		Sensitivity (CI) (%)	NPV (CI) (%)
±6	0/14 (0)	N/A	100 (78–100)
7–11	7/50 (14)	29 (8–65)	90 (81–98)
12–16	7/61 (11.5)	57 (21–94)	95 (86–98)
17–21	14/66 (21.2)	93 (68–99)	98 (90–99.7)
22–26	16/36 (44.4)	63 (39–86)	77 (61–93)
27–31	19/30 (63.3)	68 (48–89)	65 (42–87)
≥32	12/13 (92.3)	50 (22–78)	14 (3–52)
TOTAL	75/270 (28)	N/A	N/A

<sup>a</sup>Maternal infection was diagnosed by seroconversion in the 270 women; 261 (97%) were given treatment with spiramycin.

<sup>b</sup>Congenital infection was diagnosed by the persistence of *Toxoplasma* immunoglobulin antibodies after 1 year of life.

CI, Confidence interval; N/A, not applicable; NPV, negative predictive value; PCR, polymerase chain reaction.

Note: The positive predictive value was 100%, regardless of gestational age.

Modified from Montoya JG, Remington JS. Management of *Toxoplasma gondii* infection during pregnancy. Clin Infect Dis. 2008;47:554–566.



In pregnant women infected with both HIV and *T. gondii*, we recommend that primary prophylaxis against *T. gondii* be introduced when their CD4 T-cell count falls below 200/mm<sup>3</sup>. If the patient is receiving TMP-SMX, additional prophylaxis is probably not necessary. Otherwise, spiramycin can be used at a dose of 3 g/day.

## ACKNOWLEDGMENTS

We gratefully acknowledge and thank Dr. Jack S. Remington, whose lifelong study of toxoplasmosis and authorship of previous editions of this chapter provided the foundation upon which the current chapter is based. We would also like to thank Dr. Alan Sher for his review of the manuscript and helpful suggestions.

## Key References

The complete reference list is available online at Expert Consult.

5. Vietzke WM, Gelderman AH, Grimley PM, et al. Toxoplasmosis complicating malignancy. Experience at the National Cancer Institute. *Cancer*. 1968;21:816–827.
7. Luft BJ, Conley F, Remington JS, et al. Outbreak of central-nervous-system toxoplasmosis in western Europe and North America. *Lancet*. 1983;1:781–784.
9. Peyron F, Wallon M, Kieffer F, et al. Toxoplasmosis. In: Wilson CB, Nizet V, Maldonado Y, et al, eds. *Remington and Klein's Infectious Diseases of the Fetus and Newborn Infant*. 8th ed. Philadelphia: Saunders/Elsevier; 2015:949–1042.
27. Su C, Khan A, Zhou P, et al. Globally diverse *Toxoplasma gondii* isolates comprise six major clades originating from a small number of distinct ancestral lineages. *Proc Natl Acad Sci USA*. 2012;109:5844–5849.
76. Bowie WR, King AS, Werker DH, et al. Outbreak of toxoplasmosis associated with municipal drinking water. *Lancet*. 1997;350:173–177.
81. Bahia-Oliveira LM, Jones JL, Azevedo-Silva J, et al. Highly endemic, waterborne toxoplasmosis in north Rio de Janeiro state, Brazil. *Emerg Infect Dis*. 2003;9:55–62.
91. Jones JL, Kruszon-Moran D, Elder S, et al. *Toxoplasma gondii* infection in the United States, 2011–2014. *Am J Trop Med Hyg*. 2018;98:551–557.
101. Abgrall S, Rabaud C, Costagliola D. Incidence and risk factors for toxoplasmic encephalitis in human immunodeficiency virus-infected patients before and during the highly active antiretroviral therapy era. *Clin Infect Dis*. 2001;33:1747–1755.
102. Jones JL, Roberts JM. Toxoplasmosis hospitalizations in the United States, 2008, and trends, 1993–2008. *Clin Infect Dis*. 2012;54:e58–e61.
156. Suzuki Y, Orellana MA, Schreiber RD, et al. Interferon- $\gamma$ : the major mediator of resistance against *Toxoplasma gondii*. *Science*. 1988;240:516–518.
213. Montoya JG, Parmley S, Liesenfeld O, et al. Use of the polymerase chain reaction for diagnosis of ocular toxoplasmosis. *Ophthalmology*. 1999;106:1554–1563.
214. Silveira C, Belfort R Jr, Muccioli C, et al. The effect of long-term intermittent trimethoprim/sulfamethoxazole treatment on recurrences of toxoplasmic retinochoroiditis. *Am J Ophthalmol*. 2002;134:41–46.
237. McCabe RE, Brooks RG, Dorfman RF, et al. Clinical spectrum in 107 cases of toxoplasmic lymphadenopathy. *Rev Infect Dis*. 1987;9:754–774.
263. Montoya JG, Giraldo LF, Efron B, et al. Infectious complications among 620 consecutive heart transplant patients at Stanford University Medical Center. *Clin Infect Dis*. 2001;33:629–640.
281. Gajurel K, Dhakal R, Montoya JG. *Toxoplasma* prophylaxis in haematopoietic cell transplant recipients: a review of the literature and recommendations. *Curr Opin Infect Dis*. 2015;28:283–292.
283. Sumi M, Norose K, Hikosaka K, et al. Clinical characteristics and computed tomography findings of pulmonary toxoplasmosis after hematopoietic stem cell transplantation. *Int J Hematol*. 2016;104:729–740.
294. Antinori A, Larussa D, Cingolani A, et al. Prevalence, associated factors, and prognostic determinants of AIDS-related toxoplasmic encephalitis in the era of advanced highly active antiretroviral therapy. *Clin Infect Dis*. 2004;39:1681–1691.
299. van Bilsen WPH, van den Berg CHSB, Rijnders BJA, et al. Immune reconstitution inflammatory syndrome associated with toxoplasmic encephalitis in HIV-infected patients. *AIDS*. 2017;31:1415–1424.
300. Furrer H, Opravil M, Bernasconi E, et al. Stopping primary prophylaxis in HIV-1-infected patients at high risk of *Toxoplasma* encephalitis. Swiss HIV Cohort Study. *Lancet*. 2000;355:2217–2218.
301. Kirk O, Reiss P, Uberti-Foppa C, et al. Safe interruption of maintenance therapy against previous infection with four common HIV-associated opportunistic pathogens during potent antiretroviral therapy. *Ann Intern Med*. 2002;137:239–250.
315. Jones JL, Bonetti V, Hollan GN, et al. Ocular toxoplasmosis in the United States: recent and remote infections. *Clin Infect Dis*. 2015;60:271–273.
317. Aleixo AL, Curi AL, Benchimol EI, et al. Toxoplasmic retinochoroiditis: clinical characteristics and visual outcome in a prospective study. *PLoS Negl Trop Dis*. 2016;10:e0004685.
319. Silveira C, Muccioli C, Holland GN, et al. Ocular involvement following an epidemic of *Toxoplasma gondii* infection in Santa Isabel do Ivaí, Brazil. *Am J Ophthalmol*. 2015;159:1013–1021.e3.
332. Hohlfield P, Daffos F, Thulliez P, et al. Fetal toxoplasmosis: outcome of pregnancy and infant follow-up after in utero treatment. *J Pediatr*. 1989;115:765–769.
- 334a. Mandelbrot L, Kieffer F, Sitta R, et al. Prenatal therapy with pyrimethamine + sulfadiazine vs spiramycin to reduce placental transmission of toxoplasmosis: a multicenter, randomized trial. *Am J Obstet Gynecol*. 2018;219:386.e1–386.e9.
- 334b. Montoya JG. Systematic screening and treatment of toxoplasmosis during pregnancy: is the glass half full or half empty? *Am J Obstet Gynecol*. 2018;219:315–319.
335. Dunn D, Wallon M, Peyron F, et al. Mother-to-child transmission of toxoplasmosis: risk estimates for clinical counselling. *Lancet*. 1999;353:1829–1833.
337. Cortina-Borja M, Tan HK, Wallon M, et al. Prenatal treatment for serious neurological sequelae of congenital toxoplasmosis: an observational prospective cohort study. *PLoS Med*. 2010;7:e1000351.
350. Wallon M, Garweg JG, Abrahamowicz M, et al. Ophthalmic outcomes of congenital toxoplasmosis followed until adolescence. *Pediatrics*. 2014;133:e601–e608.
359. Dhakal R, Gajurel K, Pomares C, et al. Significance of a positive *Toxoplasma* immunoglobulin M test result in the United States. *J Clin Microbiol*. 2015;53:3601–3605.
368. Rougier S, Montoya JG, Peyron F. Lifelong persistence of *Toxoplasma* cysts: a questionable dogma? *Trends Parasitol*. 2017;33:414.
382. Montoya JG, Remington JS. Management of *Toxoplasma gondii* infection during pregnancy. *Clin Infect Dis*. 2008;47:554–566.
392. Pomares C, Montoya JG. Laboratory diagnosis of congenital toxoplasmosis. *J Clin Microbiol*. 2016;54:2448–2454.
419. Robert-Gangneux F, Sterkers Y, Yera H, et al. Molecular diagnosis of toxoplasmosis in immunocompromised patients: a three-year multicenter retrospective study. *J Clin Microbiol*. 2015;53:1677–1684.
426. Porter SB, Sande M. Toxoplasmosis of the central nervous system in the acquired immunodeficiency syndrome. *N Engl J Med*. 1992;327:1643–1648.
435. Luft BJ, Hafner R, Korzun AH, et al. Toxoplasmic encephalitis in patients with the acquired immunodeficiency syndrome. *N Engl J Med*. 1993;329:995–1000.
443. Navia BA, Petito CK, Gold JW, et al. Cerebral toxoplasmosis complicating the acquired immune deficiency syndrome: clinical and neuropathological findings in 27 patients. *Ann Neurol*. 1986;19:224–238.
480. Katlama C, De Wit S, O'Doherty E, et al. Pyrimethamine-clindamycin vs. pyrimethamine-sulfadiazine as acute and long-term therapy for toxoplasmic encephalitis in patients with AIDS. *Clin Infect Dis*. 1996;22:268–275.
485. Torre D, Speranza F, Martegani R, et al. A retrospective study of treatment of cerebral toxoplasmosis in AIDS patients with trimethoprim-sulphamethoxazole. *J Infect*. 1998;37:15–18.
487. Torres R, Weinberg W, Stansell J, et al. Atovaquone for salvage treatment and suppression of toxoplasmic encephalitis in patients with AIDS. *Clin Infect Dis*. 1997;24:422–429.
488. Chirgwin K, Hafner R, Lepout C, et al. Randomized phase II trial of atovaquone with pyrimethamine or sulfadiazine for treatment of toxoplasmic encephalitis in patients with acquired immunodeficiency syndrome: ACTG 237/ANRS 039 Study. AIDS Clinical Trials Group 237/Agence Nationale de Recherche sur le SIDA, Essai 039. *Clin Infect Dis*. 2002;34:1243–1250.
499. Podzamczar D, Miró J, Bolao F, et al. Twice-weekly maintenance therapy with sulfadiazine-pyrimethamine to prevent recurrent toxoplasmic encephalitis in patients with AIDS. *Ann Intern Med*. 1995;123:175–180.
501. Heald A, Flepp M, Chave J-P, et al. Treatment for cerebral toxoplasmosis protects against *Pneumocystis carinii* pneumonia in patients with AIDS. *Ann Intern Med*. 1991;115:760–763.
504. Panel on Opportunistic Infections in HIV-Infected Adults and Adolescents. Guidelines for the Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents: Recommendations from the Centers for Disease Control and Prevention, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. [http://aidsinfo.nih.gov/contentfiles/lvguidelines/adult\\_oi.pdf](http://aidsinfo.nih.gov/contentfiles/lvguidelines/adult_oi.pdf). Accessed December 30, 2017.
512. Katlama C, Mouton B, Gourdon D, et al. Atovaquone as long-term suppressive therapy for toxoplasmic encephalitis in patients with AIDS and multiple drug intolerance. *AIDS*. 1996;10:1107–1112.
513. Zolopa A, Andersen J, Powderly W, et al. Early antiretroviral therapy reduces AIDS progression/death in individuals with acute opportunistic infections: a multicenter randomized strategy trial. *PLoS ONE*. 2009;4:e5575.
514. Carr A, Tindall B, Brew BJ, et al. Low-dose trimethoprim-sulfamethoxazole prophylaxis for toxoplasmic encephalitis in patients with AIDS. *Ann Intern Med*. 1992;117:106–111.
517. Girard P-M, Landman R, Gaubert C, et al. Dapsone-pyrimethamine compared with aerosolized pentamidine as a primary prophylaxis against *Pneumocystis carinii* pneumonia and toxoplasmosis in HIV infection. *N Engl J Med*. 1993;328:1514–1520.
519. Gajurel K, Dhakal R, Montoya JG. Toxoplasmosis in hematopoietic cell transplant recipients. *Transpl Infect Dis*. 2017;19.
537. Soheilian M, Sadoughi MM, Ghajarnia M, et al. Prospective randomized trial of trimethoprim/sulfamethoxazole versus pyrimethamine and sulfadiazine in the treatment of ocular toxoplasmosis. *Ophthalmology*. 2005;112:1876–1882.
547. Silveira C, Muccioli C, Nussenblatt R, et al. The effect of long-term intermittent trimethoprim/sulfamethoxazole treatment on recurrences of toxoplasmic retinochoroiditis: 10 years of follow-up. *Ocul Immunol Inflamm*. 2015;23:246–247.
558. Wallon M, Peyron F, Cornu C, et al. Congenital *Toxoplasma* infection: monthly prenatal screening decreases transmission rate and improves clinical outcome at age 3 years. *Clin Infect Dis*. 2013;56:1223–1231.
561. Maldonado YA, Read JS, Committee on Infectious Diseases. Diagnosis, treatment, and prevention of congenital toxoplasmosis in the United States. *Pediatrics*. 2017;139:e20163860.



## References

- Montoya JG, Liesenfeld O. Toxoplasmosis. *Lancet*. 2004;363:1965–1976.
- Nicollé C, Manceaux L. Sur une infection a corps de Leishman (ou organismes voisins) du gondi. *C R Seances Acad Sci*. 1908;146:207–209.
- Janku J. Pathogenesa a patologicka anatomie tak nazvaneho vrozeneho kolobomu zlute skvrny v oku normalne velikem a mikrophthalmickem s nalezem parazitu v sitnici. *Cas Lek Ces*. 1923;62:1021–1027, 54–59, 81–85, 111–115, 138–144.
- Wolf A, Cowen D, Paige BH. Toxoplasmic encephalomyelitis, III. A new case of granulomatous encephalomyelitis due to a protozoan. *Am J Pathol*. 1939;15:657–694.
- Vietzke WM, Gelderman AH, Grimley PM, et al. Toxoplasmosis complicating malignancy. Experience at the National Cancer Institute. *Cancer*. 1968;21:816–827.
- Israelski DM, Remington JS. Toxoplasmosis in the non-AIDS immunocompromised host. In: Remington J, Swartz M, eds. *Current Clinical Topics in Infectious Diseases*. Vol. 13. London: Blackwell Scientific; 1993:322–356.
- Luft BJ, Conley F, Remington JS, et al. Outbreak of central-nervous-system toxoplasmosis in western Europe and North America. *Lancet*. 1983;1:781–784.
- Lykins J, Wang K, Wheeler K, et al. Understanding toxoplasmosis in the United States through “large data” analyses. *Clin Infect Dis*. 2016;63:468–475.
- Peyron F, Wallon M, Kieffer F, et al. Toxoplasmosis. In: Wilson CB, Nizet V, Maldonado Y, et al, eds. *Remington and Klein's Infectious Diseases of the Fetus and Newborn Infant*. 8th ed. Philadelphia: Saunders/Elsevier; 2015:949–1042.
- Ajzenberg D. 1995–2015: it is time to celebrate 20 years of (intensive) genotyping of *Toxoplasma gondii* strains. *Future Microbiol*. 2015;10:689–691.
- Cai Y, Shen J. Modulation of host immune responses to *Toxoplasma gondii* by microRNAs. *Parasite Immunol*. 2017;39.
- Ngô HM, Zhou Y, Lorenzi H, et al. *Toxoplasma* modulates signature pathways of human epilepsy, neurodegeneration & cancer. *Sci Rep*. 2017;7:11496.
- Flegel J. Predictors of *Toxoplasma gondii* infection in Czech and Slovak populations: the possible role of cat-related injuries and risky sexual behavior in the parasite transmission. *Epidemiol Infect*. 2017;145:1351–1362.
- Wakefield D, Cunningham ET Jr, Pavesio C, et al. Controversies in ocular toxoplasmosis. *Ocul Immunol Inflamm*. 2011;19:2–9.
- Derouin F, Pelloux H. Prevention of toxoplasmosis in transplant patients. *Clin Microbiol Infect*. 2008;14:1089–1101.
- Dubey JP, Lindsay DS, Speer CA. Structures of *Toxoplasma gondii* tachyzoites, bradyzoites, and sporozoites and biology and development of tissue cysts. *Clin Microbiol Rev*. 1998;11:267–299.
- Sibley LD, Boothroyd JC. Virulent strains of *Toxoplasma gondii* comprise a single clonal lineage. *Nature*. 1992;359:82–85.
- Howe DK, Honoré S, Derouin F, et al. Determination of genotypes of *Toxoplasma gondii* strains isolated from patients with toxoplasmosis. *J Clin Microbiol*. 1997;35:1411–1414.
- Fuentes I, Rubio JM, Ramirez C, et al. Genotypic characterization of *Toxoplasma gondii* strains associated with human toxoplasmosis in Spain: direct analysis from clinical samples. *J Clin Microbiol*. 2001;39:1566–1570.
- Ajzenberg D, Cogne N, Paris L, et al. Genotype of 86 *Toxoplasma gondii* isolates associated with human congenital toxoplasmosis, and correlation with clinical findings. *J Infect Dis*. 2002;186:684–689.
- Xia J, Cheng XY, Wang XJ, et al. Association between *Toxoplasma gondii* types and outcomes of human infection: a meta-analysis. *Acta Microbiol Immunol Hung*. 2017;64:229–244.
- Boothroyd JC, Grigg ME. Population biology of *Toxoplasma gondii* and its relevance to human infection: do different strains cause different disease? *Curr Opin Microbiol*. 2002;5:438–442.
- Shobab L, Pleyer U, Johnsen J, et al. *Toxoplasma* serotype is associated with development of ocular toxoplasmosis. *J Infect Dis*. 2013;208:1520–1528.
- Fekkar A, Ajzenberg D, Bodaghi B, et al. Direct genotyping of *Toxoplasma gondii* in ocular fluid samples from 20 patients with ocular toxoplasmosis: predominance of type II in France. *J Clin Microbiol*. 2011;49:1513–1517.
- Lehmann T, Marcet PL, Graham DH, et al. Globalization and the population structure of *Toxoplasma gondii*. *Proc Natl Acad Sci USA*. 2006;103:11423–11428.
- Darde ML. *Toxoplasma gondii*, “new” genotypes and virulence. *Parasite*. 2008;15:366–371.
- Su C, Khan A, Zhou P, et al. Globally diverse *Toxoplasma gondii* isolates comprise six major clades originating from a small number of distinct ancestral lineages. *Proc Natl Acad Sci USA*. 2012;109:5844–5849.
- Minot S, Melo MB, Li F, et al. Admixture and recombination among *Toxoplasma gondii* lineages explain global genome diversity. *Proc Natl Acad Sci USA*. 2012;109:13458–13463.
- Lorenzi H, Khan A, Behnke MS, et al. Local admixture of amplified and diversified secreted pathogenesis determinants shapes mosaic *Toxoplasma gondii* genomes. *Nat Commun*. 2016;7:10147.
- Fritz HM, Buchholz KR, Chen X, et al. Transcriptomic analysis of *Toxoplasma* development reveals many novel functions and structures specific to sporozoites and oocysts. *PLoS ONE*. 2012;7:e29998.
- Fritz HM, Bowyer PW, Bogoy M, et al. Proteomic analysis of fractionated *Toxoplasma* oocysts reveals clues to their environmental resistance. *PLoS ONE*. 2012;7:e29955.
- Radke JR, Striepen B, Guerini MN, et al. Defining the cell cycle for the tachyzoite stage of *Toxoplasma gondii*. *Mol Biochem Parasitol*. 2001;115:165–175.
- Dubey JP. Advances in the life cycle of *Toxoplasma gondii*. *Int J Parasitol*. 1998;28:1019–1024.
- Barragan A, Sibley LD. Transseptal migration of *Toxoplasma gondii* is linked to parasite motility and virulence. *J Exp Med*. 2002;195:1625–1633.
- Lachenmaier SM, Deli MA, Meissner M, et al. Intracellular transport of *Toxoplasma gondii* through the blood-brain barrier. *J Neuroimmunol*. 2011;232:119–130.
- Hakimi MA, Olias P, Sibley LD. *Toxoplasma* effectors targeting host signaling and transcription. *Clin Microbiol Rev*. 2017;30:615–645.
- Carruthers V, Boothroyd JC. Pulling together: an integrated model of *Toxoplasma* cell invasion. *Curr Opin Microbiol*. 2007;10:83–89.
- Boothroyd JC, Dubremetz JE. Kiss and spit: the dual roles of *Toxoplasma* rhoptries. *Nat Rev Microbiol*. 2008;6:79–88.
- McFadden GI. The apicoplast. *Protozoa*. 2011;248:641–650.
- Camps M, Arrizabalaga G, Boothroyd J. An rRNA mutation identifies the apicoplast as the target for clindamycin in *Toxoplasma gondii*. *Mol Microbiol*. 2002;43:1309–1318.
- Yeh E, DeRisi JL. Chemical rescue of malaria parasites lacking an apicoplast defines organelle function in blood-stage *Plasmodium falciparum*. *PLoS Biol*. 2011;9:e1001138.
- Conley FK, Jenkins KA, Remington JS. *Toxoplasma gondii* infection of the central nervous system. Use of the peroxidase-antiperoxidase method to demonstrate *Toxoplasma* in formalin fixed, paraffin embedded tissue sections. *Hum Pathol*. 1981;12:690–698.
- Watts E, Zhao Y, Dhara A, et al. Novel approaches reveal that *Toxoplasma gondii* bradyzoites within tissue cysts are dynamic and replicating entities in vivo. *MBio*. 2015;6:e01155–15.
- Remington JS, Cavanaugh EN. Isolation of the encysted form of *Toxoplasma gondii* from human skeletal muscle and brain. *N Engl J Med*. 1965;273:1308–1310.
- Dubey J, Thayer D. Killing of different strains of *Toxoplasma gondii* tissue cysts by irradiation under defined conditions. *J Parasitol*. 1994;80:764–767.
- Dubey J, Kotula A, Sharar A, et al. Effect of high temperature on infectivity of *Toxoplasma gondii* tissue cysts in pork. *J Parasitol*. 1990;76:201.
- Jacobs L, Remington JS, Melton ML. The resistance of the encysted form of *Toxoplasma gondii*. *J Parasitol*. 1960;46:11–21.
- El-Nawawi FA, Tawfik MA, Shaapan RM. Methods for inactivation of *Toxoplasma gondii* cysts in meat and tissues of experimentally infected sheep. *Foodborne Pathog Dis*. 2008;5:687–690.
- Ferguson DJP, Graham DI, Hutchinson WM. Pathological changes in the brains of mice infected with *Toxoplasma gondii*: a histological immunocytochemical and ultrastructural study. *Int J Exp Pathol*. 1991;72:463–474.
- Sims TA, Hay J, Talbot IC. An electron microscope and immunohistochemical study of the intracellular location of *Toxoplasma* tissue cysts within the brains of mice with congenital toxoplasmosis. *Br J Exp Pathol*. 1989;70:317–325.
- Jones TC, Bienz KA, Erb P. In vitro cultivation of *Toxoplasma gondii* cysts in astrocytes in the presence of gamma interferon. *Infect Immun*. 1986;51:146–156.
- Ferguson DJ, Hutchinson WM, Pettersen E. Tissue cyst rupture in mice chronically infected with *Toxoplasma gondii*. *Parasitol Res*. 1989;75:599–603.
- Koshy AA, Dietrich HK, Christian DA, et al. *Toxoplasma* co-opts host cells it does not invade. *PLoS Pathog*. 2012;8:e1002825.
- Dubey JP, Jones JL. *Toxoplasma gondii* infection in humans and animals in the United States. *Int J Parasitol*. 2008;38:1257–1278.
- Neu HC. Toxoplasmosis transmitted at autopsy. *JAMA*. 1967;202:284–285.
- Dubey J. Toxoplasmosis. *Am Vet Med Assoc*. 1994;205:1593–1598.
- Bevins SN, Carver S, Boydston EE, et al. Three pathogens in sympatric populations of pumas, bobcats, and domestic cats: implications for infectious disease transmission. *PLoS ONE*. 2012;7:e31403.
- Desmonts G, Couvreur J, Alison F, et al. Epidemiological study on toxoplasmosis: the influence of cooking slaughter-animal meat on the incidence of human infection. *Rev Fr Etud Clin Biol*. 1965;10:952–958, [in French].
- Dattoli VC, Veiga RV, Cunha SS, et al. Oocyst ingestion as an important transmission route of *Toxoplasma gondii* in Brazilian urban children. *J Parasitol*. 2011;97:1080–1084.
- Hill D, Coss C, Dubey JP, et al. Identification of a sporozoite-specific antigen from *Toxoplasma gondii*. *J Parasitol*. 2011;97:328–337.
- Jones JL, Dubey JP. Foodborne toxoplasmosis. *Clin Infect Dis*. 2012;55:845–851.
- Hill DE, Dubey JP. *Toxoplasma gondii* prevalence in farm animals in the United States. *Int J Parasitol*. 2013;43:107–113.
- Hill D, Dubey JP. *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clin Microbiol Infect*. 2002;8:634–640.
- Remington JS. Toxoplasmosis and congenital infection. *Intrauterine Infect*. 1968;4:47–56.
- Dubey JP. Isolation of *Toxoplasma gondii* from a naturally infected beef cow. *J Parasitol*. 1992;78:151–153.
- Aspinall TV, Marlee D, Hyde JE, et al. Prevalence of *Toxoplasma gondii* in commercial meat products as monitored by polymerase chain reaction—food for thought? *Int J Parasitol*. 2002;32:1193–1199.
- Hill DE, Haley C, Wagner B, et al. Seroprevalence of and risk factors for *Toxoplasma gondii* in the US swine herd using sera collected during the National Animal Health Monitoring Survey (Swine 2006). *Zoonoses Public Health*. 2010;57:53–59.
- Dubey JP, Hill DE, Rozeboom DW, et al. High prevalence and genotypes of *Toxoplasma gondii* isolated from organic pigs in northern USA. *Vet Parasitol*. 2012;188:14–18.
- Dubey JP, Gamble HR, Hill D, et al. High prevalence of viable *Toxoplasma gondii* infection in market weight pigs from a farm in Massachusetts. *J Parasitol*. 2002;88:1234–1238.
- Dubey JP, Sundar N, Hill D, et al. High prevalence and abundant atypical genotypes of *Toxoplasma gondii* isolated from lambs destined for human consumption in the USA. *Int J Parasitol*. 2008;38:999–1006.
- Riemann HP, Meyer ME, Theis JH, et al. Toxoplasmosis in an infant fed unpasteurized goat milk. *J Pediatr*. 1975;87:573–576.
- Sacks JJ, Roberto RR, Brooks NF. Toxoplasmosis infection associated with raw goat's milk. *JAMA*. 1982;248:1728–1732.
- Contopoulos-Ioannidis D, Talucod J, Maldonado Y, et al. Seasonal variation of acute toxoplasmic lymphadenopathy in the United States. *Epidemiol Infect*. 2015;143:1893–1897.
- Morin L, Lobry JR, Peyron F, et al. Seasonal variations in acute toxoplasmosis in pregnant women in the Rhône-Alpes region (France). *Clin Microbiol Infect*. 2012;18:E401–E403.
- Dubey JP, Graham DH, Blackston CR, et al. Biological and genetic characterisation of *Toxoplasma gondii* isolates from chickens (*Gallus domesticus*) from São Paulo, Brazil: unexpected findings. *Int J Parasitol*. 2002;32:99–105.
- Bowie WR, King AS, Werker DH, et al. Outbreak of toxoplasmosis associated with municipal drinking water. *Lancet*. 1997;350:173–177.
- Luft BJ, Remington JS. Acute *Toxoplasma* infection among family members of patients with acute lymphadenopathic toxoplasmosis. *Arch Intern Med*. 1984;144:53–56.
- Masur H, Jones TC, Lempert JA, et al. Outbreak of toxoplasmosis in a family and documentation of acquired retinochoroiditis. *Am J Med*. 1978;64:396–402.
- Contopoulos-Ioannidis D, Wheeler KM, Ramirez R, et al. Clustering of *Toxoplasma gondii* infections within families of congenitally infected infants. *Clin Infect Dis*. 2015;61:1815–1824.
- Miller MA, Gardner IA, Kreuder C, et al. Coastal freshwater runoff is a risk factor for *Toxoplasma gondii*

- infection of southern sea otters (*Enhydra lutris nereis*). *Int J Parasitol.* 2002;32:997–1006.
81. Bahia-Oliveira LM, Jones JL, Azevedo-Silva J, et al. Highly endemic, waterborne toxoplasmosis in north Rio de Janeiro state, Brazil. *Emerg Infect Dis.* 2003;9:55–62.
  82. Baldrussón S, Karanis P. Waterborne transmission of protozoan parasites: review of worldwide outbreaks—an update 2004–2010. *Water Res.* 2011;45:6603–6614.
  83. Jones JL, Dubey JP. Waterborne toxoplasmosis—recent developments. *Exp Parasitol.* 2010;124:10–25.
  84. Dumetre A, Le Bras C, Baffet M, et al. Effects of ozone and ultraviolet radiation treatments on the infectivity of *Toxoplasma gondii* oocysts. *Vet Parasitol.* 2008;153:209–213.
  85. Lindsay DS, Collins MV, Mitchell SM, et al. Sporulation and survival of *Toxoplasma gondii* oocysts in seawater. *J Eukaryot Microbiol.* 2003;50(suppl):687–688.
  86. Wainwright KE, Miller MA, Barr BC, et al. Chemical inactivation of *Toxoplasma gondii* oocysts in water. *J Parasitol.* 2007;93:925–931.
  87. Vaudaux JD, Muccioli C, James ER, et al. Identification of an atypical strain of *Toxoplasma gondii* as the cause of a waterborne outbreak of toxoplasmosis in Santa Isabel do Ivaí, Brazil. *J Infect Dis.* 2010;202:1226–1233.
  88. Conrad PA, Miller MA, Kreuder C, et al. Transmission of *Toxoplasma*: clues from the study of sea otters as sentinels of *Toxoplasma gondii* flow into the marine environment. *Int J Parasitol.* 2005;35:1155–1168.
  89. Smith KL, Wilson M, Hightower AW, et al. Prevalence of *Toxoplasma gondii* antibodies in U.S. military recruits in 1989: comparison with data published in 1965. *Clin Infect Dis.* 1996;23:1182–1183.
  90. Rosso F, Les JT, Agudelo A, et al. Prevalence of infection with *Toxoplasma gondii* among pregnant women in Cali, Colombia, South America. *Am J Trop Med Hyg.* 2008;78:504–508.
  91. Jones JL, Kruszon-Moran D, Elder S, et al. *Toxoplasma gondii* Infection in the United States, 2011–2014. *Am J Trop Med Hyg.* 2018;98:551–557.
  92. Gargatá MJ, Ferreira I, Vilares A, et al. *Toxoplasma gondii* seroprevalence in the Portuguese population: comparison of three cross-sectional studies spanning three decades. *BMJ Open.* 2016;6:e011648.
  93. Foroutan-Rad M, Majidi H, Dalvand S, et al. Toxoplasmosis in blood donors: a systematic review and meta-analysis. *Transfus Med Rev.* 2016;30:116–122.
  94. Siegel SE, Lunde MN, Gelderman AH, et al. Transmission of toxoplasmosis by leukocyte transfusion. *Blood.* 1971;37:388–394.
  95. Ryning FW, McLeod R, Maddox JC, et al. Probable transmission of *Toxoplasma gondii* by organ transplantation. *Ann Intern Med.* 1979;90:47–49.
  96. Mele A, Paterson PJ, Prentice HG, et al. Toxoplasmosis in bone marrow transplantation: a report of two cases and systematic review of the literature. *Bone Marrow Transplant.* 2002;29:691–698.
  97. Slavin MA, Meyers JD, Remington JS, et al. *Toxoplasma gondii* infection in marrow transplant recipients: a 20 year experience. *Bone Marrow Transplant.* 1994;13:549–557.
  98. Lepot C, Chêne G, Morlat P, et al. Pyrimethamine for primary prophylaxis of toxoplasmic encephalitis in patients with human immunodeficiency virus infection: a double-blind, randomized trial. *J Infect Dis.* 1996;173:91–97.
  99. San-Andrés FJ, Rubio R, Castilla J, et al. Incidence of acquired immunodeficiency syndrome-associated opportunistic diseases and the effect of treatment on a cohort of 1115 patients infected with human immunodeficiency virus, 1989–1997. *Clin Infect Dis.* 2003;36:1177–1185.
  100. Luft BJ, Remington JS. Toxoplasmic encephalitis in AIDS (AIDS commentary). *Clin Infect Dis.* 1992;15:211–222.
  101. Abgrall S, Rabaud C, Costagliola D. Incidence and risk factors for toxoplasmic encephalitis in human immunodeficiency virus-infected patients before and during the highly active antiretroviral therapy era. *Clin Infect Dis.* 2001;33:1747–1755.
  102. Jones JL, Roberts JM. Toxoplasmosis hospitalizations in the United States, 2008, and trends, 1993–2008. *Clin Infect Dis.* 2012;54:e58–e61.
  103. Jones JL, Sehgal M, Maguire JH. Toxoplasmosis-associated deaths among human immunodeficiency virus-infected persons in the United States, 1992–1998. *Clin Infect Dis.* 2002;34:1161.
  104. Clumeck N. Some aspects of the epidemiology of toxoplasmosis and pneumocystosis in AIDS in Europe. *Eur J Clin Microbiol Infect Dis.* 1991;10:177–178.
  105. Zumla A, Savva D, Wheeler RB, et al. *Toxoplasma* serology in Zambian and Ugandan patients infected with the human immunodeficiency virus. *Trans R Soc Trop Med Hyg.* 1991;85:227–229.
  106. Belanger F, Derouin F, Grangeot-Keros L, et al. Incidence and risk factors of toxoplasmosis in a cohort of human immunodeficiency virus-infected patients: 1988–1995. HEMOCO and SEROCO Study Groups. *Clin Infect Dis.* 1999;28:575–581.
  107. Dankner WM, Lindsey JC, Levin MJ. Correlates of opportunistic infections in children infected with the human immunodeficiency virus managed before highly active antiretroviral therapy. *Pediatr Infect Dis.* 2001;20:40–48.
  108. Lucas S, Hounnou A, Peacock C, et al. The mortality and pathology of HIV infection in a West African city. *AIDS.* 1993;7:1569–1579.
  109. Saavedra A, Campinha-Bacote N, Hajjar M, et al. Causes of death and factors associated with early mortality of HIV-infected adults admitted to Korle-Bu Teaching Hospital. *Pan Afr Med J.* 2017;27:48.
  110. Wallace MR, Rossetti RJ, Olson PE. Cats and toxoplasmosis risk in HIV-infected adults. *JAMA.* 1993;269:76–77.
  111. Ruskin J, Remington JS. Toxoplasmosis in the compromised host. *Ann Intern Med.* 1976;84:193–199.
  112. Chintakuntlawar A, Kidd M, Al-Kali A, et al. Toxoplasmosis in patients with hematologic malignancies. *Leuk Lymphoma.* 2015;56:536–538.
  113. Jones TC, Yeh S, Hirsch JG. The interaction between *Toxoplasma gondii* and mammalian cells. I. Mechanism of entry and intracellular fate of the parasite. *J Exp Med.* 1972;136:1157–1172.
  114. Murray HW, Nathans CF, Cohn ZA. Macrophages oxygen-dependent antimicrobial activity. IV. Role of endogenous scavengers of oxygen intermediates. *J Exp Med.* 1980;152:1610–1624.
  115. Sibley LD, Weidner E, Krahenbuhl JL. Phagosome acidification blocked by intracellular *Toxoplasma gondii*. *Nature.* 1985;315:416–419.
  116. Adams LB, Hibbs JB Jr, Taintor RR, et al. Microbiostatic effect of murine-activated macrophages for *Toxoplasma gondii*. *J Immunol.* 1990;144:2725–2729.
  117. Pfefferkorn ER. Interferon gamma blocks the growth of *Toxoplasma gondii* in human fibroblasts by inducing the host cells to degrade tryptophan. *Proc Natl Acad Sci USA.* 1984;81:908–912.
  118. Schreiber RD, Feldman HA. Identification of the activator system for antibody to *Toxoplasma* as the classical complement pathway. *J Infect Dis.* 1980;141:366–369.
  119. Hunn JP, Feng CG, Sher A, et al. The immunity-related GTPases in mammals: a fast-evolving cell-autonomous resistance system against intracellular pathogens. *Mamm Genome.* 2011;22:43–54.
  120. Degrandi D, Konermann C, Beuter-Gunia C, et al. Extensive characterization of IFN-induced GTPases mGBP1 to mGBP10 involved in host defense. *J Immunol.* 2007;179:7729–7740.
  121. Remington JS, Jacobs L, Kaufman HE. Studies on chronic toxoplasmosis: the relation of infective dose to residual infection and to the possibility of congenital transmission. *Am J Ophthalmol.* 1958;46:261–267.
  122. Brown AS. Further evidence of infectious insults in the pathogenesis and pathophysiology of schizophrenia. *Am J Psychiatry.* 2011;168:764–766.
  123. Sugden K, Moffitt TE, Pinto L, et al. Is *Toxoplasma gondii* infection related to brain and behavior impairments in humans? evidence from a population-representative birth cohort. *PLoS ONE.* 2016;11:e0148435.
  124. Sutherland AL, Fond G, Kuin A, et al. Beyond the association. *Toxoplasma gondii* in schizophrenia, bipolar disorder, and addiction: systematic review and meta-analysis. *Acta Psychiatr Scand.* 2015;132:161–179.
  125. Hofflin JM, Conley FK, Remington JS. Murine model of intracerebral toxoplasmosis. *J Infect Dis.* 1987;155:550–557.
  126. Hofflin JM, Remington JS. Tissue culture isolation of *Toxoplasma* from blood of a patient with AIDS. *Arch Intern Med.* 1985;145:925–926.
  127. Tirard V, Niel G, Rosenheim M, et al. Diagnosis of toxoplasmosis in patients with AIDS by isolation of the parasite from the blood. *N Engl J Med.* 1991;324:632.
  128. Elbez-Rubinstein A, Ajzenberg D, Darde ML, et al. Congenital toxoplasmosis and reinfection during pregnancy: case report, strain characterization, experimental model of reinfection, and review. *J Infect Dis.* 2009;199:280–285.
  129. Hunter CA, Sibley LD. Modulation of innate immunity by *Toxoplasma gondii* virulence effectors. *Nat Rev Microbiol.* 2012;10:766–778.
  130. Denkers EY, Bizik DJ, Fox BA, et al. An inside job: hacking into Janus kinase/signal transducer and activator of transcription signaling cascades by the intracellular protozoan *Toxoplasma gondii*. *Infect Immun.* 2012;80:476–482.
  131. Reese ML, Boyle JP. Virulence without catalysis: how can a pseudokinase affect host cell signaling? *Trends Parasitol.* 2012;28:53–57.
  132. Howard JC, Hunn JP, Steinfeldt T. The IRG protein-based resistance mechanism in mice and its relation to virulence in *Toxoplasma gondii*. *Curr Opin Microbiol.* 2011;14:414–421.
  133. Rosowski EE, Lu D, Julien L, et al. Strain-specific activation of the NF-kappaB pathway by GRA15, a novel *Toxoplasma gondii* dense granule protein. *J Exp Med.* 2011;208:195–212.
  134. Ma JS, Sasai M, Ohshima J, et al. Selective and strain-specific NFAT4 activation by the *Toxoplasma gondii* polymorphic dense granule protein GRA6. *J Exp Med.* 2014;211:2013–2032.
  135. Bougdour A, Durandau E, Brenier-Pinchart MP, et al. Host cell subversion by *Toxoplasma* GRA16, an exported dense granule protein that targets the host cell nucleus and alters gene expression. *Cell Host Microbe.* 2013;13:489–500.
  136. Braun L, Brenier-Pinchart MP, Yogavel M, et al. A *Toxoplasma* dense granule protein, GRA24, modulates the early immune response to infection by promoting a direct and sustained host p38 MAPK activation. *J Exp Med.* 2013;210:2071–2086.
  137. Gay G, Braun L, Brenier-Pinchart MP, et al. *Toxoplasma gondii* TgIST co-opts host chromatin repressors dampening STAT1-dependent gene regulation and IFN-γ-mediated host defenses. *J Exp Med.* 2016;213:1779–1798.
  138. Olias P, Etheridge RD, Zhang Y, et al. *Toxoplasma* effector recruits the Mi-2/NuRD complex to repress STAT1 transcription and block IFN-γ-dependent gene expression. *Cell Host Microbe.* 2016;20:72–82.
  139. Franco M, Panas MW, Marino ND, et al. A novel secreted protein, MYR1, is central to *Toxoplasma*'s manipulation of host cells. *MBio.* 2016;7:e02231–15.
  140. Yamamoto M, Ma JS, Mueller C, et al. ATF6beta is a host cellular target of the *Toxoplasma gondii* virulence factor ROP18. *J Exp Med.* 2011;208:1533–1546.
  141. Gazzinelli RT, Mendonça-Neto R, Lihue J, et al. Innate resistance against *Toxoplasma gondii*: an evolutionary tale of mice, cats, and men. *Cell Host Microbe.* 2014;15:132–138.
  142. Witola WH, Mui E, Hargrave A, et al. NALP1 influences susceptibility to human congenital toxoplasmosis, proinflammatory cytokine response, and fate of *Toxoplasma gondii*-infected monocytes. *Infect Immun.* 2011;79:756–766.
  143. Tosh KW, Mittereder L, Bonne-Annee S, et al. The IL-12 Response of primary human dendritic cells and monocytes to *Toxoplasma gondii* is stimulated by phagocytosis of live parasites rather than host cell invasion. *J Immunol.* 2016;196:345–356.
  144. Hunter CA, Subauste C, Van Cleave V, et al. Production of gamma interferon by natural killer cells from *Toxoplasma gondii*-infected SCID mice: regulation by interleukin-10, interleukin-12, and tumor necrosis factor alpha. *Infect Immun.* 1994;62:2818–2824.
  145. Khan I, Matsuura T, Kasper L. Interleukin-12 enhances murine survival against acute toxoplasmosis. *Infect Immun.* 1994;62:1639–1642.
  146. Yap GS, Sher A. Cell-mediated immunity to *Toxoplasma gondii*: initiation, regulation and effector function. *Immunobiology.* 1999;201:240–247.
  147. Goldszmid RS, Caspar P, Rivollier A, et al. NK cell-derived interferon-gamma orchestrates cellular dynamics and the differentiation of monocytes into dendritic cells at the site of infection. *Immunity.* 2012;36:1047–1059.
  148. Dunay IR, Fuchs A, Sibley LD. Inflammatory monocytes but not neutrophils are necessary to control infection with *Toxoplasma gondii* in mice. *Infect Immun.* 2010;78:1564–1570.
  149. Dupont CD, Harms Pritchard G, Hidano S, et al. Flt3 ligand is essential for survival and protective immune responses during toxoplasmosis. *J Immunol.* 2015;195:4369–4377.
  150. Subauste CS, Wessendarp M, Sorensen RU, et al. CD40-CD40 ligand interaction is central to cell-mediated immunity against *Toxoplasma gondii*: patients with hyper IgM syndrome have a defective type 1 immune response that can be restored by soluble CD40 ligand trimer. *J Immunol.* 1999;162:6690–6700.
  151. Levy J, Espanol-Boren T, Thomas C, et al. Clinical spectrum of X-linked hyper-IgM syndrome. *J Pediatr.* 1997;131:47–54.
  152. Subauste CS, Wessendarp M, Smulian AG, et al. Role of CD40 ligand signaling in defective type 1 cytokine response in human immunodeficiency virus infection. *J Infect Dis.* 2001;183:1722–1731.
  153. Hunter CA, Remington JS. Immunopathogenesis of toxoplasmic encephalitis. *J Infect Dis.* 1994;170:1057–1067.
  154. Gazzinelli R, Hieny S, Wynn T, et al. Interleukin 12 is required for the T-lymphocyte-independent induction of



- interferon  $\gamma$  by an intracellular parasite and induces resistance in T-cell-deficient hosts. *Proc Natl Acad Sci USA*. 1993;90:6115–6119.
155. Gazzinelli R, Wysocka M, Hayashi S, et al. Parasite-induced IL-12 stimulates early IFN- $\gamma$  synthesis and resistance during acute infection with *Toxoplasma gondii*. *J Immunol*. 1994;153:2533–2543.
  156. Suzuki Y, Orellana MA, Schreiber RD, et al. Interferon- $\gamma$ : the major mediator of resistance against *Toxoplasma gondii*. *Science*. 1988;240:516–518.
  157. Suzuki Y, Conley FK, Remington JS. Importance of endogenous IFN- $\gamma$  for prevention of toxoplasmic encephalitis in mice. *J Immunol*. 1989;143:2045–2050.
  158. Suzuki Y, Conley FK, Remington JS. Treatment of toxoplasmic encephalitis in mice with recombinant gamma interferon. *Infect Immun*. 1990;58:3050–3055.
  159. Robben PM, Mordue DG, Truscott SM, et al. Production of IL-12 by macrophages infected with *Toxoplasma gondii* depends on the parasite genotype. *J Immunol*. 2004;172:3686–3694.
  160. Saeij JP, Collier S, Boyle JP, et al. *Toxoplasma* co-opts host gene expression by injection of a polymorphic kinase homologue. *Nature*. 2007;445:324–327.
  161. Sibley LD, Adams LB, Fukutomi Y, et al. Tumor necrosis factor- $\alpha$  triggers antitoxoplasmic activity of IFN- $\gamma$  primed macrophages. *J Immunol*. 1991;147:2340–2345.
  162. Langermans J, van der Hulst M, Nibbering P, et al. IFN- $\gamma$  induced L-arginine-dependent *Toxoplasma* static activity in murine peritoneal macrophages is mediated by endogenous tumor necrosis factor. *J Immunol*. 1992;148:568–574.
  163. Chang H, Peckere J, Piguat P. Role of tumour necrosis factor in chronic murine *Toxoplasma gondii* encephalitis. *Immunol Infect Dis*. 1992;2:61–68.
  164. Gazzinelli RT, Wysocka M, Hieny S, et al. In the absence of endogenous IL-10, mice acutely infected with *Toxoplasma gondii* succumb to a lethal immune response dependent on CD4<sup>+</sup> T cells and accompanied by overproduction of IL-12, IFN-gamma and TNF-alpha. *J Immunol*. 1996;157:798–805.
  165. Suzuki Y, Yang Q, Yang S, et al. IL-4 is protective against development of toxoplasmic encephalitis. *J Immunol*. 1996;157:2564–2569.
  166. Suzuki Y, Rani S, Liesenfeld O, et al. Impaired resistance to the development of toxoplasmic encephalitis in interleukin-6-deficient mice. *Infect Immun*. 1997;65:2339–2345.
  167. Kasper LH, Matsuura T, Khan IA. IL-7 stimulates protective immunity in mice against the intracellular pathogen, *Toxoplasma gondii*. *J Immunol*. 1995;155:4798–4804.
  168. Hunter CA, Neyer LA, Gabriel KE, et al. The role of CD28/B7 interaction in the regulation of NK cell responses during infection with *Toxoplasma gondii*. *J Immunol*. 1997;158:2285–2293.
  169. Kelly MN, Kolls JK, Happel K, et al. Interleukin-17/interleukin-17 receptor-mediated signaling is important for generation of an optimal polymorphonuclear response against *Toxoplasma gondii* infection. *Infect Immun*. 2005;73:617–621.
  170. Lieberman LA, Cardillo F, Owyang AM, et al. IL-23 provides a limited mechanism of resistance to acute toxoplasmosis in the absence of IL-12. *J Immunol*. 2004;173:1887–1893.
  171. Stumhofer JS, Laurence A, Wilson EH, et al. Interleukin 27 negatively regulates the development of interleukin 17-producing T helper cells during chronic inflammation of the central nervous system. *Nat Immunol*. 2006;7:937–945.
  172. Böhne W, Hesseemann J, Gross U. Reduced replication of *Toxoplasma gondii* is necessary for induction of bradyzoite-specific antigens: a possible role for nitric oxide in triggering stage conversion. *Infect Immun*. 1994;62:1761–1767.
  173. Chao C, Gekker G, Hu S, et al. Human microglial cell defense against *Toxoplasma gondii*. *J Immunol*. 1994;152:1246–1252.
  174. Khan IA, Matssuura T, Fonseka S, et al. Production of nitric oxide (NO) is not essential for protection against acute *Toxoplasma gondii* infection in IRF-1<sup>-/-</sup> mice. *J Immunol*. 1996;156:636–643.
  175. Schariton-Kersten TM, Wynn TA, Denkers EY, et al. In the absence of endogenous IFN- $\gamma$  mice develop unimpaired IL-12 responses to *Toxoplasma gondii* while failing to control acute infection. *J Immunol*. 1996;157:4045–4054.
  176. Hayashi S, Chan C, Gazzinelli R, et al. Contribution of nitric oxide to the host parasite equilibrium in toxoplasmosis. *J Immunol*. 1996;156:1476–1481.
  177. Schariton-Kersten TM, Yap G, Magram J, et al. Inducible nitric oxide is essential for host control of persistent but not acute infection with the intracellular pathogen *Toxoplasma gondii*. *J Exp Med*. 1997;185:1261–1273.
  178. Taylor GA, Feng CG, Sher A. Control of IFN-gamma-mediated host resistance to intracellular pathogens by immunity-related GTPases (p47 GTPases). *Microbes Infect*. 2007;9:1644–1651.
  179. Martens S, Parvanova I, Zerrahn J, et al. Disruption of *Toxoplasma gondii* parasitophorous vacuoles by the mouse p47-resistance GTPases. *PLoS Pathog*. 2005;1:e24.
  180. Ling YM, Shaw MH, Ayala C, et al. Vacuolar and plasma membrane stripping and autophagic elimination of *Toxoplasma gondii* in primed effector macrophages. *J Exp Med*. 2006;203:2063–2071.
  181. Pomares C, Holmes TH, Estran R, et al. Cytokine profiles in patients with toxoplasmic lymphadenitis in the setting of pregnancy. *Cytokine*. 2017;90:14–20.
  182. Pavia CS. Protection against experimental toxoplasmosis by adoptive immunotherapy. *J Immunol*. 1986;137:2985–2990.
  183. Peterson P, Gekker G, Hu S, et al. Human astrocytes inhibit intracellular multiplication of *Toxoplasma gondii* by a nitric oxide-mediated mechanism. *J Infect Dis*. 1995;171:516–518.
  184. Brown CR, McLeod R. Class I MHC genes and CD8<sup>+</sup> T cells determine cyst number in *Toxoplasma gondii* infection. *J Immunol*. 1990;145:3438–3441.
  185. Brown C, Hunter C, Estes R, et al. Definitive identification of a gene that confers resistance against *Toxoplasma* cyst burden and encephalitis. *Immunology*. 1995;85:419–428.
  186. Suzuki Y, Joh K, Orellana MA, et al. A gene(s) within the H-2D region determines the development of toxoplasmic encephalitis in mice. *Immunology*. 1991;74:732–739.
  187. Suzuki Y, Wong S-Y, Grumet FC, et al. Evidence for genetic regulation of susceptibility to toxoplasmic encephalitis in AIDS patients. *J Infect Dis*. 1996;173:265–268.
  188. Mack D, Johnson J, Roberts F, et al. HLA-class II genes modify outcome of *Toxoplasma gondii* infection. *Int J Parasitol*. 1999;29:1351–1358.
  189. Habegger de Sorrentino A, Lopez R, Motta P, et al. HLA class II involvement in HIV-associated toxoplasmic encephalitis development. *Clin Immunol*. 2005;115:133–137.
  190. Shimokawa PT, Targa LS, Yamamoto L, et al. HLA-DQA1/B1 alleles as putative susceptibility markers in congenital toxoplasmosis. *Virulence*. 2016;7:456–464.
  191. Ayo CM, Frederico FB, Siqueira RC, et al. Ocular toxoplasmosis: susceptibility in respect to the genes encoding the KIR receptors and their HLA class I ligands. *Sci Rep*. 2016;6:36632.
  192. Cordeiro CA, Moreira PR, Andrade MS, et al. Interleukin-10 gene polymorphism (-1082G/A) is associated with toxoplasmic retinochoroiditis. *Invest Ophthalmol Vis Sci*. 2008;49:1979–1982.
  193. Jamieson SE, Peixoto-Rangel AL, Hargrave AC, et al. Evidence for associations between the purinergic receptor P2X7 (P2RX7) and toxoplasmosis. *Genes Immun*. 2010;11:374–383.
  194. Peixoto-Rangel AL, Miller EN, Castellucci L, et al. Candidate gene analysis of ocular toxoplasmosis in Brazil: evidence for a role for toll-like receptor 9 (TLR9). *Mem Inst Oswaldo Cruz*. 2009;104:1187–1190.
  195. Cordeiro CA, Moreira PR, Bessa TF, et al. Interleukin-6 gene polymorphism (-174 G/C) is associated with toxoplasmic retinochoroiditis. *Acta Ophthalmol*. 2013;91:e311–e314.
  196. Bustamante J, Boisson-Dupuis S, Abel L, et al. Mendelian susceptibility to mycobacterial disease: genetic, immunological, and clinical features of inborn errors of IFN- $\gamma$  immunity. *Semin Immunol*. 2014;26:454–470.
  197. Dorfman RF, Remington JS. Value of lymph-node biopsy in the diagnosis of acute acquired toxoplasmosis. *N Engl J Med*. 1973;289:878–881.
  198. Montoya JG, Jordan R, Lingamneni S, et al. Toxoplasmic myocarditis and polyomysitis in patients with acute acquired toxoplasmosis diagnosed during life. *Clin Infect Dis*. 1997;24:676–683.
  199. Negri Gualdi C, Negriagualdi C. Infectious pseudomononucleosis in lymphomatous diseases. *Minerva Med*. 1974;65:78–83, [in Italian].
  200. Weiss L, Chen Y, Berry G, et al. Infrequent detection of *Toxoplasma gondii* genome in toxoplasmic lymphadenitis: a polymerase chain reaction study. *Hum Pathol*. 1992;23:154–158.
  201. Montoya JG, Remington JS. Toxoplasmosis of the central nervous system. In: Peterson PK, Remington JS, eds. *In Defense of the Brain: Current Concepts in the Immunopathogenesis and Clinical Aspects of CNS Infections*. Boston: Blackwell Scientific; 1997:163–188.
  202. Frenkel JK. Pathology and pathogenesis of congenital toxoplasmosis. *Bull N Y Acad Med*. 1974;50:182–191.
  203. Ferguson DJ, Bowker C, Jeffery KJ, et al. Congenital toxoplasmosis: continued parasite proliferation in the fetal brain despite maternal immunological control in other tissues. *Clin Infect Dis*. 2013;56:204–208.
  204. Wong B, Gold JWM, Brown AE, et al. Central-nervous-system toxoplasmosis in homosexual men and parenteral drug abusers. *Ann Intern Med*. 1984;100:36–42.
  205. Post MJ, Chan JG, Hensley GT, et al. Toxoplasmosis encephalitis in Haitian adults with acquired immunodeficiency syndrome: a clinical-pathologic-CT correlation. *AJNR Am J Neuroradiol*. 1983;4:155–162.
  206. Strittmatter C, Lang W, Wiestler OD, et al. The changing pattern of human immunodeficiency virus associated cerebral toxoplasmosis: a study of 46 postmortem cases. *Acta Neuropathol*. 1992;83:475–481.
  207. Parmley SF, Goebel FD, Remington JS. Detection of *Toxoplasma gondii* DNA in cerebrospinal fluid from AIDS patients by polymerase chain reaction. *J Clin Microbiol*. 1992;30:3000–3002.
  208. Gray F, Gherardi R, Wingate E, et al. Diffuse “encephalitic” cerebral toxoplasmosis in AIDS. *J Neurol*. 1989;236:273–277.
  209. Mariuz P, Bosler EM, Luft BJ. *Toxoplasma* pneumonia. *Semin Respir Infect*. 1997;12:40–43.
  210. Bretagne S, Costa J-M, Fleury-Feith J, et al. Quantitative competitive PCR with bronchoalveolar lavage fluid for diagnosis of toxoplasmosis in AIDS patients. *J Clin Microbiol*. 1995;33:1662–1664.
  211. Holland GN, O'Connor GR, Belfort R Jr, et al. Toxoplasmosis. In: Pepose JS, Holland GN, Wilhelmus KR, eds. *Ocular Infection and Immunity*. St Louis: Mosby Yearbook; 1996:1183–1223.
  212. Danise A, Cinque P, Vergani S, et al. Use of the polymerase chain reaction assays of aqueous humor in the differential diagnosis of retinitis in patients infected with human immunodeficiency virus. *Clin Infect Dis*. 1997;24:1100–1106.
  213. Montoya JG, Parmley S, Liesenfeld O, et al. Use of the polymerase chain reaction for diagnosis of ocular toxoplasmosis. *Ophthalmology*. 1999;106:1554–1563.
  214. Silveira C, Belfort R Jr, Muccioli C, et al. The effect of long-term intermittent trimethoprim/sulfamethoxazole treatment on recurrences of toxoplasmic retinochoroiditis. *Am J Ophthalmol*. 2002;134:41–46.
  215. Jones JL, Muccioli C, Belfort R Jr, et al. Recently acquired *Toxoplasma gondii* infection, Brazil. *Emerg Infect Dis*. 2006;12:582–587.
  216. Gilbert RE, Freeman K, Lago EG, et al. Ocular sequelae of congenital toxoplasmosis in Brazil compared with Europe. *PLoS Negl Trop Dis*. 2008;2:e277.
  217. de-la-Torre A, Lopez-Castillo CA, Gomez-Marin JE. Incidence and clinical characteristics in a Colombian cohort of ocular toxoplasmosis. *Eye*. 2009;23:1090–1093.
  218. Vallochi AL, Muccioli C, Martins MC, et al. The genotype of *Toxoplasma gondii* strains causing ocular toxoplasmosis in humans in Brazil. *Am J Ophthalmol*. 2005;139:350–351.
  219. Grigg ME, Ganatra J, Boothroyd JC, et al. Unusual abundance of atypical strains associated with human ocular toxoplasmosis. *J Infect Dis*. 2001;184:633–639.
  220. Gherardi R, Baudrimont M, Lionnet F, et al. Skeletal muscle toxoplasmosis in patients with acquired immunodeficiency syndrome: a clinical and pathological study. *Ann Neurol*. 1992;32:535–542.
  221. Calico I, Caballero E, Martinez O, et al. Isolation of *Toxoplasma gondii* from immunocompromised patients using tissue culture. *Infection*. 1991;19:340–342.
  222. Roldan EO, Moskowitz L, Hensley GT. Pathology of the heart in acquired immunodeficiency syndrome. *Arch Pathol Lab Med*. 1987;111:943–946.
  223. Adair OV, Randive N, Krasnow N. Isolated *Toxoplasma* myocarditis in acquired immune deficiency syndrome. *Am Heart J*. 1989;118:856–857.
  224. Pauwels A, Meyohas MC, Eliazewicz M, et al. *Toxoplasma* colitis in the acquired immunodeficiency syndrome. *Am J Gastroenterol*. 1992;87:518–519.
  225. Smart PE, Weinfeld A, Thompson NE, et al. Toxoplasmosis of the stomach: a cause of antral narrowing. *Radiology*. 1990;174:369–370.
  226. Garcia LW, Hemphill RB, Marasco WA, et al. Acquired immunodeficiency syndrome with disseminated toxoplasmosis presenting as an acute pulmonary and gastrointestinal illness. *Arch Pathol Lab Med*. 1991;115:459–463.
  227. Brion J-P, Pelloux H, Le Marc'hadour F, et al. Acute toxoplasmic hepatitis in a patient with AIDS. *Clin Infect Dis*. 1992;15:183–184.
  228. Bergin C, Murphy M, Lyons D, et al. *Toxoplasma* pneumonitis: fatal presentation of disseminated toxoplasmosis in a patient with AIDS. *Eur Respir J*. 1992;5:1018–1020.
  229. Crider SR, Horstman WG, Massey GS. *Toxoplasma* orchitis: report of a case and a review of the literature. *Am J Med*. 1988;85:421–424.



230. Groll A, Schneider M, Althoff PH, et al. Morphology and clinical significance of AIDS-related lesions in the adrenals and pituitary. *Dtsch Med Wschr.* 1990;115:483–488.
231. Patrick AL, Roberts LA, Burton EN, et al. Focal and segmental glomerulosclerosis in the acquired immunodeficiency syndrome. *West Indian Med J.* 1986;1986:200–202.
232. Renold C, Sugar A, Chave J-P, et al. *Toxoplasma* encephalitis in patients with the acquired immunodeficiency syndrome. *Medicine (Baltimore).* 1992;71:224–239.
233. Remington JS. Toxoplasmosis in the adult. *Bull N Y Acad Med.* 1974;50:211–227.
234. Demar M, Aizenberg D, Maubon D, et al. Fatal outbreak of human toxoplasmosis along the Maroni River: epidemiological, clinical, and parasitological aspects. *Clin Infect Dis.* 2007;45:e88–e95.
235. Demar M, Hommel D, Djossou F, et al. Acute toxoplasmoses in immunocompetent patients hospitalized in an intensive care unit in French Guiana. *Clin Microbiol Infect.* 2012;18:E221–E231.
236. Anand R, Jones CW, Ricks JH, et al. Acute primary toxoplasmosis in travelers returning from endemic countries. *J Travel Med.* 2012;19:57–60.
237. McCabe RE, Brooks RG, Dorfman RF, et al. Clinical spectrum in 107 cases of toxoplasmic lymphadenopathy. *Rev Infect Dis.* 1987;9:754–774.
238. Remington JS, Barnett CG, Meikel M, et al. Toxoplasmosis and infectious mononucleosis. *Arch Intern Med.* 1962;110:744–753.
239. Wong WK, Upton A, Thomas MG. Neuropsychiatric symptoms are common in immunocompetent adult patients with *Toxoplasma gondii* acute lymphadenitis. *Scand J Infect Dis.* 2013;45:357–361.
240. Couvreur J, Thulliez P. Acquired toxoplasmosis with ocular or neurologic involvement. *Presse Med.* 1996;25:438–442. [in French].
241. Montoya JG, Remington JS. Toxoplasmic chorioretinitis in the setting of acute acquired toxoplasmosis. *Clin Infect Dis.* 1996;23:277–282.
242. Delair E, Monnet D, Grabar S, et al. Respective roles of acquired and congenital infections in presumed ocular toxoplasmosis. *Am J Ophthalmol.* 2008;146:851–855.
243. Burnett AJ, Shortt SG, Isaac-Renton J, et al. Multiple cases of acquired toxoplasmosis retinitis presenting in an outbreak. *Ophthalmology.* 1998;105:1032–1037.
244. Toporovski J, Romano S, Hartmann S, et al. Nephrotic syndrome associated with toxoplasmosis: report of seven cases. *Rev Inst Med Trop São Paulo.* 2012;54:61–64.
245. Montoya JG, Remington JS. Studies on the serodiagnosis of toxoplasmic lymphadenitis. *Clin Infect Dis.* 1995;20:781–790.
246. Cunningham T. Pancarditis in acute toxoplasmosis. *Am J Clin Pathol.* 1982;78:403–405.
247. Prado SP, Pacheco VC, Noemi IH, et al. *Toxoplasma* pericarditis and myocarditis. *Rev Chil Pediatr.* 1978;49:179–185. [in Spanish].
248. Pergola G, Cascone A, Russo M. Acute pericarditis and myocarditis by *Toxoplasma gondii* in an immunocompetent young man: a case report. *Infez Med.* 2010;18:48–52.
249. Greenlee JE, Johnson WD Jr, Campa JF, et al. Adult toxoplasmosis presenting as polymyositis and cerebellar ataxia. *Ann Intern Med.* 1975;82:367–371.
250. Paspalaki PK, Mihailidou EP, Bitsori M, et al. Polymyositis and myocarditis associated with acquired toxoplasmosis in an immunocompetent girl. *BMC Musculoskelet Disord.* 2001;2:8.
251. Palma S, Reyes H, Guzman L, et al. Dermatomyositis and toxoplasmosis. *Rev Med Chil.* 1984;111:164–167. [in Spanish].
252. Pollock JL. Toxoplasmosis appearing to be dermatomyositis. *Arch Dermatol.* 1979;115:736–737.
253. Brown AS, Schaefer CA, Quesenberry CP Jr, et al. Maternal exposure to toxoplasmosis and risk of schizophrenia in adult offspring. *Am J Psychiatry.* 2005;162:767–773.
254. Niebuhr DW, Millikan AM, Cowan DN, et al. Selected infectious agents and risk of schizophrenia among U.S. military personnel. *Am J Psychiatry.* 2008;165:99–106.
255. Torrey EF, Bartko JJ, Yolken RH. *Toxoplasma gondii* and other risk factors for schizophrenia: an update. *Schizophr Bull.* 2012;38:642–647.
256. Safa G, Darrieux L. Cerebral toxoplasmosis after rituximab therapy. *JAMA Intern Med.* 2013;173:924–926.
257. Stive O, Wiendl H. Iatrogenic immunosuppression with biologics in MS: expecting the unexpected? *Neurology.* 2009;73:1346–1347.
258. Lassoued S, Zabraniecki L, Marin F, et al. Toxoplasmic chorioretinitis and antitumor necrosis factor treatment in rheumatoid arthritis. *Semin Arthritis Rheum.* 2007;36:262–263.
259. Montoya JG, Gomez CA. Toxoplasmosis after solid organ transplantation. In: Ljungman P, Snyderman D, Boeckh M, eds. *Transplant Infections.* New York: Springer; 2016.
260. United Network for Organ Sharing. Toxoplasmosis Testing of All Donors Is Now Required. April 6, 2017. <https://transplantpro.org/news/opos/coming-soon-opos-should-plan-for-required-toxoplasmosis-testing-of-all-donors/>. Accessed November 16, 2018.
261. Luft BJ, Naot Y, Araujo FG, et al. Primary and reactivated *Toxoplasma* infection in patients with cardiac transplants. Clinical spectrum and problems in diagnosis in a defined population. *Ann Intern Med.* 1983;99:27–31.
262. Fernandez-Sabe N, Cervera C, Farinas MC, et al. Risk factors, clinical features, and outcomes of toxoplasmosis in solid-organ transplant recipients: a matched case-control study. *Clin Infect Dis.* 2012;54:355–361.
263. Montoya JG, Giraldo LF, Efron B, et al. Infectious complications among 620 consecutive heart transplant patients at Stanford University Medical Center. *Clin Infect Dis.* 2001;33:629–640.
264. Wreghitt TG, Gray JJ, Pavel P, et al. Efficacy of pyrimethamine for the prevention of donor-acquired *Toxoplasma gondii* infection in heart and heart-lung transplant patients. *Transplant Int.* 1992;5:197–200.
265. Wreghitt TG, McNeil K, Roth C, et al. Antibiotic prophylaxis for the prevention of donor-acquired *Toxoplasma gondii* infection in transplant patients [letter; comment]. *J Infect.* 1995;31:253–254.
266. Dávila V, Roncancio-Villamil G, Correa LA, et al. Disseminated toxoplasmosis in a heart transplant patient despite co-trimoxazole prophylaxis: a case report. *Biomedica.* 2017;37:303–307.
267. Renoult E, Georges E, Biava MF, et al. Toxoplasmosis in kidney transplant recipients: report of six cases and review. *Clin Infect Dis.* 1997;24:625–634.
268. Hamza A, Anderson IJ, Al-Khafaji B. Respiratory distress of unknown etiology in a transplant recipient: think toxoplasmosis! *Autops Case Rep.* 2017;7:37–41.
269. Filloy A, Garcia-Garcia O, Fernandez-Lorente L. Chorioretinitis as the first sign of acquired toxoplasmosis transmitted from donor following kidney transplantation: case report and review of the literature. *Ocul Immunol Inflamm.* 2013;21:27–28.
270. Assi MA, Rosenblatt JE, Marshall WF. Donor-transmitted toxoplasmosis in liver transplant recipients: a case report and literature review. *Transpl Infect Dis.* 2007;9:132–136.
271. Miltgen G, Paillot J, Vanlemmens C, et al. Fatal acute respiratory distress by *Toxoplasma gondii* in a *Toxoplasma* seronegative liver transplant recipient. *New Microbiol.* 2016;39:160–162.
272. Singer MA, Hagler WS, Grossniklaus HE. *Toxoplasma gondii* retinochoroiditis after liver transplantation. *Retina.* 1993;13:40–45.
273. Botterel F, Ichai P, Feray C, et al. Disseminated toxoplasmosis, resulting from infection of allograft, after orthotopic liver transplantation: usefulness of quantitative PCR. *J Clin Microbiol.* 2002;40:1648–1650.
274. Martino R, Bretagne S, Rovira M, et al. Toxoplasmosis after hematopoietic stem cell transplantation. Report of a 5-year survey from the Infectious Diseases Working Party of the European Group for Blood and Marrow Transplantation. *Bone Marrow Transplant.* 2000;25:1111–1114.
275. Schuster MG, Cleveland AA, Dubberke ER, et al. Infections in hematopoietic cell transplant recipients: results from the Organ Transplant Infection Project, a multicenter, prospective, cohort study. *Open Forum Infect Dis.* 2017;4:ofx050.
276. Conrad A, Le Maréchal M, Dupont D, et al. A matched case-control study of toxoplasmosis after allogeneic hematopoietic stem cell transplantation: still a devastating complication. *Clin Microbiol Infect.* 2016;22:636–641.
277. Decembrino N, Comelli A, Genco F, et al. Toxoplasmosis disease in paediatric hematopoietic stem cell transplantation: do not forget it still exists. *Bone Marrow Transplant.* 2017;52:1326–1329.
278. Martino R, Bretagne S, Einsele H, et al. Early detection of *Toxoplasma* infection by molecular monitoring of *Toxoplasma gondii* in peripheral blood samples after allogeneic stem cell transplantation. *Clin Infect Dis.* 2005;40:67–78.
279. Matsuo Y, Takeishi S, Miyamoto T, et al. Toxoplasmosis encephalitis following severe graft-vs.-host disease after allogeneic hematopoietic stem cell transplantation: 17 yr experience in Fukuoka BMT group. *Eur J Haematol.* 2007;79:317–321.
280. Caner A, Donmez A, Doskaya M, et al. Determining *Toxoplasma* high-risk autologous and allogeneic hematopoietic stem cell transplantation patients by systematic pre-transplant PCR screening of stem cell originated buffy coat. *Parasitology Int.* 2012;61:565–571.
281. Gajurel K, Dhakal R, Montoya JG. *Toxoplasma* prophylaxis in hematopoietic cell transplant recipients: a review of the literature and recommendations. *Curr Opin Infect Dis.* 2015;28:283–292.
282. Sing A, Leitritz L, Roggenkamp A, et al. Pulmonary toxoplasmosis in bone marrow transplant recipients: report of two cases and review. *Clin Infect Dis.* 1999;29:429–433.
283. Sumi M, Norose K, Hikosaka K, et al. Clinical characteristics and computed tomography findings of pulmonary toxoplasmosis after hematopoietic stem cell transplantation. *Int J Hematol.* 2016;104:729–740.
284. Liesenfeld O, Wong SY, Remington JS. Toxoplasmosis in the setting of AIDS. In: Bartlett JG, Merigan TC, Bolognesi D, eds. *Textbook of AIDS Medicine.* 2nd ed. Baltimore: Williams & Wilkins; 1999:225–259.
285. Oksenhender E, Cadranet J, Sarfati C, et al. *Toxoplasma gondii* pneumonia in patients with the acquired immunodeficiency syndrome. *Am J Med.* 1990;88(5N):18N–21N.
286. Mehren M, Burns PJ, Mamani F, et al. Toxoplasmic myelitis mimicking intramedullary spinal cord tumor. *Neurology.* 1988;38:1648–1650.
287. Herskovitz S, Siegel SE, Schneider AT, et al. Spinal cord toxoplasmosis in AIDS. *Neurology.* 1989;39:1552–1553.
288. Overhage JM, Greist A, Brown DR. Conus medullaris syndrome resulting from *Toxoplasma gondii* infection in a patient with the acquired immunodeficiency syndrome. *Am J Med.* 1990;89:814–815.
289. Rabaud C, May T, Lucet JC, et al. Pulmonary toxoplasmosis in patients infected with human immunodeficiency virus: a French national study. *Clin Infect Dis.* 1996;23:1249–1254.
290. Derouin F, Sarfati C, Beauvais B, et al. Prevalence of pulmonary toxoplasmosis in HIV-infected patients. *AIDS.* 1990;4:1036.
291. Holland G, Engstrom R Jr, Glasgow B, et al. Ocular toxoplasmosis in patients with acquired immunodeficiency syndrome. *Am J Ophthalmol.* 1988;106:653–667.
292. Heinemann MH, Gold JM, Maisel J. Bilateral *Toxoplasma* retinochoroiditis in a patient with acquired immune deficiency syndrome. *Retina.* 1986;6:224–227.
293. Louie JK, Hsu LC, Osmond DH, et al. Trends in causes of death among persons with acquired immunodeficiency syndrome in the era of highly active antiretroviral therapy. San Francisco, 1994–1998. *J Infect Dis.* 2002;186:1023–1027.
294. Antinori A, Larussa D, Cingolani A, et al. Prevalence, associated factors, and prognostic determinants of AIDS-related toxoplasmic encephalitis in the era of advanced highly active antiretroviral therapy. *Clin Infect Dis.* 2004;39:1681–1691.
295. Ghosn J, Paris L, Aizenberg D, et al. Atypical toxoplasmic manifestation after discontinuation of maintenance therapy in a human immunodeficiency virus type 1-infected patient with immune recovery. *Clin Infect Dis.* 2003;37:e112–e114.
296. Sendi P, Sachers F, Drechsler H, et al. Immune recovery vitritis in an HIV patient with isolated toxoplasmic retinochoroiditis. *AIDS.* 2006;20:2237–2238.
297. Martin-Blondel G, Alvarez M, Delobel P, et al. Toxoplasmic encephalitis IRIS in HIV-infected patients: a case series and review of the literature. *J Neurol Neurosurg Psychiatry.* 2011;82:691–693.
298. Pfeffer G, Prout A, Hooge J, et al. Biopsy-proven immune reconstitution syndrome in a patient with AIDS and cerebral toxoplasmosis. *Neurology.* 2009;73:321–322.
299. van Bilsen WPH, van den Berg CHSB, Rijnders BJA, et al. Immune reconstitution inflammatory syndrome associated with toxoplasmic encephalitis in HIV-infected patients. *AIDS.* 2017;31:1415–1424.
300. Furrer H, Opravil M, Bernasconi E, et al. Stopping primary prophylaxis in HIV-1-infected patients at high risk of *Toxoplasma* encephalitis. Swiss HIV Cohort Study. *Lancet.* 2000;355:2217–2218.
301. Kirk O, Reiss P, Uberti-Foppa C, et al. Safe interruption of maintenance therapy against previous infection with four common HIV-associated opportunistic pathogens during potent antiretroviral therapy. *Ann Intern Med.* 2002;137:239–250.
302. Miro JM, Lopez JC, Podzamczar D, et al. Discontinuation of primary and secondary *Toxoplasma gondii* prophylaxis is safe in HIV-infected patients after immunological restoration with highly active antiretroviral therapy: results of an open, randomized, multicenter clinical trial. *Clin Infect Dis.* 2006;43:79–89.
303. Stout JE, Lai JC, Giner J, et al. Reactivation of retinal toxoplasmosis despite evidence of immune response to highly active antiretroviral therapy. *Clin Infect Dis.* 2002;35:e37–e39.
304. Miro J, Esteve A, Furrer H, et al. Safety of stopping primary *T. gondii* prophylaxis with suppressed viremia

- and CD4>100. Conference on Retroviruses and Opportunistic Infections. February 22–25, 2016; Boston, Massachusetts.
305. Khan A, Su C, German M, et al. Genotyping of *Toxoplasma gondii* strains from immunocompromised patients reveals high prevalence of type I strains. *J Clin Microbiol*. 2005;43:5881–5887.
  306. Holland GN. Reconsidering the pathogenesis of ocular toxoplasmosis. *Am J Ophthalmol*. 1999;128:502–505.
  307. London NJ, Hovakimyan A, Cubillan LD, et al. Prevalence, clinical characteristics, and causes of vision loss in patients with ocular toxoplasmosis. *Eur J Ophthalmol*. 2011;21:811–819.
  308. Silveira C, Belfort R Jr, Burnier M Jr, et al. Acquired toxoplasmic infection as the cause of toxoplasmic retinochoroiditis in families. *Am J Ophthalmol*. 1988;106:362–364.
  309. Glasner PD, Silveira C, Kruszon-Moran D, et al. An unusually high prevalence of ocular toxoplasmosis in Southern Brazil. *Am J Ophthalmol*. 1992;114:136–144.
  310. Silveira C, Belfort R Jr, Muccioli C, et al. A follow-up study of *Toxoplasma gondii* infection in southern Brazil. *Am J Ophthalmol*. 2001;131:351–354.
  311. Nussenblatt R, Belfort R Jr. Ocular toxoplasmosis. *JAMA*. 1994;271:302–307.
  312. Bosch-Driessen LE, Berendschot TT, Ongkosuwo JV, et al. Ocular toxoplasmosis: clinical features and prognosis of 154 patients. *Ophthalmology*. 2002;109:869–878.
  313. Bosch-Driessen LH, Plaisier MB, Stijlma JS, et al. Reactivations of ocular toxoplasmosis after cataract extraction. *Ophthalmology*. 2002;109:41–45.
  314. Holland GN, Crespi CM, ten Dam-van Loon N, et al. Analysis of recurrence patterns associated with toxoplasmic retinochoroiditis. *Am J Ophthalmol*. 2008;145:1007–1013.
  315. Jones JL, Bonetti V, Hollan GN, et al. Ocular toxoplasmosis in the United States: recent and remote infections. *Clin Infect Dis*. 2015;60:271–273.
  316. Delair E, Latkany P, Noble AG, et al. Clinical manifestations of ocular toxoplasmosis. *Ocul Immunol Inflamm*. 2011;19:91–102.
  317. Aleixo AL, Curi AL, Benchimol EI, et al. Toxoplasmic retinochoroiditis: clinical characteristics and visual outcome in a prospective study. *PLoS Negl Trop Dis*. 2016;10:e0004685.
  318. Faridi A, Yeh S, Suhler EB, et al. Retinal detachment associated with ocular toxoplasmosis. *Retina*. 2015;35:358–363.
  319. Silveira C, Muccioli C, Holland GN, et al. Ocular involvement following an epidemic of *Toxoplasma gondii* Infection in Santa Isabel do Ivaí, Brazil. *Am J Ophthalmol*. 2015;159:1013–1021.e3.
  320. Johnson MW, Greven CM, Jaffe GJ, et al. Atypical, severe toxoplasmic retinochoroiditis in elderly patients. *Ophthalmology*. 1997;104:48–57.
  321. Kong JT, Grigg ME, Uyetake L, et al. Serotyping of *Toxoplasma gondii* infections in humans using synthetic peptides. *J Infect Dis*. 2003;187:1484–1495.
  322. Boyer K, Hill D, Mui E, et al. Unrecognized ingestion of *Toxoplasma gondii* oocysts leads to congenital toxoplasmosis and causes epidemics in North America. *Clin Infect Dis*. 2011;53:1081–1089.
  323. Gavinet MF, Robert F, Firtion G, et al. Congenital toxoplasmosis due to maternal reinfection during pregnancy. *J Clin Microbiol*. 1997;35:1276–1277.
  324. Hennequin C, Dureau P, N'Guyen L, et al. Congenital toxoplasmosis acquired from an immune woman. *Pediatr Infect Dis J*. 1997;16:75–76.
  325. Vogel N, Kirisits M, Michael E, et al. Congenital toxoplasmosis transmitted from an immunologically competent mother infected before conception. *Clin Infect Dis*. 1996;23:1055–1060.
  326. Minkoff H, Remington JS, Holman S, et al. Vertical transmission of *Toxoplasma* by human immunodeficiency virus-infected women. *Am J Obstet Gynecol*. 1997;176:555–559.
  327. Desmonts G, Couvreur J. Congenital toxoplasmosis. A prospective study of the offspring of 542 women who acquired toxoplasmosis during pregnancy. In: Thalhammer O, Pollak A, Baumgarten K, eds. *Pathophysiology of Congenital Disease. Perinatal Medicine (6th European Congress, Vienna)*. Stuttgart: Georg Thieme Publishers; 1979:51–60.
  328. Koppe JG, Loewer-Sieger DH, De Roeve-Bonnet H. Results of 20-year follow-up of congenital toxoplasmosis. *Am J Ophthalmol*. 1986;101:248–249.
  329. Wilson CB, Remington JS, Stagno S, et al. Development of adverse sequelae in children born with subclinical congenital *Toxoplasma* infection. *Pediatrics*. 1980;66:767–774.
  330. Delhaes L, Aizenberg D, Sicot B, et al. Severe congenital toxoplasmosis due to a *Toxoplasma gondii* strain with an atypical genotype: case report and review. *Prenat Diagn*. 2010;30:902–905.
  331. Forestier F. Fetal diseases, prenatal diagnoses and practical measures. *Presse Med*. 1991;20:1448–1454, [in French].
  332. Hohlfeld P, Daffos F, Thulliez P, et al. Fetal toxoplasmosis: outcome of pregnancy and infant follow-up after in utero treatment. *J Pediatr*. 1989;115:765–769.
  333. Hohlfeld P, Daffos F, Costa J-M, et al. Prenatal diagnosis of congenital toxoplasmosis with polymerase-chain-reaction test on amniotic fluid. *N Engl J Med*. 1994;331:695–699.
  334. Thiebaut R, Leproust S, Chene G, et al. Effectiveness of prenatal treatment for congenital toxoplasmosis: a meta-analysis of individual patients' data. *Lancet*. 2007;369:115–122.
  - 334a. Mandelbrot L, Kieffer F, Sitta R, et al. Prenatal therapy with pyrimethamine + sulfadiazine vs spiramycin to reduce placental transmission of toxoplasmosis: a multicenter, randomized trial. *Am J Obstet Gynecol*. 2018;219:386.e1–386.e9.
  - 334b. Montoya JG. Systematic screening and treatment of toxoplasmosis during pregnancy: is the glass half full or half empty? *Am J Obstet Gynecol*. 2018;219:315–319.
  335. Dunn D, Wallon M, Peyron F, et al. Mother-to-child transmission of toxoplasmosis: risk estimates for clinical counselling. *Lancet*. 1999;353:1829–1833.
  336. Hotop A, Hlobil H, Gross U. Efficacy of rapid treatment initiation following primary *Toxoplasma gondii* infection during pregnancy. *Clin Infect Dis*. 2012;54:1545–1552.
  337. Cortina-Borja M, Tan HK, Wallon M, et al. Prenatal treatment for serious neurological sequelae of congenital toxoplasmosis: an observational prospective cohort study. *PLoS Med*. 2010;7:e1000351.
  338. Olariu TR, Remington JS, McLeod R, et al. Severe congenital toxoplasmosis in the United States: clinical and serologic findings in untreated infants. *Pediatr Infect Dis J*. 2011;30:1056–1061.
  339. Dubey JP, Lago EG, Gennari SM, et al. Toxoplasmosis in humans and animals in Brazil: high prevalence, high burden of disease, and epidemiology. *Parasitology*. 2012;139:1375–1424.
  340. Rico-Torres CP, Vargas-Villavicencio JA, Correa D. Is *Toxoplasma gondii* type related to clinical outcome in human congenital infection? Systematic and critical review. *Eur J Clin Microbiol Infect Dis*. 2016;35:1079–1088.
  341. Hutson SL, Wheeler KM, McLone D, et al. Patterns of hydrocephalus caused by congenital *Toxoplasma gondii* infection associated with parasite genetics. *Clin Infect Dis*. 2015;61:1831–1834.
  342. Romand S, Chosson M, Franck J, et al. Usefulness of quantitative polymerase chain reaction in amniotic fluid as early prognostic marker of fetal infection with *Toxoplasma gondii*. *Am J Obstet Gynecol*. 2004;190:797–802.
  343. Yamamoto L, Targa LS, Sumita LM, et al. Association of parasite load levels in amniotic fluid with clinical outcome in congenital toxoplasmosis. *Obstet Gynecol*. 2017;130:335–345.
  344. Couvreur J, Desmonts G, Tournier G, et al. A homogeneous series of 210 cases of congenital toxoplasmosis in 0 to 11-month-old infants detected prospectively. *Ann Pediatr (Paris)*. 1984;31:815–819, [in French].
  345. De Roeve-Bonnet H, Koppe JG, Loewer-Sieger DH. Follow-up of children with congenital *Toxoplasma* infection and children who become serologically negative after 1 year of age, all born in 1964–1965. In: Thalhammer O, Pollak A, Baumgarten K, eds. *Pathophysiology of Congenital Disease. Perinatal Medicine (6th European Congress, Vienna)*. Stuttgart: Georg Thieme Publishers. 1979:61–75.
  346. Faucher B, Garcia-Meric P, Franck J, et al. Long-term ocular outcome in congenital toxoplasmosis: a prospective cohort of treated children. *J Infect*. 2012;64:104–109.
  347. Berrebi A, Assouline C, Bessieres MH, et al. Long-term outcome of children with congenital toxoplasmosis. *Am J Obstet Gynecol*. 2010;203:552.e1–552.e6.
  348. Phan L, Kasza K, Jalbrzikowski J, et al. Longitudinal study of new eye lesions in children with toxoplasmosis who were not treated during the first year of life. *Am J Ophthalmol*. 2008;146:375–384.
  349. Phan L, Kasza K, Jalbrzikowski J, et al. Longitudinal study of new eye lesions in treated congenital toxoplasmosis. *Ophthalmology*. 2008;115:553–559.e8.
  350. Wallon M, Garweg JG, Abrahamowicz M, et al. Ophthalmic outcomes of congenital toxoplasmosis followed until adolescence. *Pediatrics*. 2014;133:e601–e608.
  351. Mitchell CD, Erlich SS, Mastrucci MT, et al. Congenital toxoplasmosis occurring in infants perinatally infected with human immunodeficiency virus 1. *Pediatr Infect Dis J*. 1990;9:512–518.
  352. Costa F, Ko AI. Zika virus and microcephaly: where do we go from here? *Lancet Infect Dis*. 2018;18:236–237.
  353. Nowakowska D, Colon I, Remington JS, et al. Genotyping of *Toxoplasma gondii* by multiplex PCR and peptide-based serological testing of samples from infants in Poland diagnosed with congenital toxoplasmosis. *J Clin Microbiol*. 2006;44:1382–1389.
  354. McLeod R, Boyer KM, Lee D, et al. Prematurity and severity are associated with *Toxoplasma gondii* alleles (NCCCTS, 1981–2009). *Clin Infect Dis*. 2012;54:1595–1605.
  355. Boyer KM, Holfels E, Roizen N, et al. Risk factors for *Toxoplasma gondii* infection in mothers of infants with congenital toxoplasmosis: implications for prenatal management and screening. *Am J Obstet Gynecol*. 2005;192:564–571.
  356. Cook AJ, Gilbert RE, Buffolano W, et al. Sources of *Toxoplasma* infection in pregnant women: European multicentre case-control study. European Research Network on Congenital Toxoplasmosis. *BMJ*. 2000;321:142–147.
  357. Montoya JG. Laboratory diagnosis of *Toxoplasma gondii* infection and toxoplasmosis. *J Infect Dis*. 2002;185:S73–S82.
  358. Miller MJ, Aronson WJ, Remington JS. Late parasitemia in asymptomatic acquired toxoplasmosis. *Ann Intern Med*. 1969;71:139–145.
  359. Dhakal R, Gajurel K, Pomares C, et al. Significance of a positive *Toxoplasma* immunoglobulin M test result in the United States. *J Clin Microbiol*. 2015;53:3601–3605.
  360. Liesenfeld O, Montoya JG, Kinney S, et al. Effect of testing for IgG avidity in the diagnosis of *Toxoplasma gondii* infection in pregnant women: experience in a US reference laboratory. *J Infect Dis*. 2001;183:1248–1253.
  361. Liesenfeld O, Montoya JG, Thathineni NJ, et al. Confirmatory serologic testing for acute toxoplasmosis and rate of induced abortions among women reported to have positive *Toxoplasma* immunoglobulin M antibody titers. *Am J Obstet Gynecol*. 2001;184:140–145.
  362. Montoya JG, Liesenfeld O, Kinney S, et al. VIDAS test for avidity of *Toxoplasma*-specific immunoglobulin G for confirmatory testing of pregnant women. *J Clin Microbiol*. 2002;40:2504–2508.
  363. Sabin AB, Feldman HA. Dyes as microchemical indicators of a new immunity phenomenon affecting a protozoan parasite (*Toxoplasma*). *Science*. 1948;108:660–663.
  364. Balsari A, Poli G, Molina V, et al. ELISA for *Toxoplasma* antibody detection: a comparison with other serodiagnostic tests. *J Clin Pathol*. 1980;33:640–643.
  365. Walls KW, Bullock SL, English DK. Use of the enzyme-linked immunosorbent assay (ELISA) and its microadaptation for the serodiagnosis of toxoplasmosis. *J Clin Microbiol*. 1977;5:273–277.
  366. Walton BC, Benchoff BM, Brooks WH. Comparison of the indirect fluorescent antibody test and methylene blue dye test for detection of antibodies to *Toxoplasma gondii*. *Am J Trop Med Hyg*. 1966;15:149–152.
  367. Thulliez P, Remington JS, Santoro F, et al. A new agglutination test for the diagnosis of acute and chronic *Toxoplasma* infection. *Pathol Biol (Paris)*. 1986;34:173–177.
  368. Rougier S, Montoya JG, Peyron F. Lifelong persistence of *Toxoplasma* cysts: a questionable dogma? *Trends Parasitol*. 2017;33:414.
  369. Araujo FG, Barnett EV, Gentry LO, et al. False-positive anti-*Toxoplasma* fluorescent-antibody tests in patients with antinuclear antibodies. *Appl Microbiol*. 1971;22:270–275.
  370. Desmonts G, Remington JS. Direct agglutination test for diagnosis of *Toxoplasma* infection: method for increasing sensitivity and specificity. *J Clin Microbiol*. 1980;11:562–568.
  371. Dannemann BR, Vaughan WC, Thulliez P, et al. Differential agglutination test for diagnosis of recently acquired infection with *Toxoplasma gondii*. *J Clin Microbiol*. 1990;28:1928–1933.
  372. Montoya JG, Berry A, Rosso F, et al. The differential agglutination test as a diagnostic aid in cases of toxoplasmic lymphadenitis. *J Clin Microbiol*. 2007;45:1463–1468.
  373. Hedman K, Lappalainen M, Seppala I, et al. Recent primary *Toxoplasma* infection indicated by a low avidity of specific IgG. *J Infect Dis*. 1989;159:736–739.
  374. Lappalainen M, Koskela P, Koskineniemi M, et al. Toxoplasmosis acquired during pregnancy: improved serodiagnosis based on avidity of IgG. *J Infect Dis*. 1993;167:691–697.
  375. Montoya JG, Huffman HB, Remington JS. Evaluation of the immunoglobulin G avidity test for diagnosis of



- toxoplasmic lymphadenopathy. *J Clin Microbiol.* 2004;42:4627–4631.
376. Pelloux H, Brun E, Vernet G, et al. Determination of anti-*Toxoplasma gondii* immunoglobulin G avidity: adaptation to the Vidas system (bioMérieux). *Diagn Microbiol Infect Dis.* 1998;32:69–73.
  377. Villard O, Breit L, Cimon B, et al. Comparison of four commercially available avidity tests for *Toxoplasma gondii*-specific IgG antibodies. *Clin Vaccine Immunol.* 2013;20:197–204.
  378. Bobic B, Sibalic D, Djurkovic-Djakovic O. High levels of IgM antibodies specific for *Toxoplasma gondii* in pregnancy 12 years after primary *Toxoplasma* infection. *Gynecol Obstet Invest.* 1991;31:182–184.
  379. Liesenfeld O, Press C, Montoya JG, et al. False-positive results in immunoglobulin M (IgM) *Toxoplasma* antibody tests and importance of confirmatory testing: the Platelia toxo IgM test. *J Clin Microbiol.* 1997;35:174–178.
  380. Wilson M, Ware DA, Walls KW. Evaluation of commercial serodiagnostic kits for toxoplasmosis. *J Clin Microbiol.* 1987;25:2262–2265.
  381. U.S. Food and Drug Administration. *Public Health Advisory: Center for Devices and Radiological Health. Limitations of Toxoplasma IgM Commercial Test Kits, July 25, 1997.* Rockville, MD: U.S. Food and Drug Administration; 1997:1–3. <http://www.fda.gov/cdrh/toxophahtml>.
  382. Montoya JG, Remington JS. Management of *Toxoplasma gondii* infection during pregnancy. *Clin Infect Dis.* 2008;47:554–566.
  383. Welch PC, Masur H, Jones TC, et al. Serologic diagnosis of acute lymphadenopathic toxoplasmosis. *J Infect Dis.* 1980;142:256–264.
  384. Naot Y, Barnett EV, Remington JS. Method for avoiding false-positive results occurring in immunoglobulin M enzyme-linked immunosorbent assays due to presence of both rheumatoid factor and antinuclear antibodies. *J Clin Microbiol.* 1981;14:73–78.
  385. Filice GA, Yeager AS, Remington JS. Diagnostic significance of immunoglobulin M antibodies to *Toxoplasma gondii* detected after separation of immunoglobulin M from immunoglobulin G antibodies. *J Clin Microbiol.* 1980;12:336–342.
  386. Naot Y, Remington JS. An enzyme-linked immunosorbent assay for detection of IgM antibodies to *Toxoplasma gondii*: use for diagnosis of acute acquired toxoplasmosis. *J Infect Dis.* 1980;142:757–766.
  387. Naot Y, Desmonts G, Remington JS. IgM enzyme-linked immunosorbent assay test for the diagnosis of congenital *Toxoplasma* infection. *J Pediatr.* 1981;98:32–36.
  388. Siegel JP, Remington JS. Comparison of methods for quantitating antigen-specific immunoglobulin M antibody with a reverse enzyme-linked immunosorbent assay. *J Clin Microbiol.* 1983;18:63–70.
  389. Brenier-Pinchart MP, Fricker-Hidalgo H, Dard C, et al. Impact of heat-inactivation on anti-*Toxoplasma* IgM antibody levels. *Clin Chem Lab Med.* 2017;55:e291–e293.
  390. Remington JS, Eimstad WM, Araujo FG. Detection of immunoglobulin M antibodies with antigen-tagged latex particles in an immunosorbent assay. *J Clin Microbiol.* 1983;17:939–941.
  391. Plantaz D, Goullier A, Jouk PS, et al. Value of the immunosorbent agglutination assay (ISAGA) in the early diagnosis of congenital toxoplasmosis. *Pediatr.* 1987;42:387–391. [in French].
  392. Pomares C, Montoya JG. Laboratory diagnosis of congenital toxoplasmosis. *J Clin Microbiol.* 2016;54:2448–2454.
  393. Decoster A, Slizewicz B, Simon J, et al. Platelia-toxo IgA, a new kit for early diagnosis of congenital toxoplasmosis by detection of anti-P30 immunoglobulin A antibodies. *J Clin Microbiol.* 1991;29:2291–2295.
  394. Pinon JM, Thoannes H, Pouletty PH, et al. Detection of IgA specific for toxoplasmosis in serum and cerebrospinal fluid using a non-enzymatic IgA-capture assay. *Diagn Immunol.* 1986;4:223–227.
  395. Stepick-Biek P, Thulliez P, Araujo FG, et al. IgA antibodies for diagnosis of acute congenital and acquired toxoplasmosis. *J Infect Dis.* 1990;162:270–273.
  396. Murat JB, Souvignat A, Fricker-Hidalgo H, et al. Assessment of the IgA immunosorbent agglutination assay for the diagnosis of congenital toxoplasmosis on a series of 145 toxoplasmic seroconversions. *Clin Vaccine Immunol.* 2015;22:456–458.
  397. Pinon JM, Toubas D, Marx C, et al. Detection of specific immunoglobulin E in patients with toxoplasmosis. *J Clin Microbiol.* 1990;28:1739–1743.
  398. Wong SY, Hadju M-P, Ramirez R, et al. The role of specific immunoglobulin E in diagnosis of acute *Toxoplasma* infection and toxoplasmosis. *J Clin Microbiol.* 1993;31:2952–2959.
  399. Poirriez J, Toubas D, Marx-Chemia C, et al. Isotypic characterization of anti-*Toxoplasma gondii* antibodies in 18 cases of congenital toxoplasmic chorioretinitis. *Acta Ophthalmol (Copenh).* 1988;67:164–168.
  400. Foudrinier F, Villena I, Jausaud R, et al. Clinical value of specific immunoglobulin E detection by enzyme-linked immunosorbent assay in cases of acquired and congenital toxoplasmosis. *J Clin Microbiol.* 2003;41:1681–1686.
  401. Grover CM, Thulliez P, Remington JS, et al. Rapid prenatal diagnosis of congenital *Toxoplasma* infection by using polymerase chain reaction and amniotic fluid. *J Clin Microbiol.* 1990;28:2297–2301.
  402. Cinque P, Scarpellini P, Vago L, et al. Diagnosis of central nervous system complications in HIV-infected patients: cerebrospinal fluid analysis by the polymerase chain reaction. *AIDS.* 1997;11:1–17.
  403. Dupouy-Camet J, Lavareda de Souza L, Maslo C, et al. Detection of *Toxoplasma gondii* in venous blood from AIDS patients by polymerase chain reaction. *J Clin Microbiol.* 1993;31:1866–1869.
  404. Fricker-Hidalgo H, Brenier-Pinchart MP, Schaal JP, et al. Value of *Toxoplasma gondii* detection in one hundred thirty-three placentas for the diagnosis of congenital toxoplasmosis. *Pediatr Infect Dis J.* 2007;26:845–846.
  405. Robert-Gangneux F, Dupretz P, Yvenou C, et al. Clinical relevance of placenta examination for the diagnosis of congenital toxoplasmosis. *Pediatr Infect Dis J.* 2010;29:33–38.
  406. Sterkers Y, Varlet-Marie E, Marty P, et al. Diversity and evolution of methods and practices for the molecular diagnosis of congenital toxoplasmosis in France: a 4-year survey. *Clin Microbiol Infect.* 2010;16:1594–1602.
  407. Torres E, Rivera R, Cardona N, et al. Evaluation of IgG anti-*Toxoplasma* avidity and PCR in the postnatal diagnosis of congenital toxoplasmosis. *Pediatr Infect Dis J.* 2013;32:693–695.
  408. Dannemann BR, Israelski DM, Leoung GS, et al. *Toxoplasma* serology, parasitemia and antigenemia in patients at risk for toxoplasmic encephalitis. *AIDS.* 1991;5:1363–1365.
  409. Pelloux H, Dupouy-Camet J, Derouin F, et al; Group B-TS. A multicentre prospective study for the polymerase chain reaction detection of *Toxoplasma gondii* DNA in blood samples from 186 AIDS patients with suspected toxoplasmic encephalitis. *AIDS.* 1997;11:1888–1890.
  410. Patrat-Delon S, Gangneux JP, Lavoue S, et al. Correlation of parasite load determined by quantitative PCR to clinical outcome in a heart transplant patient with disseminated toxoplasmosis. *J Clin Microbiol.* 2010;48:2541–2545.
  411. Silveira C, Vallochi AL, Rodrigues da Silva U, et al. *Toxoplasma gondii* in the peripheral blood of patients with acute and chronic toxoplasmosis. *Br J Ophthalmol.* 2011;95:396–400.
  412. Santos FF, Nascimento H, Muccioli C, et al. Detection of *Toxoplasma gondii* DNA in peripheral blood and aqueous humor of patients with toxoplasmic active focal necrotizing retinochoroiditis using real-time PCR. *Arq Bras Ophthalmol.* 2015;78:356–358.
  413. Bastien P. Molecular diagnosis of toxoplasmosis. *Trans R Soc Trop Med Hyg.* 2002;96(suppl 1):S205–S215.
  414. Jones CD, Okhravi N, Adamson P, et al. Comparison of PCR detection methods for B1, P30, and 18S rDNA genes of *T. gondii* in aqueous humor. *Invest Ophthalmol Vis Sci.* 2000;41:634–644.
  415. Costa JM, Bretagne S. Variation of B1 gene and AF146527 repeat element copy numbers according to *Toxoplasma gondii* strains assessed using real-time quantitative PCR. *J Clin Microbiol.* 2012;50:1452–1454.
  416. Kasper DC, Sadeghi K, Prusa AR, et al. Quantitative real-time polymerase chain reaction for the accurate detection of *Toxoplasma gondii* in amniotic fluid. *Diagn Microbiol Infect Dis.* 2009;63:10–15.
  417. Belaz S, Gangneux JP, Dupretz P, et al. A 10-year retrospective comparison of two target sequences, REP-529 and B1, for *Toxoplasma gondii* detection by quantitative PCR. *J Clin Microbiol.* 2015;53:1294–1300.
  418. Morelle C, Varlet-Marie E, Brenier-Pinchart MP, et al. Comparative assessment of a commercial kit and two laboratory-developed PCR assays for the molecular diagnosis of congenital toxoplasmosis. *J Clin Microbiol.* 2012;50:3977–3982.
  419. Robert-Gangneux F, Sterkers Y, Yera H, et al. Molecular diagnosis of toxoplasmosis in immunocompromised patients: a three-year multicenter retrospective study. *J Clin Microbiol.* 2015;53:1677–1684.
  420. Meers S, Lagrou K, Theunissen K, et al. Myeloablative conditioning predisposes patients for *Toxoplasma gondii* reactivation after allogeneic stem cell transplantation. *Clin Infect Dis.* 2010;50:1127–1134.
  421. Levy RM, Bredesen DE, Rosenblum ML. Opportunistic central nervous system pathology in patients with AIDS. *Ann Neurol.* 1988;23:S7–S12.
  422. Frenkel JK, Piekarski G. The demonstration of *Toxoplasma* and other organisms by immunofluorescence: a pitfall [editorial]. *J Infect Dis.* 1978;138:265–266.
  423. Sun T, Greenspan J, Tenenbaum M, et al. Diagnosis of cerebral toxoplasmosis using fluorescein-labeled anti-*Toxoplasma* monoclonal antibodies. *Am J Surg Pathol.* 1986;10:312–316.
  424. Cerezo L, Alvarez M, Price G. Electron microscopic diagnosis of cerebral toxoplasmosis. Case report. *J Neurosurg.* 1985;63:470–472.
  425. Luft BJ, Billingham M, Remington JS. Endomyocardial biopsy in the diagnosis of toxoplasmic myocarditis. *Transplant Proc.* 1986;18:1871–1873.
  426. Porter SB, Sande M. Toxoplasmosis of the central nervous system in the acquired immunodeficiency syndrome. *N Engl J Med.* 1992;327:1643–1648.
  427. Haverkos HW. Assessment of therapy for *Toxoplasma* encephalitis. The TE Study Group. *Am J Med.* 1987;82:907–914.
  428. Levy RM, Rosenbloom S, Perrett LV. Neuroradiologic findings in AIDS: a review of 200 cases. *AJNR Am J Neuroradiol.* 1986;147:977–983.
  429. Post MJ, Kursunoglu SJ, Hensley GT, et al. Cranial CT in acquired immunodeficiency syndrome: spectrum of diseases and optimal contrast enhancement technique. *AJR Am J Roentgenol.* 1985;145:929–940.
  430. Levy RM, Mills CM, Posin JP, et al. The efficacy and clinical impact of brain imaging in neurologically symptomatic AIDS patients: a prospective CT/MRI study. *J Acquir Immune Defic Syndr.* 1990;3:461–471.
  431. Post MJ, Chan JC, Hensley GT, et al. *Toxoplasma* encephalitis in Haitian adults with acquired immunodeficiency syndrome: a clinical-pathologic-CT correlation. *AJR Am J Roentgenol.* 1983;140:861–868.
  432. Ciricillo SF, Rosenblum ML. Use of CT and MR imaging to distinguish intracranial lesions and to define the need for biopsy in AIDS patients. *J Neurosurg.* 1990;73:720–724.
  433. Holloway RG, Mushlin AI. Intracranial mass lesions in acquired immunodeficiency syndrome: using decision analysis to determine the effectiveness of stereotactic brain biopsy. *Neurology.* 1996;46:1010–1015.
  434. De La Paz R, Enzmann D. Neuroradiology of acquired immunodeficiency syndrome. In: Rosenblum ML, ed. *AIDS and the Nervous System.* New York: Raven Press; 1988:121–154.
  435. Luft BJ, Hafner R, Korzun AH, et al. Toxoplasmic encephalitis in patients with the acquired immunodeficiency syndrome. *N Engl J Med.* 1993;329:995–1000.
  436. Harris TM, Smith RR, Bognanno JR, et al. Toxoplasmic myelitis in AIDS: gadolinium-enhanced MR. *J Comput Assist Tomogr.* 1990;14:809–811.
  437. García-García C, Castillo-Álvarez F, Azcona-Gutiérrez JM, et al. Spinal cord toxoplasmosis in human immunodeficiency virus infection/acquired immunodeficiency syndrome. *Infect Dis (Lond).* 2015;47:277–282.
  438. O'Doherty MJ, Barrington SF, Campbell M, et al. PET scanning and the human immunodeficiency virus-positive patient. *J Nucl Med.* 1997;38:1575–1583.
  439. Naddaf SY, Akisik MF, Aziz M, et al. Comparison between 201Tl-chloride and 99Tc(m)-sestamibi SPET brain imaging for differentiating intracranial lymphoma from non-malignant lesions in AIDS patients. *Nucl Med Commun.* 1998;19:47–53.
  440. Ramsey RG, Gean AD. Central nervous system toxoplasmosis. *Neuroimag Clin N Am.* 1997;7:171–186.
  441. Rosenfeld SS, Hoffman JM, Coleman RE, et al. Studies of primary central nervous system lymphoma with [18F]-fluorodeoxyglucose (FDG) PET. *J Nucl Med.* 1992;33:532–536.
  442. Giancola ML, Rizzi EB, Schiavo R, et al. Reduced value of thallium-201 single-photon emission computed tomography in the management of HIV-related focal brain lesions in the era of highly active antiretroviral therapy. *AIDS Res Hum Retroviruses.* 2004;20:584–588.
  443. Navia BA, Pettito CK, Gold JW, et al. Cerebral toxoplasmosis complicating the acquired immune deficiency syndrome: clinical and neuropathological findings in 27 patients. *Ann Neurol.* 1986;19:224–238.
  444. Orefice G, Carrieri PB, de Marinis T, et al. Use of the intrathecal synthesis of anti-*Toxoplasma* antibodies in the diagnostic assessment and in the follow-up of AIDS patients with cerebral toxoplasmosis. *Acta Neurol (Napoli).* 1990;12:79–81.
  445. Potasman I, Resnick L, Luft BJ, et al. Intrathecal production of antibodies against *Toxoplasma gondii* in patients with toxoplasmic encephalitis and the acquired immunodeficiency syndrome (AIDS). *Ann Intern Med.* 1988;108:49–51.
  446. Fricker-Hidalgo H, Bulabois CE, Brenier-Pinchart MP, et al. Diagnosis of toxoplasmosis after allogeneic stem cell



- transplantation: results of DNA detection and serological techniques. *Clin Infect Dis*. 2009;48:e9–e15.
447. Bossolasco S, Cinque P, Ponzoni M, et al. Epstein-Barr virus DNA load in cerebrospinal fluid and plasma of patients with AIDS-related lymphoma. *J Neurovirol*. 2002;8:432–438.
  448. Holliman RE, Johnson JD, Savva D. Diagnosis of cerebral toxoplasmosis in association with AIDS using the polymerase chain reaction. *Scand J Infect Dis*. 1990;22:243–244.
  449. Roth A, Roth B, Höffken G, et al. Application of the polymerase chain reaction to diagnosis of pulmonary toxoplasmosis in immunocompromised patients. *Eur J Clin Microbiol Infect Dis*. 1992;11:1177–1181.
  450. Fardeau C, Romand S, Rao NA, et al. Diagnosis of toxoplasmic retinochoroiditis with atypical clinical features. *Am J Ophthalmol*. 2002;134:196–203.
  451. Westeneng AC, Rothova A, de Boer JH, et al. Infectious uveitis in immunocompromised patients and the diagnostic value of polymerase chain reaction and Goldmann-Witmer coefficient in aqueous analysis. *Am J Ophthalmol*. 2007;144:781–785.
  452. Garweg JG, de Groot-Mijnes JD, Montoya JG. Diagnostic approach to ocular toxoplasmosis. *Ocul Immunol Inflamm*. 2011;19:255–261.
  453. Bretagne S, Costa JM, Keutzi M, et al. Late toxoplasmosis evidenced by PCR in marrow transplant recipient. *Bone Marrow Transplant*. 1995;15:809–811.
  454. Turunen HJ, Leinikki PO, Saari KM. Demonstration of intraocular synthesis of immunoglobulin G *Toxoplasma* antibodies for specific diagnosis of toxoplasmic chorioretinitis by enzyme immunoassay. *J Clin Microbiol*. 1983;17:988–992.
  455. Rothova A, de Boer JH, Ten Dam-van Loon NH, et al. Usefulness of aqueous humor analysis for the diagnosis of posterior uveitis. *Ophthalmology*. 2008;115:306–311.
  456. Montoya JG, Huffman HB, Remington JS. Use of the VIDAS Toxo IgG Avidity test for the diagnosis of toxoplasmic lymphadenopathy. Presented at 40th Annual Meeting of the Infectious Diseases Society of America (IDSA). Chicago, October 24–27, 2002.
  457. Wallon M, Franck J, Thulliez P, et al. Accuracy of real-time polymerase chain reaction for *Toxoplasma gondii* in amniotic fluid. *Obstet Gynecol*. 2010;115:727–733.
  458. Sterkers Y, Pratlong F, Albaba S, et al. Novel interpretation of molecular diagnosis of congenital toxoplasmosis according to gestational age at the time of maternal infection. *J Clin Microbiol*. 2012;50:3944–3951.
  459. Remington JS, Araujo FG, Desmonts G. Recognition of different *Toxoplasma* antigens by IgM and IgG antibodies in mothers and their congenitally infected newborns. *J Infect Dis*. 1985;152:1020–1024.
  460. Rilling V, Dietz K, Krczal D, et al. Evaluation of a commercial IgG/IgM Western blot assay for early postnatal diagnosis of congenital toxoplasmosis. *Eur J Clin Microbiol Infect Dis*. 2003;22:174–180.
  461. Gallego-Marin C, Henao AC, Gomez-Marín JE. Clinical validation of a Western blot assay for congenital toxoplasmosis and newborn screening in a hospital in Armenia (Quindío) Colombia. *J Trop Pediatr*. 2006;52:107–112.
  462. Tridapalli E, Capretti M, Farneti G, et al. Congenital toxoplasmosis: the importance of the Western blot method to avoid unnecessary therapy in potentially infected newborns. *Acta Paediatr*. 2008;97:1298–1300.
  463. L'Ollivier C, Wallon M, Faucher B, et al. Comparison of mother and child antibodies that target high-molecular-mass *Toxoplasma gondii* antigens by immunoblotting improves neonatal diagnosis of congenital toxoplasmosis. *Clin Vaccine Immunol*. 2012;19:1326–1328.
  464. Cazenave J, Forestier F, Bessieres M, et al. Contribution of a new PCR assay to the prenatal diagnosis of congenital toxoplasmosis. *Prenat Diagn*. 1992;12:119–127.
  465. Fuentes I, Rodriguez M, Domingo CJ, et al. Urine sample used for congenital toxoplasmosis diagnosis by PCR. *J Clin Microbiol*. 1996;34:2368–2371.
  466. van de Ven E, Melchers W, Galama J, et al. Identification of *Toxoplasma gondii* infections by BI gene amplification. *J Clin Microbiol*. 1991;19:2120–2124.
  467. Sterkers Y, Ribot J, Albaba S, et al. Diagnosis of congenital toxoplasmosis by polymerase chain reaction on neonatal peripheral blood. *Diagn Microbiol Infect Dis*. 2001;7:174–176.
  468. Olariu TR, Remington JS, Montoya JG. Polymerase chain reaction in cerebrospinal fluid for the diagnosis of congenital toxoplasmosis. *Pediatr Infect Dis J*. 2014;33:566–570.
  469. Guerina N, Hsu H-W, Meissner H, et al. Neonatal serologic screening and early treatment for congenital *Toxoplasma gondii* infection. *N Engl J Med*. 1994;330:1858–1863.
  470. Jara M, Hsu HW, Eaton RB, et al. Epidemiology of congenital toxoplasmosis identified by population-based newborn screening in Massachusetts. *Pediatr Infect Dis J*. 2001;20:1132–1135.
  471. Cohn J, McMeeking A, Cohen W, et al. Evaluation of the policy of empiric treatment of suspected *Toxoplasma* encephalitis in patients with the acquired immunodeficiency syndrome. *Am J Med*. 1989;86:521–527.
  472. Carbone LG, Bendixen B, Appel GB. Sulfadiazine-associated obstructive nephropathy occurring in a patient with the acquired immunodeficiency syndrome. *Am J Kidney Dis*. 1988;12:72–75.
  473. Reboli AC, Mandler HD. Encephalopathy and psychosis associated with sulfadiazine in two patients with AIDS and CNS toxoplasmosis. *Clin Infect Dis*. 1992;15:556–557.
  474. Leoung GS, Stanford JE, Giordano MF, et al. Trimethoprim-sulfamethoxazole (TMP-SMZ) dose escalation versus direct rechallenge for *Pneumocystis carinii* pneumonia prophylaxis in human immunodeficiency virus-infected patients with previous adverse reaction to TMP-SMZ. *J Infect Dis*. 2001;184:992–997.
  475. Soffritti S, Ricci G, Prete A, et al. Successful desensitization to trimethoprim-sulfamethoxazole after allogeneic haematopoietic stem cell transplantation: preliminary observations. *Med Pediatr Oncol*. 2003;40:271–272.
  476. Coppola S, Angarano G, Monno L, et al. Adverse effects of clindamycin in the treatment of cerebral toxoplasmosis in AIDS patients. Program and Abstracts of 7th International Conference on AIDS. Florence, Italy, June 16–21, 1991:265.
  477. Torre D, Casari S, Speranza F, et al. Randomized trial of trimethoprim-sulfamethoxazole versus pyrimethamine-sulfadiazine for therapy of toxoplasmic encephalitis in patients with AIDS. Italian Collaborative Study Group. *Antimicrob Agents Chemother*. 1998;42:1346–1349.
  478. Alavi SM, Alavi L. Treatment of toxoplasmic lymphadenitis with co-trimoxazole: double-blind, randomized clinical trial. *Int J Infect Dis*. 2010;14(suppl 3):e67–e69.
  479. Carey RM, Kimball AC, Armstrong D, et al. Toxoplasmic lymphitis. Clinical experiences in a cancer hospital. *Am J Med*. 1973;54:30–38.
  480. Katlama C, De Wit S, O'Doherty E, et al. Pyrimethamine-clindamycin vs. pyrimethamine-sulfadiazine as acute and long-term therapy for toxoplasmic encephalitis in patients with AIDS. *Clin Infect Dis*. 1996;22:268–275.
  481. Luft BJ, Remington JS. AIDS commentary. Toxoplasmic encephalitis. *J Infect Dis*. 1988;157:1–6.
  482. Dannemann BR, McCutchan JA, Israelski DA, et al. Treatment of toxoplasmic encephalitis in patients with AIDS: a randomized trial comparing pyrimethamine plus clindamycin to pyrimethamine plus sulfadiazine. *Ann Intern Med*. 1992;116:33–43.
  483. Canessa A, Del Bono V, De Leo P, et al. Cotrimoxazole therapy of *Toxoplasma gondii* encephalitis in AIDS patients. *Eur J Clin Microbiol Infect Dis*. 1992;11:125–130.
  484. Solbreux P, Sonnet J, Zech F. A retrospective study about the use of cotrimoxazole as diagnostic support and treatment of suspected cerebral toxoplasmosis in AIDS. *Acta Clin Belg*. 1990;45:85–96.
  485. Torre D, Speranza F, Martegani R, et al. A retrospective study of treatment of cerebral toxoplasmosis in AIDS patients with trimethoprim-sulphamethoxazole. *J Infect*. 1998;37:15–18.
  486. Kovacs JA. Efficacy of atovaquone in treatment of toxoplasmosis in patients with AIDS. *Lancet*. 1992;340:637–638.
  487. Torres R, Weinberg W, Stansell J, et al. Atovaquone for salvage treatment and suppression of toxoplasmic encephalitis in patients with AIDS. *Clin Infect Dis*. 1997;24:422–429.
  488. Chirgwin K, Hafner R, Leport C, et al. Randomized phase II trial of atovaquone with pyrimethamine or sulfadiazine for treatment of toxoplasmic encephalitis in patients with acquired immunodeficiency syndrome: ACTG 237/ANRS 039 Study. AIDS Clinical Trials Group 237/Agence Nationale de Recherche sur le SIDA, Essai 039. *Clin Infect Dis*. 2002;34:1243–1250.
  489. Calderon MM, Penzak SR, Pau AK, et al. Efavirenz but not atazanavir/ritonavir significantly reduces atovaquone concentrations in HIV-infected subjects. *Clin Infect Dis*. 2016;62:1036–1042.
  490. Fernandez-Martin J, Leport C, Morlat P, et al. Pyrimethamine-clarithromycin for therapy of acute *Toxoplasma* encephalitis in patients with AIDS. *Antimicrob Agents Chemother*. 1991;35:2049–2052.
  491. Chaisson RE, Benson CA, Dube MP, et al. Clarithromycin therapy for bacteremic *Mycobacterium avium* complex disease. A randomized, double-blind, dose-ranging study in patients with AIDS. AIDS Clinical Trials Group Protocol 157 Study Team. *Ann Intern Med*. 1994;121:905–911.
  492. Cohn DL, Fisher EJ, Peng GT, et al. A prospective randomized trial of four three-drug regimens in the treatment of disseminated *Mycobacterium avium* complex disease in AIDS patients: excess mortality associated with high-dose clarithromycin. Terry Bein Community Programs for Clinical Research on AIDS. *Clin Infect Dis*. 1999;29:125–133.
  493. Ward DJ. Dapsone/pyrimethamine for the treatment of toxoplasmic encephalitis. Program and Abstracts of 8th International Conference on AIDS. Amsterdam, July 19–24, 1992:B133.
  494. Pope-Pegram L, Gathe J Jr, Bohn B, et al. Treatment of presumed central nervous system toxoplasmosis with doxycycline. Program and Abstracts of 7th International Conference on AIDS. Florence, Italy, June 16–21, 1991:188.
  495. Turett G, Pierone G, Masci J, et al. Failure of doxycycline in the treatment of cerebral toxoplasmosis. Program and Abstracts of 6th International Conference on AIDS. San Francisco, June 20–24, 1990:Th.B.479.
  496. Saba J, Morlat P, Raffi F, et al. Pyrimethamine plus azithromycin for treatment of acute toxoplasmic encephalitis in patients with AIDS. *Eur J Clin Microbiol Infect Dis*. 1993;12:853–856.
  497. Jacobson JM, Hafner R, Remington J, et al. Dose-escalation, phase I/II study of azithromycin and pyrimethamine for the treatment of toxoplasmic encephalitis in AIDS. *AIDS*. 2001;15:583–589.
  498. Pedrol E, Gonzales-Clemente JM, Gatell JM, et al. Central nervous system toxoplasmosis in AIDS patients: efficacy of an intermittent maintenance therapy. *AIDS*. 1990;4:511–517.
  499. Podzamczar D, Miró J, Bolao F, et al. Twice-weekly maintenance therapy with sulfadiazine-pyrimethamine to prevent recurrent toxoplasmic encephalitis in patients with AIDS. *Ann Intern Med*. 1995;123:175–180.
  500. Podzamczar D, Miro JM, Ferrer E, et al. Thrice-weekly sulfadiazine-pyrimethamine for maintenance therapy of toxoplasmic encephalitis in HIV-infected patients. Spanish Toxoplasmosis Study Group. *Eur J Clin Microbiol Infect Dis*. 2000;19:89–95.
  501. Heald A, Flepp M, Chave J-P, et al. Treatment for cerebral toxoplasmosis protects against *Pneumocystis carinii* pneumonia in patients with AIDS. *Ann Intern Med*. 1991;115:760–763.
  502. Duval X, Pajot O, Le Moing V, et al. Maintenance therapy with cotrimoxazole for toxoplasmic encephalitis in the era of highly active antiretroviral therapy. *AIDS*. 2004;18:1342–1344.
  503. Remington JS, Vilde JL, Antunes F, et al. Clindamycin for toxoplasmic encephalitis in AIDS. *Lancet*. 1991;338:1142–1143.
  504. Panel on Opportunistic Infections in HIV-Infected Adults and Adolescents. Guidelines for Guidelines for the Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents: Recommendations from the Centers for Disease Control and Prevention, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. [http://aidsinfo.nih.gov/contentfiles/lvguidelines/adult\\_oai.pdf](http://aidsinfo.nih.gov/contentfiles/lvguidelines/adult_oai.pdf). Accessed December 30, 2017.
  505. Köppen S, Grunewald T, Jautzke G, et al. Prevention of *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis in human immunodeficiency virus infected patients: a clinical approach comparing aerosolized pentamidine and pyrimethamine/sulfadoxine. *Clin Invest*. 1992;70:508–512.
  506. Clotet B, Sirena G, Romeu J, et al. Twice-weekly dapsone-pyrimethamine for preventing PCP and cerebral toxoplasmosis. *AIDS*. 1991;5:601–602.
  507. Opravil M, Hirschel B, Lazzarin A, et al. Once-weekly administration of dapsone/pyrimethamine vs. aerosolized pentamidine as a combined prophylaxis for *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis in human immunodeficiency virus-infected patients. *Clin Infect Dis*. 1995;20:531–541.
  508. Torres R, Barr M, Thorn M, et al. Randomized trial of dapsone and aerosolized pentamidine for the prophylaxis of *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis. *Am J Med*. 1993;95:573–583.
  509. Navin TR, Miller KD, Satriale RF, et al. Adverse reactions associated with pyrimethamine-sulfadoxine prophylaxis for *Pneumocystis carinii* infections in AIDS. *Lancet*. 1985;1:1332.
  510. Maslo C, Matheron S, Saimot AG. Cerebral toxoplasmosis: assessment of maintenance therapy. Program and Abstracts of 8th International Conference on AIDS. Amsterdam, July 19–24, 1992:B123.
  511. de Gans J, Portegies P, Reiss P, et al. Pyrimethamine alone as maintenance therapy for central nervous system

- toxoplasmosis in 38 patients with AIDS. *J Acquir Immune Defic Syndr*. 1992;5:137-142.
512. Katlama C, Mouthon B, Gourdon D, et al. Atovaquone as long-term suppressive therapy for toxoplasmic encephalitis in patients with AIDS and multiple drug intolerance. *AIDS*. 1996;10:1107-1112.
  513. Zolopa A, Andersen J, Powderly W, et al. Early antiretroviral therapy reduces AIDS progression/death in individuals with acute opportunistic infections: a multicenter randomized strategy trial. *PLoS ONE*. 2009;4:e5575.
  514. Carr A, Tindall B, Brew BJ, et al. Low-dose trimethoprim-sulfamethoxazole prophylaxis for toxoplasmic encephalitis in patients with AIDS. *Ann Intern Med*. 1992;117:106-111.
  515. Bozzette S, Finkelstein D, Spector S, et al. A randomized trial of three antipneumocystis agents in patients with advanced human immunodeficiency virus infection. *N Engl J Med*. 1995;332:693-699.
  516. Podzamczar D, Salazar A, Jiménez J, et al. Intermittent trimethoprim-sulfamethoxazole compared with dapsone-pyrimethamine for the simultaneous primary prophylaxis of *Pneumocystis pneumonia* and toxoplasmosis in patients infected with HIV. *Ann Intern Med*. 1995;122:755-761.
  517. Girard P-M, Landman R, Gaubert C, et al. Dapsone-pyrimethamine compared with aerosolized pentamidine as a primary prophylaxis against *Pneumocystis carinii* pneumonia and toxoplasmosis in HIV infection. *N Engl J Med*. 1993;328:1514-1520.
  518. Schurmann D, Bergmann F, Albrecht H, et al. Effectiveness of twice-weekly pyrimethamine-sulfadoxine as primary prophylaxis of *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis in patients with advanced HIV infection. *Eur J Clin Microbiol Infect Dis*. 2002;21:353-361.
  519. Gajurel K, Dhakal R, Montoya JG. Toxoplasmosis in hematopoietic cell transplant recipients. *Transpl Infect Dis*. 2017;19.
  520. Zeller V, Truffot C, Agher R, et al. Discontinuation of secondary prophylaxis against disseminated *Mycobacterium avium* complex infection and toxoplasmic encephalitis. *Clin Infect Dis*. 2002;34:662-667.
  521. Mussini C, Pezzotti P, Govoni A, et al. Discontinuation of primary prophylaxis for *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis in human immunodeficiency virus type I-infected patients: the changes in opportunistic prophylaxis study. *J Infect Dis*. 2000;181:1635-1642.
  522. Miro JM Stopping Secondary TE Prophylaxis in Suppressed Patients with CD4 100-200 Is Not Safe. Conference on Retroviruses and Opportunistic Infections. Poster 646. February 13-16, 2017; Seattle, Washington.
  523. Baden LR, Katz JT, Franck L, et al. Successful toxoplasmosis prophylaxis after orthotopic cardiac transplantation with trimethoprim-sulfamethoxazole. *Transplantation*. 2003;75:339-343.
  524. Durant J, Hazime F, Charles M, et al. Prevention of *Pneumocystis carinii* pneumonia and of cerebral toxoplasmosis by roxithromycin in HIV-infected patients. *Infection*. 1995;23:S33-S38.
  525. Jacobson M, Besch C, Child C, et al. Primary prophylaxis with pyrimethamine for toxoplasmic encephalitis in patients with advanced human immunodeficiency virus disease: results of a randomized trial. *J Infect Dis*. 1994;169:384-394.
  526. Klinker H, Langmann P, Richter E. Pyrimethamine alone as prophylaxis for cerebral toxoplasmosis in patients with advanced HIV infection. *Infection*. 1996;4:324-328.
  527. Raffi F, Struillou L, Ninin E, et al. Breakthrough cerebral toxoplasmosis in patients with AIDS who are being treated with clarithromycin. *Clin Infect Dis*. 1995;20:1076-1077.
  528. Lepout C, Vilde JL, Katlama C, et al. Failure of spiramycin to prevent neurotoxoplasmosis in immunosuppressed patients. *Med Clin North Am*. 1986;70:677-692.
  529. Jacobson M, Besch C, Child C, et al. Toxicity of clindamycin as prophylaxis for AIDS-associated toxoplasmic encephalitis. *Lancet*. 1992;339:333-334.
  530. Cocherneau-Massin I, LeHoang P, Lautier-Frau M, et al. Ocular toxoplasmosis in human immunodeficiency virus-infected patients. *Am J Ophthalmol*. 1992;114:130-135.
  531. Friedman AH, Orellana J, Gagliuso DJ, et al. Ocular toxoplasmosis in AIDS patients. *Trans Am Ophthalmol Soc*. 1990;88:63-88.
  532. Schnapp L, Geaghan S, Campagna A, et al. *Toxoplasma gondii* pneumonitis in patients infected with the human immunodeficiency virus. *Arch Intern Med*. 1992;152:1073-1076.
  533. Kim SJ, Scott IU, Brown GC, et al. Interventions for *Toxoplasma* retinochoroiditis: a report by the American Academy of Ophthalmology. *Ophthalmology*. 2013;120:371-378.
  534. Holland GN, Lewis KG. An update on current practices in the management of ocular toxoplasmosis. *Am J Ophthalmol*. 2002;134:102-114.
  535. Holland GN. Ocular toxoplasmosis: a global reassessment, Part I. Epidemiology and course of disease. *Am J Ophthalmol*. 2003;136:973-988.
  536. Holland GN. Ocular toxoplasmosis: a global reassessment, Part II. Disease manifestations and management. *Am J Ophthalmol*. 2004;137:1-17.
  537. Soheilian M, Sadoughi MM, Ghajarnia M, et al. Prospective randomized trial of trimethoprim/sulfamethoxazole versus pyrimethamine and sulfadiazine in the treatment of ocular toxoplasmosis. *Ophthalmology*. 2005;112:1876-1882.
  538. Baharivand N, Mahdaviard A, Fouladi RF. Intravitreal clindamycin plus dexamethasone versus classic oral therapy in toxoplasmic retinochoroiditis: a prospective randomized clinical trial. *Int Ophthalmol*. 2013;33:39-46.
  539. Soheilian M, Ramezani A, Azimzadeh A, et al. Randomized trial of intravitreal clindamycin and dexamethasone versus pyrimethamine, sulfadiazine, and prednisolone in treatment of ocular toxoplasmosis. *Ophthalmology*. 2011;118:134-141.
  540. Souza CE, Nascimento H, Lima A, et al. Intravitreal injection of sulfamethoxazole and trimethoprim associated with dexamethasone as an alternative therapy for ocular toxoplasmosis. *Ocul Immunol Inflamm*. 2017;1-4.
  541. Choudhury H, Jindal A, Pathengay A, et al. The role of intravitreal trimethoprim/sulfamethoxazole in the treatment of *Toxoplasma* retinochoroiditis. *Ophthalmic Surg Lasers Imaging Retina*. 2015;46:137-140.
  542. Pearson PA, Piracha AR, Sen HA, et al. Atovaquone for the treatment of *Toxoplasma* retinochoroiditis in immunocompetent patients. *Ophthalmology*. 1999;106:148-153.
  543. Bosch-Driessen LH, Verbraak FD, Suttrop-Schulten MS, et al. A prospective, randomized trial of pyrimethamine and azithromycin vs pyrimethamine and sulfadiazine for the treatment of ocular toxoplasmosis. *Am J Ophthalmol*. 2002;134:34-40.
  544. Lashay A, Mirshahi A, Parandini N, et al. A prospective randomized trial of azithromycin versus trimethoprim/sulfamethoxazole in treatment of toxoplasmic retinochoroiditis. *J Curr Ophthalmol*. 2016;29:120-125.
  545. Jasper S, Vedula SS, John SS, et al. Corticosteroids as adjuvant therapy for ocular toxoplasmosis. *Cochrane Database Syst Rev*. 2017;(1):CD007417.
  546. Oray M, Ozdal PC, Cebeci Z, et al. Fulminant ocular toxoplasmosis: the hazards of corticosteroid monotherapy. *Ocul Immunol Inflamm*. 2016;24:637-646.
  547. Silveira C, Muccioli C, Nussenblatt R, et al. The effect of long-term intermittent trimethoprim/sulfamethoxazole treatment on recurrences of toxoplasmic retinochoroiditis: 10 years of follow-up. *Ocul Immunol Inflamm*. 2015;23:246-247.
  548. Fernandes Felix JP, Cavalcanti Lira RP, Cosimo AB, et al. Trimethoprim-sulfamethoxazole versus placebo in reducing the risk of toxoplasmic retinochoroiditis recurrences: a three-year follow-up. *Am J Ophthalmol*. 2016;170:176-182.
  549. Grigg ME, Dubey JP, Nussenblatt RB. Ocular toxoplasmosis: lessons from Brazil. *Am J Ophthalmol*. 2015;159:999-1001.
  550. Valentini P, Buonsenso D, Barone G, et al. Spiramycin/cotrimoxazole versus pyrimethamine/sulfonamide and spiramycin alone for the treatment of toxoplasmosis in pregnancy. *J Perinatol*. 2015;35:90-94.
  551. Forestier F, Daffos F, Rainaut M, et al. Suivi thérapeutique foetomaternel de la spiramycine en cours de grossesse. [Fetomaternal therapeutic follow-up of spiramycin during pregnancy]. *Arch Fr Pediatr*. 1987;44:539-544.
  552. Gilbert R, Gras L, European Multicentre Study on Congenital Toxoplasmosis. Effect of timing and type of treatment on the risk of mother to child transmission of *Toxoplasma gondii*. *BJOG*. 2003;110:112-120.
  553. Gilbert RE, Gras L, Wallon M, et al. Effect of prenatal treatment on mother to child transmission of *Toxoplasma gondii*: retrospective cohort study of 554 mother-child pairs in Lyon, France. *Int J Epidemiol*. 2001;30:1303-1308.
  554. Gilbert R, Dunn D, Wallon M, et al. Ecological comparison of the risks of mother-to-child transmission and clinical manifestations of congenital toxoplasmosis according to prenatal treatment protocol. *Epidemiol Infect*. 2001;127:113-120.
  555. Foulon W, Villena I, Stray-Pedersen B, et al. Treatment of toxoplasmosis during pregnancy: a multicenter study of impact on fetal transmission and children's sequelae at age 1 year. *Am J Obstet Gynecol*. 1999;180:410-415.
  556. Thulliez P. Commentary: efficacy of prenatal treatment for toxoplasmosis: a possibility that cannot be ruled out. *Int J Epidemiol*. 2001;30:1315-1316.
  557. Moncada PA, Montoya JG. Toxoplasmosis in the fetus and newborn: an update on prevalence, diagnosis and treatment. *Expert Rev Anti Infect Ther*. 2012;10:815-828.
  558. Wallon M, Peyron F, Cornu C, et al. Congenital *Toxoplasma* infection: monthly prenatal screening decreases transmission rate and improves clinical outcome at age 3 years. *Clin Infect Dis*. 2013;56:1223-1231.
  559. Prusa AR, Kasper DC, Pollak A, et al. The Austrian Toxoplasmosis Register, 1992-2008. *Clin Infect Dis*. 2015;60:e4-e10.
  560. Garweg JG, Scherrer J, Wallon M, et al. Reactivation of ocular toxoplasmosis during pregnancy. *BJOG*. 2005;112:241-242.
  561. Maldonado YA, Read JS, Committee on Infectious Diseases. Diagnosis, treatment, and prevention of congenital toxoplasmosis in the United States. *Pediatrics*. 2017;139:e20163860.
  562. McAuley J, Boyer KM, Patel D, et al. Early and longitudinal evaluations of treated infants and children and untreated historical patients with congenital toxoplasmosis: the Chicago Collaborative Treatment Trial. *Clin Infect Dis*. 1994;18:38-72.
  563. Chemla C, Villena I. Sulfadoxine-pyrimethamine combination in congenital toxoplasmosis. *Pediatr Infect Dis J*. 2017;36:349-350.
  564. Teil J, Dupont D, Charpiat B, et al. Treatment of congenital toxoplasmosis: safety of the sulfadoxine-pyrimethamine combination in children based on a method of causality assessment. *Pediatr Infect Dis J*. 2016;35:634-638.
  565. McGee T, Wolters C, Stein L, et al. Absence of sensorineural hearing loss in treated infants and children with congenital toxoplasmosis. *Otolaryngol Head Neck Surg*. 1992;106:75-80.
  566. Mets MB, Hofels E, Boyer KM, et al. Eye manifestations of congenital toxoplasmosis. *Am J Ophthalmol*. 1996;122:309-324.
  567. Roizen N, Swisher CN, Stein MA, et al. Neurologic and developmental outcome in treated congenital toxoplasmosis. *Pediatrics*. 1995;95:11-20.
  568. Wreghitt TG, Hakim M, Gray JJ, et al. Toxoplasmosis in heart and heart lung transplant recipients. *J Clin Pathol*. 1989;42:194-199.
  569. Gajurel K, Gomez CA, Dhakal R, et al. Failure of primary atovaquone prophylaxis for prevention of toxoplasmosis in hematopoietic cell transplant recipients. *Transpl Infect Dis*. 2016;18:446-452.
  570. Peyron F, McLeod R, Ajzenberg D, et al. Congenital toxoplasmosis in France and the United States: one parasite, two diverging approaches. *PLoS Negl Trop Dis*. 2017;11:e0005222.
  571. Daffos F, Forestier F, Capella-Pavlovsky M, et al. Prenatal management of 746 pregnancies at risk for congenital toxoplasmosis. *N Engl J Med*. 1988;318:271-275.
  572. Desmonts G, Daffos F, Forestier F, et al. Prenatal diagnosis of congenital toxoplasmosis. *Lancet*. 1985;1:500-504.

## SHORT VIEW SUMMARY

**Definition**

- Giardiasis, caused by the protozoan *Giardia lamblia*, is a common cause of sporadic, endemic, and epidemic diarrhea throughout the world.
- Infected persons can have acute diarrhea with malaise, cramping and bloating, chronic diarrhea with malabsorption, or asymptomatic infection. Asymptomatic infection is most common in children, particularly in low-income settings, and may contribute to poor nutrition.

**Epidemiology**

- *Giardia* is one of the most widely distributed enteric parasites.
- In the United States there are approximately 17,000 documented infections annually, with estimated infections of more than 1.2 million; it is most frequently reported in children aged 1 to 9 years and adults aged 40 to 54 years.
- In low-income countries, *Giardia* infects nearly all children by the age of 10 years.
- Transmission of this fecal-oral parasite is most common through person-to-person transmission and through untreated or

inadequately treated surface water, and occasionally via contaminated food.

- Although specific human genotypes (assemblages A and B) are found in some animals, epidemics of giardiasis caused by animals have not been reliably observed. Animals typically harbor other assemblages.

**Diagnosis**

- A clinical syndrome of diarrhea and or other gastrointestinal symptoms lasting 7 to 10 days, and often associated with weight loss without fever, is helpful in distinguishing giardiasis from other enteric infections.
- Useful epidemiologic risk factors are a history of travel, drinking untreated surface water, having young children in daycare, or engaging in sexual practices with the potential for fecal-oral transmission.
- Stool examination for ova and parasites is the traditional method of diagnosis.
- Antigen detection is widely available and is more sensitive and specific than stool examination for ova and parasites. Polymerase chain reaction assays are becoming more widely used and can identify the assemblage.

**Therapy**

- Tinidazole is the drug of choice, taken in a single dose (see Table 279.3).
- Alternatives are metronidazole, nitazoxanide, and albendazole.
- Potential adverse events associated with treatment should be discussed with the patient.
- Treatment during pregnancy requires special considerations (see "Therapy").
- Post-*Giardia* irritable bowel syndrome, lactose intolerance, or another organism or cause should be considered if treatment appears to fail. If parasites are present, then re-treatment with a drug of a different class is usually successful.

**Prevention**

- Prevention of giardiasis requires the proper handling and treatment of drinking and recreational water supplies, good personal hygiene, and protection of food supplies from contamination.
- Bringing water to a boil, filtration, and careful halogenation can be effective measures to treat small volumes of water.

**DESCRIPTION OF THE PATHOGEN**


*Giardia* was probably described in the late 1600s, when van Leeuwenhoek likely discovered it in his own stool. It was in the early 1900s that the parasite received the genus name *Giardia*. The designated species name for the human parasite has been *lamblia*; *intestinalis* and *duodenalis* are also used. The genus *Giardia* is in the category of intestinal flagellates in the division Protozoa. The life cycle of *G. lamblia* has two stages: the motile dividing trophozoite present in the intestine of the host, and the environmentally resistant infectious cyst excreted in the feces. The trophozoite is 9 to 21  $\mu\text{m}$  long and 5 to 15  $\mu\text{m}$  wide (Fig. 279.1A). It has a convex dorsal surface and a flat ventral surface containing a disk with contractile proteins used for attachment (sucking or adhesive disk). There are four pairs of posteriorly directed flagella that are involved in locomotion and perhaps attachment.<sup>1-3</sup> The protozoan has two anteriorly placed diploid nuclei, each with a prominent central karyosome and a complete copy of the genome.<sup>4</sup> In stained preparations, the nuclei create the characteristic facelike image. Each of the two *Giardia* nuclei maintains a high degree of homozygosity, most likely through fusion and exchange of DNA during encystation.<sup>5,6</sup> Median bodies—tight collections of microtubules—are visualized transversely in a clawlike manner in *G. lamblia* and have been used to identify morphologically distinct types of *Giardia*. Of the *Giardia* spp., only *G. lamblia* has been successfully cultured in vitro.<sup>7</sup> Growth<sup>8</sup> is enhanced by the presence of biliary lipids

and a high concentration of cysteine, consistent with the predilection of *Giardia* for colonizing the upper small bowel. The trophozoite divides by longitudinal binary fission and has a doubling time in culture of 6 to 12 hours. It is an aerotolerant anaerobe that scavenges oxygen to enhance survival.<sup>9</sup> *Giardia* uses glucose as the major source of carbohydrate energy, metabolizing it to the end products of acetate, ethanol, alanine, and carbon dioxide; adenosine triphosphate (ATP) is generated during this process.<sup>8</sup> Metabolism of arginine via the arginine dihydrolase pathway is another mechanism for ATP generation.

The differentiation of *Giardia* into species traditionally depended on morphology and the host of origin, with a number of species described: *G. lamblia* in humans, *Giardia muris* in mice, *Giardia agilis* in amphibians, *Giardia psittaci* in parakeets, and *Giardia microti* in voles and muskrats (Table 279.1). However, molecular studies later revealed morphologically identical *Giardia* that were found in both humans and animals. The earliest molecular typing of these *Giardia* isolates divided them into at least three groups,<sup>10</sup> which were subsequently renamed and expanded into eight genotypes or referred to as assemblages A to H based on sequence differences in housekeeping genes<sup>11,12</sup> (see Table 279.1).<sup>12,13,14</sup> Assemblages A (often subtype AII) and B are associated with most but not all<sup>15</sup> human infections and a number of nonhuman mammalian hosts.<sup>11,13,14</sup> Their biology, biochemistry, and molecular mechanisms differ and fit criteria to be separate species.<sup>16,17</sup> Occasional infections with non-A, non-B assemblages, including assemblage E, have been reported in humans.<sup>15</sup> Improved sequencing methods using a broader sampling of isolates will better define genetic diversity and range of *Giardia* genotypes.<sup>12,15,18</sup>

<sup>a</sup>All material in this chapter is in the public domain, with the exception of any borrowed figures or tables.





**FIG. 279.1** *Giardia lamblia* trophozoite (A) and cyst (B) are demonstrated in a trichrome stain of fecal material. Note the prominent nuclei in the trophozoite. In the cyst, centrally located axonemes, a clawlike median body, and two nuclei can be seen.

**TABLE 279.1** Typing of *Giardia* Species

NAME	GENOTYPE	HOST RANGE
<i>Giardia agilis</i>		Amphibians
<i>Giardia ardeae</i>		Birds
<i>Giardia microti</i>		Muskrats and voles
<i>Giardia muris</i>		Rodents
<i>Giardia psittaci</i>		Birds
<i>Giardia lamblia</i>	Assemblage A	Humans, primates, dogs, cats, cattle, sheep, deer, rodents
	Assemblage B	Humans, primates, dogs, cattle, horses, beaver
	Assemblage C	Dogs
	Assemblage D	Dogs
	Assemblage E	Cattle, goats, sheep, pig
	Assemblage F	Cats
	Assemblage G	Rodents
	Assemblage H	Marine vertebrates

Modified from Feng Y, Xiao L. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. Clin Microbiol Rev. 2011;24:110–140.

Different *G. lamblia* genotypes have unique capacities to establish infection and to cause disease in animal models and experimental human infections, but whether specific genotypes differ in virulence in human natural infections<sup>8,12,19–22</sup> is unclear.

### Some Unique Biologic Aspects

The genome of the prototypic assemblage A isolate, WB (<http://giardiadb.org>),<sup>23</sup> is approximately 11.7 Mb, distributed over five chromosomes. Analysis of its small subunit ribosomal RNA<sup>8,24</sup> sequence and genes of certain pathways<sup>25</sup> indicate that it is one of the earliest-branching eukaryotes.<sup>23</sup> The genome is compact and simplified, including DNA synthesis, transcription, RNA processing, and cell-cycle machinery. The parasite contains only a few introns, extremely short promoters, sometimes 60 bp or less, overlapping transcriptional controlling elements, and mitosomes, remnants of earlier mitochondria.<sup>26,27</sup> Pathways to synthesize phospholipids, fatty acids, cholesterol, and purine and pyrimidine nucleosides are reduced or lacking, so these essential metabolic building blocks are largely scavenged from the small intestine milieu. There are only enzymes sufficient for remodeling but not synthesis of lipid membrane components.<sup>8,23</sup> Except for alanine, amino acids are taken up from the environment.

Because *Giardia* is an early-branching, highly adapted organism and an important disease-causing parasite, many aspects of its unique biology

have been extensively studied and reviewed elsewhere.<sup>8</sup> However, a few important examples of its particularities include surface antigenic variation, a unique vesicular transport system devoid of a Golgi apparatus, and the processes of encystation and excystation.

*Giardia* is the only known intestinal dwelling organism to undergo frequent surface antigenic variation.<sup>28,29</sup> A single variant-specific surface protein (VSP) is expressed on the surface of *Giardia* and, after several generations, is replaced by another VSP.<sup>29–32</sup> VSPs are a family of about 250 cysteine-rich proteins with similar motifs and general structure and a conserved transmembrane carboxyl-terminal sequence.<sup>30,33</sup> Although all VSPs are transcribed, only one is expressed. The others VSPs are possibly eliminated by interference RNA mechanisms.<sup>32,34</sup> Antigenic variation occurs during infection of humans and animals, in vitro in culture, and during encystation.<sup>8,30</sup> Although it is important in evading the host immune response, certain VSPs are favored or disallowed even in the absence of an adaptive immune response.<sup>35</sup> Specific VSPs seem to have unique physical-chemical characteristics that allow them to be biologically selected and expressed or disfavored and not expressed, depending on the intestinal environment and host.<sup>35</sup>

*G. lamblia* trophozoites encyst to form fibrillary, oval, thin-walled cysts 8 to 12  $\mu$ m long and 7 to 10  $\mu$ m wide (see Fig. 279.1B). *Giardia* encystation is a model system for vesicular transport and developmental biology because *Giardia* lacks a typical Golgi apparatus that is an essential component of vesicular transport in most higher organisms.<sup>5,36–38</sup> Encystation induced in vitro<sup>39</sup> is a complex process involving many of *Giardia*'s basic processes.<sup>5,40,41,42</sup> The early phase is characterized by the formation of highly characteristic encystment vesicles (ESVs), in which cyst wall proteins (CWPs) 1 through 3 and other molecules are processed.<sup>36,43,44</sup> The late phase of encystation consists of transport of the CWPs to the cell surface, their assembly with *N*-acetylgalactosamine (GalNAc) into the cyst wall, and then nuclear division and DNA replication without cell division so that cysts are generated with a ploidy of 16N.<sup>5,38,39,41,45</sup> At ultrastructural analysis, the cyst wall contains CWPs and GalNAc polymer in insoluble fibrils.<sup>46</sup>

Excystation is a highly coordinated process that is initiated when cysts are exposed to environmental stimuli, such as gastric acid and pancreatic enzymes.<sup>5,47</sup> The process is complex and well described morphologically but not well understood on a molecular level.<sup>48–50</sup> An excyzoite containing four nuclei (4N each) is released; this divides twice without further DNA replication, resulting in four daughter trophozoites.<sup>5,51</sup>

### EPIDEMIOLOGY

*Giardia* is widely distributed. In the United States, *G. lamblia* has been demonstrated in 4% to 7% of submitted stool specimens, making it the most commonly identified intestinal parasite. US. epidemiologic surveys of giardiasis from 2011 to 2012 found that 15,223 to 16,868 cases were reported annually, with the case rates in different states varying from 1.7 to 29.2 cases per 100,000 persons and an overall mean of 5.8.<sup>52</sup> Because of substantial underreporting, a current estimate is that more than 1.2 million cases may occur annually, accounting for an estimated

\$34 million dollars in annual hospitalization costs.<sup>53,54</sup> Only 1.3% of cases were associated with a detected outbreak.<sup>52</sup> *Giardia* was most frequently reported in children 1 to 9 years old and to a lesser extent in adults 40 to 54 years old, and during the early summer through early fall months<sup>52</sup>; these findings are similar to those in Canada and other high-income countries.<sup>55</sup> In the United States, coenteric infections with *Giardia* are uncommon (2.2%).<sup>56</sup>

In low-income regions of the world, *Giardia* is one of the first enteric pathogens to infect infants.<sup>57–60,61</sup> In these settings, incidence ranges from 37.7% to 96.4% in the first 2 years of life, and 40% of these children have persistent (>8 weeks) infection. Peak prevalence rates of 15% to 30% occur in children younger than 10 years.<sup>62,63,64,65,66</sup> Nearly all children in these settings become infected, and *Giardia* is frequently detected together with other viral, bacterial, and parasitic pathogens.<sup>60,67</sup>

Natural acquisition of the parasite occurs through fecal-oral ingestion of *Giardia* cysts through contaminated water, food, or person-to-person transmission. *Giardia* cysts survive well in the environment, particularly in cold water. *Giardia* was the most frequent cause of outbreaks of diarrhea resulting from drinking water in the United States from 1971 to 2006<sup>68</sup>; however, the risk from drinking water dropped substantially in the 1980s and continues to steadily decline<sup>69</sup> as improved measures for water treatment have been implemented. Recreational water,<sup>70</sup> primarily in treated swimming venues and lakes, is also a risk for *Giardia*. Surface water can become contaminated by human or animal sources<sup>69,71</sup> so backpackers who do not adequately treat water can become infected. Most waterborne outbreaks can be traced to untreated or improperly treated water or faulty purification systems.<sup>68,72,73</sup> Food handler contamination at restaurants due to poor hand hygiene, particularly with produce, has been linked to the majority of foodborne outbreaks.<sup>69,74,75</sup>

Person-to-person transmission occurs in groups with close contact and an increase in fecal contamination, especially between young children and their parents and siblings, in men who have sex with men, and between young children in daycare centers. Historically, the prevalence of *Giardia* in children in daycare has been as high as 20% to 50%,<sup>76</sup> which seems to have decreased in the United States lately. Many young children are asymptomatic, but they can spread the infection within their homes and contribute to secondary spread within their communities.<sup>77,78</sup>

*Giardia* was the pathogen identified most frequently in returning travelers and immigrants seeking medical care and diagnosed with infectious gastrointestinal disease—diagnosed at a rate of 31 per 1000 ill returned adult travelers.<sup>79</sup> Travel to the Indian subcontinent was the highest-risk destination.

Many mammalian hosts such as gerbils, mice, beavers, sheep, cattle, dogs, and cats (see Table 279.1)<sup>11,12,14,80,81</sup> have harbored or been experimentally infected with *Giardia* morphologically similar to, and in some instances identical to, the genotypes infecting humans. Genetic subtyping indicates the low zoonotic potential for most animal isolates.<sup>12,18</sup> The strongest epidemiologic link for animal isolates associated with outbreaks of human infection were Canadian waterborne outbreaks of giardiasis linked to beavers.<sup>82,83</sup>

## **PATHOGENESIS AND IMMUNE RESPONSE**

*Giardia* was once thought to be a harmless commensal, but its association with symptomatic diarrhea, malabsorption in children, disease after waterborne outbreaks, travel, and experimental human infections<sup>19</sup> has clearly established its pathogenicity. Pathophysiologic mechanisms need to explain the variety of symptoms and clinical outcomes that occur in exposed and infected populations, the high level of asymptomatic infections in endemic regions compared with common symptomatic epidemic disease in primarily nonendemic intermittently exposed populations, and the variety of infection rates and symptoms in singular populations. Although there is a large body of experimental data, as a group the studies are particularly difficult to interpret because of the variability of the models and methods employed. Most nonhuman experimental infections have used a variety of rodent and small mammal models with either *G. muris* in mice or rats or different human *Giardia* isolates in mice, gerbils, or rats. Frequently, the isolates used were not well characterized. Many of the findings, particularly those from studies of immunity, are based on murine host responses and extrapolated to

human infections. The emerging picture reveals a complex interaction between a genetically diverse, adaptable group of closely related, morphologically similar organisms and their hosts.

Infection occurs after oral ingestion of as few as 10 to 25 cysts.<sup>84</sup> The ability to establish infection and to cause diarrhea varies among isolates, assemblages, and possibly VSP expression patterns.<sup>8,19–21,35,85</sup> After excystation, trophozoites adhere to the brush border of the epithelium via their ventral disk<sup>1</sup> and multiply in the upper small bowel. Adherence can be modified or prevented by means of a number of mechanisms.<sup>86</sup> The parasite may avoid being expelled by mechanical means.<sup>87</sup> Small bowel biopsies in most cases show little inflammation. However, some may demonstrate villous shortening and atrophy<sup>23,88</sup>; and the degree of histologic abnormality correlates with severity of clinical manifestations.<sup>89–91</sup> Among the postulated mechanisms of trophozoite pathogenesis in the small intestine—mucosal invasion, elaboration of an enterotoxin, and disruption of the intestinal epithelial brush border and function—there is little reproducible evidence that mucosal invasion or production of an enterotoxin occurs. *G. lamblia* does disrupt brush border functional enzymes, including disaccharidases in vitro in human small intestinal epithelial cell monolayers.<sup>92</sup> Trophozoites also disrupt tight junctions, increase permeability, and induce apoptosis.<sup>93–96,97</sup> Many of these effects are caspase-3 dependent.<sup>93,94</sup> Epithelial barrier dysfunction and apoptosis have also been seen in duodenal biopsy specimens from persons with giardiasis and some murine models of persistent infection.<sup>98,99</sup>

The effects of *Giardia* on epithelial cells in vitro appear nutrient dependent. High glucose concentrations can counteract caspase-3–dependent epithelial cell apoptosis through upregulation of the sodium-dependent glucose cotransporter in response to *Giardia* proteases.<sup>100</sup> *Giardia*-mediated arginine consumption reduces epithelial cell proliferation.<sup>101</sup> Arginine consumption may favor parasite colonization by suppressing arginine-dependent nitric oxide synthetase in the host, thus limiting the protective effect of nitric oxide.<sup>102,103</sup> These interactions may partially explain dietary influences on epithelial cell responses and *G. lamblia* persistence in mice.<sup>99,104</sup>

Host immunity plays a role in clearance of the parasite,<sup>51</sup> protection against reinfection, and pathogenesis.<sup>51</sup> In part, the findings are dependent on the model system that is used. Immune responses in humans, especially during acute illness, are not well studied, and much of the information is extrapolated from mouse and gerbil models of infection with *G. muris* or *G. lamblia*. There are differences in the immune responses of mice to infection with a rodent *G. muris* compared with infections with *G. lamblia* isolated from humans.<sup>99</sup> Immune responses and course of infection also differ among *G. lamblia* isolates of different assemblages.<sup>105,106</sup> Time-dependent clearance of *G. lamblia* in murine models coincides with upregulation of intestinal genes related to  $\alpha$ -defensin function, mast cell proteases, and antibody production.<sup>107</sup> Early mechanisms of control of *G. lamblia*-infected mice include interleukin-6 (IL-6).<sup>108,109</sup> This effect may be mediated via IL-6–producing mast cells and subsequent increased intestinal motility.<sup>110–112</sup> In addition, dendritic cells producing IL-6 can reverse the inability of IL-6–deficient mice to clear *G. lamblia*.<sup>113</sup> Innate immune-mediated signaling through TLR2 may counteract parasite clearance and in turn promote infection.<sup>114</sup> T-cell responses control infection, whereas B-cell responses appear to be more important at later time points.<sup>115</sup>

Several observations indicate that partial protective immunity can develop to *Giardia* infection. Protective immunity to reinfection develops in most rodent and gerbil infection models; partial immunity also develops in some situations in humans.<sup>19,35,105</sup> Lower rates of symptomatic disease were found in long-term residents of endemic areas of North America than in visitors or short-term residents.<sup>116,117</sup> In contrast, reinfection is common, particularly in children, in endemic settings.<sup>59,60,64,118</sup> Previously infected volunteers can be reinfected with the identical isolate after treatment but at a lower intensity and without symptoms.<sup>19</sup> Possible reasons for the ability to become reinfected are exposure to a different genotype, expression of a different VSP by an individual isolate, age and nutritional status of the host, and inability to develop intestinal immunity after a single infection.

Both B-cell and T-cell immunity appear to play essential roles in protective immunity,<sup>110</sup> but which arm of the immune system is dominant

depends on the infection model studied. A systemic antibody response occurs in individuals with *Giardia*. Although serum immunoglobulin M (IgM) and IgG antibodies develop and with complement can be lethal to *Giardia* trophozoites,<sup>29</sup> it is likely that gastrointestinal IgA antibodies secreted into the lumen where trophozoites are located play a more important role.<sup>119–121</sup> Failure to develop IgA against specific *Giardia* antigens has been suggested to correlate with chronic giardiasis in humans.<sup>122</sup> Evidence supporting a role for humoral antibodies, specifically IgA responses, in the control of *Giardia* infection initially has come from the increase in chronic giardiasis in patients with common variable immunodeficiency (CVID).<sup>123</sup> Because these patients may also have cellular immune defects, this association neither supports nor refutes humoral mechanisms. Patients living with acquired immunodeficiency syndrome (AIDS) do not have an increased susceptibility to *Giardia*, although they uncommonly develop severe disease that is difficult to treat.<sup>124,125</sup> More recent studies have demonstrated that natural exposure in immunocompetent adults leads to development of CD4<sup>+</sup> memory T-cell responses.<sup>126,127</sup> These cells produce greater IL-17A after stimulation with *Giardia* antigens in exposed compared with unexposed individuals.<sup>126</sup>

In mice, immunization with the full repertoire of VSPs expressed by a single isolate can lead to protection against homologous *Giardia* challenge.<sup>128</sup> This effect coincides with the development of broad anti-VSP antibodies, but the mechanisms are likely multifactorial. Either complete absence of IgA or defective transcellular transport due to deletion of the polymeric immunoglobulin receptor (pIgR) is associated with inability to resolve infection in humans.<sup>110,119,121,129</sup> The mechanism by which IgA prevents or helps to clear infection is probably by binding to trophozoites and preventing a critical adherence step. However, there is no evidence that IgA can kill trophozoites, and studies comparing IgA-expressing or total B-cell-deficient mice have demonstrated that B-cell-mediated clearance of *Giardia* is not limited to IgA production.<sup>121</sup>

More recently, IL17A has been identified as a key regulator of host defense responses against *Giardia* in animal models. Consequences of impaired IL-17A signaling through the cognate IL17RA receptor include defective transcellular IgA transport through pIgR together with reductions in several antimicrobial peptides.<sup>130,131</sup> Similar to humans, IL17A-producing CD4<sup>+</sup> T cells increase in mice following *Giardia* challenge. However, CD4<sup>+</sup> T-cell-depleted mice are unable to clear parasites despite robust IL17A upregulation, likely from innate sources.<sup>130</sup>

The role for T-cell responses in parasite control and pathogenesis is substantial. Athymic, T-cell-deficient mice are unable to clear infection with *G. muris* or *G. lamblia* until the mice are reconstituted with lymphoid cells, particularly the CD4<sup>+</sup> helper T lymphocyte.<sup>115,132,133</sup> After reconstitution, animals develop an abnormal intestinal histologic appearance that parallels the changes seen in some humans with giardiasis: flattening of villi, crypt hypertrophy, and a mononuclear cell infiltration of the submucosa.<sup>89,134</sup> Similarly, CD4<sup>+</sup> T-cell neutralization prolongs infection but also attenuates brush border defects in *Giardia*-infected mice.<sup>106</sup>

The complex host immune responses to *Giardia*, and immunomodulatory properties of the parasite itself, influence mucosal responses and disease manifestations of other pathogens in vitro and in animal models. The cathepsin B-like cysteine protease produced by some *Giardia* strains is capable of cleaving mucosal-derived IL-8, limiting neutrophil chemotaxis, and attenuating proinflammatory intestinal responses to other stimuli.<sup>104,135,136</sup> In a murine malnutrition model, sequential coinfection with *Giardia* and then enteroaggregative *Escherichia coli* enhanced weight loss and partially altered the mucosal responses observed during enteroaggregative *E. coli* infection alone.<sup>137</sup>

There is increasing recognition that resident microbiota influence both *Giardia* susceptibility and pathogenesis. Singer and Nash made the first sentinel observation that murine microbiota are capable of protecting against infection even in severely immunocompromised mice.<sup>138</sup> In addition, some *Lactobacillus* strains limit growth of *Giardia* through bile-salt hydrolases.<sup>139</sup> Microbiota, however, also contribute to *Giardia* pathogenesis. Jejunal aspirate cultures from patients with symptomatic giardiasis are more pathogenic to germ-free mice than axenic trophozoites (assemblage A).<sup>140</sup> These manifestations may be the result of direct microbial interactions. *Giardia* alters the intestinal bacterial composition throughout the gut of mice<sup>141</sup> and perturbs gut microbial

amino acid metabolism in vivo.<sup>137</sup> Culturing otherwise commensal bacteria in the presence of *Giardia* or its products functionally promotes their pathogenic potential.<sup>142,143</sup> In some models, increases in mucosal-associated bacteria and myeloid inflammation persist beyond parasite clearance.<sup>97,144</sup> In contrast, antibiotics that reduce intestinal bacteria result in diminished CD8<sup>+</sup> T-cell activity and disaccharidase deficiency and prevent host growth impairment after *Giardia* challenge, even during malnutrition.<sup>137,145</sup> Whether and to what extent these clearly relevant interactions between *Giardia* and other microbes influence outcomes in humans, including post-*Giardia* infection sequelae, require further investigation.

## CLINICAL MANIFESTATIONS

The clinical manifestations of giardiasis are broad and range from asymptomatic cyst passage to acute diarrhea and dehydration necessitating hospitalization. Infections may be short or prolonged, resulting in spruelike symptoms of chronic diarrhea with malabsorption. The proportion of individuals manifesting symptoms in a population appears to be influenced by *Giardia* prevalence and possibly age at first exposure, with significant variability in nonendemic versus endemic populations, the host nutritional and immune status and genetic makeup, coinfection with other enteric organisms, and the assemblage or strain of the prevalent *Giardia*. The natural history of *Giardia* infections in nonendemic populations is uncertain because most patients have not been rigorously followed without treatment and because of the demonstrated variability of different *Giardia* genotypes to cause disease. Patients with or without symptoms may self-cure or become chronically infected, sometimes lasting years.<sup>146</sup>

After the ingestion of *G. lamblia* cysts, there is an incubation period of 1 to 2 weeks before the onset of symptoms. The prepatent and incubation periods after instillation of 50,000 axenically grown trophozoites into the small intestine can be as short as 5 and 6 days, respectively.<sup>19</sup> Symptoms may occur before detection of cysts.<sup>19</sup> Symptomatic giardiasis is characterized by the acute onset of diarrhea, abdominal cramps, bloating, and flatulence (Table 279.2). The patient usually has malaise, nausea, and anorexia and may report sulfuric belching. Foul-smelling flatus and stools are common complaints, as is weight loss. Vomiting and upper gastrointestinal symptoms occur less frequently but may sometimes be a dominant symptom, leading to a delay in diagnosis. Fever and tenesmus are atypical and should raise suspicions for an alternative diagnosis. Initially, stools may be profuse and watery; the median number of stools per day has been as high as nine.<sup>147</sup> In short-term travelers, giardiasis is an infrequent cause of diarrhea compared with bacterial causes; it is more common in long-term travelers. Gross blood, pus, mucus, and

TABLE 279.2 Symptoms of Giardiasis

	PERCENTAGE	RANGE
Diarrhea	89	64–100
Malaise	84	72–97
Flatulence	74	35–97
Foul-smelling, greasy stools	72	57–79
Abdominal cramps	70	44–85
Bloating	69	42–97
Nausea	68	59–79
Anorexia	64	41–82
Weight loss	64	56–73
Vomiting	27	17–36
Fever	13	0–21
Urticaria	9	4–14
Constipation	9	0–17

Data from Hill DR. Giardiasis. Issues in diagnosis and management. Infect Dis Clin North Am. 1993;7:503–525.



microscopic polymorphonuclear white blood cells (WBCs) are absent, and, if present, suggest another or concomitant diagnosis.

One of the most important distinguishing features of giardiasis is the prolonged duration of diarrhea, which is commonly intermittent with waxing and waning symptoms and usually is accompanied by weight loss. At the time of presentation, most patients have been symptomatic for more than 1 week to 10 days.<sup>147</sup> Weight loss of about 10 pounds occurs in more than 50% of patients with prolonged symptoms.<sup>148</sup> Unusual reported associations include urticaria, other skin rashes, reactive arthritis,<sup>149</sup> eye complaints, biliary tract disease, and gastric infection.<sup>147</sup> These are uncommon, and it is unclear whether a casual relationship exists. Gastric infection occurs in the presence of achlorhydria and has been seen in conjunction with *Helicobacter pylori* in both adult and child populations.<sup>150,151</sup>

Although most persons with giardiasis have a relatively benign course, some can have an acute severe illness necessitating hospitalization. Children and pregnant women were particularly affected in the United States,<sup>152</sup> but not in a similar study of those hospitalized in Scotland.<sup>153</sup>

Patients who develop chronic diarrhea may report weight loss, profound malaise, lassitude, occasional headache, and diffuse abdominal and epigastric discomfort, often exacerbated by eating. Stools, passed frequently in small volume, commonly because of malabsorption, may be greasy and foul smelling or frothy and yellowish. Waxing and waning symptoms over months is characteristic. Infected children can present with spruelike disease and failure to thrive.<sup>91,154</sup> Consequently, malabsorption of vitamins A and B<sub>12</sub>, protein, D-xylose, and iron develops.<sup>155,156</sup> Lactase is the most common disaccharidase deficiency, occurring in 20% to 40% of cases,<sup>157</sup> with post-*Giardia* lactose intolerance sometimes persisting for several weeks after treatment. This is often confused with relapse or reinfection. Some individuals may be intolerant to specific fermentable oligosaccharides, disaccharides, monosaccharides, and polyols.<sup>158</sup>

Post-*Giardia* irritable bowel syndrome (IBS) is increasingly recognized in nonendemic settings. Both IBS-like symptoms and chronic fatigue have reportedly lasted for years after primary infection in a previously unexposed nonendemic population in Norway.<sup>159</sup> These *Giardia*-exposed patients with persisting symptoms have developed elevated serum sCD40L, a finding that warrants confirmation in other populations.<sup>160</sup> Within a US insurance database, a single episode of giardiasis increased the likelihood of a new IBS diagnosis within the next year (hazard ratio [HR], 4.8; 95% confidence interval [CI], 3.6–6.4) that was only slightly attenuated after adjustment for anxiety, depression, and health care utilization (HR, 3.9; 95% CI, 2.9–5.4).<sup>161</sup>

*Giardia* is a nearly universal infection in children residing in low-income regions of the world. Despite the high prevalence of *Giardia* infection in these children, the vast majority of infections are asymptomatic; the variability in manifestations has been perplexing.<sup>162</sup> Although many *Giardia* infections in this setting do not result in acute diarrhea, there is evidence that the parasite is associated with persistent diarrhea and nutritional abnormalities. In a review and meta-analysis of studies describing *Giardia* in children younger than 5 years, *Giardia* was not seen to cause acute diarrhea (odds ratio [OR], 0.60; 95% CI, 0.38–0.94;  $P < .03$ ) but was associated with diarrhea lasting at least 14 days (persistent diarrhea; OR, 3.2; 95% CI, 1.50–6.76;  $P < .001$ ).<sup>163</sup> In several subsequent case-controls studies of pathogen-attributable causes of pediatric diarrhea in these settings, *Giardia* was detected more often in controls than cases of moderate-to-severe acute diarrhea,<sup>164,165</sup> a finding that has not been universally observed<sup>160</sup> and merits more analysis. Unlike viruses, *Cryptosporidium*, and other gastrointestinal pathogens associated with acute diarrhea,<sup>166</sup> there is no clear relationship between quantitative *Giardia* burden or stool inflammatory markers and symptomatic infection in these children.<sup>160,167</sup>

Persistent *Giardia* infection, either with or without diarrhea, could be part of the larger condition of childhood environmental enteric dysfunction in low-income regions. Studies over the last 20 years have documented stunting in Brazilian and Ecuadorian children infected with *Giardia*,<sup>59,168</sup> poor intestinal permeability in Nepali children,<sup>169</sup> low weight-for-age and height-for-age in Brazilian children with persistent symptomatic giardiasis,<sup>63,170,171</sup> underweight status in Rwandan children,<sup>172</sup> significant wasting in Malaysian and Indian children,<sup>62,173,174</sup> and decreased

cognitive function in Peruvian children with multiple episodes of giardiasis.<sup>58</sup> In a prospective study of 629 children in Bangladesh where 74% of children had had a *Giardia* exposure by 2 years of life, *Giardia* in the first 6 months of life, regardless of symptoms, was associated with decreased length-for-age scores.<sup>61</sup> In a larger multicenter study including >2000 children, early *Giardia* exposures and persistent infections were also associated with impaired growth attainment by 2 years of life.<sup>60</sup> The findings of these studies suggest that treatment of infection might help to reverse some of the deleterious effects.<sup>59,175–177</sup> However, persistent infections and reinfections occur so rapidly that repeated therapy is currently impractical and not indicated.<sup>60,64,178,179–180</sup>

In contrast, longitudinal studies in Bangladeshi children<sup>181,182</sup> and Peruvian children<sup>64</sup> did not demonstrate effects on well-being or nutrition, and in a setting where children were well-nourished, asymptomatic infection did not have apparent deleterious effects.<sup>183,184</sup> Furthermore, many children and adults will be simultaneously infected with other bacterial, viral, and parasitic infections, and there is the potential for adverse synergy.<sup>185,186</sup>

## DIAGNOSIS

The diagnosis of giardiasis should be considered in all patients with prolonged diarrhea, particularly if it is associated with malabsorption or weight loss, a history of recent travel to an endemic area, ingestion or exposure to untreated water sources, the presence of small children in the home who attend daycare centers, or sexual risk factors. In the usual patient with uncomplicated giardiasis, the WBC count is normal without eosinophilia, and gastrointestinal imaging is nondiagnostic. The diagnosis depends on detection of the parasite or its specific antigen or DNA in the feces.

Microscopic detection of cysts or trophozoites in feces for ova and parasites (O&P) is the traditional way to diagnose most gastrointestinal parasites. The O&P test is time-consuming and costly; the sensitivity is dependent on the number of organisms, the skill of the microscopist, the amount of feces examined, and the time taken to examine the stool. A 90% chance or greater of detecting a true *Giardia* infection requires three stool examinations over multiple days.<sup>148,187,188</sup> Tests requiring little training with high sensitivity and specificity and ease of performance such as assays to detect the *Giardia* soluble specific cyst wall antigen(s) of *Giardia* or cyst detection with immunofluorescence are widely available,<sup>189,190</sup> reproducible, and rapidly performed.<sup>191</sup> Polymerase chain reaction (PCR)-based diagnostic testing,<sup>12,192,193,194,195</sup> with *Giardia* usually included as one target in a larger array of gastrointestinal pathogens, has become commercially available in the United States. Comparisons between the *Giardia* antigen tests and PCR testing indicate that PCR testing is at least as sensitive and in some formats twice as sensitive<sup>166</sup> as antigen detection tests.

The detection of *Giardia* antigen in stools has proved to be extremely useful in diagnosing clinically significant infections. These assays are most helpful when giardiasis is the leading consideration. They are often less expensive than an O&P examination and are 85% to 98% sensitive and 90% to 100% specific.<sup>189,190,196–200</sup> Various methods to detect *Giardia* trophozoites in the small intestine were previously used to establish or confirm the diagnosis. Because of the availability of highly sensitive noninvasive tests, small intestinal sampling is rarely required.

Testing for systemic anti-*Giardia* antibody is neither generally available nor useful in the individual case because of persistence of antibodies from prior infections. Detection of IgG antibodies has been useful in seroepidemiologic studies.<sup>201–204</sup>

## THERAPY

Susceptibility testing of *Giardia* is not performed because of the difficulty in establishing cultures from clinical specimens and the lack of standardization of sensitivity testing.<sup>205</sup> Drug resistance can be induced in vitro<sup>206</sup>; however, the significance of this is unclear because some isolates that appear clinically resistant are susceptible in vitro, and vice versa. Most information on drug efficacy, therefore, is provided by clinical trials and cumulative experience.<sup>205,207,208,209</sup> The drug of choice for the therapy of giardiasis in the United States is tinidazole (Tindamax; Mission Pharmacal, San Antonio, TX) (Table 279.3). Like metronidazole, it is a member of the nitroimidazole family. It has high efficacy (approximately

**TABLE 279.3 Treatment of Giardiasis**

DRUG (FDA PREGNANCY CATEGORY) <sup>a</sup>	DOSAGE	
	ADULT	PEDIATRIC
Tinidazole (C)	2 g, single dose	50 mg/kg, single dose (maximum, 2 g)
Metronidazole <sup>b</sup> (B)	250 mg tid × 5–7 days	5 mg/kg tid × 7 days
Nitazoxanide (B)	500 mg bid × 3 days	Age 12–47 mo: 100 mg bid × 3 days Age 4–11 yr: 200 mg bid × 3 days
Albendazole <sup>b</sup> (C)	400 mg qd × 5 days	15 mg/kg/day × 5–7 days (maximum, 400 mg)
Paromomycin <sup>b</sup> (NC)	500 mg tid × 5–10 days	30 mg/kg/day in 3 doses × 5–10 days
Quinacrine <sup>c</sup> (C)	100 mg tid × 5–7 days	2 mg/kg tid × 7 days
Furazolidone <sup>c</sup> (C)	100 mg qid × 7–10 days	1.25–2 mg/kg qid × 10 days

<sup>a</sup>FDA pregnancy categories: B, Animal reproduction studies have failed to demonstrate a risk to the fetus, and there are no adequate and well-controlled studies in pregnant women. C, Animal reproduction studies have shown an adverse effect on the fetus, and there are no adequate and well-controlled studies in humans, but potential benefits may warrant use of the drug in pregnant women despite potential risks.

<sup>b</sup>Not an FDA-approved indication. Maximum of 8.8 mg/kg/day and not recommended for <1 month of age.

<sup>c</sup>No longer produced in the United States; may be obtained from some compounding pharmacies.

FDA, US Food and Drug Administration; NC, not categorized.

90%) and a favorable side effect profile, and because of a longer half-life than metronidazole, it can be given once daily.<sup>205</sup> Tinidazole is effective as a single dose for both children and adults and was approved in 2004.<sup>210</sup> Although the US Food and Drug Administration (FDA) has never approved metronidazole for giardiasis, there are over 50 years of experience in using it.

Metronidazole is given in divided doses for 5 to 7 days, with an efficacy of 80% to 95%. Adverse effects for tinidazole and metronidazole are similar: a metallic taste; nausea; dizziness; headache; and, rarely, reversible neutropenia, peripheral neuropathy, or seizures. When taken with alcohol, they can produce a disulfiram-like effect. High-dose, short-course regimens of metronidazole have lower efficacy rates and are sometimes poorly tolerated. There have been concerns about potential mutagenicity of metronidazole; however, this has not been documented in humans.<sup>211</sup> There have been reports of increasing nitroimidazole treatment failures.<sup>207</sup>

The mechanism of action of metronidazole and presumably tinidazole is reductive activation of the nitro group by acceptance of electrons from parasite ferredoxins, although other pathways to activate the nitro group may be involved.<sup>212–214</sup> Nitazoxanide (Alinia; Romark Laboratories, Tampa, FL) was approved in 2003 for use in pediatric giardiasis and cryptosporidiosis and, more recently, for giardiasis treatment in adults.<sup>215</sup> There is limited clinical experience with this drug compared with the nitroimidazoles; efficacy is in the range of 65% to 85% with a 3-day treatment course.<sup>216,217</sup> Its mechanism of action is inhibition of pyruvate-ferredoxin oxidoreductases; however, there does not appear to be cross-resistance with metronidazole-resistant parasites.<sup>218</sup> Its main side effect is gastrointestinal upset.

Meta-analyses of a limited number of trials of albendazole (400 mg for 5 days), a benzimidazole, compared with standard regimens of metronidazole have shown comparable results.<sup>208,219</sup> Albendazole is attractive because it is usually better tolerated than the nitroimidazoles, has efficacy against many intestinal helminths, and is used in low-income regions to decrease intestinal parasitism. However, more clinical experience will be needed to determine its place in therapy.

Quinacrine and furazolidone are no longer manufactured in the United States. Quinacrine has an efficacy of more than 90%, and can be obtained through compounding pharmacies; it is given in divided doses for 5 to 7 days (see Table 279.3). The most common side effects are bitter taste, nausea, vomiting, and abdominal cramping. Yellow

discoloration of the skin, urine, and sclerae can occasionally occur. Exfoliative dermatitis is rare. Psychosis occurs uncommonly, and we avoid its use in patients with a significant psychiatric history. Furazolidone has a success rate of about 80% to 85% but may cause gastrointestinal side effects and brown urine and may cause mild hemolysis in glucose-6-phosphate dehydrogenase-deficient individuals.

For patients in whom one drug course fails or who infrequently relapse, a switch to a drug from a different class is generally effective.<sup>205,220</sup> For the unusual patient who is not cured with single-drug therapies, combination treatment, often with metronidazole and quinacrine, can be effective.<sup>125,221,222</sup> A randomized trial comparing metronidazole alone versus metronidazole plus the probiotic *Saccharomyces boulardii* demonstrated higher cure rates in the combination group.<sup>223</sup> When refractory infection is suspected, it is important to establish true persistence by obtaining a repeat stool examination to distinguish it from post-*Giardia* IBS or lactose intolerance. For pregnant women with giardiasis, there is no consistently recommended therapy because of the theoretical adverse effects of anti-*Giardia* drugs on the fetus.<sup>205</sup> When disease is mild and hydration and nutrition can be maintained, therapy can be delayed until after delivery, or at least until after the first trimester. If treatment is necessary, paromomycin, an oral nonabsorbable aminoglycoside,<sup>224</sup> can be tried. In limited clinical experience, this approach has an efficacy rate of 60% to 70%.<sup>224</sup> Paromomycin is given in divided doses for 5 to 10 days (see Table 279.3). Metronidazole has been used extensively in pregnancy for the treatment of trichomoniasis. The teratogenic effect appears to be minimal and, if present, is greatest during the first trimester, when both metronidazole and tinidazole should not be used.<sup>205,225</sup> If therapy cannot be avoided, then metronidazole, tinidazole, and nitazoxanide can probably be used safely in the last two trimesters of pregnancy. Table 279.3 lists the FDA pregnancy categories for each of the agents.

There is no established role for nonpharmaceutical therapies. A number of studies have found various effects on *Giardia* infections, disease, or interactions with other gut pathogens as a result of micronutrient administration.<sup>167,226–228</sup> The interpretation and generality of the effects are unclear and require confirmation. However, the findings of these studies suggest a complex interaction of micronutrients with the pathogenicity of enteric organisms.

A defined management approach to chronic post-*Giardia* infection chronic sequelae in nonendemic populations has yet to be established.<sup>229</sup>

## PREVENTION

The prevention of giardiasis requires proper handling and treatment of water used for communities and good personal hygiene. Although halogenation (e.g., chlorine and iodine) alone is sufficient to kill *G. lamblia* cysts, important variables, such as water temperature, clarity, pH, and contact time, alter the efficacy of chlorine, and higher chlorine levels (4–6 mg/L) may be required.<sup>72,230</sup> Thus, in addition to chlorination, public water supplies intended for drinking should also be subjected to coagulation-flocculation, sedimentation, and filtration.<sup>69</sup>

Travelers to low-income regions of the world or to wilderness areas should consider all water potentially contaminated because of the wide array of animal and human reservoirs of giardiasis. Bringing water to a boil is sufficient to kill all protozoal cysts; at high altitudes, boiling for a minute is reasonable. If boiling is impossible, halogenation is generally effective for *Giardia*, but *Cryptosporidium* is resistant.<sup>72,230</sup> There are many chlorine-based (e.g., chlorine bleach: 5% to 6%, 2 to 4 drops/L) or iodine-based (e.g., tincture of iodine: 2%, 5 drops/L) preparations in liquid or tablet form.<sup>230</sup> Contact time should be increased for water that is cold, and the concentration of halogen should be increased for turbid water. Personal water filters with pores of an “absolute” micron size of 0.2 to 1.0  $\mu\text{m}$  can be used.<sup>230</sup> Filtered water should also be halogenated to kill enteric viruses if they are considered a risk. Uncooked foods that may have been washed or prepared in contaminated water should be avoided.

Giardiasis outbreaks in daycare centers can be difficult to resolve. Some recommend that only symptomatic children be treated; however, infected children transmit *Giardia* to parents and family members, with further potential spread to the community. Therefore, each situation requires an individual decision. If strict hand washing and treatment of