


MOTIVATION

- ▶ General background
 - ▶ The terminology of mathematics
 - ▶ Formalized by the branch of math called ‘Number Theory’.
- ▶ Why do economists’ care?
 - ▶ Economists express values, sets, and concepts using numerical (quantitative) values.
- ▶ Application in this career
 - ▶ Throughout your whole experience.

NUMBERS



1. COMMON NUMBER SETS

Real Numbers


Rational $\frac{5}{3}$ 0.63 $0.0\overline{12}$	Irrational
Integers $\{\dots, -2, -1, 0, 1, 2, \dots\}$	$\sqrt{3}$ π 0.10010001...
Whole $\{0, 1, 2, 3, \dots\}$	
Natural $\{1, 2, 3, \dots\}$	

2. REAL NUMBERS

- ▶ Two binary operators for the Reals
 - ▶ Addition: $a + b$
 - ▶ Multiplication: $a \cdot b$
- ▶ Properties:
 - ▶ Commutative: $\forall a, b \in \mathbb{R}, a + b = b + a.$
 - ▶ Commutative: $\forall a, b, c \in \mathbb{R}, (a + b) + c = a + (b + c).$
 - ▶ Zero Exists: $\forall a \in \mathbb{R}, a + 0 = a.$
 - ▶ Negation Exists: $\forall a \in \mathbb{R}, \exists(-a) \in \mathbb{R} : a + (-a) = 0.$
 - ▶ Distributive: $\forall a, b, c \in \mathbb{R}, a \cdot (b + c) = (a \cdot b) + (a \cdot c).$
- ▶ Reciprocal: $\forall a \in \mathbb{R}, \frac{1}{a}$



3. ABSOLUTE VALUE AND NUMBER LINE

- ▶ Absolute Value: $|\pm a| = a$.
- ▶ Properties:
 - ▶ $\forall a > 0 \in \mathbb{R}, |a| = a$.
 - ▶ $\forall a = 0 \in \mathbb{R}, |a| = 0$.
 - ▶ $\forall a < 0 \in \mathbb{R}, |a| = -a$.
 - ▶ $|ab| = |a||b|$.
 - ▶ $|a|^2 = |a^2|$.
 - ▶ If $|a| \leq c \iff -c \leq a \leq c$.




4. TRIANGLE INEQUALITY

- $\forall a, b \in \mathbb{R} \rightarrow |a + b| \leq |a| + |b|.$
- Corollaries:
 - $||a| - |b|| \leq |a - b|.$
 - $|a - b| \leq |a| + |b|.$
- Visual aid [Click Me].



5. NEIGHBORHOODS

- ▶ Let $a, \varepsilon \in \mathbb{R}, \varepsilon > 0$.
- ▶ Let the ε -neighborhood of a be the set $V_\varepsilon(a) := \{x \in \mathbb{R} : |x - a| < \varepsilon\}$.
- ▶ I.e., x is a value within the neighborhood of a such that it is within ε distance from a .



PRACTICE: NUMBERS

1.

MOTIVATION

- ▶ General background
 - ▶ How we map sets onto other sets.
 - ▶ The most common to represent relationships between variables.
- ▶ Why do economists' care?
 - ▶ This is primary way that we represent preferences, utility, and production.
 - ▶ Functions are at the core of how theorems are represented.
- ▶ Application in this career
 - ▶ Both throughout the development and presentation of proofs, as well as in estimating equations from data.

MOTIVATION

- ▶ General background
 - ▶ How we map sets onto other sets.
 - ▶ The most common to represent relationships between variables.
- ▶ Why do economists' care?
 - ▶ This is primary way that we represent preferences, utility, and production.
 - ▶ Functions are at the core of how theorems are represented.
- ▶ Application in this career
 - ▶ Both throughout the development and presentation of proofs, as well as in estimating equations from data.

OVERVIEW

1. Relations
2. Correspondences and Functions
3. Injective & Surjective Functions
4. Composition of Functions
5. Inverse Functions
6. Image and Pre-image
7. Homogeneity
8. Level, Superior & Inferior Sets
9. Euler's Theorem
10. Quasiconcavity & Quasiconvexity
11. Concavity & Convexity
12. Continuity
13. Upper- and Lower-Hemicontinuity
14. Brouwer's Fixed-point Theorem
15. Kakutani's Fixed-point Theorem

1. RELATIONS

- ▶ A collection of ordered-pairs (s, t) has a binary relation $s\mathcal{R}t$ between sets S and T .
 - ▶ Reflexive: $\forall x \in S, x\mathcal{R}x$.
 - ▶ Symmetric: $\forall x, y \in S, x\mathcal{R}y \implies y\mathcal{R}x$.
 - ▶ Complete: $\forall x, y \in S \rightarrow x\mathcal{R}y \vee y\mathcal{R}x$.
 - ▶ Transitive: $\forall x, y, z \in \mathcal{R}, x\mathcal{R}y \wedge y\mathcal{R}z \implies x\mathcal{R}z$.
- ▶ Equivalent Relation ($=$): Is reflexive, symmetric, and transitive.
- ▶ Common Relations:
 - ▶ Equal: $=$
 - ▶ Equivalent: \equiv
 - ▶ Better than: $>$
 - ▶ Less than: $<$


2. CORRESPONDENCES AND FUNCTIONS

- ▶ Correspondence: A relation that associates each element of one set (the domain) to the elements of another set (the range).
- ▶ Function: A relation that associates each element in the domain to a single, unique element of the range.
- ▶ Onto: Every element in the range is mapped into a point in the domain.
- ▶ One-to-one: Every element in the range is assigned only a single point in the domain.

$$f : D \rightarrow R$$


3. INJECTIVE & SURJECTIVE FUNCTIONS

- ▶ Function $f : A \rightarrow B$:
 - ▶ Injective (one-to-one):
 $\forall a_1, a_2 \in A, a_1 \neq a_2 \implies f(a_1) \neq f(a_2).$
 - ▶ Surjective (onto B): $\forall b \in B, \exists a \in A : f(a) = b.$
 - ▶ Bijective: Both injective and surjective.




NOT a
Function

A has many B




General
Function

B can have many A




Injective
(not surjective)

B can't have many A



Surjective
(not injective)

Every B has some A



Bijective
(injective, surjective)

A to B, perfectly

4. COMPOSITION OF FUNCTIONS

- ▶ $f : A \rightarrow B$ and $g : B \rightarrow C$, then

$$g \circ f(x) = g(f(x)) : A \rightarrow C$$

- ▶ Follows from associative property of functions:

$$(h \circ g) \circ f = h \circ (g \circ f)$$

- ▶ If f and g are surjective, then $g \circ f$ is surjective.

5. INVERSE FUNCTIONS

- If $f : A \rightarrow B$ is bijective, then the inverse function is $f^{-1} : B \rightarrow A$.


$$f^{-1} \circ f(x) = x$$

$$f \circ f^{-1}(x) = x$$

- $f : A \rightarrow B$ is bijective iff the inverse f^{-1} is a function $f^{-1} : B \rightarrow A$.


6. IMAGE AND PRE-IMAGE

- ▶ Let $f : A \rightarrow B$.
- ▶ Image: If $X \subseteq A$ is set $f(X) = \{f(x) : x \in X\} \subseteq B$.
- ▶ Pre-image: If $Y \subseteq B$ is set $f^{-1}(Y) = \{x \in A : f(x) \in Y\} \subseteq A$.



8. LEVEL, SUPERIOR, & INFERIOR SETS

- ▶ Level set: $L(y^0) \equiv \{x|x \in D, f(x)=y^0\}$.
- ▶ Superior set: $S(y^0) \equiv \{x|x \in D, f(x) \geq y^0\}$.
- ▶ Inferior set: $I(y^0) \equiv \{x|x \in D, f(x) \leq y^0\}$.



9. EULER'S THEOREM

- ▶ Let $f(x_1, x_2, \dots, x_N)$ be homogeneous of degree r , and differentiable.


$$\nabla f(\bar{x}) \cdot \bar{x} = \sum_{n=1}^N \frac{\partial f(\bar{x}_1, \dots, \bar{x}_N)}{\partial x_n} \bar{x}_n = rf(\bar{x}_1, \dots, \bar{x}_N)$$

- ▶ **Application:** In production theory, Euler's theorem states that a production function homogeneous of degree 1 (CRS) with factors paid their marginal product will have no surplus or deficit in total product.


10. QUASICONCAVITY & QUASICONVEXITY

- ▶ Quasiconcavity: $\forall x_1, x_2 \in D, f : D \rightarrow R$ iff
$$f(tx_1 + (1 - t)x_2) \geq \min[f(x_1), f(x_2)] \forall t \in [0, 1]$$
- ▶ Quasiconvexity: $\forall x_1, x_2 \in D, f : D \rightarrow R$ iff
$$f(tx_1 + (1 - t)x_2) \leq \max[f(x_1), f(x_2)] \forall t \in [0, 1]$$
- ▶ These become **strict** when the inequalities hold for all $x_1 \neq x_2$.


QUASICONCAVE AND QUASICONVEX



QUASICONCAVE BUT NOT QUASICONVEX




QUASICONCAVE IN TWO DIMENSIONS




QUASICONCAVE IN TWO DIMENSIONS

- $f : D \rightarrow \mathbb{R}$ is quasiconcave iff $S(y)$ is a **convex** set for all $y \in \mathbb{R}$.




QUASICONVEX IN TWO DIMENSIONS



QUASICONVEX IN TWO DIMENSIONS

- $f : D \rightarrow \mathbb{R}$ is quasiconvex iff $I(y)$ is a **convex** set for all $y \in \mathbb{R}$.



11. CONCAVITY & CONVEXITY

- ▶ f is defined on a convex subset
 $D \subset \mathbb{R}^n \forall x_1, x_2 \in D, \forall t \in [0, 1]:$

- ▶ Concave:


$$f(tx_1 + (1 - t)x_2) \geq tf(x_1) + (1 - t)f(x_2)$$

- ▶ Convex:


$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2)$$

- ▶ Strict concavity or convexity when the inequality holds.


CONCAVITY



CONVEXITY





CONCAVE UP (CONVEX) & CONCAVE DOWN



12. CONTINUITY

- Continuous: $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ at $x_0 \in \mathbb{R}^m$ if whenever $\{x_n\}_{n=1}^\infty$ is a sequence in \mathbb{R}^m which converges to x_0 , **then** the sequence $\{f(x_n)\}_{n=1}^\infty$ in \mathbb{R}^n converges to $f(x_0)$.


$$\forall \varepsilon > 0, \exists \delta > 0 : \forall x \in A, [|x - x_0| < \delta] \implies [|f(x) - f(x_0)| < \varepsilon]$$




13. UPPER- AND LOWER-HEMICONTINUITY

- ▶ $A \subset \mathbb{R}^n$ and a closed set $Y \subset \mathbb{R}^n$.
 - ▶ Upper Hemicontinuous: Correspondence $f : A \rightarrow Y$ if it has a closed graph and the images of compact sets are bounded.
 $\forall B \subset A, f(B) = \{y \in Y : y \in f(x) \exists x \in B\}$ is bounded.
- ▶ Lower Hemicontinuous: Correspondence $f : A \rightarrow Y$ if for every sequence $x^m \rightarrow x \in A$ with $x^m \in A \forall m$, and every $y \in f(x)$, we can find a sequence $y^m \rightarrow y$ and an integer $M : y^m \in f(x^m) \forall m > M$.
- ▶ Continuous: Both upper- and lower-hemicontinuous.

13. UPPER-HEMICONTINUITY




UHC




Not UHC


13. UPPER-HEMICONTINUITY



13. UPPER- AND LOWER-HEMICONTINUITY




Not upper hemicontinuous at x_1
Not upper hemicontinuous at x_2
Lower hemicontinuous



Not lower hemicontinuous at x_1	
Not lower hemicontinuous at x_2	
Upper hemicontinuous	


13. NOT UPPER- AND LOWER-HEMICONTINUITY



Not upper hemicontinuous at x_1 and x_2
Not lower hemicontinuous at x_1 and x_2



14. BROUWER'S FIXED-POINT THEOREM

- ▶ Suppose $D \subset \mathbb{R}^m$ is non-empty, compact, convex set, and $f : D \rightarrow D$ is a **continuous function**.
- ▶ Then $f(\cdot)$ has a fixed point, e.g., there is an $x \in D : x = f(x)$.



15. KAKUTANI'S FIXED-POINT THEOREM

- ▶ Suppose $D \subset \mathbb{R}^m$ is non-empty, compact, convex set, and $f : D \rightarrow D$ is a **upper-hemicontinuous correspondence** with the property $f(x) \subset D$ is non-empty and convex for all $x \in D$.
- ▶ Then $f(\cdot)$ has a fixed point, e.g., there is an $x \in D : x = f(x)$.



PRACTICE: FUNCTIONS

1.

REVIEW OF NUMBERS

1. Real Numbers
2. Common Number Sets
3. Absolute Value and Number Line
4. Triangle Inequality
5. Neighborhoods

REVIEW OF FUNCTIONS

1. Relations
2. Correspondences and Functions
3. Injective & Surjective Functions
4. Composition of Functions
5. Inverse Functions
6. Image and Pre-image
7. Homogeneity
8. Level, Superior & Inferior Sets
9. Euler's Theorem
10. Quasiconcavity & Quasiconvexity
11. Concavity & Convexity
12. Continuity
13. Upper- and Lower-Hemicontinuity
14. Brouwer's Fixed-point Theorem
15. Kakutani's Fixed-point Theorem

