

Lecture 09

Optimization Day 1

Ryan McWay[†]

[†] *Applied Economics,
University of Minnesota*

Mathematics Review Course, Summer 2023
University of Minnesota
August 17th, 2023

LAST LECTURE REVIEW

- ▶ Numbers:
 - ▶ Triangle Inequality
 - ▶ Neighborhoods
- ▶ Functions:
 - ▶ Homogeneity
 - ▶ Euler's Theorem
 - ▶ Quasiconcavity & Quasiconvexity
 - ▶ Concavity & Convexity
 - ▶ Continuity
 - ▶ Upper- and Lower-Hemicontinuity
 - ▶ Brouwer's Fixed-point Theorem
 - ▶ Kakutani's Fixed-point Theorem

REVIEW ASSIGNMENT


1. Problem Set 08 solutions are available on Github.
2. Any issues or problems **You** would like to discuss?

DAILY ICEBREAKER

- ▶ Attendance via prompt:
 - ▶ Name
 - ▶ Program and track
 - ▶ Daily icebreaker subject...



OPTIMIZATION



Unconstrained Optimization

MOTIVATION

- ▶ General background
 - ▶ Simplest form of optimization.
 - ▶ Solution involves finding critical points of the function.
- ▶ Why do economists' care?
 - ▶ Used in simple applications for testing first order conditions and second order conditions.
- ▶ Application in this career
 - ▶ Unconstrained optimization is primarily used to determine FOC and SOC.

MOTIVATION

- ▶ General background
 - ▶ Simplest form of optimization.
 - ▶ Solution involves finding critical points of the function.
- ▶ Why do economists' care?
 - ▶ Used in simple applications for testing first order conditions and second order conditions.
- ▶ Application in this career
 - ▶ Unconstrained optimization is primarily used to determine FOC and SOC.

MOTIVATION

- ▶ General background
 - ▶ Simplest form of optimization.
 - ▶ Solution involves finding critical points of the function.
- ▶ Why do economists' care?
 - ▶ Used in simple applications for testing first order conditions and second order conditions.
- ▶ Application in this career
 - ▶ Unconstrained optimization is primarily used to determine FOC and SOC.

OVERVIEW

1. First Order Conditions
2. Second Order Conditions
3. Global Min and Max

1. FIRST ORDER CONDITIONS

- ▶ Let $F : U \rightarrow \mathbb{R}$ be a differentiable defined on subset U of \mathbb{R}^n .
- ▶ If $x^* \in \mathbb{R}^n$ is a local minimum or local maximum of $F(\cdot)$ and x^* is an interior point of U , then:


$$\nabla F(x^*) = 0 \text{ or } \frac{\partial F(x^*)}{\partial x_n} \forall n$$

- ▶ The FOC can be summarized by the Jacobian matrix.

$$J = \nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \cdots & \frac{\partial f}{\partial x_n} \end{bmatrix}$$

2. SECOND ORDER CONDITIONS

- ▶ Let $F : U \rightarrow \mathbb{R}$ be C^2 whose domain is an open set $U \in \mathbb{R}^n$.
- ▶ Suppose $\nabla F(x^*) = 0$.
 - ▶ If $D^2f(x^*)$ is negative (positive) definite, then x^* is a strict local max (min).
 - ▶ If $D^2f(x^*)$ is indefinite, then x^* is neither a local max nor min.
- ▶ Suppose x^* is a local max (min) of F .
 - ▶ Then, $\nabla F(x^*) = 0$ and the symmetric $n \times n$ matrix $D^2f(x^*)$ is negative (positive) **semi-definite**.



The graph of the indefinite form $Q_3(x_1, x_2) = x_1^2 - x_2^2$.

2. SECOND ORDER CONDITIONS

- The SOC is captured by the Hessian matrix.

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}$$

3. GLOBAL MIN AND MAX

- ▶ Any point x^* of a concave (convex) function $f(\cdot)$ satisfying $\nabla F(x^*) = 0$.
- ▶ Note this may be a boundary point (corner solution) or an interior point (critical point).

| $Df(x^*)$ | $D^2f(x^*)$ | Max/Min |
|-----------|------------------------|---------------------------|
| $= 0$ | Negative Semi-definite | Local Max |
| $= 0$ | Positive Semi-definite | Local Min |
| $= 0$ | Indefinite | Saddle point or Inflexion |

PRACTICE: UNCONSTRAINED OPTIMIZATION

1.

(Equality) Constrained Optimization

MOTIVATION

- ▶ General background
 - ▶ Optimization of a function considering a constraint from other function(s).
 - ▶ Leads to ‘interior solutions’.
- ▶ Why do economists’ care?
 - ▶ This is the most typical case for optimization.
- ▶ Application in this career
 - ▶ Used throughout microeconomics.

MOTIVATION

- ▶ General background
 - ▶ Optimization of a function considering a constraint from other function(s).
 - ▶ Leads to ‘interior solutions’.
- ▶ Why do economists’ care?
 - ▶ This is the most typical case for optimization.
- ▶ Application in this career
 - ▶ Used throughout microeconomics.

MOTIVATION

- ▶ General background
 - ▶ Optimization of a function considering a constraint from other function(s).
 - ▶ Leads to ‘interior solutions’.
- ▶ Why do economists’ care?
 - ▶ This is the most typical case for optimization.
- ▶ Application in this career
 - ▶ Used throughout microeconomics.

OVERVIEW

1. Lagrangian Method
2. Second Order Conditions
3. Bordered Hessian

1. LAGRANGIAN METHOD

- ▶ Let f and h be C^1 functions.
- ▶ Suppose $x^* = (x_1^*, x_2^*)$ is a **solution** to the problem:

$$\begin{aligned} \max & f(x_1, x_2) \\ \text{s.t. } & h(x_1, x_2) = c \end{aligned}$$

- ▶ Consider (x_1^*, x_2^*) are **not** critical points of h .
- ▶ Then μ^* is a real number such that (x_1^*, x_2^*, μ^*) **is** a critical point of the following Lagrangian function.

$$L(x_1, x_2, \mu) \equiv f(x_1, x_2) - \mu[h(x_1, x_2) - c]$$


- ▶ That is...

$$\frac{\partial L}{\partial x_1} = 0, \quad \frac{\partial L}{\partial x_2} = 0, \quad \frac{\partial L}{\partial \mu} = 0$$

1. LAGRANGIAN METHOD

► Intuition:

$$\mu = -\frac{\frac{\partial f}{\partial x_1}(x^*)}{\frac{\partial f}{\partial x_2}(x^*)} = -\frac{\frac{\partial h}{\partial x_1}(x^*)}{\frac{\partial h}{\partial x_2}(x^*)}$$



1. LAGRANGIAN METHOD

- So, we have two equations represented by $\nabla f(x^*) = \mu^* \nabla h(x^*)$.

$$\frac{\partial f}{\partial x_1}(x^*) - \mu^* \frac{\partial h}{\partial x_1}(x^*) = 0$$

$$\frac{\partial f}{\partial x_2}(x^*) - \mu^* \frac{\partial h}{\partial x_2}(x^*) = 0$$

1. LAGRANGIAN METHOD

- Consider there may be **many** constraints in the constraint set C_h with local max/min $x^* \in C_h$.

$$C_h = \{x = (x_1, \dots, x_n) : (h_1(x) = a_1), \dots, h_m(x) = a_m)\}$$

- If x^* is not the critical point of $h = (h_1, \dots, h_m)$ (i.e., $\text{rank}(Dh(x^*)) < m$) – unique value for all constraints), then there are μ_1^*, \dots, μ_m^* real numbers such that $(x_1^*, \dots, x_n^*, \mu_1^*, \dots, \mu_m^*)$ is the critical point of the Lagrangian function:

$$L(x^*, \mu^*) \equiv f(x) - \mu_1[h(x) - a_1] - \dots - \mu_m[h(x) - a_m]$$

- Alternatively, all partials for x_n and μ_m are set to zero.

2. SECOND ORDER CONDITIONS

- ▶ To ensure a maximum, we need to know that the second differential of the objective function f at the critical point is **decreasing along the constraint**.
- ▶ Let $y = f(x_1, x_2(x_1))$ be the value of the objective function subject to the constraint.
- ▶ By the implicit function theorem:

$$\frac{dx_2}{dx_1} = \frac{\partial h / \partial x_1}{\partial h / \partial x_2}$$

- ▶ By chain rule:

$$\frac{dy}{dx_1} = \frac{\partial f}{\partial x_1} + \frac{\partial f}{\partial x_2} \frac{\partial x_2}{\partial x_1} = \frac{\partial f}{\partial x_1} + \frac{\partial f}{\partial x_2} \frac{\partial h / \partial x_1}{\partial h / \partial x_2}$$

3. BORDERED

- ▶ The sufficient condition for a critical point is:

$$\frac{d^2y}{dx_1^2} < 0$$

- ▶ It can be shown that

$$\frac{d^2y}{dx_1^2} = \frac{-1}{(\partial h / \partial x_2)^2} \bar{D}$$

- ▶ Where \bar{D} is the **bordered Hessian** of L .
- ▶ The determinant of the bordered Hessian tests for quasiconcavity or convexity to ensure a local max/min
- ▶ Negative definite assuming $\nabla f(x) \neq 0 \forall x$ and $f(\cdot)$ is strictly quasiconcave.

3. BORDERED

- Local minima (x_1^*, x_2^*) if $B(x_1^*, x_2^*, \lambda^*) < 0$. Local maximum (x_1^*, x_2^*) if $B(x_1^*, x_2^*, \lambda^*) > 0$.

$$B(x_1, x_2, \lambda) = \left| \begin{bmatrix} 0 & \frac{\partial h}{\partial x_1} & \frac{\partial h}{\partial x_2} \\ \frac{\partial h}{\partial x_1} & \frac{\partial^2 L}{\partial x_1^2} & \frac{\partial^2 L}{\partial x_1 \partial x_2} \\ \frac{\partial h}{\partial x_2} & \frac{\partial^2 L}{\partial x_2 \partial x_1} & \frac{\partial^2 L}{\partial x_2^2} \end{bmatrix} \right| > 0$$

PRACTICE: (EQUALITY) CONSTRAINED OPTIMIZATION

1.

UNCONSTRAINED OPTIMIZATION

1. First Order Conditions
2. Second Order Conditions
3. Global Min and Max

(EQUALITY) CONSTRAINED OPTIMIZATION

1. Lagrangian Method
2. Second Order Conditions
3. Bordered Hessian

ASSIGNMENT

- ▶ Assignment:
 - ▶ Problem Set 09 (PS09)
 - ▶ Solution set will be available following end of Lecture 10
- ▶ Struggling?
 1. Read the ‘Encouraged Reading’
 2. Review ‘Supplementary material’
 3. Reach out directly