

## Obesity and Other Cancers

Lin Yang, Bettina F. Drake, and Graham A. Colditz

Lin Yang, Bettina F. Drake, and Graham A. Colditz, Washington University School of Medicine and Siteman Cancer Center, St Louis, MO; and Lin Yang, Center for Public Health, Medical University of Vienna, Vienna, Austria.

Published online ahead of print at [www.jco.org](http://www.jco.org) on November 7, 2016.

Authors' disclosures of potential conflicts of interest are found in the article online at [www.jco.org](http://www.jco.org). Author contributions are found at the end of this article.

Corresponding author: Graham A. Colditz, MD, DrPH, Washington University School of Medicine, 660 S Euclid Ave, Campus Box 8100, St Louis, MO 63110; e-mail: colditzg@wustl.edu.

© 2016 by American Society of Clinical Oncology

0732-183X/16/3435w-4231w/\$20.00

DOI: 10.1200/JCO.2016.68.4837

### A B S T R A C T

#### Purpose

Evidence on overweight, obesity, and an increased risk of cancer continues to accumulate and was updated in the 2016 handbook on weight control from the International Agency for Research on Cancer (IARC). The underlying primary data, together with dose-response meta-analysis and, finally, pooled analysis of individual participant data, add insight into the relation between obesity and cancer risk and prognosis. We summarize the evidence for mortality from prostate cancer, hematologic malignancies, and kidney cancer.

#### Methods

We reviewed pooled analysis of rare end points across cohorts, regardless of primary results reported from the individual studies, further reducing risk of publication bias. Of these cancer sites, only kidney cancer was included in the IARC 2002 report, although mortality from prostate cancer and hematologic malignancies was noted in the American Cancer Society prospective cohort study in 2003. The 2016 update from the IARC added details for prostate and hematologic malignancies, classifying the evidence as sufficient to conclude that avoiding excess body fatness lowers the risk of multiple myeloma but found that the evidence for it lowering the risk of prostate cancer mortality or diffuse large B-cell lymphoma was limited.

#### Results

A higher body mass index is associated with an increased risk of advanced prostate cancer and prostate cancer mortality and is associated with worse survival in most subtypes of hematologic malignancies, in a dose-response fashion. Evidence for kidney cancer is built mostly on retrospective data, which supports an obesity paradox in patients with the clear cell variant; however, population-based cohort data indicate that a higher cohort-entry body mass index is associated with worse kidney cancer-specific survival.

#### Conclusion

Together, these data add support to the evidence for a growing cancer burden caused by adiposity in both early adult and later adult life, yet leave open the question of the means of weight management after diagnosis as a strategy to improve survival.

*J Clin Oncol* 34:4231-4237. © 2016 by American Society of Clinical Oncology

### INTRODUCTION

Here we review the current research addressing the role of adiposity/obesity in mortality and prognostic outcomes in prostate, hematologic, and renal cancers. Excess weight or adiposity is a common risk factor for cancer, progression, and nonsurvival; this provides a unique opportunity to address a modifiable risk factor through primary and secondary interventions. Evidence reviewed by the International Agency for Research on Cancer (IARC) and the World Cancer Research Fund support the finding that being overweight or obese is an established cause of several cancers including breast, endometrium, esophagus

(adenocarcinoma), renal, and colon and rectum.<sup>1</sup> Since the IARC report in 2002, evidence has expanded to suggest an association between a higher body mass index (BMI) and advanced prostate or prostate cancer mortality and several hematologic cancers.<sup>2</sup> The 2016 IARC update on body fatness and cancer concluded that the evidence was sufficient to conclude that avoiding excess body fatness lowers the risk of multiple myeloma but evidence for its lowering the risk of prostate cancer mortality and diffuse large B-cell lymphoma is limited. Thus, for these cancer sites, the committee could not rule out bias and confounding as contributing to the positive associations observed.<sup>2</sup> In addition, excessive weight is a risk factor for cancer mortality overall.<sup>3</sup> To

first summarize the current evidence on cause, we placed the greatest emphasis on prospective cohort studies reported as pooled analyses of individual participant data. This approach reduces variation in analytic approaches among studies. Furthermore, for rare cancers such as multiple myeloma, it allows all cohorts to contribute end points regardless of whether they have published results or not, hence reducing the publication bias that can distort meta-analyses limited to results already

published in the literature. In this review, we summarize studies on obesity and the incidence of prostate, hematologic, and renal cancers. We are not aware of any randomized controlled trials of weight-loss interventions after diagnosis of cancer at these sites. We summarize the observational evidence on adiposity/obesity and cancer prognosis in Table 1, and describe associations for each included cancer site in greater detail.

**Table 1.** Summary of Observational Studies on Adiposity/Obesity and Cancer Prognosis

Cancer type	Prospective Cohort*	Patient Cohort† and Retrospective Cohort‡
Prostate cancer		
Cancer-specific survival	Meta-analysis <sup>4</sup> : RR, 1.15 (95% CI, 1.06 to 1.25) per 5 kg/m <sup>2</sup> increased BMI	Meta-analysis <sup>4</sup> : RR, 1.20 (95% CI, 0.99 to 1.46) per 5 kg/m <sup>2</sup> increased BMI
Hematologic malignancies		
Lymphoma		
Cancer-specific survival	Meta-analysis <sup>5</sup> : RR, 1.14 (95% CI, 1.04 to 1.26) per 5 kg/m <sup>2</sup> increased BMI	
Leukemia		
Overall survival	Meta-analysis <sup>6</sup> : RR, 1.29 (95% CI, 1.11 to 1.49) in obese (BMI ≥ 30 kg/m <sup>2</sup> ) v normal-weight (18.5 kg/m <sup>2</sup> ≤ BMI < 25 kg/m <sup>2</sup> ) patients RR <sub>men</sub> , 1.45 (95% CI, 1.22 to 1.72) RR <sub>women</sub> , 1.14 (95% CI, 0.99 to 1.33)	Pooled patient cohort <sup>7</sup> : HR, 1.72 (95% CI, 1.15 to 2.58) in obese (BMI ≥ 30 kg/m <sup>2</sup> ) v nonobese (BMI < 30 kg/m <sup>2</sup> ) patients with APL
Relapse		Cohort of patients with APL <sup>8</sup> : HR, 2.45 (95% CI, 1.00 to 5.99) in overweight/obese (BMI ≥ 25 kg/m <sup>2</sup> ) v under/normal-weight (BMI < 25 kg/m <sup>2</sup> ) patients
Pediatric leukemia		
Overall survival		Meta-analysis <sup>9</sup> : HR, 1.30 (95% CI, 1.16 to 1.46) in obese (BMI > 95%) v nonobese (BMI ≤ 95%) patients
Event-free survival		Meta-analysis <sup>9</sup> : HR, 1.46 (95% CI, 1.29 to 1.64) obese (BMI > 95%) v nonobese (BMI ≤ 95%) patients
Multiple myeloma		
Cancer-specific survival	Meta-analysis <sup>10</sup> : RR, 1.15 (95% CI, 1.04 to 1.27) in overweight (25.0 kg/m <sup>2</sup> ≤ BMI < 30 kg/m <sup>2</sup> ) patients; RR, 1.54 (95% CI, 1.35 to 1.76) in obese (BMI ≥ 30 kg/m <sup>2</sup> ) v normal-weight (18.5 kg/m <sup>2</sup> ≤ BMI < 25 kg/m <sup>2</sup> ) patients Pooled analysis <sup>11</sup> : HR, 1.09 (95% CI, 1.03 to 1.16) per 5 kg/m <sup>2</sup> increased cohort-entry BMI; HR, 1.22 (95% CI, 1.09 to 1.35) per 5 kg/m <sup>2</sup> increased early-adulthood BMI; HR, 1.06 (95% CI, 1.02 to 1.10) per 5 cm increased waist circumference.	
Kidney cancer		
Overall survival		Patient cohort <sup>12</sup> : HR, 0.50 (95% CI, 0.31 to 0.81) in obese (30 kg/m <sup>2</sup> ≤ BMI < 35 kg/m <sup>2</sup> ) patients; HR, 0.24 (95% CI, 0.09 to 0.60) in severely obese (BMI ≥ 35 kg/m <sup>2</sup> ) v nonobese (BMI < 30 kg/m <sup>2</sup> ) patients Meta-analysis of retrospective studies <sup>13</sup> : HR, 0.57 (95% CI, 0.43 to 0.76) in highest v lowest BMI category
Cancer-specific survival	Million Women cohort <sup>14</sup> : RR, 1.65 (95% CI, 1.28 to 2.13) per 10 kg/m <sup>2</sup> increased BMI	Patient cohort <sup>15</sup> : HR, 0.87 (95% CI, 0.82 to 0.94) per 1 kg/m <sup>2</sup> increased BMI in patients with clear cell variant; HR, 1.32 (95% CI, 1.03 to 1.70) per 1 kg/m <sup>2</sup> increased BMI in patients with chromophobe variant; no association in patients with papillary variant Meta-analysis of retrospective studies <sup>13</sup> : HR, 0.59 (95% CI, 0.48 to 0.74) in highest v lowest BMI category
Recurrence-free survival		Meta-analysis of retrospective studies <sup>13</sup> : HR, 0.49 (95% CI, 0.30 to 0.81) in highest v lowest BMI category

Abbreviations: APL, acute promyelocytic leukemia; BMI, body mass index; HR, hazard ratio; RR, relative risk.

\*Cancer-free at cohort entry, BMI measured before diagnosis.

†BMI measured at diagnosis.

‡BMI measured after diagnosis.

## OBESITY AND INCIDENCE OF PROSTATE, HEMATOLOGIC, AND RENAL CANCERS

There is no clear link between obesity and overall prostate cancer incidence. A meta-analysis of 27 cohorts estimated the association between obesity and prostate cancer risk, reporting a relative risk of 1.03 [95% CI, 1.00 to 1.07] per 5 kg/m<sup>2</sup> increase in BMI ( $P = .11$ ) with varied results across cohort studies.<sup>16</sup> A growing body of research indicates that obesity is associated with an increased risk of advanced prostate cancer.<sup>17,18</sup> The IARC update concluded that the evidence for an association between excess body weight and fatal prostate cancer was limited.<sup>2</sup> Meta-analytic data of prospective cohort studies suggest a modest but consistent direct effect of BMI on the incidence of lymphoma, multiple myeloma, and adult leukemia.<sup>16,5,10,19</sup> Risk increases in a dose-response fashion. It is also important to note that the adverse impact of BMI on the risk of lymphoma and multiple myeloma may start growing early, during young adulthood. On the basis of a review of the evidence, the IARC stated in 2016 that the evidence was sufficient to conclude that an absence of excess body fatness lowers the risk of multiple myeloma. Mounting evidence for kidney cancer continues to show increasing risk with increasing BMI in a dose-response fashion.<sup>16,20,21</sup>

## OBESITY AND PROSTATE CANCER PROGNOSIS

A growing body of research indicates that obesity is associated with worse pathologic outcomes in prostate cancer; however, evidence of an association has varied across cohort studies. Analysis of 57 European and American prospective studies shows an overall RR of prostate cancer mortality of 1.13 (95% CI, 1.02 to 1.24) for a 5 kg/m<sup>2</sup> increase in BMI.<sup>22</sup> In contrast, in an analysis of Asian cohorts, obesity was not related to prostate cancer mortality.<sup>23</sup> Obesity is linked to hormonal imbalances and hormonal factors that may influence prostate cancer progression.<sup>24,25</sup> Several mechanisms that may contribute to the growth of prostate cancer, including adiponectin levels<sup>26</sup> and inflammatory mediators,<sup>27</sup> have also been explored for relations with survival and mortality.

Epidemiologic studies show that obesity is associated with an increased risk of prostate cancer aggressiveness,<sup>28-32</sup> progression,<sup>32-36</sup> and cancer-specific mortality.<sup>27,32,37-39</sup> Chalfin et al<sup>40</sup> found that obese men are more likely to have worse pathology and a higher risk of recurrence of their cancer. Furthermore, Asmar et al<sup>41</sup> and Ly et al<sup>42</sup> reported that increased BMI was significantly associated with a higher risk of biochemical failure.

Incidence rates for prostate cancer are 60% higher for African-American men,<sup>43</sup> who also have the highest prostate cancer mortality. Prostate cancer accounts for 44% of the overall cancer mortality disparity between African-American and white men.<sup>44</sup> Studies suggest that obesity may contribute to this disparity in prostate cancer outcomes.<sup>31,4,45-47</sup> African-American men have higher rates of obesity compared with white men.<sup>48</sup> Obesity is also associated with both prostate tumor biology and a delayed prostate-specific antigen diagnosis of prostate cancer. Studies have also found that in men with low-risk disease, obese African-American men have a higher risk of recurrence compared with obese white men.<sup>49</sup>

Although physical activity, especially vigorous activity, is inversely related to the incidence of aggressive prostate cancer<sup>50-56</sup> and may also be associated with a decreased overall incidence of prostate cancer,<sup>57</sup> associations between obesity and prostate cancer seem to be independent of the level of physical activity.

Continued study is warranted to clarify the associations reported to date and to identify additional factors that may modify the relationship between obesity and prostate cancer prognosis. Short follow-up and lack of control for smoking are major limitations in many of these studies, especially in data from clinical studies. Other neglected areas of study include the timing of BMI measurements (eg, 1, 2, or 5 years before prostate cancer diagnosis) and the ability to assess ethnic disparities, including the association between obesity and lethal prostate cancer in African-American and other minority groups.

## OBESITY AND HEMATOLOGIC MALIGNANCIES PROGNOSIS

Epidemiologic evidence from prospective cohort studies indicates dose-response associations between excessive body weight and an increased risk of mortality from several subtypes of hematologic malignancies.<sup>16</sup>

The most recent meta-analysis of BMI and lymphoma by Larsson and Wolk<sup>5</sup> summarized available prospective data through May 2011, including five studies ( $n = 3,407$  cases) of BMI in relation to non-Hodgkin lymphoma (NHL) mortality. A dose-response association between increasing BMI and increased NHL mortality was observed. Per 5 kg/m<sup>2</sup> increase in BMI, the RR of NHL mortality was 1.14 (95% CI, 1.04 to 1.26). Conversely, some studies have reported that a higher BMI is associated with better survival outcomes in retrospective data.<sup>58,59</sup> In a more recent study, Hwang et al<sup>60</sup> reported data for Asian patients with diffuse large B-cell lymphoma and noted worse overall survival (hazard ratio [HR], 1.29; 95% CI, 1.08 to 7.95) and worse progression-free survival (HR, 2.59; 95% CI, 1.06 to 6.35) in the obese. Similarly, Leo et al<sup>61</sup> observed an association between obesity and worse NHL-specific survival in patients of diverse ethnicities (HR, 1.77; 95% CI, 1.30 to 2.41). One prospective study in Taiwan included a measure of central obesity (defined as waist circumference  $\geq 90$  cm in men,  $\geq 80$  cm in women) and reported worse NHL-specific survival in centrally obese patients (HR, 2.16; 95% CI, 1.41 to 3.31) after adjusting for BMI.<sup>62</sup> Across these studies, a worse prognostic outcome in underweight patients was reported, consistent with the impact of the disease process on weight.<sup>60,63,64</sup>

Multiple myeloma is relatively rare compared with lymphoma or leukemia; it has historically poor survival, with a 5-year survival rate of  $< 50\%$  in the United States.<sup>65</sup> Both overweight and obesity are associated with multiple myeloma mortality. Wallin and Larsson<sup>10</sup> combined available data restricted to prospective cohort studies published up to 2011. Data from five prospective cohorts ( $n = 1,845$  cases) suggested worse overall survival in both overweight (RR, 1.15; 95% CI, 1.04 to 1.27) and obese (RR, 1.54; 95% CI, 1.35 to 1.76) patients with multiple myeloma. These associations, assessed as a dose-response per 5 kg/m<sup>2</sup> increase in BMI, gave a 21% increased risk of multiple myeloma mortality per 5 kg/m<sup>2</sup> increase in BMI. The most updated pooled analysis of individual participant data from 20 prospective studies<sup>11</sup> extended the

## OBESITY AND RENAL CANCER PROGNOSIS

analysis on multiple myeloma mortality, including the association of early-adulthood BMI and weight distribution. BMI at cohort entry (RR, 1.09 [95% CI, 1.03 to 1.16] per 5 kg/m<sup>2</sup>) and higher early-adulthood BMI (RR, 1.22 [95% CI, 1.09 to 1.35] per 5 kg/m<sup>2</sup>) were both associated with increased multiple myeloma mortality. In this combined analysis, waist circumference (HR, 1.06 [95% CI, 1.02 to 1.10] per 5 cm) was also associated with mortality, suggesting the deleterious impact of central obesity. The association between early-adulthood BMI and mortality was stronger in women (HR, 1.95 [95% CI, 1.33 to 2.86] for BMI > 25 kg/m<sup>2</sup> compared with BMI of 18.5 to 25 kg/m<sup>2</sup>).

Leukemia accounts for 2.5% of all cancer cases globally.<sup>66</sup> Among its four major types, acute lymphoblastic leukemia (ALL) is the most common tumor in children, whereas other subtypes (chronic lymphocytic leukemia, acute myeloid leukemia, and chronic myeloid leukemia) occur mostly later in adult life. In adult populations, obesity is associated with a higher risk of leukemia mortality (RR, 1.29; 95% CI, 1.11 to 1.49) on the basis of data from six prospective cohort studies (n = 2,358 deaths) up to 2011.<sup>6</sup> The association between elevated BMI and leukemia mortality was linear in both men (1.9% per kg/m<sup>2</sup>, *P* = .10) and women (1.2% per kg/m<sup>2</sup>, *P* = .01). Several studies were conducted with retrospective design after the meta-analysis. Wenzell et al<sup>67</sup> reported improved overall survival in obese patients with leukemia, but others reported null findings.<sup>68,69</sup> A recent analysis that pooled data from four Cancer and Leukemia Group B (Alliance) clinical trials of patients with acute myeloid leukemia, including acute promyelocytic leukemia (APL), linked obesity to worse overall survival (HR, 1.72; 95% CI, 1.15 to 2.58) and disease-free survival (HR, 1.53; 95% CI, 1.03 to 2.27) in APL, but not in non-APL, suggesting a possible distinct biologic relation between obesity in APL.<sup>7</sup>

Studies of childhood leukemia have been focused mostly on ALL, the most common tumor in children. Data on other subtypes of childhood leukemia are scarce. Amankwah et al<sup>9</sup> conducted a meta-analysis synthesizing data from 11 studies up to 2015 to access the association between obesity at diagnosis and pediatric (< 21 years old) acute leukemia survival and relapse. They reported worse overall survival (HR, 1.30; 95% CI, 1.16 to 1.46) and worse event-free survival (HR, 1.46; 95% CI, 1.29 to 1.64) in patients who were obese at diagnosis. When the analysis was restricted to ALL only, both these associations persisted and were stronger (HR, 2.25 [95% CI, 1.33 to 3.82] for overall survival; and HR, 1.49 [95% CI, 1.30 to 1.71] for event-free survival).

Current evidence suggests a modest but consistent direct effect of BMI on poor prognostic outcomes in lymphoma, multiple myeloma, and leukemia. Risk increases in a dose-response manner in several subtypes of hematologic malignancies. It is also important to note that the adverse impact of BMI on multiple myeloma prognosis may reflect etiologic risk that starts growing early, during young adulthood. Separating the etiologic effect, disease severity at diagnosis, and outcomes requires more evidence from randomized interventions. To date, the small number of prospective studies and the lack of adjustment for possible confounding factors such as treatment, smoking, and comorbidities may bias the estimates reported for mortality as a result of hematologic malignancies.

Renal cancer (predominantly renal cell carcinoma [RCC]) accounts for approximately 2% of new cancer diagnoses. The IARC review in 2002 included evidence from four cohort studies and concluded a consistently positive association,<sup>1</sup> as did the World Cancer Research Fund review in 2007<sup>70</sup> and the subsequent update in 2015. The IARC 2016 update continues to classify RCC as caused by obesity.<sup>2</sup> Obesity is, however, associated with improved prognosis in patients with RCC, supported by a 2013 systematic review. This meta-analysis included data from 20 studies and reported improved overall (HR, 0.57; 95% CI, 0.43 to 0.76), cancer-specific (HR, 0.59; 95% CI, 0.48 to 0.74), and recurrence-free (HR, 0.49; 95% CI, 0.30 to 0.81) survival in patients with higher BMI than in those with lower BMI.<sup>13</sup> Notably, all these studies used a retrospective design. Data from prospective cohorts of patients are generally sparse. A 2016 study analyzed data from a prospective randomized trial of 845 localized high-risk patients with RCC and reported that a higher BMI was associated with improved survival.<sup>12</sup> The UK Million Women Study cohort found worse kidney-specific survival (RR, 1.65; 95% CI, 1.28 to 2.13) for every 10 kg/m<sup>2</sup> increment in cohort entry BMI, and the risk with increasing BMI was stronger in never-smokers.<sup>71</sup> Recently, a Korean patient cohort of 2,769 cases investigated the association between BMI and prognosis of nonmetastatic RCC by histologic subtype.<sup>15</sup> This study reported that a higher BMI ( $\geq 23$  kg/m<sup>2</sup>) was associated with improved cancer-specific survival (HR, 0.87) in patients with the clear cell variant, but worse cancer-specific (HR, 1.32) and recurrence-free (HR, 1.32) survival in patients with the chromophobe variant. No association was seen in patients with the papillary variant.<sup>15</sup> The association between obesity and RCC prognosis thus may differ by histologic subtype, although further study is needed to confirm these associations.

Current evidence on RCC prognosis is based mostly on data generated by retrospective studies, with few data available from prospective cohorts. These data support an obesity paradox in patients with RCC with the clear cell variant. That is, although obesity is an established risk factor for RCC, being obese at diagnosis seems to be associated with a favorable outcome compared with the outcomes of patients with normal weight at diagnosis. Some studies have reported that obese patients are more likely to be diagnosed with favorable clinical characteristics, including lower-stage disease, lower Fuhrman grade, and smaller tumors, compared with normal-weight patients.<sup>13</sup> Further support from gene expression analyses demonstrates that obese patients are less likely to have a tumor with fatty acid synthase (FASN), which encodes the proteins necessary for tumor growth.<sup>72</sup>

## OBESITY AND CANCER PROGNOSIS MECHANISMS

The mechanisms of obesity in relation to cancer development and prognosis are poorly understood on the basis of current studies. Furthermore, determining how much of an outcome is attributable to obesity, treatment, or the underlying disease subtypes/biology is difficult. It has been proposed that chemotherapy dosing that is based on body surface area in obese patients may be biased and may



result in poor survival. Hourdequin et al<sup>73</sup> conducted a meta-analysis and reported similar or lower levels of toxicity in obese patients compared with normal-weight patients when they received body surface area–based chemotherapy dosing, and no differences in survival outcomes were seen. Obesity reflects increased adipose tissue, which secretes a range of adipokines that may play a role in cancer development. These adipokines include inflammatory cytokines such as tumor necrosis factor- $\alpha$  and interleukin-6, which likely cause a chronic inflammatory micro-environment that also promotes cancer progression.<sup>74</sup> Such inflammation may lead to the activation of transcription factor nuclear factor- $\kappa$ B, which is linked to several cancers, including lymphoma.<sup>75</sup> Adipose tissue secretes leptin and adiponectin, which are both linked to prostate cancer and RCC.<sup>75</sup> Other pathways associated with obesity, such as hyperinsulinemia, insulin-like growth factor signaling, and lipid levels, have been explored in relation to cancer progression. For example, downgraded expression of FASN was found in obese patients with RCC and was associated with an improved survival.<sup>72</sup> In patients with prostate cancer, overweight men carrying a variant FASN allele had a poorer prognosis compared with men of normal weight.<sup>76</sup> These possible pathways require further research to address the potential response to weight loss after cancer diagnosis and possible improvement in survival outcomes. The cancer sites in this review have less evidence for a benefit of weight loss after diagnosis than is currently accumulating for breast and colorectal cancer.

### CLINICAL RECOMMENDATIONS

At the current time, the rapidly rising prevalence of obesity in all age groups may contribute to a large future cancer burden. The American Society of Clinical Oncology established a multipronged initiative and committed to reducing the impact of obesity on cancer.<sup>77</sup> The delivery of effective and efficient counseling on weight-management strategies, however, might be challenged by the poor understanding of the relationship between obesity and cancer progression. Cancer survivors are encouraged to achieve and maintain a healthy weight. Guidelines from the American Cancer Society<sup>78</sup> and the American College of Sports Medicine<sup>79</sup> suggest that cancer survivors should follow the physical activity guidelines for Americans, with specific exercise programming adaptations on the basis of disease- and treatment-related adverse effects (ie,  $\geq 150$  minutes per week of moderate-intensity physical

activity). In addition, cancer survivors are encouraged to achieve a high intake of vegetables, fruits, and whole grains and to avoid high-calorie foods and beverages. It is still unclear what impact specific lifestyle components (energy intake/diet or energy expenditure/physical activity) that contribute to obesity have in relation to cancer progression. With limited prospective longitudinal data on weight change in relation to prognostic outcomes, and in the absence of randomized controlled trials, cancer survivors should probably not be advised to voluntarily lose weight to reduce cancer recurrence or mortality. Yet maintaining a healthy lifestyle through physical activity, healthy diet, and avoiding weight gain may lead to general health benefits beyond cancer-specific outcomes.

### FUTURE DIRECTIONS

Future research to advance the understanding of weight management in the cancer care continuum can help document the time course of weight loss and improved outcomes after diagnosis and identify pathways that may be addressed more directly through drug therapies. Given the association between obesity and cancer incidence and mortality, shifting the perspective of clinical oncology practice to include weight management as a component of primary and secondary prevention of cancer is increasingly important.

### AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at [www.jco.org](http://www.jco.org).

### AUTHOR CONTRIBUTIONS

**Conception and design:** Lin Yang, Graham A. Colditz

**Administrative support:** Graham A. Colditz

**Collection and assembly of data:** All authors

**Data analysis and interpretation:** Lin Yang, Graham A. Colditz

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

### REFERENCES

1. International Agency for Research on Cancer: Weight Control and Physical Activity. Lyon, France, International Agency for Research on Cancer, 2002
2. Lauby-Secretan B, Scoccianti C, Loomis D, et al: Body fatness and cancer—viewpoint of the IARC Working Group. *N Engl J Med* 375:794-798, 2016
3. Calle EE, Rodriguez C, Walker-Thurmond K, et al: Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med* 348:1625-1638, 2003
4. Cao Y, Ma J: Body mass index, prostate cancer-specific mortality, and biochemical recurrence: A systematic review and meta-analysis. *Cancer Prev Res (Phila)* 4:486-501, 2011
5. Larsson SC, Wolk A: Body mass index and risk of non-Hodgkin's and Hodgkin's lymphoma: A meta-analysis of prospective studies. *Eur J Cancer* 47:2422-2430, 2011
6. Castillo JJ, Reagan JL, Ingham RR, et al: Obesity but not overweight increases the incidence and mortality of leukemia in adults: A meta-analysis of prospective cohort studies. *Leuk Res* 36:868-875, 2012
7. Castillo JJ, Mulkey F, Geyer S, et al: Relationship between obesity and clinical outcome in adults with acute myeloid leukemia: A pooled analysis from four CALGB (alliance) clinical trials. *Am J Hematol* 91:199-204, 2016
8. Breccia M, Mazzarella L, Bagnardi V, et al: Increased BMI correlates with higher risk of disease relapse and differentiation syndrome in patients with acute promyelocytic leukemia treated with the AIDA protocols. *Blood* 119:49-54, 2012
9. Amankwah EK, Saenz AM, Hale GA, et al: Association between body mass index at diagnosis and pediatric leukemia mortality and relapse: A systematic review and meta-analysis. *Leuk Lymphoma* 57:1140-1148, 2016
10. Wallin A, Larsson SC: Body mass index and risk of multiple myeloma: A meta-analysis of prospective studies. *Eur J Cancer* 47:1606-1615, 2011
11. Teras LR, Kitahara CM, Birmann BM, et al: Body size and multiple myeloma mortality: A pooled

analysis of 20 prospective studies. *Br J Haematol* 166:667-676, 2014

12. Donin NM, Pantuck A, Klöpfer P, et al: Body mass index and survival in a prospective randomized trial of localized high-risk renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 25:1326-1332, 2016
13. Choi Y, Park B, Jeong BC, et al: Body mass index and survival in patients with renal cell carcinoma: A clinical-based cohort and meta-analysis. *Int J Cancer* 132:625-634, 2013
14. Reeves GK, Pirie K, Beral V, et al: Cancer incidence and mortality in relation to body mass index in the Million Women Study: Cohort study. *BMJ* 335: 1134, 2007
15. Lee WK, Hong SK, Lee S, et al: Prognostic value of body mass index according to histologic subtype in nonmetastatic renal cell carcinoma: A large cohort analysis. *Clin Genitourin Cancer* 13: 461-468, 2015
16. Renehan AG, Tyson M, Egger M, et al: Body-mass index and incidence of cancer: A systematic review and meta-analysis of prospective observational studies. *Lancet* 371:569-578, 2008
17. De Nunzio C, Albisinni S, Freedland SJ, et al: Abdominal obesity as risk factor for prostate cancer diagnosis and high grade disease: A prospective multicenter Italian cohort study. *Urol Oncol* 31: 997-1002, 2013
18. Tewari R, Rajender S, Natu SM, et al: Significance of obesity markers and adipocytokines in high grade and high stage prostate cancer in North Indian men - a cross-sectional study. *Cytokine* 63:130-134, 2013
19. Larsson SC, Wolk A: Overweight and obesity and incidence of leukemia: A meta-analysis of cohort studies. *Int J Cancer* 122:1418-1421, 2008
20. Adams KF, Leitzmann MF, Albanes D, et al: Body size and renal cell cancer incidence in a large US cohort study. *Am J Epidemiol* 168:268-277, 2008
21. Pischon T, Lahmann PH, Boeing H, et al: Body size and risk of renal cell carcinoma in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Int J Cancer* 118:728-738, 2006
22. Whitlock G, Lewington S, Sherliker P, et al: Body-mass index and cause-specific mortality in 900 000 adults: Collaborative analyses of 57 prospective studies. *Lancet* 373:1083-1096, 2009
23. Fowke JH, McLerran DF, Gupta PC, et al: Associations of body mass index, smoking, and alcohol consumption with prostate cancer mortality in the Asia Cohort Consortium. *Am J Epidemiol* 182: 381-389, 2015
24. Burton AJ, Tilling KM, Holly JM, et al: Metabolic imbalance and prostate cancer progression. *Int J Mol Epidemiol Genet* 1:248-271, 2010
25. Presti JC, Jr: Obesity and prostate cancer. *Curr Opin Urol* 15:13-16, 2005
26. Li H, Stampfer MJ, Mucci L, et al: A 25-year prospective study of plasma adiponectin and leptin concentrations and prostate cancer risk and survival. *Clin Chem* 56:34-43, 2010
27. Ma J, Li H, Giovannucci E, et al: Prediagnostic body-mass index, plasma C-peptide concentration, and prostate cancer-specific mortality in men with prostate cancer: A long-term survival analysis. *Lancet Oncol* 9:1039-1047, 2008
28. MacInnis RJ, English DR: Body size and composition and prostate cancer risk: Systematic review and meta-regression analysis. *Cancer Causes Control* 17:989-1003, 2006
29. Bañez LL, Hamilton RJ, Partin AW, et al: Obesity-related plasma hemodilution and PSA concentration among men with prostate cancer. *JAMA* 298:2275-2280, 2007
30. Loeb S, Yu X, Nadler RB, et al: Does body mass index affect preoperative prostate specific antigen velocity or pathological outcomes after radical prostatectomy? *J Urol* 177:102-106, discussion 106, 2007
31. Su LJ, Arab L, Steck SE, et al: Obesity and prostate cancer aggressiveness among African and Caucasian Americans in a population-based study. *Cancer Epidemiol Biomarkers Prev* 20:844-853, 2011
32. Allott EH, Masko EM, Freedland SJ: Obesity and prostate cancer: Weighing the evidence. *Eur Urol* 63:800-809, 2013
33. Strom SS, Kamat AM, Gruschus SK, et al: Influence of obesity on biochemical and clinical failure after external-beam radiotherapy for localized prostate cancer. *Cancer* 107:631-639, 2006
34. Davies BJ, Smaldone MC, Sadetsky N, et al: The impact of obesity on overall and cancer specific survival in men with prostate cancer. *J Urol* 182:112-117, 2009; discussion 117
35. Bassett WW, Cooperberg MR, Sadetsky N, et al: Impact of obesity on prostate cancer recurrence after radical prostatectomy: Data from CaPSURE. *Urology* 66:1060-1065, 2005
36. Freedland SJ, Aronson WJ, Kane CJ, et al: Impact of obesity on biochemical control after radical prostatectomy for clinically localized prostate cancer: A report by the Shared Equal Access Regional Cancer Hospital database study group. *J Clin Oncol* 22: 446-453, 2004
37. Giovannucci E, Liu Y, Platz EA, et al: Risk factors for prostate cancer incidence and progression in the health professionals follow-up study. *Int J Cancer* 121:1571-1578, 2007
38. Rodriguez C, Freedland SJ, Deka A, et al: Body mass index, weight change, and risk of prostate cancer in the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* 16:63-69, 2007
39. Wright ME, Chang SC, Schatzkin A, et al: Prospective study of adiposity and weight change in relation to prostate cancer incidence and mortality. *Cancer* 109:675-684, 2007
40. Chalfin HJ, Lee SB, Jeong BC, et al: Obesity and long-term survival after radical prostatectomy. *J Urol* 192:1100-1104, 2014
41. Asmar R, Beebe-Dimmer JL, Korgavkar K, et al: Hypertension, obesity and prostate cancer biochemical recurrence after radical prostatectomy. *Prostate Cancer Prostatic Dis* 16:62-66, 2013
42. Ly D, Reddy CA, Klein EA, et al: Association of body mass index with prostate cancer biochemical failure. *J Urol* 183:2193-2199, 2010
43. American Cancer Society. *Cancer Facts & Figures 2008*. Atlanta, GA, American Cancer Society, 2008
44. American Cancer Society. *Cancer Facts & Figures for African Americans 2011-2012*. Atlanta, GA, American Cancer Society, 2011
45. Motamedinia P, Korets R, Spencer BA, et al: Body mass index trends and role of obesity in predicting outcome after radical prostatectomy. *Urology* 72:1106-1110, 2008
46. Keto CJ, Aronson WJ, Terris MK, et al: Obesity is associated with castration-resistant disease and metastasis in men treated with androgen deprivation therapy after radical prostatectomy: Results from the SEARCH database. *BJU Int* 110:492-498, 2012
47. Joshi CE, Mondul AM, Menke A, et al: Weight gain is associated with an increased risk of prostate cancer recurrence after prostatectomy in the PSA era. *Cancer Prev Res (Phila)* 4:544-551, 2011
48. Flegal KM, Carroll MD, Ogden CL, et al: Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 303:235-241, 2010
49. Caire AA, Sun L, Polascik TJ, et al: Obese African-Americans with prostate cancer (T1c and a prostate-specific antigen, PSA, level of <10 ng/mL) have higher-risk pathological features and a greater risk of PSA recurrence than non-African-Americans. *BJU Int* 106:1157-1160, 2010
50. Johnsen NF, Tjønneland A, Thomsen BL, et al: Physical activity and risk of prostate cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort. *Int J Cancer* 125:902-908, 2009
51. Giovannucci E, Leitzmann M, Spiegelman D, et al: A prospective study of physical activity and prostate cancer in male health professionals. *Cancer Res* 58:5117-5122, 1998
52. Giovannucci EL, Liu Y, Leitzmann MF, et al: A prospective study of physical activity and incident and fatal prostate cancer. *Arch Intern Med* 165: 1005-1010, 2005
53. Nilsen TI, Romundstad PR, Vatten LJ: Recreational physical activity and risk of prostate cancer: A prospective population-based study in Norway (the HUNT study). *Int J Cancer* 119:2943-2947, 2006
54. Patel AV, Rodriguez C, Jacobs EJ, et al: Recreational physical activity and risk of prostate cancer in a large cohort of U.S. men. *Cancer Epidemiol Biomarkers Prev* 14:275-279, 2005
55. Orsini N, Bellocco R, Bottai M, et al: A prospective study of lifetime physical activity and prostate cancer incidence and mortality. *Br J Cancer* 101:1932-1938, 2009
56. Wolin KY, Stoll C: Physical activity and urologic cancers. *Urol Oncol* 30:729-734, 2012
57. Liu Y, Hu F, Li D, et al: Does physical activity reduce the risk of prostate cancer? A systematic review and meta-analysis. *Eur Urol* 60:1029-1044, 2011
58. Carson KR, Bartlett NL, McDonald JR, et al: Increased body mass index is associated with improved survival in United States veterans with diffuse large B-cell lymphoma. *J Clin Oncol* 30:3217-3222, 2012
59. Weiss L, Melchardt T, Habringer S, et al: Increased body mass index is associated with improved overall survival in diffuse large B-cell lymphoma. *Ann Oncol* 25:171-176, 2014
60. Hwang HS, Yoon DH, Suh C, et al: Body mass index as a prognostic factor in Asian patients treated with chemioimmunotherapy for diffuse large B cell lymphoma, not otherwise specified. *Ann Hematol* 94:1655-1665, 2015
61. Leo QJ, Ollberding NJ, Wilkens LR, et al: Obesity and non-Hodgkin lymphoma survival in an ethnically diverse population: The Multiethnic Cohort study. *Cancer Causes Control* 25:1449-1459, 2014
62. Chu DM, Wahlqvist ML, Lee MS, et al: Central obesity predicts non-Hodgkin's lymphoma mortality and overall obesity predicts leukemia mortality in adult Taiwanese. *J Am Coll Nutr* 30:310-319, 2011
63. Park S, Han B, Cho JW, et al: Effect of nutritional status on survival outcome of diffuse large B-cell lymphoma patients treated with rituximab-CHOP. *Nutr Cancer* 66:225-233, 2014
64. Navarro WH, Loberiza FR, Jr., Bajorunaite R, et al: Effect of body mass index on mortality of patients with lymphoma undergoing autologous hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 12:541-551, 2006

65. Siegel RL, Miller KD, Jemal A: Cancer statistics, 2016. *CA Cancer J Clin* 66:7-30, 2016
66. Rodriguez-Abreu D, Bordoni A, Zucca E: Epidemiology of hematological malignancies. *Ann Oncol* 18:i3-i8, 2007
67. Wenzell CM, Gallagher EM, Earl M, et al: Outcomes in obese and overweight acute myeloid leukemia patients receiving chemotherapy dosed according to actual body weight. *Am J Hematol* 88:906-909, 2013
68. Tavitian S, Denis A, Vergez F, et al: Impact of obesity in favorable-risk AML patients receiving intensive chemotherapy. *Am J Hematol* 91:193-198, 2016
69. Medeiros BC, Othus M, Estey EH, et al: Impact of body-mass index on the outcome of adult patients with acute myeloid leukemia. *Haematologica* 97:1401-1404, 2012
70. World Cancer Research Fund. Food, Nutrition, Physical Activity, and the Prevention of Cancer: A Global Perspective. Washington, DC, AICR, 2007
71. Beral V, Bull D, Green J, et al: Ovarian cancer and hormone replacement therapy in the Million Women Study. *Lancet* 369:1703-1710, 2007
72. Hakimi AA, Furberg H, Zabor EC, et al: An epidemiologic and genomic investigation into the obesity paradox in renal cell carcinoma. *J Natl Cancer Inst* 105:1862-1870, 2013
73. Hourdequin KC, Schpero WL, McKenna DR, et al: Toxic effect of chemotherapy dosing using actual body weight in obese versus normal-weight patients: A systematic review and meta-analysis. *Ann Oncol* 24:2952-2962, 2013
74. Grivnenikov SI, Greten FR, Karin M: Immunity, inflammation, and cancer. *Cell* 140:883-899, 2010
75. Khandekar MJ, Cohen P, Spiegelman BM: Molecular mechanisms of cancer development in obesity. *Nat Rev Cancer* 11:886-895, 2011
76. Nguyen PL, Ma J, Chavarro JE, et al: Fatty acid synthase polymorphisms, tumor expression, body mass index, prostate cancer risk, and survival. *J Clin Oncol* 28:3958-3964, 2010
77. Ligibel JA, Alfano CM, Courneya KS, et al: American Society of Clinical Oncology position statement on obesity and cancer. *J Clin Oncol* 32:3568-3574, 2014
78. Rock CL, Doyle C, Demark-Wahnefried W, et al: Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin* 62:243-274, 2012
79. Schmitz KH, Courneya KS, Matthews C, et al: American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 42:1409-1426, 2010

## Participate in ASCO's Practice Guidelines Implementation Network and Influence Cancer Care

ASCO members are invited to serve in the society's Practice Guidelines Implementation Network (PGIN), a network of oncology professionals who raise awareness of ASCO's evidence-based recommendations on cancer care.

Participation in PGIN provides an opportunity for members to positively influence the way that clinical oncology is delivered now and in the future. PGIN members have the opportunity to:

- Participate in Guideline Panels and Advisory Groups
- Aid in developing and reviewing Guidelines and Guideline Clinical Tools and Resources
- Serve as an "ambassador" to state societies
- Better implement Guidelines

To learn how you can participate, visit [asco.org/guidelines](http://asco.org/guidelines), or contact [PGIN@asco.org](mailto:PGIN@asco.org).



## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

### Obesity and Other Cancers

*The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [jco.ascopubs.org/site/ifc](http://jco.ascopubs.org/site/ifc).*

**Lin Yang**

No relationship to disclose

**Graham A. Colditz**

No relationship to disclose

**Bettina F. Drake**

No relationship to disclose