


# ***Cepaea nemoralis* snails are less likely to trap nematodes within their shells in urban areas - supplementary material**

Maxime Dahirel, Hannah Reyn , Katrien De Wolf, Dries Bonte


## **S1 - Variation in human population density and in built-up surfaces between Degrees of Urbanisation**

We show here (**Figs S1-1 and S1-2**) the relationships in our sites between the Degree of Urbanisation classes (Eurostat (European Commission) 2021) and the corresponding values of human population density and built-up cover, all from 1000 m resolution rasters.

- For both population density and built-up, the differences between the classes scale non-linearly, with the difference between the average Urban Centre site and the average “intermediate” site being much larger than the difference between the average intermediate and rural sites;
- Urban Centre sites contain a much broader range of variation in either variable, compared to intermediate and rural sites;
- Interestingly, the differences in built-up surfaces between the three Degree of Urbanisation classes map almost exactly to built-up thresholds designed and used independently in a series of urban ecology studies from Belgian sites (e.g. Merckx, Kaiser, and Van Dyck 2018; Piano et al. 2020) (**Fig. S1-2**). Degree of Urbanisation classes can therefore (*for our data*) be interpreted as encoding both variation in density and in built-up surfaces.



**Figure S1-1.** Relationship between Degree of Urbanisation classes and human population densities, for all 36 visited sites (including sites with no snails found).



**Figure S1-2.** Relationship between Degree of Urbanisation classes and built-up surfaces, for

all 36 visited sites (including sites with no snails found). Shaded areas and corresponding labels mark the limits of the three urban classes used in studies from the Belgian SPEEDY project, including for instance Merckx, Kaiser, and Van Dyck (2018) or Piano et al. (2020).


## S2 - Intensity of nematodes trapped in shells

Shells that do contain nematodes may also vary in the numbers of nematodes they harbour (in the context of active infections, this is termed “intensity,” Bush et al. 1997). However, given the relatively low number of shells with nematodes (104) especially relative to the number of sites, and given the expected high dispersion of these numbers (e.g. Rae 2017), we do not consider our intensity dataset to be sufficiently large to be given consideration in the main text. The partial analysis of these data we present below is therefore intended for indicative purposes only.

Using the subset of shells where nematodes were detected, we ran a negative binomial GLMM on the number of nematodes per shell - 1, with the same fixed and random effect structure as the best model for nematode prevalence (see **main text**). We find no clear indication that these numbers differed between urbanisation levels (**Table S2-1, Fig. S2-1**), or in relation to shell phenotype (**Table S2-1**).

**Table S2-1.** Analysis of Deviance for the effect of urbanisation and shell phenotype on the number of nematodes per shell (on the subset of shells containing at least one nematode).

variable	X <sup>2</sup>	Df	<i>p</i>
Degree of Urbanisation	0.15	2	0.93
City	0.73	2	0.69
Shell background colour	0.77	2	0.68
Number of bands	0.60	1	0.44
Band fusion	0.59	1	0.44



**Figure S2-1.** Relationship between the Degree of Urbanisation in 1000m grid cells and the number of nematodes *in shells that contain at least one nematode*. Grey dots are observed values; white dots (and error bars) are estimated marginal means from the best model (and their 95% confidence intervals), with the effects of city identity and phenotypic traits averaged out. Note that the  $y$  axis is on the log scale.

## References

- Bush, Albert O., Kevin D. Lafferty, Jeffrey M. Lotz, and Allen W. Shostak. 1997. "Parasitology Meets Ecology on Its Own Terms: Margolis Et Al. Revisited." *The Journal of Parasitology* 83 (4): 575–83. <https://doi.org/10.2307/3284227>.
- Eurostat (European Commission). 2021. *Applying the Degree of Urbanisation: A Methodological Manual to Define Cities, Towns and Rural Areas for International Comparisons : 2021 Edition*. LU: Publications Office of the European Union.
- Merckx, Thomas, Aurélien Kaiser, and Hans Van Dyck. 2018. "Increased Body Size Along Urbanization Gradients at Both Community and Intraspecific Level in Macro-Moths." *Global Change Biology* 24 (8): 3837–48. <https://doi.org/10.1111/gcb.14151>.
- Piano, Elena, Caroline Souffreau, Thomas Merckx, Lisa F. Baardsen, Thierry Backeljau, Dries Bonte, Kristien I. Brans, et al. 2020. "Urbanization Drives Cross-Taxon Declines in Abundance and Diversity at Multiple Spatial Scales." *Global Change Biology* 26 (3): 1196–1211. <https://doi.org/10.1111/gcb.14934>.

Rae, R. 2017. “The Gastropod Shell Has Been Co-Opted to Kill Parasitic Nematodes.” *Scientific Reports* 7 (1): 4745. <https://doi.org/10.1038/s41598-017-04695-5>.