

# Central Limit Theorem and Introduction to Inference

Sumanta Basu

# Reading

- Reading: Textbook Sections 3.1, 4.1-4.2
- Recommended Exercise: 4.1, 4.5, 4.7, 4.9, 4.13

Prelim 1 will **NOT** include readings from Chapter 4 of textbook.

# Central Limit Theorem (CLT)

# Populations and samples

**Goal:** Draw scientific inference about a specific aspect of a target population from a *representative* sample

# Example: 2012 Cherry Blossom 10-Mile Run

**Target population:** all runners who finished the run in 2012.

Research question: What is the average age of this population?

There were 16,924 people who finished... would be hard to ask them all!

Draw **simple random sample** (SRS).

# Drawing sample

## Options?

- before race starts, ask 50 people?
- send mass email with survey?
- give away energy bars to anyone who will answer your survey?
- be part of race and approach people you encounter along the way?
- stand by finish line for 30 minutes and ask whoever completes race during that time?

# Drawing sample

None of those are good. Remember lectures on sampling!

None give a **SRS**.

# Estimating mean age from sample

Let  $\mu$  be the target population's mean age:

$$\mu = \frac{\text{age}_1 + \text{age}_2 + \cdots + \text{age}_{16,924}}{16,924}$$

This is unknown!

But let's select a SRS of size 10:

$$\bar{x} = \frac{31 + 30 + 48 + 41 + 30 + 33 + 25 + 28 + 33 + 39}{10} = 33.8$$

Do we “trust” our answer?

Is  $\bar{x} = 33.8$  the same thing as  $\mu$ ?



# The R in SRS is Random

When we draw an SRS, we are doing something **random**.

$\bar{x} = 33.8$  is a **realization** of this random process!

**Key idea:** Imagine if we were to repeat this sampling process again and again.

# SRS

Recall definition of **SRS**:

Choose  $n$  units from target population at random so that each possible subset of size  $n$  is equally likely to be chosen.

Image we repeated experiment:

- draw 10 people
- calculate  $\bar{x}$

We might get 34.0 and 29.3... different realizations of a **random variable**.

# Sample mean from SRS

33.8, 34.0, 29.3 are *realizations* of a **random variable**, call it  $\bar{X}_{10}$ .

$\bar{X}_{10}$  in words - *“draw a SRS of 10 people and average their ages”*

Why is it a random variable?

- a **random variable** is a numerical summary of a random outcome.
- “draw a SRS of 10 people” is a random experiment
- “average their ages” - sample mean of the ages of the random people drawn is a numerical summary

# Sample mean as a random variable

We can write:

$$\bar{X}_{10} = \frac{X_1 + \cdots + X_{10}}{10}$$

$X_i$  = age of the  $i$  th person drawn in SRS (also a random variable!)

Suppose my SRS gives

31 30 48 41 30 33 25 28 33 39

What is  $X_5$ ?

# Sample mean as a random variable


What is  $\bar{X}_{10}$ 's distribution?

If we had entire population of ages, we could simulate...

...actually in this case we do have the entire population's ages.


# Population of ages

**Histogram of ages**




# Drawing from this population

PMF of  $X_1$ , a draw from population



# Drawing from this population


Realization of an SRS of size 10...






# Drawing from this population

Realization of an SRS of size 10...




# Drawing from this population

Realization of an SRS of size 10...




# Drawing from this population

Realization of an SRS of size 10...




# Drawing from this population

Realization of an SRS of size 10...




# Drawing from this population

Realization of an SRS of size 10...




# Drawing from this population

Realization of an SRS of size 10...




# Sample mean's distribution

By Monte Carlo simulation, we can get  $\bar{X}_{10}$ 's distribution:




# Comments



- when you compute a statistic from a sample using SRS, that is *a realization of a random variable*
- Red shows the distribution of the statistic  $\bar{X}_{10}$
- this distribution is called the **sampling distribution** of the statistic
- could only simulate sampling distribution since we had entire population's data (not realistic)




# Observations




- **sampling distribution** of  $\bar{X}_{10}$  seems to be centered at population mean,  $\mu$  (shown in black)
- variance of  $\bar{X}_{10}$  is smaller than variance of population
- shape looks normal!!
- that's surprising since population distribution was not!


# Effect of sample size?




# Effect of sample size?




# Effect of sample size?




# Effect of sample size?




# Effect of sample size?



# Effect of sample size?




# Effect of sample size?





# Effect of sample size?



# What we've observed

Let  $\bar{X}_n$  be the sample mean of a SRS of size  $n$  from a population with mean  $\mu$  and standard deviation  $\sigma$ . Then,  $\bar{X}_n$  is a random variable with

$$E(\bar{X}_n) = \mu$$

and

$$\text{SD}(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$$

## And what's more...

When  $n$  is “large enough”,  $\bar{X}_n$  is approximately normal!!

$\bar{X}_n$  is approximately  $N(\mu, \sigma/\sqrt{n})$

# Central Limit Theorem

If  $X_1, \dots, X_n$  are independent draws from a distribution with mean  $\mu$  and standard deviation  $\sigma$ , then for large  $n$ , the sample mean  $\bar{X}_n$  is approximately normal with mean  $\mu$  and standard deviation  $\sigma/\sqrt{n}$ :

$$\bar{X}_n \approx N\left(\mu, \frac{\sigma}{\sqrt{n}}\right)$$

- remarkable since individual  $X_i$  's don't have to look at all like a normal distribution
- how large should  $n$  be? Depends, but if distribution of  $X_i$  's is not strongly skewed, say  $n \geq 30$

# Example

Suppose  $X_1, \dots, X_n$  are independent coin flips, i.e.,  $X_i \sim \text{Bernoulli}(p)$ .

The **sample proportion**, sometimes written  $\hat{p}_n$ , is just  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ .

Recall  $E(X_i) = p$  and  $\text{Var}(X_i) = p(1 - p)$ .

CLT tells us

$$\hat{p}_n \approx N(p, \sqrt{p(1 - p)/n})$$

# Galton Bean Machine

[A dramatic video of a beautiful phenomenon](#)



Prelim 1 will cover topics only  
upto this point, not the next  
slides

# Inference

**Sampling distribution:** A probabilistic description of how the observed values of a numerical summary statistic (e.g., sample mean) behave under repeated SRS.

This concept underlies all basic statistical inference procedures – its importance cannot be overstated!

*In practice:* we only collect one sample.

**Question:** how can we combine the information from a single SRS about a population parameter with our knowledge of sampling distributions in order to perform statistical inference?



# Two primary goals


1. A **confidence interval** - a range of plausible values for a (population) parameter, based on the data obtained from our observed sample.
2. A **hypothesis (or significance) test** - an assessment of whether the observed value of a statistic computed using the sample data is consistent with or divergent from some hypothesized value of the (population) parameter.

*Note:* these get at “*what is  $\mu$ ?*” better than just reporting a single **point estimate** (e.g.,  $\bar{x} = 33.8$ )

# Confidence Interval (CI)


# Point estimate

**Realization of an SRS of size 10...**



# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate

## Realization of an SRS of size 10...




- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$



# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate


Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Interval estimate

Realization of an SRS of size 10...



- interval is calculated based on sample (centered at  $\bar{X}_n$ )
- thus it is random... remember we're looking at just one realization
- interval here is  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$

# Question

What is the probability that  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$  includes  $\mu$ ?

## [1] 0.9583


(used Monte Carlo to answer this.)

What happens if interval narrower?

$P(\bar{X}_n - 2.0 \leq \mu \leq \bar{X}_n + 2.0) = ?$


## [1] 0.4608

# Monte Carlo simulation in a picture



- 100 realizations of the random interval  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$
- 97 out of 100 "cover"  $\mu$

# Monte Carlo simulation in a picture



- 100 realizations of the random interval  $[\bar{X}_n - 2.0, \bar{X}_n + 2.0]$
- 48 out of 100 "cover"  $\mu$

# Confidence Interval

The random interval  $[\bar{X}_n - 3.2, \bar{X}_n + 3.2]$  is called a 96% **confidence interval**. Here, 96% is said to be its **confidence level**.

$$P([\bar{X}_n - 3.2, \bar{X}_n + 3.2] \text{ includes } \mu) = 96\%$$