




Planar Graphs and Partial k -Trees

Philip T. Henderson

Supervisor: Therese Biedl


pthender@cs.uwaterloo.ca



Overview


- Terminology
- Nested Satisfiability
- Matryoshka Graphs
- Decompositions of Bounded Treewidth

Planar Graphs




- Vertices as points
- Edges as continuous curves
- Can be drawn in the plane without edge crossing

Partial k -Trees




- Subclass of all graphs: every graph is a partial k -tree for some k
- Many NP-hard problems (e.g. SAT, independent set, etc.) are solvable in linear time for constant k

Tree Decomposition



A **tree decomposition** of graph G is a tree with labels on nodes such that:


Tree Decomposition



A **tree decomposition** of graph G is a tree with labels on nodes such that:

- every vertex appears in a label


Tree Decomposition



A **tree decomposition** of graph G is a tree with labels on nodes such that:

- every vertex appears in a label
- for any edge (u, v) , u and v are in a label


Tree Decomposition




A **tree decomposition** of graph G is a tree with labels on nodes such that:

- every vertex appears in a label
- for any edge (u, v) , u and v are in a label
- for any vertex v , the labels containing v form a tree

Partial k -Trees




- Width = maximum label size – 1
- Partial k -tree: graph has tree decomposition of width $\leq k$
- Equivalently: graph has treewidth $\leq k$



Part I: Nested SAT

- SAT = satisfiability: classic NP-hard problem
- Knuth '90: Nested SAT can be solved in linear-time
- Kratochvíl and Křivánek '93: extends to MaxSAT and co-nested SAT
- We show: Nested SAT graphs are partial 3-trees
- This unifies and generalizes the results above

Nested Satisfiability



SAT (in CNF):


- variables X_1, \dots, X_n
- clauses C_1, \dots, C_m

SAT graph:

- vertex for each X_i and C_j
- edge (X_i, C_j) iff $X_i \in C_j$ or $\overline{X_i} \in C_j$
- edges (X_i, X_{i+1})

Nested Satisfiability


Nested SAT: SAT that has LH-drawing with variables on the line and clauses in the half-space




LH-Drawing

Given: graph G with vertex partition $V_1 \cup V_2$, an **LH-drawing** is a planar drawing such that


- vertices in V_1 are on the **line**
- vertices in V_2 are in the **half-space** above
- all edges are y -monotonic curves



Clause Hierarchy and Visible Variables




Clause Hierarchy and Visible Variables



- Visible variables: can add clause-variable edge and still have an LH-drawing


Clause Hierarchy and Visible Variables



- Visible variables: can add clause-variable edge and still have an LH-drawing
- Clause hierarchy: LH-drawing imposes tree structure on clauses


Clause Paths

Define tree decomposition for single clause and its visible variables




Clause Paths

Define tree decomposition for single clause and its visible variables




Clause Paths

Define tree decomposition for single clause and its visible variables




Clause Paths

Define tree decomposition for single clause and its visible variables




Clause Paths

Define tree decomposition for single clause and its visible variables





Nested SAT Tree Decomposition

Combine clause paths using clause hierarchy tree




Nested SAT Tree Decomposition

Combine clause paths using clause hierarchy tree




Nested SAT Tree Decomposition

Combine clause paths using clause hierarchy tree




Nested SAT Tree Decomposition

Combine clause paths using clause hierarchy tree




Nested SAT Tree Decomposition

Combine clause paths using clause hierarchy tree



Nested SAT Tree Decomposition


Combine clause paths using clause hierarchy tree



⇒ Nested SAT graphs are partial 3-trees

Part II: Generalizing Nested SAT Graphs

Can add edges and remain a partial 3-tree:



- Edges to visible variables
- Edges in clause hierarchy


LH-drawing satisfies:

- All faces incident to the line

Can we push this farther?


Clause Trees

- For each clause, can add edges from visible variables to rightmost variable



Clause Trees


- Modification: each clause can have an outerplanar[†] graph on its visible variables



† Outerplanar = planar and all vertices are on the outer-face


Clause Trees

- Modification: each clause can have an outerplanar[†] graph on its visible variables



† Outerplanar = planar and all vertices are on the outer-face


Matryoshka Graphs



Maximal Matryoshka graph has planar drawing with:

- vertices on or above line
- every face incident to line
- edges above: y -monotone
- edges below: endpoints have common neighbour in half-space

Matryoshka graph: subgraph of maximal Matryoshka graph



Characterizing Matryoshka Graphs

From above: Matryoshka graphs are partial 3-trees


- Can we characterize them?
- How do they differ from planar partial 3-trees?

Theorem All maximal Matryoshka graphs have a Hamiltonian cycle
⇒ Not all planar partial 3-trees are Matryoshka graphs

Binary Planar 3-Trees

Proof by picture for **binary planar 3-trees**:

- Start with K_4
- Repeatedly subdivide at most two faces of a K_4 in graph




Every maximal Matryoshka graph is a binary planar 3-tree

Binary Planar 3-Trees

Proof by picture for **binary planar 3-trees**:

- Start with K_4
- Repeatedly subdivide at most two faces of a K_4 in graph




Every maximal Matryoshka graph is a binary planar 3-tree

Binary Planar 3-Trees

Proof by picture for **binary planar 3-trees**:

- Start with K_4
- Repeatedly subdivide at most two faces of a K_4 in graph




Every maximal Matryoshka graph is a binary planar 3-tree

Binary Planar 3-Trees

Proof by picture for **binary planar 3-trees**:

- Start with K_4
- Repeatedly subdivide at most two faces of a K_4 in graph




Every maximal Matryoshka graph is a binary planar 3-tree

Binary Planar 3-Trees

Proof by picture for **binary planar 3-trees**:


- Start with K_4
- Repeatedly subdivide at most two faces of a K_4 in graph



Every maximal Matryoshka graph is a binary planar 3-tree

Superclasses and Subclasses

- Matryoshka graphs \subset binary planar 3-trees
- Matryoshka graphs \supset Halin and IO-graphs
- Matryoshka graphs $\not\supset$ planar partial 2-trees[†]




[†] Recent result, does not appear in thesis



Summary of Matryoshka Graphs


- Strict subset of binary planar 3-trees
- Generalizes Halin graphs, IO-graphs, Nested SAT graphs
- No efficient algorithm for recognition yet exists
- Not closed under minors, so minor theory does not help



Part III: Planar Graph Decompositions

- Chartrand-Geller-Hedetniemi conjecture '71:
Planar graphs can be decomposed into two outerplanar graphs
- Know they can be decomposed as:
 - two partial 2-trees (SP-graphs)
 - three partial 1-trees (forests)

We study: Can any planar graph be decomposed into a forest and a partial k -tree?




Motivation

Motivation I: Matryoshka graphs are partial $(1, 2)$ -trees (forest plus SP-graph)

Motivation II: Could lead to better algorithms


- Can efficiently solve many NP-hard problems on the partial k -tree, then modify solution using forest
- Gives a simple 2-approximation algorithm for weighted vertex cover and weighted independent set

Hamiltonian Planar Graphs



Theorem Sub-Hamiltonian planar graphs are partial $(O(\log n), 1)$ -trees.


Hamiltonian Planar Graphs



Theorem Sub-Hamiltonian planar graphs are partial $(O(\log n), 1)$ -trees.

- Inside: outerplanar


Hamiltonian Planar Graphs



Theorem Sub-Hamiltonian planar graphs are partial $(O(\log n), 1)$ -trees.

- Inside: outerplanar \Rightarrow partial 2-tree


Hamiltonian Planar Graphs



Theorem Sub-Hamiltonian planar graphs are partial $(O(\log n), 1)$ -trees.

- Inside: outerplanar \Rightarrow partial 2-tree
- Outside: 3 outerplanar sections


Hamiltonian Planar Graphs



Theorem Sub-Hamiltonian planar graphs are partial $(O(\log n), 1)$ -trees.


- Inside: outerplanar \Rightarrow partial 2-tree
- Outside: 3 outerplanar sections
- Each outerplanar section has dual binary tree

Dual Tree Contraction



Theorem A sub-Hamiltonian planar graph G is a partial $(O(\log n), 1)$ -tree.


Dual Tree Contraction



Theorem A sub-Hamiltonian planar graph G is a partial $(O(\log n), 1)$ -tree.

Proof idea: In dual tree, contract edge to taller child

Dual Tree Contraction




Theorem A sub-Hamiltonian planar graph G is a partial $(O(\log n), 1)$ -tree.

Proof idea: In dual tree, contract edge to taller child

- This deletes a forest in G
- Contracted dual tree has $O(\log n)$ height

Dual Tree Contraction



Theorem A sub-Hamiltonian planar graph G is a partial $(O(\log n), 1)$ -tree.


Proof idea: In dual tree, contract edge to taller child

- This deletes a forest in G
- Contracted dual tree has $O(\log n)$ height

\Rightarrow Rest of G is partial $O(\log n)$ -tree

Special Case: Bounded Degrees


Theorem Every max-deg-3 graph can be decomposed into a matching and a partial 2-tree.




- A **matching** is a set of edges, no two of which are adjacent
- Petersen's Theorem handles 2-edge-connected 3-regular graphs: a perfect matching exists
- Remaining graph has max-deg-2: partial 2-tree

Special Case: Bounded Degrees

Theorem Every max-deg-3 graph can be decomposed into a matching and a partial 2-tree.




- Bridges don't hurt: handle 2-edge-connected components separately
- For each component: can find a matching that reaches all deg-3 vertices
- Remaining graph has max-deg-2: partial 2-tree




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $O(1)$ -tree?




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $O(1)$ -tree?
- No!




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $O(\log n)$ -tree?




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $O(\log n)$ -tree?
- No!




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?



Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- No!




Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar

Does This Generalize?


- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar



Does This Generalize?


- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar

For any matching:



Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar




For any matching:

- Diamonds contract to an edge

Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar




For any matching:

- Diamonds contract to an edge

Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar




For any matching:

- Diamonds contract to an edge
- Bowties contract to a vertex

Does This Generalize?

- Can every max-deg- d graph be decomposed into a matching and a partial $o(\sqrt{n})$ -tree?
- **No!** Not even for $d = 4$, not even when planar



For any matching:

- Diamonds contract to an edge
- Bowties contract to a vertex

Obtain $k \times k$ grid \Rightarrow not a partial $(k - 1)$ -tree

Summary

Planar graphs as partial $(k, 1)$ -trees

- Sub-Hamiltonian graphs are partial $(O(\log n), 1)$ -trees
 - Does it hold for some constant $k \geq 3$?
- Cannot easily extend to all planar graphs
 - Does it hold for some $k \in o(\sqrt{n})$?



Any Questions?

Thanks to:

- Angèle Hamel for her help during Therese's sabbatical
- Rick Mabry for sending me correspondence with Paul Seymour on related material
- Fellow graduate students Reinhold Burger, Niel de Beaudrap, Magdalena Georgescu, Graeme Kemkes, Brendan Lucier, Zach Olesh, Alex Stewart, and Luke Tanur for various helpful discussions