

# Bootstrap-Assisted Inference for Generalized Grenander-type Estimators

Matias D. Cattaneo<sup>1</sup>, Michael Jansson<sup>2</sup>, and Kenichi Nagasawa<sup>3</sup>

April 2023

---

<sup>1</sup>Princeton University.

<sup>2</sup>UC-Berkeley.

<sup>3</sup>University of Warwick.

## Setup: Grenander-Type Estimators [Westling & Carone, 2020]

Monotone function estimators  $\widehat{\theta}_n(x)$  at interior point  $x$  exhibit:

$$n^{q/(1+2q)}(\widehat{\theta}_n(x) - \theta_0(x)) \rightsquigarrow \frac{1}{\partial\Phi_0(x)}\partial_{-}\text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_x^q)(0),$$

$$\mathcal{M}_x^q(v) = \frac{\partial^q\theta_0(x)\partial\Phi_0(x)}{(1+q)!}v^{1+q}, \quad q = \min\{j \in \mathbb{N} : \partial^j\theta_0(x) \neq 0\}.$$

as  $n \rightarrow \infty$ , where:

- ▶  $\theta_0(\cdot)$  is density, regression or hazard function, among other possibilities.
- ▶  $\mathcal{G}_x$  is zero-mean Gaussian process (nonstationary, rough cov kernel).
- ▶  $\mathcal{M}_x^q$  is non-random drift function (usually quadratic, but not always).
- ▶  $\Phi_0$  unknown non-decreasing and càdlàg function.
- ▶  $\text{GCM}_I(f)$  is greatest convex minorant of function  $f$  on interval  $I$ .

**Motivation:** Conducting inference can be challenging.

## Setup: Grenander-Type Estimators [Westling & Carone, 2020]

$$n^{q/(1+2q)}(\widehat{\theta}_n(x) - \theta_0(x)) \rightsquigarrow \frac{1}{\partial\Phi_0(x)} \partial_{-} \text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_x^q)(0),$$

$$\mathcal{M}_x^q(v) = \frac{\partial^q \theta_0(x) \partial \Phi_0(x)}{(1+q)!} v^{1+q}, \quad q = \min \{j \in \mathbb{N} : \partial^j \theta_0(x) \neq 0\}.$$

### Examples:

- ▶ Isotonic density [Grenander, 1956] and extensions (censoring, covariates).
- ▶ Isotonic regression [Brunk, 1970] and extensions (censoring, covariates).
- ▶ Monotone Hazard [Rao, 1970] and extensions (censoring, covariates).
- ▶ Current Status [Ayer et al. 1955] and extensions (censoring, covariates).

**Problem:** bootstrap inconsistent [Kosorok, 2008; Sen, Banerjee & Woodroffe, 2010].

- ▶ Restore bootstrap validity: modifying the distribution used when resampling (subsampling,  $m$ -out-of- $n$  bootstrap, smooth bootstrap).
- ▶ This paper: Modifying (“reshaping”) functional form estimator [CJN, 2020].

# Leading Example (Today): Isotonic Density Estimation

## Model:

- ▶  $X_1, \dots, X_n$  i.i.d. with support  $[0, 1]$ .
- ▶  $F(x) = \mathbb{P}[X_i \leq x]$  absolutely continuous.
- ▶  $\partial F(x) = f(x)$  monotone (e.g., non-decreasing).

**Estimand:**  $\theta_0(x) = f(x)$  for interior point  $x$ .

**Estimator:** for  $\mathcal{F}$  the class of non-decreasing densities supported on  $[0, 1]$ ,

$$\hat{\theta}_n(\cdot) = \arg \max_{f \in \mathcal{F}} \sum_{i=1}^n \log f(X_i)$$

$$\implies \hat{\theta}_n(x) = \partial_- \text{GCM}_{[0,1]}(\hat{\Gamma}_n)(x), \quad \hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$$

**Asy Dist:** under regularity,  $q = 1$  and  $\Phi_0(x) = x$ ,

$$\sqrt[3]{n}(\hat{\theta}_n(x) - \theta_0(x)) \rightsquigarrow \partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_{x,1})(0) \sim \sqrt[3]{4f(x)\partial f(x)} \underset{v \in \mathbb{R}}{\operatorname{argmin}} \{ \mathcal{G}(v) + v^2 \}$$

## Second Example: Isotonic Regression Estimation

**Model:**

- ▶  $(Y_1, X_1), \dots, (Y_n, X_n)$  i.i.d. with  $X_i$  on support  $I$ , and  $F(x) = \mathbb{P}[X_i \leq x]$ .
- ▶  $\mu(x) = \mathbb{E}[Y_i | X_i = x]$  differentiable and monotone (e.g., non-decreasing).

**Estimand:**  $\theta_0(x) = \mu(x)$  for interior point  $x$ .

**Estimator:** for  $\mathcal{F}$  the class of non-decreasing functions supported on  $I$ ,

$$\begin{aligned}\hat{\theta}_n(x) &= \operatorname{argmin}_{f \in \mathcal{F}} \sum_{i=1}^n (Y_{(i)} - f(X_{(i)}))^2 \\ \implies \hat{\theta}_n(x) &= \left( \partial_{-\text{GCM}}_{[0,1]}(\hat{\Gamma}_n \circ \hat{\Phi}_n^-) \right) \circ \hat{\Phi}_n^-(x), \\ \hat{\Gamma}_n(x) &= \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x) Y_i, \quad \hat{\Phi}_n^- = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x).\end{aligned}$$

**Asy Dist:** under regularity,  $q = 1$  and  $\Phi_0(x) = F(x)$ ,

$$\sqrt[3]{n}(\hat{\theta}_n(x) - \theta_0(x)) \rightsquigarrow \frac{1}{\partial \Phi_0(x)} \partial_{-\text{GCM}}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_{x,1})(0) \sim \sqrt[3]{\frac{4\sigma^2(x)\partial\mu(x)}{f(x)}} \operatorname{argmin}_{v \in \mathbb{R}} \{\mathcal{G}(v) + v^2\}$$

# Third Example: Covariate-Adjusted Isotonic Regression

## Model:

- ▶  $(Y_1, X_1, \mathbf{Z}'_1), \dots, (Y_n, X_n, \mathbf{Z}'_n)$  i.i.d. with  $X_i$  on support  $I$ , and  $F(x) = \mathbb{P}[X_i \leq x]$ .
- ▶  $\mu(x) = \mathbb{E}[\mathbb{E}[Y_i | X_i = x, \mathbf{Z}_i]]$  differentiable and monotone (e.g., non-decreasing).

**Estimand:**  $\theta_0(\mathbf{x}) = \mathbb{E}[\mathbb{E}[Y_i | X_i = \mathbf{x}, \mathbf{Z}_i]]$ .

## Estimator:

$$\hat{\theta}_n(\mathbf{x}) = \left( \partial_- \text{GCM}_{[0,1]}(\hat{\Gamma}_n \circ \hat{\Phi}_n^-) \right) \circ \hat{\Phi}_n^-(\mathbf{x})$$

$$\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x) \left[ \frac{Y_i - \hat{\mu}(X_i, \mathbf{Z}_i)}{\hat{f}(\mathbf{Z}_i | X_i)} + \frac{1}{n} \sum_{j=1}^n \hat{\mu}(X_i, \mathbf{Z}_j) \right], \quad \hat{\Phi}_n^-(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x).$$

**Asy Dist:** under regularity,  $q = 1$  and  $\Phi_0(x) = F(x)$ ,

$$\sqrt[3]{n}(\hat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})) \rightsquigarrow \frac{1}{\partial \Phi_0(\mathbf{x})} \partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x},1})(0) \sim c_0(\mathbf{x}) \underset{v \in \mathbb{R}}{\operatorname{argmin}} \{ \mathcal{G}(v) + v^2 \}$$

- ▶ Causal inference and program evaluation: Westling, Gilbert & Carone (2020, JRSSB).

# Generalized Grenander-Type Estimators: Framework

**Estimand:**  $\theta_0(\mathbf{x})$  at interior point  $\mathbf{x}$ , a monotone function.

**Estimator:**

$$\widehat{\theta}_n(\mathbf{x}) = [\partial_- \text{GCM}_{J_n}(\widehat{\Gamma}_n \circ \widehat{\Phi}_n^-)] \circ \widehat{\Phi}_n(\mathbf{x}),$$

for some (application specific)  $\widehat{\Gamma}_n \rightarrow_{\mathbb{P}} \Gamma_0$  and  $\widehat{\Phi}_n \rightarrow_{\mathbb{P}} \Phi_0$ .

**Asy Dist:**

$$n^{\mathfrak{q}/(1+2\mathfrak{q})}(\widehat{\theta}_n(\mathbf{x}) - \theta_0(\mathbf{x})) \rightsquigarrow \frac{1}{\partial \Phi_0(\mathbf{x})} \partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_{\mathbf{x}} + \mathcal{M}_{\mathbf{x}}^{\mathfrak{q}})(0),$$

$$\mathcal{M}_{\mathbf{x}}^{\mathfrak{q}}(v) = \frac{\partial^{\mathfrak{q}} \theta_0(\mathbf{x}) \partial \Phi_0(x)}{(1 + \mathfrak{q})!} v^{1+\mathfrak{q}}, \quad \mathfrak{q} = \min \{j \in \mathbb{N} : \partial^j \theta_0(\mathbf{x}) \neq 0\}.$$

**Our Goals:**

- ▶ Develop valid bootstrap-assisted & automatic distributional approximations.
- ▶ Develop valid inference procedures: e.g., confidence intervals for  $\theta_0(\mathbf{x})$ .

## Isotonic Density Estimation: Distribution Theory

- ▶  $\theta_0(x) = f(x)$ , a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .

### Switching Relation:

$$\mathbb{P} \left[ \sqrt[3]{n} (\hat{\theta}_n(x) - \theta_0(x)) > t \right] = \mathbb{P} \left[ \arg \max_{v \in S_n} \left\{ \hat{G}_{x,n}(v) + t \hat{L}_{x,n}(v) + M_{x,n}(v) \right\} < 0 \right],$$

where

$$\hat{G}_{x,n}(v) = -n^{2/3} \left[ \hat{\Gamma}_n(x + vn^{-1/3}) - \Gamma_0(x + vn^{-1/3}) - \hat{\Gamma}_n(x) - \Gamma_0(x) \right]$$

$$\hat{L}_{x,n}(v) = v$$

$$M_{x,n}(v) = -n^{2/3} \left[ \Gamma_0(x + vn^{-1/3}) - \Gamma_0(x) - \theta_0(x)vn^{-1/3} \right]$$

and

$$S_n = [-xn^{1/3}, \infty) \quad \text{with} \quad \mathbb{1}(v \in S_n) \rightarrow \mathbb{1}(v \in I)$$



## Isotonic Density Estimation: Distribution Theory

- ▶  $\theta_0(x) = f(x)$ , a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .

### Switching Relation:

$$\mathbb{P} \left[ \sqrt[3]{n}(\hat{\theta}_n(x) - \theta_0(x)) > t \right] = \mathbb{P} \left[ \arg \max_{v \in S_n} \left\{ \hat{G}_{x,n}(v) + t\hat{L}_{x,n}(v) + M_{x,n}(v) \right\} < 0 \right],$$

where

$$\hat{G}_{x,n}(v) = -n^{2/3} \left[ \hat{\Gamma}_n(x + vn^{-1/3}) - \Gamma_0(x + vn^{-1/3}) - \hat{\Gamma}_n(x) - \Gamma_0(x) \right] \rightsquigarrow \mathcal{G}_x(v)$$

$$\hat{L}_{x,n}(v) = v$$

$$M_{x,n}(v) = -n^{2/3} \left[ \Gamma_0(x + vn^{-1/3}) - \Gamma_0(x) - \theta_0(x)vn^{-1/3} \right] \rightarrow -\mathcal{M}_x^q(v) = -\frac{\partial f(x)}{2}v^2$$

which implies

$$\arg \max_{v \in S_n} \left\{ \hat{G}_{x,n}(v) + t\hat{L}_{x,n}(v) + M_{x,n}(v) \right\} \rightsquigarrow \arg \max_{v \in I} \left\{ \mathcal{G}_x(v) + tv - \mathcal{M}_x^q(v) \right\}$$

## Isotonic Density Estimation: Distribution Theory

$$\begin{aligned}\mathbb{P}\left[\sqrt[3]{n}(\widehat{\theta}_n(x) - \theta_0(x)) > t\right] &= \mathbb{P}\left[\arg\max_{v \in S_n} \left\{\widehat{G}_{x,n}(v) + t\widehat{L}_{x,n}(v) + M_{x,n}(v)\right\} < 0\right] \\ &\rightarrow \mathbb{P}\left[\arg\max_{v \in I} \left\{\mathcal{G}_x(v) + tv - \mathcal{M}_x^q(v)\right\} < 0\right] \\ &= \mathbb{P}\left[\partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_x^q)(0) > t\right]\end{aligned}$$

where

$$\mathcal{M}_x^q = \frac{\partial f(x)}{2} v^2$$

► In this stylized example,

$$q = 1, \quad \text{Phi}_0(x) = x, \quad \mathcal{G}_x(v) = \sqrt{f(x)}\mathcal{W}(v),$$

with  $\mathcal{W}(\cdot)$  two-sided Wiener process.

**Next:** let's investigate what happens when we apply the nonparametric bootstrap...

## Isotonic Density Estimation: Bootstrapping

- ▶  $\theta_0(x) = f(x)$  a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n^*(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n^*)(x)$  with  $\hat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i^* \leq x)$ .

### Switching Relation:

$$\mathbb{P}^* \left[ \sqrt[3]{n}(\hat{\theta}_n^*(x) - \hat{\theta}_n(x)) > t \right] = \mathbb{P}^* \left[ \arg \max_{v \in S_n^*} \left\{ \hat{G}_{x,n}^*(v) + t \hat{L}_{x,n}^*(v) + M_{x,n}^*(v) \right\} < 0 \right],$$

where

$$\hat{G}_{x,n}^*(v) = -n^{2/3} \left[ \hat{\Gamma}_n^*(x + vn^{-1/3}) - \hat{\Gamma}_n(x + vn^{-1/3}) - \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x) \right]$$

$$\hat{L}_{x,n}^*(v) = v$$

$$M_{x,n}^*(v) = -n^{2/3} \left[ \hat{\Gamma}_n(x + vn^{-1/3}) - \hat{\Gamma}_n(x) - \hat{\theta}_n(x)vn^{-1/3} \right]$$

and

$$S_n^* = [-xn^{1/3}, \infty) \quad \text{with} \quad \mathbb{1}(v \in S_n^*) \rightarrow \mathbb{1}(v \in I)$$

## Isotonic Density Estimation: Bootstrapping

- ▶  $\theta_0(x) = f(x)$  a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n^*(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n^*)(x)$  with  $\hat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i^* \leq x)$ .

### NP Bootstrap Failure:

$$\arg \max_{v \in S_n^*} \left\{ \hat{G}_{x,n}^*(v) + t \hat{L}_{x,n}^*(v) + M_{x,n}^*(v) \right\} \not\rightsquigarrow_{\mathbb{P}} \arg \max_{v \in I} \{ \mathcal{G}_{x,\alpha}(v) + tv + \mathcal{M}_{x,\alpha}(v) \}$$

because

$$\hat{G}_{x,n}^*(v) = -n^{2/3} [\hat{\Gamma}_n^*(x + vn^{-1/3}) - \hat{\Gamma}_n(x + vn^{-1/3}) - \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x)] \rightsquigarrow_{\mathbb{P}} \mathcal{G}_x(v)$$

$$\hat{L}_{x,n}^*(v) = v$$

$$M_{x,n}^*(v) = -n^{2/3} [\hat{\Gamma}_n(x + vn^{-1/3}) - \hat{\Gamma}_n(x) - \hat{\theta}_n(x)vn^{-1/3}] \not\rightsquigarrow_{\mathbb{P}} -\mathcal{M}_{x,n}(v) = -\frac{\partial f(x)}{2}v^2$$

- ▶ **Recall:** for Asy Dist, we instead had

$$M_{x,n}(v) = -n^{2/3} [\Gamma_0(x + vn^{-1/3}) - \Gamma_0(x) - \theta_0(x)vn^{-1/3}] \rightarrow -\mathcal{M}_{x,n}(v) = -\frac{\partial f(x)}{2}v^2$$

## Isotonic Density Estimation: Recap and Intuition

- ▶  $\theta_0(x) = f(x)$  a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .

### Asymptotic Distribution:

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}[\sqrt[3]{n}(\hat{\theta}_n(x) - \theta_0(x)) \leq t] - \mathbb{P}[\partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_x^q)(0) \leq t] \right| \rightarrow 0$$

**NP bootstrap:**  $\hat{\theta}_n^*(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n^*)(x)$  with  $\hat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i^* \leq x)$  is invalid,

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}^*[\sqrt[3]{n}(\hat{\theta}_n^*(x) - \hat{\theta}_n(x)) \leq t] - \mathbb{P}[\sqrt[3]{n}(\hat{\theta}_n(x) - \theta_0(x)) \leq t] \right| \not\rightarrow_{\mathbb{P}} 0$$

**This paper:** consistency can be achieved by reshaping  $\hat{\Gamma}_n^*$ .

- ▶ Intuition:
  - ▶ around  $x$ ,  $\hat{\Gamma}_n(x)$  has mean  $\Gamma_0(x) \approx \Gamma_0(x) + f(x)(x - x) + \frac{1}{2} \partial f(x)(x - x)^2$
  - ▶ whereas the mean of  $\hat{\Gamma}_n^*(x)$  under the bootstrap distribution is given by  $\hat{\Gamma}_n(x)$ .
- ▶ Reshaping: Let  $\partial \tilde{f}_n(x)$  denote a consistent estimator of  $\partial f(x)$ , then

$$\tilde{\Gamma}_n^*(x) = \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x) + \hat{\theta}_n(x)(x - x) + \frac{1}{2} \partial \tilde{f}_n(x)(x - x)^2$$

## Isotonic Density Estimation: Bootstrap Consistency

- ▶  $\theta_0(x) = f(x)$  a monotone density with  $\Gamma_0(x) = \mathbb{P}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n)(x)$  with  $\hat{\Gamma}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i \leq x)$ .
- ▶  $\hat{\theta}_n^*(x) = \partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n^*)(x)$  with  $\hat{\Gamma}_n^*(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(X_i^* \leq x)$ .
- ▶  $\tilde{\theta}_n^*(x) = \partial_- \text{GCM}_{J_n}(\tilde{\Gamma}_n^*)(x)$  with

$$\tilde{\Gamma}_n^*(x) = \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x) + \hat{\theta}_n(x)(x - x) + \frac{1}{2} \partial \tilde{f}_n(x)(x - x)^2.$$

### Bootstrap-Assisted Validity:

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}^*[\tilde{\theta}_n^*(x) - \hat{\theta}_n(x) \leq t] - \mathbb{P}[\hat{\theta}_n(x) - \theta_0(x) \leq t] \right| \xrightarrow{\mathbb{P}} 0$$

provided that

$$\partial \tilde{f}_n(x) \rightarrow_{\mathbb{P}} \partial f(x).$$

- ▶ The consistency requirement is mild and easy to achieve automatically.
- ▶ Confidence intervals based on kernel estimator perform well in simulations.

## Generalized Grenander-Type Estimators: General Case

- ▶  $\theta_0(x)$  at interior point  $x$ , a monotone function.
- ▶  $\hat{\theta}_n(x) = [\partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n \circ \hat{\Phi}_n^-)] \circ \hat{\Phi}_n(x)$ , with  $(\hat{\Gamma}_n, \hat{\Phi}_n) \rightarrow_{\mathbb{P}} (\Gamma_0, \Phi_0)$ .

**Asymptotic Distribution:** using their Generalized Switching Relation,

$$n^{q/(1+2q)} (\hat{\theta}_n(x) - \theta_0(x)) \rightsquigarrow \frac{1}{\partial \Phi_0(x)} \partial_- \text{GCM}_{\mathbb{R}}(\mathcal{G}_x + \mathcal{M}_x^q)(0),$$

$$\mathcal{M}_x^q(v) = \frac{\partial^q \theta_0(x) \partial \Phi_0(x)}{(1+q)!} v^{1+q}, \quad q = \min \{j \in \mathbb{N} : \partial^j \theta_0(x) \neq 0\}.$$

**Bootstrap-assisted Inference** using reshaped estimator:

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}^*[\tilde{\theta}_n^*(x) - \hat{\theta}_n(x) \leq t] - \mathbb{P}[\hat{\theta}_n(x) - \theta_0(x) \leq t] \right| \xrightarrow{\mathbb{P}} 0$$

where

$$\tilde{\theta}_n^*(x) = [\partial_- \text{GCM}_{J_n}(\tilde{\Gamma}_n^* \circ (\hat{\Phi}_n^*)^-)] \circ \hat{\Phi}_n^*(x)$$

with

$$\tilde{\Gamma}_n^*(x) = \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x) + \hat{\theta}_n(x) \hat{\Phi}_n(x - x) + \tilde{\mathcal{M}}_{x,q}(x - x),$$

and provided that

$$\tilde{\mathcal{M}}_{x,q}(v) \rightarrow_{\mathbb{P}} \mathcal{M}_x^q(v).$$

# Grenander-Type Estimators: Bootstrap-assisted Inference

**Valid percentile CI** for monotone function  $\theta_0(x)$  at interior point  $x$ :

$$I_{1-a}^*(x) = \left[ \hat{\theta}_n(x) - q_{1-a/2}^* \quad , \quad \hat{\theta}_n(x) - q_{a/2}^* \right]$$

where

$$\blacktriangleright \hat{\theta}_n(x) = [\partial_- \text{GCM}_{J_n}(\hat{\Gamma}_n \circ \hat{\Phi}_n^-)] \circ \hat{\Phi}_n(x), \quad (\hat{\Gamma}_n, \hat{\Phi}_n) \rightarrow_{\mathbb{P}} (\Gamma_0, \Phi_0).$$

$$\blacktriangleright q_a^* = \inf\{t \in \mathbb{R} : \mathbb{P}^*[\tilde{\theta}_n^*(x) - \hat{\theta}_n(x) \leq t] \geq a\},$$

$$\tilde{\theta}_n^*(x) = [\partial_- \text{GCM}_{J_n}(\tilde{\Gamma}_n^* \circ (\hat{\Phi}_n^*)^-)] \circ \hat{\Phi}_n^*(x),$$

$$\tilde{\Gamma}_n^*(x) = \hat{\Gamma}_n^*(x) - \hat{\Gamma}_n(x) + \hat{\theta}_n(x)\hat{\Phi}_n(x - x) + \tilde{\mathcal{M}}_{x,q}(x - x),$$

$$\tilde{\mathcal{M}}_{x,q}(v) \rightarrow_{\mathbb{P}} \mathcal{M}_x^q(v).$$

**Key outstanding issue:** How to construct  $\tilde{\mathcal{M}}_{x,q}(v)$ ?



## Bootstrap-assisted Inference: Drift Estimation

**Recall:**  $\widehat{\theta}_n(x) = [\partial_- \text{GCM}_{J_n}(\widehat{\Gamma}_n \circ \widehat{\Phi}_n^-)] \circ \widehat{\Phi}_n(x)$ ,  $(\widehat{\Gamma}_n, \widehat{\Phi}_n) \rightarrow_{\mathbb{P}} (\Gamma_0, \Phi_0)$ ,

$$\mathcal{M}_x^q(v) = \frac{\partial^q \theta_0(x) \partial \Phi_0(x)}{(1+q)!} v^{1+q}, \quad q = \min \{j \in \mathbb{N} : \partial^j \theta_0(x) \neq 0\}.$$

- ▶ Sometimes,  $\partial^q \theta_0(x)$  and  $\partial \Phi_0(x)$  easy to characterize (if  $q$  is known!) and estimate.
  - ▶ For example,  $\pi_0(x) = \partial f(x)$  and  $\partial \Phi_0(x) = 1$  in Isotonic Density Estimation.
- ▶ In general, if  $q$  is known,  $\tilde{\mathcal{M}}_{x,q}(v)$  based on numerical derivative estimator:

$$\frac{\partial^q \theta_n(\widehat{x}) \partial \widehat{\Phi}_n(x)}{(q+1)!} = \epsilon_n^{-(q+1)} \sum_{k=1}^{q+1} (-1)^{k+q+1} \binom{q+1}{k} [\widehat{\Upsilon}_n(x + k\epsilon_n) - \widehat{\Upsilon}_n(x)]$$

where

- ▶  $\widehat{\Upsilon}_n(v) = \widehat{\Gamma}_n(v) - \widehat{\theta}_n(x) \widehat{\Phi}_n(v)$  and  $\epsilon_n > 0$  is a (small) tuning parameter.
- ▶  $\tilde{\mathcal{M}}_{x,q}(v) \rightarrow_{\mathbb{P}} \mathcal{M}_x^q(v)$  requires  $\epsilon_n \rightarrow 0$  and  $n\epsilon_n^{1+2q} \rightarrow \infty$ .
- ▶ Under additional conditions, MSE-optimal  $\epsilon_n$  can be obtained.
- ▶ Possible to develop estimator  $\tilde{\mathcal{M}}_{x,q}(v)$  adaptive to unknown  $q \leq \bar{q}$ .

	DGP 1				DGP 2				DGP 3			
	$\hat{\mathcal{D}}_{1,n}$	$\hat{\mathcal{D}}_{3,n}$	Coverage	Length	$\hat{\mathcal{D}}_{1,n}$	$\hat{\mathcal{D}}_{3,n}$	Coverage	Length	$\hat{\mathcal{D}}_{1,n}$	$\hat{\mathcal{D}}_{3,n}$	Coverage	Length
<b>Standard</b>			0.832	0.370			0.835	0.516			0.904	0.028
<b>m-out-of-n</b>												
$m = \lceil n^{1/2} \rceil$			0.900	0.413			0.909	0.583			0.940	0.031
$m = \lceil n^{2/3} \rceil$			0.872	0.399			0.879	0.556			0.921	0.029
$m = \lceil n^{4/5} \rceil$			0.856	0.391			0.862	0.544			0.913	0.029
<b>Reshaped</b>												
Oracle	1.000	0.000	0.942	0.393	1.00	0.000	0.943	0.549	0.000	1.000	0.943	0.029
ND known $\mathbf{q}$	1.045	0.000	0.950	0.396	1.04	0.000	0.944	0.543	0.000	1.012	0.935	0.028
ND robust	1.045	0.633	0.951	0.398	1.04	0.981	0.953	0.556	0.014	1.012	0.959	0.030

## Discussion and Conclusion

- ▶ Nonparametric bootstrap fails for Grenander-Type Estimators.
- ▶ Other valid resampling methods available change the bootstrap distribution.
- ▶ This paper:
  - ▶ Employs standard nonparametric bootstrap.
  - ▶ Reshapes estimator to deal with bootstrap inconsistency.
- ▶ Our method applies to many problems in econ, stats, biostats, ML and beyond:
  - ▶ Isotonic density [Grenander, 1956] and extensions (censoring, covariates).
  - ▶ Isotonic regression [Brunk, 1970] and extensions (censoring, covariates).
  - ▶ Monotone Hazard [Rao, 1970] and extensions (censoring, covariates).
  - ▶ Current Status [Ayer et al. 1955] and extensions (censoring, covariates).
- ▶ **Coming soon:** Smoothed pairwise maximum rank correlation and related problems.
  - ▶  $\sqrt{n}$ -consistent U-process optimizer but with Chernoff-type distribution