- 30. Knoop, L. L., and Baker, S. J. (2000) The splicing factor U1C represses EWS/FLI-mediated transactivation. *J Biol Chem* **275**, 24865-24871
- 31. Armknecht, S., Boutros, M., Kiger, A., Nybakken, K., Mathey-Prevot, B., and Perrimon, N. (2005) High-throughput RNA interference screens in Drosophila tissue culture cells. *Methods in enzymology* **392**, 55-73
- 32. Soeller, W. C., Poole, S. J., and Kornberg, T. (1988) In vitro transcription of the Drosophila engrailed gene. *Genes Dev* **2**, 68-81
- Wohlschlegel, J. A. (2009) Identification of SUMO-conjugated proteins and their SUMO attachment sites using proteomic mass spectrometry. *Methods Mol Biol* **497**, 33-49
- 34. Zhou, R., Mohr, S., Hannon, G. J., and Perrimon, N. (2013) Inducing RNAi in Drosophila cells by transfection with dsRNA. *Cold Spring Harbor protocols* **2013**, 461-463
- 35. Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S. L. (2013) TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome biology* **14**, R36
- 36. Love, M. I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 550
- 37. Herrmann, C., Van de Sande, B., Potier, D., and Aerts, S. (2012) i-cisTarget: an integrative genomics method for the prediction of regulatory features and cis-regulatory modules. *Nucleic Acids Res* **40**, e114
- 38. Lyne, R., Smith, R., Rutherford, K., Wakeling, M., Varley, A., Guillier, F., Janssens, H., Ji, W., McLaren, P., North, P., Rana, D., Riley, T., Sullivan, J., Watkins, X., Woodbridge, M., Lilley, K., Russell, S., Ashburner, M., Mizuguchi, K., and Micklem, G. (2007) FlyMine: an integrated database for Drosophila and Anopheles genomics. *Genome biology* 8, R129
- 39. Brantjes, H., Roose, J., van De Wetering, M., and Clevers, H. (2001) All Tcf HMG box transcription factors interact with Groucho-related co-repressors. *Nucleic Acids Res* **29**, 1410-1419
- 40. Daniels, D. L., and Weis, W. I. (2005) Beta-catenin directly displaces Groucho/TLE repressors from Tcf/Lef in Wnt-mediated transcription activation. *Nature structural & molecular biology* **12**, 364-371
- 41. Bouazoune, K., and Brehm, A. (2006) ATP-dependent chromatin remodeling complexes in Drosophila. *Chromosome Res* **14**, 433-449
- 42. Flores-Saaib, R. D., and Courey, A. J. (2000) Analysis of Groucho-histone interactions suggests mechanistic similarities between Groucho- and Tup1-mediated repression. *Nucleic Acids Res* **28**, 4189-4196
- 43. Sekiya, T., and Zaret, K. S. (2007) Repression by Groucho/TLE/Grg proteins: genomic site recruitment generates compacted chromatin in vitro and impairs activator binding in vivo. *Mol Cell* 28, 291-303
- 44. Edmondson, D. G., Smith, M. M., and Roth, S. Y. (1996) Repression domain of the yeast global repressor Tup1 interacts directly with histones H3 and H4. *Genes Dev* **10**, 1247-1259
- 45. Patel, S. B., and Bellini, M. (2008) The assembly of a spliceosomal small nuclear ribonucleoprotein particle. *Nucleic Acids Res* **36**, 6482-6493
- 46. Will, C. L., and Luhrmann, R. (2011) Spliceosome structure and function. *Cold Spring Harb Perspect Biol* **3**
- 47. Fisher, A. L., Ohsako, S., and Caudy, M. (1996) The WRPW motif of the hairy-related basic helix-loop-helix repressor proteins acts as a 4-amino-acid transcription repression and protein-protein interaction domain. *Mol Cell Biol* **16**, 2670-2677
- Thorne, N., Inglese, J., and Auld, D. S. (2010) Illuminating insights into firefly luciferase and other bioluminescent reporters used in chemical biology. *Chemistry & biology* **17**, 646-657
- 49. Kaul, A., Schuster, E., and Jennings, B. H. (2014) The Groucho co-repressor is primarily recruited to local target sites in active chromatin to attenuate transcription. *PLoS genetics* **10**, e1004595