

# Lecture 4

## Simple Solutions of the 1-D Transport Equation

### 1 The 1-D Monoenergetic Transport Equation

Consider the 1-D slab geometry, monoenergetic, transport equation with isotropic scattering:

$$\mu \frac{\partial \psi}{\partial x}(x, \mu) + \sigma_t \psi(x, \mu) = \frac{\sigma_s}{4\pi} \phi(x) + \frac{Q(x)}{4\pi}, \quad (1)$$

where

$$\phi = 2\pi \int_{-1}^{+1} \psi(x, \mu) d\mu \quad (2)$$

and  $\mu = \cos \theta$ , where  $\theta$  is illustrated in Fig. 1. The boundary conditions for this equation define  $\psi(x_L, \mu)$  for  $\mu > 0$ , and  $\psi(x_R, \mu)$  for  $\mu < 0$ .

#### 1.1 Common Boundary Conditions

Vacuum Boundary Conditions:

$$\psi(x_L, \mu) = 0, \text{ for } \mu > 0. \quad (3a)$$

Reflective Boundary Conditions:

$$\psi(x_L, \mu) = \psi(x_L, -\mu), \text{ for } \mu > 0. \quad (3b)$$




Figure 1: The direction variable,  $\theta$ . Note that the directional dependence is assumed to be azimuthally symmetric.

Periodic Boundary Conditions:

$$\psi(x_L, \mu) = \psi(x_R, \mu) \quad , \text{for } \mu > 0. \quad (3c)$$

Source Boundary Conditions:

$$\psi(x_L, \mu) = \psi(\mu) \quad , \text{for } \mu > 0. \quad (3d)$$

## 2 Pure Absorber Solutions

Consider the following problem.

$$\psi(x_L, \mu) = f(\mu) \quad , \text{for } \mu > 0, \quad (4a)$$

$$\psi(x_R, \mu) = 0 \quad , \text{for } \mu < 0, \quad (4b)$$

$$\sigma_s = 0, \quad (4c)$$

$$Q = 0. \quad (4d)$$

The corresponding transport equation is

$$\mu \frac{\partial \psi}{\partial x} + \sigma_a \psi = 0. \quad (4e)$$

Dividing Eq. (4e) by  $\mu$ , we get

$$\frac{\partial \psi}{\partial x} + \frac{\sigma_a}{\mu} \psi = 0. \quad (5)$$

Note that we have a simple first-order ODE for each value of  $\mu$ . The solution is a simple exponential. To see this, we first multiply Eq. (5) by  $e^{\frac{\sigma_a x}{\mu}}$ :

$$e^{\frac{\sigma_a x}{\mu}} \frac{\partial \psi}{\partial x} + e^{\frac{\sigma_a x}{\mu}} \frac{\sigma_a}{\mu} \psi = 0. \quad (6)$$

Equation (6) can be re-expressed as

$$\frac{\partial}{\partial x} \left[ e^{\frac{\sigma_a x}{\mu}} \psi(x, \mu) \right]. \quad (7)$$

So

$$e^{\frac{\sigma_a x}{\mu}} \psi(x, \mu) = c, \quad (8)$$

is a solution, or equivalently,

$$\psi(x, \mu) = c e^{-\frac{\sigma_a x}{\mu}}. \quad (9)$$

The boundary conditions determine  $c$ .

$$\psi(x_L, \mu) = c e^{-\frac{\sigma_a}{\mu} x_L} = f(\mu), \quad (10)$$

so

$$c = f(\mu) e^{\frac{\sigma_a}{\mu} x_L}. \quad (11)$$

Thus our complete solution is

$$\begin{aligned} \psi(x, \mu) &= f(\mu) e^{-\frac{\sigma_a}{\mu} (x - x_L)}, \text{ for } \mu > 0, \\ &= 0, \text{ for } \mu < 0. \end{aligned} \quad (12)$$

Each ray is exponentially attenuated proportional to the distance that it travels from  $x_L$  to  $x$ :




Figure 2: The distance  $s$  given  $x$  and  $\mu$ .

### 3 The Formal Solution with Scattering

Consider the following problem.

$$\psi(x_L, \mu) = f(\mu), \quad (13a)$$

$$\psi(x_R, \mu) = g(\mu), \quad (13b)$$

$$Q(x) = 0. \quad (13c)$$

The equation to be solved is

$$\mu \frac{\partial \psi}{\partial x} + \sigma_t \psi = \frac{\sigma_s}{4\pi} \phi(x), \quad (13d)$$

Use the integrating factor approach again:

$$\frac{\partial \psi}{\partial x} + \frac{\sigma_t}{\mu} \psi = \frac{\sigma_s}{4\pi\mu} \phi, \quad (14)$$

$$e^{\frac{\sigma_t x}{\mu}} \frac{\partial \psi}{\partial x} + e^{\frac{\sigma_t x}{\mu}} \frac{\sigma_t}{\mu} \psi = \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t x}{\mu}} \phi, \quad (15)$$

$$\frac{\partial}{\partial x} \left[ e^{\frac{\sigma_t x}{\mu}} \psi \right] = \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t x}{\mu}} \phi, \quad (16)$$

$$\psi(x, \mu) e^{\frac{\sigma_t x}{\mu}} - \psi(x_L, \mu) e^{\frac{\sigma_t x_L}{\mu}} = \int_{x_L}^x \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t x'}{\mu}} \phi(x') dx' \quad , \text{ for } \mu > 0, \quad (17a)$$

$$\psi(x_R, \mu) e^{\frac{\sigma_t x_R}{\mu}} - \psi(x, \mu) e^{\frac{\sigma_t x}{\mu}} = \int_x^{x_R} \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t x'}{\mu}} \phi(x') dx' \quad , \text{ for } \mu < 0, \quad (17b)$$

So the final solution is

$$\psi(x, \mu) = \psi(x_L, \mu) e^{-\frac{\sigma_t}{\mu}(x-x_L)} + \int_{x_L}^x \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x'-x)} \phi(x') dx' \quad , \text{ for } \mu > 0, \quad (18a)$$

and

$$\psi(x, \mu) = \psi(x_R, \mu) e^{\frac{\sigma_t}{\mu}(x_R - x)} - \int_x^{x_R} \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x' - x)} \phi(x') dx' \quad , \text{ for } \mu < 0. \quad (18b)$$

Note that if you know  $\phi(x)$ , you need only perform an integral to get the solution, but  $\phi(x)$  is actually an integral of  $\psi(x, \mu)$ . This suggests that you might be able to iterate to a solution. Specifically, you can use the order-of-scatter or Neumann series technique:

First we calculate the uncollided flux:

$$\psi^{(o)}(x, \mu) = \psi(x_L, \mu) e^{-\frac{\sigma_t}{\mu}(x - x_L)} \quad , \text{ for } \mu > 0, \quad (19a)$$

and

$$\psi^{(o)}(x, \mu) = \psi(x_R, \mu) e^{\frac{\sigma_t}{\mu}(x_R - x)} \quad , \text{ for } \mu < 0. \quad (19b)$$

Next we calculate the first-scattered flux:

$$\psi^{(1)}(x, \mu) = \int_{x_L}^x \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x' - x)} \phi^{(o)}(x') dx' \quad , \text{ for } \mu > 0, \quad (20a)$$

and

$$\psi^{(1)}(x, \mu) = - \int_x^{x_R} \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x' - x)} \phi^{(o)}(x') dx' \quad , \text{ for } \mu < 0. \quad (20b)$$

where

$$\phi^{(o)}(x) = 2\pi \int_{-1}^{+1} \psi^{(o)}(x, \mu) d\mu. \quad (21)$$

⋮

Continue on

$\vdots$

You calculate the  $n + 1$ 'th-scattered angular flux using the  $n$ 'th-scattered scalar flux:

$$\psi^{(n+1)} = \int_{x_L}^x \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x'-x)} \phi^{(n)}(x') dx' \quad , \text{ for } \mu > 0, \quad (22a)$$

and

$$\psi^{(n+1)} = - \int_x^{x_R} \frac{\sigma_s}{4\pi\mu} e^{\frac{\sigma_t}{\mu}(x'-x)} \phi^{(n)}(x') dx' \quad , \quad \mu < 0. \quad (22b)$$

Finally, add up the contributions to obtain the total angular flux:

$$\psi(x, \mu) = \sum_{n=0}^{\infty} \psi^{(n)}(x, \mu). \quad (23)$$

This is very closely related to a basic iteration technique used in  $S_n$  codes.