

Table A1. *Summary of the data of the fictive case*

	A1	A2	A3
$n_i$	41.00	21.00	31.00
$\bar{X}$	24	23	27
$s^2$	81.75	10.075	38.40

## Appendix 1: The Mathematical Development of the $F$ -test, $W$ -test, and $F^*$ -test: Numerical Example

A summary is presented in Table A1. The complete example is available on Github. The DV is a score that can vary from 0 to 40. The IV is a three-level factor A (levels =  $A_1$ ,  $A_2$  and  $A_3$ ).

The global mean (i.e. the mean of the global dataset) is a weighted mean of the group means:

$$\frac{(41 * 24) + (21 * 23) + (31 * 27)}{41 + 21 + 31} = \frac{2304}{93} \approx 24.77$$

The  $F$ -test statistic and degrees of freedom are computed by applying formulas (1), (2) and (3):

$$F = \frac{\frac{1}{3-1} [41 * (24 - \frac{2304}{93})^2 + 21 * (23 - \frac{2304}{93})^2 + 31 * (27 - \frac{2304}{93})^2]}{\frac{1}{93-3} [(41-1) * 81.75 + (21-1) * 10.075 + (31-1) * 38.4]} \approx 2.377$$

$$df_n = 3 - 1 = 2$$

$$df_d = 93 - 3 = 90$$

The  $F^*$ -test and his degrees of freedom are computed by applying formulas 4, 5 and 6:

$$F^* = \frac{41 * (24 - \frac{2304}{93})^2 + 21 * (23 - \frac{2304}{93})^2 + 31 * (27 - \frac{2304}{93})^2}{(1 - \frac{41}{93}) * 81.75 + (1 - \frac{21}{93}) * 10.075 + (1 - \frac{31}{93}) * 38.4} \approx 3.088$$

$$df_n = 3 - 1 = 2$$

$$df_d = \frac{1}{\frac{(\frac{(1-\frac{41}{93}) * 81.75}{\sum_{j=1}^k (1-\frac{n_j}{N}) s_j^2})^2}{41-1} + \frac{(\frac{(1-\frac{21}{93}) * 10.075}{\sum_{j=1}^k (1-\frac{n_j}{N}) s_j^2})^2}{21-1} + \frac{(\frac{(1-\frac{31}{93}) * 38.4}{\sum_{j=1}^k (1-\frac{n_j}{N}) s_j^2})^2}{31-1}} \approx 81.149$$

$$\text{Where } \sum_{j=1}^k (1 - \frac{n_j}{N}) * s_j^2 \approx 79.11$$

Finally, the  $W$ -test and his degrees of freedom are computed in applying formulas 7, 8 and 9:

$$W = \frac{\frac{1}{3-1} [\frac{41}{81.75} (24 - \bar{X}')^2 + \frac{21}{10.075} (23 - \bar{X}')^2 + \frac{31}{38.4} (27 - \bar{X}')^2]}{\frac{2(3-2)}{3^2-1} [(\frac{1}{41-1})(1 - \frac{41}{81.75})^2 + (\frac{1}{21-1})(1 - \frac{21}{10.075})^2 + (\frac{1}{31-1})(1 - \frac{31}{38.4})^2] + 1} \approx 4.606$$

Where:

$$w = \sum_{j=1}^k w_j \approx 3.39$$

$$\bar{X}' = \frac{\sum_{j=1}^k (w_j \bar{x}_j)}{w} \approx 24.1$$

$$df_n = 3 - 1$$

$$df_d = \frac{3^2 - 1}{3[\frac{(1-\frac{w_j}{w})^2}{41-1} + \frac{(1-\frac{w_j}{w})^2}{21-1} + \frac{(1-\frac{w_j}{w})^2}{31-1}]} \approx 59.32$$

One should notice that in this example, the biggest sample size has the biggest variance. As previously mentioned, it means that the  $F$ -test will be too conservative, because the  $F$  value decreases. The  $F^*$ -test will also be a little too conservative, even if the test is less affected than the  $F$ -test. As a consequence:  $W > F^* > F$ .

## Appendix 2: Justification for the choice of distributions in simulations

The set of simulations described in the article was repeated for 7 distributions. We used R commands to generate data from different distributions:

- $k$  normal distributions (Figure A2.1): in order to assess the Type I error rate and power of the different tests under the assumption of normality, data were generated by means of the function “rnorm” (from the package “stats”; “R: The Normal Distribution,” 2016).
- $k$  double exponential distributions (Figure A2.2): In order to assess the impact of high kurtosis on the Type I error rate and power of all tests, data were generated by means of the function “rdoublex” (from the package “smoothest”; “R: The double exponential (Laplace) distribution,” 2012).
- $k$  mixed normal distributions (Figure A2.3): In order to assess the impact of extremely high kurtosis on the Type I error rate and power of all tests, regardless of variance, data were generated by means of the function “rmixnorm” (from the package “bda”; Wang & Wang, 2015).
- $k$  normal right skewed distributions (Figure A2.4): In order to assess the impact of moderate skewness on the Type I error rate and power, data were generated by means of the function “rsnorm” (from the package “fGarch”; “R: Skew Normal Distribution,” 2017). The normal skewed distribution was chosen because it is the only skewed distribution where the standard deviation ratio can vary without having an impact on skewness.
- $k-1$  normal left skewed distributions (Figure A2.5) and 1 normal right skewed distribution (Figure A2.5): In order to assess the impact of unequal shapes, in terms of skewness, on the Type I error rate and power, when data have moderate skewness, data were generated by means of the functions “rsnorm” (from the package “fGarch”; “R: Skew Normal Distribution,” 2017).
- $k-1$  chi-squared distributions with two degrees of freedom (See Figure A2.7), and one normal right skewed distribution (Figure A2.5): In order to assess the impact of high asymmetry on the Type I error rate and power,  $k-1$  distributions were generated by means of the functions “rchisq” (“R: The (non-central) Chi-squared Distribution,” 2016). The last distribution was generated by means of “rsnorm” in order to follow a normal right skewed distribution with a mean of 2 (from the package “fGarch”; “R: Skew Normal Distribution,” 2017).




Figure 1: centered normal probability density function, as a function of the population SD




Figure 2: centered double exponential probability density function, as a function of the population SD

- $k-1$  chi-squared distributions with two degrees of freedom (See Figure A2.7), and one normal left skewed distribution (Figure A2.5): In order to assess the impact of unequal shapes, in terms of skewness, on Type I error rate and power when distributions have extreme skewness,  $k-1$  distributions were generated by means of the functions “rchisq” (“R: The (non-central) Chi-squared Distribution,” 2016). The last distribution was generated by means of “rsnorm” in order to follow a normal right skewed distribution with a mean of 2 (from the package “fGarch”; “R: Skew Normal Distribution,” 2017)

```
# {r "", echo=FALSE, fig.width = 15,fig.height=8,out.width = '400px',fig.cap = "centered
normal right #skewed probability density function, as a function of the population SD"}
# knitr::include_graphics("Rmarkdown folder/Rmarkdown inputs/") #

# {r "", echo=FALSE, fig.width = 15,fig.height=8,out.width = '400px',fig.cap = "centered
normal left #skewed probability density function, as a function of the population SD"}
# knitr::include_graphics("Rmarkdown folder/Rmarkdown inputs/") #
```




Figure 3: centered mixed normal probability density function, as a function of the population SD




Figure 4: chi-squared with 2 degrees of freedom probability density function, as a function of the population SD