

CLINICAL TRIAL RESULTS REPORT (RAP-504)

1. PARTICIPANT DISPOSITION

A total of 687 participants were screened, and 650 were randomized (325 to ArthroBalance-X, 325 to Placebo). The completion rate was 97.2%. Demographics: Baseline mean DAS28-CRP was 5.8 across all groups, with mean disease duration of 7.2 years.

2. EFFICACY ANALYSIS (PRIMARY)

The ArthroBalance-X group showed rapid and sustained disease improvement with remission observed in 48% by Week 24 ($p < 0.001$) compared to 8% in the placebo group. Mean DAS28-CRP reduction was 3.2 points in the treatment group vs. 0.4 points in placebo.

Conclusion: The trial successfully met its primary endpoint, exceeding the pre-specified target of 45% remission achievement.

3. SECONDARY OUTCOMES

ACR Response: ACR20/50/70 response rates were 76%/62%/48% in ArthroBalance-X vs. 12%/4%/2% in placebo ($p < 0.001$).

Low Disease Activity: 68% of ArthroBalance-X participants achieved minimal disease activity (MDA) criteria vs. 14% in placebo.

Functional Improvement: Mean HAQ-DI reduction was 0.68 points in treatment group vs. 0.08 points in placebo.

Time to Remission: Median time to DAS28 remission was 8 weeks in ArthroBalance-X arm.

4. SAFETY PROFILE

ArthroBalance-X demonstrated a manageable safety profile consistent with TNF- α inhibitor class. The most common adverse events (AEs) were:

- Upper respiratory tract infections: 13.2% (Predicted: 12-15%)
- Injection site reactions: 9.1% (Predicted: 8-10%)
- Herpes zoster reactivation: 2.8% (Predicted: 2-3%)

No cases of tuberculosis reactivation, serious fungal infections, or malignancies were reported. One case of non-serious hepatitis B reactivation was observed and managed successfully. No

treatment-related deaths occurred.