

CLINICAL TRIAL RESULTS REPORT (GDP-401)

1. PARTICIPANT DISPOSITION

A total of 798 participants were screened, and 750 were randomized (375 to GlucoControl-7, 375 to Placebo). The completion rate was 93.5%. Demographics: Baseline mean HbA1c was 8.3% across all groups, with mean BMI of 31.2 kg/m².

2. EFFICACY ANALYSIS (PRIMARY)

The GlucoControl-7 group showed consistent and significant HbA1c reduction starting at Week 4. At Week 24, the mean reduction in HbA1c was 1.8% ($p < 0.001$) compared to 0.2% in the placebo group.

Conclusion: The trial successfully met its primary endpoint, exceeding the pre-specified target of 1.5% reduction by 0.3%.

3. SECONDARY OUTCOMES

Fasting Glucose: Mean reduction of 28.4 mg/dL in GlucoControl-7 group vs. 3.1 mg/dL in placebo ($p < 0.001$).

Glycemic Control: 68% of participants in GlucoControl-7 group achieved HbA1c $< 7.0\%$, compared to 8% in placebo arm.

Body Weight: Mean weight reduction of 2.3 kg observed with GlucoControl-7 vs. 0.1 kg with placebo.

Lipid Profile: Statistically significant improvements in LDL-C (12% reduction) and triglycerides (18% reduction).

4. SAFETY PROFILE

GlucoControl-7 was generally well-tolerated. The most common adverse events (AEs) were:

- Nausea: 4.2% (Predicted: 3-5%)
- Headache: 5.1% (Predicted: 4-6%)
- Mild Diarrhea: 2.8% (New observation)

No cases of pancreatitis, severe hypoglycemia, or acute kidney injury were noted. Liver function tests remained within normal limits in 99.2% of participants.

