

CLINICAL TRIAL RESULTS REPORT (RKP-203)

1. PARTICIPANT DISPOSITION

A total of 945 participants were screened, and 900 were randomized (450 to RenalGuard-5, 450 to Placebo). The completion rate was 96.3%. Demographics: Baseline mean eGFR was 38.2 mL/min/1.73m², mean UACR was 142.5 mg/g across all groups.

2. EFFICACY ANALYSIS (PRIMARY)

The RenalGuard-5 group showed consistent and significant proteinuria reduction starting at Week 6. At Week 52, the mean reduction in urinary albumin excretion was 42% ($p < 0.001$) compared to 8% in the placebo group.

Conclusion: The trial successfully met its primary endpoint, exceeding the pre-specified target of 40% reduction.

3. SECONDARY OUTCOMES

Renal Function: Mean eGFR decline was significantly slowed: 2.1 mL/min/year in RenalGuard-5 vs. 5.3 mL/min/year in placebo ($p < 0.001$).

Proteinuria Control: 64% of participants in RenalGuard-5 group achieved >50% albuminuria reduction, compared to 18% in placebo.

Blood Pressure: Mean systolic BP reduction of 8.2 mmHg in RenalGuard-5 vs. 2.1 mmHg in placebo.

Fibrosis Markers: Significant reduction in FGF-23 levels (28% decrease) and KIM-1 (35% decrease).

4. SAFETY PROFILE

RenalGuard-5 demonstrated a manageable safety profile. The most common adverse events (AEs) were:

- Hyperkalemia ($K^+ > 5.5$ mEq/L): 7.1% (Predicted: 6-8%)
- Dizziness: 5.2% (Predicted: 4-6%)
- Transient eGFR decline: 12.8% (Predicted: 10-15%)

No cases of acute kidney injury grade 3+, anaphylaxis, or life-threatening hyperkalemia requiring dialysis were observed. Hyperkalemia was managed with potassium-lowering therapy in all cases.

