

1 The Missing Baryon Problem: Investigating lost matter using the 2 Universe's baby picture

3 ¹*The University of Melbourne, Victoria, Australia*

4 1 The Cosmic Microwave Background

5 Fifty five years after its accidental discovery, the first image of a baby universe has still not given up all its secrets.
6 The afterglow of the Big Bang, known as the Cosmic Microwave Background (CMB), is the only faint remnant of
7 the early universe currently still visible in the sky, and it has been central in ushering in a golden age of cosmology.

8 The CMB is made up of microwaves which carry very low power, and the characteristic splotches in it
9 provide key insights into conditions in the early universe. This allows astronomers to determine characteristics,
10 such as the relative proportions of matter, dark matter, dark energy, and other fundamental parameters. These
11 splotches yield the most precise measures of the age, geometry, and composition of the universe to date.

12 In more recent years, a number of experiments have been done specifically to examine the properties of
13 the CMB, including the Atacama Cosmology Telescope in the mountains of Chile, the South Pole Telescope,
14 and the European Space Agency's *Planck* Satellite. These experiments have broadened the scope of the original
15 detection, leading to the collection of an unprecedented amount of data, and applications across a number of areas
16 of astronomy.

17 2 The Missing Baryon Problem

18 The CMB has allowed scientists to determine with a very high level of confidence the composition of the universe.
19 The universe is broadly comprised of light, regular matter, and the infamous dark matter and dark energy, the
20 precise ratios of which are very tightly constrained.

21 However when astronomers look at the sky, and measure the amount of matter they see with telescopes, it
22 becomes apparent that only about half of the universe's regular matter - known as 'baryons' - is present in the
23 light emitted from stars and galaxies. This begs the question, where is all the missing matter?

24 It was clear that the matter must be there, since it is very clear to discern between dark matter and baryons
25 in physical models. The universe could not have evolved to be what we see today if that fraction wasn't correct.
26 But it was also apparent that only about 10 percent of baryons exist in galaxies themselves, with some 30 to 40
27 percent existing in gas clouds surrounding galaxies. This still leaves a large fraction unaccounted for.

28 One possible answer arises from simulations. Astronomers can model the evolution of the universe by
29 tracking individual clumps of gas, and when they evolve them forward in time, they notice something interesting.
30 The gas that doesn't collapse to form galaxies stretches out into long filaments.

31 These filaments contain very small amounts of matter, superheated to temperatures between thousands and
32 millions of degrees, and which is spread out over the vast reaches of space. The combination of its speed and
33 these distances makes these filaments very sparse - only a few atoms per cubic meter.

34 Detecting these filaments has proven to be a difficult task, since the scarcity of the gas makes it very unlikely
35 that there will be any observable signal. The only possible method of direct detection of this matter involves using
36 a quasar, a galaxy with a very active super-massive black hole at its centre, as a backlight. Unfortunately this is
37 an incredibly rare event, since it requires us to find just such an object, and for its beam to pass through both a
38 filament and to be captured by a telescope.

39 **3 A new use for the oldest light in the universe**

40 One possible solution is to use the CMB itself. At a fundamental level, the CMB is just light, sitting at the edge
41 of the observable universe, and light is interactive. It is distorted by the matter in between it and our detectors in

42 a number of different ways.

43 Scientists appeal first to a phenomenon known as Compton Scattering. The subject of the 1927 Nobel Prize,
44 and named after its discoverer, Arthur Holly Compton, this effect describes how light can exchange some of its
45 energy with electrons, and vice versa. Since there is hot gas in between the CMB and us, there will be sections
46 of the sky which will have absorbed some energy from the intervening gas, and look more energetic than its
47 surroundings.

48 A second possible tool for analysis is to take advantage of the distortion that matter makes in space-time, an
49 effect known as Gravitational Lensing. This distortion leaves characteristic traces in the light that we see, making
50 it appear rotated when we compare it to the background light.

51 On their own, the signals obtained by these techniques are too small for us to detect for a single filament,
52 there is just not enough matter. We can however, use this fact to our advantage, since by definition, the noise in
53 the signal is random. This means that if you add up enough of it, it will eventually zero out. Conversely, the
54 signals we are looking for should stack on top of each other. Therefore, we can make a direct detection if we add
55 multiple images on top of each other. We expect the noise to cancel out, leaving just the signal.

56 **4 Tracing the Large Scale Structure**

57 With this in mind, scientists know that they must use locations where they expect to find the missing matter in
58 order to make a detection of any kind. To do so, they look to comprehensive galaxy surveys, such as the Sloan
59 Digital Sky Survey (SDSS) and the Dark Energy Survey (DES), which have created some of the most detailed
60 three-dimensional maps of the universe we have to date.

61 Once the galaxies are located, they are paired up based on whether they are likely to have a filament con-
62 necting them, and located in the CMB. The signals are then stacked on top of each other, hopefully revealing the

63 missing baryons.

64 This search has been shown to work with low resolution images of the CMB, revealing the missing matter
65 with relative certainty, however there is still more to be done to confirm this. By accessing higher resolution
66 images, scientists hope to confirm the location of the missing baryon fraction, and so remove any shred of doubt
67 as to the validity of current cosmological models.

68