

Geriatrics Practice Parameter

Mark Grant

2022-10-14T00:00:00-05:00

Table of contents

Preface	3
1 Introduction	4
2 Potentially Inappropriate Medications (KQ5)	5
2.0.1 Interventions of Interest	5
3 Summary	6
References	7

Preface

This is a Quarto book.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

1 + 1

[1] 2

1 Introduction

This is a book created from markdown and executable code.

See Knuth (1984) for additional discussion of literate programming.

```
1 + 1
```

```
[1] 2
```

2 Potentially Inappropriate Medications (KQ5)

Among geriatric patients undergoing surgery and anesthesia, do commonly used potentially inappropriate medications administered during the perioperative period increase the risk of postoperative delirium or other adverse outcomes?

2.0.1 Interventions of Interest

- Anticholinergics
- Antipsychotics
- Benzodiazepines
- Corticosteroids
- Ketorolac and other NSAIDs
- H2-receptor antagonists
- Nonbenzodiazepine benzodiazepine receptor agonist hypnotics: eszopiclone, zaleplon, zolpidem

3 Summary

In summary, this book has no content whatsoever.

`1 + 1`

[1] 2

References

Knuth, Donald E. 1984. “Literate Programming.” *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.