

ENRICHED BISIMULATIONS

MATTHEW DiMEGLIO
(Joint work with Bryce Clarke)

APPLIED CATEGORY THEORY 2023

LABELLED TRANSITION SYSTEMS

Used to describe the possible behaviours of discrete processes.

LABELLED TRANSITION SYSTEMS

Used to describe the possible behaviours of discrete processes.

S
set of states

A
set of actions

$\xrightarrow{\alpha} \subseteq S \times S$
transition relation
for each $\alpha \in A$

LABELLED TRANSITION SYSTEMS


Used to describe the possible behaviours of discrete processes.

S
set of states

A
set of actions

$\xrightarrow{\alpha} \subseteq S \times S$
transition relation
for each $\alpha \in A$

EXAMPLE: Vending Machine



$$S_1 = \{On, Off\}$$


$$A = \{coin, biscuit, chocolate\}$$

STRONG BISIMULATION




(S and T are A -labelled transition systems)

STRONG BISIMULATION




(S and T are A -labelled transition systems)

STRONG BISIMULATION



(S and T are A -labelled transition systems)

STRONG BISIMULATION



(S and T are A -labelled transition systems)

FUNCTIONAL STRONG BISIMULATIONS*

3

A function $f:S \rightarrow T$ whose graph is a strong bisimulation.

(S and T are A -labelled transition systems)

*Also called abstraction homomorphisms and pure morphisms

FUNCTIONAL STRONG BISIMULATIONS*

A function $f:S \rightarrow T$ whose graph is a strong bisimulation.

PROPOSITION:

spans of functional
strong bisimulations

$$\begin{array}{ccc} S & \xleftarrow{f_1} & X \\ & & \xrightarrow{f_2} T \\ S & \xleftarrow{\pi_1} & R \\ & & \xrightarrow{\pi_2} T \end{array}$$

split surjection



strong bisimulations

$$\begin{array}{c} \text{Im}\langle f_1, f_2 \rangle \subseteq S \times T \\ R \subseteq S \times T \end{array}$$



(S, T and X are A -labelled transition systems)

*Also called abstraction homomorphisms and pure morphisms

\mathcal{V} -ENRICHED ASYMMETRIC LENS

$$F: A \rightarrow B$$

$$A(a,a') \xrightarrow{F_{a,a'}} B(Fa,Fa')$$

hom maps

$$A_{\text{ob}} \xrightarrow{F} B_{\text{ob}}$$

object map

$$B(Fa,b') \xrightarrow{\sum_{a' \in F^{-1}\{b'\}} F_{a,a'}} A(a,a')$$

lifting maps

| \mathcal{V} | \mathcal{V} -ENRICHED CATEGORY | \mathcal{V} -ENRICHED ASYMMETRIC LENS |
|----------------------|----------------------------------|---|
| Set | category | delta lens |
| wSet | weighted category | weighted lens |
| $([0,\infty], \geq)$ | metric space | weak submetry |

\mathcal{V} is a distributive monoidal category

A -labelled transition systems = $\mathcal{P}(A)$ -enriched graphs

S

set of states

$\xrightarrow{\alpha} \subseteq S \times S$

transition relation
for each $\alpha \in A$

$S = \mathcal{S}_{\text{ob}}$

\mathcal{S}_{ob}

set of objects

$s \xrightarrow{\alpha} s' \iff \alpha \in \mathcal{S}(s, s')$

$\mathcal{S}(s, s') \in \mathcal{P}(A)$

hom object
for all $s, s' \in S$

A -labelled transition systems = $\mathcal{P}(A)$ -enriched graphs

functional strong bisimulations = asymmetric lenses

A function $f: S \rightarrow T$ such that

(1) $s \xrightarrow{\alpha} s'$ implies $fs \xrightarrow{\alpha} fs'$,

(2) $fs \xrightarrow{\alpha} t'$ implies exists s' with $s \xrightarrow{\alpha} s'$ and $fs' = t'$.

A function $f: S_{\text{ob}} \rightarrow T_{\text{ob}}$ such that

(1) $S(s, s') \subseteq T(fs, fs')$,

(2) $T(fs, t') \subseteq \bigcup_{s' \in f^{-1}\{t'\}} S(s, s')$.

A -labelled transition systems = $\mathcal{P}(A)$ -enriched graphs

functional strong bisimulations = asymmetric lenses

strong bisimulations = ?

\mathcal{V} -ENRICHED BISIMULATION

A relation $R \subseteq A_{ob} \times B_{ob}$ together with morphisms

$$A(a, a') \xrightarrow[\sum_{b': (a', b') \in R}]{} \sum B(b, b') \quad \text{and} \quad \sum A(a, a') \xleftarrow[\sum_{a': (a', b') \in R}]{} B(b, b')$$

that are compatible with identities and composites.

\mathcal{V} is a distributive monoidal category


\mathcal{V} -ENRICHED BISIMULATION

A relation $R \subseteq A_{\text{ob}} \times B_{\text{ob}}$ together with morphisms

$$A(a, a') \xrightarrow{\vec{R}_{a, a', b}} \sum_{b' : (a', b') \in R} B(b, b') \quad \text{and} \quad \sum_{a' : (a', b') \in R} A(a, a') \xleftarrow{\vec{R}_{a, b, b'}} B(b, b')$$

that are compatible with identities and composites.

These don't compose!



\mathcal{V} is a distributive monoidal category


\mathcal{V} -ENRICHED BISIMULATION

A relation $R \subseteq A_{\text{ob}} \times B_{\text{ob}}$ together with morphisms

$$A(a, a') \xrightarrow{\vec{R}_{a, a', b}} \sum_{b' : (a', b') \in R} B(b, b') \quad \text{and} \quad \sum_{a' : (a', b') \in R} A(a, a') \xleftarrow{\vec{R}_{a, b, b'}} B(b, b')$$

that are compatible with identities and composites.

These don't compose!



\mathcal{V} is a distributive monoidal category

\mathcal{V} -ENRICHED BISIMULATION

A relation $R \subseteq A_{\text{ob}} \times B_{\text{ob}}$ together with morphisms

$$A(a, a') \xrightarrow{\vec{R}_{a, a', b}} \sum_{b': (a', b') \in R} B(b, b') \quad \text{and} \quad \sum_{a': (a', b') \in R} A(a, a') \xleftarrow{\vec{R}_{a, b, b'}} B(b, b')$$

that are compatible with identities and composites.

These don't compose!



\mathcal{V} is a distributive monoidal category

\mathcal{V} -ENRICHED BISIMULATION

A relation $R \subseteq A_{\text{ob}} \times B_{\text{ob}}$ such that

$$A(a, a') \leq \bigvee_{b': (a', b') \in R} B(b, b') \quad \text{and} \quad \bigvee_{a': (a', b') \in R} A(a, a') \leq B(b, b').$$

These do compose:

$$(a, b) \in R \subseteq A_{\text{ob}} \times B_{\text{ob}}$$

$$(b, c) \in S \subseteq B_{\text{ob}} \times C_{\text{ob}}$$

$$A(a, a') \leq \bigvee_{b': (a', b') \in R} B(b, b') \leq \bigvee_{b': (a', b') \in R} \bigvee_{c': (b, c') \in S} C(c, c') \leq \bigvee_{c': (a', c') \in R; S} C(c, c')$$

For graphs: \mathcal{V} is a suplattice

For categories: \mathcal{V} is a quantale

EXAMPLES

strong bisimulation of
 A -labelled transition systems $=$ bisimulation of
 $P(A)$ -enriched graphs

weak bisimulation of
 $A \sqcup \{\tau\}$ -labelled transition systems $=$ bisimulation of
 $P(A^*)$ -enriched categories

bisimulation of
Kripke frames $=$ bisimulation of
 $\mathbf{2}$ -enriched graphs

PROPOSITION:

spans of \mathcal{V} -enriched
asymmetric lenses

split surjection \rightarrow

\mathcal{V} -enriched
bisimulations

$$A_{ob} \xleftarrow{f_1} X_{ob} \xrightarrow{f_2} B_{ob} \quad \mapsto$$

$$\text{Im}\langle f_1, f_2 \rangle$$

$$A_{ob} \xleftarrow{\pi_1} R_{ob} \xrightarrow{\pi_2} B_{ob}$$

$$R \subseteq A_{ob} \times B_{ob}$$

$$R_{ob} = R \quad R\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} a' \\ b' \end{pmatrix}\right) = A(a, a') \wedge B(b, b') \Leftarrow$$

PROPOSITION:

spans of \mathcal{V} -enriched
asymmetric lenses

split surjection \rightarrow

\mathcal{V} -enriched
bisimulations

$$\mathcal{A}_{\text{ob}} \xleftarrow{f_1} \mathcal{X}_{\text{ob}} \xrightarrow{f_2} \mathcal{B}_{\text{ob}}$$

\mapsto

$$\text{Im}\langle f_1, f_2 \rangle$$

$$\mathcal{A}_{\text{ob}} \xleftarrow{\pi_1} \mathcal{R}_{\text{ob}} \xrightarrow{\pi_2} \mathcal{B}_{\text{ob}}$$

$$\mathcal{R}_{\text{ob}} = R \quad R((\begin{smallmatrix} a \\ b \end{smallmatrix}), (\begin{smallmatrix} a' \\ b' \end{smallmatrix})) = \mathcal{A}(a, a') \wedge \mathcal{B}(b, b')$$

$$R \subseteq \mathcal{A}_{\text{ob}} \times \mathcal{B}_{\text{ob}}$$

$(a, b) \in \text{Im}\langle f_1, f_2 \rangle \iff f_1 x = a \text{ and } f_2 x = b \text{ for some } x \in \mathcal{X}_{\text{ob}}$

$$\mathcal{A}(a, a') = \mathcal{A}(f_1 x, a') \leq \bigvee_{x' \in f_1^{-1} a'} \mathcal{X}(x, x') \leq \bigvee_{x' \in f_1^{-1} a'} \mathcal{B}(f_2 x, f_2 x') = \bigvee_{\substack{b' : (a', b') \\ \in \text{Im}\langle f_1, f_2 \rangle}} \mathcal{B}(b, b')$$

PROPOSITION:

spans of \mathcal{V} -enriched
asymmetric lenses

split surjection \rightarrow

\mathcal{V} -enriched
bisimulations

$$\mathcal{A}_{\text{ob}} \xleftarrow{f_1} \mathcal{X}_{\text{ob}} \xrightarrow{f_2} \mathcal{B}_{\text{ob}}$$

\mapsto

$$\text{Im}\langle f_1, f_2 \rangle$$

$$\mathcal{A}_{\text{ob}} \xleftarrow{\pi_1} \mathcal{R}_{\text{ob}} \xrightarrow{\pi_2} \mathcal{B}_{\text{ob}}$$

$$\mathcal{R}_{\text{ob}} = R \quad R((\begin{smallmatrix} a \\ b \end{smallmatrix}), (\begin{smallmatrix} a' \\ b' \end{smallmatrix})) = A(a, a') \wedge B(b, b')$$

$$R \subseteq \mathcal{A}_{\text{ob}} \times \mathcal{B}_{\text{ob}}$$

$$R(r, r') = A(\pi_1 r, \pi_1 r') \wedge B(\pi_2 r, \pi_2 r') \leq A(\pi_1 r, \pi_1 r')$$

$$\begin{aligned} A(\pi_1 r, a') &= A(\pi_1 r, a') \wedge A(\pi_1 r, a') \leq A(\pi_1 r, a') \wedge \bigvee_{b': (a', b') \in R} B(\pi_2 r, b') \\ &= \bigvee_{b': (a', b') \in R} A(\pi_1 r, a') \wedge B(\pi_2 r, b') = \bigvee_{r' \in \pi_1^{-1} a'} R(r, r') \end{aligned}$$

PROPOSITION:

spans of \mathcal{V} -enriched
asymmetric lenses

split surjection \rightarrow

\mathcal{V} -enriched
bisimulations

$$A_{ob} \xleftarrow{f_1} X_{ob} \xrightarrow{f_2} B_{ob}$$

\mapsto

$$\text{Im}\langle f_1, f_2 \rangle$$

$$A_{ob} \xleftarrow{\pi_1} R_{ob} \xrightarrow{\pi_2} B_{ob}$$

$$R_{ob} = R \quad R\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} a' \\ b' \end{pmatrix}\right) = A(a, a') \wedge B(b, b')$$

$$R \subseteq A_{ob} \times B_{ob}$$

local reflection*

$$\boxed{\mathcal{V}\text{-}\underline{\text{LensSpan}}(A, B)} \quad \xrightleftharpoons{\perp} \quad \mathcal{V}\text{-}\underline{\text{Bisim}}(A, B)$$

bicategory when \mathcal{V} is
locally completely distributive

*See Walker's PhD thesis

SUMMARY

Enriched bisimulations

- generalise several common kinds of bisimulation
- are equivalence classes of spans of enriched asymmetric lenses
- are only defined for thin bases of enrichment

QUESTIONS

- What are the bisimulations for other common quantales?
- What other parts of bisimulation theory generalise?

<https://mdimeglio.github.io>
<https://bryceclarke.github.io>