

# CATEGORICAL HILBERT THEORY

MATTHEW DiMEGLIO




PhD student of Chris Heunen at the  
University of Edinburgh


My PhD  
research



My PhD  
research

Category  
Theory





Quantum  
Foundations

Category  
Theory


Functional  
Analysis

My PhD  
research

- 1 Limits are limits
- 2 Categorical Hilbert theory
- 3 Dilators

# Background


Hilbert spaces are vector spaces with geometry  
(encoded by a complete inner product)



Hilbert spaces are vector spaces with geometry  
(encoded by a complete inner product)

$$\|x\| = \sqrt{\langle x|x \rangle}$$


Lengths



Hilbert spaces are vector spaces with geometry  
(encoded by a complete inner product)

$$\|x\| = \sqrt{\langle x|x \rangle}$$

Lengths




$$\cos \theta = \frac{\text{Re} \langle x|y \rangle}{\|x\| \|y\|}$$

Angles

Hilbert spaces are vector spaces with geometry  
(encoded by a complete inner product)

$$\|x\| = \sqrt{\langle x|x \rangle}$$

Lengths



$$\cos \theta = \frac{\operatorname{Re} \langle x|y \rangle}{\|x\| \|y\|}$$

Angles

e.g.  $\mathbb{C}, \mathbb{C}^2, \dots, \ell^2(\mathbb{N})$

Adjointables are maps  $f: X \rightarrow Y$  between Hilbert spaces with an adjoint  $f^*: Y \rightarrow X$

$$\langle y | fx \rangle = \langle f^*y | x \rangle$$

Adjoints are maps  $f: X \rightarrow Y$  between Hilbert spaces with an adjoint  $f^*: Y \rightarrow X$

$$\langle y | fx \rangle = \langle f^*y | x \rangle$$

e.g. All linear maps  $\mathbb{C}^m \rightarrow \mathbb{C}^n$   
(adjoint is conjugate transpose)

Adjointables are maps  $f: X \rightarrow Y$  between Hilbert spaces with an adjoint  $f^*: Y \rightarrow X$


$$\langle y | fx \rangle = \langle f^*y | x \rangle$$

e.g. All linear maps  $\mathbb{C}^m \rightarrow \mathbb{C}^n$   
(adjoint is conjugate transpose)

They form a category **Adjointable**


Contractions are linear maps between  
Hilbert spaces that decrease lengths

Contractions are linear maps between Hilbert spaces that decrease lengths



$$\|f\mathbf{x}\| \leq \|\mathbf{x}\|$$

Contractions are linear maps between Hilbert spaces that decrease lengths




$$\|f_x\| \leq \|x\|$$

They form a subcategory **Contraction** of **Adjointable**


Isometries are maps between  
Hilbert spaces that preserve geometry

Isometries are maps between  
Hilbert spaces that preserve geometry



$$\|s\mathbf{x}\| = \|\mathbf{x}\|$$

Isometries are maps between  
Hilbert spaces that preserve geometry



$$\|sx\| = \|x\|$$


They form a subcategory **Isometry of Contraction**

1


Limits are limits

Limits in analysis are about  
approximating elements

Limits in analysis are about  
approximating elements



Limits in analysis are about  
approximating elements




$$\lim_{n \rightarrow \infty} 2^{-n} = 0$$

Codirected limits in category theory  
are about approximating objects

in the category Contraction

Codirected limits in category theory  
are about approximating objects



in the category Contraction

Codirected limits in category theory  
are about approximating objects

$$\begin{array}{ccccccc} x_1 & \longleftarrow & (x_1, x_2) & \longleftarrow & (x_1, x_2, x_3) & & \\ \mathbb{C} & \longleftarrow & \mathbb{C}^2 & \longleftarrow & \mathbb{C}^3 & \longleftarrow & \dots \end{array}$$

$$\lim_{n \in \mathbb{N}} \mathbb{C}^n = \ell^2(\mathbb{N}) = \left\{ x \in \mathbb{C}^{\mathbb{N}} \mid |x_1|^2 + |x_2|^2 + \dots < \infty \right\}$$

in the category Contraction

# AXIOMS FOR THE CATEGORY OF HILBERT SPACES

CHRIS HEUNEN AND ANDRE KORNELL

ABSTRACT. We provide axioms that guarantee a category is equivalent to that of continuous linear functions between Hilbert spaces. The axioms are purely categorical and do not presuppose any analytical structure. This addresses a question about the mathematical foundations of quantum theory raised in reconstruction programmes such as those of von Neumann, Mackey, Jauch, Piron, Abramsky, and Coecke.

No analytic  
assumptions

## AXIOMS FOR THE CATEGORY OF HILBERT SPACES

CHRIS HEUNEN AND ANDRE KORNELL

ABSTRACT. We provide axioms that guarantee a category is equivalent to that of continuous linear functions between Hilbert spaces. The axioms are purely categorical and do not presuppose any analytical structure. This addresses a question about the mathematical foundations of quantum theory raised in reconstruction programmes such as those of von Neumann, Mackey, Jauch, Piron, Abramsky, and Coecke.

No analytic  
assumptions

$\mathbb{R}$  or  $\mathbb{C}$  from codirected limits  
via black-box: Solèr's theorem

## AXIOMS FOR THE CATEGORY OF HILBERT SPACES

CHRIS HEUNEN AND ANDRE KORNELL

ABSTRACT. We provide axioms that guarantee a category is equivalent to that of continuous linear functions between Hilbert spaces. The axioms are purely categorical and do not presuppose any analytical structure. This addresses a question about the mathematical foundations of quantum theory raised in reconstruction programmes such as those of von Neumann, Mackey, Jauch, Piron, Abramsky, and Coecke.

No analytic assumptions

$\mathbb{R}$  or  $\mathbb{C}$  from codirected limits  
via black-box: Solèr's theorem

## AXIOMS FOR THE CATEGORY OF HILBERT SPACES

CHRIS HEUNEN AND ANDRE KORNELL

ABSTRACT. We provide axioms that guarantee a category is equivalent to that of continuous linear functions between Hilbert spaces. The axioms are purely categorical and do not presuppose any analytical structure. This addresses a question about the mathematical foundations of quantum theory raised in reconstruction programmes such as those of von Neumann, Mackey, Jauch, Piron, Abramsky, and Coecke.

What is the deeper connection between  
these two kinds of limits?


# KEY IDEA

Given real numbers  $0 < a_1 \leq a_2 \leq \dots \leq 1$

in the category Contraction

# KEY IDEA

Given real numbers  $0 < a_1 \leq a_2 \leq \dots \leq 1$




$$\|\chi_1\|^2 = a_1 \quad \|\chi_2\|^2 = a_2 \quad \dots$$

in the category Contraction

# KEY IDEA

Given real numbers  $0 < a_1 \leq a_2 \leq \dots \leq 1$




$$\lim_{n \in \mathbb{N}} X_n$$

$$\|x_1\|^2 = a_1 \quad \|x_2\|^2 = a_2 \quad \dots$$

in the category Contraction

# KEY IDEA

Given real numbers  $0 < a_1 \leq a_2 \leq \dots \leq 1$




$$\|x_1\|^2 = a_1 \quad \|x_2\|^2 = a_2 \quad \dots$$

in the category Contraction

# KEY IDEA

Given real numbers  $0 < a_1 \leq a_2 \leq \dots \leq 1$




$$\|x_1\|^2 = a_1 \quad \|x_2\|^2 = a_2 \quad \dots$$

$$\|x\|^2 = \sup_{n \in \mathbb{N}} a_n$$

in the category Contraction

# KEY IDEA

Given operators  $0 < a_1 \leq a_2 \leq \dots \leq 1$  on a Hilbert space  $A$



$$\chi_1^* \chi_1 = a_1 \quad \chi_2^* \chi_2 = a_2 \quad \dots$$

$$\chi^* \chi = \sup_{n \in \mathbb{N}} a_n$$

in the category Contraction

This idea yields

- insightful new proofs of the characterisations of the categories of Hilbert spaces

This idea yields

- insightful new proofs of the characterisations of the categories of Hilbert spaces
- the first characterisation of a category of finite-dimensional Hilbert spaces

DAGGER CATEGORIES AND THE COMPLEX NUMBERS:  
AXIOMS FOR THE CATEGORY OF FINITE-DIMENSIONAL  
HILBERT SPACES AND LINEAR CONTRACTIONS

MATTHEW DI MEGLIO AND CHRIS HEUNEN

ABSTRACT. We characterise the category of finite-dimensional Hilbert spaces and linear contractions using simple category-theoretic axioms that do not refer to norms, continuity, dimension, or real numbers. Our proof directly relates limits in category theory to limits in analysis, using a new variant of the classical characterisation of the real numbers instead of Solèr's theorem.

This idea yields

- insightful new proofs of the characterisations of the categories of Hilbert spaces
- the first characterisation of a category of finite-dimensional Hilbert spaces

Relevant for  
quantum  
computing

DAGGER CATEGORIES AND THE COMPLEX NUMBERS:  
AXIOMS FOR THE CATEGORY OF FINITE-DIMENSIONAL  
HILBERT SPACES AND LINEAR CONTRACTIONS

MATTHEW DI MEGLIO AND CHRIS HEUNEN

ABSTRACT. We characterise the category of finite-dimensional Hilbert spaces and linear contractions using simple category-theoretic axioms that do not refer to norms, continuity, dimension, or real numbers. Our proof directly relates limits in category theory to limits in analysis, using a new variant of the classical characterisation of the real numbers instead of Solèr's theorem.

2

# Categorical Hilbert theory

# CATEGORICAL REFORMULATIONS

Theory	Categorical setting
homological algebra	abelian categories
probability theory	Markov categories
differential geometry	tangent categories

Reformulating a theory category theoretically can

- unify and generalise known results,
- reveal new results,
- simplify it, making it more accessible.

# CATEGORICAL REFORMULATIONS

Theory

homological algebra

probability theory

differential geometry

functional analysis

Categorical setting


abelian categories

Markov categories


tangent categories

?

# $R^*$ -categories are a new categorical abstraction of algebraic aspects of Hilbert spaces



$R^*$ -categories are a new categorical abstraction  
of algebraic aspects of Hilbert spaces



$M^*$ -categories also include analytic aspects  
Articles in preparation

Let  $\mathbf{C}$  be a  $*$ -category with

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjoints}$  then

- (1) a zero object,

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjoints}$  then

(1) a zero object, ——————  $[0] = \{0\}$

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjoints}$  then

(1) a zero object, ——————  $[0] = \{0\}$

(2) binary orthonormal biproducts

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

(1) a zero object, ——————  $[0] = \{0\}$

(2) binary orthonormal biproducts, ——————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

(1) a zero object, ——————  $[0] = \{0\}$

(2) binary orthonormal biproducts, ——————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$

(3) isometric equalisers, such that

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

(1) a zero object,

$$0 = \{0\}$$

(2) binary orthonormal biproducts,

$$\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$$

(3) isometric equalisers, such that

$$\begin{cases} \text{Eq}(f,g) = \{x \in X : fx = gx\} \\ \text{with restricted inner product} \end{cases}$$

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

- (1) a zero object, ——————  $[0 = \{0\}]$
- (2) binary orthonormal biproducts, ——————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$
- (3) isometric equalisers, such that  $\begin{cases} Eq(f,g) = \{x \in X : fx = gx\} \\ \text{with restricted inner product} \end{cases}$
- (4) all isometries are kernels,

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjoints}$  then

(1) a zero object,

$$0 = \{0\}$$

(2) binary orthonormal biproducts

$$\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$$

(3) isometric equalisers, such that

$$\begin{cases} Eq(f,g) = \{x \in X : fx = gx\} \\ \text{with restricted inner product} \end{cases}$$

(4) all isometries are kernels,

$$Ker(f) = Eq(f, 0)$$

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjoints}$  then

- (1) a zero object, —————  $[0 = \{0\}]$
- (2) binary orthonormal biproducts —————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$
- (3) isometric equalisers, such that  
with restricted inner product
- (4) all isometries are kernels, —————  $\begin{cases} \text{Ker}(f) = \text{Eq}(f, 0) \\ m^*m = 1 \end{cases}$

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

- (1) a zero object, —————  $[0 = \{0\}]$
- (2) binary orthonormal biproducts —————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$
- (3) isometric equalisers, such that  
with restricted inner product
- (4) all isometries are kernels, —————  $\begin{cases} \text{Ker}(f) = \text{Eq}(f, 0) \\ m^*m = 1 \end{cases}$
- (5)  $\text{Isometry}(\mathbf{C})$  has directed colimits.

Let  $\mathbf{C}$  be a  $*$ -category with

If  $\mathbf{C} = \text{Adjointable}$  then

- (1) a zero object, —————  $[0 = \{0\}]$
- (2) binary orthonormal biproducts —————  $\begin{cases} X \oplus Y = \{(x,y) : x \in X, y \in Y\} \\ \langle (x,y) | (u,v) \rangle = \langle x|u \rangle + \langle y|v \rangle \end{cases}$
- (3) isometric equalisers, such that  
with restricted inner product
- (4) all isometries are kernels —————  $\begin{cases} \text{Ker}(f) = \text{Eq}(f, 0) \\ m^*m = 1 \end{cases}$
- (5)  $\text{Isometry}(\mathbf{C})$  has directed colimits. —————  $e.g. \ell^2(\mathbb{N})$

Let  $\mathbf{C}$  be a  $*$ -category with


- (1) a zero object,
- (2) binary orthonormal biproducts,
- (3) isometric equalisers, such that
- (4) all isometries are kernels,
- (5)  $\mathbf{Isometry}(\mathbf{C})$  has directed colimits.

Let  $\mathbf{C}$  be a  $*$ -category with

- (1) a zero object,
- (2) binary orthonormal biproducts,
- (3) isometric equalisers, such that
- (4) all isometries are kernels,
- (5)  $\mathbf{Isometry}(\mathbf{C})$  has directed colimits.

$R^*$ -category

Let  $\mathbf{C}$  be a  $*$ -category with

- (1) a zero object,
  - (2) binary orthonormal biproducts,
  - (3) isometric equalisers, such that
  - (4) all isometries are kernels,
  - (5)  $\mathbf{Isometry}(\mathbf{C})$  has directed colimits.
- 

R $*$ -category


M $*$ -category

Let  $\mathbf{C}$  be a  $*$ -category with

- (1) a zero object,
- (2) binary orthonormal biproducts,
- (3) isometric equalisers, such that
- (4) all isometries are kernels,
- (5)  $\text{Isometry}(\mathbf{C})$  has directed colimits.

## THEOREM:

If  $\mathbf{C}$  has a simple separator  $A$  then  $\mathbf{C}(A, A)$  is  $\mathbb{R}, \mathbb{C}$  or  $\mathbb{H}$  and



# THEORY

- Additivity (-1 exists)

# THEORY

- Additivity (-1 exists)
- Canonical partial order and inner products  
 $f^*f \geq 0$        $\langle f | g \rangle = f^*g$

# THEORY

- Additivity (-1 exists)
- Canonical partial order and inner products  
 $f^*f \geq 0$        $\langle f | g \rangle = f^*g$
- Symmetry (If  $a \geq 1$  then  $a$  invertible)

# THEORY

- Additivity (-1 exists)
- Canonical partial order and inner products  
 $f^*f \geq 0$        $\langle f | g \rangle = f^*g$
- Symmetry (If  $a \geq 1$  then  $a$  invertible)
- Contractions (Morphisms  $f$  with  $f^*f \leq 1$ )

# THEORY


- Additivity (-1 exists)
- Canonical partial order and inner products  
 $f^*f \geq 0$        $\langle f | g \rangle = f^*g$
- Symmetry (If  $a \geq 1$  then  $a$  invertible)
- Contractions (Morphisms  $f$  with  $f^*f \leq 1$ )
- Monotone completeness  
(Bounded increasing nets have suprema)

3

Dilators

# DEFINITION:


A codilation of  $f: X \rightarrow Y$  is  
a cospan of isometries



such that  $s_2^*s_1 = f$ .

# DEFINITION:

A codilation of  $f: X \rightarrow Y$  is a cospan of isometries



A codilator of  $f$  is an initial codilation of  $f$ .

such that  $s_2^*s_1 = f$ .

# DEFINITION:

A codilation of  $f: X \rightarrow Y$  is a cospan of isometries

$$\begin{array}{ccc} & S & \\ S_1 \nearrow & & \swarrow S_2 \\ X & & Y \end{array}$$


such that  $S_2^* S_1 = f$ .

A codilator of  $f$  is an initial codilation of  $f$ .

$$\begin{array}{ccc} & X \boxplus_f Y & \\ i_1 \nearrow & & \swarrow i_2 \\ X & & Y \end{array}$$


# DEFINITION:

A codilation of  $f: X \rightarrow Y$  is a cospan of isometries




such that  $s_2^* s_1 = f$ .

A codilator of  $f$  is an initial codilation of  $f$ .




# DEFINITION:

A codilation of  $f: X \rightarrow Y$  is a cospan of isometries



such that  $s_2^*s_1 = f$ .

A codilator of  $f$  is an initial codilation of  $f$ .



# PROPOSITION:

Every morphism in **Contraction** has a codilator

$$X \xrightarrow{f} Y$$

Inspired by minimal unitary dilations

# PROPOSITION:


Every morphism in **Contraction** has a codilator

$$\begin{array}{ccc} & X \oplus Y & \\ \swarrow & & \uparrow \\ [ \sqrt{1-f^*f} & f ] & \quad [ 0 & 1 ] \\ X & \xrightarrow{f} & Y \end{array}$$

Inspired by minimal unitary dilations

# PROPOSITION:


Every morphism in **Contraction** has a codilator



Inspired by minimal unitary dilations

# PROPOSITION:

Every morphism in **Contraction** has a codilator



Inspired by minimal unitary dilations

Other  $*$ -categories with codilators of all morphisms:

- Sets and partial bijections
- Sets and bitotal relations
- Finite probability spaces and stochastic maps  
( $f^*$  is the Bayesian inverse of  $f$ )

Monotone completeness  
of  $M^*$ -categories

Dilators are useful

Louis Lemonnier  
Semantics for symmetric  
pattern matching

Universal property for  $\oplus$   
in Contraction

Monotone completeness  
of  $M^*$ -categories

Dilators are useful

Louis Lemonnier  
Semantics for symmetric  
pattern matching

Universal property for  $\oplus$   
in Contraction

**PROPOSITION:**  
Isometry → Contraction preserves directed colimits

# PROPOSITION:

Isometry → Contraction preserves directed colimits

# PROOF:

$$X_1 \rightarrow X_2 \rightarrow \dots$$


**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits

**PROOF:**




**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits

**PROOF:**




**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits

**PROOF:**




**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits

**PROOF:**




**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits

**PROOF:**



**PROPOSITION:**  
Isometry  $\rightarrow$  Contraction preserves directed colimits


**PROOF:**



# PROPOSITION:

Isometry → Contraction preserves directed colimits

## PROOF:



# SUMMARY

- Limits are limits
- $R^*$ - and  $M^*$ -categories  
for Hilbert theory
- Dilators relate isometries  
and contractions

<https://mdimeglio.github.io>

m.dimeglio@ed.ac.uk

# SUMMARY

- Limits are limits
- $R^*$ - and  $M^*$ -categories for Hilbert theory
- Dilators relate isometries and contractions

# NEXT STEPS

- Contractions as relations in **Isometry**

<https://mdimeglio.github.io>

m.dimeglio@ed.ac.uk

# SUMMARY

- Limits are limits
- $R^*$ - and  $M^*$ -categories for Hilbert theory
- Dilators relate isometries and contractions

# NEXT STEPS

- Contractions as relations in **Isometry**
- Axioms for **Isometry**

<https://mdimeglio.github.io>

m.dimeglio@ed.ac.uk

# SUMMARY

- Limits are limits
- $R^*$ - and  $M^*$ -categories for Hilbert theory
- Dilators relate isometries and contractions

# NEXT STEPS

- Contractions as relations in **Isometry**
- Axioms for **Isometry**
- Axioms for a category of probability spaces

<https://mdimeglio.github.io>

m.dimeglio@ed.ac.uk

# Reformulating a theory category theoretically can

- **unify and generalise known results,**  
e.g. Discrete and continuous cases of the Blackwood–Sherman–Stein theorem  
on statistical experiments unified for the first time via Markov categories
- **reveal new results,**  
e.g. convergence in mean for backward martingales indexed by an arbitrary net  
was proved for the first time (according to the authors) using dagger categories
- **simplify it, making it more accessible.**  
e.g. to theoretical computer scientists who already know category theory

# PROPOSITION: Every morphism in FinPS has a dilator

bloom-shrieg factorisation  
“The information loss  
of a stochastic map”

coisometric  
 $\Leftrightarrow$   
deterministic

